LA--10467-MS

DE86 001226

The Beta Spectra of $^{90}_{38}\text{Sr}$ and $^{90}_{39}\text{Y}$

Joseph J. Devaney

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

THE BETA SPECTRA OF $^{90}_{38}\text{Sr}$ AND $^{90}_{39}\text{Y}$

by

Joseph J. Devaney

ABSTRACT

Using as a base the beta energy spectra provided by T. R. England, the number spectra of $^{90}_{38}\text{Sr}$ and $^{90}_{39}\text{Y}$ were corrected for the Coulomb effects on the first forbidden shape factor using the tables of M. E. Rose, and for higher order effects using the experimentally determined corrections of H. Daniel et al. We, therefore, present a fully corrected beta number spectrum of $^{90}_{38}\text{Sr}$ and its daughter $^{90}_{39}\text{Y}$. The half-lives and end-point energies are also listed.

I. INTRODUCTION

This note gives the beta spectra of ^{90}Sr and its daughter ^{90}Y . We give the spectra fully corrected for Coulomb and relativistic effects, as well as higher order effects ($< 4\%$), as of 1964. It is, however, not completely clear that corrections of less than, say, 10% are significant. The data do seem to justify accuracies of better than 2%, and calculations (i.e. MC type) can be made using the data, to accuracies of 5% or better, so we decided to do the full calculation and let the user take his preference. We also have computed the spectra on a perhaps needlessly fine energy mesh in order to match the basic spectral data of T. England. Again the user has, therefore, a full choice; he may use the full detail or condense it as he sees fit. The isotope $^{90}_{38}\text{Sr}$ decays* 100% by negatrons, half-life 28.6Y, $Q = 546.0$ keV, to the ground state of $^{90}_{39}\text{Y}$, i.e., no gammas. The isotope $^{90}_{39}\text{Y}$, 64.1 h, decays¹ 99.9885% by negatrons to the ground state of $^{90}_{40}\text{Zr}$, $Q = 2279.2$ keV. We neglect the 0.0115%

*Information provided by T. R. England, Group T-2, Los Alamos National Laboratory (July 1984).

transition by β^- decay to the 1760.7 keV excited state of ^{90}Zr . No 1760.7 keV γ is observed, (e-pairs, 2 γ 's, and conversion electrons are observed). Thus the spectrum of $^{90}\text{Y} \beta^-$ given here is for 100%, $Q = 2279.2$ keV decay. Both decays, $^{90}\text{Sr} \beta^-$ and $^{90}\text{Y} \beta^-$, are first forbidden unique.^{1,2} The ground state of the granddaughter, $^{90}_{40}\text{Zr}$, is stable.¹

If N_i is the amount of isotope i , then the amount of $^{90}_{38}\text{Sr}$ is N_{38} , etc., and after time t

$$N_{38} = N_{38-0} \exp(-\lambda_{38}t) \quad , \quad (1)$$

where N_{38-0} is the original amount of ^{90}Sr . λ_{38} is the ^{90}Sr decay constant which is related to the half-life, $T_{1/2}$, by:

$$\lambda = \ln 2 / T_{1/2} \quad . \quad (2)$$

From the buildup and decay of N_{39} we have:

$$dN_{39}/dt = \lambda_{38} N_{38} - \lambda_{39} N_{39} \quad . \quad (3)$$

Substituting Eq.(1) and solving yields

$$N_{39} = N_{38-0} \left(\frac{\lambda_{38}}{\lambda_{39} - \lambda_{38}} \right) (e^{-\lambda_{38}t} - e^{-\lambda_{39}t}) \quad , \quad (4)$$

as is well known. The $^{90}_{38}\text{Sr}$ and $^{90}_{39}\text{Y}$ activities are

$$A_{38} = \lambda_{38} N_{38} \quad (5a)$$

and

$$A_{39} = \lambda_{39} N_{39} \quad , \quad (5b)$$

whence using Eqs. (1) and (4), we get

$$A_{39} = \left(\frac{\lambda_{39}}{\lambda_{39} - \lambda_{38}} \right) A_{38} [1 - e^{-(\lambda_{39} - \lambda_{38})t}] \quad . \quad (6)$$

At 28.6 y half-life (250708. h), $\lambda_{38} = \lambda_{39} (64.1/250708) = 2.557 \times 10^{-4} \lambda_{39}$ so that $\lambda_{39} - \lambda_{39} \approx \lambda_{39}$ to 0.026% and Eq. (6) becomes

$$\begin{aligned} A_{39} &\approx A_{38} [1 - e^{-\lambda_{39}t}] = A_{38} [1 - 2^{(T_{\frac{1}{2}} - 39/t)}] \\ &\approx A_{38} [1 - (0.5)^{(T_{\frac{1}{2}} - 39/t)}] . \end{aligned} \quad (7)$$

Thus, at $t = 64.1$ h, the activity of $^{90}_{39}\text{Y}$ is already 50% of $^{90}_{39}\text{Sr}$ and it is 90% A_{39} at 8.87 days and $A_{38} = 99\% A_{39}$ at 17.75 days and 99.9% at 26.62 days. The times, t , are the times after a pure sample of ^{90}Sr is isolated.

When $A_{38} \approx A_{39}$ to the extent desired, ^{90}Y and ^{90}Sr are said to be in "secular equilibrium."

The beta spectrum of a sample of ^{90}Sr in secular equilibrium with its daughter ^{90}Y is simply obtained by adding the two spectra. The result normalizes to a total of two (2) electrons per ^{90}Sr decay, i.e. at the rate:

$$A_{38} (\approx A_{39}) = (\ln 2 / T_{\frac{1}{2}} - 38) N_{38} , \quad (8)$$

with the electron rate being $2A_{38}$.

II. BRIEF GENERAL REMARKS ON THE THEORY OF BETA DECAY

Negatron (i.e. β^-) decay is the following process:

$$n \rightarrow p + e^- + \bar{\nu} , \quad (9)$$

where n is a neutron, p a proton, e^- a negative electron, and $\bar{\nu}$ an anti-neutrino. For negatron decay of a nucleus, the process (9) takes place within the nucleus and is, therefore, limited by the energetically available nuclear states. The basic idea giving the energy spectrum of the electron is as simple as possible: the spectrum is just that given by the phase space (i.e. momentum and configuration space) of the products. Although the total decay energy is fixed, the spectrum is continuous because in a three-body decay the conservation of momentum and energy is not restrictive enough to fix all

energies and directions as momentum and energy do for two-body decay. The simple phase space result gives, therefore, an expression of the form

$$(dN/dp_e dp_{\bar{\nu}}) = \text{const } p_e^2 dp_e d\Omega_e dV_e p_{\bar{\nu}}^2 dp_{\bar{\nu}} d\Omega_{\bar{\nu}} dV_{\bar{\nu}} \quad , \quad (10)$$

where p is momentum, Ω solid angle, and V configuration volume for each of the electrons and antineutrinos.

The volume element in three-dimensional momentum space is of course $p^2 dp d\Omega$. If one is interested only in the momentum spectrum of the electron, integration and manipulation of Eq. (10) yield

$$(dN/dp_e) = \text{const. } p_e^2 (E_0 - E_e)^2 \quad , \quad (11)$$

where E_0 is the total decay energy and E_e the electron energy. Eq. (11) is

the basic reason why β -spectra are plotted in the form $\sqrt{\frac{dN}{dp_e} \cdot \frac{1}{p_e^2 F}}$ versus E_e

which plot is then a straight line. F is a function of Z (atomic number) and E to account for some of the things this simple discussion has left out. Such plots are called Fermi or Kurie (with a K!) plots.

Unfortunately, for the user of beta-spectra, life is not so simple. There are three basic reasons:

1. We have not accounted for the Coulomb interaction between the electron and the proton in Eq. (9),
2. The decay is not that of Eq. (9) but rather that of Eq. (12) below with much more complicated parent and daughter,

3. The antineutrino is always relativistic and the electron is effectively relativistic above, say, 20 keV or so.

Item 3 brings the usual complicated algebra of relativity, but 1 and 2 bring a host of complications. In the first place, the daughter nucleus has charge $(Z + 1)e$ and so the negatron kinetic energy is reduced. Thus the β^- spectra are shifted to lower energies by this "Coulomb correction." Second the spins, orbital angular momentum, and parity of the initial and final nuclear states may not match up in the optimum way so leading to less likely

decay probabilities and also to different spectra. Indeed there are classes of this sort of thing called allowed (roughly 0 orbital angular momentum), first forbidden (orbital angular momentum 1), second forbidden, etc. Crudely we may think of nonzero electron orbital angular momentum as providing a centrifugal force on the electron which in quantum mechanics equates to a low probability of having a wave-function in the nucleus where the electron must be born and so a lower probability of β -decay. The orbital angular momentum modifies the phase space and so the electron distribution. Again these modifications must be Coulomb and relativistically corrected.

The matrix elements of the beta transitions are products (and sums thereof) of matrix elements in the nuclear and leptonic spaces. It turns out that there are only five relativistically invariant interaction Hamiltonians:³ Scalar, Vector, Tensor, Axial or Psuedo-vector, and Psuedo-scalar. It further happens that these nuclear elements are reducible to only two dominant terms for allowed transitions: the Fermi (Scalar or Vector) or the Gamov-Teller (Tensor or Axial-vector) corresponding to the transition operators 1 and σ respectively. The "unique" forbidden transitions are those which arise from the presence of only Gamov-Teller couplings and depend on only one nuclear moment³ $(\int \beta_0 \cdot \vec{p} \, y_{lm}(\vec{r}))$ and have only one spectrum shape factor. The decays of both ^{90}Sr and ^{90}Y are first forbidden unique.

The spectra of ^{90}Sr and ^{90}Y are, therefore, determined unless there are higher order contributions,² i.e. "regular" third forbidden contributions, weak magnetism terms, and perhaps other terms. These we shall account for from experiment.² They are small (< 2% to < 4%, ^{90}Y and ^{90}Sr , respectively).

The most convenient and extensive source of beta spectra known to this writer is the tables of Tal England, Los Alamos Group T-2,* which are based upon ENDF/B-IV data. His calculation uses a simple yet accurate approximation for the spectra as reported in Stamatelatos and England.⁴ For most purposes, use of the England Table is accurate enough. It does provide screened Coulomb corrections for the allowed part and is relativistic. The data are given in the form of group β energy values, i.e. $E(dN/dE) \Delta E$, versus the group energy

*Provided by T. R. England, Group T-2, Los Alamos National Laboratory (July 1984).

boundaries. Division by the group mid-point energies yields the particle number distribution, $\Delta N = (dN/dE) \Delta E$, to good accuracy as checked against the normalization. Further corrections applicable to the England work are the Coulomb corrections to the forbidden shape factor and higher order corrections. We make these corrections below (Sections III and IV). They amount to about $\pm 10\%$ at the spectrum ends for ^{90}Sr , 0 in the middle; and -3 and $+5\%$ for ^{90}Y , also 0 in the middle. It is not completely clear that users will want to calculate to the above accuracies or that the data can be trusted to those accuracies, although the Daniel et al.² data appear to be good to about $\pm 1.5\%$ with statistical errors usually about $\pm 0.4\%$ or so.

The Coulomb correction to the first forbidden shape replaces the first forbidden factor,^{3,4} $(p^2 + q^2)$, by the factor, $(L_0 q^2 + 9 L_1)$, where L_0 and L_1 are given in various tables.^{5,6} In Appendix A we show that the L_1 of Dzhelepov⁶ is the (L_1/L_0) of Rose.⁵ Daniel² and Dzhelepov⁶ have absorbed L_0 into the function $F(z,w)$, i.e. $F(z,w)$ of Refs. 2 and 6 is equal to L_0 times the usual $F(z,w)$. The quantity p is the relativistic electron momentum and q is the relativistic antineutrino momentum.

III. THE β^- SPECTRUM OF $^{90}_{38}\text{Sr}$

Our base reference spectra are those of T. R. England,* which are given in the form of energy group boundaries and beta energy per group, i.e. $E \cdot (dN/dE) \Delta E$, where N is the number of betas (or their probability), and E is beta kinetic energy. To form the number-density spectrum, $\Delta N = (dN/dE) \Delta E$, we divide the energy per group by the group mid-energy to form:

$$\Delta N = [E(dN/dE)\Delta E]/E_{\text{mid-energy}} \quad (13)$$

These are listed as a function of $E_{\text{mid-energy}}$ in column 3 of Table I. The sum, which should equal 1, gives a measure of the overall accuracy (0.6%) of the method. Column 1 gives the group beta energy boundaries and column 2 gives the group mid-point energies.

*Provided by T. R. England, Group T-2, Los Alamos National Laboratory (July 1984).

TABLE I

⁹⁰Sr β -Spectrum Analysis and Results, ($E_0 = 546$ keV)

Energy Group Boundaries (keV)	Group Mid-energy (keV)	$\Delta N = (dN/dE) \Delta N$ group = (England)/ E_{mid}	Unique first forbidden correc- tion factor (Coulomb)	Higher Order correction factor	ΔN corrected unnormalized	ΔN corrected normalized
0						
	5	0.02486	1.0557	1.0367	0.02721	0.02700
10						
	15	0.03324	1.0505	1.03556	0.03616	0.03588
20						
	25	0.03313	1.0456	1.0344	0.03583	0.03555
30						
	35	0.03304	1.0404	1.0332	0.03552	0.03524
40						
	45	0.03294	1.0353	1.0321	0.03520	0.03492
50						
	55	0.03283	1.0306	1.0309	0.03488	0.03461
60						
	65	0.03269	1.0260	1.0298	0.03454	0.03427
70						
	75	0.03252	1.0210	1.0286	0.03415	0.03388
80						
	85	0.03231	1.0162	1.0274	0.03373	0.03347
90						
	95	0.03205	1.0113	1.0263	0.03326	0.03300
100						
	125	0.15441	0.9971	1.0228	0.15747	0.15624
150						
	175	0.14098	0.9768	1.0170	0.14005	0.13895
200						
	225	0.12249	0.9590	1.0112	0.11876	0.11783
250						
	275	0.10019	0.9425	1.0054	0.09494	0.09420
300						
	325	0.07573	0.9392	0.9996	0.07110	0.07054
350						
	375	0.05113	0.9311	0.9938	0.04731	0.04694
400						
	425	0.02873	0.9261	0.9880	0.02629	0.02608
450						
	475	0.011188	0.9240	0.9823	0.010155	0.010076
500						
	525	0.001476	0.9232	0.9765	0.001331	0.001321
550						
	SUM	1.00593			1.007886	0.99999

Column 4 gives the Coulomb corrections to the unique, first-forbidden shape factor from the expression

$$[q^2 + 9(L_1/L_{O-Rose})]/(q^2 + p^2) \quad , \quad (14)$$

where the quantities L_1 and L_0 were interpolated from the tables of Rose.⁵ Eq. (14) is not normalized. [Strictly: Eq. (13) is not adjusted so that its product with ΔN is normalized.] We will compute our completely corrected spectrum and then normalize it.

Column 5 gives the higher order corrections, C_1 , found experimentally by Daniel et al.² and given in the form (normalization adjusted):

$$C_1 = 1.0965 (1 - 0.054 W) \quad , \quad (15)$$

where W is the full relativistic beta energy in terms of the electron rest mass, $m_e c^2$:

$$W = 1 + (E/m_e c^2) \quad . \quad (16)$$

The two corrections, columns 4 and 5, are multiplied together and then multiplied by the number spectrum of column 3 to yield the corrected but unnormalized spectrum of column 6. Column 7 gives the normalized spectrum and is our result for the beta number spectrum, $\Delta N = (dN/dE) \Delta E$, of ^{90}Sr , including all corrections known to us.

Note that the spectrum is relatively flat at low beta energies and drops off above 200 keV. The combined corrections range from + 8.6% at 5 keV to -1.4% at 175 keV to - 10.5% at 525 keV. Thus, if, say, $\pm 9\%$ is significant, we recommend the use of column 7 versus column 2 (ΔN vs. E_{mid}) for the spectrum, otherwise column 3 versus 2 (ΔN vs. E_{mid}) is satisfactory.

IV. THE β^- SPECTRUM OF $^{90}_{39}\text{Y}$

Again the base reference spectrum is that of England. Divided by the group mid-point energy (column 2, Table II), the resulting number spectrum per Eq. (13) is found in Table II, column 3. The overall accuracy of the method is indicated by the sum of column 3 entries and is 0.1%.

Using Eq. (14) the (unnormalized) correction factors due to Coulomb corrections to the first forbidden shape factor are given in column 4, Table II.

The experimental higher order corrections from Daniel et al.² are given by (normalization adjusted)

$$C_1 = 1.0270 (1 - 0.0072 W) , \quad (17)$$

with the energy, W , given by Eq. (16). We note that another source for the corrections to the β -spectrum of ^{90}Y is that of Langer, Spejewski, and Wortman⁷ (also 1964). The Langer et al. correction is given as the full correction to the allowed shape, i.e. it includes factors of the form $(q^2 + p^2)$ times Eq. (14) times Eq. (17).

As for ^{90}Sr , the two corrections, columns 4 and 5, are multiplied together and then are multiplied by the number spectrum of column 3 to yield the corrected but unnormalized spectrum of column 6. Column 7 gives the normalized spectrum and is, therefore, our result for the fully corrected beta number (or probability) spectrum, $\Delta N = (dN/dE) \Delta E$, of ^{90}Y . The combined spectrum of ^{90}Sr and ^{90}Y in secular equilibrium, per ^{90}Sr disintegration is the sum of columns 7 of Tables I and II.

Note, in contrast to ^{90}Sr , that the ^{90}Y spectrum has a maximum more in the center at 825 keV; thus the corrections already smaller than for ^{90}Sr are of relatively less impact, since they are largest where the number densities are least, namely, at the ends of high and low beta energy. The combined ^{90}Y corrections to the England derived number spectrum of column 3, Table II, are + 5.1% at 5 keV beta energy, + 4.2% at 175 keV, + 2.2% at 525 keV, + 0.1% at 825 keV, -2.3% at 1275 keV, -3.3% at 1675 keV, -3.0% at 1975 keV, -3.2% at 2075 keV, and -2.3% at 2275 keV.

ACKNOWLEDGMENTS

It is most pleasant to acknowledge the extraordinary helpfulness I received in preparing this report. I have already detailed the assistance of Professors Rolf M. Steffins and Herbert Daniel in Appendix A. To them I would like to add: T. R. England, B. J. Dropesky, M. E. Bunker, and H. Smith.

TABLE II

⁹⁰Y β -Spectrum Analysis and Results, ($E_0 = 2279.2$ keV)

Energy Group Boundaries (keV)	Group Mid-energy (keV)	$\Delta N = (dN/dE) \Delta N$ group $= (\text{England})/E_{\text{mid}}$	Unique first forbidden correc- tion factor (Coulomb)	Higher Order correction factor	ΔN corrected unnormalized	ΔN corrected normalized
0						
10	5	0.001828	1.0033	1.0195	0.001871	0.001921
20	15	0.002562	1.0030	1.0194	0.002620	0.002690
30	25	0.002627	1.0028	1.0192	0.002685	0.002756
40	35	0.002697	1.0024	1.0191	0.002755	0.002828
50	45	0.002722	1.00215	1.0190	0.002779	0.002853
60	55	0.002848	1.00185	1.0188	0.002907	0.002984
70	65	0.002927	1.0016	1.0187	0.002986	0.003065
80	75	0.003007	1.00125	1.0185	0.003066	0.003148
90	85	0.003088	1.00098	0.0184	0.003148	0.003232
100	95	0.003170	1.0007	1.0182	0.003228	0.003314
150	125	0.017149	0.9998	1.0178	0.017451	0.017916
200	175	0.019185	0.9981	1.0171	0.019476	0.019994
250	225	0.021180	0.99643	1.0163	0.021450	0.022021
300	275	0.023104	0.9943	1.0156	0.023331	0.023948
350	325	0.024934	0.9927	1.0149	0.025121	0.025790
400	375	0.026655	0.9902	1.0142	0.026768	0.027481
450	425	0.028251	0.9879	1.0135	0.028285	0.029038
500	475	0.029710	0.9858	1.0127	0.029661	0.030451
550	525	0.031022	0.9834	1.0120	0.030873	0.031695
600	575	0.032177	0.9808	1.0113	0.031915	0.032765

TABLE II (cont) 90Y β

Energy Group Boundaries (keV)	Group Mid-energy (keV)	$\Delta N = (dN/dE) \Delta N$ group = (England)/ E_{mid}	Unique first forbidden correc- tion factor (Coulomb)	Higher Order correction factor	ΔN corrected unnormalized	ΔN corrected normalized
	625	0.033166	0.97823	1.0106	0.032787	0.033660
650	675	0.033983	0.9756	1.0098	0.033480	0.034371
700	725	0.034622	0.9730	1.0091	0.033994	0.034899
750	775	0.035080	0.9704	1.0084	0.034327	0.035241
800	825	0.035355	0.9678	1.0077	0.034479	0.035397
850	875	0.035445	0.9654	1.0069	0.034456	0.035373
900	925	0.035351	0.9630	1.0062	0.034254	0.035166
950	975	0.035074	0.9607	1.0055	0.033881	0.034762
1000	1025	0.034618	0.9586	1.0048	0.033343	0.034231
1050	1075	0.033987	0.9566	1.0040	0.032644	0.033513
1100	1125	0.033187	0.9546	1.0033	0.031785	0.032631
1150	1175	0.032223	0.95295	1.0026	0.030787	0.031607
1200	1225	0.031106	0.95156	1.0019	0.029655	0.030444
1250	1275	0.029844	0.95037	1.0012	0.028396	0.029152
1300	1325	0.028448	0.9493	1.0004	0.027017	0.027736
1350	1375	0.026930	0.9486	0.9997	0.025538	0.026218
1400	1425	0.025303	0.9480	0.9990	0.023963	0.024601
1450	1475	0.023583	0.9475	0.9983	0.022306	0.022900
1500	1525	0.021784	0.9471	0.9975	0.020581	0.021129
1550	1575	0.019925	0.94683	0.9968	0.018805	0.019306
1600	1625	0.018023	0.94678	0.9961	0.016997	0.017449
1650	1675	0.015098	0.9468	0.9954	0.015171	0.015575
1700	1725	0.014181	0.9471	0.9946	0.013359	0.013715

TABLE II (cont) 90Y β

Energy Group Boundaries (keV)	Group Mid-energy (keV)	$\Delta N = (dN/dE) \Delta N$ group = (England)/E _{mid}	Unique first forbidden correc- tion factor (Coulomb)	Higher Order correction factor	ΔN corrected unnormalized	ΔN corrected normalized
1750						
	1775	0.012334	0.9476	0.9939	0.011616	0.011925
1800						
	1825	0.010399	0.9483	0.9932	0.009795	0.010056
1850						
	1875	0.008603	0.9501	0.9925	0.008112	0.008328
1900						
	1925	0.006901	0.9520	0.9918	0.006515	0.006688
1950						
	1975	0.005318	0.9531	0.9910	0.005024	0.005158
2000						
	2025	0.003885	0.9532	0.9903	0.003667	0.003765
2050						
	2075	0.002630	0.95283	0.9896	0.002479	0.002545
2100						
	2125	0.001583	0.9552	0.9889	0.001495	0.001535
2150						
	2175	0.000777	0.95875	0.9881	0.000736	0.000755
2200						
	2225	0.000244	0.9619	0.9874	0.000232	0.000238
2250						
	2275	0.0000174	0.96455	0.9867	0.0000166	0.0000170
2300						
TOTAL SUM		1.001007			0.9740686	0.999997

APPENDIX A

COMPARISON OF FIRST FORBIDDEN COULOMB CORRECTIONS

Our reference² for higher order corrections referred to Dzhelepov and Zyrianova⁶ for the Coulomb corrections to the first forbidden spectrum shape factor. Unfortunately the Dzhelepov book was not to be had in Los Alamos and, therefore, not available for some weeks. Yet the Daniel et al. reference² specifically advised that their Coulomb correction was different than usual and only to be had from Dzhelepov. By dint of considerable asking around, this writer located a former student (H. Smith, Q-1) of one of the world's experts in beta decay who retired from Purdue University to Tesuque, N.M., and is presently a consultant to Group P-2 of the Los Alamos National Laboratory. This expert, Prof. Rolf M. Steffen, although he gave most of his library to Purdue, found that he had kept the Dzhelepov and Zyrianova book because it was a personal gift from Dzhelepov, and most graciously volunteered to change his consulting day in order to bring the book to Los Alamos for me the very next day. I am deeply grateful to Prof. Steffen for his kindness.

In addition, serendipitously, one contact, Bruce Dropesky, INC-11, advised me that H. Daniel (principal author, Ref. 2) was then consulting at the Los Alamos Meson Physics Facility. A conversation with Prof. Daniel elicited the fact that the Dzhelepov tables were listed under the parent rather than (the actual Coulomb field that the beta sees) of the daughter. Moreover, he suggested that Landolt-Bornstein (Tabelle im Physik und Chemie) also contains Coulomb corrections. I am also grateful to Prof. Daniel for his help.

Because the tables of Dzhelepov and Zyrianova (No. 24) are not readily available, and because the data are spaced far apart in energy (1, 100, 400, 1000 keV), it was advantageous to relate their table to the more detailed and readily available table of Rose.⁵ With Dzhelepov's book in hand and plus the obvious fact there that the quantity q^2 had no coefficient in the first forbidden shape factor normally given as $(L_0 q^2 + 9L_1)$, it was clear that Dzhelepov's L_1 from $(q^2 + 9L_1)$ satisfied

$$L_{1\text{Dzhelepov}}^+ = (L_1/L_0)_{\text{Rose}} \quad , \quad (\text{A1})$$

which we checked from the respective tables^{5,6} for Z_{daughter} (Rose) = 40 and Z_{parent} (Dzhelepov⁺) = 39. The results, in Table AI, agree to the limits of

the tabular accuracy, except at 1 keV which we ascribe to slightly different calculational algorithms in a sensitive (high Coulomb field) but unimportant-for-us part of the spectrum. With the confidence substantiated by Table AI, we used the tables of Rose⁵ plus the experiments of Daniel et al.² to correct England, the results of which are found in Sections III and IV.

TABLE AI

COMPARISON OF COULOMB CORRECTION TERM, L_1 , FROM DZHELEPOV AND L.N.ZYRIANORA
AND L_1/L_0 FROM M. E. ROSE, $Z_{\text{DAUGHTER}} = 40$ (DZHELEPOV + LIST ACCORDING TO
PARENT, $Z = 39$ HERE), FOR β^- DECAY

β^- energy (keV)	β^- momentum ($m_e c$)	L_1 Dzhelepov ⁺	L_1/L_0 Rose
1	0.0626	8.67(3)	8.59(-3) at momentum 0.1
100	0.656	4.92 ₅ (-2)	4.92(-2)
400	1.476	0.223	0.224
1000	2.783	0.798	0.798

REFERENCES:

1. D. C. Kocher, "A = 90," Nuclear Data Sheets, 16, (1975) 55.
2. H. Daniel, G. Th. Kaschl, H. Schmitt, and K. Springer, "Shapes of Allowed and First-Forbidden β -Ray Spectra: I^{114} , K^{42} , Rb^{86} , Sr^{90} , and Y^{90} ," Phys. Rev., 136B (1964) 1240.
3. E. Konopinski, "The Theory of Forbidden β -Decay" in K. Siegbahn, ed, Beta- and Gamma Ray Spectroscopy North-Holland, (Amsterdam, 1955).
4. M. G. Stamatelatos and T. R. England, "Beta-Energy Averaging and Beta Spectra," Los Alamos Scientific Laboratory report, LA-6445-MS, (August 1976).
5. M. E. Rose, Appendix III of Ref. 2 above, "Tables of Forbidden β -Decay Functions." or M. E. Rose, C. L. Perry and N. M. Dismuke, "Tables for the Analysis of Allowed and Forbidden Beta Transitions," Oak Ridge National Laboratory report ORNL-1459, (February 1953).
6. B. S. Dzhelepov and L. N. Zyrianova, The Influence of the Atomic Electron Field on The Beta Decay, Akad. NAUK SSSR, (Moscow, 1956). Note that values are listed in terms of the parent nucleus rather than the usual daughter.*
7. L. M. Langer, E. H. Spejewski, and D. E. Wortman, "Precise Shape Measurements of Beta Spectra of Y^{90} and Y^{91} ," Phys. Rev., 135B (1964) 581.

*Information provided by H. Daniel, Visiting Scientist at Los Alamos. Physik Department, E18, Fakultät für Physik, Technische Universität München, West Germany.