DOE/ER/54087---T1

DE92 016218

FINAL TECHNICAL REPORT

for

Received by (ISI)

JUN 26 1992

LARGE ORBIT MAGNETIC CONFINEMENT SYSTEMS

FOR ADVANCED FUSION FUELS

DOE Contract No. DE-FG03-90ER54087

April 1, 1990 - February 29, 1992

Principal Investigator

Norman Rostoker

Department of Physics

University of California

Irvine, California 92717

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *ep*

TABLE OF CONTENTS

	Page
1. Summary	i
2. Confinement of Non-Adiabatic Vlasov-Maxwellian Plasmas Brett Spivey, Ph.D. Thesis 1992	ii
3. Magnetic Fusion with High Energy Self-Colliding Ion Beams, N. Rostoker, F. Wessel, B. Maglich and A. Fisher, Invited Paper at BEAMS '92 Conference, May 25-29, 1992.	80

SUMMARY

The objective of the grant/contract was to illuminate the problem of magnetic confinement for plasmas where the majority of ions have large gyro-orbits and do not obey adiabatic particle dynamics. The electrons are adiabatic. (There are many papers on plasmas where large orbit ions are a minority).

We considered a class of equilibria where large orbit ions dominate; the equilibria are rigorous solutions of the Vlasov/Maxwell equations. For a simple case - the infinitely long, low beta, rotating plasma a complete stability analysis was carried out. This problem was the basis of the first paper on finite Larmor radius stabilization [Rosenbluth, Krall and Rostoker, Nuclear Fusion, 1962 Supplement, Part 1, 143 (1962)]. In that paper an expansion in $\epsilon = \bar{a}_i / r_o$ was carried out to the first significant order beyond M.H.D. [$\bar{a}_i$ is the mean ion gyroradius and r_o is the characteristic radius of the plasma]. In this report the same problem is solved to all orders in ϵ . While this case is of limited applicability to experiments it is rigorous and without approximations, so that it can be used to verify approximations to be developed for more complex and useful cases. The application of the results to date to small fusion reactors is described in the second paper which was written after the termination of the contract, but is based in part on material developed during the contract period.

Abstract of the Dissertation

Confinement of Non-adiabatic Vlasov-Maxwellian Plasmas

by

Brett Spivey

Doctor of Philosophy in Physics

University of California, Irvine 1992

Professor Norman Rostoker, Chair

Adiabatic plasma physics, whereby particles gyrate tightly around magnetic field lines, constitutes a well studied area of plasma physics. On the other hand, plasmas in which a significant number of particles have orbits comparable to the confinement scale have many interesting properties which are not yet well understood. In particular, high energy particles appear to ignore turbulence[1], and do not behave like fluids. MHD equations cannot generally be adapted to modeling of non-adiabatic plasma; for example, the field reversed configuration is predicted in the MHD model to have a tilt mode instability which is not experimentally observed[2]. We consider in this dissertation the general case of arbitrary orbit size. Specifically, the solutions which we consider are drifted Maxwellian distributions which satisfy the Vlasov equation, and we call these Vlasov-Maxwellian (or V-M) solutions. The Maxwellian form for the distribution function is chosen for its immunity to ion-ion collisional diffusion, which becomes more significant as Larmor radius is increased.

We impose the condition that all Maxwellian ion species have common temperature and drift velocity so that the ion-ion collision operator collapses to zero. In section 2 we derive self-consistent equilibria for V-M plasmas. First we show that all V-M plasmas must be rigid rotors. Next we derive the V-M equilibrium equation for a single ion species. Finally we numerically evaluate several solutions in finite geometry. Solutions are found which are field reversed and have high density contrast. In section 3 an electrostatic linear stability analysis is carried out for a restricted class of V-M plasmas. The cases considered involve cylindrical geometry and are finite in the radial direction. A dispersion relation is obtained which is sufficiently simple that a complete investigation of finite orbit stability can be carried out. Both analytic and numerical stability analyses are presented.

Contents

1	Introduction	1
2	Vlasov-Maxwellian Equilibria	5
2.1	Steady state Vlasov-Maxwellian solutions	6
2.2	Numerical calculation of solutions	7
2.3	Low-beta equilibrium	10
3	Stability of Low-beta Vlasov-Maxwellian Solutions	13
3.1	Integration over orbits	13
3.1.1	Orbit Equations	14
3.1.2	Closure by forming charge density	17
3.1.3	Eigenmode solutions	18
3.1.4	Radial modes	20
3.2	Evaluation of Stability from Eigenvalue Equation	20
3.2.1	Evaluation of ϵ	25
3.2.2	Evaluation for $ \omega \rightarrow \infty$	26
3.2.3	Electron dielectric constant	27
3.2.4	Plasma behavior near electron plasma waves	28
3.3	Analytical Approximate Methods for Moderate Frequencies	30
3.3.1	Adiabatic Modes	35
3.3.2	Low Temperature, Large Orbit Modes	38
3.3.3	Regular Modes	39
3.3.4	Wall Modes	41
3.4	Numerical Solution	43
4	Conclusions	47
4.1	Rigid Rotor Condition	47

4.2	Equilibrium Solutions	47
4.3	General Stability Observations	48
4.4	High Frequency Instability	49
4.5	Classification of Instabilities	49
4.6	Connection with Previous Results for MHD Modes	50
4.7	Suggested Future Research	52
A	Appendix	56
B	Appendix	66

*Conference paper Cycled
Separately.*

List of Figures

1-1	Comparison between small and large orbit dynamics.	2
2-1	Shallow solution near bifurcation; density at maximum = e^3 times density at wall.	11
2-2	Deep solution near bifurcation; density at maximum = $e^{2.1}$ times density at wall.	11
2-3	Deep solution away from bifurcation; density at maximum = $e^{5.8}$ times density at wall.	11
3-1	Typical orbits vs. the parameter α	16
3-2	Radial eigenmode values ν vs. m , $r_w = \infty$	21
3-3	Radial eigenmode values ν vs. m , $r_w = 4r_0$	21
3-4	Characteristics of $\epsilon(\omega)$. Illustrated is a typical case for the real part of ϵ	23
3-5	Contour used for Nyquist integral.	24
3-6	$\omega_{\pm}$ versus m and ν . $\omega_{\pm}$ is reported in units of $\Delta\omega_e$	29

List of Tables

3-1	The peak interaction function $\mathcal{A}$ vs. z_e , B , C . δ is picked so that $\mathcal{A}$ is real and positive. The symbol *** indicates no solution exists for $-5 < \delta < 5$	34
3-2	Most negative value of $\mathcal{F}$ vs. α_i . The values of ν correspond to the case $r_w = 4r_0$	42

1 Introduction

Magnetic confinement has been studied for nearly 40 years. Almost all systems considered are based on the premise that a particle's Larmor radius is small compared to the size of the system and therefore that all particles obey adiabatic dynamics. The physical difference between small and large orbit particles is considerable. Illustrated at the top of Figure 1-1 is a Tokamak with small orbit particles which obey adiabatic dynamics. To first approximation the particles follow helical field lines; indeed that is why they are confined. A Betatron is also illustrated in Figure 1-1. The large orbit non-adiabatic particles are confined by the indicated magnetic field, and move across the magnetic field lines. To confine small orbit particles in a Tokamak, it is essential that the safety factor q be greater than unity,

$$q = \frac{2\pi}{\Delta\theta} = \frac{rB_t}{RB_p} > 1$$

where $\Delta\theta$ is the rotational transform illustrated in Figure 1-1, r is the minor radius, R the major radius, B_t the toroidal magnetic field, and B_p the poloidal magnetic field. In contrast, the confinement of large orbit particles with $q < 1$ has been observed for electrons[3,4,5] and for ions[6].

There have been many investigations of the stability of plasmas with a small number of large orbit particles in a background of adiabatic ions and electrons. These investigations began with the Astron[7] and high energy electrons forming an E-layer. High energy ions were also investigated[8,9,10,11,12] in a dense plasma background. More recently there has been interest in the effect of energetic alpha particles produced from D-T fusion on the stability of a burning plasma.

In this dissertation we consider a plasma where most of the ions are non-adiabatic. Experimentally such a plasma has been studied[13,14,15,16] for small values of the quantity $\beta = 8\pi nkT/B^2$. A simplified model of a high-beta field

BETATRON - LARGE ORBIT
NON-ADIABATIC PARTICLES

$$B_{\uparrow} = 0$$

TOKAMAK - SMALL ORBIT
ADIABATIC PARTICLES

$$B_{\uparrow} = B_p \left(\frac{R}{r} \right) \left(\frac{2\pi}{\Delta\theta} \right)$$

Figure 1-1: Comparison between small and large orbit dynamics.

reversed configuration with large orbit ions has been investigated theoretically[17]. The absence of a high density adiabatic plasma makes a considerable difference, as an adiabatic background plasma produces a very high dielectric constant perpendicular to the magnetic field, $\epsilon_{\perp} \approx 1 + 4\pi n m_i c^2 / B^2 \gg 1$. If there are few adiabatic ions, then $\epsilon_{\perp} \approx 1$.

The study of field reversed configurations using small orbit MHD theory produces erroneous results. The configuration is predicted to be tilt mode unstable which is not observed in many experiments. This has been explained in terms of large orbit ions[18,19].

In section 2 we find self-consistent steady-state equilibria for plasmas which have drifted Maxwellian distributions and satisfy the Vlasov equation. We call these Vlasov-Maxwellian (or V-M) solutions. First we show that all V-M plasmas must be rigid rotors. Next we derive the V-M equilibrium equation for a single ion species. Finally we numerically evaluate several solutions in finite geometry. Solutions are found which are field reversed and have high density contrast.

In Section 3 we treat a subset of V-M solutions, low-beta ($\beta = 8\pi n k T / B^2$), which are especially amenable to stability analysis. These are low density cases for which self magnetic fields are insignificant. The model which we consider is infinite in the axial direction with only electrostatic perturbations. First the modes of oscillation $\delta\rho(\vec{x})$ and $\delta\phi(\vec{x})$ are found using the method of integration over orbits. Charge neutrality is used to produce an integral equation which defines the complex frequencies for each of these modes. We analytically but approximately find the unstable modes from this integral equation, including a few specific cases which trace back to earlier results. Finally we numerically evaluate stability in a more exact fashion for a large number of cases.

The model in section 3 was used previously[20] to investigate finite Larmor radius effects. The earlier analysis consisted of an expansion in r_L/r_0 where r_L is the mean

Larmor radius and r_0 is the characteristic radial dimension of the plasma. The expansion was carried out to the first significant order beyond MHD and only for the flute-interchange mode $k_z = 0$. In this dissertation all orders in r_L are retained, finite k_z is considered, and all unstable modes are investigated.

2 Vlasov-Maxwellian Equilibria

In this section we will show the surprising result that all V-M equilibrium distributions are rigid rotors, and then calculate the V-M distribution function for a number of cases.

The initial distribution function $f_j(\vec{x}, \vec{v})$ satisfies the Boltzmann equation

$$\frac{df_j}{dt} \equiv \dot{f}_j + (\vec{v} \cdot \nabla) f_j + \frac{q_j}{m_j} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right) \cdot \nabla_v f_j = (\text{collision terms}). \quad (2.1)$$

If we choose the Maxwellian form for the distribution function,

$$f_j = n_j(\vec{x}) \exp [-m_j(\vec{v} - \vec{u}_j(\vec{x}))^2 / 2T_j(\vec{x})], \quad (2.2)$$

then self-collisions between particles of common $\vec{u}$ and T do not affect the distribution. In particular, for electrons plus ions with a common $\vec{u}$ and T ,

$$\frac{df_j}{dt} = (\text{electron - ion collision terms}). \quad (2.3)$$

The electron-ion collision terms manifest themselves most significantly in the form of resistive instabilities. We will assume for this treatment that the e-i collision rate ($\approx \omega_p / 4n\lambda_D^3$) is small compared to the frequencies for the collective modes. When we neglect the right hand side of equation 2.3 we are left with

$$\frac{df_j}{dt} = 0,$$

which is the Vlasov equation.

2.1 Steady state Vlasov-Maxwellian solutions

Our imposed requirement of Maxwellian distribution functions, with electron-ion collisions neglected, yields the steady state solutions f_j which satisfy

$$(\vec{v} \cdot \nabla) f_j + \frac{q_j}{m_j} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right) \cdot \nabla_v f_j = 0 \quad (2.4)$$

and

$$f_j = n_j(\vec{x}) \exp [-m_j(\vec{v} - \vec{u}_j(\vec{x}))^2 / 2T_j(\vec{x})] \quad (2.5)$$

The only other condition which we impose a priori is $\vec{u}_i(\vec{x}) = \vec{u}(\vec{x})$ and $T_i(\vec{x}) = T(\vec{x})$ for all ion species. Substitution yields (dropping the subscripts)

$$\vec{v} \cdot \nabla (\ln n) + \frac{m(\vec{v} - \vec{u})^2}{2T^2} \vec{v} \cdot \nabla T + \frac{m(\vec{v} - \vec{u})}{T} \cdot \left[(\vec{v} \cdot \nabla) \vec{u} - \frac{q}{m} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right) \right] = 0 \quad (2.6)$$

This is a polynomial equation of order 3 in $\vec{v}$. For the equality to hold for all $\vec{v}$, each coefficient must individually vanish.

First consider the coefficient of $v^2 \vec{v}$:

$$(mv^2/2T^2)(\vec{v} \cdot \nabla) T = 0, \quad (2.7)$$

which implies $T(\vec{x}) = \text{constant}$. Substitution back leaves

$$(\vec{v} \cdot \nabla)(\ln n) + \frac{m}{T}(\vec{v} - \vec{u}) \cdot (\vec{v} \cdot \nabla) \vec{u} - \frac{q}{T} \vec{E} \cdot (\vec{v} - \vec{u}) + \frac{q}{T} \left(\frac{\vec{v}}{c} \times \vec{B} \right) \cdot \vec{u} = 0 \quad (2.8)$$

Next gather terms of order $\vec{v} \vec{v}$:

$$(m/T) \vec{v} \cdot (\nabla \vec{u}) \cdot \vec{v} = 0, \quad (2.9)$$

which implies $\nabla_j u_k + \nabla_k u_j = 0$. This is solved in general by $\vec{u} = \vec{\omega} \times \vec{r} + \vec{u}_0$ where $\vec{\omega}$ and $\vec{u}_0$ are constants. This happens to be the general form for a rigid rotor solution. Thus we have the interesting result that the only solutions which both satisfy Vlasov's equation and have a Maxwellian distribution are rigid rotors.

Without loss of generality, we can pick the reference frame where $\vec{u}_0 = 0$. Then the equation for order $\vec{v}$ is

$$(\vec{v} \cdot \nabla)(\ln n) - \frac{m}{T}(\vec{\omega} \times \vec{r}) \cdot (\vec{\omega} \times \vec{v}) - \frac{q}{T}\vec{E} \cdot \vec{v} + \frac{q}{T}(\frac{\vec{v}}{c} \times \vec{B}) \cdot (\vec{\omega} \times \vec{r}) = 0 \quad (2.10)$$

with solution

$$n = n_0 \exp [m\omega^2 r^2 / 2T - q\phi/T + qA_\theta \omega r / cT] \quad (2.11)$$

where r is the cylindrical radial coordinate, $\vec{\omega} = \omega_z$, $\vec{E} = -\nabla\phi$, and $\vec{B} = \nabla \times \vec{A}$.

The equation of order v^0 and is $\vec{E} \cdot \vec{\omega} \times \vec{r} = 0$ which yields $E_\theta = 0$ since we have picked $\vec{\omega} = \omega_z$.

2.2 Numerical calculation of solutions

In this section we will compute equilibria which satisfy the equations derived in the previous section, and the Maxwell's equation

$$\nabla \times \vec{B} = \frac{4\pi\vec{j}}{c} \quad (2.12)$$

By restricting ourselves to a single ion species, we can tabulate the five equations to be simultaneously solved:

Maxwellian for electrons:

$$n_e = n_{oe} \exp [m_e \omega_e^2 r^2 / 2T_e + e\phi/T_e - eA_\theta \omega_e r / cT_e] \quad (2.13)$$

Maxwellian for ions:

$$n_i = n_{oi} \exp [m_i \omega_i^2 r^2 / 2T_i - Ze\phi/T_i + ZeA_\theta \omega_i r / cT_i] \quad (2.14)$$

Equation 2.12:

$$\left[\frac{\partial}{\partial r} \frac{1}{r} \frac{\partial}{\partial r} r + \frac{\partial^2}{\partial z^2} \right] A_\theta = -4\pi j / c \quad (2.15)$$

Charge neutrality:

$$n_e - Zn_i = 0 \quad (2.16)$$

Definition of current:

$$j = -en_e r \omega_e + Z e n_i r \omega_i \quad (2.17)$$

We are able to use equations 2.13 and 2.14 to eliminate ϕ , and use $n_e = Z n_i$:

$$T_i \ln(n_e / Z n_{oi}) + Z T_e \ln(n_e / n_{oe}) = Z m_e \omega_e^2 r^2 / 2 + m_i \omega_i^2 r^2 / 2 + (Z e A_\theta r / c)(\omega_i - \omega_e), \quad (2.18)$$

or solving for A_θ in terms of n_e ,

$$r A_\theta = \frac{c(T_i + Z T_e)}{Z e(\omega_i - \omega_e)} \ln n_e + K_1 r^2 + K_2 \quad (2.19)$$

where K_1 and K_2 are constants which will later be eliminated. Now substitute this expression into equation 2.15:

$$\begin{aligned} \left[r \frac{\partial}{\partial r} \frac{1}{r} \frac{\partial}{\partial r} + \frac{\partial^2}{\partial z^2} \right] \left[\frac{c(T_i + Z T_e)}{Z e(\omega_i - \omega_e)} \ln n_e + K_1 r^2 + K_2 \right] &= -4\pi j r / c \\ &= -4\pi r^2 e n_e (\omega_i - \omega_e) / c \end{aligned} \quad (2.20)$$

or equivalently,

$$n_e = \left[\frac{-c^2(T_e + T_i/Z)}{4\pi e^2(\omega_i - \omega_e)^2} \right] \left[\frac{1}{r} \frac{\partial}{\partial r} \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial z^2} \right] \ln n_e. \quad (2.21)$$

This is the equation which constrains the density distribution for these Vlasov-Maxwellian rigid rotor plasmas.

The boundary condition we choose is to fix the potential ϕ at the outer wall and endcaps, corresponding to a conductive containment vessel. When we solve for n_e in terms of ϕ , we see that

$$\ln n_e \propto \phi + K_3 r^2 + K_4, \quad (2.22)$$

therefore $\ln n_e$ has the form $K_3 r^2 + K_4$ at the boundary. The values of K_3 and K_4 give a complete description of the boundary condition. Note that this description is valid for arbitrary containment vessel shape as long as cylindrical symmetry is maintained.

For numerical treatment, convert to dimensionless form:

$$e^{-\psi} = \left[\frac{\partial^2}{\partial \rho^2} + \frac{1}{4a^2} \frac{1}{\rho} \frac{\partial^2}{\partial \zeta^2} \right] \psi ; \quad (2.23)$$

the relationship between ψ and electron density is

$$\psi = -\ln n_e + \ln \frac{c^2(T_e + T_i/Z)}{\pi r_{\max}^4 e^2 (\omega_i - \omega_e)^2} , \quad (2.24)$$

and the boundary condition at the outer wall and endcaps assuming $\phi = 0$ has the form of equation 2.22:

$$\psi = -\ln \frac{\pi n_0 r_{\max}^4 e^2 (\omega_i - \omega_e)^2}{c^2(T_e + T_i/Z)} - \frac{1}{2} m \omega_i^2 r_{\max}^2 \frac{\omega_e}{T \omega_e + Z \omega_i T_e} \rho . \quad (2.25)$$

In the last three expressions

n_0 = electron density at the core,

a = aspect ratio,

$r_{\max}$ = radius of container,

ρ = radial coordinate $r^2/r_{\max}^2$,

ζ = axial coordinate $z/z_{\max}$.

The numerical integration of these equations is actually quite difficult because the solutions bifurcate - that is, a single boundary condition can produce multiple solutions, while a different boundary condition might produce no solutions. For this case, the "shallow" solution is straightforward, whereas the "deep" solution is difficult to integrate. Here "shallow" refers to the low density solution, and "deep" refers to the high density solution. Both solutions are generated from the same boundary conditions, but the deep solution is contained primarily by its self-magnetic field and so increases to high density.

The techniques we used for numerical integration follow those reported in [21] and so will not be described here. Although we actually evaluated several cases,

typical solutions obtained are shown in Figures 2-1, 2-2, and 2-3. Plotted in each case is ψ vs. z on the horizontal axis and r^2 on the vertical axis. The bottom edge is the axis of the container, the top edge is the outer wall, and the right edge is the endcap. The aspect ratio of the container is 2. The contours of ψ are equally spaced, varying from minimum to maximum, and so represent logarithmic electron density contours. The contour spacing is maximum minus minimum divided by 10.

Solutions 2-1 and 2-2 are the shallow and deep solutions corresponding to a particular boundary condition. The fact that these two solutions are somewhat similar indicates that the boundary condition is close to the bifurcation point where the solutions converge to each other.

Away from the bifurcation point very deep solutions can be obtained (solution 2-3). These deep solutions are the most interesting solutions, as the wall density is much lower than the core density; in this case we have achieved a significant level of containment. It is also interesting that the profile has become somewhat independent of the wall geometry. These deep solutions may represent an interesting plasma configuration.

A very similar treatment was reported in [22] to study astron equilibria. A later treatment[23] presented solutions and discussion for exponential-rigid-rotor distribution functions, which turn out to be equivalent to V-M solutions, but without the natural choice of boundary conditions used in this treatment.

2.3 Low-beta equilibrium

In the case where self magnetic fields are negligible, the current term in equation 2.20 can be ignored which leaves

$$\left[\frac{1}{r} \frac{\partial}{\partial r} \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial z^2} \right] \ln n_e = 0. \quad (2.26)$$

Figure 2-1: Shallow solution near bifurcation; density at maximum = e^{-8} times density at wall.

Figure 2-2: Deep solution near bifurcation; density at maximum = $e^{2.1}$ times density at wall.

Figure 2-3: Deep solution away from bifurcation; density at maximum = $e^{5.8}$ times density at wall.

For the case of no z -dependence in the system, the solution becomes

$$n_e = \exp(-r^2/2r_0^2) . \quad (2.27)$$

This form will be used in section 3 to analyze stability of Vlasov-Maxwellian low-beta plasmas.

3 Stability of Low-beta Vlasov-Maxwellian Solutions

We treat a subset of V-M solutions, low-beta ($\beta = 8\pi nkT/B^2$), which are especially amenable to stability analysis. These are low density cases for which self magnetic fields are insignificant. A previous treatment[20] analysed stability for these equilibria by treating finite Larmor radius effects as an expansion to first order in orbit radius. Here we derive solutions which are accurate to all orders, and in addition include non-zero k_z .

Low-beta V-M plasmas can be described by a constant magnetic field in the z -direction $B_z \equiv m_j c \Omega_j / q_j$. In this case the only permissible self-consistent solutions have only radial dependence with $E_r \equiv -WBr/c$. The resulting distribution function is

$$f_j = f_{oj} \exp [(m_j r^2 / 2T_j)(\omega_j^2 - W\Omega_j + \omega_j \Omega_j) - m_j (v - \omega_j \times r)^2 / 2T_j] \quad (3.1)$$

which has a gaussian density profile with one sigma radius of

$$\frac{1}{r_0^2} = \frac{m_j}{T_j} (-\omega_j^2 + W\Omega_j - \omega_j \Omega_j) . \quad (3.2)$$

3.1 Integration over orbits

We perform an electrostatic linearized stability analysis using the method of integration over orbits. Start with the linearized Vlasov equation in the low-beta limit ($\delta B \rightarrow 0$):

$$\delta \dot{f}_j + (\vec{v} \cdot \nabla) \delta f_j + \frac{q_j}{m_j} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right) \cdot \nabla_v \delta f_j + \frac{q_j}{m_j} \delta \vec{E} \cdot \nabla_v f_j = 0 \quad (3.3)$$

and make the substitutions

$$\delta \vec{E} \equiv -\nabla \phi ,$$

$$\delta f_j \equiv g_j f_j ,$$

$$f_j = f_{oj} \exp [-(r^2/2r_0^2) - m_j(\vec{v} - \vec{\omega}_j \times \vec{r})^2/2T_j] ,$$

which gives the result

$$\frac{d}{dt} g_j \equiv \dot{g}_j + (\vec{v} \cdot \nabla) g_j + \frac{q_j}{m_j} \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right) \cdot \nabla g_j = -\frac{q_j}{T_j} (\vec{v} - \vec{\omega}_j \times \vec{r}) \cdot \nabla \phi \quad (3.4)$$

or equivalently,

$$\frac{d}{dt} \left[g_j + \frac{q_j}{T_j} \phi \right] = \frac{q_j}{T_j} (\omega_j \times r) \cdot \nabla \phi + \frac{q_j}{T_j} \dot{\phi} . \quad (3.5)$$

We consider perturbations of the form $\phi(r) e^{i\omega t + im\theta + ikz}$ and perform the time integration, which is actually an integration over orbits:

$$g_j = -\frac{q_j}{T_j} \phi + \frac{q_j}{T_j} i(\omega + m\omega_j) \int_{-\infty}^{t_0} \phi[t] dt . \quad (3.6)$$

Here the time integration is actually an integration over the unperturbed orbit history of each particle. We denote ϕ at this position by $\phi[t]$. This result yields the perturbation in the distribution function g_j in terms of the perturbed potential ϕ . In order to calculate the integral, however, we must first find the unperturbed orbits.

3.1.1 Orbit Equations

The particle orbit differential equations are $\ddot{x} = \Omega \dot{y} - \Omega W x$, $\ddot{y} = -\Omega \dot{x} - \Omega W y$, and $\ddot{z} = 0$ which have solutions

$$\begin{aligned} \zeta[t] \equiv r[t] e^{i\theta[t]} &= a e^{i\omega_a t} + b e^{i\omega_b t} \\ z[t] &= v_z t + z[0] \end{aligned} \quad (3.7)$$

where

$$\begin{aligned}\omega_a &= -\frac{1}{2} \left(\Omega_j + \sqrt{\Omega_j^2 + 4\Omega_j W} \right) \\ \omega_b &= -\frac{1}{2} \left(\Omega_j - \sqrt{\Omega_j^2 + 4\Omega_j W} \right) .\end{aligned}$$

The higher frequency $\omega_a \sim -\Omega$ corresponds to the gyrofrequency, while the lower frequency $\omega_b \sim W$ corresponds to the $E \times B$ drift frequency. In particular, the constants a and b can be defined in terms of the values at $t = 0$:

$$\begin{aligned}\zeta[t] &= \zeta[0] \left[\frac{\omega_b}{\omega_b - \omega_a} e^{i\omega_a t} + \frac{\omega_a}{\omega_a - \omega_b} e^{i\omega_b t} \right] \\ &+ v_\perp[0] \left[\frac{i}{\omega_b - \omega_a} e^{i\omega_a t} + \frac{i}{\omega_a - \omega_b} e^{i\omega_b t} \right] .\end{aligned}\tag{3.8}$$

At this point it is convenient to define

$$\alpha_j = \frac{\omega_{bj} - \omega_j}{\omega_{bj} - \omega_{aj}} , \quad \Delta\omega_j = \omega_{aj} - \omega_{bj}$$

so that

$$\begin{aligned}\zeta[t] &= \zeta[0] (\alpha e^{i\omega_a t} + (1 - \alpha) e^{i\omega_b t}) \\ &+ (v_\perp[0] - i\omega_j \zeta[0]) (-i e^{i\omega_a t} + i e^{i\omega_b t}) / \Delta\omega .\end{aligned}\tag{3.9}$$

The relevance of α is depicted in Figure 3-1. The orbits described by these formulas are similar to the orbit of the earth's moon around the sun, with a constant v_z added. Small α implies adiabatic-like orbits, α near 1 implies low temperature betatron-like orbits, and intermediate α implies high temperature orbits which are not localized to a particular radius. Also of interest is the term $(v_\perp - i\omega_j \zeta)$ which is the velocity in the drifted reference frame. If α is near 0, this term is small and $\zeta[t] \approx \zeta[0] e^{i\omega_b t}$. If α is near 1, this velocity is again small and $\zeta[t] \approx \zeta[0] e^{i\omega_a t}$. For intermediate values of α all of the terms become significant.

Also note the important relation

$$-\omega_j^2 + W\Omega_j - \omega_j\Omega_j = -(\omega_a - \omega_j)(\omega_b - \omega_j) = \alpha(1 - \alpha)\Delta\omega_j^2 ,\tag{3.10}$$

Figure 3-1: Typical orbits vs. the parameter α .

when combined with equation 3.2 gives

$$T_j = m_j r_0^2 \alpha_j (1 - \alpha_j) \Delta \omega_j^2 . \quad (3.11)$$

We now have the results necessary to integrate equation 3.6, to find g_j in terms of ϕ . We find in particular eigensolutions which satisfy the additional restriction on g_j of plasma charge neutrality.

3.1.2 Closure by forming charge density

Charge neutrality is the condition needed to generate valid solutions. To form the charge density, we integrate the distribution function over (v_r, v_θ, v_z) at the space-time point $(t = 0, r[0], \theta[0], z[0])$:

$$\rho_j = q_j \int d^2 v_\perp dv_z f_j g_j . \quad (3.12)$$

Without loss of generality, we can let $\theta[0] = 0$ and $z[0] = 0$ so that $r \equiv r[0] = \zeta[0]$.

Substitute for f_j and g_j :

$$\rho_j = \frac{q_j^2}{T_j} \left\{ -n_j(r) \phi(r) + i(\omega + m\omega_j) \int_{-\infty}^{t_0} dt \int d^2 v_\perp dv_z f_{0j} e^{-\left(\frac{r^2}{2r_0^2}\right) - \frac{m_j}{2T_j} (\vec{v} - \vec{\omega}_j \times \vec{r})^2} \phi[t] \right\} \quad (3.13)$$

where we define

$$\phi[t] \equiv \phi(r[t]) e^{im\theta[t] + ik_z z[t] + i\omega t} .$$

The potential at the particle $\phi[t]$ is reexpressed in terms of ζ and t :

$$\phi[t] = \phi(|\zeta[t]|) \frac{\zeta^m[t]}{|\zeta[t]|^m} e^{ik_z v_z t + i\omega t} .$$

Make the change of variable

$$v_\perp \rightarrow v_\perp + \vec{\omega}_j \times \vec{r} = v_\perp + i\omega_j \zeta ,$$

and define $\zeta[t] \equiv \zeta[0]S_j + v_\perp R_j$, where

$$S_j \equiv [\alpha e^{i\omega_a t} + (1 - \alpha) e^{i\omega_b t}]$$

$$R_j \equiv [-i e^{i\omega_a t} + i e^{i\omega_b t}] / \Delta \omega .$$

This allows us to reexpress ρ_j as

$$\rho_j = \frac{n_j q_j^2}{T_j} \left\{ -\phi(r) + i(\omega + m\omega_j) \int_{-\infty}^0 dt \int d^2 v_{\perp} dv_z A_j^3 e^{-\frac{m_j}{2T_j}(|v_{\perp}|^2 + v_z^2) + i k_z v_z t + i \omega t} \frac{\zeta^m}{|\zeta|^m} \phi(|\zeta|) \right\} \quad (3.14)$$

where A_j^3 is a normalization factor defined by

$$\int d^2 v_{\perp} dv_z A_j^3 \exp \left[-\frac{m_j}{2T_j}(|v_{\perp}|^2 + v_z^2) \right] = 1.$$

For closure, we must use $\nabla^2 \phi = -4\pi \Sigma \rho_j$ which defines solutions for ϕ and ρ when combined with equation 3.14. However, notice

$$\nabla^2 \phi = \mathcal{O} \left(\frac{1}{r_0^2} \phi \right)$$

and

$$4\pi \rho_j = \mathcal{O} \left(\frac{n_j q_j^2}{T_j} \phi \right) = \frac{4\pi n_j m_j c^2}{\alpha_j (1 - \alpha_j) B^2} \mathcal{O} \left(\frac{1}{r_0^2} \phi \right)$$

where we have used equation 3.11 to substitute for T_j . We take $4\pi n_j m_j c^2 / B^2 \gg 1$ for reasonably interesting densities so that $|4\pi \rho_j| \gg |\nabla^2 \phi|$, and the dominant condition becomes $\rho_e \approx -\rho_i$. We can therefore with sufficient accuracy simply impose charge neutrality $\Sigma \rho_j = 0$ and ignore $\nabla^2 \phi$.

3.1.3 Eigenmode solutions

Here we show that the eigenmodes $\phi(r) = r^m M(-\nu, m+1, r^2/2r_0^2)$ as defined in [24] form the solutions to $\Sigma \rho_j = 0$ for ρ_j given by equation 3.14. The expansion for Kummer's function is

$$M = \sum_{n=0}^{\infty} \frac{(-\nu)_n}{(m+1)_n} \frac{1}{n!} (r^2/2r_0^2)^n \quad (3.15)$$

where

$$(x)_n \equiv x(x+1) \cdots (x+n-1).$$

Substituting this expansion and performing the v_z integral in equation 3.14 yields

$$\rho_j = \frac{n_j q_j^2}{T_j} \sum_{n=0}^{\infty} \frac{(-\nu)_n}{(m+1)_n n! (2r_0^2)^n} \cdot \left\{ -r^{m+2n} + i(\omega + m\omega_j) \int_{-\infty}^0 dt e^{-\frac{T_j k_z^2 t^2}{2m_j} + i\omega t} \int d^2 v_{\perp} A_j^2 e^{-\frac{m_j |v_{\perp}|^2}{2T_j}} \zeta^m[t] |\zeta[t]|^{2n} \right\} \quad (3.16)$$

The $v_{\perp}$ integral can now be performed:

$$\begin{aligned} & \int d^2 v_{\perp} A_j^2 \exp\left(-\frac{m_j}{2T_j} |v_{\perp}|^2\right) \zeta^m[t] |\zeta[t]|^{2n} \\ &= \int d^2 v_{\perp} A_j^2 \exp\left(-\frac{m_j}{2T_j} |v_{\perp}|^2\right) (rS_j + v_{\perp} R_j)^{m+n} (rS_j^* + v_{\perp}^* R_j^*)^n \\ &= \sum_{p=0}^n (rS_j)^m |rS_j|^{2(n-p)} \left(\frac{2T_j}{m_j}\right)^p |R_j|^{2p} \frac{n!(m+n)!}{(m+n-p)!p!(n-p)!} \end{aligned} \quad (3.17)$$

Substituting this back into equation 3.16 and using the sum rule

$$\sum_{n=0}^{\infty} \sum_{p=0}^n g_{n,p} = \sum_{n=0}^{\infty} \sum_{p=0}^{\infty} g_{n+p,p}$$

yields

$$\rho_j = \frac{n_j q_j^2}{T_j} \sum_{n=0}^{\infty} \frac{r^{m+2n}}{(2r_0^2)^n} \left\{ \frac{-(-\nu)_n}{(m+1)_n n!} + i(\omega + m\omega_j) \int_{-\infty}^0 dt e^{-\frac{T_j}{2m_j} k_z^2 t^2 + i\omega t} \Sigma \right\} \quad (3.18)$$

where we have defined

$$\Sigma \equiv \sum_{p=0}^{\infty} S_j^m |S_j|^{2n} |R_j|^{2p} \left(\frac{2T_j}{m_j}\right)^p \frac{(-\nu)_{n+p} m!}{(m+n)! p! n! (2r_0^2)^p}.$$

Now use the identity

$$|R_j|^2 = (1 - |S_j|^2) \left(\frac{m_j r_0^2}{T_j}\right) \quad (3.19)$$

and insert into Σ from equation 3.18 so that we can reexpress this sum as

$$\begin{aligned} \Sigma &= S_j^m |S_j|^{2n} \frac{m!}{(m+n)! n!} \frac{\Gamma(-\nu+n)}{\Gamma(-\nu)} \sum_{p=0}^{\infty} (1 - |S_j|^2)^p \frac{\Gamma(-\nu+n+p)}{\Gamma(-\nu+n)p!} \\ &= S_j^m |S_j|^{2n} \frac{(-\nu)_n}{(m+1)_n n!} \end{aligned} \quad (3.20)$$

Substitution back into equation 3.18 yields finally

$$\rho_j = r^m \left[\sum_{n=0}^{\infty} \frac{r^{2n}}{(2r_0^2)^n} \frac{(-\nu)_n}{(m+1)_n n!} \right] \epsilon_j = \phi(r) \epsilon_j \quad (3.21)$$

where the constants ϵ_j will be used in the following sections to determine stability, and are given by

$$\epsilon_j \equiv \frac{n_j q_j^2}{T_j} \left\{ -1 + i(\omega + m\omega_j) \int_{-\infty}^0 dt e^{-\frac{T_j}{2m_j} k_z^2 t^2 + i\omega t} S_j^m |S_j|^{2\nu} \right\}. \quad (3.22)$$

Charge neutrality $\sum \rho_j = 0$ becomes $\phi(r) \sum \epsilon_j = 0$, and so we finally have the eigenvalue equation

$$0 = \sum_j \frac{n_j q_j^2}{T_j} \left\{ -1 + i(\omega + m\omega_j) \int_{-\infty}^0 dt e^{-\frac{T_j}{2m_j} k_z^2 t^2 + i\omega t} S_j^m |S_j|^{2\nu} \right\} \quad (3.23)$$

with $S_j = \alpha e^{i\omega_{aj}t} + (1 - \alpha)e^{i\omega_{bj}t}$. Valid solutions are those for which ϕ satisfies boundary conditions; we consider a conductive wall for which $\phi(r_w) = 0$ so that the eigenmode equation is

$$M(-\nu, m+1, r_w^2/2r_0^2) = 0. \quad (3.24)$$

The radial modes are those values of ν which satisfy this equation. There are an infinite number of radial modes for any value of m and r_w , and all have $\nu > 0$.

3.1.4 Radial modes

We evaluate the values of ν which result from application of the radial mode condition $M = 0$, by using the identity

$$M(-\nu, m+1, z) = \frac{m!}{\Gamma(m+\nu+1)} e^z z^{-m/2} \int_0^\infty e^{-t} t^{\nu+m/2} J_m(\sqrt{2zt}) dt.$$

Plotted in Figure 3-2 are the radial modes for the case $r_w = \infty$, in which case ν takes on non-negative integer values. When the wall is brought in, the values of ν become more widely spaced, particularly for large m values. The case of $r_w = 4r_0$ is shown in Figure 3-3.

3.2 Evaluation of Stability from Eigenvalue Equation

In order to determine stability we need to analyze the solutions to equation 3.23. We call the terms in the sum ϵ_j "dielectric constants," as they are in a sense "modal

Figure 3-2: Radial eigenmode values ν vs. m , $r_w = \infty$.

Figure 3-3: Radial eigenmode values ν vs. m , $r_w = 4r_0$.

dielectric constants" although not true dielectric constants. An unstable solution is one for which there exist a growing ($\mathcal{I}(\omega) < 0$) root to the charge neutrality condition,

$$\epsilon(\omega) = \sum_j \epsilon_j(\omega) = 0. \quad (3.25)$$

Typically ϵ_i consists of a series of peaks separated by $\sim \Omega_i$, while ϵ_e consists of a much broader central peak with two other irrelevant peaks at $\pm \Omega_e$. A typical case is shown in Figure 3-4 which is an illustration of the general behavior of the real part of ϵ . Clearly, unstable regions tend to occur around the peaks of ϵ_i , with stability determined by the relationship between ϵ_e and the envelope of ϵ_i .

To help restrict the search for the zeros of ϵ , we turn to Nyquist's theorem:

$$\oint dz \frac{f'(z)}{f(z)} = 2\pi i \cdot (\# \text{zeros} - \# \text{poles}) \text{ of } f \text{ in contour.} \quad (3.26)$$

Since the absolute value of the integral is clearly bound from above by

$$\int_{-\infty}^0 dt e^{-\frac{T_i}{2m_j} k_z^2 t^2} < \infty, \quad (3.27)$$

there can be no poles. Therefore

$$\oint dz \frac{\epsilon'(z)}{\epsilon(z)} = 0 \quad (3.28)$$

for stability, where the contour is taken as shown in Figure 3-5. Likewise, an unstable solution has

$$\oint dz \frac{\epsilon'(z)}{\epsilon(z)} = 2\pi i \cdot (\text{positive integer}). \quad (3.29)$$

From this point on we will restrict ourselves to an analysis of only two species, electrons and ions. Nyquist's integral then becomes

$$\oint dz \frac{\epsilon'_e(z) + \epsilon'_i(z)}{\epsilon_e(z) + \epsilon_i(z)} = \oint d[\ln(\epsilon_e + \epsilon_i)] = \oint dz \frac{\epsilon'_e}{\epsilon_e} + \oint d \left[\ln \left(1 + \frac{\epsilon_i}{\epsilon_e} \right) \right]. \quad (3.30)$$

We write the integral in this form because ϵ_e is usually of larger magnitude than ϵ_i . The electrons are in thermal equilibrium with themselves, and so cannot become

Figure 3-4: Characteristics of $\epsilon(\omega)$. Illustrated is a typical case for the real part.

Figure 3-5: Contour used for Nyquist integral.

unstable by themselves, and so

$$\oint dz \frac{\epsilon'_e}{\epsilon_e} = 0 \quad (3.31)$$

(for that matter, $\oint dz (\epsilon'_i/\epsilon_i) = 0$ also). We are left with evaluating

$$\oint d \left[\ln \left(1 + \frac{\epsilon_i}{\epsilon_e} \right) \right] \quad (3.32)$$

which is much easier to work with than the original Nyquist's integral. We can determine its behavior by counting the number of times the argument $1 + \epsilon_i/\epsilon_e$ crosses the branch cut of the logarithm along the negative real axis. The difference in number of clockwise vs. counterclockwise crossings is then the number of roots. Upon evaluation, we find that for threshold cases the number of crossings is typically small so that this becomes a very effective means to search for instability.

3.2.1 Evaluation of ϵ

We evaluate stability by searching the contour previously described for occurrences of $\epsilon_i/\epsilon_e = (\text{Real}\#) \leq -1$. To evaluate the integrals $\epsilon_j(\omega)$, transform equation 3.23 slightly:

$$\epsilon_j = \frac{n_j q_j^2}{T_j} \left\{ -1 + i(\omega + m\omega_j) \int_{-\infty}^0 dt e^{-\frac{T_j}{2m_j} k_z^2 t^2} e^{i(\omega + m\omega_j)t} Z_j^m |Z_j|^{2\nu} \right\}, \quad (3.33)$$

where $Z_j = \alpha_j e^{i(1-\alpha_j)\Delta\omega_j t} + (1 - \alpha_j) e^{-i\alpha_j \Delta\omega_j t}$.

To explore this integral, we must expand $Z^m |Z|^{2\nu}$ as a power series. Presented here is the derivation for $\frac{\alpha}{1-\alpha} < 1$; the corresponding solution for $\frac{\alpha}{1-\alpha} > 1$ is also given at the end of the derivation. We write Z in a form which can be expanded:

$$Z = (1 - \alpha) e^{-i\alpha \Delta\omega t} \left(1 + \frac{\alpha}{1 - \alpha} e^{i\Delta\omega t} \right). \quad (3.34)$$

Now we can expand the powers of Z :

$$Z^{m+\nu} = (1 - \alpha)^{m+\nu} e^{-i\alpha(m+\nu)\Delta\omega t} \sum_{p=0}^{\infty} \left(\frac{\alpha}{1 - \alpha} e^{i\Delta\omega t} \right)^p \binom{m+\nu}{p} \quad (3.35)$$

and

$$Z^{*\nu} = (1 - \alpha)^\nu e^{i\alpha\nu\Delta\omega t} \sum_{q=0}^{\infty} \left(\frac{\alpha}{1 - \alpha} e^{-i\Delta\omega t} \right)^q \binom{\nu}{q} \quad (3.36)$$

where

$$\binom{x}{p} \equiv \frac{\Gamma(x+1)}{p! \Gamma(x+1-p)}.$$

This expansion is justified for $|\frac{\alpha}{1-\alpha}| < 1$ due to analyticity of x^ν for $\mathcal{R}(x) > 0$ if we pick the branch cut along the negative real axis, since $\mathcal{R}(1 - \frac{\alpha}{1-\alpha} e^{i\Delta\omega t})$ is always greater than zero for $|\frac{\alpha}{1-\alpha}| < 1$. We perform the time integration by making use of the identity

$$\int_{-\infty}^0 dt e^{-\frac{T_j}{2m_j} k_z^2 t^2} e^{i(\omega + m\omega_j)t} e^{ist} = \frac{\sqrt{\pi}}{\sigma} w[-(\omega + m\omega_j + s)/\sigma] \quad (3.37)$$

where, as in [25]

$$\begin{aligned} w(z) &= e^{-z^2} \operatorname{erfc}(-iz) \\ \sigma &= k_z v^{th} \equiv k_z \sqrt{2T_j/m_j}. \end{aligned}$$

We obtain, for $\alpha < \frac{1}{2}$,

$$\epsilon = \frac{nq^2}{T} \left\{ -1 + i(\omega + m\omega_j)(1 - \alpha)^{m+2\nu} \frac{\sqrt{\pi}}{\sigma} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{\alpha}{1 - \alpha} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} w(z_{<}) \right\} \quad (3.38)$$

where $z_{<} \equiv -(\omega + m\omega_j + (p - q)\Delta\omega - \alpha m\Delta\omega)/\sigma$.

The corresponding expression for $\alpha > \frac{1}{2}$ is

$$\epsilon = \frac{nq^2}{T} \left\{ -1 + i(\omega + m\omega_j)\alpha^{m+2\nu} \frac{\sqrt{\pi}}{\sigma} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{1 - \alpha}{\alpha} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} w(z_{>}) \right\} \quad (3.39)$$

where $z_{>} \equiv -(\omega + m\omega_j - (p - q)\Delta\omega + (1 - \alpha)m\Delta\omega)/\sigma$.

3.2.2 Evaluation for $|\omega| \rightarrow \infty$

We examine ϵ for $|\omega| \rightarrow \infty$ to verify that this part of the contour does not contribute to the Nyquist integral. Since $\mathcal{I}(\omega) \leq 0$, we can use the following asymptotic

expansion for $w(z)$:

$$w(z) \rightarrow \frac{i}{\sqrt{\pi}z} \left[1 + \sum_{m=1}^{\infty} \frac{1 \cdot 3 \cdots (2m-1)}{(2z^2)^m} \right]. \quad (3.40)$$

This is valid for $z \rightarrow \infty$, and $-\pi/4 < \arg z < 5\pi/4$. This expansion substituted into the expressions for ϵ gives, to lowest order in $1/\omega$,

$$\epsilon \rightarrow \frac{nq^2}{T} \frac{\Delta\omega^2}{\omega^2} \alpha(1-\alpha)(m+2\nu+k_x^2 r_0^2) \quad (3.41)$$

valid for any α within $0 < \alpha < 1$. We are now able to evaluate ϵ_i/ϵ_e on this part of the contour:

$$\frac{\epsilon_i(\omega)}{\epsilon_e(\omega)} = \frac{n_i q_i^2 T_e \Delta\omega_i^2 \alpha_i(1-\alpha_i)}{n_e q_e^2 T_i \Delta\omega_e^2 \alpha_e(1-\alpha_e)} = -\frac{q_i m_e}{q_e m_i} > 0. \quad (3.42)$$

Clearly there are no branch crossings here so we can from this point limit our discussions to the real axis of ω .

3.2.3 Electron dielectric constant

Use of the identities,

$$\alpha_e(1-\alpha_e) = \frac{T_e}{m_e r_0^2 \Delta\omega_e^2} = \frac{T_e m_e c^2}{r_0^2 q_e^2 B^2} = \left[\frac{q_i^2 T_e m_e}{q_e^2 T_i m_i} \right] \alpha_i(1-\alpha_i) \ll 1, \quad (3.43)$$

shows that $\alpha_e \ll 1$ since the term in brackets is very small. We therefore can derive an expansion for small α_e :

$$\epsilon_e \sim \frac{n_e q_e^2}{T_e} \left\{ -1 + i(\omega + m\omega_e)(1 - (m+2\nu)\alpha_e) \frac{\sqrt{\pi}}{\sigma_e} [w(z_e) + \alpha_e(m+\nu)w(z_+) + \alpha_e \nu w(z_-)] \right\} \quad (3.44)$$

where

$$\begin{aligned} z_e &\equiv -(\omega + m\omega_e - \alpha_e m \Delta\omega_e)/\sigma_e, \\ z_+ &\equiv -(\omega + m\omega_e + \Delta\omega_e - \alpha_e m \Delta\omega_e)/\sigma_e, \\ z_- &\equiv -(\omega + m\omega_e - \Delta\omega_e - \alpha_e m \Delta\omega_e)/\sigma_e. \end{aligned}$$

There are three peaks at $m\omega_e - \alpha_e m \Delta\omega_e$ and $m\omega_e - \alpha_e m \Delta\omega_e \pm \Delta\omega_e$. Away from these peaks we can use the asymptotic expansion for $w(z)$ which, upon evaluation, yields

$$\epsilon_e(\text{off peaks}) \sim \frac{n_e q_e^2}{T_e} \alpha_e \Delta\omega_e^2 \frac{(m + 2\nu + k_z^2 r_0^2)(\omega - \omega_+)(\omega - \omega_-)}{(\omega + m\omega_e - \alpha_e m \Delta\omega_e)^2 [(\omega + m\omega_e - \alpha_e m \Delta\omega_e)^2 - \Delta\omega_e^2]} \quad (3.45)$$

where

$$\omega_{\pm} = -m\omega_e + \alpha_e m \Delta\omega_e + \frac{1}{2} \Delta\omega_e \frac{m \pm \sqrt{m^2 + 4k_z^2 r_0^2 (k_z^2 r_0^2 + m + 2\nu)}}{k_z^2 r_0^2 + m + 2\nu}$$

are electron waves analogous to the plasma waves described in [26], and would be the only modes if the ion mass were infinite. The frequencies $\omega_{\pm}$ are plotted vs. m and ν in Figure 3-6. Note that these zeros are always located in between the pairs of peaks. These zeros are important as they are a possible source of instability.

3.2.4 Plasma behavior near electron plasma waves

We now investigate the real axis of ω when $|\omega|$ is large compared to $\Delta\omega_i$. In this regime ϵ_e is always much larger than ϵ_i except at the electron plasma waves $\omega_{\pm}$, so that we can limit our discussion to these two values for ω . The question of stability is determined by the imaginary part of ϵ_i , as the effect of the real part of ϵ_i is just a perturbation of the location of the real frequencies $\omega_{\pm}$. Using the fact that $\mathcal{R}(w(z)) = e^{-z^2}$, we can write for $\alpha < 1/2$,

$$\mathcal{I}(\epsilon_i) = \frac{n_i q_i^2}{T_i} (\omega + m\omega_i) (1 - \alpha_i)^{m+2\nu} \frac{\sqrt{\pi}}{\sigma_i} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \left(\frac{\alpha_i}{1 - \alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} e^{-z^2} \quad (3.46)$$

where we define $z \equiv -(\omega + m\omega_i + (p - q)\Delta\omega_i - \alpha_i m \Delta\omega_i) / \sigma_i$.

This has a significant contribution around the plasma waves $\omega_{\pm}$ only for $\omega_{\pm} + m\omega_i + (p - q)\Delta\omega_i - \alpha_i m \Delta\omega_i \sim 0$ (due to the gaussian term), and since $\omega_{\pm}$ is large compared to $\Delta\omega_i$ then either p or q must be large. The smallest value for $p + q$ which is significant is then $p + q \sim |\omega_{\pm} / \Delta\omega_i|$. Therefore, due to the term $\left(\frac{\alpha_i}{1 - \alpha_i} \right)^{p+q}$,

Figure 3-6: $\omega_{\pm}$ versus m and ν . $\omega_{\pm}$ is reported in units of $\Delta\omega_e$.

$\mathcal{I}(\epsilon_i)$ is exponentially negligible unless α_i is very close to $1/2$. The corresponding expression for $\alpha > 1/2$ is $\left(\frac{1-\alpha_i}{\alpha_i}\right)^{p+q}$ which gives the same result.

Note that when the terms $\binom{m+\nu}{p}$ and $\binom{\nu}{q}$ are non-zero for large p and q , then ν cannot be an integer. Non-integer ν implies that the finite walls are affecting the solution. We therefore can conclude that the plasma is stable for high frequencies, $\omega \sim \mathcal{O}(\Delta\omega_e)$, except for possibly an unusual mode with finite walls where α_i is very close to $1/2$ (to within a fraction of a percent).

3.3 Analytical Approximate Methods for Moderate Frequencies

In this section we consider modes which are at moderate frequencies, $\omega \sim \mathcal{O}(\Delta\omega_i)$, which is the regime where ϵ_e and ϵ_i are comparable, and therefore where most of the unstable modes are found. We find these unstable modes by searching for occurrences of $\epsilon_i = \beta\epsilon_e$ where β is real and ≤ -1 . Here we will make approximations which are motivated by the numerical calculations which are presented in the next section. Although the numerical results provide the definitive conclusions, an understanding of the physics is better provided in this section.

The fundamental assumption in this section is that the ion dielectric constant must divide into well separated peaks for instability (in other words, $\Delta\omega_i > \sigma_i$). This is well supported by the numerical calculations. We are therefore able to treat each peak separately and independently from the others. We first express the dielectric constants in a slightly different form by using the following identities:

$$w'(z) = -2zw(z) + \frac{2i}{\sqrt{\pi}}, \quad (3.47)$$

$$1 = (1-\alpha)^{m+2\nu} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{\alpha}{1-\alpha}\right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q}, \quad (3.48)$$

so that we can write the dielectric constants as

$$\epsilon_{\alpha < 1/2} = \frac{nq^2}{T} \sum_{\substack{p=0 \\ q=0}}^{\infty} (1-\alpha)^{m+2\nu} \left(\frac{\alpha}{1-\alpha} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} \left[\frac{i\sqrt{\pi}}{2} w'(z_{<}) + (\alpha m - p + q) \frac{\Delta\omega}{\sigma} i\sqrt{\pi} w(z_{<}) \right], \quad (3.49)$$

$$\epsilon_{\alpha > 1/2} = \frac{nq^2}{T} \sum_{\substack{p=0 \\ q=0}}^{\infty} \alpha^{m+2\nu} \left(\frac{1-\alpha}{\alpha} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} \left[\frac{i\sqrt{\pi}}{2} w'(z_{>}) + (\alpha m - m + p - q) \frac{\Delta\omega}{\sigma} i\sqrt{\pi} w(z_{>}) \right], \quad (3.50)$$

with

$$z_{<} = -(\omega + m\omega_j + (p-q)\Delta\omega - \alpha m\Delta\omega)/\sigma,$$

$$z_{>} = -(\omega + m\omega_j - (p-q)\Delta\omega + (1-\alpha)m\Delta\omega)/\sigma.$$

This form is useful because now each term of the sum is clearly a localized quantity which drops off as $\mathcal{O}(1/\omega)$ for ω large. Since we are assuming in this section that ϵ_i has well separated peaks, we can consider each peak in isolation from the other peaks. These peaks are centered around

$$\omega_s = -m\omega_i - s\Delta\omega_i + \alpha_i m\Delta\omega_i \quad (3.51)$$

where s is the harmonic number (refer back to Figure 3-4 for an illustration of these peaks). We transform to these new coordinates, replacing $\omega \equiv \omega_s - \delta\sigma_i$, where δ is a measure of deviation from peak center. With this substitution, we get

$$z_{i<} = \delta, \quad \text{with } s = p - q$$

$$z_{i>} = \delta, \quad \text{with } s = m - p + q$$

and with this definition we get for z_e ,

$$z_e = -(-m\omega_i + \alpha_i m\Delta\omega_i + m\omega_e - \alpha_e m\Delta\omega_e - s\Delta\omega_i - \delta\sigma_i)/\sigma_e. \quad (3.52)$$

The expression for z_e can be simplified by using the identity

$$\omega_j = -\frac{1}{2}(\Omega_j + \Delta\omega_j(1 - 2\alpha_j)) = W + \alpha_j\Delta\omega_j + W^2/\Delta\omega_j + \mathcal{O}(W^3), \quad (3.53)$$

and so

$$z_e \approx mW^2/\Delta\omega_i\sigma_e + s\Delta\omega_i/\sigma_e + \delta\sigma_i/\sigma_e. \quad (3.54)$$

We rewrite our dielectric constants in our simplified approximate form:

$$\epsilon_i \sim \frac{n_i q_i^2}{T_i} \mathcal{F}(\alpha_i, m, \nu, s) \left[\frac{i\sqrt{\pi}}{2} w'(\delta) + (\alpha_i m - s) \frac{\Delta\omega_i}{\sigma_i} i\sqrt{\pi} w(\delta) \right] \quad (3.55)$$

where $\mathcal{F}$ is defined as the envelope of ϵ_i ,

$$\begin{aligned} \mathcal{F}(\alpha_i < 1/2, m, \nu, s) &= \sum_{\substack{p=0 \\ q=0}}^{\infty} (1 - \alpha_i)^{m+2\nu} \left(\frac{\alpha_i}{1 - \alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} \delta_{s, p-q} \\ \mathcal{F}(\alpha_i > 1/2, m, \nu, s) &= \sum_{\substack{p=0 \\ q=0}}^{\infty} \alpha_i^{m+2\nu} \left(\frac{1 - \alpha_i}{\alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} \delta_{m-s, p-q} \end{aligned} \quad (3.56)$$

The electron dielectric constant is dominated by its central peak in the regime we are now examining. We therefore consider only the $p = 0, q = 0$ peak:

$$\epsilon_e \sim \frac{n_e q_e^2}{T_e} \left[\frac{i\sqrt{\pi}}{2} w'(z_e) + \frac{\alpha_e m \Delta\omega_e}{\sigma_e} i\sqrt{\pi} w(z_e) \right] \quad (3.57)$$

with

$$z_e = mW^2/\Delta\omega_i\sigma_e + s\Delta\omega_i/\sigma_e + \delta\sigma_i/\sigma_e.$$

We form the ratio $\beta = \epsilon_i/\epsilon_e$ to search for β real and ≤ -1 :

$$\beta = \frac{\epsilon_i}{\epsilon_e} \sim \frac{\frac{n_i q_i^2}{T_i} \mathcal{F}(\alpha_i, m, \nu, s)}{\frac{n_e q_e^2}{T_e}} \frac{\frac{i\sqrt{\pi}}{2} w'(\delta) + (\alpha_i m - s) \frac{\Delta\omega_i}{\sigma_i} i\sqrt{\pi} w(\delta)}{\frac{i\sqrt{\pi}}{2} w'(z_e) + \frac{\alpha_e m \Delta\omega_e}{\sigma_e} i\sqrt{\pi} w(z_e)}. \quad (3.58)$$

We must first search over δ to find the proper value which makes β real, and then search over s to see if any of these values are ≤ -1 . Before proceeding, note

$$\frac{\frac{n_i q_i^2}{T_i}}{\frac{n_e q_e^2}{T_e}} = \frac{T_e}{T_i/Z}$$

and

$$\frac{\alpha_e m \Delta \omega_e}{\sigma_e} = -\frac{T_e}{T_i/Z} m \alpha_i (1 - \alpha_i) \frac{\Delta \omega_i}{\sigma_e}.$$

Now we define the peak interaction term

$$\mathcal{A}(z_e, \mathcal{B}, \mathcal{C}) \equiv \frac{i \frac{\sqrt{\pi}}{2} w'(\delta) + C i \sqrt{\pi} w(\delta)}{-i \frac{\sqrt{\pi}}{2} w'(z_e) + B i \sqrt{\pi} w(z_e)} \quad (3.59)$$

with

$$\mathcal{B} = \frac{T_e}{T_i/Z} m \alpha_i (1 - \alpha_i) \frac{\Delta \omega_i}{\sigma_e} \quad (3.60)$$

$$\mathcal{C} = (\alpha_i m - s) \frac{\Delta \omega_i}{\sigma_i} \quad (3.61)$$

where we are implicitly defining δ which makes $\mathcal{A}$ real and positive or negative, depending on the sign of $\mathcal{F}$. This leaves

$$\beta = -\frac{T_e}{T_i/Z} \mathcal{F}(\alpha_i, m, \nu, s) \mathcal{A}(z_e, \mathcal{B}, \mathcal{C}). \quad (3.62)$$

For $\mathcal{F} > 0$, we must find values for $\mathcal{B}$ and $\mathcal{C}$ so that $\mathcal{A}$ is positive. We have evaluated the function $\mathcal{A}$ numerically by varying δ from -5 to 5 and picking occurrences of real, positive $\mathcal{A}$. The results are tabulated in Table 3-1.

Note that $\mathcal{B} + x_e$ and $\mathcal{C} - \delta$ must both be of the same sign for a positive solution. Since

$$(\mathcal{B} + x_e) \frac{\sigma_e}{\sigma_i} + (\mathcal{C} - \delta) = \frac{m W^2}{\Delta \omega_i \sigma_i} + \frac{T_e}{T_i/Z} m \alpha_i (1 - \alpha_i) \frac{\Delta \omega_i}{\sigma_i} + \frac{\alpha_i m \Delta \omega_i}{\sigma_i} \geq 0,$$

both terms must individually be positive. Here we have adopted the sign convention $\Delta \omega_i > 0$.

We make some observations regarding $\mathcal{F}$, which was defined in equation 3.56. If ν is an integer, then $\mathcal{F}$ is always positive because all non-zero terms in the sum are positive. Negative terms can occur only when ν is not an integer, and either $q > \nu$ or $p > m + \nu$. These negative terms are small, and so negative $\mathcal{F}$ instability is an unlikely occurrence.

Z_e	B	C										
		-1.0	-0.5	-0.2	0.0	0.2	0.5	1.0	2.0	5.0	10.0	20.0
0.1	0.0	***	***	***	0.2	0.3	0.5	0.9	1.7	4.2	8.3	16.6
0.2	0.0	***	***	0.0	0.2	0.4	0.7	1.2	2.2	5.4	10.8	21.5
0.5	0.0	***	***	0.2	0.5	0.8	1.2	2.0	3.6	8.6	17.1	34.3
1.0	0.0	***	***	0.7	1.1	1.5	2.1	3.2	5.7	13.6	27.0	53.8
1.5	0.0	***	1.1	1.9	2.3	2.7	3.4	4.7	7.8	18.5	36.5	73.2
2.0	0.0	***	3.7	4.3	4.6	4.9	5.4	6.6	10.1	23.3	46.2	92.0
2.5	0.0	7.5	8.3	8.5	8.6	8.8	9.0	9.5	12.6	28.2	55.7	111.9
3.0	0.0	14.3	14.3	14.3	14.4	14.4	14.4	14.4	16.0	33.2	65.6	130.8
0.0	0.1	***	***	***	0.2	0.3	0.5	0.9	1.7	4.1	8.1	16.2
0.2	0.1	***	***	0.1	0.3	0.5	0.8	1.4	2.5	6.2	12.2	24.4
0.5	0.1	***	***	0.2	0.5	0.8	1.2	2.0	3.6	8.6	17.0	33.9
1.0	0.1	***	***	0.7	1.1	1.4	1.9	2.9	5.0	11.9	23.6	47.0
1.5	0.1	***	1.1	1.7	2.0	2.3	2.8	3.8	6.2	14.6	29.1	57.9
2.0	0.1	***	3.1	3.4	3.6	3.8	4.2	4.9	7.4	17.1	33.9	67.5
2.5	0.1	5.7	6.1	6.2	6.2	6.3	6.4	6.8	8.6	19.1	38.0	75.8
3.0	0.1	9.4	1.0	0.1	***	***	***	9.5	10.3	21.1	41.6	83.0
-0.1	0.2	***	***	***	0.2	0.3	0.5	0.9	1.6	4.0	7.9	15.8
0.0	0.2	***	***	0.0	0.2	0.4	0.7	1.1	2.1	4.9	9.9	19.7
0.5	0.2	***	***	0.3	0.5	0.8	1.2	1.9	3.4	8.2	16.2	32.4
1.0	0.2	***	***	0.7	1.0	1.2	1.7	2.5	4.3	10.3	20.5	40.9
1.5	0.2	***	1.0	1.5	1.7	1.9	2.3	3.1	5.1	11.9	23.8	47.3
2.0	0.2	***	2.6	2.8	3.0	3.1	3.4	3.9	5.8	13.3	26.4	52.5
2.5	0.2	4.5	4.8	4.8	4.9	4.9	5.0	5.2	6.5	14.3	28.3	56.5
3.0	0.2	7.1	0.8	0.1	***	***	***	7.1	7.6	15.3	30.2	60.2
-0.4	0.5	***	***	***	0.1	0.3	0.5	0.8	1.5	3.6	7.1	14.2
-0.3	0.5	***	***	0.0	0.2	0.3	0.5	0.9	1.8	4.2	8.4	16.7
0.0	0.5	***	***	0.1	0.3	0.5	0.7	1.2	2.2	5.4	10.7	21.3
0.5	0.5	***	***	0.3	0.5	0.7	1.0	1.5	2.7	6.4	12.8	25.5
1.0	0.5	***	***	0.6	0.8	0.9	1.2	1.7	3.0	7.0	14.0	28.0
1.5	0.5	***	0.8	1.1	1.2	1.3	1.6	2.0	3.2	7.5	14.9	29.7
2.0	0.5	1.2	1.8	1.9	1.9	2.0	2.1	2.4	3.4	7.8	15.5	30.9
2.5	0.5	2.8	2.9	2.9	3.0	3.0	3.0	3.1	3.7	8.1	16.0	31.8
3.0	0.5	1.2	0.4	0.0	***	***	***	***	4.2	8.3	16.4	32.6
-0.9	1.0	***	***	***	0.1	0.2	0.4	0.6	1.2	2.9	5.8	11.6
-0.8	1.0	***	***	***	0.1	0.2	0.4	0.7	1.3	3.2	6.4	12.8
-0.5	1.0	***	***	0.0	0.2	0.3	0.5	0.8	1.5	3.7	7.4	14.8
0.0	1.0	***	***	0.1	0.3	0.4	0.6	0.9	1.7	4.1	8.1	16.2
0.5	1.0	***	***	0.2	0.4	0.5	0.7	1.0	1.8	4.3	8.5	16.9
1.0	1.0	***	0.2	0.4	0.5	0.6	0.8	1.1	1.9	4.4	8.7	17.4
1.5	1.0	***	0.6	0.7	0.8	0.9	1.0	1.2	1.9	4.5	8.9	17.7
2.0	1.0	0.9	1.1	1.2	1.2	1.3	1.3	1.4	2.0	4.6	9.0	17.9
2.5	1.0	1.7	1.8	1.8	1.8	1.8	1.8	1.8	2.1	4.6	9.1	18.2
3.0	1.0	0.7	0.3	0.0	***	***	***	***	2.4	4.7	9.2	18.3

Table 3-1: The peak interaction function $\mathcal{A}$ vs. z_e , B , C . δ is picked so that $\mathcal{A}$ is real and positive. The symbol *** indicates no solution exists for $-5 < \delta < 5$.

We now have the groundwork completed to evaluate β for a variety of cases. We divide into the following groups:

1. adiabatic modes: $\mathcal{F} > 0$, $(m + 2\nu)\alpha_i \ll 1$,
2. low temperature, large orbit modes: $\mathcal{F} > 0$, $(m + 2\nu)(1 - \alpha_i) \ll 1$,
3. regular modes: $\mathcal{F} > 0$, $(m + 2\nu)\alpha_i(1 - \alpha_i) = \mathcal{O}(1)$,
4. wall modes: $\mathcal{F} < 0$

We will analyze these one at a time in the following sections. The wall modes also include the high frequency mode discussed earlier in section 3.2.4.

3.3.1 Adiabatic Modes

Here the assumption is $\alpha_i(m + 2\nu) \ll 1$. For this case, $\mathcal{F} \sim 1$ when $s = 0$ but $\mathcal{F} \sim 0$ otherwise. Since $\mathcal{F}$ must be large for instability, s is forced to zero, and the arguments of $\mathcal{A}$ become:

$$z_e = mW^2/\Delta\omega_i\sigma_e + \delta\sigma_i/\sigma_e \sim mW^2/\Delta\omega_i\sigma_e ,$$

$$\mathcal{B} = \frac{T_e}{T_i/Z} m\alpha_i(1 - \alpha_i) \frac{\Delta\omega_i}{\sigma_e} ,$$

$$\mathcal{C} = \alpha_i m \frac{\Delta\omega_i}{\sigma_i} \leq \mathcal{O}(1) .$$

For this range of parameters we can approximate $\mathcal{A}$ by

$$\mathcal{A} \sim \frac{2z_e^2 - 3}{1 + 2z_e\mathcal{B}}$$

for $z_e \geq 2$, and $\mathcal{A}$ small otherwise. We therefore have for instability,

$$1 \leq \frac{T_e}{T_i/Z} (2z_e^2 - 3)/(1 + 2z_e\mathcal{B})$$

or

$$\left(\frac{T_i/Z}{T_e} + 3 \right) \sigma_e^2 \leq 2m^2W^2(W^2/\Delta\omega_i^2 - \alpha_i) \quad (3.63)$$

with $mW^2/\Delta\omega_i\sigma_e \geq 2$. Since $s = 0$, then $\omega \approx -mW$, which indicates a low frequency mode.

Clearly, for the MHD case, $\sigma_e = 0$ and $\alpha_i = 0$ so the mode is always unstable. Note, however, that even a very small value for α_i corresponding to a small but non-zero temperature stabilizes this mode.

The MHD case is unique in that it suffers from another unstable mode which is considerably removed from the $s = 0$ peaks of ϵ_i and ϵ_e . In order to evaluate this mode we must back up to equation 3.49. Since we are not on the peak, there are actually three significant ϵ_i terms rather than just one, namely $p = q = 0$; $p = 1, q = 0$; and $p = 0, q = 1$:

$$\begin{aligned} \epsilon_{i,\alpha \sim 0} \sim & \frac{n_i q_i^2}{T_i} \left[\frac{i\sqrt{\pi}}{2} w'(\delta) + \alpha_i m \frac{\Delta\omega_i}{\sigma_i} i\sqrt{\pi} w(\delta) \right] \\ & - \frac{n_i q_i^2}{T_i} \alpha_i (m + \nu) \frac{\Delta\omega_i}{\sigma_i} i\sqrt{\pi} w(\delta - \Delta\omega_i/\sigma_i) \\ & + \frac{n_i q_i^2}{T_i} \alpha_i \nu \frac{\Delta\omega_i}{\sigma_i} i\sqrt{\pi} w(\delta + \Delta\omega_i/\sigma_i) \end{aligned} \quad (3.64)$$

where $\omega \approx -mW - \delta\sigma_i$. The expression which we have used for ϵ_e , equation 3.57, is adequate in this case, where

$$z_e = mW^2/\Delta\omega_i\sigma_e + \delta\sigma_i/\sigma_e.$$

Terms of order $\mathcal{O}(\Omega_i/\Omega_e)$ have been dropped. Since we are off peak, we can use the asymptotic expansion for w :

$$w(z) \sim \frac{i}{\sqrt{\pi}z}, \quad w'(z) \sim \frac{-i}{\sqrt{\pi}z^2},$$

therefore our dielectric constant becomes

$$\begin{aligned} \epsilon \sim & \frac{n_i q_i^2}{T_i} \left[\alpha_i k_z^2 r_0^2 \left(\frac{\Delta\omega_i}{\sigma_i \delta} \right)^2 - \alpha_i m \frac{\Delta\omega_i}{\sigma_i \delta} + \alpha_i (m + \nu) \frac{\Delta\omega_i}{\sigma_i \delta - \Delta\omega_i} - \alpha_i \nu \frac{\Delta\omega_i}{\sigma_i \delta + \Delta\omega_i} \right] \\ & + \frac{n_e q_e^2}{T_e} \left[\alpha_e m \frac{-\Delta\omega_e}{\sigma_i \delta + mW^2/\Delta\omega_i} + \alpha_e k_z^2 r_0^2 \left(\frac{\Delta\omega_e}{\sigma_i \delta + mW^2/\Delta\omega_i} \right)^2 \right]. \end{aligned} \quad (3.65)$$

The coefficient of the last term must be accurately expanded:

$$\frac{n_e q_e^2}{T_e} \alpha_e \Delta \omega_e \approx -\frac{n_i q_i^2}{T_i} \alpha_i \Delta \omega_i \left(1 - \frac{2W}{\Delta \omega_i}\right)$$

which is a higher order approximation than used previously. The eigenvalue is the root of $\epsilon = 0$, so we find the mode by solving for

$$0 = \frac{k_z^2 r_0^2}{x^2} - \frac{m}{x} + \frac{(m + \nu)}{x - 1} - \frac{\nu}{x + 1} + \left(1 - \frac{2W}{\Delta \omega_i}\right) \left(\frac{m}{x + mW^2/\Delta \omega_i^2} - \frac{k_z^2 r_0^2 \Delta \omega_e / \Delta \omega_i}{(x + mW^2/\Delta \omega_i^2)^2}\right) \quad (3.66)$$

where $x = \sigma_i \delta / \Delta \omega_i \approx -(mW + \omega) / \Delta \omega_i$. There are five roots to this equation; if we set $k_z r_0 \leq \mathcal{O}(W / \sqrt{|\Delta \omega_e \Delta \omega_i|})$ and assume that $x = \mathcal{O}(W / \Delta \omega_i)$, then we can drop higher order terms in W to yield the quadratic equation

$$0 = (2\nu + m)x^2 + 2mWx/\Delta \omega_i + m^2W^2/\Delta \omega_i^2 + k_z^2 r_0^2 \Delta \omega_e / \Delta \omega_i \quad (3.67)$$

with solutions

$$x = \frac{-mW/\Delta \omega_i}{m + 2\nu} \left[1 \pm \sqrt{1 - (m + 2\nu)(1 - k_z^2 r_0^2 |\Delta \omega_e \Delta \omega_i| / m^2 W^2)}\right] \quad (3.68)$$

For $\nu = 0$ and $k_z = 0$ this is seen as the MHD instability,

$$\omega = -mW + W \left[1 \pm \sqrt{1 - m}\right] \quad (3.69)$$

This mode stabilizes for $k_z r_0 \geq mW / \sqrt{|\Delta \omega_e \Delta \omega_i|}$, and so is of interest only for k_z very small.

The MHD result was extended to finite Larmor radius in [20] by letting W become small, of order α_i , although only $k_z = 0$ was considered. This requires expansion of the dielectric constant through order α_i^2 . The prescribed expansion with $k_z = 0$ yields a voluminous order 5 polynomial equation. We simplify considerably when we assume $x = \mathcal{O}(W / \Delta \omega_i)$ and $W / \Delta \omega_i = \mathcal{O}(\alpha_i)$; the order 5 equation then reduces to the quadratic equation

$$0 = (2\nu + m)x^2 + 2mWx/\Delta \omega_i - \alpha_i m(2\nu + m - 1)x + m^2W^2/\Delta \omega_i^2 \quad (3.70)$$

which is similar to Equation 3.67. The solutions to this quadratic equation are

$$x = m \frac{\frac{\alpha_i(2\nu+m-1)}{2} - \frac{W}{\Delta\omega_i} \pm \sqrt{\left(\frac{\alpha_i(2\nu+m-1)}{2} - \frac{W}{\Delta\omega_i}\right)^2 - \frac{(2\nu+m)W^2}{\Delta\omega_i^2}}}{2\nu+m} \quad (3.71)$$

or equivalently,

$$(2\nu+m) \frac{\omega + mW}{mW} = - \frac{\alpha_i(2\nu+m-1)\Delta\omega_i}{2W} + 1 \pm \sqrt{\left(-\frac{\alpha_i(2\nu+m-1)\Delta\omega_i}{2W} + 1\right)^2 - (2\nu+m)} \quad (3.72)$$

which is exactly the result obtained in [20] when the notation is brought into correspondence. The mode stabilizes for $2\nu+m > 1$ when

$$\frac{\alpha_i\Delta\omega_i}{2W} > \frac{1}{\sqrt{2\nu+m-1}}$$

or

$$-\frac{\alpha_i\Delta\omega_i}{2W} > \frac{1}{\sqrt{2\nu+m+1}}.$$

For the case $\Delta\omega_i/W < 0$, the mode stabilizes when $\alpha_i > |W/\Delta\omega_i|$. For the case $\Delta\omega_i/W > 0$, the wall causes ν to take on a small value $\nu_{\min}^{m=1}$ for the lowest radial mode of $m = 1$, so we must require $\alpha_i > 2W/\Delta\omega_i \nu_{\min}^{m=1}$ for stability. The case $\Delta\omega_i/W > 0$ is therefore much more difficult to stabilize by increasing α_i . When $2\nu+m < 1$, we know from section 3.1.4 that $2\nu+m \approx 0$ so that the mode is stable except for the special case $\alpha_i\Delta\omega_i/2W = -1$, for which non-zero ν (wall effect) allows instability.

3.3.2 Low Temperature, Large Orbit Modes

Here the assumption is $(1 - \alpha_i)(m + 2\nu) \ll 1$. For this case, $\mathcal{F} \sim 1$ when $s = m$ but $\mathcal{F} \sim 0$ otherwise. Since $\mathcal{F}$ must be large for instability, s is forced to m , and the arguments of $\mathcal{A}$ become:

$$z_e \sim \frac{m\Delta\omega_i}{\sigma_e},$$

$$\mathcal{B} = \frac{T_e}{T_i/Z} m \alpha_i (1 - \alpha_i) \frac{\Delta \omega_i}{\sigma_e},$$

$$\mathcal{C} = -(1 - \alpha_i) m \frac{\Delta \omega_i}{\sigma_i} \leq \mathcal{O}(1).$$

For this range of parameters we can again approximate $\mathcal{A}$ by

$$\mathcal{A} \sim \frac{2z_e^2 - 3}{1 + 2z_e \mathcal{B}}$$

for $z_e \geq 2$, and $\mathcal{A}$ small otherwise. We therefore have for instability,

$$1 \leq \frac{T_e}{T_i/Z} (2z_e^2 - 3) / (1 + 2z_e \mathcal{B})$$

or

$$\left(\frac{T_i/Z}{T_e} + 3 \right) \leq 2 \left(\frac{m \Delta \omega_i}{\sigma_e} \right)^2 \quad (3.73)$$

with $m \Delta \omega_i / \sigma_e \geq 2$. Here $\omega \approx m \Omega_i$ so that this is a relatively high frequency mode.

3.3.3 Regular Modes

Here the assumption is $(m + 2\nu) \alpha_i (1 - \alpha_i) \sim \mathcal{O}(1)$. For this case, $\mathcal{F}$ is not so rapidly varying vs. s as in the previous cases, and there are at least two values of s for which $\mathcal{F}$ is reasonably large. Because of this, the argument $\mathcal{C} = (\alpha_i m - s) \Delta \omega_i / \sigma_i$ optimizes at a fairly large value. For large $\mathcal{C}$, we can use a different approximation for $\mathcal{A}$ accurate in this regime:

$$\mathcal{A} \sim \mathcal{C} \frac{.6 + 2(z_e + \mathcal{B})}{1 + 2\mathcal{B}(z_e + \mathcal{B})}. \quad (3.74)$$

Since $\mathcal{C} > 0$, then $s < \alpha_i m \leq m - 1$. Also, since $z_e + \mathcal{B} > 0$, then $s \geq 0$. As a function over s , $\mathcal{F}$ is maximized at $s \sim \alpha_i m$. We find that for $1 < s < m - 1$, s is accurately approximated in the vicinity of its maximum by

$$\mathcal{F} \sim \frac{\exp [-(\alpha_i m - s)^2 / 2(m + 2\nu) \alpha_i (1 - \alpha_i)]}{\sqrt{2\pi(m + 2\nu) \alpha_i (1 - \alpha_i)}}. \quad (3.75)$$

This is true because $\mathcal{F}$ is a product of binomial series, which are known in general to approach gaussian form. For $s = 0, 1, m - 1$ this approximation is no longer

accurate, and we write out explicit terms, assuming α_i small for $s = 0$ or 1 and $(1 - \alpha_i)$ small for $s = m - 1$:

$$\mathcal{F}(s = 0) \sim (1 - \alpha_i)^{m+2\nu} \quad (3.76)$$

$$\mathcal{F}(s = 1) \sim (1 - \alpha_i)^{m+2\nu-1}(m + \nu)\alpha_i \quad (3.77)$$

$$\mathcal{F}(s = m - 1) \sim \alpha_i^{m+2\nu-1}(m + \nu)(1 - \alpha_i) . \quad (3.78)$$

These are special cases to be investigated separately. For the case $1 < s < m - 1$, we must optimize vs. s to find the most unstable mode. The inequality for instability is

$$1 \leq \frac{T_e}{T_i/Z} \mathcal{F} \cdot (\alpha_i m - s) \frac{\Delta\omega_i}{\sigma_i} \frac{.6 + 2(z_e + \mathcal{B})}{1 + 2\mathcal{B}(z_e + \mathcal{B})} . \quad (3.79)$$

We treat the last quotient as slowly varying, and optimize vs. s the term

$$\mathcal{F} \cdot (\alpha_i m - s) = \frac{\exp [(\alpha_i m - s)^2 / 2(m + 2\nu)\alpha_i(1 - \alpha_i)]}{\sqrt{2\pi(m + 2\nu)\alpha_i(1 - \alpha_i)}} (\alpha_i m - s) . \quad (3.80)$$

This expression is maximum for $s = \alpha_i m - \sqrt{(m + 2\nu)\alpha_i(1 - \alpha_i)}$ and its value at maximum is .24, independent of all parameters.

We will summarize our various inequalities vs. s , but first define a new function

$$\mathcal{K}(z_e) \equiv \frac{T_e}{T_i/Z} \frac{\Delta\omega_i}{\sigma_i} \frac{.6 + 2(z_e + \mathcal{B})}{1 + 2\mathcal{B}(z_e + \mathcal{B})}$$

which simplifies the expressions:

$$s = 0, \omega \approx -mW$$

$$1 \leq \mathcal{K} \left[\frac{mW^2}{\Delta\omega_i \sigma_e} \right] (1 - \alpha_i)^{m+2\nu} \alpha_i m , \quad (3.81)$$

$$s = 1, \omega \approx \Omega_i$$

$$1 \leq \mathcal{K} \left[\frac{\Delta\omega_i}{\sigma_e} \right] (1 - \alpha_i)^{m+2\nu-1} (m + \nu) \alpha_i (\alpha_i m - 1) , \quad (3.82)$$

$$1 < s < m - 1, \omega \approx \alpha_i m \Omega_i$$

$$1 \leq .24\mathcal{K} \left[\left(\alpha_i m - \sqrt{(m + 2\nu)\alpha_i(1 - \alpha_i)} \right) \frac{\Delta\omega_i}{\sigma_e} \right] , \quad (3.83)$$

$$s = m - 1, \omega \approx (m - 1)\Omega_i$$

$$1 \leq \mathcal{K} \left[(m - 1) \frac{\Delta\omega_i}{\sigma_e} \right] \alpha_i^{m+2\nu-1} (m + \nu)(1 - \alpha_i)(1 - (1 - \alpha_i)m). \quad (3.84)$$

If any of these inequalities hold, the mode is unstable. In all cases

$$\mathcal{B} = \frac{T_e}{T_i/Z} m \alpha_i (1 - \alpha_i) \frac{\Delta\omega_i}{\sigma_e} \quad (3.85)$$

which is often of negligible magnitude for threshold cases of β close to -1.

In general, all of these solutions are unstable for large m or small k_z , and stable for large k_z .

3.3.4 Wall Modes

We define wall modes as those for which $\mathcal{F}$ is negative; $\mathcal{F}$ can be negative only if ν is not an integer, which implies finite wall radius. We investigate negative $\mathcal{F}$ by explicitly calculating Equation 3.56 and searching for the largest negative magnitude versus s . The results for the case $r_w = 4r_0$ are listed in Table 3-2. Various combinations of m and ν in this case were brought over from section 3.1.4. Only values of $\alpha < .5$ are shown, because the minimum of $\mathcal{F}(\alpha)$ equals the minimum of $\mathcal{F}(1 - \alpha)$.

One can see that these negative values of $\mathcal{F}$ are of very small magnitude and therefore stabilizing. It is unlikely therefore that there are unstable wall modes at moderate frequencies. For high frequencies, the electron dielectric constant passes through zero, and the modes described in section 3.2.4 may be unstable.

		alpha				
		.01	.03	.1	.3	.5
m=0	nu=0.0	0	0	0	0	0
m=1	nu=.02	-9.9e-7	-9.1e-6	-1.1e-4	-1.1e-3	-1.6e-3
m=2	nu=.06	-2.8e-6	-2.5e-5	-2.7e-4	-1.9e-3	-1.1e-3
m=3	nu=.13	-5.6e-6	-4.9e-5	-4.8e-4	-2.2e-3	-7.0e-4
m=4	nu=.26	-9.4e-6	-8.0e-5	-7.2e-4	-2.2e-3	-4.3e-4
m=5	nu=.44	-1.2e-5	-9.8e-5	-7.9e-4	-1.7e-3	-2.2e-4
m=0	nu=1.08	-1.3e-8	-3.7e-7	-1.4e-5	-4.4e-4	-1.5e-3
m=1	nu=1.22	-3.5e-8	-9.3e-7	-3.3e-5	-6.7e-4	-1.1e-3
m=2	nu=1.45	-5.9e-8	-1.5e-6	-4.8e-5	-6.4e-4	-5.6e-4
m=0	nu=2.51	-4.e-10	-3.1e-8	-3.5e-6	-1.9e-4	-5.7e-4

Table 3-2: Most negative value of $\mathcal{F}$ vs. α_i . The values of ν correspond to the case $r_w = 4r_0$.

3.4 Numerical Solution

Here we treat the general case numerically, including non-zero ν . Recall the electron dielectric constant

$$\epsilon_e = \frac{n_e q_e^2}{T_e} \left\{ -1 + i(\omega + m\omega_e)(1 - (m + 2\nu)\alpha_e) \frac{\sqrt{\pi}}{\sigma_e} (w(z_0) + \alpha_e(m + \nu)w(z_+) + \alpha_e \nu w(z_-)) \right\}$$

where

$$\begin{aligned} z_0 &\equiv -(\omega + m\omega_e - \alpha_e m \Delta\omega_e) / \sigma_e, \\ z_+ &\equiv -(\omega + m\omega_e + \Delta\omega_e - \alpha_e m \Delta\omega_e) / \sigma_e, \\ z_- &\equiv -(\omega + m\omega_e - \Delta\omega_e - \alpha_e m \Delta\omega_e) / \sigma_e, \end{aligned}$$

and ion dielectric constants

$$\begin{aligned} \epsilon_{i,<} &= \frac{n_i q_i^2}{T_i} \left\{ -1 + i(\omega + m\omega_i)(1 - \alpha_i)^{m+2\nu} \frac{\sqrt{\pi}}{\sigma_i} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \left(\frac{\alpha_i}{1 - \alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} w(z_{<}) \right\} \\ \epsilon_{i,>} &= \frac{n_i q_i^2}{T_i} \left\{ -1 + i(\omega + m\omega_i) \alpha_i^{m+2\nu} \frac{\sqrt{\pi}}{\sigma_i} \sum_{p=0}^{\infty} \sum_{q=0}^{\infty} \left(\frac{1 - \alpha_i}{\alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} w(z_{>}) \right\} \end{aligned}$$

with

$$\begin{aligned} z_{<} &= -(\omega + m\omega_i + (p - q)\Delta\omega_i - m\alpha_i \Delta\omega_i) / \sigma_i, \\ z_{>} &= -(\omega + m\omega_i - (p - q)\Delta\omega_i + (1 - \alpha_i)m\Delta\omega_i) / \sigma_i. \end{aligned}$$

In the manner corresponding to the analytical derivations we let ω be expressed as

$$\omega = -m\omega_i + m\alpha_i \Delta\omega_i - n\sigma_i \cdot \text{scale} \quad (3.86)$$

where n is the new frequency index. Scale is picked small enough so that the frequency is adequately sampled. The relevant terms become

$$\frac{\omega + m\omega_e}{\sigma_e} = \frac{m\alpha_e\Delta\omega_e - m\omega^2/\Delta\omega_i - n\sigma_i \cdot \text{scale}}{\sigma_e}, \quad (3.87)$$

$$-\frac{\omega + m\omega_e - m\alpha_e\Delta\omega_e}{\sigma_e} = \frac{mW^2/\Delta\omega_i + n\sigma_i \cdot \text{scale}}{\sigma_e}, \quad (3.88)$$

$$\frac{\omega + m\omega_i}{\sigma_i} = \frac{m\alpha_i\Delta\omega_i - n\sigma_i \cdot \text{scale}}{\sigma_i}, \quad (3.89)$$

$$-\frac{\omega + m\omega_i - m\alpha_i\Delta\omega_i}{\sigma_i} = n \cdot \text{scale}. \quad (3.90)$$

We define the following constants (and retain the $\Delta\omega_i > 0$ convention):

$$\begin{aligned} \frac{\alpha_e\Delta\omega_e}{\sigma_e} &= C_1 < 0, \\ \frac{W^2}{\Delta\omega_i\sigma_i \cdot \text{scale}} &= n_1 \geq 0, \\ \frac{\sigma_i \cdot \text{scale}}{\sigma_e} &= C_2 > 0, \\ \frac{\alpha_i\Delta\omega_i}{\sigma_i} &= C_3 > 0, \\ \frac{\Delta\omega_e}{\sigma_e} &= C_4 < 0, \\ \frac{\Delta\omega_i}{\text{scale} \cdot \sigma_i} &= n_2 > 0. \end{aligned} \quad (3.91)$$

We next consider specifically the sums

$$\Sigma_{<} = (1 - \alpha_i)^{m+2\nu} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{\alpha_i}{1 - \alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} w((n - (p - q)n_2) \cdot \text{scale})$$

$$\Sigma_{>} = \alpha_i^{m+2\nu} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{1 - \alpha_i}{\alpha_i} \right)^{p+q} \binom{m+\nu}{p} \binom{\nu}{q} w((n - (m - p + q)n_2) \cdot \text{scale}).$$

We can efficiently calculate Σ by noting the recursion relation

$$\Sigma(m + 1, \nu, n) = \alpha_i \Sigma(m, \nu, n - n_2) + (1 - \alpha_i) \Sigma(m, \nu, n)$$

and recursively calculating $\Sigma(m, \nu, n)$ from $\Sigma(0, \nu, n)$. Since the overall normalization nq^2/T has no relevance by itself, we form the quotient

$$\frac{n_i q_i^2 / T_i}{n_e q_e^2 / T_e} = \frac{T_e}{T_i / Z} \quad (3.92)$$

and apply this to the ion dielectric constant. We call this C_5 . We are now left with

$$\begin{aligned} \epsilon_e = & -1 + i[mC_1 - (n + mn_1)C_2][1 - (m + 2\nu)\alpha_e]\sqrt{\pi} \\ & (w((n + mn_1)C_2) + \alpha_e(m + \nu)w((n + mn_1)C_2 - C_4) \\ & + \alpha_e \nu w((n + mn_1)C_2 + C_4))) \end{aligned} \quad (3.93)$$

$$\epsilon_i = C_5 \left\{ -1 + i(mC_3 - n \cdot \text{scale})\sqrt{\pi}\Sigma(m, \nu, n) \right\} \quad (3.94)$$

with

$$\Sigma(m + 1, \nu, n) = \alpha_i \Sigma(m, \nu, n - n_2) + (1 - \alpha_i) \Sigma(m, \nu, n), \quad (3.95)$$

$$\Sigma_{<}(0, \nu, n) = (1 - \alpha_i)^{2\nu} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{\alpha_i}{1 - \alpha_i} \right)^{p+q} \binom{\nu}{p} \binom{\nu}{q} w((n - (p - q)n_2) \cdot \text{scale})$$

$$\Sigma_{>}(0, \nu, n) = \alpha_i^{2\nu} \sum_{\substack{p=0 \\ q=0}}^{\infty} \left(\frac{1 - \alpha_i}{\alpha_i} \right)^{p+q} \binom{\nu}{p} \binom{\nu}{q} w(n - (p - q)n_2) \cdot \text{scale}.$$

Finally, we must search for occurrences of $\epsilon_i = \beta \epsilon_e$ with β real and ≤ -1 . When we find such an occurrence, we report the relevant parameters.

The program written to perform this numerical investigation is called STAB5 and is listed in Appendix A along with the calculated results. STAB5 steps through values of α_i , ν , and $\Delta\omega_i/\sigma_i$, and finds the first m for which an instability occurs. Invariably, larger values of m produce ever increasing numbers of instabilities. In this case we set $\Delta\omega_e/\Delta\omega_i = -4000$ and $ZT_e/T_i = 0.2$. Also reported is s and δ .

An additional constraint must be imposed to filter out spurious numerical artifacts, due to the fact that the infinite sums for non-integer ν cannot be exactly evaluated. Specifically, values of ϵ_e which are purely real to one part in 10^6 or better are discarded. This does have the unfortunate drawback of excluding the modes predicted in section 3.2.4.

The results presented in Appendix A agree reasonably well with the predictions from section 3.3. One noteworthy point is the weak dependence of ν on instability. We exploit this with a modified program STAB5A which evaluates stability at various values of $\Delta\omega_e/\Delta\omega_i$ and ZT_e/T_i , but at $\nu = 0$ only. These results are listed in Appendix B.

4 Conclusions

4.1 Rigid Rotor Condition

The first interesting and somewhat surprising result is that all V-M distributions are rigid rotors (section 2.1). In other words, the rotational drift velocity must be of the form $\langle v_\theta \rangle = r\omega_j$ where ω_j are constants, or the distribution function will not satisfy the Vlasov equation with $\dot{f}_j = 0$. We are not aware of any previous derivations of this result. Naturally, the rigid rotor condition must break down as the plasma approaches the wall, but may be attainable for the high density collisional portions of the plasma.

4.2 Equilibrium Solutions

The deep V-M equilibrium solutions (section 2.2) seem to be different from those experimentally observed. They may represent an interesting plasma configuration, but no prediction about their stability is possible at this point. The high density contrast is particularly attractive. A point in favor of stability for these solutions is their high k_z equivalent content, which was found in section 3 to be stabilizing in the low-beta case.

An interesting feature of the deep solutions is their localized nature, which was dubbed in [23] as the “bicycle wheel” configuration. These localized regions seem to be fairly independent of the boundary conditions. A curious puzzle is the fact that the unbound 1-D solution (solved from the same equations with the z -dependence removed) cannot be obtained as the limit of the 2-D finite solutions and hence appear to be unphysical. Elongated solutions have previously been obtained[28,29] that do approach an unbound 1-D solution, but only for a case which is purely adiabatic and incompatible with the Maxwellian condition.

4.3 General Stability Observations

Here we will briefly summarize the stability results from section 3. The independent parameters are taken to be m , ν , k_z , Ω_e/Ω_i , ZT_e/T_i , W , and α :

m : higher azimuthal mode number invariably causes more instability. Linear theory is not really relevant at high mode number (since high spatial frequencies tend to rapidly saturate), and so we searched only to $m = 10$.

ν : radial mode number seems to have little effect on stability for the regimes we numerically searched. This is due to the symmetry of the radial modes, which do not shift the maxima of the ion dielectric constant. The azimuthal modes in contrast do shift the maxima to higher frequencies.

k_z : low values of $k_z v_i^{th}/\Omega_i$ are always destabilizing. It is problematical how to interpret k_z , but we treat k_z as being defined by the spacing l between the conductive walls at the endcaps. For that case $k_z^{max} = \pi/l$. Of course, the plasma is not confined from the walls in that configuration, and any interpretation must make an assumption about axial confinement. The effect of k_z is to provide diffusion and hence damp the modes.

Ω_e/Ω_i : higher values are slightly more stable, but there is little effect over the range searched. The values we used corresponded to H, D, and He-3.

ZT_e/T_i : higher values are destabilizing. This makes physical sense, because larger electron temperatures broaden and consequently reduce the amplitude of the low-frequency electron dielectric constant peak.

W : the $E \times B$ electron drift rotation frequency is only important for α small, in which case the plasma is most stable at $W = 0$.

α : when α_i is very near 0, the plasma can be unstable for sufficiently large values of W . When α_i exceeds a small value, it stabilizes this tilt mode and enters its most stabilizing regime. As α_i increases, it gradually destabilizes the plasma.

Unfortunately, there are no particularly simple scaling laws which are generally applicable. Section 3.3 gives the best indication of parameter scaling.

4.4 High Frequency Instability

It is predicted in section 3.2.4 that there may be an unusual high frequency instability when $\alpha = 1/2$. It is due to an unusually large high frequency ion dielectric constant which may be obtainable at very high harmonic numbers of Ω_i for this particular configuration. Notice that in the integral in equation 3.22, S_i passes through 0 only for the unique case $\alpha = 1/2$, which forces the integrand through a branch point if $\nu \neq (\text{integer})$. No analogue to this instability has been found in any other reference.

4.5 Classification of Instabilities

This dissertation presents the first complete investigation of electrostatic instabilities in a plasma configuration which has cylindrical geometry. We therefore must be satisfied with comparing the modes which we have discovered to electrostatic modes propagating in a planar gradient, which have been studied previously. The classification of these instabilities is determined by the effect which produces the instability.

The unstable modes which we have found at the $s = 0$ harmonic reside within both the electron and ion drift frequency peaks, and are consequently called "drift instabilities". These drift waves propagate with and are driven by the particles moving at the drift velocity. There is a natural correspondence between these modes and (for example) the "Low-frequency Drift Waves Propagating Oblique to B_0 " in [27] which are planar modes. That treatment and others [30] neglect the harmonics of Ω_i , but present the conditions for drift instability. Both sources suffer from the limiting assumption that the argument of the w-function is either very

large or very small. In contrast, we find that intermediate values for the argument in the peaks of ϵ_i are the most important, which adds considerable complexity to the calculation. The drift instability is sometimes called the “universal instability”.

Reference [30] also discusses planar modes for which $\omega/k_z > v_e^{th}$ so that the frequency is off of the electron drift peak, and again other harmonics of Ω_i are neglected. This corresponds to the off-peak MHD modes which we derived in section 3.3.1. These modes are termed “interchange instabilities” or sometimes “flute-interchange instabilities” since they seem to exist only for k_z very near 0. These waves propagate faster than the drift velocity.

A general discussion of drift waves and their properties can be found in [31]. This reference considers interchange modes to be a subset of drift modes.

In section 3 we found modes with $s > 0$ which become unstable due to harmonics of the ion cyclotron frequency at $\omega_i + s\Omega_i$. This effect was studied for the planar case in [32], which derives conditions for the “intersection” of the electron drift peak with the ion cyclotron harmonic peaks, similar to our treatment in 3.3.2. The earlier treatment used arguments for the w -function that were limited to the asymptotic regions, however. Modes which lie on these ion cyclotron harmonics are most reasonably termed “drift-cyclotron instabilities” or just “cyclotron instabilities”.

We have found in this dissertation that as Larmor radius is increased, the drift instabilities stabilize but are replaced with cyclotron instabilities.

4.6 Connection with Previous Results for MHD Modes

It is interesting to note that the treatment in section 3.3.1 predicts both the tilt mode and its stabilization (the tilt mode is defined as the electrostatic $m = 1$, $k_z > 0$ mode). We found that the tilt mode is a drift instability which stabilizes when

$$\alpha_i > W^2/\Omega_i^2 ,$$

or equivalently,

$$v_i^{th} > \sqrt{2}r_0|W| = \sqrt{2}\langle v_i^D \rangle ,$$

where $\langle v_i^D \rangle$ is the average ion drift velocity. This condition is met for most reasonable plasmas, which may be why the tilt mode is not experimentally observed. This drift mode is completely different from the classical MHD interchange instability, which was derived later in that same section. We found that the interchange mode stabilizes when

$$k_z r_0 > mW / \sqrt{|\Delta\omega_e \Delta\omega_i|} .$$

Since this is relevant only for very small k_z , it is therefore not a tilt mode. This mode was never observed in any of our numerical results because the extremely small values of k_z required to see this mode would first cause all of the drift modes to become unstable and dominate the calculation. We believe that the drift instabilities dominate the interchange instabilities in physical systems as well .

The stabilization of the interchange instability by non-zero α_i was first derived in [20] to include finite Larmor radius effects, for the case α and W small and $k_z = 0$. We similarly derived this result in section 3.3.1 to connect with the previous literature. An interesting new result was found when we considered non-integer ν (wall effect) which causes increased instability for certain modes, depending on the relative rotation of the electrons and ions. For $\Delta\omega_i/W < 0$, the mode stabilizes when $\alpha_i > |W/\Delta\omega_i|$, except for the case $\alpha_i \Delta\omega_i / 2W = -1$. For $\Delta\omega_i/W > 0$, however, the mode stabilizes at $\alpha_i > 2W/\Delta\omega_i \nu_{min}^{m=1}$ which is a much larger value.

The interchange instability seems to be specific to very small k_z and is unique in that it does not fall on a peak. The implication is that these modes cannot be extrapolated beyond a narrow window of parameters.

Both $k_z = 0$ and $\alpha = 0$ represent pathological cases, as for both cases the dielectric constant peaks are of infinitesimal width and consequently the drift instabilities are sometimes erroneously ignored. In section 3 we find that most of the unstable modes are actually on these peaks, including the tilt mode instability.

4.7 Suggested Future Research

The natural problem to tackle would seem to be a stability analysis of the high-beta, deep solutions of section 2.2. This has the potential for producing a very exciting result if stability is found for some set of parameters. There are two major challenges, however. First, these solutions are high-beta, and so the perturbation in the magnetic field δB must be included. This complicates the algebra considerably. Second, the equilibria are found numerically, and consequently the orbits must also be found numerically. This removes any hope of an analytic treatment. Perhaps particle simulation of the plasma would be a more practical approach. Another possible approach might be use of a variational principle, as in reference [17], which might prove fruitful due to the Maxwellian form of the distribution functions.

A more modest problem would be to track the zeros of the eigenvalue equation as parameters are varied, thereby obtaining growth rates rather than just stability. This also allows the modes to be continuously followed, allowing a more clear definition of the modes. A challenge here is to usefully tabulate the results over the large parameter space.

References

- [1] W. W. Heidbrink, et al., Phys. Fluids B, Vol.3, 3167 (1991).
- [2] M. Tuszewski, Nuclear Fusion, Vol.28, No.11, 2033 (1988).
- [3] H. Fleischmann, et al., Phys. Fluids, Vol.17, 2226 (1974).
- [4] H. Fleischmann, et al., Phys. Fluids, Vol.19, 728 (1976).
- [5] A. Mohri, Nuclear Fusion, Vol.25, 1299 (1985).
- [6] R. Sieman, et al., Fusion Technology, Vol.9, 13 (1986).
- [7] N. Christophilos, Proc. 2nd United Nations Intl. Conf. on the Peaceful Uses of Atomic Energy, (UN, Geneva 1958), p.279
- [8] R. Lovelace, Phys. Fluids, Vol.21, 1389 (1978).
- [9] R. Lovelace, Phys. Fluids, Vol.22, 591 (1979).
- [10] R. Lovelace, Phys. Fluids, Vol.22, 977 (1980).
- [11] R. Sudan, M. Rosenbluth, Phys. Rev. Letters, Vol.36, 972 (1976).
- [12] J. Finn, R. Sudan, Phys. Rev. Letters, Vol.41, 695 (1978).
- [13] P. Bell, et al., IAEA Conference Proceedings Culham Sept. 6-10, Vol.2, 77 (1965).
- [14] B. Maglich, Nuc. Inst. and Methods, Vol.151, 1 (1978).
- [15] B. Maglich, Nuc. Inst. and Methods A, Vol.271, 13 (1988).
- [16] D. Al-Salameh, et al., Phys. Rev. Letters, Vol.54, 796 (1985).

- [17] H. Wang, H. Berk, R. Lovelace, N. Rostoker, Phys. Fluids B, Vol.3, No.11, 2973 (1991).
- [18] D. Barnes, et al., Phys. Fluids, Vol.29, 2616 (1986).
- [19] D. Barnes, et al., Workshop on Compact Torus Research Los Alamos Nov. 6-8 (1989).
- [20] M. N. Rosenbluth, N. A. Krall, N. Rostoker, Nuclear Fusion: 1962 Supplement, Part 1, 143 (1962).
- [21] B. Mardner, H. Weitzner, Plasma Physics, Vol.12, 435 (1970).
- [22] D. V. Anderson, J. Killeen, M. E. Resnick, Phys. Fluids, Vol.15, No.2, 351 (1972).
- [23] D. A. Larrabee, R. V. Lovelace, H. H. Fleischmann, Nuclear Fusion, Vol.19, No.4, 499 (1979).
- [24] M. Abramowitz, I. Stegun, Handbook of Mathematical Functions, (Dover, New York 1968), p.504
- [25] Ibid., p.297
- [26] N. Krall, A. Trivelpiece, Principles of Plasma Physics, (San Francisco Press, San Francisco 1986), p.11
- [27] Ibid., p.430
- [28] R. Spencer, D. Hewitt, Phys. Fluids, Vol.25, 1365 (1982).
- [29] D. Hewitt, R. Spencer, Phys. Fluids, Vol.26, 1299 (1983).
- [30] K. Miyamoto, Plasma Physics for Nuclear Fusion, (MIT Press, Cambridge 1976), p.371

- [31] N. Krall, *Advances in Plasma Physics*, eds. A. Simon, W. B. Thompson, (Interscience, New York 1968), p.153
- [32] A.Mikhailovskii, A.Timofeev, *Soviet Physics JETP*, Vol.17, No.3, 626 (1963).

A Appendix

The program written to perform the numerical investigation is called STAB5 and is listed here along with the calculated results. STAB5 steps through values of α_i , ν , and $\Delta\omega_i/\sigma_i$, and finds the first m for which an instability occurs. Invariably, larger values of m produce ever increasing numbers of instabilities. In this case we set $\Delta\omega_e/\Delta\omega_i = -4000$ and $ZT_e/T_i = 0.2$. Also reported is s and δ .


```

program stab5
parameter(nmax=10001,nmax2=5001,mmax=10)
complex w,wi(nmax),we_center(nmax),we_left(nmax),we_right(nmax)
complex eps_i(nmax),eps_e(nmax),sum_i(nmax)
real alparm(12),dwparm(4),anuparm(11)
data alparm/.01,.02,.05,.1,.2,.4,.6,.8,.9,.95,.98,.99/
data dwparm/5.,10.,20.,40./
data anuparm/0.,.5,1.,1.5,2.,2.5,3.,3.5,4.,4.5,5./
open(1,file='stab5.dat',access='sequential')

c
c
c
set parameters

n1=0
scale=.1
c5=.2
dwe_by_dwi=-4000.
do 300 idw=1,4
dwi_by_si=dwparm(idw)
do 300 ialp=1,12
alpha_i=alparm(ialp)
do 300 inu=1,11
anu=anuparm(inu)

c
c
c
derived parameters

alpha_e=-c5*alpha_i*(1.0-alpha_i)/dwe_by_dwi
se_by_si=sqrt(-c5*dwe_by_dwi)
c1=alpha_e*dwe_by_dwi*dwi_by_si/se_by_si
c2=scale/se_by_si
c3=alpha_i*dwi_by_si
c4=dwe_by_dwi*dwi_by_si/se_by_si
n2=nint(dwi_by_si/scale)
sqrtpi=sqrt(3.1415926)

c
c
c
form w functions

do 10 n=1,nmax
w1(n)=w((n-nmax2)*scale)
we_center(n)=w((n-nmax2)*c2)
we_left(n)=w((n-nmax2)*c2-c4)
we_right(n)=w((n-nmax2)*c2+c4)
continue

c
c
c
form ion sum for m=0

call sumform(w1,sum_i,nmax,alpha_i,anu,n2)

c
c
c
form dielectric constants

do 100 m=0,mmax
do 30 n=1,nmax

c
eps_i(n)=
& c5*(-1.0+cmplx(0.0,1.0)*(m*c3-(n-nmax2)*scale)*sqrtpi*sum_i(n))
c
nshift=n+m*n1
if(nshift.gt.nmax) nshift=nmax

c
eps_e(n)=-1.0+cmplx(0.0,1.0)*(m*c1-(nshift-nmax2)*c2)*sqrtpi*
& (1.0-(m+2.0*anu)*alpha_e)*
& (we_center(nshift)+alpha_e*(m+anu)*we_left(nshift)+
& alpha_e*anu*we_right(nshift))
c
nshift=n-n2
if(nshift.ge.1.and.nshift.le.nmax) then
wi(n)=alpha_i*sum_i(nshift)+(1.0-alpha_i)*sum_i(n)
else

```

```

                                wi(n)=sum_i(n)
endif
30  continue
c
do 47 n=1,nmax
47  sum_i(n)=wi(n)
c
call eval(eps_e,eps_i,nmax,nmax2,m,anu,dwi_by_si,alpha_i,scale,iflag)
if(iflag.eq.1) go to 300
100 continue
write(1,1000) anu,dwi_by_si,alpha_i
write(*,1000) anu,dwi_by_si,alpha_i
1000 format(1x,'***',1x,f4.1,1x,f5.1,1x,f4.2)
300 continue
close(1)
end

complex function w(x)
complex w
pi=3.1415926
fact=1./sqrt(pi)
if(x.lt.10.) then
    parm=exp(-x*x)
else
    parm=0.
endif
if(abs(x).lt.6.) go to 600
aml=1.
a0=0.
bml=0.
a=1.
n=1
fo=0.
60  an=x*a0+a*aml
    bn=x+a*bml
    fn=an/bn
    if(abs(1-fo/fn).lt.1.e-7) go to 500
    fo=fn
    n=n+1
    a=-.5*(n-1)
    aml=a0/bn
    a0=fn
    bml=1./bn
    go to 60
500 w=cmlpx(parm,fn*fact)
    return
600 sum=1.
    n=1
    term=x*x
610 sumn=sum+term/(2*n+1)
    term=term*x*x/(n+1)
    n=n+1
    if(abs(term/sumn).lt.1.e-7) go to 1000
    sum=sumn
    go to 610
1000 w=parm*cmlpx(1.,x*2.*fact*sumn)
    end

subroutine eval(eps_e,eps_i,nmax,nmax2,m,anu,dwi_by_si,alpha_i,
scale,iflag)
complex eps_i(nmax),eps_e(nmax),ctempl,ctemp2
iflag=0
do 10 i=1,nmax-1
phase=aimag(eps_e(i))/real(eps_e(i))

```

```

if(abs(phase).lt.1.e-6) go to 10
ctemp1=eps_i(i)/eps_e(i)
ctemp2=eps_i(i+1)/eps_e(i+1)
a1=sign(1.,aimag(ctemp1))*sign(1.,aimag(ctemp2))
a2=.5*real(ctemp1)+.5*real(ctemp2)
if(a1.gt.0.or.a2.gt.-1.0) go to 10
iflag=1
is=nint(scale*(i+.5-nmax2)/dwi_by_si)
delt=scale*(i+.5-nmax2)-is*dwi_by_si
write(1,100) m,anu,dwi_by_si,alpha_i,is,delt
write(*,100) m,anu,dwi_by_si,alpha_i,is,delt
format(1x,i3,1x,f4.1,1x,f5.1,1x,f4.2,1x,i3,1x,f5.1)
100 continue
10 end

subroutine sumform(wi,sum_i,nmax,alpha_i,anu,n2)
complex wi(nmax),sum_i(nmax)
c
if(alpha_i.lt.0.5) then
a=alpha_i
else
a=1.0-alpha_i
endif
aratio=a/(1.0-a)
c
do 20 n=1,nmax
j=0
sum_i(n)=0.0
comb=1.0
aprod=1.0
21 ip=n+n2*j
if(ip.gt.nmax) go to 20
sum_i(n)=sum_i(n)+wi(ip)*aprod*comb
aprod=aprod*aratio
comb=comb*(anu-j)/float(j+1)
j=j+1
go to 21
20 continue
c
do 22 n=1,nmax
22 wi(n)=sum_i(n)
c
do 24 n=1,nmax
j=0
sum_i(n)=0.0
comb=1.0
aprod=1.0
23 ip=n-n2*j
if(ip.lt.1) go to 24
sum_i(n)=sum_i(n)+wi(ip)*aprod*comb
aprod=aprod*aratio
comb=comb*(anu-j)/float(j+1)
j=j+1
go to 23
24 continue
c
const=(1.-a)**(2.0*anu)
do 25 n=1,nmax
25 sum_i(n)=sum_i(n)*const
c
end

```

first unst. m	nu	omega over sigm	alph	s	delt
>10	0.0	5.0	0.01		
>10	1.0	5.0	0.01		
>10	2.0	5.0	0.01		
>10	3.0	5.0	0.01		
>10	4.0	5.0	0.01		
>10	5.0	5.0	0.01		
>10	0.0	5.0	0.02		
>10	1.0	5.0	0.02		
>10	2.0	5.0	0.02		
>10	3.0	5.0	0.02		
>10	4.0	5.0	0.02		
>10	5.0	5.0	0.02		
>10	0.0	5.0	0.05		
>10	1.0	5.0	0.05		
>10	2.0	5.0	0.05		
>10	3.0	5.0	0.05		
>10	4.0	5.0	0.05		
>10	5.0	5.0	0.05		
>10	0.0	5.0	0.10		
>10	1.0	5.0	0.10		
>10	2.0	5.0	0.10		
>10	3.0	5.0	0.10		
>10	4.0	5.0	0.10		
>10	5.0	5.0	0.10		
>10	0.0	5.0	0.20		
>10	1.0	5.0	0.20		
>10	2.0	5.0	0.20		
>10	3.0	5.0	0.20		
>10	4.0	5.0	0.20		
>10	5.0	5.0	0.20		
>10	0.0	5.0	0.40		
>10	1.0	5.0	0.40		
>10	2.0	5.0	0.40		
>10	3.0	5.0	0.40		
>10	4.0	5.0	0.40		
>10	5.0	5.0	0.40		
>10	0.0	5.0	0.60		
>10	1.0	5.0	0.60		
>10	2.0	5.0	0.60		
>10	3.0	5.0	0.60		
>10	4.0	5.0	0.60		
>10	5.0	5.0	0.60		
>10	0.0	5.0	0.80		
>10	1.0	5.0	0.80		
>10	2.0	5.0	0.80		
>10	3.0	5.0	0.80		
>10	4.0	5.0	0.80		
>10	5.0	5.0	0.80		
>10	0.0	5.0	0.90		
>10	1.0	5.0	0.90		
>10	2.0	5.0	0.90		
>10	3.0	5.0	0.90		
>10	4.0	5.0	0.90		
>10	5.0	5.0	0.90		

first unst. m	nu	omega over sigm	alph	s	delt
>10	0.5	5.0	0.01		
>10	1.5	5.0	0.01		
>10	2.5	5.0	0.01		
>10	3.5	5.0	0.01		
>10	4.5	5.0	0.01		
>10	0.5	5.0	0.02		
>10	1.5	5.0	0.02		
>10	2.5	5.0	0.02		
>10	3.5	5.0	0.02		
>10	4.5	5.0	0.02		
>10	0.5	5.0	0.05		
>10	1.5	5.0	0.05		
>10	2.5	5.0	0.05		
>10	3.5	5.0	0.05		
>10	4.5	5.0	0.05		
>10	0.5	5.0	0.10		
>10	1.5	5.0	0.10		
>10	2.5	5.0	0.10		
>10	3.5	5.0	0.10		
>10	4.5	5.0	0.10		
>10	0.5	5.0	0.20		
>10	1.5	5.0	0.20		
>10	2.5	5.0	0.20		
>10	3.5	5.0	0.20		
>10	4.5	5.0	0.20		
>10	0.5	5.0	0.40		
>10	1.5	5.0	0.40		
>10	2.5	5.0	0.40		
>10	3.5	5.0	0.40		
>10	4.5	5.0	0.40		
>10	0.5	5.0	0.60		
>10	1.5	5.0	0.60		
>10	2.5	5.0	0.60		
>10	3.5	5.0	0.60		
>10	4.5	5.0	0.60		
>10	0.5	5.0	0.80		
>10	1.5	5.0	0.80		
>10	2.5	5.0	0.80		
>10	3.5	5.0	0.80		
>10	4.5	5.0	0.80		
>10	0.5	5.0	0.90		
>10	1.5	5.0	0.90		
>10	2.5	5.0	0.90		
>10	3.5	5.0	0.90		
>10	4.5	5.0	0.90		

first unst.	nu m	omega over sigm	alph	s	delt
>10	0.0	5.0	0.95		
>10	1.0	5.0	0.95		
>10	2.0	5.0	0.95		
>10	3.0	5.0	0.95		
>10	4.0	5.0	0.95		
>10	5.0	5.0	0.95		
>10	0.0	5.0	0.98		
>10	1.0	5.0	0.98		
>10	2.0	5.0	0.98		
>10	3.0	5.0	0.98		
>10	4.0	5.0	0.98		
>10	5.0	5.0	0.98		
>10	0.0	5.0	0.99		
>10	1.0	5.0	0.99		
>10	2.0	5.0	0.99		
>10	3.0	5.0	0.99		
>10	4.0	5.0	0.99		
>10	5.0	5.0	0.99		
>10	0.0	10.0	0.01		
>10	1.0	10.0	0.01		
>10	2.0	10.0	0.01		
>10	3.0	10.0	0.01		
>10	4.0	10.0	0.01		
>10	5.0	10.0	0.01		
>10	0.0	10.0	0.02		
>10	1.0	10.0	0.02		
>10	2.0	10.0	0.02		
>10	3.0	10.0	0.02		
>10	4.0	10.0	0.02		
>10	5.0	10.0	0.02		
>10	0.0	10.0	0.05		
>10	1.0	10.0	0.05		
>10	2.0	10.0	0.05		
>10	3.0	10.0	0.05		
>10	4.0	10.0	0.05		
>10	5.0	10.0	0.05		
>10	0.0	10.0	0.10		
>10	1.0	10.0	0.10		
>10	2.0	10.0	0.10		
>10	3.0	10.0	0.10		
>10	4.0	10.0	0.10		
>10	5.0	10.0	0.10		
>10	0.0	10.0	0.20		
>10	1.0	10.0	0.20		
>10	2.0	10.0	0.20		
>10	3.0	10.0	0.20		
>10	4.0	10.0	0.20		
>10	5.0	10.0	0.20		
10	0.0	10.0	0.40	3	0.3
>10	1.0	10.0	0.40		
7	1.5	10.0	0.40	-13	0.8
>10	2.5	10.0	0.40		
>10	3.5	10.0	0.40		
>10	4.5	10.0	0.40		

first unst.	nu m	omega over sigm	alph	s	delt
>10	0.5	5.0	0.95		
>10	1.5	5.0	0.95		
>10	2.5	5.0	0.95		
>10	3.5	5.0	0.95		
>10	4.5	5.0	0.95		
>10	0.5	5.0	0.98		
>10	1.5	5.0	0.98		
>10	2.5	5.0	0.98		
>10	3.5	5.0	0.98		
>10	4.5	5.0	0.98		
>10	0.5	5.0	0.99		
>10	1.5	5.0	0.99		
>10	2.5	5.0	0.99		
>10	3.5	5.0	0.99		
>10	4.5	5.0	0.99		
>10	0.5	10.0	0.01		
>10	1.5	10.0	0.01		
>10	2.5	10.0	0.01		
>10	3.5	10.0	0.01		
>10	4.5	10.0	0.01		
>10	0.5	10.0	0.02		
>10	1.5	10.0	0.02		
>10	2.5	10.0	0.02		
>10	3.5	10.0	0.02		
>10	4.5	10.0	0.02		
>10	0.5	10.0	0.05		
>10	1.5	10.0	0.05		
>10	2.5	10.0	0.05		
>10	3.5	10.0	0.05		
>10	4.5	10.0	0.05		
>10	0.5	10.0	0.10		
>10	1.5	10.0	0.10		
>10	2.5	10.0	0.10		
>10	3.5	10.0	0.10		
>10	4.5	10.0	0.10		
>10	0.5	10.0	0.20		
>10	1.5	10.0	0.20		
>10	2.5	10.0	0.20		
>10	3.5	10.0	0.20		
>10	4.5	10.0	0.20		
>10	0.5	10.0	0.40		
7	1.5	10.0	0.40	-13	-1.5
>10	2.0	10.0	0.40		
>10	3.0	10.0	0.40		
>10	4.0	10.0	0.40		
>10	5.0	10.0	0.40		

first unst. m	nu	omega over sigm	alph	s	delt
7	0.0	10.0	0.60	3	0.3
7	1.0	10.0	0.60	3	0.2
7	2.0	10.0	0.60	3	0.2
7	3.0	10.0	0.60	3	0.2
7	4.0	10.0	0.60	3	0.2
8	5.0	10.0	0.60	3	0.3
5	0.0	10.0	0.80	3	0.2
5	1.0	10.0	0.80	3	0.2
5	2.0	10.0	0.80	3	0.2
5	3.0	10.0	0.80	3	0.2
5	4.0	10.0	0.80	3	0.1
5	5.0	10.0	0.80	3	0.1
5	0.0	10.0	0.90	4	0.5
4	1.0	10.0	0.90	3	0.1
4	2.0	10.0	0.90	3	0.1
4	3.0	10.0	0.90	3	0.1
5	4.0	10.0	0.90	3	0.2
5	4.5	10.0	0.90	3	0.2
5	5.0	10.0	0.90	3	0.2
5	0.0	10.0	0.95	4	0.6
5	1.0	10.0	0.95	4	0.6
5	2.0	10.0	0.95	4	0.6
4	3.0	10.0	0.95	3	0.1
4	4.0	10.0	0.95	3	0.1
4	5.0	10.0	0.95	3	0.1
7	0.0	10.0	0.98	6	1.5
6	1.0	10.0	0.98	5	1.0
6	2.0	10.0	0.98	5	1.0
6	3.0	10.0	0.98	5	1.0
6	4.0	10.0	0.98	5	1.0
5	5.0	10.0	0.98	4	0.6
7	0.0	10.0	0.99	7	-0.7
7	1.0	10.0	0.99	7	-0.7
7	2.0	10.0	0.99	7	-0.7
7	3.0	10.0	0.99	7	-0.7
7	4.0	10.0	0.99	7	-0.7
7	4.5	10.0	0.99	7	-0.7
7	5.0	10.0	0.99	7	-0.8
>10	0.0	20.0	0.01		
>10	1.0	20.0	0.01		
>10	2.0	20.0	0.01		
>10	3.0	20.0	0.01		
>10	4.0	20.0	0.01		
>10	5.0	20.0	0.01		
>10	0.0	20.0	0.02		
>10	1.0	20.0	0.02		
>10	2.0	20.0	0.02		
>10	3.0	20.0	0.02		
>10	4.0	20.0	0.02		
>10	5.0	20.0	0.02		
>10	0.0	20.0	0.05		
>10	1.0	20.0	0.05		
>10	2.0	20.0	0.05		
>10	3.0	20.0	0.05		
>10	4.0	20.0	0.05		

first unst. m	nu	omega over sigm	alph	s	delt
7	0.5	10.0	0.60	3	0.3
7	1.5	10.0	0.60	3	0.2
7	2.5	10.0	0.60	3	0.2
7	3.5	10.0	0.60	3	0.2
7	4.5	10.0	0.60	3	0.2
5	0.5	10.0	0.80	3	0.2
5	1.5	10.0	0.80	3	0.2
5	2.5	10.0	0.80	3	0.2
5	3.5	10.0	0.80	3	0.2
5	4.5	10.0	0.80	3	0.1
4	0.5	10.0	0.90	3	0.1
4	1.5	10.0	0.90	3	0.1
4	2.5	10.0	0.90	3	0.1
4	3.5	10.0	0.90	3	0.1
5	4.0	10.0	0.90	4	0.5
5	4.5	10.0	0.90	4	0.5
5	5.0	10.0	0.90	4	0.5
5	0.5	10.0	0.95	4	0.6
5	1.5	10.0	0.95	4	0.6
5	2.5	10.0	0.95	4	0.6
4	3.5	10.0	0.95	3	0.1
4	4.5	10.0	0.95	3	0.1
7	0.5	10.0	0.98	6	1.5
6	1.5	10.0	0.98	5	1.0
6	2.5	10.0	0.98	5	1.0
6	3.5	10.0	0.98	5	1.0
6	4.5	10.0	0.98	5	1.0
7	0.5	10.0	0.99	7	-0.7
7	1.5	10.0	0.99	7	-0.7
7	2.5	10.0	0.99	7	-0.7
7	3.5	10.0	0.99	7	-0.7
7	4.5	10.0	0.99	6	1.5
7	5.0	10.0	0.99	6	1.5
>10	0.5	20.0	0.01		
>10	1.5	20.0	0.01		
>10	2.5	20.0	0.01		
>10	3.5	20.0	0.01		
>10	4.5	20.0	0.01		
>10	0.5	20.0	0.02		
>10	1.5	20.0	0.02		
>10	2.5	20.0	0.02		
>10	3.5	20.0	0.02		
>10	4.5	20.0	0.02		
>10	0.5	20.0	0.05		
>10	1.5	20.0	0.05		
>10	2.5	20.0	0.05		
>10	3.5	20.0	0.05		
>10	4.5	20.0	0.05		

first unst. m	nu	omega over sigm	alph	s	delt
>10	5.0	20.0	0.05		
>10	0.0	20.0	0.10		
>10	1.0	20.0	0.10		
>10	2.0	20.0	0.10		
>10	3.0	20.0	0.10		
>10	4.0	20.0	0.10		
>10	5.0	20.0	0.10		
7	0.0	20.0	0.20	1	-0.2
6	0.5	20.0	0.20	-6	2.1
8	1.5	20.0	0.20	-5	-0.7
8	1.5	20.0	0.20	1	-0.2
6	2.5	20.0	0.20	-6	-1.9
8	3.0	20.0	0.20	1	-0.2
8	3.5	20.0	0.20	-5	-0.1
8	4.0	20.0	0.20	1	-0.2
6	4.5	20.0	0.20	-6	1.4
4	0.0	20.0	0.40	1	-0.2
4	0.5	20.0	0.40	-6	2.7
4	1.0	20.0	0.40	1	-0.2
4	2.0	20.0	0.40	1	-0.2
4	2.5	20.0	0.40	-6	2.0
4	3.0	20.0	0.40	1	-0.3
5	3.5	20.0	0.40	-5	0.7
5	4.0	20.0	0.40	1	-0.2
4	4.5	20.0	0.40	-6	1.9
3	0.0	20.0	0.60	1	-0.2
3	1.0	20.0	0.60	1	-0.2
3	2.0	20.0	0.60	1	-0.2
3	3.0	20.0	0.60	1	-0.2
3	4.0	20.0	0.60	1	-0.3
3	5.0	20.0	0.60	1	-0.3
2	0.0	20.0	0.80	1	-0.3
2	1.0	20.0	0.80	1	-0.3
2	2.0	20.0	0.80	1	-0.3
2	3.0	20.0	0.80	1	-0.3
2	4.0	20.0	0.80	1	-0.3
2	5.0	20.0	0.80	1	-0.4
2	0.0	20.0	0.90	1	-0.3
2	1.0	20.0	0.90	1	-0.3
2	2.0	20.0	0.90	1	-0.3
2	3.0	20.0	0.90	1	-0.3
2	4.0	20.0	0.90	1	-0.3
2	5.0	20.0	0.90	1	-0.3
3	0.0	20.0	0.95	2	0.7
3	1.0	20.0	0.95	2	0.7
2	2.0	20.0	0.95	1	-0.3
2	3.0	20.0	0.95	1	-0.3
2	4.0	20.0	0.95	1	-0.3
2	5.0	20.0	0.95	1	-0.3
4	0.0	20.0	0.98	3	1.5
4	1.0	20.0	0.98	3	1.5
3	2.0	20.0	0.98	2	0.7
3	3.0	20.0	0.98	2	0.7
3	4.0	20.0	0.98	2	0.7
3	5.0	20.0	0.98	2	0.7

first unst. m	nu	omega over sigm	alph	s	delt
>10	0.5	20.0	0.10		
>10	1.5	20.0	0.10		
>10	2.5	20.0	0.10		
>10	3.5	20.0	0.10		
>10	4.5	20.0	0.10		
6	0.5	20.0	0.20	-6	-2.6
8	1.0	20.0	0.20	1	-0.2
8	1.5	20.0	0.20	-5	0.3
8	2.0	20.0	0.20	1	-0.2
6	2.5	20.0	0.20	-6	1.3
8	3.5	20.0	0.20	-5	-0.3
8	3.5	20.0	0.20	1	-0.2
6	4.5	20.0	0.20	-6	-2.0
9	5.0	20.0	0.20	1	-0.1
4	0.5	20.0	0.40	-6	-3.2
4	0.5	20.0	0.40	1	-0.2
4	1.5	20.0	0.40	1	-0.2
4	2.5	20.0	0.40	-6	-2.5
4	2.5	20.0	0.40	1	-0.2
5	3.5	20.0	0.40	-5	-1.2
5	3.5	20.0	0.40	1	-0.2
4	4.5	20.0	0.40	-6	-2.4
5	5.0	20.0	0.40	1	-0.2
3	0.5	20.0	0.60	1	-0.2
3	1.5	20.0	0.60	1	-0.2
3	2.5	20.0	0.60	1	-0.2
3	3.5	20.0	0.60	1	-0.3
3	4.5	20.0	0.60	1	-0.3
2	0.5	20.0	0.80	1	-0.3
2	1.5	20.0	0.80	1	-0.3
2	2.5	20.0	0.80	1	-0.3
2	3.5	20.0	0.80	1	-0.3
2	4.5	20.0	0.80	1	-0.4
2	0.5	20.0	0.90	1	-0.3
2	1.5	20.0	0.90	1	-0.3
2	2.5	20.0	0.90	1	-0.3
2	3.5	20.0	0.90	1	-0.3
2	4.5	20.0	0.90	1	-0.3
3	0.5	20.0	0.95	2	0.7
2	1.5	20.0	0.95	1	-0.3
2	2.5	20.0	0.95	1	-0.3
2	3.5	20.0	0.95	1	-0.3
2	4.5	20.0	0.95	1	-0.3
4	0.5	20.0	0.98	3	1.5
3	1.5	20.0	0.98	2	0.7
3	2.5	20.0	0.98	2	0.7
3	3.5	20.0	0.98	2	0.7
3	4.5	20.0	0.98	2	0.7

first unst. m	nu	omega over sigm	alph	s	delt
4	0.0	20.0	0.99	4	-0.8
4	1.0	20.0	0.99	4	-0.8
4	1.5	20.0	0.99	4	-0.8
4	2.0	20.0	0.99	4	-0.8
4	2.5	20.0	0.99	4	-0.8
4	3.0	20.0	0.99	4	-0.8
4	3.5	20.0	0.99	4	-0.8
4	4.0	20.0	0.99	4	-0.8
4	4.5	20.0	0.99	4	-0.8
4	5.0	20.0	0.99	4	-0.8

>10	0.0	40.0	0.01		
>10	1.0	40.0	0.01		
>10	2.0	40.0	0.01		
>10	3.0	40.0	0.01		
>10	4.0	40.0	0.01		
>10	5.0	40.0	0.01		

>10	0.0	40.0	0.02		
>10	1.0	40.0	0.02		
>10	2.0	40.0	0.02		
>10	3.0	40.0	0.02		
>10	4.0	40.0	0.02		
>10	5.0	40.0	0.02		

8	0.0	40.0	0.05	0	-2.0
8	1.0	40.0	0.05	0	-2.0
8	2.0	40.0	0.05	0	-2.0
8	3.0	40.0	0.05	0	-2.0
9	4.0	40.0	0.05	0	-1.9
9	5.0	40.0	0.05	0	-1.9

4	0.0	40.0	0.10	0	-2.0
4	1.0	40.0	0.10	0	-2.0
5	1.5	40.0	0.10	-3	1.4
5	2.0	40.0	0.10	0	-1.9
5	2.5	40.0	0.10	0	-1.9
6	3.5	40.0	0.10	0	-1.6
6	4.5	40.0	0.10	0	-1.6

3	0.0	40.0	0.20	0	-1.8
3	1.0	40.0	0.20	0	-1.8
3	1.5	40.0	0.20	-3	2.8
3	2.0	40.0	0.20	0	-1.8
4	3.0	40.0	0.20	0	-1.6
4	4.0	40.0	0.20	0	-1.6
4	5.0	40.0	0.20	0	-1.6

2	0.0	40.0	0.40	0	-1.8
0	0.5	40.0	0.40	2	2.4
2	1.5	40.0	0.40	-3	-3.6
2	1.5	40.0	0.40	0	-1.8
2	2.5	40.0	0.40	0	-1.8
3	3.5	40.0	0.40	0	-1.5
3	4.0	40.0	0.40	0	-1.5
2	4.5	40.0	0.40	-3	8.5
3	5.0	40.0	0.40	1	0.6

2	0.0	40.0	0.60	0	-1.8
0	0.5	40.0	0.60	-2	-2.4
2	1.0	40.0	0.60	0	-1.8
2	1.5	40.0	0.60	-3	-3.4
2	1.5	40.0	0.60	0	-1.8

first unst. m	nu	omega over sigm	alph	s	delt
4	0.5	20.0	0.99	4	-0.8
4	1.5	20.0	0.99	3	1.5
4	2.0	20.0	0.99	3	1.5
4	2.5	20.0	0.99	3	1.5
4	3.0	20.0	0.99	3	1.5
4	3.5	20.0	0.99	3	1.5
4	4.0	20.0	0.99	3	1.5
4	4.5	20.0	0.99	3	1.5
4	5.0	20.0	0.99	3	1.5

>10	0.5	40.0	0.01		
>10	1.5	40.0	0.01		
>10	2.5	40.0	0.01		
>10	3.5	40.0	0.01		
>10	4.5	40.0	0.01		

>10	0.5	40.0	0.02		
>10	1.5	40.0	0.02		
>10	2.5	40.0	0.02		
>10	3.5	40.0	0.02		
>10	4.5	40.0	0.02		

8	0.5	40.0	0.05	0	-2.0
8	1.5	40.0	0.05	0	-2.0
8	2.5	40.0	0.05	0	-2.0
9	3.5	40.0	0.05	0	-1.9
9	4.5	40.0	0.05	0	-1.9

4	0.5	40.0	0.10	0	-2.0
5	1.5	40.0	0.10	-3	-2.9
5	1.5	40.0	0.10	0	-1.9
5	2.5	40.0	0.10	-3	1.5
5	3.5	40.0	0.10	0	-1.9
5	4.0	40.0	0.10	-3	1.5
6	5.0	40.0	0.10	0	-1.6

3	0.5	40.0	0.20	0	-1.8
3	1.5	40.0	0.20	-3	-3.4
3	1.5	40.0	0.20	0	-1.8
3	2.5	40.0	0.20	0	-1.8
4	3.5	40.0	0.20	0	-1.6
4	4.5	40.0	0.20	0	-1.6

0	0.5	40.0	0.40	-2	-2.4
2	1.0	40.0	0.40	0	-1.8
2	1.5	40.0	0.40	-3	3.0
2	2.0	40.0	0.40	0	-1.8
2	3.0	40.0	0.40	0	-1.8
3	3.5	40.0	0.40	1	0.7
3	4.0	40.0	0.40	1	0.7
3	5.0	40.0	0.40	0	-1.5

2	0.0	40.0	0.60	1	0.7
0	0.5	40.0	0.60	2	2.4
2	1.0	40.0	0.60	1	0.7
2	1.5	40.0	0.60	-3	2.8
2	1.5	40.0	0.60	1	0.7

first unst. m	nu	omega over sigm	alph	s	delt
2	2.0	40.0	0.60	0	-1.8
2	2.5	40.0	0.60	0	-1.8
2	3.0	40.0	0.60	0	-1.8
2	3.5	40.0	0.60	0	-1.8
2	4.0	40.0	0.60	0	-1.8
2	4.5	40.0	0.60	0	-1.8
2	5.0	40.0	0.60	0	-1.8
2	0.0	40.0	0.80	1	0.8
2	1.0	40.0	0.80	1	0.8
2	2.0	40.0	0.80	1	0.8
2	3.0	40.0	0.80	1	0.8
2	4.0	40.0	0.80	1	0.7
2	5.0	40.0	0.80	1	0.7
2	0.0	40.0	0.90	1	0.7
2	1.0	40.0	0.90	1	0.7
2	2.0	40.0	0.90	1	0.7
2	3.0	40.0	0.90	1	0.7
2	4.0	40.0	0.90	1	0.7
2	5.0	40.0	0.90	1	0.7
2	0.0	40.0	0.95	1	0.7
2	1.0	40.0	0.95	1	0.7
2	2.0	40.0	0.95	1	0.7
2	3.0	40.0	0.95	1	0.7
2	4.0	40.0	0.95	1	0.7
2	5.0	40.0	0.95	1	0.7
2	0.0	40.0	0.98	1	0.7
2	1.0	40.0	0.98	1	0.7
2	2.0	40.0	0.98	1	0.7
2	3.0	40.0	0.98	1	0.7
2	4.0	40.0	0.98	1	0.7
2	5.0	40.0	0.98	1	0.7
2	0.0	40.0	0.99	2	-0.8
2	1.0	40.0	0.99	2	-0.8
2	2.0	40.0	0.99	1	0.7
2	2.5	40.0	0.99	1	0.7
2	3.0	40.0	0.99	1	0.7
2	3.5	40.0	0.99	1	0.7
2	4.0	40.0	0.99	1	0.7
2	4.5	40.0	0.99	1	0.7
2	5.0	40.0	0.99	1	0.7

first unst. m	nu	omega over sigm	alph	s	delt
2	2.0	40.0	0.60	1	0.7
2	2.5	40.0	0.60	1	0.7
2	3.0	40.0	0.60	1	0.6
2	3.5	40.0	0.60	1	0.6
2	4.0	40.0	0.60	1	0.6
2	4.5	40.0	0.60	1	0.6
2	5.0	40.0	0.60	1	0.6
2	0.5	40.0	0.80	1	0.8
2	1.5	40.0	0.80	1	0.8
2	2.5	40.0	0.80	1	0.8
2	3.5	40.0	0.80	1	0.8
2	4.5	40.0	0.80	1	0.7
2	0.5	40.0	0.90	1	0.7
2	1.5	40.0	0.90	1	0.7
2	2.5	40.0	0.90	1	0.7
2	3.5	40.0	0.90	1	0.7
2	4.5	40.0	0.90	1	0.7
2	0.5	40.0	0.95	1	0.7
2	1.5	40.0	0.95	1	0.7
2	2.5	40.0	0.95	1	0.7
2	3.5	40.0	0.95	1	0.7
2	4.5	40.0	0.95	1	0.7
2	0.5	40.0	0.98	1	0.7
2	1.5	40.0	0.98	1	0.7
2	2.5	40.0	0.98	1	0.7
2	3.5	40.0	0.98	1	0.7
2	4.5	40.0	0.98	1	0.7
2	0.5	40.0	0.99	2	-0.8
2	1.5	40.0	0.99	2	-0.8
2	2.0	40.0	0.99	2	-0.8
2	2.5	40.0	0.99	2	-0.8
2	3.0	40.0	0.99	2	-0.8
2	3.5	40.0	0.99	2	-0.8
2	4.0	40.0	0.99	2	-0.8
2	4.5	40.0	0.99	2	-0.8
2	5.0	40.0	0.99	2	-0.8

B Appendix

In Appendix A we observed the weak dependence of ν on instability. We exploit this with a modified program STAB5A which evaluates stability at various values of $\Delta\omega_e/\Delta\omega_i$ and ZT_e/T_i , but at $\nu = 0$ only.

```

program stab5a
parameter(nmax=10001,nmax2=5001,mmax=10)
complex w,wi(nmax),we_center(nmax),we_left(nmax),we_right(nmax)
complex eps_i(nmax),eps_e(nmax),sum_i(nmax)
real alparm(12),dwparm(4),anuparm(11),dweparm(3),c5parm(4)
data alparm/.01,.02,.05,.1,.2,.4,.6,.8,.9,.95,.98,.99/
data dwparm/5.,10.,20.,40./
data dweparm/-3000.,-4000.,-6000./
data c5parm/.1,.2,.5,1./
open(1,file='stab5a.dat',access='sequential')

c
c
c   set parameters

n1=0
anu=0.
scale=.1
do 300 ic5=1,4
c5=c5parm(ic5)
do 300 idwe=1,3
dwe_by_dwi=dweparm(idwe)
do 300 idw=1,4
dwi_by_si=dwparm(idw)
do 300 ialp=1,12
alpha_i=alparm(ialp)

c
c
c   derived parameters

alpha_e=-c5*alpha_i*(1.0-alpha_i)/dwe_by_dwi
se_by_si=sqrt(-c5*dwe_by_dwi)
c1=alpha_e*dwe_by_dwi*dwi_by_si/se_by_si
c2=scale/se_by_si
c3=alpha_i*dwi_by_si
c4=dwe_by_dwi*dwi_by_si/se_by_si
n2=nint(dwi_by_si/scale)
sqrtpi=sqrt(3.1415926)

c
c
c   form w functions

do 10 n=1,nmax
w1(n)=w((n-nmax2)*scale)
we_center(n)=w((n-nmax2)*c2)
we_left(n)=w((n-nmax2)*c2-c4)
we_right(n)=w((n-nmax2)*c2+c4)
10 continue

c
c
c   form ion sum for m=0, anu=0

do 15 n=1,nmax
15 sum_i(n)=wi(n)

c
c
c   form dielectric constants

do 100 m=0,mmax
do 30 n=1,nmax

c
c   eps_i(n)=
& c5*7-1.0+cmplx(0.0,1.0)*(m*c3-(n-nmax2)*scale)*sqrtpi*sum_i(n)

c
nshift=n+m*n1
if(nshift.gt.nmax) nshift=nmax

c
c   eps_e(n)=-1.0+cmplx(0.0,1.0)*(m*c1-(nshift-nmax2)*c2)*sqrtpi*
& (1.0-(m+2.0*anu)*alpha_e)*
& (we_center(nshift)+alpha_e*(m+anu)*we_left(nshift)+
& alpha_e*anu*we_right(nshift))

c
nshift=n-n2

```

```

        if(nshift.ge.1.and.nshift.le.nmax) then
            wi(n)=alpha_i*sum_i(nshift)+(1.0-alpha_i)*sum_i(n)
        else
            wi(n)=sum_i(n)
        endif
        continue
30      c
        do 47 n=1,nmax
47      sum_i(n)=wi(n)
        c
        call eval(eps_e,eps_i,nmax,nmax2,m,anu,dwi_by_si,alpha_i,scale,iflag,
        & dwe_by_dwi,c5)
        if(iflag.eq.1) go to 300
100      continue
        write(1,1000) anu,dwi_by_si,alpha_i,dwe_by_dwi,c5
        write(*,1000) anu,dwi_by_si,alpha_i,dwe_by_dwi,c5
1000     format(1x,'***',1x,f4.1,1x,f5.1,1x,f4.2,1x,f7.0,1x,f4.2)
300     continue
        close(1)
        end

        complex function w(x)
        complex w
        pi=3.1415926
        fact=1./sqrt(pi)
        if(x.lt.10.) then
            parm=exp(-x*x)
        else
            parm=0.
        endif
        if(abs(x).lt.6.) go to 600
        aml=1.
        a0=0.
        bml=0.
        a=1.
        n=1
        fo=0.
60      an=x*a0+a*aml
        bn=x+a*bml
        fn=an/bn
        if(abs(1-fo/fn).lt.1.e-7) go to 500
        fo=fn
        n=n+1
        a=-.5*(n-1)
        aml=a0/bn
        a0=fn
        bml=1./bn
        go to 60
500     w=cmlpx(parm,fn*fact)
        return
600     sum=1.
        n=1
        term=x*x
        sumn=sum+term/(2*n+1)
610     term=term*x*x/(n+1)
        n=n+1
        if(abs(term/sumn).lt.1.e-7) go to 1000
        sum=sumn
        go to 610
1000    w=parm*cmlpx(1.,x*2.*fact*sumn)
        end

        subroutine eval(eps_e,eps_i,nmax,nmax2,m,anu,dwi_by_si,alpha_i,
        & scale,iflaq,dwe_by_dwi,c5)

```

```

complex eps_i(nmax),eps_e(nmax),ctempl,ctemp2
iflag=0
do 10 i=1,nmax-1
phase=aimag(eps_e(i))/real(eps_e(i))
if(abs(phase).lt.1.e-6) go to 10
ctempl=eps_i(i)/eps_e(i)
ctemp2=eps_i(i+1)/eps_e(i+1)
a1=sign(1.,aimag(ctempl))*sign(1.,aimag(ctemp2))
a2=.5*real(ctempl)+.5*real(ctemp2)
if(a1.gt.0.or.a2.gt.-1.0) go to 10
iflag=1
is=nint(scale*(i+.5-nmax2)/dwi_by_si)
delt=scale*(i+.5-nmax2)-is*dwi_by_si
write(1,100) m,anu,dwi_by_si,alpha_i,dwe_by_dwi,c5,is,delt
write(*,100) m,anu,dwi_by_si,alpha_i,dwe_by_dwi,c5,is,delt
format(1x,i3,1x,f4.1,1x,f5.1,1x,f4.2,1x,f7.0,1x,f4.2,i3,1x,f5.1)
continue
end
100
10

```

first unst.	nu	omega over sigm	alph	omega e over omega i	Te 2 over Ti	s	delt
>10	0.0	5.0	0.01	-3000.	0.10		
>10	0.0	5.0	0.02	-3000.	0.10		
>10	0.0	5.0	0.05	-3000.	0.10		
>10	0.0	5.0	0.10	-3000.	0.10		
>10	0.0	5.0	0.20	-3000.	0.10		
>10	0.0	5.0	0.40	-3000.	0.10		
>10	0.0	5.0	0.60	-3000.	0.10		
>10	0.0	5.0	0.80	-3000.	0.10		
>10	0.0	5.0	0.90	-3000.	0.10		
10	0.0	5.0	0.95	-3000.	0.10	9	1.7
10	0.0	5.0	0.98	-3000.	0.10	10	-1.0
10	0.0	5.0	0.99	-3000.	0.10	10	-0.5
>10	0.0	10.0	0.01	-3000.	0.10		
>10	0.0	10.0	0.02	-3000.	0.10		
>10	0.0	10.0	0.05	-3000.	0.10		
>10	0.0	10.0	0.10	-3000.	0.10		
>10	0.0	10.0	0.20	-3000.	0.10		
>10	0.0	10.0	0.40	-3000.	0.10		
8	0.0	10.0	0.60	-3000.	0.10	4	1.8
6	0.0	10.0	0.80	-3000.	0.10	4	1.7
5	0.0	10.0	0.90	-3000.	0.10	4	1.6
5	0.0	10.0	0.95	-3000.	0.10	4	1.7
5	0.0	10.0	0.98	-3000.	0.10	5	-1.0
5	0.0	10.0	0.99	-3000.	0.10	5	-0.5
>10	0.0	20.0	0.01	-3000.	0.10		
>10	0.0	20.0	0.02	-3000.	0.10		
>10	0.0	20.0	0.05	-3000.	0.10		
>10	0.0	20.0	0.10	-3000.	0.10		
8	0.0	20.0	0.20	-3000.	0.10	1	0.4
4	0.0	20.0	0.40	-3000.	0.10	1	0.4
3	0.0	20.0	0.60	-3000.	0.10	1	0.4
2	0.0	20.0	0.80	-3000.	0.10	1	0.3
3	0.0	20.0	0.90	-3000.	0.10	2	1.8
3	0.0	20.0	0.95	-3000.	0.10	2	1.8
3	0.0	20.0	0.98	-3000.	0.10	3	-1.2
3	0.0	20.0	0.99	-3000.	0.10	3	-0.6
>10	0.0	40.0	0.01	-3000.	0.10		
>10	0.0	40.0	0.02	-3000.	0.10		
>10	0.0	40.0	0.05	-3000.	0.10		
9	0.0	40.0	0.10	-3000.	0.10	0	-1.6
6	0.0	40.0	0.20	-3000.	0.10	0	-1.5
6	0.0	40.0	0.20	-3000.	0.10	1	1.9
3	0.0	40.0	0.40	-3000.	0.10	1	1.8
2	0.0	40.0	0.60	-3000.	0.10	1	1.8
2	0.0	40.0	0.80	-3000.	0.10	1	1.9
2	0.0	40.0	0.90	-3000.	0.10	1	1.9
2	0.0	40.0	0.95	-3000.	0.10	1	1.8
2	0.0	40.0	0.98	-3000.	0.10	2	-4.0
>10	0.0	40.0	0.99	-3000.	0.10		
>10	0.0	5.0	0.01	-4000.	0.10		
>10	0.0	5.0	0.02	-4000.	0.10		
>10	0.0	5.0	0.05	-4000.	0.10		
>10	0.0	5.0	0.10	-4000.	0.10		
>10	0.0	5.0	0.20	-4000.	0.10		
>10	0.0	5.0	0.40	-4000.	0.10		
>10	0.0	5.0	0.60	-4000.	0.10		
>10	0.0	5.0	0.80	-4000.	0.10		
>10	0.0	5.0	0.90	-4000.	0.10		
>10	0.0	5.0	0.95	-4000.	0.10		
>10	0.0	5.0	0.98	-4000.	0.10		
>10	0.0	5.0	0.99	-4000.	0.10		
>10	0.0	10.0	0.01	-4000.	0.10		
>10	0.0	10.0	0.02	-4000.	0.10		

first unst.	nu	omega over	alph	omega e over	Te 2 over	s	delt
m		sigm		omega i	Ti		
>10	0.0	10.0	0.05	-4000.	0.10		
>10	0.0	10.0	0.10	-4000.	0.10		
>10	0.0	10.0	0.20	-4000.	0.10		
>10	0.0	10.0	0.40	-4000.	0.10		
10	0.0	10.0	0.60	-4000.	0.10	5	2.0
6	0.0	10.0	0.80	-4000.	0.10	4	1.3
5	0.0	10.0	0.90	-4000.	0.10	4	1.2
6	0.0	10.0	0.95	-4000.	0.10	5	1.9
6	0.0	10.0	0.98	-4000.	0.10	6	-1.2
6	0.0	10.0	0.99	-4000.	0.10	6	-0.6
>10	0.0	20.0	0.01	-4000.	0.10		
>10	0.0	20.0	0.02	-4000.	0.10		
>10	0.0	20.0	0.05	-4000.	0.10		
>10	0.0	20.0	0.10	-4000.	0.10		
9	0.0	20.0	0.20	-4000.	0.10	1	0.3
4	0.0	20.0	0.40	-4000.	0.10	1	0.2
3	0.0	20.0	0.60	-4000.	0.10	1	0.2
2	0.0	20.0	0.80	-4000.	0.10	1	0.1
3	0.0	20.0	0.90	-4000.	0.10	2	1.4
3	0.0	20.0	0.95	-4000.	0.10	2	1.4
3	0.0	20.0	0.98	-4000.	0.10	3	-1.2
3	0.0	20.0	0.99	-4000.	0.10	3	-0.6
>10	0.0	40.0	0.01	-4000.	0.10		
>10	0.0	40.0	0.02	-4000.	0.10		
>10	0.0	40.0	0.05	-4000.	0.10		
10	0.0	40.0	0.10	-4000.	0.10	0	-1.6
6	0.0	40.0	0.20	-4000.	0.10	0	-1.6
6	0.0	40.0	0.20	-4000.	0.10	1	1.5
3	0.0	40.0	0.40	-4000.	0.10	1	1.5
2	0.0	40.0	0.60	-4000.	0.10	1	1.4
2	0.0	40.0	0.80	-4000.	0.10	1	1.5
2	0.0	40.0	0.90	-4000.	0.10	1	1.5
2	0.0	40.0	0.95	-4000.	0.10	1	1.5
2	0.0	40.0	0.98	-4000.	0.10	2	-3.3
2	0.0	40.0	0.98	-4000.	0.10	2	-1.6
2	0.0	40.0	0.99	-4000.	0.10	2	-0.8
>10	0.0	5.0	0.01	-6000.	0.10		
>10	0.0	5.0	0.02	-6000.	0.10		
>10	0.0	5.0	0.05	-6000.	0.10		
>10	0.0	5.0	0.10	-6000.	0.10		
>10	0.0	5.0	0.20	-6000.	0.10		
>10	0.0	5.0	0.40	-6000.	0.10		
>10	0.0	5.0	0.60	-6000.	0.10		
>10	0.0	5.0	0.80	-6000.	0.10		
>10	0.0	5.0	0.90	-6000.	0.10		
>10	0.0	5.0	0.95	-6000.	0.10		
>10	0.0	5.0	0.98	-6000.	0.10		
>10	0.0	5.0	0.99	-6000.	0.10		
>10	0.0	10.0	0.01	-6000.	0.10		
>10	0.0	10.0	0.02	-6000.	0.10		
>10	0.0	10.0	0.05	-6000.	0.10		
>10	0.0	10.0	0.10	-6000.	0.10		
>10	0.0	10.0	0.20	-6000.	0.10		
>10	0.0	10.0	0.40	-6000.	0.10		
>10	0.0	10.0	0.60	-6000.	0.10		
8	0.0	10.0	0.80	-6000.	0.10	6	1.7
7	0.0	10.0	0.90	-6000.	0.10	6	1.6
7	0.0	10.0	0.95	-6000.	0.10	6	1.8
7	0.0	10.0	0.98	-6000.	0.10	7	-1.5
7	0.0	10.0	0.99	-6000.	0.10	7	-0.7
>10	0.0	20.0	0.01	-6000.	0.10		
>10	0.0	20.0	0.02	-6000.	0.10		
>10	0.0	20.0	0.05	-6000.	0.10		

first unst.	nu	omega over	alpha	omega e over	Te 2 over	s	delt
m		sigm		omega i	Ti		
>10	0.0	20.0	0.10	-6000.	0.10		
9	0.0	20.0	0.20	-6000.	0.10	1	0.0
5	0.0	20.0	0.40	-6000.	0.10	1	0.0
3	0.0	20.0	0.60	-6000.	0.10	1	-0.1
3	0.0	20.0	0.80	-6000.	0.10	2	0.9
3	0.0	20.0	0.90	-6000.	0.10	2	0.9
4	0.0	20.0	0.95	-6000.	0.10	3	1.9
4	0.0	20.0	0.98	-6000.	0.10	4	-1.6
4	0.0	20.0	0.99	-6000.	0.10	4	-0.8
>10	0.0	40.0	0.01	-6000.	0.10		
>10	0.0	40.0	0.02	-6000.	0.10		
>10	0.0	40.0	0.05	-6000.	0.10		
>10	0.0	40.0	0.10	-6000.	0.10		
6	0.0	40.0	0.20	-6000.	0.10	1	1.0
3	0.0	40.0	0.40	-6000.	0.10	1	1.0
2	0.0	40.0	0.60	-6000.	0.10	1	1.0
2	0.0	40.0	0.80	-6000.	0.10	1	1.0
2	0.0	40.0	0.90	-6000.	0.10	1	1.0
2	0.0	40.0	0.95	-6000.	0.10	1	1.0
2	0.0	40.0	0.98	-6000.	0.10	2	-1.6
2	0.0	40.0	0.99	-6000.	0.10	2	-0.8
>10	0.0	5.0	0.01	-3000.	0.20		
>10	0.0	5.0	0.02	-3000.	0.20		
>10	0.0	5.0	0.05	-3000.	0.20		
>10	0.0	5.0	0.10	-3000.	0.20		
>10	0.0	5.0	0.20	-3000.	0.20		
>10	0.0	5.0	0.40	-3000.	0.20		
>10	0.0	5.0	0.60	-3000.	0.20		
>10	0.0	5.0	0.80	-3000.	0.20		
>10	0.0	5.0	0.90	-3000.	0.20		
10	0.0	5.0	0.95	-3000.	0.20	9	0.8
>10	0.0	5.0	0.98	-3000.	0.20		
>10	0.0	5.0	0.99	-3000.	0.20		
>10	0.0	10.0	0.01	-3000.	0.20		
>10	0.0	10.0	0.02	-3000.	0.20		
>10	0.0	10.0	0.05	-3000.	0.20		
>10	0.0	10.0	0.10	-3000.	0.20		
>10	0.0	10.0	0.20	-3000.	0.20		
8	0.0	10.0	0.40	-3000.	0.20	2	0.0
7	0.0	10.0	0.60	-3000.	0.20	3	0.5
5	0.0	10.0	0.80	-3000.	0.20	3	0.4
4	0.0	10.0	0.90	-3000.	0.20	3	0.3
5	0.0	10.0	0.95	-3000.	0.20	4	0.9
6	0.0	10.0	0.98	-3000.	0.20	5	1.4
6	0.0	10.0	0.99	-3000.	0.20	6	-0.6
>10	0.0	20.0	0.01	-3000.	0.20		
>10	0.0	20.0	0.02	-3000.	0.20		
>10	0.0	20.0	0.05	-3000.	0.20		
>10	0.0	20.0	0.10	-3000.	0.20		
7	0.0	20.0	0.20	-3000.	0.20	1	0.0
4	0.0	20.0	0.40	-3000.	0.20	1	0.0
3	0.0	20.0	0.60	-3000.	0.20	1	0.0
2	0.0	20.0	0.80	-3000.	0.20	1	-0.1
2	0.0	20.0	0.90	-3000.	0.20	1	-0.1
3	0.0	20.0	0.95	-3000.	0.20	2	1.0
4	0.0	20.0	0.98	-3000.	0.20	3	1.9
4	0.0	20.0	0.98	-3000.	0.20	4	-2.5
4	0.0	20.0	0.98	-3000.	0.20	4	-1.6
3	0.0	20.0	0.99	-3000.	0.20	3	-0.6
>10	0.0	40.0	0.01	-3000.	0.20		
>10	0.0	40.0	0.02	-3000.	0.20		
7	0.0	40.0	0.05	-3000.	0.20	0	-2.2
4	0.0	40.0	0.10	-3000.	0.20	0	-2.0

first unst. m	nu	omega over sigm	alpha	omega e over omega i	Te 2 over Ti	s	delt
3	0.0	40.0	0.20	-3000.	0.20	0	-1.6
2	0.0	40.0	0.40	-3000.	0.20	0	-1.6
2	0.0	40.0	0.60	-3000.	0.20	0	-1.6
2	0.0	40.0	0.60	-3000.	0.20	1	1.0
2	0.0	40.0	0.80	-3000.	0.20	1	1.1
2	0.0	40.0	0.90	-3000.	0.20	1	1.0
2	0.0	40.0	0.95	-3000.	0.20	1	1.0
2	0.0	40.0	0.98	-3000.	0.20	1	1.0
2	0.0	40.0	0.98	-3000.	0.20	2	-2.5
2	0.0	40.0	0.98	-3000.	0.20	2	-1.6
2	0.0	40.0	0.99	-3000.	0.20	2	-0.8
>10	0.0	5.0	0.01	-4000.	0.20		
>10	0.0	5.0	0.02	-4000.	0.20		
>10	0.0	5.0	0.05	-4000.	0.20		
>10	0.0	5.0	0.10	-4000.	0.20		
>10	0.0	5.0	0.20	-4000.	0.20		
>10	0.0	5.0	0.40	-4000.	0.20		
>10	0.0	5.0	0.60	-4000.	0.20		
>10	0.0	5.0	0.80	-4000.	0.20		
>10	0.0	5.0	0.90	-4000.	0.20		
>10	0.0	5.0	0.95	-4000.	0.20		
>10	0.0	5.0	0.98	-4000.	0.20		
>10	0.0	5.0	0.99	-4000.	0.20		
>10	0.0	10.0	0.01	-4000.	0.20		
>10	0.0	10.0	0.02	-4000.	0.20		
>10	0.0	10.0	0.05	-4000.	0.20		
>10	0.0	10.0	0.10	-4000.	0.20		
>10	0.0	10.0	0.20	-4000.	0.20		
10	0.0	10.0	0.40	-4000.	0.20	3	0.3
7	0.0	10.0	0.60	-4000.	0.20	3	0.3
5	0.0	10.0	0.80	-4000.	0.20	3	0.2
5	0.0	10.0	0.90	-4000.	0.20	4	0.5
5	0.0	10.0	0.95	-4000.	0.20	4	0.6
7	0.0	10.0	0.98	-4000.	0.20	6	1.5
7	0.0	10.0	0.99	-4000.	0.20	7	-0.7
>10	0.0	20.0	0.01	-4000.	0.20		
>10	0.0	20.0	0.02	-4000.	0.20		
>10	0.0	20.0	0.05	-4000.	0.20		
>10	0.0	20.0	0.10	-4000.	0.20		
7	0.0	20.0	0.20	-4000.	0.20	1	-0.2
4	0.0	20.0	0.40	-4000.	0.20	1	-0.2
3	0.0	20.0	0.60	-4000.	0.20	1	-0.2
2	0.0	20.0	0.80	-4000.	0.20	1	-0.3
2	0.0	20.0	0.90	-4000.	0.20	1	-0.3
3	0.0	20.0	0.95	-4000.	0.20	2	0.7
4	0.0	20.0	0.98	-4000.	0.20	3	1.5
4	0.0	20.0	0.99	-4000.	0.20	4	-0.8
>10	0.0	40.0	0.01	-4000.	0.20		
>10	0.0	40.0	0.02	-4000.	0.20		
8	0.0	40.0	0.05	-4000.	0.20	0	-2.0
4	0.0	40.0	0.10	-4000.	0.20	0	-2.0
3	0.0	40.0	0.20	-4000.	0.20	0	-1.8
2	0.0	40.0	0.40	-4000.	0.20	0	-1.8
2	0.0	40.0	0.60	-4000.	0.20	0	-1.8
2	0.0	40.0	0.60	-4000.	0.20	1	0.7
2	0.0	40.0	0.80	-4000.	0.20	1	0.8
2	0.0	40.0	0.90	-4000.	0.20	1	0.7
2	0.0	40.0	0.95	-4000.	0.20	1	0.7
2	0.0	40.0	0.98	-4000.	0.20	1	0.7
2	0.0	40.0	0.99	-4000.	0.20	2	-0.8
>10	0.0	5.0	0.01	-6000.	0.20		
>10	0.0	5.0	0.02	-6000.	0.20		
>10	0.0	5.0	0.05	-6000.	0.20		

first	unst.	nu	omga	alph	omega e	Te 2	s	delt
m			over	over	over	over		
			sigm		omega i	Ti		
>10	0.0		5.0	0.10	-6000.	0.20		
>10	0.0		5.0	0.20	-6000.	0.20		
>10	0.0		5.0	0.40	-6000.	0.20		
>10	0.0		5.0	0.60	-6000.	0.20		
>10	0.0		5.0	0.80	-6000.	0.20		
>10	0.0		5.0	0.90	-6000.	0.20		
>10	0.0		5.0	0.95	-6000.	0.20		
>10	0.0		5.0	0.98	-6000.	0.20		
>10	0.0		5.0	0.99	-6000.	0.20		
>10	0.0		10.0	0.01	-6000.	0.20		
>10	0.0		10.0	0.02	-6000.	0.20		
>10	0.0		10.0	0.05	-6000.	0.20		
>10	0.0		10.0	0.10	-6000.	0.20		
>10	0.0		10.0	0.20	-6000.	0.20		
>10	0.0		10.0	0.40	-6000.	0.20		
7	0.0		10.0	0.60	-6000.	0.20	3	0.0
6	0.0		10.0	0.80	-6000.	0.20	4	0.3
5	0.0		10.0	0.90	-6000.	0.20	4	0.2
6	0.0		10.0	0.95	-6000.	0.20	5	0.6
8	0.0		10.0	0.98	-6000.	0.20	7	1.3
8	0.0		10.0	0.99	-6000.	0.20	8	-0.9
>10	0.0		20.0	0.01	-6000.	0.20		
>10	0.0		20.0	0.02	-6000.	0.20		
>10	0.0		20.0	0.05	-6000.	0.20		
>10	0.0		20.0	0.10	-6000.	0.20		
8	0.0		20.0	0.20	-6000.	0.20	1	-0.4
4	0.0		20.0	0.40	-6000.	0.20	1	-0.4
3	0.0		20.0	0.60	-6000.	0.20	1	-0.4
2	0.0		20.0	0.80	-6000.	0.20	1	-0.5
2	0.0		20.0	0.90	-6000.	0.20	1	-0.5
3	0.0		20.0	0.95	-6000.	0.20	2	0.3
4	0.0		20.0	0.98	-6000.	0.20	3	1.1
4	0.0		20.0	0.99	-6000.	0.20	4	-0.9
>10	0.0		40.0	0.01	-6000.	0.20		
>10	0.0		40.0	0.02	-6000.	0.20		
8	0.0		40.0	0.05	-6000.	0.20	0	-2.0
4	0.0		40.0	0.10	-6000.	0.20	0	-2.2
3	0.0		40.0	0.20	-6000.	0.20	0	-1.9
2	0.0		40.0	0.40	-6000.	0.20	0	-1.9
2	0.0		40.0	0.60	-6000.	0.20	0	-1.9
2	0.0		40.0	0.60	-6000.	0.20	1	0.4
2	0.0		40.0	0.80	-6000.	0.20	1	0.4
2	0.0		40.0	0.90	-6000.	0.20	1	0.4
2	0.0		40.0	0.95	-6000.	0.20	1	0.4
3	0.0		40.0	0.98	-6000.	0.20	2	1.8
2	0.0		40.0	0.99	-6000.	0.20	2	-0.9
>10	0.0		5.0	0.01	-3000.	0.50		
>10	0.0		5.0	0.02	-3000.	0.50		
>10	0.0		5.0	0.05	-3000.	0.50		
>10	0.0		5.0	0.10	-3000.	0.50		
>10	0.0		5.0	0.20	-3000.	0.50		
>10	0.0		5.0	0.40	-3000.	0.50		
>10	0.0		5.0	0.60	-3000.	0.50		
9	0.0		5.0	0.80	-3000.	0.50	6	-0.3
9	0.0		5.0	0.90	-3000.	0.50	7	-0.1
8	0.0		5.0	0.95	-3000.	0.50	7	-0.3
10	0.0		5.0	0.98	-3000.	0.50	9	0.1
>10	0.0		5.0	0.99	-3000.	0.50		
>10	0.0		10.0	0.01	-3000.	0.50		
>10	0.0		10.0	0.02	-3000.	0.50		
>10	0.0		10.0	0.05	-3000.	0.50		
8	0.0		10.0	0.10	-3000.	0.50	0	-2.0
5	0.0		10.0	0.20	-3000.	0.50	0	-1.9

first unst. m	nu	omega over sigm	alph	omega e over omega i	Te 2 over Ti	s	delt
4	0.0	10.0	0.40	-3000.	0.50	1	-1.1
3	0.0	10.0	0.60	-3000.	0.50	1	-1.1
3	0.0	10.0	0.80	-3000.	0.50	2	-0.8
3	0.0	10.0	0.90	-3000.	0.50	2	-0.7
4	0.0	10.0	0.95	-3000.	0.50	3	-0.3
5	0.0	10.0	0.98	-3000.	0.50	4	0.1
7	0.0	10.0	0.99	-3000.	0.50	6	0.8
>10	0.0	20.0	0.01	-3000.	0.50		
>10	0.0	20.0	0.02	-3000.	0.50		
6	0.0	20.0	0.05	-3000.	0.50	0	-2.4
4	0.0	20.0	0.10	-3000.	0.50	0	-2.0
2	0.0	20.0	0.20	-3000.	0.50	0	-2.0
2	0.0	20.0	0.40	-3000.	0.50	0	-1.8
2	0.0	20.0	0.60	-3000.	0.50	0	-1.8
2	0.0	20.0	0.60	-3000.	0.50	1	-0.7
2	0.0	20.0	0.80	-3000.	0.50	1	-0.5
2	0.0	20.0	0.90	-3000.	0.50	1	-0.6
2	0.0	20.0	0.95	-3000.	0.50	1	-0.6
3	0.0	20.0	0.98	-3000.	0.50	2	0.2
4	0.0	20.0	0.99	-3000.	0.50	3	0.8
4	0.0	20.0	0.99	-3000.	0.50	4	-1.2
>10	0.0	40.0	0.01	-3000.	0.50		
8	0.0	40.0	0.02	-3000.	0.50	0	-2.2
3	0.0	40.0	0.05	-3000.	0.50	0	-2.3
2	0.0	40.0	0.10	-3000.	0.50	0	-2.0
1	0.0	40.0	0.20	-3000.	0.50	0	-2.0
1	0.0	40.0	0.40	-3000.	0.50	0	-1.8
1	0.0	40.0	0.60	-3000.	0.50	0	-1.8
1	0.0	40.0	0.80	-3000.	0.50	0	-2.0
2	0.0	40.0	0.90	-3000.	0.50	1	0.3
2	0.0	40.0	0.95	-3000.	0.50	1	0.2
2	0.0	40.0	0.98	-3000.	0.50	1	0.2
2	0.0	40.0	0.99	-3000.	0.50	1	0.2
2	0.0	40.0	0.99	-3000.	0.50	2	-1.2
>10	0.0	5.0	0.01	-4000.	0.50		
>10	0.0	5.0	0.02	-4000.	0.50		
>10	0.0	5.0	0.05	-4000.	0.50		
>10	0.0	5.0	0.10	-4000.	0.50		
>10	0.0	5.0	0.20	-4000.	0.50		
>10	0.0	5.0	0.40	-4000.	0.50		
>10	0.0	5.0	0.60	-4000.	0.50		
>10	0.0	5.0	0.80	-4000.	0.50		
10	0.0	5.0	0.90	-4000.	0.50	8	-0.2
10	0.0	5.0	0.95	-4000.	0.50	9	-0.3
>10	0.0	5.0	0.98	-4000.	0.50		
>10	0.0	5.0	0.99	-4000.	0.50		
>10	0.0	10.0	0.01	-4000.	0.50		
>10	0.0	10.0	0.02	-4000.	0.50		
>10	0.0	10.0	0.05	-4000.	0.50		
9	0.0	10.0	0.10	-4000.	0.50	0	-2.0
6	0.0	10.0	0.20	-4000.	0.50	0	-1.8
5	0.0	10.0	0.40	-4000.	0.50	1	-1.0
3	0.0	10.0	0.60	-4000.	0.50	1	-1.2
4	0.0	10.0	0.80	-4000.	0.50	2	-0.7
3	0.0	10.0	0.90	-4000.	0.50	2	-0.8
4	0.0	10.0	0.95	-4000.	0.50	3	-0.4
5	0.0	10.0	0.98	-4000.	0.50	4	-0.1
7	0.0	10.0	0.99	-4000.	0.50	6	0.5
>10	0.0	20.0	0.01	-4000.	0.50		
>10	0.0	20.0	0.02	-4000.	0.50		
6	0.0	20.0	0.05	-4000.	0.50	0	-2.4
4	0.0	20.0	0.10	-4000.	0.50	0	-2.0
2	0.0	20.0	0.20	-4000.	0.50	0	-2.2

first unst. m	nu	omega over sigm	alpha	omega e over omega i	Te Z over Ti	s	delt
2	0.0	20.0	0.40	-4000.	0.50	0	-1.8
2	0.0	20.0	0.60	-4000.	0.50	0	-1.8
2	0.0	20.0	0.60	-4000.	0.50	1	-0.8
2	0.0	20.0	0.80	-4000.	0.50	1	-0.7
2	0.0	20.0	0.90	-4000.	0.50	1	-0.7
2	0.0	20.0	0.95	-4000.	0.50	1	-0.7
3	0.0	20.0	0.98	-4000.	0.50	2	0.0
4	0.0	20.0	0.99	-4000.	0.50	3	0.6
>10	0.0	40.0	0.01	-4000.	0.50		
8	0.0	40.0	0.02	-4000.	0.50	0	-2.2
3	0.0	40.0	0.05	-4000.	0.50	0	-2.4
2	0.0	40.0	0.10	-4000.	0.50	0	-2.0
1	0.0	40.0	0.20	-4000.	0.50	0	-2.2
1	0.0	40.0	0.40	-4000.	0.50	0	-1.8
1	0.0	40.0	0.60	-4000.	0.50	0	-1.8
1	0.0	40.0	0.80	-4000.	0.50	0	-2.0
2	0.0	40.0	0.90	-4000.	0.50	1	0.1
2	0.0	40.0	0.95	-4000.	0.50	1	0.0
2	0.0	40.0	0.98	-4000.	0.50	1	0.0
3	0.0	40.0	0.99	-4000.	0.50	2	1.2
3	0.0	40.0	0.99	-4000.	0.50	3	-2.0
3	0.0	40.0	0.99	-4000.	0.50	3	-1.3
>10	0.0	5.0	0.01	-6000.	0.50		
>10	0.0	5.0	0.02	-6000.	0.50		
>10	0.0	5.0	0.05	-6000.	0.50		
>10	0.0	5.0	0.10	-6000.	0.50		
>10	0.0	5.0	0.20	-6000.	0.50		
>10	0.0	5.0	0.40	-6000.	0.50		
>10	0.0	5.0	0.60	-6000.	0.50		
>10	0.0	5.0	0.80	-6000.	0.50		
>10	0.0	5.0	0.90	-6000.	0.50		
>10	0.0	5.0	0.95	-6000.	0.50		
>10	0.0	5.0	0.98	-6000.	0.50		
>10	0.0	5.0	0.99	-6000.	0.50		
>10	0.0	10.0	0.01	-6000.	0.50		
>10	0.0	10.0	0.02	-6000.	0.50		
>10	0.0	10.0	0.05	-6000.	0.50		
10	0.0	10.0	0.10	-6000.	0.50	0	-2.0
6	0.0	10.0	0.20	-6000.	0.50	0	-1.9
5	0.0	10.0	0.40	-6000.	0.50	1	-1.2
3	0.0	10.0	0.60	-6000.	0.50	1	-1.3
4	0.0	10.0	0.80	-6000.	0.50	2	-0.8
3	0.0	10.0	0.90	-6000.	0.50	2	-1.0
4	0.0	10.0	0.95	-6000.	0.50	3	-0.6
6	0.0	10.0	0.98	-6000.	0.50	5	-0.1
8	0.0	10.0	0.99	-6000.	0.50	7	0.4
>10	0.0	20.0	0.01	-6000.	0.50		
>10	0.0	20.0	0.02	-6000.	0.50		
7	0.0	20.0	0.05	-6000.	0.50	0	-2.2
4	0.0	20.0	0.10	-6000.	0.50	0	-2.2
2	0.0	20.0	0.20	-6000.	0.50	0	-2.3
2	0.0	20.0	0.40	-6000.	0.50	0	-1.9
2	0.0	20.0	0.60	-6000.	0.50	0	-1.9
2	0.0	20.0	0.60	-6000.	0.50	1	-1.0
2	0.0	20.0	0.80	-6000.	0.50	1	-0.8
2	0.0	20.0	0.90	-6000.	0.50	1	-0.9
2	0.0	20.0	0.95	-6000.	0.50	1	-0.9
3	0.0	20.0	0.98	-6000.	0.50	2	-0.3
4	0.0	20.0	0.99	-6000.	0.50	3	0.2
>10	0.0	40.0	0.01	-6000.	0.50		
8	0.0	40.0	0.02	-6000.	0.50	0	-2.3
3	0.0	40.0	0.05	-6000.	0.50	0	-2.5
2	0.0	40.0	0.10	-6000.	0.50	0	-2.0

first unst. m	nu	omega over sigm	alph	omega e over omega i	Te 2 over Ti	s	delt
1	0.0	40.0	0.20	-6000.	0.50	0	-2.3
1	0.0	40.0	0.40	-6000.	0.50	0	-1.9
1	0.0	40.0	0.60	-6000.	0.50	0	-1.9
1	0.0	40.0	0.80	-6000.	0.50	0	-2.2
2	0.0	40.0	0.90	-6000.	0.50	1	-0.2
2	0.0	40.0	0.95	-6000.	0.50	1	-0.2
2	0.0	40.0	0.98	-6000.	0.50	1	-0.2
3	0.0	40.0	0.99	-6000.	0.50	2	0.7
>10	0.0	5.0	0.01	-3000.	1.00		
>10	0.0	5.0	0.02	-3000.	1.00		
>10	0.0	5.0	0.05	-3000.	1.00		
10	0.0	5.0	0.10	-3000.	1.00	0	-2.0
8	0.0	5.0	0.20	-3000.	1.00	0	-1.8
6	0.0	5.0	0.40	-3000.	1.00	1	-1.4
6	0.0	5.0	0.60	-3000.	1.00	2	-1.1
7	0.0	5.0	0.80	-3000.	1.00	4	-0.8
7	0.0	5.0	0.90	-3000.	1.00	5	-0.7
6	0.0	5.0	0.95	-3000.	1.00	5	-0.9
8	0.0	5.0	0.98	-3000.	1.00	7	-0.6
>10	0.0	5.0	0.99	-3000.	1.00		
>10	0.0	10.0	0.01	-3000.	1.00		
>10	0.0	10.0	0.02	-3000.	1.00		
7	0.0	10.0	0.05	-3000.	1.00	0	-2.3
4	0.0	10.0	0.10	-3000.	1.00	0	-2.2
2	0.0	10.0	0.20	-3000.	1.00	0	-2.4
2	0.0	10.0	0.40	-3000.	1.00	0	-2.0
2	0.0	10.0	0.60	-3000.	1.00	0	-1.9
2	0.0	10.0	0.80	-3000.	1.00	1	-1.3
2	0.0	10.0	0.90	-3000.	1.00	1	-1.4
3	0.0	10.0	0.95	-3000.	1.00	2	-0.9
4	0.0	10.0	0.98	-3000.	1.00	3	-0.6
6	0.0	10.0	0.99	-3000.	1.00	5	-0.1
>10	0.0	20.0	0.01	-3000.	1.00		
8	0.0	20.0	0.02	-3000.	1.00	0	-2.4
4	0.0	20.0	0.05	-3000.	1.00	0	-2.2
2	0.0	20.0	0.10	-3000.	1.00	0	-2.2
1	0.0	20.0	0.20	-3000.	1.00	0	-2.3
1	0.0	20.0	0.40	-3000.	1.00	0	-2.0
1	0.0	20.0	0.60	-3000.	1.00	0	-1.9
1	0.0	20.0	0.80	-3000.	1.00	0	-2.2
2	0.0	20.0	0.90	-3000.	1.00	1	-0.8
2	0.0	20.0	0.95	-3000.	1.00	1	-0.9
2	0.0	20.0	0.98	-3000.	1.00	1	-0.9
3	0.0	20.0	0.99	-3000.	1.00	2	-0.3
8	0.0	40.0	0.01	-3000.	1.00	0	-2.4
4	0.0	40.0	0.02	-3000.	1.00	0	-2.4
2	0.0	40.0	0.05	-3000.	1.00	0	-2.2
1	0.0	40.0	0.10	-3000.	1.00	0	-2.2
1	0.0	40.0	0.20	-3000.	1.00	0	-1.8
1	0.0	40.0	0.40	-3000.	1.00	0	-1.5
1	0.0	40.0	0.60	-3000.	1.00	0	-1.5
1	0.0	40.0	0.80	-3000.	1.00	0	-1.6
1	0.0	40.0	0.90	-3000.	1.00	0	-2.0
2	0.0	40.0	0.95	-3000.	1.00	1	-0.2
2	0.0	40.0	0.98	-3000.	1.00	1	-0.2
2	0.0	40.0	0.99	-3000.	1.00	1	-0.2
>10	0.0	5.0	0.01	-4000.	1.00		
>10	0.0	5.0	0.02	-4000.	1.00		
>10	0.0	5.0	0.05	-4000.	1.00		
>10	0.0	5.0	0.10	-4000.	1.00		
8	0.0	5.0	0.20	-4000.	1.00	0	-1.8
7	0.0	5.0	0.40	-4000.	1.00	1	-1.3
6	0.0	5.0	0.60	-4000.	1.00	2	-1.2

first unst. m	nu	omega over sigm	alph	omega e over omega i	Te Z over Ti	s	delt
7	0.0	5.0	0.80	-4000.	1.00	4	-0.9
7	0.0	5.0	0.90	-4000.	1.00	5	-0.9
7	0.0	5.0	0.95	-4000.	1.00	6	-0.9
9	0.0	5.0	0.98	-4000.	1.00	8	-0.6
>10	0.0	5.0	0.99	-4000.	1.00		
>10	0.0	10.0	0.01	-4000.	1.00		
>10	0.0	10.0	0.02	-4000.	1.00		
7	0.0	10.0	0.05	-4000.	1.00	0	-2.4
4	0.0	10.0	0.10	-4000.	1.00	0	-2.3
2	0.0	10.0	0.20	-4000.	1.00	0	-2.4
2	0.0	10.0	0.40	-4000.	1.00	0	-2.0
2	0.0	10.0	0.60	-4000.	1.00	0	-2.0
2	0.0	10.0	0.80	-4000.	1.00	1	-1.4
2	0.0	10.0	0.90	-4000.	1.00	1	-1.4
3	0.0	10.0	0.95	-4000.	1.00	2	-1.0
4	0.0	10.0	0.98	-4000.	1.00	3	-0.7
6	0.0	10.0	0.99	-4000.	1.00	5	-0.2
>10	0.0	20.0	0.01	-4000.	1.00		
8	0.0	20.0	0.02	-4000.	1.00	0	-2.5
4	0.0	20.0	0.05	-4000.	1.00	0	-2.2
2	0.0	20.0	0.10	-4000.	1.00	0	-2.3
1	0.0	20.0	0.20	-4000.	1.00	0	-2.4
1	0.0	20.0	0.40	-4000.	1.00	0	-2.0
1	0.0	20.0	0.60	-4000.	1.00	0	-2.0
1	0.0	20.0	0.80	-4000.	1.00	0	-2.3
2	0.0	20.0	0.90	-4000.	1.00	1	-0.9
2	0.0	20.0	0.95	-4000.	1.00	1	-1.0
3	0.0	20.0	0.98	-4000.	1.00	2	-0.4
3	0.0	20.0	0.99	-4000.	1.00	2	-0.4
8	0.0	40.0	0.01	-4000.	1.00	0	-2.4
4	0.0	40.0	0.02	-4000.	1.00	0	-2.4
2	0.0	40.0	0.05	-4000.	1.00	0	-2.2
1	0.0	40.0	0.10	-4000.	1.00	0	-2.3
1	0.0	40.0	0.20	-4000.	1.00	0	-1.8
1	0.0	40.0	0.40	-4000.	1.00	0	-1.6
1	0.0	40.0	0.60	-4000.	1.00	0	-1.6
1	0.0	40.0	0.80	-4000.	1.00	0	-1.8
1	0.0	40.0	0.90	-4000.	1.00	0	-2.0
2	0.0	40.0	0.95	-4000.	1.00	1	-0.3
2	0.0	40.0	0.98	-4000.	1.00	1	-0.4
2	0.0	40.0	0.99	-4000.	1.00	1	-0.4
>10	0.0	5.0	0.01	-6000.	1.00		
>10	0.0	5.0	0.02	-6000.	1.00		
>10	0.0	5.0	0.05	-6000.	1.00		
>10	0.0	5.0	0.10	-6000.	1.00		
>10	0.0	5.0	0.20	-6000.	1.00		
7	0.0	5.0	0.40	-6000.	1.00	1	-1.4
8	0.0	5.0	0.60	-6000.	1.00	3	-1.1
8	0.0	5.0	0.80	-6000.	1.00	5	-1.0
8	0.0	5.0	0.90	-6000.	1.00	6	-0.9
10	0.0	5.0	0.95	-6000.	1.00	9	-0.9
10	0.0	5.0	0.98	-6000.	1.00	9	-0.7
>10	0.0	5.0	0.99	-6000.	1.00		
>10	0.0	10.0	0.01	-6000.	1.00		
>10	0.0	10.0	0.02	-6000.	1.00		
7	0.0	10.0	0.05	-6000.	1.00	0	-2.5
4	0.0	10.0	0.10	-6000.	1.00	0	-2.4
2	0.0	10.0	0.20	-6000.	1.00	0	-2.5
2	0.0	10.0	0.40	-6000.	1.00	0	-2.0
2	0.0	10.0	0.60	-6000.	1.00	0	-2.0
2	0.0	10.0	0.80	-6000.	1.00	1	-1.5
2	0.0	10.0	0.90	-6000.	1.00	1	-1.5
3	0.0	10.0	0.95	-6000.	1.00	2	-1.2

first unst. m	nu	omega over sigm	alph	omega e over omega i	Te 2 over Ti	s	delt
5	0.0	10.0	0.98	-6000.	1.00	4	-0.7
7	0.0	10.0	0.99	-6000.	1.00	6	-0.3
>10	0.0	20.0	0.01	-6000.	1.00		
8	0.0	20.0	0.02	-6000.	1.00	0	-2.5
4	0.0	20.0	0.05	-6000.	1.00	0	-2.3
2	0.0	20.0	0.10	-6000.	1.00	0	-2.3
1	0.0	20.0	0.20	-6000.	1.00	0	-2.5
1	0.0	20.0	0.40	-6000.	1.00	0	-2.0
1	0.0	20.0	0.60	-6000.	1.00	0	-2.0
1	0.0	20.0	0.80	-6000.	1.00	0	-2.4
2	0.0	20.0	0.90	-6000.	1.00	1	-1.1
2	0.0	20.0	0.95	-6000.	1.00	1	-1.1
3	0.0	20.0	0.98	-6000.	1.00	2	-0.6
4	0.0	20.0	0.99	-6000.	1.00	3	-0.2
8	0.0	40.0	0.01	-6000.	1.00	0	-2.5
4	0.0	40.0	0.02	-6000.	1.00	0	-2.5
2	0.0	40.0	0.05	-6000.	1.00	0	-2.3
1	0.0	40.0	0.10	-6000.	1.00	0	-2.3
1	0.0	40.0	0.20	-6000.	1.00	0	-1.9
1	0.0	40.0	0.40	-6000.	1.00	0	-1.6
1	0.0	40.0	0.60	-6000.	1.00	0	-1.6
1	0.0	40.0	0.80	-6000.	1.00	0	-1.9
1	0.0	40.0	0.90	-6000.	1.00	0	-2.2
2	0.0	40.0	0.95	-6000.	1.00	1	-0.5
2	0.0	40.0	0.98	-6000.	1.00	1	-0.6
2	0.0	40.0	0.99	-6000.	1.00	1	-0.6

END

**DATE
FILMED**

8/14/92