DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Appendix B

Invited talk at the Quark Matter '83 Workshop held at Brookhaven National Laboratory, September 1983. To be published in the Proceedings.

QUARK DISTRIBUTIONS IN NUCLEI: RESULTS FROM LEPTON PROBES

CONF-8309220--1

James P. VARY

CONF-8309220--1

Physics Department, Iowa State University, Ames, IA 50011

DE84 007430

Deep inelastic lepton scattering from nuclei indicates that quark-parton distributions are significantly different from the distributions in isolated nucleons. The EMC and SLAC data on iron and deuterium targets show enhanced and suppressed scattering for kinematic regions where Bjorken $x < 1$. In addition, the SLAC data on ^3He shows strong enhancement over results with conventional nuclear theory for $x > 1$. These data and recent model interpretations are summarized. Detailed comparisons with the data are presented for the quark cluster model.

1. INTRODUCTION

Deep inelastic lepton scattering (DIS) has proven to be a powerful tool for mapping out the charged components of the substructure of the hadrons. In a similar fashion we expect that the deep inelastic lepton scattering from nuclei will show how that substructure is altered in the nuclear medium. Recent data on ^3He ¹, on ^{56}Fe and D targets^{2,3} and on ^{27}Al ⁴ targets all indicate significantly altered quark distributions within nuclei. I will present the data and discuss the qualitative features before summarizing the specific models that have appeared. From the data alone it will be easy to see that the nucleus is not simply a collection of quasifree nucleons. I will then present the quark cluster model⁵ in a simplified manner and compare it with representative data sets. At the conclusion I will offer some speculations on the implications of these results for relativistic hadron-nucleus and nucleus-nucleus collisions.

2. DIS-KINEMATICS AND DEFINITION OF VARIABLES

The conventional single-arm inclusive experiments provide the DIS data. The initial and final lepton four-momenta are $k = (\vec{k}, E) \approx E(\hat{k}, 1)$ and $k' = (\vec{k}', E') \approx E'(\hat{k}', 1)$ respectively. The virtual photon four-momentum is $q = k - k' = (\vec{q}, E - E') \equiv (\vec{q}, \nu)$. The lab scattering angle is θ and it is conventional to define $Q^2 \equiv -q^2 = 4EE'\sin^2\theta/2$. The scattering occurs from a target nucleus of A nucleons (N neutrons and Z protons) whose mass M is approximated as Am , where m is the nucleon mass.

AC02-82ER40068

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

To obtain a perspective on these variables, consider the final state invariant mass squared W^2 which must be greater than or equal to the invariant initial mass squared $A^2 m^2$. It can be evaluated in the lab frame with $P = (0, M)$ and $P' = P + q = (\vec{q}, M + v)$ for the initial and final target four momenta respectively.

$$W^2 = P'^2 = (P + q)^2 = M^2 + 2Pq + q^2$$

or

$$W^2 = A^2 m^2 + 2Amv - Q^2 \geq A^2 m^2 = P^2. \quad (2.1)$$

From this the elastic limit in terms of experimental observables is

$$Q^2 = 4EE' \sin^2 \theta / 2 = 2Amv \quad (\text{elastic limit}). \quad (2.2)$$

In general then

$$Q^2 < 2Amv \quad (2.3)$$

and it is convenient to introduce the standard Bjorken x variable

$$x \equiv \frac{Q^2}{2m\nu} \quad (2.4)$$

which by virtue of Eqn (2.3) can range over

$$0 < x < A \quad (2.5)$$

for nuclear targets.

It is clear that DIS in the region $x > 1$ will be sensitive to effects which originate from the many-nucleon nature of the nucleus since there is no cross section from an isolated stationary nucleon ($A = 1$ in Eqn (2.5)) for $x > 1$. A typical list of many body effects would include: Fermi motion, meson exchange currents, delta isobars, clusters, short range correlations and others. I shall attempt to distinguish experimental consequences of these

various mechanisms and to draw attention to the fact that different kinematic regions, $x < 1$ vs $x > 1$, will be sensitive to different mechanisms.

The data from an arbitrary target can be cast in terms of the structure functions $\nu W_2(\nu, Q^2)$ and $\nu W_1(\nu, Q^2)$ via

$$\frac{\nu}{\sigma_M} \frac{d^2\sigma}{d\Omega dE'} = \nu W_2(\nu, Q^2) + \nu W_1(\nu, Q^2) \tan^2 \theta/2 \quad (2.6)$$

where the Mott cross section is

$$\sigma_M = \frac{4\alpha^2 E'^2 \cos^2 \theta/2}{Q^4} \quad (2.7)$$

and α is the fine structure constant. For the data I will discuss θ is sufficiently small to neglect the second term in Eqn (2.6).

DIS and the quark-parton interpretation of data on protons and deuterons has provided one of the strongest supports for QCD. Many excellent summaries of DIS and the relationship with QCD have appeared. The most recent comprehensive treatment to my knowledge, is by Buras⁷ and contains references to the other review articles. For pedantic reasons, I will adopt a simplified treatment which gains respectability as Q^2 gets very large. All calculations I report here with the quark-cluster model however are performed in a considerably more detailed framework and are discussed more adequately in References 5, 6, and 8.

In order to be mindful of scales involved it is useful to note that the de Broglie relation $Q = 2\pi/\lambda$ implies that for $Q^2 > 0.64 \text{ GeV}^2/c^2$ the virtual photon is probing distances $\lambda \leq$ nucleon diameter. For $Q^2 \sim 1 \text{ GeV}^2/c^2$ and above one therefore expects the DIS data to reflect the behavior of the substructure of the hadrons. Before discussing specific models it is appropriate to review the recent data provided by References 1-4.

3. OVERVIEW OF THE ELECTRON AND MUON DATA

The data from the EMC² on Fe and D targets has prompted a great amount of theoretical and experimental activity within the last year. In Fig. 1(a) I show the ratio R of structure functions per nucleon $(\nu W_2/56)_{\text{Fe}}/(\nu W_2/2)_{\text{D}}$ as measured by the EMC² ($9 < Q^2 < 170 \text{ GeV}^2/c^2$) and by the Rochester-SLAC-MIT experiment³ ($3 < Q^2 < 20 \text{ GeV}^2/c^2$). For comparison I present in Fig. 1(b) the ratio of aluminum to deuterium structure functions per nucleon as measured in

another Rochester-SLAC-MIT experiment⁴ ($2 < Q^2 < 20 \text{ GeV}^2/c^2$). There are several immediate observations.

1. If there were no effects of the nuclear medium ("many-body effects") on the nucleon structure function or if the many body effects were the same in Fe, Al and D then $R = 1$ would result. Both possibilities are ruled out by the $R \neq 1$ data. There are many body effects and they are different in Fe, Al and D.
2. The differences in $R(\text{Fe}/\text{D})$ and $R(\text{Al}/\text{D})$ indicate the possibility of a density effect (Fe has a higher average density than Al - see below) or of a radius effect or both.
3. The data at small x and Q^2 shown here only for $R(\text{Al}/\text{D})$ follow a trend attributed to shadowing⁴. I will not discuss this feature.
4. The apparent independence of Q^2 seen by comparing the two sets of $R(\text{Fe}/\text{D})$ data is a puzzle. This implies that the Q^2 evolution of the nuclear structure function per nucleon is independent of A .
5. The enhanced scattering for $x < 0.3$ and the suppressed scattering for $0.3 < x < 0.8$ is suggestive of a "softening" of the quark momentum distributions in the heavier nuclei. This feature is evident in most models discussed below and, in many cases, is associated with an assumed larger color singlet volume in which the quarks may move in the nucleus.

There is extensive data on ^3He ^{1,9} and some data on ^4He ⁹ from other SLAC experiments which concentrate on the $x > 1$ region. The ^3He data¹ was used in an earlier effort to demonstrate the role of six quark clusters in nuclei.⁵ Here, I show one representative data set for $\nu W_2(^3\text{He})$ corresponding to $E = 7.959 \text{ GeV}$ in Fig. 2 along with the best currently available calculation within conventional nuclear theory¹⁰. Additional comparisons may be found in References 1, 6, 8 and 10. The calculations used as a backup are the quasi-elastic nuclear structure functions $\nu W_2^{q\text{-el}}$ based upon Faddeev wavefunctions calculated¹⁰ for the Paris potential¹¹. Two observations are stimulated by the comparison in Fig. 2 and by the other ^3He data sets and calculations.

1. These high precision data are available for $1 < Q^2 < 4 \text{ GeV}^2/c^2$ and show enhanced scattering when compared with the most recent $\nu W_2^{q\text{-el}}$ in the region $x > 1$.
2. Meson exchange current and delta resonance corrections typically contribute little in this large x region. The comparison suggests many-

body cooperative mechanisms are to work since the Faddeev wavefunctions already account for the strong short range correlations.

I will summarize by claiming that the data on the various targets discussed in this section indicate significant many-body effects are present in nuclei. Furthermore, the effects seen for $x > 1$ are stronger than expected from currently available nuclear theory calculations. Later I will discuss a report that the $x < 1$ data is also stronger than expected from currently available nuclear theory calculations. Then I will give an overview of the various models before showing in some detail how the quark cluster model⁵ accounts for the qualitative features of the $x < 1$ data and the quantitative behavior of the $x > 1$ data.

4. SURVEY OF MODELS

I will divide the discussion into two areas based on the fact that the EMC data has sparked not only a resurgence of interest in some earlier models and earlier data but also the generation of new models.

4.1 Pre-1983 Data and Models

An early set of theoretical efforts¹²⁻¹⁴ predicted an "anti-shadowing" phenomenon where $R > 1$ would be found. At about the same time a 7-GeV muon experiment on various nuclear targets¹⁵ and two electron scattering experiments on nuclei at SLAC¹⁶⁻¹⁷ were reported that demonstrated shadowing ($R < 1$) was somewhat stronger and persisted to larger x than expected on the basis of model calculations. More recently, a Fermilab muon experiment at 209 GeV has also observed shadowing with nuclear targets with higher Q^2 than previously measured but still at small x . The persistence of the $R < 1$ behavior in the 7 GeV muon data to larger than expected values of x was explained by a quark-parton tube model by Berlad, Dar and Eilat¹⁹. This model also explains many features of high energy hadron-nucleus data. While the quark-parton tube model produces a dip in R similar to that shown in Fig. 1 it did not predict the $R > 1$ behavior at $x \lesssim 0.3$.

Other earlier models were stimulated by hadron-nucleus data²⁰⁻²¹ and also postulated collective mechanisms. Krzywicki²⁰ discussed the implications of his model for nuclear structure functions and qualitatively predicts features now observed in Fig. 1. The origin of these features is common to several models and will be discussed further in the context of the quark cluster model⁵ below.

Baym²² pointed out that, given our knowledge of nuclear densities and nucleon substructure, percolation of quarks and gluons could occur in nuclei. This would have a similar effect on structure functions as the models of references 19-21. Another approach which maintains the integrity of individual nucleons but parameterizes the (otherwise unknown) high momentum components of the nuclear wavefunctions has been proposed by Frankfurt and Strickman²³ and by Chemtob²⁴. Some features of earlier lepton and hadron data on nuclei were also successfully interpreted within these models^{23,24}. Noble²⁵ was motivated by a relativistic quantum field theory of interacting fermion and boson fields solved in the mean field approximation to suggest that the nucleon size and mass changes substantially within the nucleus. He pointed out that several consistency checks would be satisfied and that this could explain a number of outstanding difficulties with lower energy ($E < 1$ GeV) inelastic electron scattering experiments on nuclei.

All the models I have discussed above appeared before the EMC data. Next, I will report on papers that involve some of these models or introduce new models to explain the EMC data.

4.2. Applications of existing models to EMC data and new models

Jaffe²⁶ showed in some detail the modifications of the sea and valence quark distributions that are implied by the measured $R(\text{Fe}/\text{D})$. He also drew attention to a number of mechanisms such as percolation, meson currents, short range correlations and 6-quark clusters which would alter the quark distribution in the direction of the data.

A number of authors have given quantitative results²⁷⁻³⁰ for six and nine quark cluster effects. The degree of success in describing the data depends on the probability taken for finding a quark in a six-quark cluster in ^{56}Fe and on the assumptions about the sea quark behavior in a six-quark cluster. In general, the larger the six-quark cluster probability and/or the more the sea quarks in the six-quark cluster are shifted to smaller x then the better the description of the EMC data. Carlson and Havens²⁷ take 30%, Pirner and Vary use²⁸ 16% and Date³⁰ uses 23% for the six-quark cluster probability in ^{56}Fe . The approaches used to obtain these percentages reflect different assumptions. Carlson and Havens take 30% since this fits the EMC data while Pirner and Vary take 16% which arose from their earlier fits⁵ to the ^3He data. Date obtains the six-quark cluster probability from studying the mass number dependence of large transverse momentum production in hadron-nucleus

and in alpha-alpha collisions and from studying the mass number dependence of massive lepton pair production³¹. I will present new calculations below which take density effects into account and show that the fits to the ^3He data imply approximately 18% and 4% for the six and nine-quark cluster probabilities in ^{56}Fe respectively^{6,8}.

Several authors³²⁻³⁶ have presented calculations of $R(\text{Fe}/D)$ involving the role of pions or delta resonances in nuclei. Here again, the degree of success in describing the EMC data depends on the magnitude of the pion exchange current or magnitude of delta resonances or both. At least one effort³⁵ is based on a microscopic many body calculation³⁷ and is therefore tied to cross checks with other observables. Their conclusion is: "Although features of the data are reproduced qualitatively, we are unable to approach a quantitative description of the magnitude of the experimental effect at small x unless we postulate an enhanced pion density of about 0.4 extra pions per nucleon (roughly three times that provided by conventional nuclear models), and/or make assumptions about the small x behavior of the meson structure function which are inconsistent with available data"³⁵. Had this effort succeeded with relative ease to describe $R(\text{Fe}/D)$ I believe there would now be little motivation to study the $x < 1$ region further in the hopes of finding distinct quark effects in nuclei.

A novel picture of color conductivity in nuclei has been proposed by Nachtmann and Pirner³⁸. Loosely speaking they propose that the nucleus appears to undergo a transition from a color insulator to a color conductor as Q^2 increases. They present arguments in analogy with the Anderson model of the metal-insulator phase transition³⁹. In this picture the structure function of the nucleus at high Q^2 becomes a function only of $Q^2 R_A^2$ where R_A is the rms radius of the nucleus. Thus data from nucleons and nuclei are simply related and this picture provides an excellent description of the EMC data. Data from other nuclear targets at high Q^2 will be needed to confirm this model. If confirmed the model will present a new test for QCD since it presents a new scale $Q^2 R_A^2$ in addition to the scale provided by $x = Q^2/2m\nu$. Note that Nachtmann and Pirner only argue their model is valid at high Q^2 so their model is not in obvious contradiction with the apparent Q^2 independence of $R(\text{Fe}/D)$ as discussed above in section II of this paper. That is, the color conductivity model may be compared with the EMC data (Q^2 large) but not with the SLAC data (Q^2 small).

Two efforts⁴⁰⁻⁴¹ which propose that the nucleon's size and mass are altered in the nuclear medium can also explain $R(\text{Fe}/\text{D})$. This proposal is also based on altering the scale of color confinement in nuclei but does not involve the nuclear radius. Their analyses lead to an increase of 10-20% in the radius of a nucleon "bag" and an associated 10-20% decrease in the nucleon's mass to successfully describe the EMC data. This picture implies nuclei are even closer to percolation²² than analyses from isolated nucleons would suggest. It is interesting to note that the sign of these changes is the same as the proposal by Noble²⁵ but somewhat smaller in magnitude. Noble had proposed a 30% increase in the nucleon's radius and associated decrease in the mass inside a nucleus. The fact that the changes proposed are similar even though the underlying field theories are so different merits close scrutiny itself.

A model in which alpha clusters in nuclei explain this EMC effect has been proposed by Faissner and Kim⁴². In their model $20 \pm 5\%$ of the quarks in iron are in 12-quark clusters (alpha particles) and the remaining quarks are in nucleons.

Another approach developed by Furmanski and Krzywicki⁴³ treats the entire nucleus as a single bag or 3A-quark cluster and extends the model of anomalous nuclear enhancement introduced earlier by Krzywicki²⁰. Like many of the models summarized here this model reproduces the qualitative features of the EMC data but does not quantitatively reproduce the precise behavior.

Recently, Baldin⁴⁴ has shown how the EMC effect can be explained in terms of his model for the cumulative effect²¹ in hadron-nucleus interactions. A feature common to the cumulative model, the tube model and the various cluster models is the enhanced cross sections for $x > 1$. The degree to which these various models explain the $x > 1$ ^3He and ^4He data^{1,9} would therefore prove an interesting comparison. It is the support for high momentum components that implies (through normalization conditions) that $x < 1$ distributions are also altered. Another common feature is the qualitative behavior of these alterations in the $x < 1$ region which will be discussed below with the specific case of the quark cluster model⁵.

Fredriksson⁴⁵ has developed an explanation of the EMC effect based on increasing the radius of an assumed diquark substructure of the nucleon⁴⁶ within the nuclear medium. He finds that increasing the diquark radius by 10-45% in iron would reproduce the data. As in a number of the other models

discussed above there is an increase in some characteristic distance over which quarks are free to move within the nucleus which provides the description of the qualitative features in the data.

The above summary is most likely incomplete and will be happily updated in future publications if any authors missing would kindly provide manuscripts.

It is clear from the above discussion there are many models which appear capable of explaining a subset of the $x < 1$ data²⁻⁴. Where appropriate these models should also be applied to other data available^{1,9}. With the remaining time I would like to give an overview of the quark cluster model⁵ and its application of the $x < 1$ and the $x > 1$ data sets.

5. QUARK CLUSTER MODEL AND FITS TO ^3He DATA

The chief assumptions of this model^{5,6,8} may be summarized for describing DIS from a nucleus.

1. A photon absorbed by a nucleus at high Q^2 is absorbed by a quark through a quasi-free process when viewed in the infinite momentum frame.
2. The quark is a constituent of a quasi-free color singlet cluster in the nucleus.
3. A 3-quark (3q) cluster (nucleon) is assigned a critical radius R_c .

Clusters with $i = 6, 9, 12, \dots, 3A$ quarks are defined by the number of 3q clusters joined by links of length $\leq 2R_c$.

The critical radius R_c is taken as a free parameter and adjusted to fit the ^3He data¹. There are two interesting limiting situations. If $R_c = 0$ provides the best fit then no quark clusters larger than nucleons are formed and the "standard model" of the nucleus survives. If R_c is large, say $\sim 1.2-1.5$ Fm then percolation occurs and the nucleus has a high probability of being found in a $3A$ -q cluster configuration. We actually find $R_c \approx 0.5$ Fm which implies appreciable 6-q cluster probability in the nucleus.

I will now present the model details in a simplified version which is valid at very large Q^2 . For proper accounting of thresholds and masses see references 5, 6, and 8.

The model is easily visualized in the infinite momentum frame (referred to as the "Breit frame") defined by requiring that the quark which absorbs the

photon have longitudinal momentum $+p$ before and $-p$ after the interaction. Thus $Q = 2p$ and recalling the four vectors introduced in section II we have

$$2PQ = 2P(2p) = 2Amv \quad (5.1)$$

or

$$2mv = \frac{4Pp}{A} \quad (5.2)$$

Then

$$x = \frac{Q^2}{2mv} = \frac{4p^2}{\frac{4Pp}{A}} = A \frac{p}{P} \quad (5.3)$$

or

$$p = \frac{x}{A} P \quad (5.4)$$

where $\frac{x}{A}$ is the fraction of total nuclear momentum P carried by the struck quark. In this frame I neglect the transverse motion of the constituents and I find the probability a photon is absorbed at this Q^2 is proportional to finding a quark with $\frac{x}{A} P$ momentum in the nucleus. More precisely

$$vW_2(v, Q^2) = \sum_j \text{quarks} e_j^2 \frac{x}{A} \mathcal{P}_j(x) \quad (5.5)$$

where we sum over all quarks in the nucleus, e_j is the charge on quark j and $\mathcal{P}_j(x)$ is the probability that quark j carries fraction $\frac{x}{A}$ of nuclear four momentum P . I assume a properly weighted average of up and down quark distributions and write

$$vW_2(v, Q^2) \approx \sum_j \text{quarks} e_j^2 \frac{x}{A} \mathcal{P}(x) \quad (5.6)$$

According to the model assumptions⁵ given above I assume the quarks are found in an i -quark cluster ($i = 3, 6, 9, \dots, 3A$) within the nucleus with probability $\tilde{p}_i$ so that

$$\mathcal{P}(x) = \sum_i \text{clusters} \tilde{p}_i \bar{\mathcal{P}}_i(x) \quad (5.7)$$

where $\bar{P}_i(x)$ is the x distribution of quarks from an i -q cluster in the nucleus. That is

$$\bar{P}_i(x) = \int_0^A dy \int_0^{i/3} du n_{q/i}(u) N_{i/A}(y) \delta\left(\frac{u}{i/3} \cdot \frac{y}{A} - \frac{x}{A}\right) \quad (5.8)$$

which is the convolution of the probability $\bar{n}_{q/i}(u)$ that a quark carries momentum fraction $u/(i/3)$ of the cluster's momentum fraction yP with the probability $N_{i/A}(y)$ the cluster carries momentum fraction yP of the total nuclear momentum P . The delta function selects those probability products which give a quark at the required momentum $\frac{x}{A} P$.

For 3-q clusters the distributions from best fits to data and from QCD evolution equations are employed⁴⁷. The sum in Eqn (5.7) is terminated at the 9-q cluster contribution. Then, using $n_{q/i}(3u/i) = (i/3) \bar{n}_{q/i}(u)$, the quark distributions for 6-q and 9-q clusters are taken from counting rules⁴⁸ and Regge behavior to be:

$$n_{q/6}(v) = 1.762(v)^{-1/2}(1-v)^9 \quad (5.9)$$

$$n_{q/9}(v) = 2.239(v)^{-1/2}(1-v)^{15}. \quad (5.10)$$

The characteristic features of the $\bar{n}_{q/i}$ can be seen in a plot of $u\bar{n}_{q/i}(u) = v n_{q/i}(v)$ as a function of $u = iv/3$ as shown in Fig. 3. For the 3-q cluster I plot the corresponding distribution from counting rules and Regge behavior

$$n_{q/3}(v) = 1.094 v^{-1/2} (1-v)^3 \quad (5.11)$$

which is a reasonable approximation to the nucleon data only at the largest Q^2 measured⁴⁷. If I neglect the role of Fermi motion of clusters which is represented by $N_{i/A}$ in Eqn (5.8) then $u = x$ and I can read several features to be expected from the quark cluster model directly from Fig. 3. First at small x values ($x < 0.1$) the 6-q and 9-q distributions are slightly enhanced over the 3-q cluster. Second, there is a broad region $0.1 < x < 0.75$ where the 6-q and 9-q distributions are depressed relative to the 3-q distribution. Finally, for $x > 0.75$ the 3-q distribution drops rapidly due to its kinematic limit at $x = 1$. These three regions of behaviour correspond roughly to the

behavior of the EMC data. That is, even before we improve the model by incorporating Fermi motion of clusters and sea quark effects we can see that by including more 6-q clusters in iron and aluminum than in deuterium we will reproduce the trend of the EMC² and SLAC^{3,4} data.

The role played by Fermi motion in most of the analyses performed to date^{5,6,8} has been minor compared to the change in quark distributions from 3-q clusters to 6-q clusters. Hence, I have adopted our simplified treatment⁵ of Fermi motion at present

$$N_{i/A}(y) = \frac{1}{(2\pi)^{1/2}\sigma} \exp\left[-\frac{(y - i/3)^2}{2\sigma^2}\right] \quad (5.12)$$

with $\sigma = 0.1152$ based upon a simplified analysis of a ${}^3\text{He}$ wave function. Other efforts in progress are based upon boosting semi-realistic nuclear wavefunctions and will be reported elsewhere⁶.

To fit the ${}^3\text{He}$ data to obtain R_c one must incorporate the contributions of the nucleon quasi elastic peak. Earlier we⁵ employed a harmonic oscillator wavefunction for vW_2^{q-el} and to obtain the cluster probabilities $\tilde{p}_i$ as a function of R_c . However, now we^{6,8} employ wavefunctions obtained from solutions^{10,49} of the Faddeev equations for the realistic Paris¹¹ and the semirealistic Malfliet-Tjon⁵⁰ potentials. Hence the final form of the structure function is

$$vW_2^{\text{tot}}(v, Q^2) = vW_2 + \tilde{p}_3 vW_2^{q-el}(v, Q^2) \quad (5.13)$$

where vW_2^{q-el} will now be taken from the results of the Hannover group¹⁰ and the $\tilde{p}_i$ will be calculated from the wavefunctions provided by the Los Alamos group⁴⁹. As long as the wavefunctions have at least some semi-realistic short range correlations we do not expect significant sensitivity of the $\tilde{p}_i$ values to varying the choice of wavefunction. To test this we have evaluated the $\tilde{p}_i$ in ${}^3\text{He}$ ^{6,8} with Faddeev wavefunctions⁴⁹ from various semi-realistic nucleon-nucleon potentials⁵⁰ and have found the $\tilde{p}_i$ as a function of R_c are primarily sensitive to the rms radius of ${}^3\text{He}$ given by the wavefunction. Thus if we take only those wavefunctions which have approximately the measured rms radius we obtain little variation in the $\tilde{p}_i$ values.

The form of the model I have presented here is simplified by neglecting scaling violations (additional Q^2 dependence in vW_2) due to finite mass

effects. The full calculations I present below include these effects and more complete details of the model may be found in references 5, 6 and 8.

We fit the ${}^3\text{He}$ data sets¹ for $7.26 < E < 14.70$ GeV since they span $0.8 < Q^2 < 4$ GeV²/c² and obtain $R_c = 0.50$ Fm. This implies $(\tilde{p}_3, \tilde{p}_6, \tilde{p}_9) = (0.88, 0.11, 0.01)$ for the quark cluster probabilities in ${}^3\text{He}$. The vW_2^{tot} for $E = 7.959$ GeV is shown in Fig. 2 as the solid curve. The 6-q cluster provides enhanced cross section in the neighborhood of $x = 2$. The cluster model results will have a smoothed behavior once the effects of transverse motion are included⁶.

In our earlier effort⁵ with harmonic oscillator wavefunctions we found a best fit with $R_c = 0.45$ Fm and somewhat larger 6-q and 9-q cluster probabilities. That there is only a small change in R_c resulting from the use of more realistic nuclear wavefunctions can be understood in terms of the interplay between vW_2 and $vW_2^{q=1}$ as short range correlations are introduced. As short range correlations become more realistic, $vW_2^{q=1}$ strengthens in the $x > 1$ region and more closely approaches the dashed curve in Fig. 2. At the same time, for fixed R_c , the $\tilde{p}_6$ and $\tilde{p}_9$ values decrease so that vW_2 for $x > 1$ weakens. The combination leaves vW_2^{tot} relatively constant for fixed R_c .

For the remainder of this presentation I will hold R_c fixed at 0.50 Fm as determined by the model fit to the ${}^3\text{He}$ data.

6. APPLICATIONS TO OTHER NUCLEI

Appropriate many body wavefunctions with short range correlations are not readily available for heavier nuclei. We therefore adopt the ${}^3\text{He}$ results but scale the overlap probabilities according to changes in the average one-body density of the nucleus under consideration relative to ${}^3\text{He}$. We take the charge densities as measured by elastic electron scattering and correct for the finite proton charge form factor to arrive at $\rho(r)$ which we assume is the same for neutrons and protons. Then we approximate the density dependence of $\tilde{p}_i$ by

$$\tilde{p}_i(R_c, A) = \tilde{p}_i(\bar{R}_c, A=3) \quad (5.14)$$

where

$$\bar{R}_c = \frac{\langle \rho \rangle_A^{1/3}}{\langle \rho \rangle_{A=3}^{1/3}} R_c \quad (5.15)$$

in which the average density of nucleus A is

$$\langle \rho \rangle_A \equiv \int \rho_A^2 d^3r / \int \rho_A d^3r . \quad (5.16)$$

For the present purposes I even use this to obtain $\tilde{p}_6$ for deuterium. I only expect a small change when I use a realistic deuteron wavefunction. With $R_c = 0.50$ Fm and the ^3He Faddeev wavefunction from the Malfliet-Tjon I-III potential I obtain $(\tilde{p}_3, \tilde{p}_6) = (0.964, 0.036)$ for deuterium and $(\tilde{p}_3, \tilde{p}_6, \tilde{p}_9) = (0.777, 0.180, 0.043)$ for ^{56}Fe .

In order to compare with $R(\text{Fe}/D)$ in the low x region we can neglect the quasi-elastic contributions but we must include sea quark effects in the clusters²⁸. We take the sea quark distribution to be the same form in all clusters²⁸ and to be that given by Buras and Gaemers⁴⁷. Putting together our assumed valence and sea quark distributions for 6-q and 9-q clusters we effectively assume that less of the momentum per nucleon is carried by valence quarks and more of the momentum per nucleon is carried by sea quarks in 6-q and 9-q clusters. We find the momentum per nucleon carried by the gluons increases in 6-q and 9-q clusters to preserve momentum balance²⁸. Our assumptions differ from those of Carlson and Havens²⁷ in that they enhance the sea distributions further and maintain the gluon momentum fraction per nucleon in 6-q clusters the same as in 3-q clusters. Also, they do not consider the role of 9-q clusters and take $(\tilde{p}_3, \tilde{p}_6) = (0.70, 0.30)$. As a result of these differences they obtain a somewhat better description of $R(\text{Fe}/D)$ than our own²⁸.

I present in Fig. 4 two sets of results for $R(\text{Fe}/D)$ obtained from our model. The solid curve corresponds to $E = 100$ GeV, $\theta = 8^\circ$ and includes the sea quark distributions appropriate to a fixed $Q^2 = 20$ (GeV/c)² and is intended to approximate the kinematic region of the EMC data². The dashed curve corresponds to $E = 20$ GeV and includes the sea quark distributions at $Q^2 = 1.8$ (GeV/c)² and approximates the kinematic region of the SLAC data³. The comparison clearly indicates this data is not completely understood within our quark-cluster model at this stage. It is particularly curious that the data do not indicate a Q^2 dependence, a point raised earlier in this presentation. At the present level of our development of the quark cluster model we can only say that the $x > 1$ data appears to be adequately described but there is need for improving our understanding of the $x < 1$ data. We are

working on a number of improvements to our model⁶ but we also leave open the possibility that the $x < 1$ data indicate the presence of other phenomena such as those summarized in section IV of this presentation.

Before closing I would like to promote the idea that more experiments in the $x > 1$ region would be especially important for refining the quark cluster model. In particular the model predicts an interesting "staircase" behavior for $R(A_1/A_2)$ where nucleus A_1 is more dense on average than nucleus A_2 . For $1 < x < 2$ a plateau is predicted at $R \approx \tilde{p}_6(A_1)/\tilde{p}_6(A_2)$, then for $2 < x < 3$ another plateau is predicted for $R \approx \tilde{p}_9(A_1)/\tilde{p}_9(A_2)$ and so on. The observation of these plateaus and their magnitudes would be a strong confirmation of the quark cluster model and would provide refined information on how multiquark clusters are formed in nuclei.

7. CONCLUSIONS

Deep inelastic lepton scattering from nuclei has already revealed that quark-parton distributions in nuclei are significantly altered from those of isolated nucleons. The alterations are so strong they may not be explained by conventional nuclear models for both the $x < 1$ region³⁵ and for the $x > 1$ region¹⁰. Many models have been introduced which purport to explain features of at least some data. These models involve different explicit or implied assumptions about how the momentum shared by quarks and gluons is altered in the nucleus. In addition the dependence on momentum transfer, atomic number, radius and density distribution are different in various models. Naturally this suggests that more data on different targets even for $x < 1$ and modest Q^2 will eliminate some models. Results from a recent SLAC experiment⁵¹ should be helpful in this context when they are available. On the other hand many of the models have yet to be applied to other existing data to which I have referred. Therefore, I exhort the model developers to address this additional data. I also suggest they give absolute cross sections (check the abundant deuterium measurements), predict the Q^2 dependence and predict the $x > 1$ behavior of their model.

I have given an overview of the quark cluster model and a sample of its results. I would like to stress the need for good data from several nuclei in the $x > 1$ region since the quark cluster picture indicates a staircase feature to $R(A_1/A_2)$ in that region.

There are additional implications of the quark cluster model for subthreshold production, lepton pair production and high p_T anomalies in hadron-nucleus and nucleus-nucleus collisions⁵². We have also shown⁵³ that the measured elastic form factor of ^3He can be explained by the quark cluster model. Although time does not permit addressing these matters here I would like to conclude with the following speculation. Within the quark cluster model we have seen that DIS explores the altered quark-parton distributions in nuclei due to many-body phenomena at short distances. These correlations are "zero temperature" quantum field theory fluctuations into a quark-gluon plasma configuration. Such fluctuations are important for the initial stages of relativistic heavy ion collisions. For example α - α collisions at $\frac{v}{c} \sim 1$ have 1 interaction in 10^5 or 10^6 where a 12-q cluster collides with another 12-q cluster. In other words two zero-temperature quark-gluon "plasma fluctuations" collide with high energy density already present in the initial state. It will be important to trace the consequences of such initial state fluctuations for predictions of quark-gluon phase transitions in relativistic nucleus-nucleus collisions.

ACKNOWLEDGEMENTS

I would like to acknowledge my principal collaborator in the quark cluster model work which I have summarized, Hans Pirner. I also acknowledge numerous fruitful discussions with my other collaborators Sid Coon and Guillermo Bozzolo. I thank Stan Brodsky and Fritz Coester for helpful information related to preparing this paper. This work was supported by USDOE contract no. DE-AC02-82ER40068, HENP Division.

REFERENCES

1. D. Day et al. Phys. Rev. Lett. 43 (1979) 1143.
2. European Muon Collaboration (EMC), J. J. Aubert et al. Phys. Lett. 123B (1983) 123.
3. A. Bodek et al. Phys. Rev. Lett. 50 (1983) 1431.
4. A. Bodek et al. Phys. Rev. Lett. 51 (1983) 534.
5. H. J. Pirner and J. P. Vary, Phys. Rev. Lett. 46 (1981) 1376.
6. H. J. Pirner and J. P. Vary, to be published.

7. A. J. Buras, Rev. Mod. Phys. 52 (1980) 199.
8. J. P. Vary and H. J. Pirner, to be published in the Proceedings of the Third International Conference on Recent Progress in Many Body Theories, Altenberg, West Germany, Lecture Notes in Physics, Springer-Verlag (Heidelberg).
9. S. Rock et al. Phys. Rev. C 26 (1982) 1592.
10. H. Meier-Hajduk, Ch. Hajduk, P. U. Sauer and W. Theis, Nucl. Phys. A 395 (1983) 332.
11. M. Lacombe et al. Phys. Rev. C 21 (1980) 861.
12. V. I. Zakharov and N. N. Nikolaev, Sov. J. Nucl. Phys. 21 (1975) 227; Phys. Lett 55B (1975) 397.
13. J. D. Bjorken, Various review talks around 1975.
14. N. N. Nikolaev, Sov. Phys. Usp. 24 (1982) 531.
15. M. May et al. Phys. Rev. Lett. 35 (1975) 407.
16. S. Stein et al. Phys. Rev. D 12 (1975) 1884.
17. J. W. Cronin et al. Phys. Rev. D 11 (1975) 3105.
18. M. S. Goodman et al. Phys. Rev Lett. 47 (1981) 293.
19. G. Berland, A. Dar and G. Eilam, Phys Rev D 22 (1980) 1547.
20. A. Krzywicki, Phys. Rev D 14 (1976) 152.
21. A. M. Baldin, in Proceedings of the 19th International Conference on High Energy Physics, Tokyo, 1978, edited by S. Homma, M. Kawaguchi, and H. Miyazawa (Phys. Soc. of Japan, Tokyo, 1979), p. 455; Fiz. Elem. Chastits At. Yadra (1977) 429 [Sov J. Part. Nucl. 8, 429 (1977)] and references therein.
22. G. Baym, Physica 96A (1979) 131.
23. L. L. Frankfurt and M. I. Strickman, Phys. Rep. 76 (1981) 215; Nucl. Phys. B 181 (1981) 22 and references therein.
24. M. Chemtob, Nucl. Phys. A 336 (1980) 299.
25. J. V. Noble, Phys. Rev. Lett. 46 (1981) 412.
26. R. L. Jaffe, Phys. Rev. Lett. 50 (1983) 228.
27. C. E. Carlson and T. J. Havens, Phys. Rev. Lett. 51 (1983) 261.
28. H. J. Pirner and J. P. Vary, Heidelberg preprint, UNI-HD-83-02 (1983).

29. A. I. Titov, Dubna preprint, E2-83-72 (1983).
30. S. Date, Waseda University, Tokyo, preprint WU-HEP-83-4.
31. S. Date and A. Nakamura, Prog. Theor. Phys. 69 (1983) 565.
32. J. Szwed, Cracow preprint, TPJU - 1/83 (1983).
33. C. H. Llewellyn Smith, Oxford Preprint (1983).
34. M. Ericson and A. W. Thomas, Cern preprint, TH 3553 (1983).
35. E. L. Berger, F. Coester and R. B. Wiringa, Argonne preprint, ANL-HEP-PR-83-24 (1983).
36. D. Stump, G. Bertsch and J. Pumplin, contribution to the Workshop on Manifestations of Hadron Substructure in Nuclear Physics, Bloomington, Indiana, October 19-21, 1983.
37. B. L. Friman, V. R. Pandharipande and R. B. Wiringa, Phys. Rev. Lett. 51 (1983) 763.
38. O. Nachtmann and H. J. Priner, Heidelberg preprint, HD-THEP-83-8 (1983).
39. P. W. Anderson, Phys. Rev. 109 (1958) 1492.
40. F. E. Close, R. G. Roberts and G. G. Ross, Rutherford preprint, RL-83-051, T-324 (1983).
41. M. Staszczel, J. Rozynek and G. Wilk, Warsaw preprint (1983).
42. H. Faissner and B. R. Kim, Aachen preprint (1983).
43. W. Furmanski and A. Krzywicki, contribution to the Sixth High Energy Heavy Ion Study and Second Workshop on Anomalons, Berkeley, California, June 1983; Orsay preprint, LPTH 83/11 (1983).
44. A. M Baldin, Dubna preprint 2-83-415 (1983).
45. S. Fredriksson, Stockholm preprint, TRITA-TFY-83-15 (1983).
46. S. Fredriksson, M. Jandel and T. I. Larsson, Z. Phys. C 14 (1982) 35; C 19 (1983) 53.
47. A. J. Buras and K. J. F. Gaemers, Nucl. Phys. B 132 (1978) 249.
48. D. Sivers, S. J. Brodsky and R. Blankenbeckler, Phys. Rep. 23C (1976) 1, and references therein.
49. J. L. Friar, B. F. Gison, E. L. Tomusiak and G. L. Payne, Phys. Rev. C 24 (1981) 665; and private communication.
50. R. A. Malfliet and J. A. Tjon, Nucl. Phys. A 127 (1969) 161.
51. R. G. Arnold et al. SLAC experiment E139 (1983).

52. G. Bozzolo, H. J. Pirner and J. P. Vary, to be published.
53. J. P. Vary, S. A. Coon and H. J. Pirner, Proceedings of the Tenth International Conference on Few Body Problems in Physics, Karlsruhe, Germany, August 21-27, 1983 (to be published).

$X \rightarrow$

Fig 1(a)

$X \rightarrow$

Fig 1(b)

Fig 2

Fig 3

Fig 4