ORNL
MASTER COPY

OAK RIDGE NATIONAL LABORATORY

operated by

UNION CARBIDE CORPORATION

for the

U.S. ATOMIC ENERGY COMMISSION

ORNL-TM- 73

COPY NO. - 4

DATE - December 5, 1961

On the Reality of the Eigen-Values for a One-group
N-region Diffusion Problem

T. W. Hildebrandt

NOTICE

RELEASE APPROVED
BY PATENT BRANCH

SIGNATURE

This document contains information of a preliminary nature and was prepared primarily for internal use at the Oak Ridge National Laboratory. It is subject to revision or correction and therefore does not represent a final report. The information is not to be abstracted, reprinted or otherwise given public dissemination without the approval of the ORNL patent branch, Legal and Information Control Department.

UNCLASSIFIED

1. In the computation of the eigen-values for multiregion diffusion problems, some assurance is required that the eigen-values, and, in particular, the eigen-values of largest or smallest modulus, are real. This note provides a sufficient condition for the reality of the eigen-values for a fairly general one-group, multiregion two-dimensional diffusion problem. The condition is also sufficient to insure the orthogonality of eigen-functions corresponding to distinct eigen-values.

2. Consider the bounded, connected plane region S , which is subdivided into N subregions S_i , $i = 1, \dots, N$, by the Jordan arcs G_{ij} , $i = 1, \dots, N$; $j = 0, 1, \dots, N$.

The arc G_{ij} is to be taken with a positive sense when viewed from the interior of the region S_i . Thus $G_{ji} = -G_{ij}$. The arc G_{i0} is the segment of the outer boundary of S which is common to the boundary of S_i .

In each region S_i , the linear differential equation with real coefficients

$$(1a) \quad \nabla^2 u_i + D_i u_i = \lambda \theta_i u_i; \quad \theta_i \geq 0, \quad i = 1, \dots, N; \quad \sum_{i=1}^N \theta_i > 0.$$

(θ_i constant in each S_i ; D_i may be a function of position only)

is to be satisfied, with the eigen-value λ to be constant over those regions S_i where $\theta_i > 0$.

UNCLASSIFIED

The following interface conditions are to be satisfied on each

$$G_{ij}, \quad i = 1, \dots, N \\ j = 1, \dots, N.$$

$$(1b) \quad \begin{cases} u_i = \alpha_{ij} u_j + \beta_{ij} \frac{\partial u_j}{\partial n} \\ \frac{\partial u_i}{\partial n} = \gamma_{ij} u_j + \delta_{ij} \frac{\partial u_j}{\partial n} \end{cases} \quad ; \quad \varphi_{ij} = (\alpha_{ij} \delta_{ij} - \beta_{ij} \gamma_{ij}) \neq 0. \\ \alpha_{ij}, \dots, \delta_{ij} \text{ real.}$$

On the outer boundary G_{i0} , $i = 1, \dots, N$, we may specify either

$$u_0 = 0 \quad \text{or} \quad \frac{\partial u_0}{\partial n} = 0, \quad \text{but not both, or, more generally, the determinant}$$

$$\varphi_{i0} = 0.$$

Note that where two subregions S_i and S_j have no common interface, a nonsingular ($\varphi_{ij} \neq 0$) condition is satisfied vacuously, since $G_{ij} = 0$; and also that $\varphi_{ii} = 1$, $G_{ii} = 0$, which is to say that the solution is to be continuous within any given subregion S_i .

Note also that $\varphi_{ij} \neq 0$ implies

$$(2) \quad \varphi_{ji} = \frac{1}{\varphi_{ij}}.$$

3. Assume that the problem has a complex eigen-value λ , and a corresponding eigen-function u . Then, because of the reality of the coefficients, the conjugates $\bar{\lambda}$ and $\bar{u}$ must also satisfy the problem:

UNCLASSIFIED

$$\left. \begin{aligned} (3a) \quad \nabla^2 u_i + D_i u_i &= \lambda \theta_i u_i \\ (3b) \quad \nabla^2 \bar{u}_i + D_i \bar{u}_i &= \bar{\lambda} \theta_i \bar{u}_i \end{aligned} \right\} \quad i = 1, \dots, N$$

Multiply (3a) by $\bar{u}_i$, and (3b) by u_i , and take the (surface) integral of the difference.

$$(\lambda - \bar{\lambda}) \theta_i \iint_{S_i} u_i \bar{u}_i d\sigma = \iint_{S_i} (\bar{u}_i \nabla^2 u_i - u_i \nabla^2 \bar{u}_i) d\sigma; \quad i = 1, \dots, N.$$

The application of Green's theorem in each subregion leads to

$$(\lambda - \bar{\lambda}) \theta_i \iint_{S_i} u_i \bar{u}_i d\sigma = \sum_{j=0}^N \int_{G_{ij}} (\bar{u}_i \frac{\partial u_i}{\partial n} - u_i \frac{\partial \bar{u}_i}{\partial n}) ds; \quad i = 1, \dots, N.$$

Multiply each equation by a constant $K_i > 0$, to be determined, and sum:

$$(4) \quad (\lambda - \bar{\lambda}) \sum_{i=1}^N K_i \theta_i \iint_{S_i} u_i \bar{u}_i d\sigma = \sum_{i=1}^N \sum_{j=0}^N K_i \int_{G_{ij}} (\bar{u}_i \frac{\partial u_i}{\partial n} - u_i \frac{\partial \bar{u}_i}{\partial n}) ds.$$

Now for $i < j$, and for $j = 0$, we make the substitution indicated by the interface conditions (1b), so that

$$\begin{aligned} K_i \int_{G_{ij}} (\bar{u}_i \frac{\partial u_i}{\partial n} - u_i \frac{\partial \bar{u}_i}{\partial n}) &= K_i \int_{G_{ij}} (\alpha_{ij} \delta_{ij} - \beta_{ij} \gamma_{ij}) (\bar{u}_j \frac{\partial u_j}{\partial n} - u_j \frac{\partial \bar{u}_j}{\partial n}) ds \\ &= K_i \varphi_{ij} \int_{G_{ij}} (\bar{u}_j \frac{\partial u_j}{\partial n} - u_j \frac{\partial \bar{u}_j}{\partial n}) ds. \end{aligned}$$

UNCLASSIFIED

UNCLASSIFIED

Since $\varphi_{i0} = 0$, all terms containing $j = 0$ vanish. The right-hand side of (4) then becomes

$$\begin{aligned}
 & \sum_{j=1}^N \sum_{i=1}^{j-1} K_i \varphi_{ij} \int_{G_{ij}} \left(\bar{u}_j \frac{\partial u_j}{\partial n} - u_j \frac{\partial \bar{u}_j}{\partial n} \right) ds + \sum_{j=1}^N \sum_{i=j}^N K_i \int_{G_{ij}} \left(\bar{u}_i \frac{\partial u_i}{\partial n} - u_i \frac{\partial \bar{u}_i}{\partial n} \right) ds \\
 &= \sum_{j=1}^N \sum_{i=1}^{j-1} K_i \varphi_{ij} \int_{G_{ij}} \left(\bar{u}_j \frac{\partial u_j}{\partial n} - u_j \frac{\partial \bar{u}_j}{\partial n} \right) ds + \sum_{i=1}^N \sum_{j=1}^{i-1} K_i \int_{G_{ij}} \left(\bar{u}_i \frac{\partial u_i}{\partial n} - u_i \frac{\partial \bar{u}_i}{\partial n} \right) ds \\
 &= \sum_{j=1}^N \sum_{i=1}^{j-1} \left[K_i \varphi_{ij} \int_{G_{ij}} + K_j \int_{G_{ji}} \right] \left(\bar{u}_j \frac{\partial u_j}{\partial n} - u_j \frac{\partial \bar{u}_j}{\partial n} \right) ds \\
 &= \sum_{j=1}^N \sum_{i=1}^{j-1} (K_i \varphi_{ij} - K_j) \int_{G_{ij}} \left(\bar{u}_j \frac{\partial u_j}{\partial n} - u_j \frac{\partial \bar{u}_j}{\partial n} \right) ds .
 \end{aligned}$$

The first equality results from interchanging the order of summation in the second term; the second by "renaming" the indices in the second term; the last from recalling that $G_{ji} = -G_{ij}$.

From the last expression, it is obvious that if

$$(5) \quad K_i \varphi_{ij} - K_j = 0 ; \quad 0 < i < j \leq N ,$$

then the right-hand side of (4) vanishes; since $u_i \bar{u}_i = |u_i|^2 > 0$, and $K_i \varphi_i > 0$ for at least one i , the left-hand side of (4) vanishes

UNCLASSIFIED

only if $\lambda = \bar{\lambda}$ in those S_i where $\theta_i \neq 0$. Equation (5) therefore furnishes a sufficient condition that the eigen-values for the problem be real: that there exist a set of N positive constants K_i such that $K_j = K_i \varphi_{ij}$ for all $0 < i < j \leq N$. Since $K_i > 0$, $K_j > 0$, $\varphi_{ij} > 0$.

4. Assuming that equation (5) holds, it is possible to develop a more succinct statement of the condition. Note that

$$K_j = K_i \varphi_{ij} = K_h \varphi_{hj} = K_i \varphi_{ih} \varphi_{hj}, \quad i < h < j.$$

Therefore

$$\varphi_{ij} = \varphi_{ih} \varphi_{hj},$$

and in view of (2),

$$(6) \quad \begin{cases} \varphi_{ih} \varphi_{hj} \varphi_{ji} = 1, & \text{for all } 0 < i, j, h \leq N \\ \varphi_{ij} > 0 & \text{all } 0 < i, j \leq N. \end{cases}$$

Theorem. A sufficient condition that the one-group, N -region, diffusion problem (1a, b) have real eigen-values is that the determinants φ_{ij} of the interface conditions (1b) satisfy (6).

5. If $\lambda, \bar{\lambda}, u, \bar{u}$ are interpreted now to represent distinct (real) eigen-values and their corresponding eigen-functions, the identical analysis, bearing in mind that now $(\lambda - \bar{\lambda}) \neq 0$, leads to the conclusion:

Corollary. Condition (6) is sufficient to insure that the eigen-functions corresponding to distinct (real) eigen-values be orthogonal over the regions S_i where $\theta_i > 0$.

6. In certain frequently occurring special cases, the condition (6) is automatically satisfied.

(a) Continuity across the interface:

$$\begin{cases} u_i = u_j \\ K_i \frac{\partial u_i}{\partial n} = K_j \frac{\partial u_j}{\partial n} \end{cases} \quad K_i > 0, \quad \text{all } 0 < i \leq N$$

$$\varphi_{ij} = K_j / K_i$$

(b) Contact resistance

$$K_i \frac{\partial u_i}{\partial n} = K_j \frac{\partial u_j}{\partial n} = h_{ij}(u_i - u_j); \quad K_i > 0 \quad \text{all } i, j$$

$$h_{ij} \neq 0$$

implies

$$\begin{cases} u_i = u_j + \frac{K_j}{h_{ij}} \frac{\partial u_j}{\partial n} \\ \frac{\partial u_i}{\partial n} = \frac{K_j}{K_i} \frac{\partial u_j}{\partial n} \end{cases}$$

$$\varphi_{ij} = K_j / K_i$$

Distribution

- 1-3. DTIE, AEC
- 4. M. J. Skinner
- 5-6. Central Research Library
- 7. Document Reference Section