50/192
6/11/92
PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

PPPL-TH91-1
UC-420,427

PPPL-TH91-1

PRINCETON UNIVERSITY
PLASMA PHYSICS LABORATORY

THEORY DIVISION
QUARTERLY REPORT*

October 1 – December 31, 1991

*This work was supported by U.S. Department of Energy Contract No. DE-AC02-76-CHO-3073.

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the:
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
(703) 487-4650

.....

PPPL THEORY NEWS is a quarterly bulletin presenting condensed versions of selected research reports recently printed and distributed by the PPPL Theory Division at PPPL. Lists of printed reports and publications, external presentations at professional meetings, and recent visitors to the group are also included. Further information may be obtained by contacting the individual authors. Copies of PPPL Reports may be requested from the R & P Office at PPPL. Our address is:

Princeton Plasma Physics Laboratory
James Forrestal Campus
P.O. Box 451
Princeton NJ 08543

Phone: (609) 243-2656
Fax: (609) 243-2662

Roscoe R. White, *Editor*
Terry Greenberg, *Copy Production & Layout*

Contents

I Edge Plasma Modeling	3
II Theory Division Computing Modernization	4
III Error Field Effects	6
IV Theory Publication List	10
V Theory Visitor List	12

I. Edge Plasma Modeling

C. F. F. Karney

The behavior of the edge of a tokamak plasma is of the utmost importance in machines such as ITER. The crucial issues are the particle and heat fluxes to the divertor and the control of impurities flowing back to the plasma. Edge physics is characterized by complex geometry, material walls, large concentrations of neutrals, etc.

The two major approaches to modeling the plasma edge are being pursued by the Modeling and ITER Physics Division. These approaches utilize either a fluid code, which follows the electron and ion fluids but which treats the neutrals approximately, or a neutrals code, which follows the neutrals using Monte-Carlo techniques in a given plasma background. Examples of edge fluid codes are B2, written by Bas Braams, and PLANET, written by Marijan Petravic. An example of a neutrals code is DEGAS, written by Dan Heifetz.

One of the major goals of the edge modeling work at PPPL is to couple B2 and DEGAS to provide a realistic model of the plasma edge. This work being performed by Daren Stotler and David Coster. The B2 code necessarily has to include neutrals. They are a source of plasma through ionization, a source of momentum through charge-exchange, etc. All of these processes are very important in the edge. However, B2 employs a rather crude diffusion model to treat the flow of neutrals. The object of coupling B2 to DEGAS is to improve the modeling of neutrals. B2 can be run with its neutral model to provide a plasma profile for use with DEGAS. DEGAS is run to provide a neutral profile and this is used when B2 is next run. The process is repeated until convergence.

The procedure, unfortunately, is not as simple as that. The chief problems are: (a) that the convergence is poor because DEGAS produces noisy profiles which can vary significantly from one iteration to the next even close to convergence; and (b) that the procedure is CPU intensively because each invocation of DEGAS may consume many minutes of Cray time. There are various suggestions for removing these difficulties which we are currently exploring.

The other problem is that the interaction of a plasma with a material surface or with a neutral species can lead to very steep plasma gradients which invalidate the fluid description of the plasma. In the case of the interaction with a material surface, a sheath is established and boundary conditions are normally imposed on the fluid code specifying sonic flow towards the plate in the plasma

region just beyond the sheath. Similarly plasma-neutral interactions can lead to steep gradients because of the sensitivity of the charge-exchange cross-section to plasma temperature. An accurate treatment of both these situations requires a kinetic treatment of the plasma. This is the thesis topic for David Coster who is writing a Fokker-Planck code with two velocity dimensions plus one spatial dimension to study the physics of these interactions.

Another area of current interest is the effect of stochastic magnetic fields on the heat load to a divertor plate. Neil Pomphrey and Allan Reiman are studying this with a simple field-line following code. We would like to be able to incorporate their results into B2.

II. Theory Division Computing Modernization

G. Rewoldt, E. Valeo, and C.F.F. Karney

Over the last two years, the beginnings of a major change have taken place in the computing capabilities of the Theory Division. At the beginning of this period, the available equipment consisted of an Apple Macintosh or an IBM personal computer with a small, monochrome screen in the office of each theorist, along with a heavily overloaded DEC Vax minicomputer in the PPPL Computer Center, to be used in accessing the CRAY supercomputers at the National Energy Research Supercomputer Center (NERSC) in Livermore, California. In the last several years, a revolution has taken place in the capabilities of desktop workstation/servers employing Reduced Instruction Set Computing (RISC) processor chips and running the UNIX operating system. They now have processing capabilities which were available previously only from mainframes and supercomputers. The Theory Division has now purchased three Sun SPARCstation workstation/servers with DOE and NSF money, and has made arrangements to use a fourth SPARCstation and an IBM RISC System/6000 model 320 workstation/server. Each of these workstation/servers (costing \$10-20k) has at least twice the processing capacity of the USC Vax (costing ~\$500k). The fastest of these workstation/servers, the IBM model 320, has run at up to 44% of the speed of a CRAY-1, or of a single CRAY-2 processor (costing more than \$2 million), on one of our benchmark programs. These workstation/servers can be used for editing and file manipulation, for pre- and post-processing, and to run most of our small and medium-size computer calculations (some of the large calculations still have to be run on the NERSC CRAYs).

The Theory Division has also purchased twenty one graphics X-terminals from NCD, IBM, and Tektronix. These allow access to any of the workstation/servers and to any of the CRAYs by means of separate windows on the X-terminal screen, using the X-windows protocol. These X-terminals can display high-resolution(1024×1280) (8-bit) color graphics as well as text. X-terminals consist essentially of just a screen, a keyboard, a mouse, and a specialized graphics processor, and communicate with "compute servers" and "file servers" over a high-speed "ethernet" network. One of the advantages of X-terminals is that they separate the function of data display from the function of data processing. This makes economic sense in the longer term, because "compute servers" like our workstation/servers are evolving much more rapidly than X-terminals. (There seems to be a new generation of workstation/servers appearing every year at the same price with twice the speed.)

Also, in the last year, we have had an "ethernet" network, which operates at a 10 megabit/second rate, installed in every office in the Theory Division. This prepares us for the modern era of "distributed computing", where different computational resources can be in different physical locations, but are all accessible from any user's office by means of a network. Fortunately, since the price for a unit of computing power (a workstation/server) is miniscule compared to the investment in a mainframe, a moderate annual capital budget allows continual upgrading, so that the Theory Division computing facilities can be maintained, on average, near the state of the art.

In the next several years, we would like to evolve to a configuration in the Theory Division with a high-resolution color X-terminal on each physicist's desk, from which they can access the workstation/servers, with their associated peripheral equipment, such as hard disk drives and printers, as well as accessing remote supercomputers, such as the NERSC CRAYs. We would also like to get a few specialized graphics workstations, which would allow us to manipulate 3-dimensional graphics images, rather than just display them. Such a data visualization capability will become more important as our computational effort turns increasingly towards solving three-dimensional plasma models. The time it takes to realize this plan will of course depend on the funding available each year for computer equipment purchases.

III. Error Field Effects

A. Reiman and D. Monticello

Over the past few years, experimental evidence has accumulated that resonant error fields of order $\delta B/B_t \approx 10^{-4}$ can produce significant effects on tokamak discharges. Some of the observations raise serious concerns for the next generation of tokamaks. Islands produced by error fields are believed to cause nonaxisymmetric heat deposition on divertor plates. By increasing the peak heat load, this effect exacerbates the already serious problem of divertor heat load. Error fields also are believed to increase plasma disruptivity through mode locking. This is an issue because the divertor plates can only tolerate a limited number of disruptions due to ablation of material from the divertor plates during disruptions. In addition to the practical issues, error field effects also provide a testing ground for some interesting physics issues. A small, resonant magnetic field perturbation affects the plasma significantly only in a narrow region about the rational surface, allowing us to test our understanding of the physics of magnetic islands.

We have studied the effects of small, externally imposed error fields using both a cylindrical, analytical model and using the PIES three-dimensional equilibrium code. Our analytical calculation, assuming narrow islands and tokamak ordering, is a generalization of the usual nonlinear tearing mode theory to include the effects of a fixed external perturbation. Details of the calculation are presented in Ref. 1. Figure 1 shows schematically what the solution looks like. The helical flux at the rational surface, $\psi(r_s)$, is plotted as a function of a profile parameter, with Δ' assumed to increase monotonically as a function of the parameter.

The dashed lines in Fig. 1 show the solution in the absence of an error field. For $\Delta' > 0$, nonaxisymmetric solutions appear, corresponding to saturated tearing modes. As $\Delta' \rightarrow 0$ from the right, the island width goes to zero. The point $\Delta' = 0$ corresponds to a bifurcation point of the solution. The solid lines show the solution for $\epsilon > 0$. The external perturbation breaks the bifurcation at $\Delta' = 0$, so that the solution now consists of two independent branches. The solution for $\Delta' < 0$ now goes smoothly into the locked mode solution for $\Delta' > 0$. This is the branch that we are interested in.

One thing that we would like to understand about the solution of Fig. 1 is whether the plasma amplifies or diminishes the error field when $\Delta' < 0$. To get

a feeling for that, we have also studied a special case where things simplify

FIG. 1. Solution for $\psi(r_s)$ with finite ϵ . The dashed line indicates the solution in the $\epsilon = 0$ limit.

somewhat. We assume that the island is in the vacuum region outside the plasma. We find a critical value $\Delta' = \Delta'_v$ at which the island width is just that due to the vacuum error field. When Δ' is sufficiently large and negative, the plasma is self-healing. When $\Delta' < 0$ approaches zero, we get amplification.

Our numerical results have been obtained with the PIES three-dimensional equilibrium code. Some of these results have been presented in Ref. 2, and a more extensive set of numerical computations will be presented in Ref. 3. Figures 2 and 3 show Poincare plots for solutions with $R/a = 3$, $\gamma = 1$, and circular cross-section, without and with an error field. (We caution that the origin of our coordinate system does not lie on the magnetic axis, so the region about the axis looks like an island, but is not an island.) The tearing mode is unstable for these parameters. In the absence of an error field there are saturated islands at the $q = 2$ and $q = 3$ rational surfaces, with the total island width across the midplane about 6%. The solution in Fig. 3 has an error field imposed. The coil current and shift were chosen to approximately reproduce the reported $n = 1$ error field spectrum on DIII-D, with an $n = 1, m = 2$ amplitude of about $5.5 \times 10^{-5} B_t$.⁴ With the imposed error field, the islands are larger and more numerous. The total island width across the midplane is now about 23%.

Our analytical calculation suggests that the presence of an error field

FIG. 2. Poincare plot for a toroidal solution with $R/a = 3$, $\gamma = 1$ and no error field.

broadens the range of plasma parameters over which deleterious tearing effects are seen. This is consistent with the observation that the error compensation coil on DIII-D effects the threshold for the onset of low density locked modes. The numerical results indicate that the saturation amplitude of a tearing instability can be sensitive to the presence of a relatively small error field. This is consistent with the general belief that locked modes tend to grow to larger amplitude and to disrupt more readily than rotating tearing modes. We conclude that experiment and theory are presenting us with a consistent picture indicating that error fields of the magnitude that exist in present day tokamaks are having significant effects. The theoretical results suggest that error field amplitudes in the next generation of tokamak experiments will affect heat deposition peaking factors on the divertor plates (through nonaxisymmetric heat deposition), and will affect disruptivity. Improved understanding of error field effects will play a role in the design of the next generation of tokamaks in terms of suggesting construction tolerances, and possibly in terms of suggesting the desirability of trim coils for tuning out residual field errors.

FIG. 3. Poincare plot for a toroidal solution with $R/a = 3$, $\gamma = 1$ and error field as described in the text.

References

- ¹A. Reiman, Phys. Fluids **3**, 2617 (1991).
- ²A. Reiman and D. Monticello, Phys. Fluids **3**, 2230 (1991).
- ³A. Reiman and D. Monticello, Three-dimensional tokamak equilibria in the presence of resonant field errors, PPPL 2818, Princeton Plasma Physics Laboratory, 1992.
- ⁴T. Scoville et al., Bull. Am. Phys. Soc. **34**, 2118 (1989).

IV. THEORY PUBLICATION LIST 10/91 - 12/91

Multispecies Transport Theory for Axisymmetric Rotating Plasmas, M. Tessarotto and R. White PPPL Report 2798 (7/23/91), Physics of Fluids Nov 91.

Flow Shear Suppression of Turbulence Using Extremely Driven Ion Bernstein and Alven Waves, Biglari, Ono, Diamond, and Goddard, PPPL Report 2811, 9th Topical Conference on RF Plasmas to be published Charleston, SC Aug 19-21, 1991

Resonant Damping of Toroidicity Induced Shear Alfvén Eigenmodes in Tokamak, Zonca and Chen PPPL Report cancelled Phys. Rev. Letters Vol. 68, p. 592, Feb 1992

A Self-Consistent Theory of Collective Alpha Particle Losses Induced by Alfvénic Turbulence Biglari and Diamond PPPL-2809 and Phys. Rev. Letts.

The Effects of Plasma Deformability on the Feedback Stabilization of Axisymmetric Modes in Tokamak Plasmas, D. Ward and S. Jardin PPPL 2805

Effective Diffusion and Nonlocal Heat Transport in a Stochastic Magnetic Field J.M. Rax and R.B. White, No PPPL Report, Phys. Rev. Letts.

Gyrokinetic Analysis of Electrostatic Microinstabilities in the Presence of Sheared Flows, M. Artun and W.M. Tang, Physics of Fluids BPPL Report 2810.

Ring Current Instabilities in the Magnetohydrodynamic Frequency Range Hasegawa and Chen No PPPL Report, Annales Geophysicae Journal

MHD Theory of Field Line Resonance in Magnetosphere, Cheng, Chang, Ling, and Tsai, PPPL Report 2814, submitted to J. of Geophysical Research

Kinetic Extensions of Magnetohydrodynamics for Axisymmetric Toroidal Plasmas, Cheng, PPPL 2604 from 1989, Physics Reports Journal (to Elsevier 10/14/91)

Nonlinear Relativistic Interaction of an Ultrashort Laser Pulse with a Cold Plasma, Rax and Fisch PPPL, Report 2817, Phys. Fluids

Three-Dimensional Tokamak Equilibria in the Presence of Resonant Field Errors, Reiman and Monticello, PPPL Report 2818, Nuclear Fusion

The Field Line Topology of a Uniform Magnetic Superposed on the Field of a Distributed Ring Current, Chance, Greene and Jensen, PPPL Report published long ago, Geophysical and Astrophysical Fluid Dynamics 11/27/91

Anomalous Energy Exchange in the gBL and QAuasilinear Theories,
Mynick, PPPL Report #2824, Physics of Fluids B

Survey of Physics Computer Codes Supported at PPPL, editor S.C. Jardin,
Technical Memo "TM" 12/18/91

Weak-and Strong-Turbulence Regimes of the Hasegawa-Mima
Equations, Ottaviani and Krommes, PPPL Report 2826 12/19/91, Phys. Rev.
Lett.

Flow Shear Induced Compton Scattering of Electron Instability, T.S.
Hahm PPPL Report 2830 1/13/92, Phys. of Fluids B

Comparison of ATF and TJ-11 Stellarator Equilibria as Computed by the 3-
D V MEC and PIES Codes., Johnson Monticello Reiman, Salas, Fraguas,
Hirschman, PPPL Report 2828 12/11/91 submitted, J. of Computer Physics
Communications

External Kink Modes as a Model for MHD Activity Associated with ELM's
J. Manickam, Phys. Fluids, PPPL Report #2819

Flow Shear Suppression of Turbulence Using Externally Driven Ion
Berstein and Alfvén Waves, Bigalri, Ono, Diamond, and Craddock, PPPL
Report #2811.

Inverse Problem for Bremsstrahlung Radiation, Voss and Fisch, PPPL
Report 2793, Journal ?

Simulations of DT Experiments in TFTR, Budny, Bell, Biglari, PPPL
Report 2808, Nucl. Fusion

V. Theory Visitor List

For Year 1991 -

- Dewar, B., The Australian National University, Australian, January 1991 and 1992.
- Lee, Lou-Chuang, College of Natural Sciences at the University of Alaska - Fairbanks, January - March.
- Tessarotto, M., University of Trieste, Italy, January, February, April, May, July, August, September 1991 and January - February 1992.
- Xia, Peking University, PRC, March - June.
- Zonca, F. ENEA, Centro Richerche, Energia, Italy, May - July.
- Fruchtmann, Weitzmann Institute fo Science, Israel, August.
- Nakamura, Kyoto University of Plasma Physics Laboratory, Japan, September, November.
- Rath, S., Princeton University, October 1990 - September 1992.
- Rax, J.M., Institute de Recherche Fondamentale, France, September 1990 - September 1992.
- Li, Royal Institute of Technology, September - December.
- Romanelli, ENEA - Centro Ricerche Energia, Italy, November.

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-Europeen, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL
 Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James,, Univ. of Alberta, CANADA
 Dr. P. Lukáč, Komenského Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE
 C. Mouttet, Lab. de Physique des Milieux ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinri, Univ. of Ioannina, GREECE
 Dr. T. Mui, Academy Bibliographic Ser., HONG KONG
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theo. Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Director, Japan Atomic Energy Research Inst., JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Inst. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 S. Mori, Technical Advisor, JAERI, JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Libby, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Dr. O. Petrus, AL I CUZA Univ., ROMANIA
 Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 Dr. L. Stenflo, Univ. of UMEA, SWEDEN
 Library, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. I Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY
 Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

**DATE
FILMED**
8/11/92