DOE/FG02/90ER14159-2

ATOMIC PHYSICS IN STRONG FIELDS

DOE/ER/14159--2

DE92 015356

PROGRESS REPORT

For Grant Period September 15, 1990 through November 14, 1992

Shih-I Chu

Principle Investigator

Department of Chemistry

The University of Kansas

Lawrence, Kansas 66045

April 1992

Prepared for
The U. S. Department of Energy
Under Grant No. DE-FG02-90ER14159

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, not any of their contractors, subcontractors, or their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately-owned rights.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

PROGRESS REPORT

GRANT NO. DE-FG02-90ER14159-2

TABLE OF CONTENTS

- I. Introduction**
- II. Work Completed and in Progress**
 - A. Microwave Driven Multiphoton Excitation Dynamics in Rydberg Atoms**
 - B. Nonadiabatic Geometric Phases of Multiphoton Transitions in Dissipative Systems and Spin-j Systems**
 - C. Nonperturbative Treatments of Atomic and Molecular Processes in Intense Laser Fields**
- III. References**
- IV. Table and Figures**
- V. Manuscripts Resulting from This Grant Period**
- VI. Conference Presentations**
- VII. Principal Investigator's Efforts**

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

I. Introduction

In the current grant period (FY 91, 92), significant progress has been made in several areas. Fifteen papers have been published or are in press. Given in the following section (Sec. II) is a summary of the major accomplishments and progress made in this period. Detailed discussion of individual subjects can be found in the reprints and preprints (listed in Sec. V) submitted to the Department of Energy.

II. Work Completed and In Progress

IIA. *Microwave-Driven Multiphoton Excitation Dynamics in Rydberg Atoms: Fast-Fourier-Transformation Time-Dependent Propagation Method*

The fast Fourier transformation (FFT) method is an attractive and increasingly popular technique¹ for numerically integrating the time-dependent Schrödinger equation without the use of basis-set expansion. It also allows the treatment of the continuum exactly. By combining the FFT algorithm along with the Floquet theorem, one has a powerful numerical method for solving the time propagation of the wavefunction in periodic fields. Recently we have explored the feasibility and reliability of the FFT method for treating the *Rydberg atom* dynamical problem.

We use the split-operator algorithm to propagate the wavefunction: ($\hbar = 1$)

$$\psi(t + \Delta t) \cong \exp[i(\hat{P}^2/4m)\Delta t] \cdot \exp[-i\hat{V}\Delta t] \cdot \exp[-i(\hat{P}^2/4m)\Delta t] \psi(t) + O[(\Delta t)^3] , \quad (1)$$

where the effect of applying the operator $\exp[-i(\hat{P}^2/4m)\Delta t]$ is evaluated in terms of discrete Fourier transforms and the FFT algorithm. The procedure is found to be reliable and efficient and provides detailed time-dependent dynamical information². We have applied the FFT method to the study of the time propagation of the wave function and coarse-grained Wigner density distribution⁴ of the problem of microwave-driven multiphoton excitation and ionization of Rydberg hydrogen atoms², a subject of considerable current interest in atomic physics and

quantum chaos.³

The Hamiltonian under consideration is the 1D atomic hydrogen (stretched in the field direction)³ in the presence of ac and dc fields, as is widely used in the study of this subject. The initial state is chosen to be at $n_0=72$, and 1024 grid points are used to discretize the coordinate space up to $25,000 a_0$. Fig. 1 displays an example of the development of the probability density $|\psi(x,t)|^2$ in space (x) and time (t). It shows that the electronic flux spreads and recollects (in x) as time advances with occasional leaking to the continuum. In addition, we have constructed and studied the coarse-grained Wigner density distribution⁴, providing a vivid visualization and further insight of the time-development of the quantum system in phase space (x,p).² We also calculated the multiphoton excitation probability by projecting $\psi(t)$ onto unperturbed hydrogenic states and averaged over the last few optical cycles. The result is found to be in good agreement with the recent experimental data of Bayfield et al⁵ (Fig. 2). This is actually the *first* calculation of microwave-driven Rydberg atom dynamics, taking into account the effect of continuum exactly. Earlier quantal studies of this problem have used primarily basis set expansion methods.³ Our preliminary study shows that the FFT time-dependent propagation method is a feasible and powerful numerical technique for probing strong-field Rydberg atom quantum dynamics. More refined time-dependent propagation procedures for 3D Rydberg atom dynamical studies will be developed and discussed in the renewal proposal.

IIB. Nonadiabatic Geometric Phases of Multiphoton Transitions in Dissipative Systems and Spin-j Systems

The study of geometric phases is a subject of much current interest in many branches of physical sciences. Our recent interest in this direction is focused on the extension of Berry's adiabatic geometric phase notion to *non*-adiabatic geometric phase, with particular attention to physical phenomena occurring in multiphoton transitions and nonlinear optical processes.

We have made theoretical developments along the following new directions: (i) We develop a generalized version of the Feynman-Vernon-Hellwarth geometric representation and a *biorthogonal* density matrix formalism^{6,7} for the description of the non-Hermitian Schrödinger equation. The theory is applied to the study of *complex* geometric quantum phases in dissipative systems. It is shown that the complex Aharonov-Anandan (AA) geometric phase is related to the complex solid angle enclosed by a complex Bloch vector trajectory $\vec{S}(t)$. General analytic formulas are obtained for the complex AA phase for a driven dissipative two-level system undergoing *multiphoton* Rabi floppings. (ii) We have recently shown that cyclic quantum evolution can be realized and the non-adiabatic Aharonov-Anandan (AA) geometric phase can be determined for any spin- j system driven by periodic fields.⁸ Two methods are extended for the study of this problem: the generalized spin-coherent-state technique and the Floquet quasienergy approach. Using the former approach, we have developed a *generalized* Bloch-sphere model and presented a SU(2) Lie-group formulation of the AA geometric phase in the spin-coherent state. We show that the AA phase is equal to j times the solid angle enclosed by the trajectory traced out by the tip of a generalized Bloch vector. General analytic formulas are obtained for the Bloch vector trajectory and the AA geometric phase in terms of external physical parameters. In addition to these findings, we have also approached the same problem from an alternative but complimentary point of view without recourse to the concept of coherent state terminology. Here we first determine the Floquet quasienergy eigenvalues and eigenvectors for the spin- j system driven by periodic fields. This in turn allows the construction of the time-evolution propagator, the total wave function, and the AA geometric phase in a more general fashion.

In both cases, general analytical expressions have been obtained which can be tested by experiments.

IIC. *Nonperturbative Treatments of Atomic and Molecular Processes in Intense Laser Fields*

IIC1. Level Shifts and Widths of Excited States of Atoms in Strong Fields

Recent experiments on multiphoton ionization (MPI) and above-threshold ionization (ATI) of atoms in strong fields show significant energy shifts and broadenings of ATI peaks⁹. Further, for laser pulse widths of less than 1 psec, the electron energy spectrum can exhibit fine structure in the individual ATI peak⁹. These observations suggest that the structure of excited states plays a significant role in determining the properties of ATI. This prompts the necessity of determining the excited-state atomic energy-level structure in the presence of strong fields.

We have recently performed new nonperturbative results of intensity-dependent *complex* quasi-energies (energies and widths) for several low-lying excited states of atomic H in strong fields¹⁰, using the L^2 non-Hermitian Floquet matrix technique (Table I shows an example). Furthermore, we have developed a new nonperturbative L^2 -technique for the accurate treatment of ac Stark shifts of *arbitrary* excited states.¹⁰ We found that all the Rydberg states in weak fields are upshifted and closely follow the quadratic field dependence described by the pondermotive potential $e^2 F^2 / 4m\omega$. Large deviations from the pondermotive shift and intricate level upshift behaviors, however, occur in strong fields.¹⁰

II.C.2. Multiple High-Order Harmonic Generation of Atoms in Intense Laser Fields

Recently high-order (up to order 33) harmonic generation has been observed when atoms are irradiated by intense ($10^{13} - 10^{15}$ W/cm²) pulsed laser fields.¹¹ The experimental data show several unexpected novel nonlinear optical phenomena which cannot be understood from the perturbation theory. To probe the underlying mechanisms responsible for multiple high-order harmonic generation, we have recently performed a classical nonperturbative treatment of the electron motion under the influence of both the Coulomb and laser fields.¹⁰ We show that the spectral analysis of classical trajectories can provide detailed insights regarding the nonlinear

interaction of atoms and laser fields. Furthermore, we have identified a general type of classical trajectory whose power spectrum mimics all the essential experimental observations.¹⁰

More recently we have extended a fast Fourier transformation - split operator technique, in 3D spherical coordinates, to the solution of the time-dependent wave functions of atomic hydrogen in intense laser fields.¹² We performed a quantal study of the high-order harmonic generation (HG) spectra of atomic hydrogen at 248 nm based on the expectation values of the induced dipole moment and acceleration.¹³ The power spectrum in dipole form is obtained from the Fourier transform of the expectation value of the time dependent induced dipole $\langle z(t) \rangle$. Similarly, the photoemission spectrum in acceleration form is determined by Fourier transforming the expectation value of the acceleration $\langle \ddot{z}(t) \rangle$. Here the mean acceleration is obtained by means of Ehrenfest's theorem,

$$\langle \ddot{z}(t) \rangle = \langle -\partial \hat{H} / \partial z \rangle = -\langle z/r^3 \rangle - E(t), \quad (2)$$

where $E(t) = Ff(t) \sin \omega t$, F is the electric field amplitude and $f(t)$ the pulse shape of the laser field. It is found that while the dipole and acceleration forms yield equivalent power spectrum at weaker fields, significant difference occurs at intense fields. An example is given in Figure 3 which shows the dipole and acceleration power spectra for a stronger field $I=7.0 \times 10^{14}$ W/cm² and at $\lambda = 248$ nm. The acceleration photoemission power spectrum (solid line) contains well resolved high-order HG peaks up to 37th order, a feature similar to those observed in the rare gas experiments. However, the dipole power spectrum (dotted line) contains a spurious background and all higher-order HG peaks are completely masked. Floquet calculation at this intensity shows large ionization width ($\Gamma/2 \approx 1.657 \times 10^{-2}$ a.u.).¹² The spurious background in the dipole spectrum can be attributed to the improper use of the dipole moment which is not vanishing at large times when ionization occurs.¹³ Previous classical calculations indicated that HG can occur only when the electron is close to the nucleus and any asymptotic

value of the dipole moment is not relevant.¹⁰ The acceleration form Eq. (2), yields the correct result since it take into account automatically and self consistently the fact that the dominant contribution to HG arises from the portion of wavefunction where the electron experiences the largest force.

II.C.3. Multiphoton and Above-Threshold Dissociation of Small Molecules in Intense Laser Fields

We have recently developed a new *complex-scaling Fourier-grid Hamiltonian* (CSFGH) method^{14,15} for accurate and efficient determination of atomic or molecular resonance states *without* the need of using basis set expansions. The method requires neither the computation of potential matrix elements nor the imposition of boundary conditions, and the eigenvectors provide directly the values of the resonance wave functions at the space grid points. The procedure is particularly valuable for excited-state problems where basis set expansion methods face the challenge. Further the method is far more efficient and accurate than the finite difference methods as demonstrated in the problem of tunneling in anharmonic oscillator.¹⁴ The simplicity and usefulness of the CSFGH method is further demonstrated by a case study of the intensity-dependent *complex* quasi-vibrational energy eigenvalues (E_R , $-\Gamma/2$) and eigenvectors associated with multiphoton and above-threshold dissociation of H_2^+ ions in the presence of intense laser fields ($I = 10^{12} - 10^{14}$ W/cm²).¹⁵ For example, using only 51 evenly distributed grid points, we were able to achieve at least 4 significant digits of accuracy for both low-lying and highly excited quasi-vibrational states ($v=0,1,2,---,16$).¹⁵

Recent advances in high-intensity experiments on multiphoton and above-threshold ionization (MPI/ATI) of atoms⁹ have stimulated growing new interest in the study of multiphoton dynamics in diatomic molecules. In particular, experiments on multiphoton dissociation (MPD) of molecular hydrogen¹⁶ have revealed two notable intense-field phenomena: (a) The molecules

can absorb more photons than necessary to dissociate the chemical bond and the kinetic-energy distribution of the dissociated fragments show the appearance of equally-spaced multiple peaks, a phenomenon termed as above-threshold dissociation (ATD); (b) The H_2^+ chemical bond can be "*softened*" in the presence of intense fields, leading to efficient photodissociation of *low-lying* vibrations levels¹⁶. The presence of additional internuclear degrees of freedom in molecules enriches greatly the nonlinear multiphoton dynamics.

Most recently, we further extend the CSFGH technique and investigate the laser-induced resonance structure and multiphoton dynamics of H_2^+ molecules. We report a novel new high-intensity phenomenon: the "*hardening*" of molecular bonds and laser-induced *stabilization* of molecules in intense fields.¹⁷ It is found that, with increasing laser intensity, the molecular structure is delicately distorted, and the vibrational quasi-energy resonances can break into two or more groups separated from each other (see Fig. 4 for an example) each of which possessing different dynamical behavior. In particular, the lifetimes of *high-lying* resonances increase with increasing laser intensity, and the molecules become more stabilized in strong fields (Fig. 5), a new phenomenon which may be termed as "bond hardening". This is in contrast to the behavior of low-lying states, which due to "bond softening",¹⁶ become more and more unstable against dissociation in intense fields. Further work in this direction will be discussed in the renewal proposal.

III. References

1. See, for example, M. D. Feit, J. A. Fleck, Jr., and Steiger, J. Comput. Phys. 47, 412 (1982); R. Kosloff and D. Kosloff, J. Chem. Phys. 79, 1823 (1983).
2. T. F. Jiang and S. I. Chu, Comput. Phys. Comm. 63, 482 (1991).
3. For a recent review, see G. Casati, B. V. Chirikov, D. L. Shepelyansky, and I. Guarneri, Phys. Rep. 154, 77 (1987).
4. See, for example, S. J. Chang and K. J. Shi, Phys. Rev. Lett. 55, 269 (1985).
5. J. E. Bayfield and D. W. Sokol, Phys. Rev. Lett. 61, 2007 (1988).
6. S. I. Chu, Z. C. Wu and E. Layton, Chem. Phys. Lett. 157, 151 (1989).
7. E. Layton and S. I. Chu, Radiation Effects and Defects in Solids, (Gordon and Breach), Vols. 122-123, pp. 73-89 (1991).
8. E. Layton, Y. Huang, and S. I. Chu, Phys. Rev. A 41, 42 (1990).
9. For a recent review, see, R. R. Freeman and P. H. Bucksbaum, J. Phys. B 24, 325 (1991).
10. S. I. Chu, K. Wang, and E. Layton, JOSA B 7, 425 (1990).
11. For a recent review, see, A. L'Huillier, K. J. Schafer, and K. C. Kulander, J. Phys. B 24, 3315 (1991).
12. T. F. Jiang and S. I. Chu, "High-Order Harmonic Generation in Atomic Hydrogen at 248 nm: Dipole vs Acceleration Spectrum," Phys. Rev. A (submitted).
13. The power spectrum based on the acceleration form is recently advocated by K. Burnett et al., Phys. Rev. A 45, 3347 (1992) in a one-dimensional model calculation.
14. S. I. Chu, Chem. Phys. Lett. 167, 155 (1990).
15. S. I. Chu, J. Chem. Phys. 94, 7901 (1991).
16. See, for example, P. H. Bucksbaum et al., Phys. Rev. Lett. 64, 1883 (1990).
17. G. Yao and S. I. Chu, "Laser-Induced Molecular Bond Hardening and Stabilization in Intense Fields," Phys. Rev. Lett. (submitted).

IV Figure Captions

- Fig. 1. Time-development of the probability density $|\psi(x,t)|^2$ for 1D atomic hydrogen subject to d.c. ($F_{dc} = 6.5$ V/cm) and microwave ($F_{ac} = 0.018/N_0^4$ (a.u.) fields at frequency $\omega = 13.0$ GHz. The atom is initially ($t=0$) prepared at the quantum state $N_0=72$.
- Fig. 2. Population versus quantum number n . Solid curve: experimental results (ref. 5); dots joined by solid lines: FFT results. Parameters same as Fig. 1.
- Fig. 3. The dipole (dotted line) and acceleration (solid line) power spectra of atomic H at $\lambda = 248$ nm and $I = 7.0 \times 10^{14}$ W/cm². At this intensity, significant ionization occurs before 20 optical cycles.
- Fig. 4. Vibrational quasi-energy level structure and dressed adiabatic potential of H_2^+ at $\lambda = 266$ nm and (a) $I = 10^{11}$ W/cm² (weak field) and (b) $I = 10^{14}$ W/cm² (strong field). It is seen that at strong fields, the vibrational quasi-energies break into two groups. The resonance states in the upper group are supported mainly by the upper adiabatic potential well labeled $|g,n=0\rangle$. These states become more stabilized at stronger fields (see also Fig. 5).
- Fig. 5. Photodissociation (half-) widths ($\Gamma/2$) of vibrational quasi-energy states (labeled by v') of H_2^+ for (a) $I=10^{11}$ W/cm² (weak field) and (b) $I=10^{14}$ W/cm² (strong field) at $\lambda = 266$ nm. Notice that the upper group resonance states become more stable and longer lived as the laser intensity increases, a phenomenon which may be termed as "bond hardening".

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

V. Manuscripts Resulting From This Grant Period

1. S. I. Chu, K. Wang, and E. Layton, "Nonperturbative treatments of level shifts of excited states and high-order harmonic generation in strong fields," *J. Opt. Soc. Am. B* **7**, 425-432 (1990).
2. E. Layton, Y. Huang, and S. I. Chu, "Cyclic quantum evolution and Aharonov-Anandan geometric phases in SU(2) spin-coherent states," *Phys. Rev. A* **41**, 42-48 (1990).
3. S. I. Chu, "Complex-Scaling Fourier-Grid Hamiltonian Method for Resonance State Problems," *Chem. Phys. Lett.* **167**, 155-157 (1990).
4. S. I. Chu, "Quantum Dynamics and Fractal Character of Quasi-Energy States in Multi-Color Fields," *Comments Atomic and Molecular Physics* **25**, 101-112 (1990).
5. T. F. Jiang and S. I. Chu, "Nonperturbative Methods for the Study of Multiphoton Dynamics in Intense Periodic and Multi-Color Laser Fields," *Computer Phys. Comm.* **63**, 482-493 (1991).
6. I. M. Hammoud and S. I. Chu, "Irregular Wavefunction Behavior in Diamagnetic Rydberg Atoms: A Dynamical SO(4,2) Group Study," *Chem. Phys. Lett.* **182**, 63-68 (1991).
7. E. Layton and S. I. Chu, "Generalized Fourier-Grid Hamiltonian Approach to the Dirac Equation," *Chem. Phys. Lett.* **186**, 100-106 (1991).
8. S. I. Chu, "Complex Quasi-Vibrational Energy Formalism for Intense-Field Multiphoton and Above-Threshold Dissociation: Complex-Scaling Fourier-Grid Hamiltonian Method," *J. Chem. Phys.* **94**, 7901-79-09 (1991).
9. S. I. Chu, "Generalized Floquet Theoretical Methods for Nonperturbative Treatments of Schrödinger and Liouville Equations in the Presence of Strong Fields," Radiation Effects and Defects in Solids, (Gordon and Breach), Vols. 122-123, pp. 57-71 (1991).
10. E. Layton and S. I. Chu, "Nonadiabatic Geometric Phases of Multiphoton Transitions in Dissipative Systems and Spin-J Systems," Radiation Effects and Defects in Solids, (Gordon and Breach), Vol. 122-123, pp. 73-89 (1991).
11. G. Yao and S. I. Chu, "Strong Field Effects in Above-Threshold Detachment of a Model Negative Ion," *J. Phys. B* **25**, 363-376 (1992).
12. S. I. Chu, "The Complex-Scaling Coupled-Channel Methods for Atomic and Molecular Resonances in Intense External Fields," to appear in Applied Many-Body Methods in Spectroscopy and Electronic Structure, edited by D. Mukherjee (Plenum, NY, 1992).

13. G. Yao and S. I. Chu, "Complex-Scaling Fourier-Grid Hamiltonian Method. III. Oscillatory Behavior of Complex Quasi-Energies and the Stability of Negative Ions in Superintense Laser Fields," Phys. Rev. A (to appear in May 1, 1992 issue).
14. T. F. Jiang and S. I. Chu, "High-Order Harmonic Generation in Atomic Hydrogen at 248 nm: Dipole vs Acceleration Spectrum," Phys. Rev. A (submitted).
15. G. Yao and S. I. Chu, "Laser-Induced Molecular Bond Hardening and Stabilization in Intense Fields," Phys. Rev. Lett. (submitted).

VI. Conference Presentations of Works Completed Under This Grant Period (1990-92)
(I. T. - Invited Talk, C. P. - Contributed Paper)

1. First International Conference on Coherent Radiation Processes in Strong Fields. Washington, D. C., June 18-22, 1990. Invited Talk. "Generalized Floquet Theoretical Approaches to Intense-Field Multiphoton and Nonlinear Optical Processes."
2. Summer School of Nonlinear Optics and Nonlinear Dynamics. (Institute of Atomic and Molecular Sciences, Academia Sinica, Taipei, Taiwan. Aug. 3 - August 17, 1990.) Co-organizer of the Summer School. Delivered a series of lectures (12 hours) on nonlinear optics at intense laser fields and classical and quantum chaos.
3. Workshop on Time-Dependent Phenomena in Atomic, Molecular, and Optical Physics. Harvard University, Cambridge, Massachusetts, March 1 - 3, 1991, Invited Talk. "Multiphoton Dynamics in Intense Periodic and Multi-Color Laser Fields."
4. APS - DAMOP National Meeting, Washington, D. C., April 22 - 24, 1991. C.P.
 - a. "Complex-Scaling Fourier-Grid Hamiltonian Method for Multiphoton Resonances"
 - b. "Fast-Fourier-Transformation Time-Dependent Propagation Method for Microwave-Driven Rydberg Atom Dynamics" (with T. F. Jiang).
 - c. "Nonadiabatic Geometric Phases in Multiphoton and Nonlinear Optical Processes" (with E. Layton and Y. Huang).
5. First International Conference on Atomic and Molecular Dynamics (Satellite of XVII ICPEAC), Taipei, Taiwan, July 5 - 8, 1991. Plenary Lecture. "Recent Theoretical Development for Nonperturbative Treatments of Atomic and Molecular Processes in Strong Fields."
6. XVII ICPEAC, Brisbane, Australia. July 10 - 16, 1991. C.P.
 - a. "Complex-Scaling Fourier-Grid Hamiltonian Method for Multi-channel Atomic and Molecular Resonances: Application to Multiphoton and Above-threshold Dissociation of Molecules in Intense Laser Fields."
 - b. "Nonperturbative Treatments of Atomic Multiphoton Processes in Intense Laser Fields." (with K. Wang, E. Layton, Y. Huang, and T. F. Jiang).
7. APS - DAMOP National Meeting, Chicago, IL, May 19 - 22, 1992. C.P.
 - a. "Multiphoton Detachment of H^- Ions," (with C. Laughlin).
 - b. "Oscillatory Behavior of Photodetachment Rates and Stabilization of Negative Ions in Superintense Laser Fields," (With G. Yao).
 - c. "Time-Dependent Propagation Methods for the Studies of Multiphoton Dynamics and Stability of Atoms in Intense Laser Fields," (with T. F. Jiang).
 - d. "Eigenstates of Chaotic Diamagnetic Rydberg Atoms in the Positive Energy Regime," (with Y. Huang and J. Wang).
 - e. "Generalized Fourier-Grid Hamiltonian Approach to the Dirac Equation: Variational Solution without Basis Set," (with E. Layton).

VII. PRINCIPAL INVESTIGATOR'S EFFORTS

During this grant period, the effort devoted to this contract by the principle investigator (S. I. Chu) has been at least the amount as indicated in the contract. During the remainder of the contract period for this year, the effort devoted by the principle investigator is expected to continue at least the amount indicated in the contract (20% academic year, 100% summer time for 2.5 months).

**DATE
FILMED**
7/23/92