Progress/Technical Report

DOE/ER/40214--1

DE92 016181

RESEARCH IN THEORETICAL PARTICLE PHYSICS

The U.S. Department of Energy
Agreement No. DE-FG02-85ER40214

May 1, 1991 - April 30, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Douglas W. McKay, Principal Investigator
Herman Munczek, Principal Investigator
John Ralston, Principal Investigator

The University of Kansas
Lawrence, KS 66045

May 1992

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *eb*

II. Technical Progress Report for "Research in Theoretical Particle Physics", 1991-1992

Publications

Published

1. "The Schwinger-Dyson Equation in QCD: Comparison of Some Approximations," H. Munczek and D. McKay, *Phys. Rev. D* **42**, 3548 (1990).
2. "A Low Energy Technicolor Lagrangian with Vector Mesons," R. Johnson, B.-L. Young, and D. McKay, *Phys. Rev. D* **42**, 3855 (1990).
3. "Patterns of Small-x QCD", D. McKay and J. Ralston, *Nuclear Physics B (Proc. Suppl.)* **18C**(1990) 86-91.
4. "Techni-Vector Meson Dominance and W, Z Boson Properties," R. Johnson, B.-L. Young and D. McKay *Phys. Rev. D* **43**, R17 (1991).
5. "Comparing Coherent Microwave Emission From LPM and B-H Showers," J.P. Ralston and D.W. McKay, AIP Conference Proceedings 220, *High Energy Gamma-Ray Astronomy*, (Ann Arbor 1990) Ed. J. Mathews (AIP, New York 1991) p. 295.
6. "Calculation of the Pion Decay Constant in the Framework of the Bethe-Salpeter Equation," P. Jain and H. Munczek, *Phys. Rev. D* **44**, 1873 (1991).
7. "Poincare Algebra and Soliton Energy in Kink Field Theory," P. Jain, *Mod. Phys. Lett. A* **6**, 2363 (1991).
8. "Applications of Semiclassical Techniques to Sine-Gordon Theory: Re-Examined," P. Jain *Mod. Phys. Lett. A* **6**, 2509 (1991).
10. "Hollow Galactic Halos of Fermionic Dark Matter," J.P. Ralston and L.L. Smith, *Astrophys. J.* **367**, 54 (1991).
11. "Color Transparency: A Manifestation of Nuclear Filtering," B. Pire and J.P. Ralston, in *Les Houches Winter School in Physics: Hadronic Physics with Multi-GeV Electrons* (B. Desplanques and D. Goutte, eds) (Nova, NY 1991).
12. "Duality Symmetry and Power Law Fading of Frustration in a Quantum Multiconnected Superconductor," S.N. Sun and J.P. Ralston, *Phys. Rev. B* **43**, 5375 (1991).
13. "Quantum Color Transparency," J.P. Ralston and B. Pire, *Phys. Rev. Lett.* **65**, 2343 (1990).

14. "Hard Scattering in a Nuclear Environment," B. Pire and J.P. Ralston, *Nuc. Phys.* **A525** (419c) 1991, proceedings of *Quark Matter 90*, edited by J.P. Blaizot, *et al.*
15. "A Transparency Scaling Law for Large Q^2 Exclusive Reactions on Nuclei," B. Pire and J.P. Ralston, *Phys. Lett.* **B256**, 523 (1991).
16. "On the Time Scale Issue in Quantum Color Transparency," (published as "Ralston and Pire Reply"), J.P. Ralston and B. Pire, *Phys. Rev. Lett.* **67**, 2112 (1991).
17. "Spin and the Independent Scattering Mechanism," J.P. Ralston and B. Pire, in *Polarized Collider Workshop* (AIP Conference Proceedings No. 223) edited by J. Collins *et al.* (AIP, NY, 1991)
18. "Lepton Pair Production in Polarized Collider Experiments as a Method to Measure the Transverse Polarization of Quarks in the Proton," J. Cortes, B. Pire and J.P. Ralston, in *Polarized Collider Workshop*, *ibid.*
20. "An Upper Bound on the Higgs Mass," D.W. McKay, *Proceedings of "Beyond the Standard Model II,"* (Norman, OK, 1990) World Scientific (1991), K. Milton, Ed.
21. "Large Technicolor Effects at the Z^0 ," B-L. Young, R. Johnson and D.W. McKay, *Proceedings of "Beyond the Standard Model II,"* *ibid.*
22. "Lorentz Covariance of Quantum Fluctuations in Quantum Field Theory", P. Jain in *Proceedings of "Beyond the Standard Model II,"* *ibid.*
23. "Lorentz Covariance of Current Matrix Elements in a Soliton Model," P. Jain, *Phys. Rev.* **D 44**, 3982 (1991).
24. "Approach to Color Transparency in the Soliton Picture of the Nucleon," P. Jain, J. Schechter and H. Weigel, *Phys. Rev.* **D45**, 1470 (1992).
25. "Color Transparency and Electronuclear Physics" J.P. Ralston and B. Pire, to be published in *Proceedings of Second Workshop on Hadronic Physics with Electrons Beyond 10 GeV* (Dourdan, France 1990) *Nuc. Phys. A* **5332**, 155C (1991).
26. "Anomalous Color Transparency," J.P. Ralston, *Physics Letters B* **269**, 439 (1991).

Review Article

27. "Colour Transparency and Nuclear Filtering," B. Pire and J.P. Ralston, Feature Article in *Nuc. Phys. News* 1(No. 4) 23-26 (1991).

In Press

28. "The Possibility of Quenching of Integer Quantum Hall Behavior with Increasing Lattice Asymmetry," S.N. Sun and J.P. Ralston, *Physical Review B*.
29. "Theory of Color Transparency" B. Pire and J.P. Ralston, in *Proceedings of 5th Workshop on Perspectives in Nuclear Physics at Intermediate Energies* (Trieste 1991) (World Scientific).
30. "Update on Quantum Color Transparency", B. Pire and J.P. Ralston, in *Proceedings of European Physics Society/Lepton Photon International Meeting* (Geneva 1991).
31. "On the Validity of Approximations for the Calculation of the Pion Decay Constant", P. Jain and H. Munczewk, *Phys. Lett. B*.
32. "Relativistic Pseudoscalar $q\bar{q}$ Bound States: Bethe-Salpeter Wave Functions and Decay Constants," H. Munczek and P. Jain, *Phys. Rev.D*.

Submitted

33. "Measuring the Transverse Polarization of Quarks in the Proton," *Zeit. Phys. C*. (Ecole Polytechnique A048.0491 (1991)).

II.B. 1990-'91 NARRATIVE ACCOUNT OF RESEARCH

Dynamical Symmetry Breaking and Schwinger-Dyson Equation

If the fermion inverse propagator's "mass term" is dynamically generated, a necessary condition for spontaneous breaking of chiral symmetry is satisfied. The Schwinger-Dyson equation for the inverse propagator provides a framework for studying dynamical chiral symmetry breaking. Various modifications to the ladder approximation (where the bare fermion-gauge boson vertex is adopted) have been used by many authors over many years to model QED, QCD, technicolor and extended technicolor dynamical mass generation. In QCD and technicolor modeling, the most frequently used approximation¹ has been a modification of the gluon propagator to include the one-loop running coupling constant effects in the ladder approximation, and then to keep only the leading term in the expansion of the integrand in its angular variables. The latter approximation amounts to the replacement in the integrand:

$$\alpha_s((k-q)^2) \rightarrow \theta(k^2 - q^2)\alpha_s(k^2) + \theta(q^2 - k^2)\alpha_s(q^2)$$

where α_s is the strong interaction running coupling constant. This “standard” approximation has the attractive feature that the leading asymptotic behavior of the solutions to the Schwinger-Dyson equation can be readily obtained analytically and agree with the renormalization group and operator product expansion results.² However, if the low- q^2 part of the mass-function is important in applications of the solutions, or if sub-leading asymptotic behavior is important, then the approximation described above is insufficient.³

In work reported by Munczek and McKay⁴ in *Physical Review D* and presented by McKay at an Ecole Polytechnique theory seminar, the “standard approximation” to the Schwinger-Dyson equation was shown to have the wrong slope at $q^2 = 0$. New approximations which fixed this problem and which included successively improved treatment of the angular integration were developed. Improvements to the solutions ranging from 5% to 30% were obtained, depending on the q^2 value. The correct leading asymptotic behavior was retained in every case.

Munczek and McKay also included the two-loop QCD corrections to the running coupling in the Schwinger-Dyson equation and showed that the resulting corrections to the sub-leading asymptotic behavior are roughly 50% compared to the one-loop α_s results. Two-loops are needed if the subleading terms are to be properly handled.

The two-loop corrections to the *regular* solutions were computed for the first time, and the behavior of the irregular solutions to the Schwinger-Dyson equation with the two-loop renormalization group improved (running) mass⁵ were found to agree up to vertex corrections not included in the Schwinger-Dyson equation. The next step in the program of a *complete* ladder approximation, coupled fermion Schwinger-Dyson and Bethe-Salpeter equation program is described in the work of Jain and Munczek below.

Consistency Bound on the Minimal Model Higgs Mass

There has been considerable recent interest on bounds on the Higgs mass in the standard model. McKay noted that the Higgs boson mass is bounded from above by $M_H \lesssim 500 \text{ GeV}$ for $m_t \lesssim 200 \text{ GeV}$.⁶ The bound follows from the consistency requirement that the quartic scalar coupling constant, λ , be real at the one-loop level in the calculation of the effective potential.⁷ The bound reads

$$\left(\frac{M_H}{M_W}\right)^4 \leq \frac{16}{3} \left\{ \left[4\pi^2/3g^2 + (g^2/\pi^2) \left(\frac{M_t}{M_W}\right)^4 \right]^2 - \frac{1}{4}(2 + \sec^4 \theta_W) \right\}$$

where θ_W is the weak mixing angle, g the $SU_L(2)$ gauge coupling constant and M_H , M_W and M_t are the Higgs, the W -boson and the top quark masses, respectively. The bound obtained

is comparable to the "triviality bound" which follows from a new physics cutoff of several TeV. McKay reported a preliminary version of this work at the *Beyond the Standard Model* meeting held in Norman, Oklahoma.

Tests of Physics Beyond the Standard Model – Large Technicolor Effects in Z_0 Physics

The LEP precision measurements of Z_0 parameters now make it possible to test detailed predictions of new physics models. Young, Johnson and McKay^{8,9} showed that an estimate of technicolor effects using generalized vector dominance leads to large deviations from standard model values for M_W/M_Z and the b -quark decay width of the Z_0 unless the technicolor model parameters are tightly constrained:

$$5 < N_d N_T < 25 ,$$

for $130 < M_t < 220 \text{ GeV}$, where N_d is the number of techni-quark doublets and N_T is the number of technicolors deduced at the 90% confidence level (by comparison with experimental limits). The technicolor models to which the tests apply are those patterned closely after QCD, and our vector dominance assumption is only good to 10–20% in QCD physics of weak currents, but the limits are good enough to rule out large technicolor groups and multi-family models nonetheless.⁹

Particle Astrophysics

Ralston and McKay have investigated coherent microwave emission as a possible probe of neutrino charged current events in the ultra high energy (UHE) region upwards of 10^{14} eV . Such emission from showers initiated in rock or ice by UHE cosmic rays from point sources may be detectable with relatively inexpensive antennas.^{10–12} The issue was raised whether the multiple scattering process (Landau, Pomeranchuk, Migdal process (LPM)) predicted to set in when e^+ and e^- are very energetic and forward as in the UHE region will modify the microwave emission characteristics. It was shown that the charge imbalance ΔQ , and therefore the power emitted, should be largely independent of whether the shower had LPM or the usual Bethe-Heitler development.¹³ It was noted that power emitted into an angular region around the critical angle, δW in a frequency interval $d\omega$ ¹⁴

$$\delta W = \frac{(\Delta Q)^2}{c^2} L \left(1 - \frac{1}{\mu^2} \right) \omega d\omega$$

gets an enhancement from the longer range of shower development, L . The comparatively longer LPM shower may give some enhancement to the efficiency for converting energy into the coherent microwave component. This work was presented by McKay at the Michigan meeting on *High Energy Gamma Ray Astronomy*.

The Schwinger-Dyson and the Bethe-Salpeter Equations: Analysis and Consequences of Improved Treatment.

In association with the groups' postdoctoral associate P. Jain, Munczek continued research¹⁵ on the consistent treatment of the Schwinger-Dyson (SD) and the Bethe-Salpeter (BS) equations. For a long time¹⁶ it has been recognized that a simultaneous, nonperturbative, solution of these equations is necessary to examine the consequences of chiral symmetry breaking, whether spontaneous or explicit. Because of the calculational complexity of the equations, most studies resort to theoretical and numerical approximations whose validity is hard to assess. Improvements on this subject done during the current grant period include:

a) *Spontaneous chiral symmetry breaking and the calculation of the pion decay constant:* In a paper by Jain and Munczek¹⁷ the SD equation with zero bare mass was solved simultaneously with the BS equation for a pseudoscalar Nambu-Goldstone meson (expanded to first order in the pion momentum p). The quantities calculated were the two components of the quark propagator $S(q) = [\not{q}A(q^2) - B(q^2)]^{-1}$ and the four components of the Bethe-Salpeter wave function,

$$\chi(p, k) = \gamma_5 \chi_0(p, k) + \gamma_5 \not{p} \chi_1(p, k) + \gamma_5 \not{k} \chi_2(p, k) + \gamma_5 [\not{k}, \not{p}] \chi_3(p, k) .$$

The Landau gauge, one-gluon exchange, coupled equations were solved iteratively and without employing the usual approximation of writing the running QCD coupling α_s in the form $\alpha_s[(q-k)^2] = \theta(k^2 - q^2)\alpha_s(k^2) + \theta(q^2 - k^2)\alpha_s(q^2)$, an approximation shown to be unreliable by previous work by Munczek and McKay⁴ described in the previous section. Instead, the general form of the wave function was expanded in 4-dimensional spherical harmonics (Gegenbauer polynomials), truncated after 4 terms, and then integrated numerically. It was also shown by Jain and Munczek that the widely used approximation $A(q^2) = 1$ is unjustified in QCD. This was true for any of a variety of infrared behaviors of $\alpha_s(q^2)$ which were probed and which gave reasonable values for f_π , the pion decay constant. Fig. 1 shows some numerical results.

Fig. 1. Momentum dependence of propagator functions $B(x)$ and $A(x)$ obtained by solving the Schwinger-Dyson equations. The complete quark propagator is given by $S(q) = [A(x)\not{q} - B(x)]^{-1}$, where $x = -q^2/\Lambda_{QCD}^2$. Curves show the effects of different models of the gluon propagator.

The calculation of f_π was done by using an exact expression for f_π involving the components of the pion BS wave function. This was then compared with the usual Pagels and Stokar (PS) approximation to the expression for f_π [which sets $A = 1$ and approximates the BS wave function by dropping a variety of terms linear in the pion momentum]. We found that the major contributions to the value of f_π come from the low momentum or infrared region where $A(q^2)$ is approximately twice the naive value. Furthermore, we found that the Pagels-Stokar approximation systematically gave values of f_π several times larger than those of our numerically accurate calculations, thus showing that such an approximation is quantitatively unreliable.¹⁸

b) Explicit chiral symmetry breaking and relativistic calculation of light and heavy pseudoscalar bound states

The work of (a) was extended by Munczek and Jain¹⁹ to the general case in which there is an explicit mass term μ_a in the SD equation for the propagator of a quark of flavor a .

The coupled SD and BS equations were solved without resorting to nonrelativistic approximations and by extending the numerical methods of the previous paper.

Qualitatively it was found that the procedure gives a unified treatment of the light and heavy quark spectroscopy. For light pseudoscalars (would-be Nambu-Goldstone bosons) it was found that the mass M_{ab} of the bound states behaved like $M_{ab}^2 \cong (\mu_a + \mu_b)C$, with C only weakly dependent on flavor and in good numerical agreement with the current algebra result that $C \simeq \langle \bar{q}q \rangle / f_\pi^2$. On the other hand for heavy quarks the result is that M_{ab} is approximately proportional to the sum of the slowly varying heavy quark masses evaluated at zero momentum, in accord with nonrelativistic limit expectations. Another qualitative result found was that for all quarks the pseudoscalar vertex function behaves asymptotically like the regular solution of the SD equation, even in the presence of explicit chiral symmetry breaking. This result is in agreement with considerations based on renormalization group analysis and operator product expansion.²⁰

In order to obtain quantitative results a model for the gluon propagator was needed. The high momentum behavior of $\alpha_s(q^2)$ was taken to be the known two-loop renormalization group results, while the low momentum region was modelled by considering confinement and potential model ideas. With four parameters in $\alpha_s(q^2)$ and given the masses μ_a it was possible to fit, or predict reasonably, masses for all the ground states of pseudoscalar spectrum (η and η' were not considered at this stage). The leptonic constants were also calculated for all the states: using f_π as input to determine the parametrization of α_s , the prediction of f_k was good and the values of all the other f 's were well within experimental limits, consistent with predictions coming from sum rule, lattice and potential model calculations.²¹

c) Extensions of Explicit Chiral Symmetry Breaking to Vector Mesons

The Bethe-Salpeter equations for vector meson states have been obtained and are presently being set up for numerical work. This and other extensions are discussed in our research program.

The Interface Between Perturbative and Non-Perturbative QCD

Several topics are being studied by John Ralston with Bernard Pire in a collaboration of ten years. During 1990-91, Ralston was on sabbatical leave at Pire's institution (Ecole

Polytechnique) where a number of new projects were begun. The general area of the work is the interface between perturbative and non-perturbative QCD. Among the specific topics being studied are:

a) Color Transparency and Nuclear Filtering

The treatment of exclusive processes in QCD has often been controversial. In exclusive processes, there is the potential for non-short-distance effects to upset the perturbative treatment even at high energy and large momentum transfer (q^2). Another problem has been the comparatively modest values of q^2 which experiments have been able to probe, and the interesting question of the role of non-perturbative effects.

Color transparency is a new research area in which all of these issues arise and can be studied. In the original formulation of the idea by Brodsky and Mueller,²² a physical picture is invoked in which large q^2 selects quarks which are close together in space in the participating hadron. The spatial extent of quark wave functions in the coordinate transverse to the momentum of a hadron is usually about one fermi, but (the argument goes) only fluctuations of the wavefunction in which the quarks' transverse separation is of order $\sqrt{1/q^2}$ will actually participate. If this is true, a hadron in such a configuration should have a measurably reduced cross section for further strong interactions.

Considering the interactions in perturbative QCD, there will be destructive interference of gluon emission from a color-singlet hadron, with the soft gluons which threaten the consistency of the perturbative QCD cancelling. It is known^{23,24} that the perturbative cross section goes like the geometrical area of the hadron: thus, a small hadron will have a correspondingly small cross section. At the same time, we know there is much in the strong interactions that is non-perturbative. It is an interesting and testable assumption that non-perturbative processes will cancel along with the perturbative ones. If "color transparency", namely the reduced interaction cross section is observed, one will see that the "old fashioned" strong interactions can be turned off under certain circumstances.

One test involves measuring "hard" exclusive processes occurring in a large nuclear target. The nucleus then serves as a test medium. One can test that the interaction cross section of a hard-struck hadron is actually reduced by measuring a reduced attenuation in the nuclear medium. A fairly model - independent test of the proposal is to look at the ratio of an exclusive proton (p) cross section in the nuclear target to the analogous process in free space: an example

would be to study the c.m. energy (s) dependence of the ratio $T(s)$ given by

$$T(s) = \frac{d\sigma/dt(pA \rightarrow p'p''(A-1))}{d\sigma/dt(pp \rightarrow p'p'')} \Big|_{\vartheta} \quad (1)$$

where the c.m. scattering angle ϑ is fixed. This ratio should increase with increasing s , as a signal of color transparency. At least, this was the first expectation for experiments.

Ralston and Pire²⁵ had worked on exclusive process since 1982 with a slightly different approach. In our picture, the short distance “quark-counting” contribution was one possibility, but could not be the whole story. As a test of the relevance of exponentiating leading log series, including what is known as Sudakov corrections, we studied the independent scattering contributions to fixed-angle $pp \rightarrow pp$ scattering. The independent scattering or “Landshoff”²⁶ terms involve momentum flow which violates the assumptions of the quark-counting rules. Indeed, while two quarks are close together in space when they scatter, they are not necessarily close to another pair of quarks that scatter, so these terms in the amplitude represent hard scattering of spatially “large” objects. We showed that in this case primitive quark amplitudes acquired energy dependent imaginary phase shifts and identified these as the perturbative version of the Coulomb phase in QCD. We argued that the energy dependence of such phases were nevertheless perturbatively calculable and could be measured in interference. Simplifying to a “two-component” model, we fit the data for pp scattering, which shows rather large oscillations (of order 50% in magnitude) about a pure power-law decrease of the cross section with energy.

The question of the origin of the oscillations was certainly not settled by this work. Brodsky and DeTerra²⁷ proposed that certain charmed dibaryon thresholds could be the explanation. Our conjecture that the imaginary phases were calculable, and could be organized as purely imaginary anomalous dimensions²⁸ was recently derived at the hadron level by Botts and Serman.²⁹ However, their calculation obtained a phase coefficient which was much too small. This caused some doubt that the idea was right, until it was realized that perhaps the wrong thing had been calculated: the most rapidly varying phase dominates the oscillations, not the phase of the asymptotic amplitude. Botts³⁰ subsequently showed that the asymptotic calculation was not accurate in any case below the unreachable TeV c.m. energy region. Thus the general structure of our model has been upheld, while the details of calculating parameters from first principles are still open.

This work became important for our interpretation of the first color transparency experiment conducted by the group of Carroll *et al.*³¹ The experiment first showed an increase of the ratio T (eq. 1) with energy, as expected if color transparency were being observed, but then an

unexpected *decrease* set in above a lab energy of about 10 GeV. This caused perplexity at first, and some suggestions (including that the data might be wrong) were made to explain it. We proposed³² that the structure in the transparency ratio was caused mainly by the oscillations of the denominator. For the numerator, which involves the scattering inside the nuclear target, we argued that the independent scattering amplitudes had been reduced. This would be expected if the nucleus were playing a more active role than that of a test medium, but also was filtering away the spatially "large" amplitudes, enriching the short-distance terms. We called this concept "nuclear filtering"; it turned out to be related to an earlier proposal for diffractive processes by Bertsch *et al.*³³

Our calculation based on attenuating the independent scattering terms reproduced the energy dependence of the ratio T with no free parameters. Moreover, we predicted that the oscillations would continue in a specified way with higher energy. In arriving at this result, we arranged the calculation to avoid model-dependence due to the nuclear physics; thus the energy dependence at fixed nuclear number A , but not the A dependence, was calculated, as shown below.

Fig. 2 (a) The energy dependence of $s^{10} d\sigma/dt(pp) |_{90^\circ}$ for high energy pp elastic scattering at 90° cm angle. (b) Prediction of oscillating transparency $T(s)$ for $A = 27$ after varying over all possible nuclear phases (upper and lower limits are shown); data from A.S. Carroll *et al.*

In subsequent work³⁴⁻³⁷ we elaborated several corollaries of the concept of nuclear filtering, and worked toward making the whole framework better defined and completely quantum mechanical. We introduced the concept of a "mini-hadron" to describe a state with the quantum numbers of a hadron, but with the quarks close together in space, and in a minimal Fock space

representation (three quarks for a proton, a quark anti-quark pair for a meson). We suggested some spin tests of the ideas, based on the logic that either the spin puzzles in exclusive processes are in the short distance sector (and then perturbative QCD is not being applied correctly) or associated with long distance processes (and then capable of being filtered away) . There will be more discussion of exclusive spin puzzles below. We also studied the case of exclusive electroproduction, where quantities which act like the electromagnetic form factor of protons inside the nucleus can be measured. These generally are not affected by the independent scattering problem and should show transparency as naively expected. We considered the possibilities of pion-knockout, and baryonic resonance production inside the nuclear targets. These results were presented at several meetings, including the 1989 *Rencontre de Moriond*, the 1989 *International Conference of the European Physical Society, Quark Matter '90*, and the 1990 *Les Houches Winter School of Physics*.

A formalism to describe “mini-hadrons” and the logic behind color transparency was published in *Physical Review Letters*.³⁸ We will review it briefly here to further define the discussion. Let $\psi(\vec{k}_T; x)$ be the light-cone Bethe-Salpeter wave function to find quarks with transverse momentum k_T and longitudinal momentum fraction x in a hadron. The amplitude M for hadrons to scatter exclusively can be expressed as an integral over the wavefunctions with a primitive scattering kernel $H(k_{Ti}, x_i)$, and is given by

$$\begin{aligned} \int \prod_{i,j} \psi_i^*(k_{Ti}; x_i) H(k_{Ti}, k_{Tj}, x_i, x_j) \psi(k_{Tj}; x_j) dx_i dx_j d^2 k_{T,i} d^2 k_{T,j} \\ \simeq \prod_{i,j} \int dx_i dx_j H(0, 0; x_i, x_j) \int_0^q d^2 k_{T,i} \psi_i^*(k_{T,i}; x_i) \int_0^q d^2 k_{T,j} \psi_j(k_{T,j}; x_j) \end{aligned} \quad (2)$$

where in the second step it is assumed that H is smooth as the transverse momentum goes to zero. The exceptions to this due to the independent scattering contributions will be discussed below. In (2) we see the emergence of the quark “distribution function” $\varphi(q^2, x)$, a quantity introduced by LePage and Brodsky³⁹ to set up the short distance dominance of the exclusive scattering. We proceed using the Fourier representation of the wave function in transverse separation (b_T) space, integrating to find

$$\begin{aligned} \varphi(q^2, x) &\equiv \int_0^q d^2 k_T \psi(k_T; x) \\ &= 2\pi q \int_0^\infty db_T b_T J_1(qb_T) \tilde{\psi}(\vec{b}_T, x) \end{aligned} \quad (3)$$

where

$$\tilde{\psi}(\vec{b}_T, x) = \int d^2 k_T e^{i\vec{b}_T \cdot \vec{k}_T} \psi(\vec{k}_T, x) . \quad (4)$$

The short distance dominance is more clear with (3), since the Bessel function selects the region $b^2 = 1/q^2$ so long as the wave function is smooth as $b^2 \rightarrow 0$.

We generalize to the nuclear case by modifying the procedure to allow for filtering of hadron by the nucleus. As the separation b of the quarks goes to zero, we use the perturbative cross section going like b^2 ; for larger b , we suppose some smoothly increasing cross section function of b^2 until the total cross section is reached. This is where non-perturbative effects set in. For the nuclear case we make an ansatz for the filtered distribution amplitude function with the formula

$$\varphi_A(q^2, x) = 2\pi q \int_0^\infty db_T b_T J_1(qb_T) f_A(b_T^2) \tilde{\psi}(\vec{b}_T, x) ; \quad (5)$$

$$f_A(b_T^2) \sim 1 - \lambda b_T^2 + \dots \quad (6)$$

where $f_A(b)$ can be related to the scattering amplitude of the struck hadron in the nuclear medium.³⁸ Now, by restricting our assumptions to (a) a factorized form for the amplitude such as (2), and (b) a cross section for interaction beginning with the power series (6), we were able to give a quantum mechanical demonstration of the semi-classical expectation of color transparency. Moreover, a new result is that the interaction with the nucleus provides an effective infra-red cutoff: filtering itself motivates the short distance argument even when q^2 may not be sufficiently high to make the perturbative argument entirely convincing (the case argued by Isgur and Llewellyn Smith⁴⁰). In effect, we claim that certain processes in a nuclear target are a cleaner QCD problem than the analogous process in free space.

These results were also useful in obtaining the next major step, which we call transparency scaling.

Transparency scaling is a property obtained from (5) assuming that the wave function is smooth as $b \rightarrow 0$. In fact, with the renormalization group one can show there are logarithmic singularities in the wave function in perturbative QCD, but these are small effects: scaling attempts to control rapidly varying dependence involving powers of $1/q^2$ and powers of A . Scaling addresses the problem of large q^2 scattering in a large nucleus, in which case there was a shortcoming in the original arguments: one cannot neglect powers of $1/q^2$ (in taking the high energy limit) if they increase with A .

Since there are two large dimensionful scales, namely the large q^2 and the large nuclear size, we looked for dependence on any dimensionless ratio of scales. From (5) we obtained a scaling law that at large q^2 and large A , the transparency ratio measured in the lab should be a function of the "scaling variable" $q^2/A^{1/3}$. This was published⁴¹ in *Physics Letters B*, and

presented in invited talks at several conferences, including the *Dourdan (France) Meeting*, the *Penn State Collider Physics Meeting*, the *5th Trieste Workshop*, and the *1991 Joint European Physical Society-International Lepton Photon Meeting* at CERN.^{42,43,44}

We anticipate that the transparency scaling law will be tested in upcoming experiments at SLAC (NE18) and Brookhaven . The Brookhaven experiment is especially exciting, because the construction of a new detector (known as EVA) custom-made to measure color transparency, along with other improvements, will increase data by 200-400 times compared to the first experiment. We suggest that it is more fundamental to test transparency scaling itself than to test the value of quantities such as the transparency ratio. There is much less model-dependence in the scaling law than in a calculation of, say, a particular cross section at a particular energy in a particular nucleus. Nevertheless, we are not certain the scaling law will apply to the experiment: the law itself serves to test the importance of the so-called "expansion time" question in the high energy limit.

The expansion time question arises when one tries to model in detail the time-evolution of the struck mini-hadrons and their interaction with the nucleus. Our own work has been conservative, starting with specific properties of leading order perturbative QCD and then working down in energy and up in complexity to describe the nuclear scattering case. Other workers^{43,44} have taken a complementary approach, in which a struck mini-hadron begins expanding immediately after the hard scattering, and therefore interacts with the nuclear test medium quickly. There is probably some truth in such a picture but there is uncertainty in the dynamical details. One realizes the need to separate out the dependence of a third scale – the energy of the hadron – which is kinematically different in its effects from the large q^2 . In obtaining our scaling law we assumed that the Lorentz boost is sufficiently big to time-dilate details of hadron expansion into unobservability. If the time scale is about the inverse of a few hundred MeV, and the boost has a Lorentz factor of order 10, the assumption should be good. Still there is controversy whether this assumption will be true at the energies of upcoming experiments.^{45,46} To separate the issues, Ralston has suggested a different experiment⁴⁷ (described below) involving π^0 photoproduction in which the question arises in a very interesting way.

A review of our work up to mid-1991 was commissioned and published by *European Nuclear Physics News*, a new periodical designed to function for the progressive nuclear physics community much as *Cern Courier* does for high energy physics. Besides the other papers already mentioned, a review was invited and is being written for the *International Journal of Modern Physics A*. During 1990-91, Ralston visited for three months to discuss these issues with the

experimental nuclear group (SEPN) of the *Saclay Nuclear Research Facility* in France. Ralston also gave invited talks at *Saclay*, at the *Rutherford Lab Christmas Meeting*, the *Saclay Meeting on Electron Physics Above 15 GeV*, as well as presenting seminars on the subject at *Zaragoza Universidad* (Zaragoza, Spain), the *Universite de Marseille* (Luminy, France), the *Universite Claude Bernard* (Lyon, France) and the *Universiti di Milano* (Milan, Italy). Ralston was discussion leader and session chairman for the *Session on Color Transparency of the 1991 Gordon Conference* held in Tilton, NH.

b) *Measuring Transverse Quark Polarization*

The question of the transverse spin structure of quarks in the parton model is an old one, but still not completely clear in the literature. A number of misconceptions were based on Feynman's treatment⁴⁸ of the higher twist structure function g_2 , measurable in inclusive electroproduction. This treatment itself would not be called "wrong", inasmuch as it predates QCD and is reasonably self consistent. However, the attempt to define transversely polarized parton distributions in terms of g_2 leads to inconsistencies. Subsequently there were (wrong) statements in the literature that all transverse spin effects are "higher twist" and impossible to describe in parton language. It turns out that the transverse polarization distributions simply represent a fundamental degree of freedom that has never been measured, and can make contributions at leading order to certain experiments. This fact in itself is not new⁴⁹ but its time may have come.

Renewed interest in the general topic of the high energy spin structure of the proton is due to the fact that the technology to measure such structure functions finally exists. The EMC measurement of the Ellis Jaffe sum rule has gathered much attention, and we anticipate there will be further measurements of spin quantities such as g_2 . This makes an updated effort to clarify these issues worthwhile.

In the field-theoretic expression of the parton model, distributions are defined in terms of certain correlation functions of the quark operators $\psi(x)$ in the proton state. The correlators which have a factor of the big momentum "P" in the high energy limit are leading order. The complete list of these is

$$\int \frac{x P^+}{2\pi} dy^- e^{iz P^+ y^-} \langle P, S | \bar{\psi}_\beta(0) \psi_\alpha(0, 0, y^-) | PS \rangle =$$

$$\left\{ x P(x, Q^2) \not{P}/2 - x \lambda h_L(x, Q^2) \not{P} \gamma_5 /2 - x h_T(x, Q^2) \gamma_5 \not{P} \not{S}_T /2 \right\}_{\alpha\beta}$$

where S_T is the transverse spin of the proton and λ twice its helicity. The last term has the interpretation of the *transverse polarization distribution* of the quarks in a transversely polarized proton. It is not the same function that is measured in deeply inelastic scattering with g_2 . However, in certain experiments where there are two transversely polarized hadrons, the distribution contributes at “leading twist”. An example is Drell-Yan lepton pair production: then one can measure the angular distribution of the pair, to find a term bilinear in the h_T distributions:

$$\frac{d\sigma}{d^4Q d\Omega} = \sum_a \frac{\alpha^2 e_a^2}{2Q^2 s} h_T^{a/A}(x_A) h_T^{\bar{a}/B}(x_B) \sin^2 \vartheta \cos(2\phi - \phi_A - \phi_B) + A \rightleftharpoons B ,$$

where ϕ is the azimuthal angle of the pair, and ϕ_A, ϕ_B describe the hadron spin orientation. Constructing an asymmetry by integrating over the angles, subtracting two “side” quadrants from two “top and bottom” quadrants according to

$$d\Delta\sigma/dQ^2 dy = \int_{-\pi/4}^{3\pi/4} d\phi + \int_{5\pi/4}^{7\pi/4} d\phi - \int_{3\pi/4}^{5\pi/4} d\phi - \int_{-\pi/4}^{\pi/4} d\phi \quad d\sigma/dQ^2 dy d\phi$$

where both protons are polarized in the x direction, one finds⁵¹

$$d\Delta\sigma/dQ^2 dy = \frac{4}{9} \sum_a \frac{\alpha^2 e_a^2}{Q^2 s} h_T^{a/A}(x_A) h_T^{\bar{a}/B}(x_B) + A \rightleftharpoons B .$$

This shows that measurement of leading order transverse spin polarization distribution is not much more difficult than measurement of the unpolarized distributions, provided one has a source of polarized beam and target.

Where could doubly polarized experiments be performed? Siberian snake technology has developed so that high energy polarized beams are a practical possibility.⁵⁰ The RHIC Spin Collaboration proposes that the Relativistic Heavy Ion Collider planned at BNL be designed to allow acceleration of high energy polarized protons.

We opened up this question at the *Penn State Polarized Collider Physics* meeting held in November, 1990.⁵¹ (Ralston also served on the External Advisory Committee for the meeting.) The issue of transverse spin had also received some discussion by Collins,⁵² and Artru and Mekhfi,⁵² and recently has been examined by Jaffe and Ji.⁵² In a longer and much more comprehensive paper⁵³ with Cortes (Zaragoza University) we explain and simplify the relation of the leading twist polarization distribution to the much more complicated problem of electroproduction. Electroproduction involves understanding the parton model at higher twist, a rather technical subject which has been clarified to some extent.⁵⁴ We were able to show that certain

measurements in Drell-Yan could be used to strengthen the interpretation of measurements of g_2 . We obtained a sum rule for the twist 3 correlator $g_n(x, Q^2)$ which when inverted reads⁵³

$$g_n(x, Q^2) = g_1(x, Q^2) + g_2(x, Q^2) + \int_x^1 \frac{dy}{y} g_1(y, Q^2) + \sum_f \frac{m_f}{M} e_f^2 \int_x^1 \frac{dy}{y} \frac{\partial h_T}{\partial y}(y, Q^2)$$

where m_f is the quark mass and M the proton mass. This shows that the measurement of g_2 does not uniquely isolate twist -3 operator matrix elements, despite some statements to the effect in the literature. Instead, one must know h_T from, *e.g.* Drell-Yan measurements. Besides the *Penn State* talk, this topic was also presented by Ralston in seminars at *Universite Paris Sud* and with the *SMC Collaboration* at CERN.

c) High Energy Hadronic Helicity Violation in Exclusive Processes

For many years there has been a perplexing pattern in data on exclusive hadron reactions that shows a violation of perturbative QCD expectations. Specifically, the helicity conservation of the theory at the quark level does not translate itself into reliable selection rules at the hadron level. There is a pattern of violation of the most simple and general results, which do not depend on details assumed of the wave functions. An example is the failure of the total hadronic helicity conservation rule given by LePage and Brodsky,³⁹ which says that for hadrons $A + B \rightarrow C + D$,

$$\lambda_A + \lambda_B = \lambda_C + \lambda_D$$

where λ_i are the helicities of the hadrons involved. This rule can be shown to be as general as the factorization prescription itself, schematically expressed in (2). Thus its violation is a serious issue, widely interpreted to mean trouble for perturbative QCD. In some cases researchers have been led to make claims that the theory itself must be wrong.

It turns out that a loophole in the arguments allows high energy hadronic helicity non-conservation, even in a theory with perfect chiral symmetry at the quark level. In our study of the independent scattering processes in transverse b -space, we found this loophole by looking at the orbital angular momentum decomposition of the hard scattering. Unlike the quark counting case, the independent scatterers do *not* project out the zero orbital angular momentum quark wave function. Instead, the independent scattering “hard kernel” remembers the scattering plane in every event; it is literally “cigar shaped”, pointing out of the scattering plane. Thus the scattering process itself, lacking rotational symmetry about the scattering momentum directions, allows many units of orbital angular momentum to participate. We believe this physical mechanism can explain every observed violation of the helicity conservation rules.

The result violates the usual factorization assumptions because the whole independent scattering amplitude factorizes in an entirely different way. Botts and Sterman²⁹ gave formalism which is very useful for seeing this, and in fact also commented on the “matchstick shape”, without connecting it to spin. Indeed, the lack of symmetry was present in the independent scattering mechanism all along.

Putting together the formalism with the short-distance properties of non-zero angular momentum states (which are prescribed by their transformation properties) we calculated the asymptotic power of energy of the helicity flip amplitudes using Mueller’s method.⁵⁵ They turn out to be fractionally suppressed, but not true “higher twist”, and potentially large.⁵⁶

These results for the meson-meson scattering case were published in a short contribution to the *Penn State Polarized Collider Physics Workshop*.⁵⁶ We pointed out that the mechanism is testable, in that one should see a calculable attenuation of helicity – flip amplitudes in experiments on nuclear targets. Work nearing completion involves finishing a more comprehensive paper, where we examine fairly complicated details of the mechanism, including intricacies associated with the Feynman x -dependence of the wave functions.

Particle Physics and Cosmology

Hollow Galactic Dark Matter Halos

Dark matter in the universe has become a central area in the interface between cosmology and particle physics. The case for “hot dark matter” has become stronger with the realization that the alternative “cold dark matter” scenario has probably run into contradictions with data on large scale structure. Attention naturally turns to the most likely candidate for hot dark matter, the neutrino. However attractive the neutrino for large scale structure, there has been a tendency to dismiss the neutrino as an explanation of dark matter, in part due to the so-called Tremaine–Gunn limit.⁵⁷ This relation, a more precise formulation of a bound first obtained by Cowsik and McClelland,⁵⁷ specifies a lower limit on the mass of neutrinos gravitationally bound in galaxy halos. One uses the known Big-Bang phase space density to limit the number density of neutrinos one can later find in the halo. A convenient form for the bound on the neutrino mass m_ν is

$$m_\nu > m_{TG} = 120 eV \left(\frac{100 km/s}{\sigma} \right)^{1/4} \left(\frac{1 kpc}{r} \right)^{1/2} g_\nu^{-1/4} ,$$

where σ is a galactic velocity dispersion, r a galaxy core size, and g_ν the number of neutrino spin states. Since the best data for dark matter is from dwarf galaxies (which have small r and σ) the relation is potentially serious.⁵⁸ Indeed, data from dwarf galaxies has been used to argue there

is a contradiction in having neutrinos explain dark matter on both the galactic and cosmological scale.⁵⁹ However, Ralston showed⁶⁰ that the “bounds” were actually model-dependent, (and therefore not really bounds), by constructing allowed solutions of the relevant equations which violated the bounds.

Exploring this result more generally, Ralston and KU graduate student Lesley Smith⁶¹ found that the special solutions were examples of a general pattern: anisotropic phase space halos often violate the bound. An anisotropic phase space can occur in a collisionless system, where, *e.g.*, the average radial and angular velocities of the particles are not equal. This complicates the mathematics a bit, and one must study the non-linear “Poisson-Vlasov” equation

$$\nabla^2\psi = 4\pi Gm_\nu \int d^3v f(x,v)$$

numerically to investigate it. Here ψ is the gravitational potential energy per mass, G is Newton’s constant, and $f(x,v)$ is the joint position-velocity space density, which depends implicitly on ψ .

An anisotropic phase space is actually expected due to angular momentum conservation. When a halo collapses, the orbital angular momentum is limited because of the limited impact parameter of particles trapped in the halo. This leads to a distorted velocity space, and in fact to a “hole” in the density distribution in the center of a halo, due to an angular momentum barrier.

Rather amazingly, the long range structure of such an anisotropic halo generically gives the proper mass density to produce the flat rotation curves observed in data. This result can be shown analytically with a calculation that accounts for the angular momentum cutoff in the velocity space. Numerical work on the non-linear Poisson-Vlasov equation was presented and proved that the analytic arguments held up. It was apparently not known before: most studies simply assumed an isotropic velocity space. Flat rotation curves were then attributed to a special “fine-tuned” distribution occurring again and again for no compelling reason.

Re-calculating the neutrino mass limit (which would also apply to any fermion, and in some cases to bosons⁶²) the following more general relation was obtained:⁶¹

$$m_\nu > K_0^{1/4} m_{TG}$$

where K_0 is a dimensionless parameter containing the typical angular momentum scale of the halo; limited angular momentum is $K_0 \ll 1$. For this case, the “bound” becomes weaker. The argument was subsequently picked up and extended by Madsen⁶² who came to the same conclusion.

The results⁶¹ were published in *Astrophysical Journal*, and presented in invited seminars at *Ohio State University*, *William and Mary College*, *Michigan State University* and *CFMC Lisboa (Lisbon)*. Ralston also contributed an invited review of dark matter for the *1991 Yearbook of the McGraw Hill Encyclopedia Of Science and Technology*.

Anisotropy in Quantum Networks and Integer Quantum Hall Behavior

Quantum networks⁶³ are a topic of interest to physicists and mathematicians interested in gauge theories. The idea is to define the gauge theory on finite one-dimensional subspaces called “wires”, and then to build a non-trivial multi-dimensional system by patching the wires at intersections called “nodes”. One reduces an infinite dimensional problem a finite dimensional one, because the equations can generally be solved along the wires, and there are just a finite number of gauge invariant conditions to be met for a finite number of nodes. Perhaps even more interesting, the physics of some systems can be tested in the laboratory by building the network out of superconducting filaments and treating the Ginzburg -Landau order parameter as a macroscopic wave function interacting with an external magnetic field.

Sun and Ralston looked at the effects of changing the network connection rules. The standard rules are known as the deGennes–Alexander equations.⁶³ One simple change which had not been investigated was to allow for anisotropy in the node patching rules of a two dimensional network. Two-dimensional electrons in a gauge (magnetic) field is an old and surprisingly rich topic, but no one had yet investigated the effects of such anisotropy in a network.

The investigation revealed a surprisingly complicated spectrum, related to the “butterfly” fractal made famous by Hofstadter.⁶⁴ With more anisotropy, we found we could regulate and apparently turn-off the fractal character, which is associated with what is known as “frustration” in condensed matter physics. This was a significant finding we published in *Physical Review B*.⁶⁵ Sun, a KU student supported by University funds, gave a talk at an APS meeting presenting the results.

Very recently, we were pleased to learn that the laboratory realization of our network had been constructed by the group of Itzler, Bojko and Chaikin (a Penn/Princeton/Cornell collaboration).⁶⁶ The group looked for the phenomenon of “fading” of the phase transition we predicted and observed it. But besides is behavior going as we predicted, the group reported a second anisotropic phenomenon that we did not expect in the network. It is not clear what this means, and it is being studied.

In the meantime, Sun and Ralston adapted the study to the area of Integer Quantum Hall behavior. This is possible because the fundamental symmetry of the network can be made the same as that of electrons in a two-dimensional periodic potential and a magnetic field. The arguments of Laughlin⁶⁷ and Thouless⁶⁷ then apply; indeed, one can turn the network argument around, and make a prediction of new phenomena in the Quantum Hall effect. We found that anisotropy in the network can regulate the Hall conductivity and turn off or “quench” the integer quantization. This is a surprising result, given that integer quantization is nowadays seen as commonplace: it is done by unwinding the topological quantum number when levels mix with increasing distortion of the lattice parameters. A paper submitted to *Physical Review B* has been accepted and will appear soon.⁶⁸ Sun has finished his studies at KU and is currently a post-doc in condensed matter theory at Georgia Tech.

Anomalous Color Transparency

Color transparency is not expected to occur in a conventional treatment of exclusive nuclear photoproduction of π^0 's. However, the anomalous $\gamma\gamma\pi^0$ vertex involved in the production via the Primakoff effect suggests a very interesting possibility. If one interprets the vertex in the most naive way, then its value comes from regions in momentum space where the quarks are far off-shell and have large transverse momenta. In parton language, the interaction favors the configuration of quarks in the pion which are spatially close together. In this case, color transparency might occur as a closely separated quark anti-quark pair is created in the nuclear electromagnetic field. The system would subsequently time-evolve into a normal pion, but only long after interacting. The signal would be an unexpectedly small attenuation of the photoproduced pion by the nuclear target.

On the other hand, one can also view the anomaly as a general sort of consistency condition, and conjecture that the perturbative quark triangle graph may be physically misleading. Indeed there is no large invariant momentum in the process to favor short distance. In that case one would not have color transparency, but instead would expect a rather normal cross section for photoproduced pions to interact with a nucleus.

Both possibilities have been worked out in a paper by Ralston to appear⁴⁷ in *Physics Letters B*. Instead of attempting to settle the problem theoretically – which probably cannot be done – the paper proposes a way to measure the spatial extent of the anomalous vertex. Small momentum transfer, in a kinematic region which will select the interesting events which are photoproduced and yet penetrate the nucleus, is used along with high energy. The energy

dependence shows a characteristic signal if the produced quark anti-quark system is anomalously small, rising with energy significantly in the region $E \gtrsim \text{GeV}$. The rise with energy would be due to the Lorentz dilation of a time scale for the quarks to become a pion; thus the experiment is relevant to the hadron expansion time issue mentioned above. There are few conceivable effects which could mimic such a signal: thus the experiment is suggested to be performed. The paper was written during Ralston's visit at CERN during the summer of 1991, and was presented at the *Saclay Meeting on Electron Physics above 15 GeV* in June, as well as in a joint seminar at the *Gordon Conference* in Tilton, NH, in July.

Quantum Treatment of Solitons

Jain has been working on the $1 + 1$ dimensional ϕ^4 and the sine Gordon field theories in an attempt to understand the behavior of the quantum solitons under Lorentz transformations. Although the systematic quantum expansion in the presence of solitons was developed and applied to these model field theories almost a decade ago,⁶⁹⁻⁷² the behavior of such solutions under boosts was not properly analyzed. This has been pointed out in ref. [73], where Jain has shown that the regularization used so far in the literature^{69,72} to calculate the soliton energy in both the ϕ^4 and the sine Gordon theory violates the Poincare algebra. It turns out, fortunately, that one can find regularizations which do not suffer from this problem. Jain has also analyzed the doublet solutions of the sine Gordon theory by using a regularization which respects the Poincare algebra. It is well known that the semiclassical procedure used by Dashen, Hasslacher and Neveu⁶⁹ yields exact results for this case. Jain has shown, however, that this procedure is not reliable in general since it does not yield renormalization-scheme-independent results even for the mass ratios of the doublet solutions. It turns out that only the weak coupling expansion, also discussed in ref. [69], yields unambiguous results after renormalization. Finally,⁷⁴ by taking the simple example of the topological current in $1 + 1$ dimensional kink field theory, Jain has demonstrated the procedure for calculating the diagonal matrix elements of current operators in the presence of solitons, while preserving fundamental symmetries like Lorentz covariance. A similar demonstration for the case of energy of the soliton has already been given in refs. [70] and [75].

Beside this interest in the quantum treatment of $1 + 1$ dimensional solitons, Jain is engaged in applying these methods to the phenomenological chiral lagrangians of strong interactions which contain only the mesons as the "fundamental" degrees of freedom. These theories are rather interesting since, as originally suggested by Skyrme⁷⁶, the baryons emerge as solitonic solutions in these models. Although a lot of progress has been made in understanding the

soliton sector of these theories,⁷⁷⁻⁷⁹ a thorough quantum treatment is still lacking. Moreover, no realistic calculations of the energy or form factors of the baryons, performed so far, have even attempted to preserve the fundamental symmetries like Lorentz covariance. Jain is interested in setting up a systematic quantum treatment of baryons, such that the Lorentz covariance is preserved to each order of the expansion. Jain and Ralston intend to investigate numerical approximations to Ralston's method,⁸⁰ which is explicitly covariant but also is fairly complicated. One of the main problems in performing such a calculation is the nonrenormalizability of these models, which will make it necessary to introduce counterterms with undetermined parameters. Such parameters, however, can be fitted phenomenologically.

Quantum Stabilization of Skyrmions

Jain is also working on the speculative idea^{81,82} that in the absence of the stabilizing Skyrme term, the Skyrme soliton may be stabilized by quantum fluctuations. The interesting aspect of this model is that it has no undetermined parameters. The only quantity that remains undetermined is the functional form of the soliton profile. In ref. [83] this profile was obtained by fitting it to the experimental proton charge density. The resulting baryon spectrum and the nucleon static properties were shown to be in better agreement with experiments in comparison to the standard Skyrme model. It can be argued that the soliton profile should be obtained by minimizing the total energy. However it turns out that it is possible to find soliton profiles that will reduce the energy to zero.^{81,84} In ref. [81], Jain and collaborators had argued that this was unphysical since such collapse of soliton was obtained only for very long-range profiles. Since only the collective modes were included for quantization, an implicit assumption in this procedure is that the motion of each part of the baryon system is instantaneously correlated with all other parts. This clearly violates relativity in a serious way if the profile is too large. The problem is closely related to a "no-go" rule for collective coordinates published earlier by Ralston.⁸⁵ A possible improvement over existing procedures is to also include the pion fluctuations around the baryons.

Color Transparency

Jain, Schecter and Weigel⁷⁹ have recently used the Skyrme model to calculate the color transparency of the nucleons taking part in nucleon nucleon elastic scattering inside nucleus at momentum transfer up to about 1.5 GeV. In order to perform the calculations, it was assumed that the nuclei taking part in elastic scattering with large momentum transfer fluctuate to a very small spatial size. This "mini proton" was modelled in the Skyrme model by quantizing the breathing mode and by assuming that the wave function of this object is a superposition of the

radial eigenstates. The subsequent time development was then determined by the usual quantum dynamics. Proceeding in this fashion, Jain *et al.* calculated the decrease in cross section of the “mini-protons” propagating through the nucleus by assuming that the cross section scales as the expectation value of the area. The result obtained in ref. [86] show that the Skyrme model can support the notion of color transparency.

Casimir effect

As discussed above, Jain has shown that a careful consideration of the behaviour of solitons under Lorentz boosts reveals that the regularization used so far in the literature^{69–72} to analyze the kink soliton and the sine Gordon soliton violates the Poincare algebra. This result was demonstrated by calculating the diagonal matrix elements of the Poincare algebra for these theories between the one soliton states. It turns out that the Poincare algebra provides us with a useful check on any calculation which involves a nontrivial subtraction of the vacuum energy. Jain is interested in applying these considerations to analyze the change in Casimir stress⁸⁷ between macroscopic objects under Lorentz transformations.

REFERENCES

1. S.H. Tye, E. Tomboulis and E. Poggio, *Phys. Rev. D* **11**, 2839 (1975); K. Higashijima, *ibid.* **29**, 1228 (1983).
2. K. Lane *Phys. Rev. D* **10**, 2605 (1974); H.D. Politzer, *Nucl. Phys.* **B117**, 397 (1976). The former reference does not claim to reproduce the correct power of $\ln p^2$ which modifies the p^{-2} behavior of the regular solution.
3. C. Roberts and B. McKellar, *Phys. Rev. D* **41**, 672 (1990).
4. H. Munczek and D. McKay, *Phys. Rev. D* **42**, 3548 (1990).
5. R. Tarrach, *Nucl. Phys.* **B183**, 384 (1981).
6. D. McKay "An Upper Bound on the Higgs Mass", "Beyond the Standard Model II", K. Milton, Ed. *World Scientific* (to be published) and seminars at Iowa State U. and the University of Mainz.
7. P. Hung, *Phys. Rev. Lett.* **42**, 873 (1979). It is not yet clear whether this bound applies when a less restrictive form of the potential is used. The potentials are reviewed by M. Sher, *Phys. Rep.* **179**, 273 (1989).
8. R. Johnson, B-L. Young and D. McKay, *Phys. Rev. D* **43**, R17 (1991).
9. Other authors reached similar conclusions at the same time using a variety of techniques, all different from Johnson, Young and McKay and from each other: M. Peskin and T. Takeuchi, *Phys. Rev. Lett.* **65**, 964 (1990); B. Holdom and J. Terning, *Phys. Lett. B* **247**, 88 (1990); M. Goldon and L. Randall, *Nucl. Phys. B*.
10. G. Askaryan, *Zh. Eksp. Teor. Fiz* **41**, 616 (1961). [*Soviet Physics JETP* **14**, 441 (1962)].
11. M. Markov and I. Zheleznykh, *Nuclear Instruments and Methods in Physics Research* **A248**, 242 (1986).
12. J. Ralston and D. McKay, "Patterns of Small- x QCD", *Nucl. Phys. B* (Proc. Suppl.) **14A**, 1990.
13. J. Ralston and D. McKay, "Comparing Coherent Microwave Emission in LPM and B-H Showers", in *High Energy Gamma Ray Astronomy*, AIP Conference Proceedings Number **220**, **295** (AIP, NY 1991).
14. H. Allen, *Progress in Elementary Particles and Cosmic Ray Physics* **10**, eds. J. Wilson and S. Wouthuysen (North Holland, 1971).

15. H.J. Munczek and A. Nemirovsky, *Phys. Rev. D* **28**, 181 (1983); H.J. Munczek, *Phys. Lett. B* **175**, 215 (1986); D.W. McKay, H.J. Munczek and B.L. Young, *Phys. Rev. D* **37**, 195 (1988).
16. Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961); M. Baker, K. Johnson, and B.W. Lee, *Phys. Rev.* **133**, B209 (1964); H. Pagels, *Phys. Rev. D* **7**, 3689 (1973); R. Jackiw and K. Johnson, *Phys. Rev. D* **8**, 2386 (1973); J.M. Cornwall and R.E. Norton, *Phys. Rev. D* **8**, 3338 (1973); E.J. Eichten and F.L. Feinberg, *Phys. Rev. D* **10**, 3254 (1974); R. Delbourgo and M.D. Scadron, *J. Phys. G* **5**, 1621 (1979).
17. P. Jain and H.J. Munczek, *Phys. Rev. D* **44**, (1991).
18. The issue of the applicability of the PS expression for f_π is of importance not only in QCD but also in other theories such as technicolor. A paper by Aoki *et al.* *Progress of Theor. Phys.* **84**, 63 (1990) of which we became aware after Ref. 17 was published claims that the PS approximation is a good one. However, this claim is based on calculations that employ the approximations to $\alpha_s(q^2)$ and to $A(q^2)$ mentioned above. We have verified numerically that both approximations are quite inaccurate for the cases considered by Aoki *et al.* That inaccuracy leads to their wrong conclusion about the PS approximation. A short comment on the subject is in preparation.
19. H.J. Munczek and P. Jain, Preprint KUHEP-91-7, submitted to *Phys. Rev. D*; Preliminary report submitted to the *Joint International Lepton-Photon Symposium*, Geneva, August 1991.
20. V.A. Miransky, *Mod. Phys. Lett. A* **5**, 1979 (1990).
21. See, for example, J.L. Rosner, *Phys. Rev. D* **42**, 3732 (1990); University of Chicago Report EFI 90-63.
22. S.J. Brodsky and A.H. Mueller, *Phys. Lett. B* **206**, 685 (1988) and references therein.
23. F. Low, *Phys. Rev. D* **12**, 163 (1975).
24. J.F. Gunion and D.E. Soper, *Phys. Rev. D* **15**, 2617 (1977); E.M. Levin and M.G. Ryskin, *Yad. Fiz.* **34**, 1114 (1981).
25. B. Pire and J.P. Ralston, *Phys. Lett. B* **117**, 233 (1982).
26. P.V. Landshoff, *Phys. Rev. D* **10**, 1024 (1974).
27. S.J. Brodsky and G. deTerra, *Phys. Rev. Lett.* **49**, 1605 (1982).

28. J.P. Ralston and B. Pire, *Phys. Rev. Lett.* **49**, 1605 (1982).
29. J. Botts and G. Sterman, *Nuc. Phys. B* **325**, 62 (1989).
30. J. Botts, Washington Preprint (1991).
31. A.S. Carroll, *et al.* *Phys. Rev. Lett.* **61**, 1698 (1988).
32. J.P. Ralston and B. Pire, *Phys. Rev. Lett.* **61**, 1823 (1988).
33. G. Bertsch, S.J. Brodsky, A.S. Goldhaber and J.E. Gunion, *Phys. Rev. Lett.* **47**, 297 (1981).
34. B. Pire and J.P. Ralston, "Coherence and Color Transparency," in *Topical Conference on Electronuclear Physics with Internal Targets* (Stanford, 1989), edited by R.G. Arnold (World Scientific, 1990).
35. J.P. Ralston and B. Pire, "Nuclear Filtering of Strong Interactions," in *24th International Rencontre de Moriond* (Les Arcs, 1989) edited by J. Tran Thanh Van (Editions Frontiers, 1990).
36. B. Pire and J.P. Ralston, "Color Transparency and Mini-Hadron Dynamics," in *European Physical Society-International Conference on High Energy Physics* (Madrid, 1989). *Nuc. Phys. B, (Proc. Suppl.)*, **16**, 254, (1990).
37. B. Pire and J.P. Ralston, "Hard Scattering in a Nuclear Environment," *Nuc. Phys.* **A525** (419c) 1991, proceedings of *Quark Matter 90*, edited by J.P. Blaizot, *et al.*
38. J.P. Ralston and B. Pire "Quantum Color Transparency," *Phys. Rev. Lett.* **65**, 2343 (1990).
39. G.P. Lepage and S.J. Brodsky, *Phys. Rev.* **D22**, 2157 (1980). See also in the review *Quantum Chromodynamics*, edited by A.H. Mueller (World Scientific, Singapore, 1990).
40. N. Isgur and C.H. Llewellyn-Smith, *Phys. Rev. Lett.* **52**, 1080 (1984); *Phys. Lett.* **B47**, 535 (1989).
41. B. Pire and J.P. Ralston, "A Transparency Scaling Law for Large Q^2 Exclusive Reactions on Nuclei," *Phys. Lett.* **B256**, 523 (1991).
42. B. Pire and J.P. Ralston, "Theory of Color Transparency" in *Proceedings of 5th Workshop on Perspectives in Nuclear Physics at Intermediate Energies* (Trieste 1991) (World Scientific, in press); J.P. Ralston and B. Pire, "Color Transparency and Electronuclear Physics"

to be published in *Proceedings of Second Workshop on Hadronic Physics with Electrons Beyond 10 GeV* (Dourdan, France 1990) *Nuc. Phys. A* (in press).

43. G.R. Farrar, H. Liu, L. Frankfurt and M. Strikman, *Phys. Rev. Lett.* **61**, 686 (1988).
44. B. Kopeliovich, *Sov.J. Part. Nucl.* **21**, 49 (1990); Jennings and G. Miller, *Phys. Lett. B* **236**, 209 (1990); *Phys. Rev. D* (in press).
45. G.R. Farrar, L. Frankfurt and M. Strikman, *Phys. Rev. Lett.* (Comments) **67**, 2111 (1991).
46. B. Pire and J.P. Ralston, *Phys. Rev. Lett.* (Comment Response) **67**, 2112 (1991).
47. J.P. Ralston, "Anomalous Color Transparency," CERN preprint TH.6139/91, *Physics Letters B* (in press).
48. R.P. Feynman, *Photon Hadron Interactions* (Benjamin, NY, 1972).
49. J.P. Ralston and D.E. Soper, *Nuc. Phys.* **B152**, 109 (1979).
50. A. Krisch, in *Polarized Collider Workshop*, (AIP Conference Proceedings No. 223) edited by J. Collins *et al.* (AIP, NY, 1991).
51. J. Cortes, B. Pire and J.P. Ralston, "Lepton Pair Production in Polarized Collider Experiments as a Method to Measure the Transverse Polarization of Quarks in the Proton," in *Polarized Collider Workshop* *ibid.*
52. X. Artu and M. Mekhfi, *Zeit. Phys.* **C45**, 669 (1990); J.C. Collins in *Polarized Collider Workshop* (*ibid.*); R.L. Jaffe and X. Ji, MIT Preprint (1991).
53. J. Cortes, B. Pire and J.P. Ralston, "Measuring the Transverse Polarization of Quarks in the Proton," Ecole Polytechnique preprint A048.0491 (1991), submitted to *Zeit. Phys. C*.
54. P.G. Ratcliff, *Nuc. Phys. B* **264**, 493 (1986). See also L. Mankiewicz and Z. Ryzak, *Phys. Rev. D* **43**, 733 (1991).
55. A.H. Mueller, *Phys. Rep.* **73**, 237 (1981).
56. J.P. Ralston and B. Pire, "Spin and the Independent Scattering Mechanism," in *Polarized Collider Workshop* (AIP Conference Proceedings No. 223) edited by J. Collins *et al.* (AIP, NY, 1991)
57. R. Cowsik and J. McClelland, *Phys. Rev. Lett.* **29**, 669 (1972); *Astrophys. J.* **180**, 7 (1973); S. Tremaine and J.E. Gunn, *Phys. Rev. Lett.* **42**, 407 (1979).

58. For a review see *Dark Matter in the Universe*, (Reidel, Dordrecht 1986) edited by J. Kormendy and G.R. Knapp.
59. D.N. Spergel, D.H. Weinberg, and J.R. Gott III, *Phys. Rev. D* **38**, 2014 (1988).
60. J.P. Ralston, *Phys. Rev. Lett.* **63**, 1038 (1989).
61. L.L. Smith and J.P. Ralston, "Hollow Galactic Halos of Fermionic Dark Matter," *Astrophys. J.* **367**, 54 (1991).
62. J. Madsen, *Astrophys. J.* **367**, 341 (1991); Aarhus preprint (1991).
63. For a review see J. Avron *et al.*, *Rev. Mod. Phys.* **60**, 873 (1988). See also P. DeGennes, *C.R. Acad. Sci. Ser. II*, **9** (1981); S. Alexander, *Phys. Rev.* **B27**, 1541 (1983).
64. D.R. Hofstadter, *Phys. Rev.* **B14**, 2239 (1976).
65. S.N. Sun and J.P. Ralston, "Duality Symmetry and Power Law Fading of Frustration in a Quantum Multi-Connected Superconductor," *Phys. Rev.* **B43**, 5375 (1991).
66. M.A. Itzler, R. Bojku and P.M. Chaikin, Penn Preprint (1991), submitted to *Phys. Rev. Lett.*
67. R.B. Laughlin, *Phys. Rev.* **B23**, 5623 (1981); D.J. Thouless *et al.* *Phys. Rev. Lett.*, **49**, 405 (1982).
68. S.N. and J.P. Ralston, "The Possibility of Quenching of Integer Quantum Hall Behavior with Increasing Lattice Asymmetry," *Phys. Rev. B* (in press).
69. R. Dashen, B. Hasslacher and A. Neveu, *Phys. Rev.* **D10**, 4130 (1974); **D11**, 3424 (1975).
70. N. H. Christ and T. D. Lee, *Phys. Rev.* **D12** 1606 (1975).
71. J.-L. Gervais, A. Jevicki and B. Sakita, *Phys. Rev.* **D12**, 1038 (1975); J. Goldstone and R. Jackiw, *Phys. Rev.* **D11**, 1486 (1975); E. Tomboulis, *Phys. Rev.* **D12**, 1678 (1975); C. Callan and D. Gross, *Nucl. Phys.* **B93**, 29 (1975).
72. See *e.g.* *Solitons and Instantons* by R. Rajaraman (1982), (Amsterdam: North Holland).
73. P. Jain, *Modern Physics Letters A* **6**, 2363 (1991); 2509 (1991).
74. P. Jain, "Lorentz Covariance of Current Matrix Elements in a Soliton Model," to be published in *Physical Review D*.
75. P. Jain, *Phys. Rev.* **D41**, 3273(BR), (1990).

76. T. Skyrme, *Proc. R. Soc. London* **A260**, 127 (1961); *Nucl. Phys.* **31**, 556 (1962).
77. Reviews include I. Zahed and G. Brown, *Phys. Rep.* **142**, 1 (1986); G. Holzwarth and B. Schwesinger, *Rep. Prog. Phys.* **49**, 825 (1986).
78. G. S. Adkins, C. R. Nappi and E. Witten, **B228**, 552 (1983); A. Balachandran, V. P. Nair, S. G. Rajeev, and A. Stern, *Phys. Rev.* **D27**, 1153 (1983).
79. I. Zahed, A. Wirzba, U.-G. Meissner, *Phys. Rev.* **D33**, 830 (1986).
80. J.P. Ralston, *Phys. Rev.* **D33**, 496 (1986); **D43**, 2003 (1986).
81. P. Jain, J. Schechter and R. Sorkin, *Phys. Rev.* **D39**, 998, (1989); **D41**, 3855(BR), 1990.
82. J. Carlson, *Nucl. Phys.* **B253**, 149 (1985); **B277**, 253 (1986).
83. P. Jain, *Phys. Rev.* **D41**, 3527(BR), (1990).
84. A. Kobayashi, H. Otsu and S. Sawada, *Phys. Rev.* **D42**, 1868 (1990); S. Sawada and K. Yang, *Phys. Rev.* **D44**, 1578 (1991).
85. J.P. Ralston, *Phys. Rev. (Rapid Comm.)* **D34**, 1245 (1986).
86. "Approach to Color Transparency in the Soliton Picture of the Nucleon," P. Jain, J. Schechter, H. Wiegel, submitted to *Physical Review D*.
87. See, for example, C. Itzykson and J. B. Zuber, *Quantum Field Theory* (McGraw-Hill, 1980); T. H. Boyer, *Annals of Physics* **56**, 474 (1970); Roger Balian and Bertrand Duplantier, *Annals of Physics* **112**, 165, (1978); Julian Schwinger, Lester L. DeRaad, Jr. and Kimball A. Milton, *Annals of Physics* **115**, 1 (1978).

END

**DATE
FILMED**

8 / 19 / 92