The Electroweak Anomaly and Current
Algebra for $K \rightarrow \gamma \ell \nu$

Kimball A. Milton^{*,†}
Department of Physics
University of California
Los Angeles, California 90024

and

Walter W. Wada^{††}
Department of Physics
The Ohio State University
Columbus, Ohio 43210

MASTER

Abstract

The process $K \rightarrow \gamma \ell \nu$ is calculated using the electroweak axial-vector anomaly with the quark color factor of 3, together with standard current-algebra techniques. The result, which generalizes that of Das, Mathur, and Okubo for the axial-vector part, is in good agreement with experiment.

*Work supported in part by the National Science Foundation.

†Address 1980-81: Department of Physics, The Ohio State University, Columbus, Ohio 43210.

††Work supported in part by the Department of Energy under contract DE-AC02-76ER01545.

DISCLAIMER

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

fcy

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

In the context of the development of what presumably is the theory of strong interactions, quantum chromodynamics, current algebra and associated questions are receiving something of a renaissance. One of the key advances in that earlier development was the understanding of the two-photon decay of the pion. It was the axial-vector anomaly^[1] which precluded the zero predicted in the soft-pion limit by Sutherland's theorem,^[2] and which in fact predicts a rate for $\pi^0 \rightarrow \gamma\gamma$ completely consistent with experiment.¹

The synthesis of electromagnetic and weak interactions^[3] provides an arena for testing extensions of this anomaly-mediated process. The electroweak anomaly, for example, should underly the vector form factor of the decay $K \rightarrow \gamma \ell \nu$. This and the attendant axial-vector form factor resulting from current algebra are particularly ripe for study on both theoretical and experimental grounds. For although there is an extensive literature^[4] on the theory of these and related amplitudes, the models are principally of a phenomenological character. Beyond the inclusion of the anomaly, what we attempt here is a generalization of the treatment by Das, Mathur, and Okubo^[5] of the axial-vector form factor for the analogous decay $\pi \rightarrow \gamma e \nu$. Thus, we are testing the sufficiency of anomaly-mediation, PCAC for the kaon, current algebra, and the concomitant soft kaon limit. Experimentally, $K \rightarrow \gamma e \nu$, which proceeds at almost the same rate as $K \rightarrow e \nu$, has now been measured over a significant portion of its phase space,^[6] so that one knows more than simply the magnitude of the amplitude. Perhaps surprisingly, agreement with experiment is excellent.

The essential matrix element of the weak vector current governing the structure-radiation² amplitude for the decay $K^+(p) \rightarrow \gamma(k) + \ell^+ + \nu$ in the soft-kaon limit ($p \rightarrow 0$) is given by

$$\begin{aligned}
\langle \gamma(k, \epsilon) | V_{\mu}^{+}(0) | K^{+}(p) \rangle &= i \int d^4x \frac{1}{(2\pi)^{\frac{3}{2}}} \frac{1}{\sqrt{2\omega_k}} e^{-ipx} \\
&\times \frac{1}{F_K} \langle \gamma(k, \epsilon) | T(V_{\mu}^{+}(0) [\partial^{\alpha} A_{\alpha}^{-}(x) - a F_{\alpha\beta}(x) {}^*W^{-\alpha\beta}(x)]) | 0 \rangle, \quad (1)
\end{aligned}$$

where $V_{\mu}^{+} (= V_{\mu}^4 - i V_{\mu}^5)$ is the weak vector current coupled to the W vector boson. The PCAC relation with the anomaly term for the interpolating field of the K^{-} has been incorporated in (1). It is given by

$$\partial^{\alpha} A_{\alpha}^{-} = m_K^2 F_K \phi_{K^{-}} + a F_{\mu\nu} {}^*W^{-\mu\nu}, \quad (2)$$

where

$$A_{\alpha}^{-} = A_{\alpha}^4 + i A_{\alpha}^5, \quad (3)$$

$${}^*W^{-\alpha\beta} = \frac{1}{2} \epsilon^{\alpha\beta\rho\sigma} (\partial_{\rho} W_{\sigma}^{-} - \partial_{\sigma} W_{\rho}^{-}), \quad (4)$$

and

$$a = \frac{\alpha \sin \theta_c}{4\sqrt{2} \pi \sin \theta_W} \quad (5)$$

in the Weinberg-Salam theory of weak interaction.^[3] Equation (5) includes the color factor of 3.

The anomaly term³ in (2) has been obtained from the quark diagrams in Fig. 1 in analogy to that for $\pi^0 \rightarrow \gamma\gamma$.^[1] The divergence of the axial vector term $\partial^{\alpha} A_{\alpha}^{-}$ in (1) leads to zero in the matrix element (1), by Sutherland's

theorem.^[2] Introducing the coupling of the W with the leptonic current,^[3] it can easily be shown that the vector decay amplitude is given by

$$M_V^{SR}(K^+ \rightarrow \gamma \ell^+ \nu) = i \frac{G_F e \sin \theta_c}{\sqrt{2}} \mathcal{V} \epsilon_{\lambda\mu\alpha\beta} k^\alpha p^\beta \epsilon^\lambda \bar{u}_\nu \gamma^\mu (1 - \gamma_5) v_\ell, \quad (6)$$

where the vector form factor $\mathcal{V}$ is

$$\mathcal{V} = \frac{1}{4\pi^2} \frac{1}{F_K}, \quad (7)$$

or numerically ($F_K = 148 \text{ MeV}^{[9]}$)

$$m_K \mathcal{V} = 0.084. \quad (8)$$

The matrix element analogous to (1) for the weak axial-vector current is

$$\langle \gamma(k, \epsilon) | A_\mu^+(0) | K^+(p) \rangle = -i \frac{1}{(2\pi)^{\frac{3}{2}}} \frac{1}{\sqrt{2}\omega_\gamma} e \epsilon^\lambda M_{\lambda\mu}(\nu, p^2), \quad (9)$$

where

$$M_{\lambda\mu}(\nu, p^2) = \int d^4x e^{ikx} \langle 0 | T(V_\lambda^{\text{em}}(x) A_\mu^+(0)) | K^+(p) \rangle, \quad (10)$$

with $\nu = (kp)$ and V_λ^{em} the electromagnetic current.

The calculation of (10) has been carried out following the method developed for the axial-vector decay amplitude for $\pi^+ \rightarrow \gamma \ell^+ \nu$ within the chiral $SU(2) \times SU(2)$ symmetry.^[5] A generalization to the chiral $SU(3) \times SU(3)$ is straightforward, and we simply state the result:

$$M_{\lambda\mu}(\nu, 0) = F_K g_{\lambda\mu} + G(\nu g_{\lambda\mu} - p_\lambda k_\mu) + \mathcal{O}(\nu^2), \quad (11)$$

where the axial vector form factor G is given by

$$G = \frac{1}{F_K} \int dm^2 \frac{\rho_A(m^2) - \rho_V(m^2)}{m^4} + \frac{1}{3} F_K \langle r^2 \rangle_K \quad (12)$$

Here, $\langle r^2 \rangle_K$ is the mean squared charge radius of the charged kaon, and ρ_A , ρ_V are the axial-vector and vector two-point spectral functions, respectively.

In the process of calculation, terms proportional to m_K^2 have been dropped.

The uncertainties introduced thereby in the axial-vector amplitude are expected to be of the order of $m_K^2/m_Q^2 \approx 0.15$. This is to be compared with the uncertainties of the order of $m_\pi^2/m_{A_1}^2 \approx 0.02$ for $\pi^+ \rightarrow \gamma e^+ \nu$ (see Ref.[5]).

As is expected, the amplitude that results from the constant term, $F_K g_{\lambda\mu}$, in (11) is the same as that which follows from the modification of the PCAC relation in the presence of the electromagnetic potential (see Fig. 2(a)), i.e.

$$A_\mu^+(x) = -F_K (\partial_\mu - i e A_\mu(x)) K^+(x) \quad (13)$$

It can be shown that this amplitude cancels with a term in the inner bremsstrahlung amplitude corresponding to the emission of the photon by the outgoing ℓ^+ (see Fig. 2(c)).

The axial-vector decay amplitude for the structure radiation, analogous to (6), is now given by

$$M_3^{SR}(K^+ \rightarrow \gamma \ell^+ \nu) = \frac{G_F e \sin \theta_c}{\sqrt{2}} G(p_\lambda k_\mu \gamma g_{\lambda\mu}) \epsilon^\lambda \bar{u}_\nu \gamma^\mu (1 - \gamma_5) v_\ell \quad (14)$$

We evaluate (12) in the vector-dominance approximation, where⁴

$$\int dm^2 \frac{\rho_V(m^2)}{m^4} = \frac{1}{3} F_K^2 \langle r^2 \rangle_K \quad (15)$$

and with $\rho_A(m^2)$ saturated with the strangeness-carrying axial-vector meson Q:

$$\rho_A(m^2) \cong G_Q^2 \delta(m^2 - m_Q^2) \quad (16)$$

with^[12]

$$G_Q^2 \cong G_{K^*}^2 \cong 2m_{K^*}^2 F_K^2 \quad (17)$$

This makes use of the (approximately experimentally satisfied) relation

$$m_Q \cong \sqrt{2} m_{K^*} \quad (18)$$

and leads to ($m_{K^*} = 892$ MeV)

$$G \cong \frac{1}{2} \frac{F_K}{m_{K^*}} = 0.046 m_K^{-1} \quad (19)$$

Combining (8) and (19) we find

$$m_K(\gamma + G) \cong 0.13 \quad (20)$$

$$\gamma \equiv G/\gamma \cong 0.55 \quad (21)$$

These compare quite favorably with the experimental results^[6]

$$m_K |\gamma + G| = 0.153 \pm 0.011 \quad (22)$$

as well as exclusion of the region

$$-1.8 < \gamma < -0.54 \quad (23)$$

Thus we see that kaon PCAC, current algebra, and the axial-vector anomaly seem to provide an adequate description of the decay $K \rightarrow \gamma e \nu$, perhaps at least as good as for the analogous decay $\pi \rightarrow \gamma e \nu$, where theoretically^[5]

$$\gamma_\pi = 0.58 \quad , \quad (24)$$

as compared to the experimental value^[13]

$$\gamma_\pi = 0.44 \pm 0.12 \quad \text{or} \quad -2.23 \pm 0.12 \quad . \quad (25)$$

Note that because Ref.[5] used CVC they could not predict the sign of γ_π , while the axial-vector anomaly yields an unambiguous sign.

Finally, for the sake of completeness we write down the inner bremsstrahlung amplitude (see Fig. 2(b) and (c)):

$$\begin{aligned} M_{(K^+ \rightarrow \gamma \ell^+ \nu)}^{IB} = & -m_{\ell} F_K \frac{G_F e \sin \theta_c}{\sqrt{2}} \bar{u}_\nu(\ell') (1 + \gamma_5) \\ & \times \left[\frac{(p \epsilon)}{(p k)} - \frac{(\ell \epsilon)}{(\ell k)} + \frac{(\gamma k)(\gamma \epsilon)}{2(\ell k)} \right] v_\ell(\ell) \quad . \end{aligned} \quad (26)$$

This is manifestly gauge invariant, as are M_V^{SR} and M_A^{SR} individually. The total decay amplitude is the sum $M_V^{SR} + M_A^{SR} + M^{IB}$. Because of the appearance of m_ℓ in (26), M^{IB} is negligible for $K \rightarrow \gamma e \nu$ but dominant for $K \rightarrow \gamma \mu \nu$.

Acknowledgements

We would like to thank M. Suzuki for helpful discussions and suggestions. One of us (KAM) is grateful to the Aspen Center for Physics and the other (WWW) to the Lawrence Berkeley Laboratory for hospitality extended to them during the course of this work.

Footnotes

1. See Ref.[1]. It was only with the advent of the concept of color that the coincidence between nucleon and quark models in this regard was recognized.
2. "Structure radiation" is distinguished from "inner bremsstrahlung" in which the photon is radiated from the external kaon or electron.
3. The non-renormalization of the anomaly in quantum electrodynamics was shown in Ref.[7]. The same for non-Abelian gauge theories was proved in Ref.[8].
4. Relation (15) is consistent with recent determinations of the charged kaon charge radius. See Refs.[10] and [11].

References

1. J. Schwinger, Phys. Rev. 82 (1951) 664; S. L. Adler, Phys. Rev. 177 (1969) 2426; J. Bell and R. Jackiw, Nuovo Cimento 60A (1969) 47;
S. L. Adler, Lectures on Elementary Particles and Quantum Field Theory, ed. S. Deser, M. Grisaru, and H. Pendleton, MIT Press, Cambridge, Mass. (1970), Vol. 1, p.1; R. Jackiw, Lectures in Current Algebra and Applications, ed. S. B. Treiman, R. Jackiw, and D. J. Gross, Princeton University Press, Princeton, New Jersey (1972), p.97.
2. D. Sutherland, Phys. Lett. 23 (1966) 384; M. Veltman, Proc. Roy. Soc. A301 (1967) 107; S. L. Adler, Phys. Rev. B137 (1965) 1022; R. A. Brandt and G. Preparata, Ann. Phys. (N.Y.) 61 (1970) 119.
3. S. Weinberg, Phys. Rev. Lett. 19 (1967) 1264; 27 (1971) 1688; A. Salam, Elementary Particle Theory, ed. N. Svartholm, Almquist, Stockholm (1968), p.367.
4. D. Yu. Bardin and E. A. Ivanov, Fiz. Elem. Chastits At. Yadra 7 (1976) 726 [English transl.: Sov. J. Part. Nucl. 7 (1976) 286].
5. T. Das, V. S. Mathur, and S. Okubo, Phys. Rev. Lett. 19 (1967) 859.
6. J. Heintze et al., Nucl. Phys. B149 (1979) 365.
7. S. L. Adler and W. A. Bardeen, Phys. Rev. 182 (1969) 1517.
8. W. A. Bardeen, Proc. XVI International Conference on High Energy Physics, Batavia, (1972), Vol. 2, p.295.
9. L. M. Chounet, J. M. Gaillard, and M. K. Gaillard, Phys. Rep. 4C (1972) 201.
10. E. Tsyganov, Proc. XIX International Conference on High Energy Physics, Tokyo (1978), p.315.
11. E. B. Dalley et al., Phys. Rev. Lett. 45 (1980) 232.

12. T. Das, V. S. Mathur, and S. Okubo, Phys. Rev. Lett. 18 (1967) 761.
13. D. A. Ortendahl, Lawrence Berkeley Laboratory Report No. LBL-5305 (unpublished); A. Stetz et al., Nucl. Phys. B138 (1978) 185.

Figure Captions

Fig. 1 Feynman diagrams for the anomaly.

Fig. 2 Feynman diagrams for the inner bremsstrahlung.

Fig. 1

Fig. 2