LA--11261-MS

DE88 009205

*A Toxicological Study of
Gadolinium Nitrate*

Jerry E. London

MASTER

ds
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

CONTENTS

	Page No.
ABSTRACT	1
I. INTRODUCTION	1
II. EXPERIMENTAL PROCEDURE	1
A. Source of Material	1
B. Skin Sensitization	1
C. Primary Skin Irritation	2
D. Eye Irritation	3
III. RESULTS AND DISCUSSION	3
A. Skin Sensitization	3
B. Primary Skin Irritation	3
C. Eye Irritation	4
REFERENCES	5

TABLES

I. Eye Irritation Responses in Rabbits Treated with Gadolinium Nitrate	4

A TOXICOLOGICAL STUDY OF GADOLINIUM NITRATE

by

Jerry E. London

ABSTRACT

The sensitization study in the guinea pig did not show gadolinium nitrate to have potential sensitizing properties. Skin application studies in the rabbit demonstrated that it was cutaneously a severe irritant. This material was considered an irritant in the rabbit eye application studies.

I. INTRODUCTION

As a part of the Toxicology Group (LS-1) applied program, gadolinium nitrate, hexahydrate was examined to define its toxic properties with the following tests: (1) skin sensitization, (2) primary skin irritation, and (3) eye (conjunctival) instillation. Acute oral toxicity has been reported to be greater than 5 g/kg.¹

II. EXPERIMENTAL PROCEDURE

A. Source of Material

The test material was supplied by WX-3 (Los Alamos National Laboratory) as white crystals. The material was stored in a tin can, with friction top at 25°C.

B. Skin Sensitization

Eight female guinea pigs, weighing 284-321 g each, were used in the treatment group. The animals were housed individually and fed commercial

laboratory stock diet ad libitum supplemented by daily cabbage, or apples because of their need for vitamin C.

The test compound solution was made to a concentration of 0.1% with saline. The compound was administered in a series of ten "sensitizing" injections into the lower back and flanks of the guinea pigs. Before each injection, the test sites were clipped free of hair with electric small-animal clippers. Injections were made randomly over the test area via a 1-cc tuberculin syringe fitted with a 25-gauge needle. The volume of the first injection was 0.05 ml and that of the other nine was 0.10 ml. At 24 h after injection, the reaction was scored for erythema (redness), height, and diameter. Redness and height were scored as described by Landsteiner and Jacobs;² the diameters of the reactions were measured in millimeters using micro calipers.

Two weeks after administration of the tenth sensitizing injection, the lower back and flanks of each guinea pig were clipped free of hair, and a challenge injection of 0.05 ml was administered. The reaction of each animal was graded 24 h later and compared with the reactions from the sensitizing injections.

C. Primary Skin Irritation

The Draize test³ was used to assess primary irritation. Six New Zealand white rabbits, weighing 2.7 - 3.8 kg each, were used in this test group. The back of each rabbit was electrically clipped free of hair 24 h before application of the compound. Two sites were abraded and two left unabraded. The compound was applied as a paste consisting of 500 mg in corn oil to each location. The test sites were covered with a gauze pad, and the entire back was covered with an adhesive plastic drape. Each rabbit was then placed in a rabbit jacket (Alice King Chatham, Los Angeles, CA). After 24 h, the wraps were removed, and each test site was visually scored for erythema and edema. Readings were recorded for 24, 48, and 72 h. The final irritation score is an average for the 24- and 72-h readings.

D. Eye Irritation

Six New Zealand rabbits, weighing 2.8 - 3.4 kg. each, were used in this facet of the study. Both eyes were checked for abnormalities before the compound was instilled. The compound (100 mg) was instilled into the conjunctival envelope of the right eye of each rabbit; the left eye served as a control. Two of the rabbits had the compound washed from the eye thirty sec after instillation; two, 5 min after instillation; and two did not have the compound washed from the eye. Each eye was graded for ocular lesions at 1 and 4 h on the day of the application and again at 24, 48, and 72 h. Of particular interest was whether the cornea, iris, and conjunctiva became inflamed. The procedure and grading system were taken from the Draize test.

III. RESULTS AND DISCUSSION

A. Skin Sensitization

The data collected for each guinea pig in the treatment group indicated that all challenge injection reactions were within the limits of the reactions during the sensitizing period. The guinea pig skin sensitization study did not show gadolinium nitrate to be a sensitizer. Each injection site developed necrosis. These areas were still present after the challenge dose.

B. Primary Skin Irritation

Gadolinium nitrate caused skin irritation on all rabbits. After 24 h, all rabbits exhibited edema and 4 rabbit backs had erythema. Two rabbits had grey-white areas. At 48 h, four rabbits displayed edema and 3 had erythema. The abraded areas on one rabbit were grey with hardened surfaces and the unabraded areas were white. Another rabbit had 2 hardened areas. After 72 h, four rabbits exhibited one or more of the following: grey or white hardened surfaces, necrosis, edema and erythema. The other 2 rabbits had edema and erythema. After 8 days, all rabbits exhibited effects of the material from slight (no hair) to deep depressed necrotic areas.

The primary irritation index was 6.6, which is considered a severe irritant.

C. Eye Irritation

All rabbits' eyes had redness, chemosis and discharge after 1 h. Three rabbits' eyes exhibited iris and cornea involvement after 4 h and 2 more rabbits experienced the same effect after 24 h. These 5 rabbits' eyes were positive for cornea and iris scores on day nine. All rabbits had heavy discharge with some redness and chemosis on day nine. Results are shown in Table I.

TABLE I

EYE IRRITATION RESPONSES IN RABBITS^a
TREATED WITH GADOLINIUM NITRATE

<u>Tissue Graded^b</u>	<u>Time to Score</u>		<u>Time to Score</u>		
	<u>1</u>	<u>4</u>	<u>1</u>	<u>2</u>	<u>3</u>
	<u>(hours)</u>		<u>(days)</u>		
<u>Wash at 30 sec</u>					
Cornea	0	12.5	17.5	12.5	5.0
Iris	0	2.5	5.0	5.0	5.0
Conjunctivae	4.0	10.0	12.0	11.0	10.0
<u>Wash at 5 min</u>					
Cornea	0	10.0	20.0	20.0	10.0
Iris	0	5.0	17.5	17.5	17.5
Conjunctivae	10.0	13.0	10.0	9.0	7.0
<u>No Wash</u>					
Cornea	0	10.0	10.0	10.0	2.5
Iris	0	4.6	2.5	2.5	2.5
Conjunctivae	10.0	12.0	10.0	8.0	8.0

---

^aTwo rabbits per wash conditions.

^bMaximum cornea response = 80; maximum iris response = 10; maximum conjunctivae responses = 20.

REFERENCES

1. Journal of Pharmaceutical Sciences, Vol. 54, 663 (1965).
2. K. Landsteiner and J. Jacobs, "Studies on the Sensitization of Animals with Simple Chemical Compounds," J. Exp. Med. 61, 643-656 (1935).
3. J. H. Draize, G. Woodward, and H.O. Calveny, "Methods for the Study of Irritation and Toxicity of Substances Applied Topically to the Skin and Mucous Membranes," J. Pharmacol. Exp. Ther. 82, 337-390 (1944).