DOE/ER/60246--5

DE92 016223

Progress Report No. DOE/ER/60246-5

Grant No. DE-FG02-84ER60246

With

Ohio State University Research Foundation

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

JMR

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

*Hydrological and Geochemical Response and Recovery in Disturbed Arctic Ecosystems -
Extension Request*

Background

Throughout the period from 1985 to 1992 the Department of Energy supported research on the hydrology and geochemistry of the headwater basin of Imnavait Creek has focused on the quantification of the input from atmospheric sources of biologically significant and other related chemical variables; the transport of these variables in surface and subsurface flow and their efflux from the basin; and the development of geochemical budgets. The acquisition of multi-year data sets (the longest and most detailed sets in the Arctic) have made it possible to define seasonal ranges and amplitudes; determine spatial and temporal relationships within the different flow compartments; to begin to model the pathways and rates of movement through and across different landscape units. The length of record has also made it possible to examine the quantity and influence of local and extra-regional additions.

In the proposed extension we seek funds to complete synthesis and the write-up of this large data base.

Progress To Date

The following paragraphs provide a general summary of research results and is followed by a listing of authored and co-authored papers and presentations resulting in part or whole from past funding.

Snow and rain are the principal forms of precipitation that add nutrients and other ions to the Imnavait Creek watershed and tussock tundra region of Alaska in general. The six-year data base for this area indicates the substantial variability in both the total ion load and its composition. The rain chemistry appears to reflect the composition of air masses principally marine from the southwest quadrant that have crossed extensive continental distances and that have acquired (in some years) elevated concentrations of NO_3^- , SO_4^{2-} , K^+ , Ca^{2+} , NH_4^+ and Na^+ , which are thought to represent forest fire additions. Since most of the fires are of natural origin such additions are considered natural. Aperiodic high

concentrations of SO_4^{2-} , Cl^- , NO_3^- are recorded in snowfall in some years and are thought to represent anthropogenic sources to the north.

Dust from the nearby Dalton Highway, aperiodically may be a source of cations, particularly Ca^{2+} in certain rain events and may be the principal source of the ion in dryfall. Dryfall is a significant source of K^+ , PO_4^{3-} and NH_4^+ .

In each of the years of record SO_4^{2-} and NO_3^- have been in excess of H^+ with H_2SO_4 being the principal source of free acidity. Only in 1988 was HNO_3 more important.

The Imnavait Creek area and the region generally does not appear to be significantly affected by anthropogenic sources. The limited comparison that can be made with other northern sites suggests it receives one of the lowest ionic inputs in North America and thus can provide an excellent base-line against which to measure future changes in air quality as it is measurable in precipitation induced by both regional and extra-regional sources.

Of the two principal forms of precipitation rainfall contributes the greatest proportion of the ionic input, but this ranges greatly from year to year (25 to 70% - 6700 to 117300 moles) and is not directly related to the total volume of rainfall.

Spatially the ionic composition of the snow ranges widely, and mean basin values of individual ions have ranged widely over the six years of measurement, particularly in the contribution of anions for which snow is the principal source in most years. The reason for this lies in the different strengths of anthropogenic sources, i.e., from the southeast industrial areas; the petroleum production area to the north and natural sources such as tropospheric dust and volcanic emissions, which are all controlled by air mass movement.

Snow melt runoff is quite different in composition than the snow due to chemical fractionation within the snow during the melt process. During overland and shallow subsurface flow the discharge composition is further modified by sorption and desorption between the solute and substrate and leaching of substrate materials. Nitrate, and to a lesser degree, sulfate, are retained by the vegetation and thawing soil and along with chloride are not significant in stream flow beyond the first several days.

The period of snow melt and its associated runoff is the most important hydrological and geochemical event of any year. It accounts for between 50 and 75 percent of the

seasonal discharge and between 60 and 100 percent of the total seasonal efflux of individual ions, much of which takes place in the early phase of the event.

With few exceptions, summer precipitation is light and tends to contribute more cations than anions. A considerable range in specific ion concentration is found among sampling periods within a given year and in mean concentrations between years. As with snow, it is the anion composition that shows the greatest temporal variability. Among the natural processes that influence this are forest fires south of the Brooks Range.

Following the recession phase of melt-off, flow may fall to very low levels or stop altogether because interflow that may be associated with thaw of the seasonally-frozen active layer is insufficient to produce a base flow. A protracted rainfall re-establishes flow that commonly increases in volume as the summer progresses. High intensity rains produce short duration peak flow events resembling in pattern to that of melt-off. The close correspondence between ionic concentration and discharge phase that characterizes the melt-off flow is not maintained. Only dissolved organic carbon and suspended solids show an increase and are probably both derived from the channel and flood plain. Trends in post melt-off concentration of cations are weak but consistent from year to year.

The Imnavait Creek tundra and plants are capable of retaining nearly all of the anions added by precipitation and consistently loses cations during the melt-off discharge and throughout the post-melt flow period; the only exception being a net retention of K^+ during post melt. The loss of cations represents a combination of biological and chemical weathering within the active layer.

The efflux of carbon ($DOC < POC < COC$) is closely tied to the discharge regime of the stream and although impressive in relation to ionic concentration is insignificant in terms of the quantity of stored soil carbon in the valley and that fixed by vegetation.

The concentrations of ions in the water increases with depth in the active layer, probably as a function of time in contact with the substrate, and local conditions such as degree of oxygen saturation, and decreased biological uptake. Mass flow studies using soil tracers and laboratory determination of hydraulic conductivity suggest vertical interchange by mass flow is slow but that lateral mass transport is rapid near the surface.

The extent of geochemical interchange between and among landscape units is most extensive during the melt-off event, but is a surface and near-surface phenomenon. As the summer progresses there may be little or no surface and near surface interchange and geochemical isolation of landscape units takes place especially in those farthest removed from positions proximal to drainages. Interflow in the deepening active layer is too slow to contribute to surface flow. Precipitation events, especially those that fill the detention storage capacity, reestablish the interchange between some or among all units for varying lengths of time. Interchange at depth in the active layer may continue well after the upper region has refrozen. The hydrological and geochemical template for the upland tundra is thus comparable to most temperate regions.

PROGRESS TO DATE

Published and In Press Papers

- Everett, K.R. and Ostendorf, B., 1988. Hydrology and geochemistry of a small drainage basin in upland tundra, northern Alaska in *Permafrost: Fifth International Conference, Proceedings*, vol. 1, K. Senneset, ed., Tapir Publ., Trondheim, Norway, pp. 574-579.
- Everett, K.R., G.M. Marion and D.L. Kane, 1989. Seasonal geochemistry of an arctic tundra drainage basin. *Holarctic Ecology*, vol. 12, pp. 279-289.
- Hinzman, L.D., D.L. Kane, R.E. Gliack and K.R. Everett, 1990. Hydrologic and thermal properties of the active layer in the Alaskan Arctic. *Cold Regions Science and Technology*, vol. 19, pp. 95-110.
- Kane, D.L., L.D. Hinzman, M.-k. Wood and K.R. Everett, 1992. Arctic hydrology and climate change. Chapter 3. in *Arctic Ecosystems in a Changing Climate*. Academic Press, Inc., New York, 37-59.
- Kane, D.L., L.D. Hinzman, C.S. Benson and K.R. Everett, 1989. Hydrology of Imnavait Creek an arctic watershed. *Holarctic Ecology*, vol. 12, 279-289.
- Marion, G.M. and K.R. Everett, 1989. The effect of nutrient and water additions on elemental mobility through small tundra watersheds. *Holarctic Ecology*, vol. 12, pp. 317-323.

- Oswood, M.W., K.R. Everett and D.M. Schell, 1989. Some physical and chemical characteristics of an arctic beaded stream. *Holarctic Ecology*, vol. 12, pp. 290-295.
- Walker, M.D., D.A. Walker and K.R. Everett, 1989. Wetland soils and vegetation, Arctic Foothills, Alaska. U.S. Department of the Interior, Fish and Wildlife Service, Research and Development Biological Report, vol. 89, No. 7, 89 pgs.

Papers in Press

- Everett, K.R. Precipitation chemistry at an Arctic Foothills site, Alaska. Chapter 10 in *Landscape Function: Implications for Ecosystem Response to Disturbance. A Case Study in Arctic Tundra*. Springer-Verlag Ecological Studies Series.
- Everett, K.R., D.L. Kane and L.D. Hinzman. Surface water chemistry and hydrology of a small Arctic drainage basin. Chapter 11 in *Landscape Function: Implications for Ecosystem Response to Disturbance. A Case Study in Arctic Tundra*. Springer-Verlag Ecological Studies Series.

Papers Submitted

- Everett, K.R. (Journal of Environmental Quality) Seasonal precipitation chemistry in northern Alaska.

Presentations with Published Abstracts

- Everett, K.R., 1989. Seasonal precipitation chemistry at Imnavait Creek, Alaska. *Proceedings, 40th Arctic Science Conference*, Fairbanks, Alaska, p. 10.
- Everett, K.R., 1986. Hydrology of a drainage basin in the Arctic Foothills, Alaska. Program Abstracts, *The Ohio Academy of Sciences, 95th Annual Meeting*, vol. 86, No. 2, pg. 50.
- Everett, K.R., Marion, G.M. and D.L. Kane, 1987. Hydrology and geochemistry of a small tundra basin. *Bulletin of Ecological Society of America*, vol. 68, No. 3, pg. 298.

END

**DATE
FILMED**

8 / 10 / 92