NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

DOE/ER/45373--4

DE92 015287

Technical Progress Report
"Single-Electron Charging Effects"
DE-FG02-91ER45373
Dec. 15, 1991

Steven T. Ruggiero
University of Notre Dame

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DMR

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

The status of our project on single-electron tunneling is, again, excellent. As outlined in our original proposal, a key goal for this project has been the development of a scanning tunneling instrument for the purpose of imaging individual particles and tunneling into these particles at high magnetic fields.

As discussed in our previous progress report, we have had continued success with the use of metal droplet layers to form single-electron tunnel systems. However, as discussed on our recent publication on the subject¹ tunnel systems containing a multitude of particles can be problematical in terms of the existence of finite particle size distributions and particle clustering.

Ag-droplet tunneling system. This is the system used at Notre Dame to tunnel into ultra-small particles. Both barriers are created by the RF sputter deposition of Al₂O₃.

In order to avoid these problems we have constructed a low temperature scanning tunneling microscope so that tunneling can proceed into single particles as shown above whereas previously tunneling occurred simultaneously into a multitude of particles.

Also as discussed in our previous report—and in our original proposal—microwave excitations of these systems have been made as a function of frequency, and can also be made in our new low temperature STM as shown below.

1. S.T. Ruggiero and J.B. Barner, "Particle-Size effects in Single-Electron Tunnel Systems," *Z. Physic*, 1991 (accepted for publication).

Schematic of microwave scanning tunneling microscope. The microscope is designed for operation at 1K in a field of 9T. The dual piezo design is used to cancel the effects of thermal expansion (to first order).

Again, STM measurements are being carried out with samples as illustrated in the first figure with the exception that the last metal and oxide layer will be eliminated (see accompanying next-year proposal). This will allow us to use silver particles as in previous work,² which also show excellent droplet-formation characteristics. At present the STM is now operational at room temperatures and we are in the process of qualifying the instrument at low temperatures (77/4.2K).

2. J.B. Barner and S.T. Ruggiero, Phys. Rev. Lett. 59, 807 (1987); J.B. Barner and S.T. Ruggiero, Phys. Rev. B 39, 2060 (1989).

The STM system is unique in that it fully computer controlled (digital), even to the extent that the computer is actually in the feedback loop of the system.³ Also, because we have used high resolution D/A boards (16 bit) the system is capable of both scanning surfaces with atomic resolution and collecting current-voltage curves for tunneling into a given spot on a surface. Shown below are recent results with the STM demonstrating its high resolution capabilities in the form of a picture of a gold surface⁴ and the current-voltage characteristic for tunneling into (in this case) an In-particle system. The final figure shown an STEM micrograph of typical In-particle sample for reference.

As discussed in our original and in the accompanying next-year proposal, of significant interest for these systems is the possibility of observing spin splitting of quantum states of the electrons (particle-in-a-box states)⁵ via resonant tunneling. These states should be visible for temperatures in the vicinity of 1K and in a magnetic field of 1-10T. The magnet for these experiments is now in the lab and ready for use on these experiments.

In summary then, over the past year we have put our digital, low temperature, microwave scanning tunneling microscope on line. The next step is (as discussed in the accompanying proposal) is the study of single-electron tunneling effects in single particles in the presence of high magnetic fields (9 T) and in the presence of microwave radiation.

3. M. Honkannen and S.T. Ruggiero, "Low Temperature Digital Scanning Tunneling Microscope for Microwave Applications," (to be sub. Rev. Sci. Instrum., 1991).

4. M. Honkannen and S.T. Ruggiero, "High Resolution Imaging of Metallic Surfaces with Digital Scanning Tunneling Microscope" (in prep. for Phys. Rev. Lett.).

5. D.V. Averin and A.N. Korotkov, J. Low Temp. Phys. 80, 173 (1990).

Shown is a current-voltage trace of an In particle using the digital STM in IV mode.

Shown is high-resolution image of a Au surface using the digital STM described in text.

Shown is a STEM micrograph of In particles used with digital STM as discussed above (large particles are 350Å in diameter).

Publications/Presentations

S. T. Ruggiero and J.B. Barner, "Particle-Size Effects In Single-Electron Tunnel Systems" (to be published, Z. Physic, 1991).

M. Honkannen and S.T. Ruggiero, "Low Temperature Digital Scanning Tunneling Microscope for Microwave Applications," (to be sub. Rev. Sci. Instrum., 1991).

M. Honkannen and S.T. Ruggiero, "High Resolution Imaging of Metallic Surfaces with Digital Scanning Tunneling Microscope (in prep. for Phys. Rev. Lett.)

S.T. Ruggiero, "Single-Electron Tunneling," presented at the NATO Advanced Institute on Single Charge Tunneling, Les Houches, France, March 8, 1991.

N.M. Jisrawi, S.J. Koester, Q. Lu, M.J. Honkannen, W. Porod, G. Bernstein, and S.T. Ruggiero, "Microwave Studies of Single Electron Charging Effects," Bull. Am. Phys. Soc. 36, 872 (1991).

**1992/1993 Continuation Proposal
"Single-Electron Charging Effects"
DE-FG02-91ER45373**

**Steven T. Ruggiero
University of Notre Dame**

As outlined in our original proposal, and discussed in the accompanying progress report for 1990/1991, a key goal in the development of this project has been the development of a scanning tunneling microscope for the purpose of imaging individual particles and tunneling into these particles at high magnetic fields.

As discussed in our progress report, we now have an STM in operation, producing both sample images and current-voltage characteristics. Having done this, one goal for the coming year will be to combine the STM with our high-field magnet to attempt the first experimental verification of energy quantization in mesoscopic systems.

The idea here¹ is that a strong magnetic field will break the degeneracy of electrons in the usual way as:

$$\epsilon_m \rightarrow \epsilon_m \pm (1-\rho)\mu H$$

where ρ parameterizes the strength of the spin-orbit interactions in the system. The important prediction, thus, is that spin splitting effects due particle quantization should be manifest in the current-voltage characteristics of these systems as a magnetic field is applied.

Because of the extremely small size of particles which can be achieved with metal droplets (the smallest investigated by our group has been 5 Å), and the fact that a single particle can be looked at once, we are in a very good position to unambiguously observe such effects. This would represent a very exciting milestone in the field of mesoscopic systems.

1. D.V. Averin and A.N. Korotkov, J. Low Temp. Phys. 80, 173 (1990)

Illustration showing STM tunneling into ultra-small-capacitance particles and associated current-voltage characteristic. The step-like behavior of the IV curve will give rise to a series of peaks in tunneling conductance.

Also as discussed in our previous report, and in our original proposal, microwave excitations of these systems have been made as a function of frequency, and can also be made in our new low temperature STM. Time permitting, we plan to continue this work in the coming year with a view toward the observation of Bloch effects in single particles.

From a technical standpoint these goals will entail the final low-temperature qualification of the STM, final check out of our high field magnet, and some minor electronic and software work.

**DATE
FILMED**

7/23/92