RECEIVED BY DTIC MAY 21 1969

NYO-3624-3
Final Contract Report

MASTER

TERRESTRIAL OCCURRENCE AND
DISTRIBUTION OF IODINE-129

Contract AT(30-1)-3624

between

Environmental Sciences Branch
Division of Biology and Medicine
U.S. Atomic Energy Commission

and

Department of Chemistry
Carnegie-Mellon University

There is no objection from the patent
point of view to the publication or
dissemination of this document:

Patent Group (Brookhaven)

By *218 Potter*

5/5
1969

RECEIVED
MAY 15 1969
PATENT GROUP (BROOKHAVEN)

Raymond R. Edwards,
Principal Investigator

Report Prepared at:
Department of Chemistry
CHAPMAN COLLEGE
Orange, California 92666

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

NYO-3624-3

TERRESTRIAL OCCURRENCE AND
DISTRIBUTION of IODINE-129

Raymond R. Edwards¹
and
Plinio Rey²

DEPARTMENT of CHEMISTRY
CARNEGIE-MELLON UNIVERSITY
PITTSBURGH, PENNSYLVANIA 15213

LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or

B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

¹Present Address: DEPARTMENT of CHEMISTRY
CHAPMAN COLLEGE
ORANGE, CALIFORNIA 92666

²Present Address: RADIATION CENTER
OREGON STATE UNIVERSITY
CORVALLIS, OREGON 97331

T A B L E   O F   C O N T E N T S

Abstract	ii
Preface	iii
I. Introduction	1
II. Iodine Geochemistry	3
III. Terrestrial I-129	6
A. Natural Production Mechanisms and Rates.	6
B. Expected I-129/I-127 Ratios from Natural Sources	9
C. Technological Production of I-129.	11
IV. Sample Collection and Processing	12
V. Analytical Procedures	14
A. Pre-Irradiation Chemistry	14
B. Irradiation	15
C. Post-Irradiation Chemistry	15
D. Radioactivity Measurements	15
VI. Results and Discussion	21
A. Evaluation of Triple Neutron-Capture Interference	21
B. I-129 in Mineral Sources of Iodine	23
C. Evaluation of Application to Geochronology	25
VII. Acknowledgements	26
References	29

ABSTRACT

This program was undertaken to explore the occurrence of the long-lived iodine-129 (half-life 1.6×10^7 years) in natural materials, expected from spontaneous and induced fission of uranium, and from cosmic-ray interaction with the atmosphere, meteorites, and similarly exposed materials, and the somewhat larger quantities introduced into the environment through atomic energy activities. This is feasible because of the extreme sensitivity of activation analysis as a means of detection and assay of this nuclide. Development of the assay techniques, and their application to the study of iodine (in geochemistry, ecology, and geochronology) have been explored.

Measurement of the I-130 activation product in iodine obtained from Pleistocene (<13 My) brines and from Chilean caliche have indicated a ratio of $I-129/I-127 \geq 5 \times 10^{-12}$ for these geologically young materials, in essential agreement with hypothetical estimates. A sample of iodine from a much older (>300 My) brine gave the anticipated negative result. A large number of additional biological and mineral specimens have been prepared for the activation analysis.

TERRESTRIAL OCCURRENCE AND
DISTRIBUTION OF IODINE-129

PREFACE

The research activities summarized in this report were carried out in the Department of Chemistry of Carnegie-Mellon University (formerly Carnegie Institute of Technology) under Contract No. AT(30-1)-3624 with the Environmental Sciences Branch, Division of Biology and Medicine, United States Atomic Energy Commission. The initial period of the contract began on December 1, 1965, resulting from a proposal submitted August 10, 1965. The final contract year began on November 1, 1967, but laboratory work was terminated during May, 1968, since the principal investigator accepted a summer appointment at Brookhaven National Laboratory, and subsequently (September, 1968) joined the faculty of Chapman College, Orange, California.

Interim reports were submitted under dates of June 30, 1966 (NYO-844-67) and June 30, 1967 (NYO-844-71) in connection with the annual report on Contract No. AT(30-1)-844 with the Chemistry Branch, Division of Research. During the period covered by these reports, the work under these two contracts was integrated under the Interdisciplinary Program in Earth and Astronomical Sciences. A progress report (NYO-3624-1) covering the first year of work was submitted under date of November 30, 1966.

This report encompasses background information from the proposal and earlier reports, describes subsequent experimental work to June 1, 1968, and summarizes experimental results for the entire period of the contract.

I. INTRODUCTION

Early interest (1,2) in the 16-million-year isotope of iodine (I-129) as a tool in cosmochemistry has persisted, with the focus of experimental effort on mass-spectrometric measurements of its stable daughter nuclide, Xe-129, principally in meteoritic matter (see, for example, the reviews by Kuroda (3) and Reynolds (4), and the report by Hohenberg et al (5) on xenon-iodine dating).

Direct measurement of the levels of iodine-129 itself in natural materials has not been pursued because of its low inherent specific activity (0.16 microcuries per milligram). The possibility of its detection by neutron activation analysis was first recognized by Purkayastha and Martin (6), who detected the nuclide in pitchblende, presumably as a product of (α, n)-neutron-induced fission of uranium-235.

Estimates of the natural levels of I-129 from spontaneous fission and from cosmic-ray production in atmospheric xenon, and an assessment of the utility of the isotope as a tracer, employing activation analysis for its detection and measurement, have been made by Edwards (7, Appendix I). Almost simultaneously, Studier et al. (8) published a detailed procedure for isolation of iodine (127 and 129) from natural materials, and described activation-analysis and activation-product measurement techniques capable of determining as little as 10^{-13} grams of I-129. Subsequent studies (9) have demonstrated the accuracy, precision, and high sensitivity of the measurement of I-129/I-127 ratios in a

variety of materials, including atmospheric precipitation, airborne particulate matter, human and other mammalian thyroid tissue, minerals, and marine flora.

Extension of this work to the study of pre-fallout and modern levels of I-129 in the dynamic geosystems and ecosystems in which iodine is a major participant has been the general scope of the program reported on here.

II. IODINE GEOCHEMISTRY

The geochemical cycle of iodine presented by Miyake and Tsunogai (10, Figure 1) appears to be generally in accord with the best available data, as presented by Goldschmidt (11) and Rankama and Sahama (12) for the atmosphere, the lithosphere, and the hydrosphere. Vinogradov (13) deals specifically with the marine biosphere, and a bibliography of publications of iodine in rocks, minerals, and soils has been published by the Chilean Iodine Educational Bureau (14).

The principal reservoir of iodine in dynamic equilibrium with the biosphere is of course the oceans, and the hydrospheric inventory value of 7×10^{16} grams is roughly in accord with accepted values for the concentration of iodine in sea water (15) and the volume of the oceans (12), $1.37 \times 10^{24} \text{ cm}^3$.

The inventory value for the atmosphere, $1.0 \times 10^{12} \text{ g}$, is considerably higher than expected from the recent work of Duce et al (16) and Lininger et al (17), who report the iodine content of aerosols over Boston, Massachusetts, and over the Hawaiian Islands to be in the range of $10^{-9} - 10^{-8} \text{ g m}^{-3}$ of air. Using the value of Verniani (18) for the atmospheric volume, $4 \times 10^{18} \text{ m}^3$, these data indicate an atmospheric inventory of only $10^9 - 10^{10}$ grams of iodine in aerosol form. It is of course possible, as suggested by Goldschmidt (11), that gaseous forms of iodine predominate in the atmosphere.

Figure 1. Inventory and annual turnover of iodine, from Miyake and Tsunogai (10).

If one assumes a steady state for the marine iodine concentration, and accepts the input values of Miyake and Tsunogai, the marine sedimentation rate can be equated to the sum of the rates of addition by volcanic activity and by weathering, 1.3×10^9 grams per year. Combined with their hydrospheric inventory value of 7.1×10^{16} grams, this implies a residence time of $\sim 5 \times 10^7$ years for iodine in the oceans. This is more than twice the mean lifetime of I-129 (2.4×10^7 years), suggesting that the secular equilibrium isotope ratio, I-129/I-127, in the marine environment (and in fresh sediments) should be established by the processes of production and radioactive decay in the terrestrial surface environment. Furthermore, the magnitude of the residence times is such that I-129 may be assumed to be identical chemically with common iodine as a consequence of chemical exchange reactions and of the relatively rapid reversible biospheric and atmospheric processes involving iodine.

III. TERRESTRIAL I-129

A. Natural Production Mechanisms and Rates

Natural production of terrestrial I-129 occurs by the processes of natural fission of heavy elements, principally spontaneous fission of U-238, and by the action of cosmic rays on stable elements, principally atmospheric xenon.

The fission contribution to the hydrosphere is made (a) by uranium in sea water, (b) by uranium in weathering rocks, and (c) through release of fission-product I-129 in volcanic processes. The global production rate of I-129 from each of these processes has been estimated (19), assuming the U-238 spontaneous fission decay constant to be $7.8 \times 10^{-17} \text{ y}^{-1}$ (20) and the spontaneous fission yield of the chain at mass 129 to be 0.012 percent, the upper limit established by Wetherill (21). The estimated values are presented in Table I.

Cosmogenic I-129 may arise (d) from direct interactions of primary or secondary cosmic radiation with atmospheric xenon, and (e) from cosmic ray- and solar proton-induced reactions in extraterrestrial matter which ultimately reaches the geosphere. The contribution from Xe spallation has been estimated (19), assuming one-third of the atmospheric xenon is exposed to an effective particle flux of $\sim 25 \text{ cm}^{-2} \text{ sec}^{-1}$, and an effective cross section for I-129 production of $\sim 10 \text{ mb}$. The "cosmic dust" contribution was estimated on the basis of Wasson's (22) values, since reiterated by Wasson, Alder, and Oeschger (23), for the radiation exposure of the material, and employing the estimated production rates of I-129 in chondrites tabulated by Kohman (24).

The resulting estimated values from these two sources are presented in Table I, and the equilibrium process is diagrammed in Figure 2.

Table I
NATURAL PRODUCTION OF I-129

Process	Matrix	Rate of Production (atoms per year)		Equilibrium Contribution to Hydrosphere (atoms)
Spontaneous fission	Sea Water	(a)	1.0×10^{17}	2.3×10^{24}
	Weathering rocks	(b)	1.5×10^{18}	3.6×10^{25}
	Volcanic matter	(c)	9.6×10^{18}	2.3×10^{26}
Cosmic ray spallation	Atmosphere (Xe)	(d)	2.4×10^{19}	5.8×10^{26}
	Extraterrestrial matter	(e)	3.7×10^{17}	8.9×10^{24}
All processes		(a)-(e)	3.6×10^{19}	8.6×10^{26}

An independent estimate of the relatively important contribution of cosmic-ray reactions on atmospheric xenon can be made by analogy with chlorine-36, which is produced principally by (n,p) reactions on Ar-36 and by less efficient reactions on the more abundant Ar-40. Lal and Peters (25) estimate the global inventory of Cl-36 to be 1.5×10^7 grams, or about 3×10^{29} atoms.

Assuming the elemental cross sections of argon and xenon for cosmic-ray production of Cl-36 and I-129, respectively, are similar, the terrestrial inventory of I-129 should be related to that of Cl-36 by:

$$N(\text{I-129}) = N(\text{Cl-36}) \frac{\tau(129)}{\tau(36)} \times \frac{(\text{Xe})}{(\text{Ar})},$$

where the τ values are the respective product mean-lives, and $(\text{Xe})/(\text{Ar})$ is the atomic abundance ratio (atmospheric) of the target noble gases ($\approx 10^{-5}$).

Figure 2. Natural Production and Loss of I-129 in the Hydrosphere Reservoir. (Production rates are in atoms/year; inventory in atoms.)

Solution of the equation yields $N(\text{I-129}) = 1.5 \times 10^{26}$ from atmospheric cosmic-ray reactions alone, corresponding to $r_a = 0.65 \times 10^{19}$ atoms per year, the same order of magnitude as the values in Table I and Figure 2. The values could be brought into perfect agreement by arbitrarily correcting for the greater isotope abundance values of the effective target isotopes in the case of xenon.

B. Expected I-129/I-127 Ratios from Natural Sources

The calculations described above indicate that the principal processes contributing to the natural terrestrial inventory of I-129 are cosmic-ray interactions with xenon in the atmosphere and volcanic release of fissiogenic I-129, which together contribute about 8×10^{26} atoms at steady state. The other processes considered contribute, in sum, only about a tenth of this quantity, about 6×10^{25} atoms.

If the total marine inventory of I-129 is taken to be 8.6×10^{26} atoms, and that of I-127, based on a concentration of $6 \times 10^{-8} \text{ g cm}^{-3}$ (15), as 3.9×10^{38} atoms, the marine ratio of I-129/I-127 is estimated to be $\sim 2.2 \times 10^{-12}$.

Since the cosmic-ray production processes will contribute directly to atmospheric precipitation, the isotope ratio will be variable and generally higher on the land mass surface and in fresh water than in the marine environment. These expected variations are depicted schematically in Figure 3.

Figure 3. Variation of the Isotope Ratio (I-129/I-127) in the Iodine Cycle.

C. Technological Production of I-129

As a fission product of uranium-235, I-129 has been produced in nuclear reactors in unknown quantities since 1942. Also unknown (and no doubt variable) is the fraction of I-129 so produced which is released to the atmosphere or (directly or indirectly) to the hydrosphere in the waste from the processing of spent nuclear fuels.

The yield of I-129 in the thermal fission of U-235 (0.9%) corresponds to a production rate of 1.25 grams per megawatt-year ($\sim 6 \times 10^{21}$ atoms per year per megawatt of fission). Thus the hydrospheric inventory of I-129 (Fig. 2) will be significantly affected only when it is contaminated with fission-product I-129 equivalent to $>10^5$ megawatt-years of operation. Incomplete mixing, however, will certainly yield abnormal I-129/I-127 ratios in the atmosphere, the sea and land surfaces, and in particular geographical locations.

The gaseous fission product Kr-85, whose yield in fission is comparable to that of I-129, has been measured in the atmosphere. Its concentration in 1964 reportedly (26) corresponded to the fission of ~ 100 tons of U-235, which would have yielded roughly 5×10^5 grams of I-129. Even if less than 10% of this were released to the environment, localized very high concentrations of I-129 would be expected.

IV. SAMPLE COLLECTION AND PROCESSING

Table II presents a summary of the current inventory of samples collected for I-129 analyses. It is evident from the table that additional specimens of marine origin uncontaminated by the technological fission-product material are needed in order to establish (or set limits on) the I-129/I-127 ratio in the natural marine environment. The two old Japanese whale thyroids listed (1912 and 1915) were obtained from preserved fetuses (~ 30 -40 cm long) in the Japanese National Museum at Tokyo. These samples are being held in reserve until the analytical procedures and measurement methods are operating at high levels of confidence and at extreme sensitivity.

Other specimens have been processed to I_2 as obtained and approximately 50 are now ready for irradiation. Most of these are in the range 10-100 mg I_2 and thus, for I-130 measurement, will require the more sensitive β - γ coincidence counting (see below). Other I_2 samples are available in gram quantities, as indicated in Table II, and are thus susceptible to the γ - γ coincidence technique described below.

Monitor (comparator) samples in the form of I_2 (I-129/I-127 $\approx 10^{-6}$) sealed in quartz have been obtained commercially.

Table II

SPECIMENS^a ACQUIRED FOR ANALYSIS FOR I-129

Type of Specimen	Geographical Source	Collection Date	Method of Extraction ^b
<u>Biological</u>			
Seaweed, Various	Maine, Nova Scotia, Prince Edward Island Zanzibar, Philippines, Malaysia, West Coast Portugal, West Coast Spain, West Coast Mexico, Peru, Chile	1961-1966	Alkaline fusion, followed by solvent extraction
Coral	Bikini	1946 (pre-test)	
Thyroid, whale	Japan Japan Australia California Bering Sea Antarctic	<u>1912</u> <u>1915</u> <u>1966</u> 1966 1966 1966-67	
Thyroid, Pharmaceutical	Commercial	1930's (Manufactured)	
<u>Mineral</u>			
Brine	California Michigan California Pennsylvania	1961, <u>1930</u> , <u>1963</u> , 1966 <u>1963</u> --	Commercial I ₂ ^c Commercial I ₂ Oxidation, solvent extraction
Iodyrite (AgI)	New South Wales New Mexico	-- --	Thermal decomposi- tion, sublimation
Chile Nitrate	Chile	1965	Commercial I ₂
<u>Atmospheric</u>			
Spanish Moss	Texas Gulf Coast	1966	Burning in O ₂ (see ref. 8)

^aNote: collection dates for pre-1945 specimens are underlined

^bFor details, see "Analytical Procedures"

^cAvailable in gram quantities.

V. ANALYTICAL PROCEDURES

A. Pre-Irradiation Chemistry

Iodine is obtained from biological specimens (seaweed, thyroid) by wet-ashing the specimen (20-50 grams wet weight) in a nickel crucible with $\text{NaOH-Na}_2\text{CO}_3$ (10-20 grams each). Following the ashing procedure, the mix is dissolved in 1 M HNO_3 to give a final concentration of 0.1 M HNO_3 and the solution is cleaned by filtration or centrifugation. Sodium bisulfite is added to reduce all iodine to I^- , then an excess of 1 M NaNO_2 is added to oxidize to I_2 . This is extracted into CCl_4 , back extracted into 0.1 M NaHSO_3 , and precipitated by an excess of Pd(II) as PdI_2 .

The PdI_2 , after drying at 110°C , is transferred to a quartz U-tube open at one end. The open end is sealed, and the PdI_2 decomposed thermally ($>350^\circ\text{C}$). The resulting I_2 vapor is condensed in the cold arm of the tube, which is then sealed off, constituting the irradiation sample.

Mineral brines yield I_2 for CCl_4 extraction by simple oxidation [by Fe(III)] in the acidified brine. The remainder of the procedure is as described above for biological specimens.

Silver iodide-rich minerals, such as iodyrite, are susceptible to thermal decomposition ($>550^\circ\text{C}$), and the resulting I_2 vapor can be condensed on a water-cooled cold finger and transferred directly to quartz for irradiation.

Mineral iodine samples received as I_2 probably require no further treatment, but generally are sublimed before analysis to reduce

the probability of formation of I-130 by reactions other than (n, γ) on I-129.

B. Irradiation

Samples of I_2 , sealed in quartz, are irradiated for 12.5 hours (one half life of I-130) in a well-moderated neutron flux. Comparator samples, containing iodine with a known I-129/I-127 ratio are irradiated in the same vessel with the unknowns.

C. Post-Irradiation Chemistry

The irradiated quartz vials containing I_2 are broken and crushed in a mortar under an excess of 1 M $NaHSO_3$ solution. Bromine-water is added to provide Br holdback, the solution is added to oxidize I^- to I_2 . This is extracted into CCl_4 , back-extracted into $NaHSO_3$ solution, and more Br_2 added. This process is repeated five times, the final $NaHSO_3$ solution is acidified to 0.5 M HNO_3 , and an excess of silver nitrate solution is added to precipitate AgI. The precipitate is collected on a tared 42-mm filter paper, washed with 1 M HNO_3 , alcohol, and ether, dried under an infra-red lamp, weighed, and mounted for counting.

D. Radioactivity Measurements

Neutron irradiation of natural iodine containing traces of I-129 leads to the formation of the 25m I-128 by (n, γ) on I-127, and the 13.2d I-126 by $(n, 2n)$ on I-127, as well as to the desired product, 12.5h I-130, formed by (n, γ) on I-129. The decay modes of these three activation products are shown schematically in Figure 4. Earlier measurements have utilized the unique γ sum (>1.7 MeV) either alone (8)

or in coincidence with β^- (9) to distinguish instrumentally the I-130 from the other products. The β - γ coincidence technique, with a fast (~ 100 nsec) coincidence circuit to minimize chance sums, is reported (9) to be 100 times more sensitive than simple γ -sum counting, because of the large background reduction obtained in the coincidence system.

A limitation of β - γ coincidence counting is imposed by self-absorption of the β^- particles in large specimens of iodine. For this reason, it was considered advisable to explore a γ - γ coincidence measurement method to determine its sensitivity for I-130 measurements in gram-size samples of irradiated iodine.

γ - γ Coincidence Counting. Inspection of Figure 4 reveals only two coincident γ -rays unique, within the resolving power of a NaI(Tl) spectrometer, to I-130 decay in the presence of I-126. These are the 1.15 MeV and the 0.538 MeV γ 's, which were chosen initially for γ - γ coincidence counting. Two 3" NaI(Tl) crystals were calibrated with Na^{22} and Zn^{65} sources, and each single-channel analyzer was set to accept pulses only in the region of one of these γ energies (1.10-1.20 MeV and 0.51-0.57 MeV). Two scalars were set to accept pulses separately from the single-channel analyzers, and the third scalar to receive only pulses coincident within ~ 100 nsec in the two single-channel circuits.

In this mode, the coincidence background was found to be 0.5-0.6 cph, negligible (in samples containing more than decigram quantities of iodine) relative to the interference level of I-126 continuum coincidence counts.

More recent measurements have employed the coincidence of the 0.538 MeV gamma ray with those in the region of 0.669 to 0.743 MeV, the latter pair having a combined intensity of 1.85 γ 's per disintegration. This has resulted in considerable improvement in counting efficiency for I-130 over that experienced where the less abundant ($\sim 20\%$) 1.15 MeV was counted in coincidence with the 0.538 MeV γ . The I-126 decay scheme shows no true coincidences which would interfere with the 0.538 - 0.669-0.743 coincident pair, although Compton coincidences are expected (Figure 4).

β - γ Coincidence Counting. Predominantly, the quantities of iodine obtained from the natural materials listed in Table II will be small compared with the requirements of the γ - γ coincidence measurement outlined for I-130. Consequently, suitable equipment for a β - γ coincidence system has been assembled. The $4\pi\beta$ - γ coincidence arrangement consists (Figure 5) of a flat ($2\frac{1}{2}$ " X $3/8$ ") flow counter (proportional or Geiger) for β 's, positioned between two 4" X 4" NaI(Tl) crystals. For I-130 counting, this arrangement could be employed with a fast (~ 200 nsec) coincidence circuit, one channel accepting sum pulses generated by total γ -absorption of >1.5 MeV, and the other channel receiving the β^- pulses.

By biasing out all energies below 1.5 MeV, as indicated in Figure 4, it will be possible to measure I-130 with little interference from I-126, since the intensity of the 1.68 MeV cascade from this isotope is less than 0.28%. The I-128 (25 min) formed during irradiation will also be completely biased in the remote possibility that some might remain present at the time of measurement. The I-130 will be counted through the 1.60 MeV and higher cascades up to 2.34 MeV as shown in

Figure 4. This arrangement would be similar in principle to that previously reported (9), but might permit a somewhat greater areal distribution of the counting sample, thus reducing counting losses due to self-absorption in the more massive samples.

VI. RESULTS AND DISCUSSION

Experiments completed have employed iodine samples of 1-4g I_2 with the I-130 measurements made by γ - γ coincidence counting of AgI following post-irradiation chemistry.

A. Evaluation of the Triple Neutron-Capture Interference

Of the nuclear reactions other than $I-129(n,\gamma)I-130$ which might lead to interference in the I-129 activation analysis, only one can occur in pure iodine:

stable $I-127(n,\gamma)_{25m}I-128(n,\gamma)1.6 \times 10^7y I-129(n,\gamma)12.5h I-130$

$$\sigma = 6.4b$$

$$\sigma = ?$$

$$\sigma = 27b$$

An experiment was performed to evaluate the unknown neutron-capture cross section of the $_{25m}I-128$, hence to permit an estimate of the levels of neutron flux, sample size, and irradiation time at which I-130 would be produced from I-127 at levels detectable in the γ - γ coincidence measurement system. A 1.5g sample of iodine (from Chilean nitrate, 1965 production) was irradiated near the core of the Argonne CP-5 reactor (flux $\sim 2 \times 10^{-13} \text{ cm}^{-2} \text{ sec}^{-1}$) for approximately two weeks, receiving a reported nvt of $\sim 2 \times 10^{19} \text{ neutrons cm}^{-2}$.

Analysis of the γ - γ coincidence decay curves, and comparison with a sample of known I-129 content irradiated simultaneously, indicated that these conditions of irradiation produced $1.1 \times 10^{-9} \text{ g I-129}$, or $\sim 0.7 \times 10^{-9} \text{ g I-129/g I-127}$. Assuming a steady-state I-128 level throughout the irradiation, and accepting the 6.4b cross section for

I-127, this production rate corresponds to a capture cross section of ~ 100 b for the ^{25}m I-128.

This value is in agreement with that reported previously (27) on material irradiated in the MIT reactor. An independent measurement has been made by Lindstrom and Arnold (28), with iodine irradiated in the MTR, who obtained a comparable value (~ 110 barns). It should be noted that none of the three irradiations was performed in a well-thermalized neutron flux, so the cross-section values obtained are for "pile" neutrons rather than thermal.

Assuming a cross section of 100 barns, solution of the rate equation for I-130 production in the experimental conditions normally employed in the I-129 analysis (flux of $\sim 5 \times 10^{12}$, irradiation time of 12.5 hours) indicates that the triple neutron capture interference would simulate a ratio I-129/I-127 of $\sim 7 \times 10^{-13}$, about a factor of three below the sensitivity limit for γ - γ coincidence measurements with few-gram samples of iodine, and a factor of 6 to 14 below the positive values observed experimentally (Table III).

B. I-129 in Mineral Sources of Iodine

As indicated in Table II, gram quantities of iodine produced before 1940 were available from Chilean nitrate (1939) and from Long Beach, California brine (1930). Commercial iodine of recent production from these sources, as well as from the Monroe, Michigan brines, is also available. A single sample of Merck I_2 , Resublimed, marketed about 1920, was also examined.

Analyses of the pre-nuclear specimens, and of a specimen of 1966 Michigan brine iodine, have been completed. Results of the analyses are presented in Table III. An error of $\pm 50\%$ is assigned to the I-129/I-127 ratio on the basis of counting statistics alone at the extremely low levels of I-129 observed in the California brine and Chilean iodine. A meaningful value of the precision of the analysis at these levels can be obtained only by repeated analysis of samples of the same specimens.

The Long Beach, California brines are reported (29) to occur in Pliocene ($< 13\text{My}$) rock formations. Assuming their iodine was in equilibrium with hydrospheric and atmospheric processes at the time of trapping, the I-129 should have decayed by less than one-half. The analytical data thus would indicate an equilibrium isotope ratio $5 \times 10^{-12} < I-129/I-127 < 2 \times 10^{-11}$ for the Pliocene. This range is within less than a factor of ten of the value $\sim 2.2 \times 10^{-12}$ estimated above for the present-day hydrospheric equilibrium value.

The Chile nitrate deposits have not been reliably dated, nor is there agreement on the source and nature of their origin (14).

The analytical data for I-129 indicate that this iodine is little, if any, older than the Long Beach brines, probably Pliocene, possibly late Miocene, and suggest that their iodine is of marine (or biological), rather than mineral origin.

The Michigan brines (29) occur in rock formations of Silurian or Lower Devonian age (~ 400 My), and thus are expected to be depleted in I-129 from natural production processes. The analytical data support this expectation. While the specimen analyzed was produced in 1966, the production process involves "blowing-out" of iodine from the brine with recycled air, so that significant contamination from technologically produced I-129 in the atmosphere is avoided.

Table III

I-129 IN MINERAL SOURCES OF IODINE

Specimen	Date Produced	Age of Deposit	I-129/I-127 (observed ratio)
Brine, Long Beach California	1930	Pliocene (< 13 My)	1×10^{-11} 1×10^{-11} 5×10^{-12}
Chile Nitrate	1939	Unknown	5×10^{-12} 4.5×10^{-12}
Michigan Brine	1966	Silurian (400 My)	2×10^{-12} 3×10^{-12}
Merck I ₂	1920	(Probably Chile nitrate)	5×10^{-12}

The data in Table III are confirmed in order of magnitude by a recent report from Hanford (30). The results reported are:

	<u>I-129/I-127</u>
Pre-1936 human thyroid	4×10^{-11}
Pre-1942 I_2O_5	4×10^{-12}
Pre-1954 iodine	2×10^{-12}

The human thyroid tissue showed a higher ratio than the chemical iodine preparations, as would be expected (Figure 3) if the latter came from marine or mineral sources.

C. Evaluation of Application to Geochronology

Repeated analyses of the available specimens of iodine uncontaminated with the fission product material consistently show a significant difference between the geologically young iodine from the California brines and the very old material from Michigan, with the Chile nitrate apparently at an intermediate level. These results suggest the possibility of distinguishing old material from young, but the measured values are uncomfortably close to the sensitivity limit of the analysis for precise dating of specimens. With improved technique and activation-product measurement, approximate dating in the range 0-20 My might be feasible.

VII. ACKNOWLEDGEMENTS

All personnel associated with the project express their appreciation to the Division of Biology and Medicine of the U. S. Atomic Energy Commission for its sponsorship and support of the research through Contract No. AT(30-1)-3624.

The facilities employed for this work were shared with work in the Department of Chemistry under Contract No. AT(30-1)-844 with the Division of Research of the AEC. We acknowledge substantial assistance rendered by personnel associated with that project.

Mr. Michael J. Toia, Research Electronics Engineer, provided instruction and assistance in the γ - γ coincidence measurements and advice on specifications for a β - γ coincidence system.

Miss Christina Ann Switalsky, Mrs. Elizabeth Fang, and Miss Beth DeWalt, Laboratory Technicians, assisted in the chemical work.

Mr. Francis J. Bomba, Health Physicist, provided assistance in connection with the reactor irradiations.

Technical services were rendered by Laboratory Mechanics Mr. Charles W. Womer and George B. Klaus and by Scientific Glassblower Mr. Jan Zandhuis.

Secretarial and clerical services during the course of the research were provided by Mrs. Janet H. Muder, Mrs. June C. Johnston, and Miss Darlene Covaleski.

Special recognition is due Miss Sandi Hill of Chapman College for preparation of this manuscript and Figures 1, 2, and 3.

We are indebted to Mr. L. Strom, Mr. G. K. Moum, and Mr. Richard Carlson of the Special Materials Division, Argonne National Laboratory, for arranging thermal neutron irradiations of iodine samples at the Argonne CP-5 reactor.

We wish to express our appreciation for use of the Nuclear Engineering Test Facility at the Wright-Patterson Air Force Base, Dayton, Ohio, and to Lt. Max F. Thompson, Jr. for making the necessary arrangements for neutron irradiations.

The following persons and organizations are gratefully acknowledged as sources of valuable information concerning specimens of natural iodine-bearing materials and of commercially produced iodine chemicals, and as donors of the specimens themselves:

Dr. G. C. Egleson, Minerals Research, Dow Chemical Company, Midland, Michigan.

Mr. Richard O. Pfeiffer, Chilean Iodine Educational Bureau, New York.

Mr. Ivo Violic, Compania Salitrera Anglo-Lautaro, Antofagasta, Chile.

Dr. Bernard Keisch, Nuclear Science and Engineering Corporation, Pittsburgh, Pa.

Dr. Allen A. Wolman, Biologist, Fish and Wildlife Service, U.S. Department of the Interior, Richmond, California.

Dr. Dale W. Rice, Biologist, Fish and Wildlife Service, U.S. Department of the Interior, Seattle, Washington.

Dr. J. L. Bannister, Division of Fisheries and Oceanography, Western Australian Regional Laboratory, CSIRO, Nedlands, W.A., Australia.

Dr. Age Jonsgard, Universitetets Biologiske Laboratorium and Statens Institutt vor Hvalforskning, Oslo, Norway.

Mr. E. Vangstein, Norges Hvalfangstforbund, Sandefjord, Norway.

Dr. Hideo Omura, The Whales Research Institute, 1-3, Fukagawa Etchuzima,
Koto-Ku, Tokyo, Japan.

Dr. F. C. Fraser, British Museum (Natural History), London, S.W. 7.

Mr. M. A. Mackintosh, Whale Research Unit, National Institute of
Oceanography, British Museum (Natural History), London, S.W. 7.

Prof. Dr. E. J. Slijper, Zoologisch Laboratorium der Universitet van
Amsterdam, Netherlands.

REFERENCES

1. S. Katcoff, O. A. Schaeffer, and J. M. Hastings, *Phys. Rev.* 82, 688 (1951).
2. H. E. Suess and H. Brown, *Phys. Rev.* 83, 1254 (1951).
3. P. K. Kuroda, "Dating Methods Based on the Extinct Radionuclides I-129 and Pu-244." pp. 259-68 of Radioactive Dating and Methods of Low-Level Counting. Vienna, IAEA (1967).
4. J. H. Reynolds, "Isotopic Abundance Anomalies in the Solar System." *Ann. Rev. Nucl. Sci.* 17, 253-316 (1967).
5. C. M. Hohenberg, F. A. Podosek, and J. H. Reynolds, *Science* 156, 233-6 (1967).
6. B. C. Purkayastha and G. R. Martin, *Can J. Chem.* 34, 293 (1956).
7. R. R. Edwards, *Science* 137, 851 (1962).
8. M. H. Studier, C. Postmus, Jr., J. Mech, R. R. Walters, and E. N. Sloth, *J. Inorg. Nucl. Chem.* 24, 755 (1962).
9. R. C. Koch and B. Keisch, in L'Analyse par Radioactivation, D. Comar, ed., pp. 47-68, Presses Universitaires de France, Vendome (1964).
10. Y. Miyake and S. Tsunogai, *J. Geophys. Res.* 68, 3989 (1963).
11. V. M. Goldschmidt, Geochemistry, Oxford, The Clarendon Press (1953).
12. K. Rankama and Th. G. Sahama, Geochemistry, The University of Chicago Press (1950).
13. A. P. Vinogradov, The Elementary Composition of Marine Organisms, Memoir Number II, Sears Foundation for Marine Research, New Haven (1953).
14. Geochemistry of Iodine, Chilean Iodine Educational Bureau, Stone House, Bishopsgate, London E. C. 2. (1956).
15. E. D. Goldberg, *Ann. Rev. Phys. Chem.* 12, 29 (1961).
16. R. A. Duce, J. T. Wasson, J. W. Winchester, and F. Burns, *J. Geophys. Res.* 68, 3943 (1963).

References-2

17. R. L. Lininger, R. A. Duce, J. W. Winchester, and W. R. Matson, J. Geophys. Res. 71, 2457 (1966).
18. F. Verniani, J. Geophys. Res. 71, 385 (1966).
19. T. P. Kohman and R. R. Edwards, NYO-3624-1 (1966).
20. M. N. Rao and P. K. Kuroda, Phys. Rev. 147, 884 (1966).
21. G. W. Wetherill, Phys. Rev. 92, 907 (1953).
22. J. T. Wasson, Icarus 2, 54 (1963).
23. J. T. Wasson, B. A. Alder, and H. Oeschger, Science 155, 446 (1966).
24. T. P. Kohman, NYO-844-71, (1967).
25. D. Lal and B. Peters, "Cosmic-Ray Produced Radioactivity on the Earth," Encyclopedia of Physics, S. Flugge, ed., XLVI/2, Springer Verlag (1967).
26. D. Ehhalt, K. O. Minnich, W. Roether, J. Schlöch, and W. Stich, Paper A, Conf. 28/P/541, 3rd ICPUAE, Geneva (1964).
27. B. Keisch, R. C. Koch, and R. R. Edwards, Nuclear Science and Engineering Corp. Report No. 102, U.S.A.E.C. Contract AT(30-1)-3082 (1963).
28. R. M. Lindstrom and J. R. Arnold, University of California, San Diego, private communication.
29. G. C. Egleson, Dow Chemical Company, private communication.
30. F. P. Brauer, letter to R. P. Epple, BNWL-CC-12216, June 27, 1967.

Paper presented at the Southwest Regional Meeting
American Chemical Society; Little Rock, Arkansas, December 1967

121 IODINE-129 AND THE HISTORY OF MATTER. Raymond R. Edwards, Department of Chemistry, Carnegie-Mellon University, Pittsburgh, Pa. 15213.

Anomalous isotopic abundances of Xe-129 in natural xenon, attributed to its formation by beta decay of 16-million year I-129, have been widely used to date significant events in the history of cosmic matter, e.g., the time period between the end of nucleosynthesis and the onset of xenon retention by stony meteorites (see, for example, Merrihue, J. Geophys. Res. 71, 263, 1966; Kuroda et al, Nature 212, 241, 1966). Recent calculations by Kohman and Edwards (NYO-3624-1, 1966) suggest that cosmogenic I-129 in meteorites may reach 2×10^{-15} g I-129/g meteorite, and imply that the process of "nucleosynthesis" required to account for the observed Xe-129 excesses may have occurred at any time during the sojourn in space of meteoritic matter, requiring only an effective flux-time product about 100 times as great as that estimated for the past ten million or so years. Terrestrial Xe-129 excesses in Te ores (Inghram and Reynolds, Phys. Rev. 78, 822, 1950) and in natural gases (Butler et al, J. Geophys. Res. 68, 3283, 1963) have been variously explained as resulting from a recent (20 million years ago) galactic explosion (Tagaki et al, J. Geophys. Res. 72, 2267, 1967), or as the decay product of primordial I-129 incorporated in the earth at the time of its formation. Estimated terrestrial production rates of I-129 by cosmic ray action on atmospheric Xe, and by spontaneous fission, suggest a natural I-129/I-127 ratio of 2×10^{-12} in well-mixed systems, an order of magnitude which has now been confirmed experimentally by neutron activation analysis of iodine from geologically young terrestrial materials. The observed Xe-129 excesses may thus reflect inhomogeneous distribution of I-129 and/or recent trapping of daughter Xe-129. (Invited Paper)

Iodine-129: Its Occurrence in Nature and Its Utility as a Tracer

Abstract. Natural production rates of the 16 million-year iodine-129 from spontaneous fission and from cosmic-ray reactions are estimated to contribute a steady-state concentration of more than 10^{-14} gram of I^{129} per gram of I^{127} to the hydrosphere, the atmosphere, and the biosphere. These concentrations are expected to be detectable by neutron-activation analysis in natural materials that concentrate iodine. Substantial additional contributions have been made by fission-produced I^{129} in fallout, serving as an intrinsic tracer for natural iodine kinetics. Iodine-129 activation analysis has advantages as a tracer technique.

Iodine-129 occurs as a long-lived fission product of uranium, with a half-life of 1.6×10^7 years, in a (thermal neutron) fission yield of about 1 percent. This iodine isotope has found little use as a tracer because of its inherently low specific activity (0.16 $\mu\text{C}/\text{mg}$) and the low energy of its radiations (150 keV β^- , 38 keV γ). The low specific radioactivity results from the long lifetime (small decay constant, λ), a property which could make I^{129} particularly useful in many kinds of tracer experiments which require long periods of time to conduct. Further, the properties of low specific activity and low radiation energy are desirable from the standpoints of safe handling and effect on the environment (for example, on molecular species exposed to the radiations). The long lifetime not only permits experiments of essentially infinite duration but also permits long-term storage of labeled compounds and samples collected for assay.

The deficiencies of I^{129} as a tracer are successfully overcome if, instead of attempting to follow it by measuring its intrinsic radioactive decay processes as one customarily does with a radioactive tracer, one subjects matrices containing it to neutron-activation analysis. The isotopic capture cross-section for thermal neutrons has been set at between 27 and 35 barns (1, 2). The (n, γ) activation product is 12.6-h I^{130} , which has

readily detectable β and γ radiations. Other thermal neutron reactions either on I^{129} , on the naturally occurring I^{127} , or on neighboring elements, do not present serious interferences.

Iodine-129 thus can be characterized quite uniquely as a nuclide occurring in nature only to the extent it exists in secular equilibrium with current formation processes, which may thus be added to a chemical, physical, or biological system in which its inherent concentration is virtually zero, to serve as a "nonradioactive" tracer (in the radiological hazard sense) for the element iodine in any of its forms or for solvents or other media (water, air, organic materials, and so forth) into which it is introduced physically or chemically.

As a fission product I^{129} is produced in current-day fission reactors in multi-kilogram quantities, readily separable from the bulk of the radiologically hazardous fission products.

After neutron activation of an iodine-containing sample to saturation with respect to the 12.6-h I^{130} , the saturation activity will be given by the production rate of I^{130} atoms, $N\sigma\phi$, where N is the number of I^{129} atoms in the sample, σ is the isotopic capture cross-section of I^{129} (30 barns), and ϕ is the thermal neutron flux per square centimeter per second. The improved measurement sensitivity for I^{129} thus achieved by measuring the activation product at $\phi = 10^{14}$ is

$$\sigma\phi/\lambda_{130} = 2.3 \times 10^6,$$

corresponding to an increase in specific activity from 0.16 microcuries to 350 millicuries per milligram of I^{129} , or to a 3.5-picrourie sample of activation product from 10^{-14} g of I^{129} .

Other neutron-activation products following iodine chemistry are not expected to interfere with I^{130} measure-

ment, except, of course, the 25-m I^{128} , which will be produced ($\phi = 10^{14}$, $\sigma = 6.3$ barns) at a rate of 3×10^{12} per second per gram of I^{127} (natural iodine), and which will thus be the dominant iodine activity in most samples immediately after irradiation.

The half-life disparity (25 min versus 756 min), however, will lead to a reduction factor of over 10^{12} for I^{128} in a 17-hour post-irradiation cooling period, during which time the I^{130} activity will be reduced by a factor of only 2.5. Such a reduction factor will bring the I^{128} activity produced in a gram of I^{127} to the same order of magnitude as the I^{130} activity produced in 10^{-13} g of I^{129} .

Detailed information on the radiation spectra in I^{130} and I^{128} decay is available from the usual sources (3). Both are negative β emitters (some β^+ in I^{128}) and each has a complex spectrum of γ -rays. However, the highest energy γ -ray in I^{128} is 0.99 Mev (0.3 percent intensity), while that in I^{130} is 1.15 Mev (40 percent intensity). The two nuclides could be distinguished on this basis alone with a scintillation detector-pulse height discriminator arrangement. Measurement of characteristic γ - γ coincidences could be employed in the presence of considerable contamination by foreign activation products.

Levels in Nature

Prior to the nuclear age I^{129} will have been introduced into the atmosphere, lithosphere, and hydrosphere, presumably in a radioactive secular equilibrium situation, from cosmic-ray production and natural fission processes. If samples of suitable size or isotopic concentration, or both, are available, it will be possible to ascertain these equilibrium levels within the limits of sensitivity described above.

Using neutron-activation analysis, Purkayastha and Martin (1) measured the I^{129} content of pitchblende (from natural fission), and established a limit of $<10^{-8}$ atom of I^{129} per atom of I^{127} in a "natural" iodine sample, source unspecified. It should be possible now to explore readily the region between this value and $I^{129}/I^{127} = 10^{-14}$. While it is not possible to predict with confidence the levels that existed before the nuclear age, it would indeed be surprising if they lie below this limit in equilibrium samples which have concentrated iodine in recent geologic times.

The principal earth-level processes contributing fission-product elements to the lithosphere and hydrosphere in

modern geologic times are the spontaneous fission processes in uranium and thorium. Neutron-induced natural fission contributes relatively little additional (1), and this only in uranium-rich minerals.

Pre-fallout iodine obtained from marine plants and animals should be representative of natural I^{129} levels, since their environment represents a relatively homogeneous equilibrium. The uranium concentration in sea water is reported as $3 \mu\text{g/liter}$ (4). The yield of I^{129} in spontaneous fission of U^{238} is unknown, but may be assumed from the recent work of Kuroda *et al.* to be of the order of 0.1 percent (5). At a U^{238} spontaneous fission rate of $4 \times 10^{-3} \text{ g}^{-1} \text{ sec}^{-1}$, the steady-state I^{129} concentration would be $2.1 \times 10^{-18} \text{ g/liter}$ of sea water. Since the total iodine concentration in sea water is $6 \times 10^{-5} \text{ g/liter}$ (4), the I^{129}/I^{127} ratio is predicted to be about 3.5×10^{-14} , near the lower limit of the range of detectability for a 1-g iodine sample, easily detectable in a 10-g sample of marine iodine. Weathering processes will contribute approximately an equal quantity of the spontaneous fission product to the marine environment.

Contributions from spontaneous fission of U^{238} might be of equal or greater magnitude as a result of a shift in the mass-yield curve.

Production of I^{129} in the upper atmosphere obviously has occurred in pre-1945 processes whose contribution can be estimated in terms of reasonably assumed parameters. Assuming sufficiently energetic neutrons, protons, and γ -rays, reactions producing I^{129} will occur in Xe^{129} , Xe^{130} , Xe^{131} , and Xe^{132} .

If it is assumed that the atmospheric atom fraction for Xe is 10^{-8} and that the effective elemental cross-section for production of I^{129} is 10 mbarn, the steady-state surface density of the activation product will be about 10^6 atoms per square centimeter of the earth's surface. If the atmospheric and lithospheric residence times are short relative to the lifetime of I^{129} , this source will contribute an additional concentration of about 10^{-14} g of I^{129} per gram of I^{127} to the oceans.

The experimental evaluation of I^{129} levels arising from these reactions will be significant in developing the entire geophysical and biospheric history of I^{129} , and perhaps in highly significant dating measurements in the several-

million-year age range.

The occurrence of I^{129} as a fission-product provides a "built-in" tracer for the fate of fission iodine through the atmosphere, the hydrosphere, and the biosphere. Based on cumulative levels of Sr^{90} in soil prior to the recent tests (about $3 \times 10^{18} \text{ atom/mi}^2$) (6), I^{129} deposition from nuclear tests, integrated to mid-1960, should approximate 10^{19} atoms per square mile (1 percent fission yield), or about 10^7 atoms per square centimeter of the earth's surface.

If reasonable biological concentration factors are assumed, this material should now appear in selected biological samples in easily detectable quantities. The absolute sensitivity of this detection is greater by a factor of 3 for fission iodine inventories than for fission Sr^{90} inventories.

Since 1945, I^{129} has been added to the upper atmosphere in several substantial increments as a result of nuclear explosions. If lithospheric and hydrospheric entry are similar to those observed for Sr^{90} , this source will have added substantially to the pre-nuclear age base line described above. Its contribution of I^{129} to the oceans as a whole will have undoubtedly exceeded that of the natural processes by a large factor, and individual samples of marine iodine will show quite high I^{129}/I^{127} ratios as a result of incomplete mixing.

Analysis of biological samples dating back to 1945 should show these increments, and, when interpreted in terms of atmospheric inventories and fallout rates, should provide valuable information on the rates of entry of I^{129} into the biosphere.

If the analytical sensitivities described above are correct, analysis of iodine-rich specimens for I^{129} should provide a convenient measure of ages in the Cenozoic range. Petroleum brines and marine sediments should provide interesting samples.

The weapon-produced material provides a built-in long-term tracer for atmospheric transport and a soluble element for measuring fresh-water flow and oceanographic mixing rates. Its occurrence in fallout provides an intrinsic tracer for the cycling of iodine through the atmosphere and the oceans and for determining the contribution of fission processes to the total atmospheric iodine concentrations (7, 8).

The ultrasensitive analytical method for I^{129} also presents an opportunity for its measurement in meteorites, in

which it will have been formed by cosmic-ray spallation reactions on cesium, barium, and rare earth elements. It thus can serve as an additional monitor of the radiation history of meteorites, in a range of time periods between that of Be^{10} (2.7×10^6 years) and that of K^{40} (1.3×10^9 years).

Tracer Applications

As a deliberately added tracer, I^{129} opens many new vistas for tracer work, particularly as it relates to the tracing of iodine in any of its chemical forms, but also as it relates to atmospheric and hydrologic tracer studies.

The advantages of the I^{129} radio-tracer-activation approach to tracer studies over conventional tracing techniques are:

1) The long half-life of I^{129} (1.6×10^7 years) permits tracer studies of essentially infinite duration.

2) The long lifetime and low radiation energy of the nuclide result in virtual elimination of the radiological health hazard associated with the use of conventional radioactive tracers.

3) Problems encountered from radiation effects on the system under study and on labeled compounds are virtually eliminated; its applicability as a second label in I^{131} - or I^{125} -labeled compounds is obvious.

4) While I^{129} can be used in a very limited range of studies as a conventional radiotracer, utilizing direct measurement of its radiations, the radio-tracer-activation approach increases the measurement sensitivity by factors greater than 10^6 , so that quantities less than 10^{-18} g may be detected.

5) The radiotracer-activation method is superior to stable isotope tracing, inasmuch as the tracer nuclide does not occur in nature in abundances even remotely comparable with those of stable nuclides, or with quantities of the long-lived tracer which may be added to the system under study.

Viewed in combination, the advantages just listed and the detection sensitivity for I^{129} have implications for many different kinds of tracer studies. When introduced into any system, I^{129} behaves essentially as a nonradioactive substance insofar as environmental effects and nuclear stability are concerned. Samples of the system or of any of its parts may be taken at any time following introduction of the tracer, and subjected at will to neutron-activation analysis. The net result of the sensitivity is that a single gram of I^{129}

may be traced through processes which result in dilution factors up to 10^{18} . This is an equivalent dilution sensitivity to that attainable with about 350 curies of any radionuclide.

For full realization of the potentialities of the tracer applications described here for I^{129} , the base-level values (natural and artificial) described above should be established experimentally. Work is now in progress in our laboratories to obtain these values in pre-1945 and current biospheric, atmospheric, and hydrospheric iodine samples (9). Simple demonstration experiments of tracer applications of I^{129} in chemical systems (solubility of lead iodide), and in biological systems (with I^{125} as a

parallel tracer) (10) have been performed successfully (11, 12).

R. R. EDWARDS

Nuclear Science and Engineering Corporation, P.O. Box 10901, Pittsburgh 36, Pennsylvania

References and Notes

1. B. C. Purkayastha and G. R. Martin, *Can. J. Chem.* **34**, 293 (1956).
2. J. D. Garrison and B. W. Roos, *Nucl. Sci. Eng.* **12**, 115 (1962).
3. D. Strominger, J. M. Hollander, G. T. Seaborg, *Rev. Mod. Phys.* **30**, 585 (1958).
4. E. D. Goldberg, *Ann. Rev. of Phys. Chem.* **12**, 29 (1961).
5. B. T. Kenna and P. K. Kuroda, *J. Inorg. Nucl. Chem.* **16**, 1 (1960).
6. H. A. Knapp, *TID-13945* (1961).
7. R. A. Duce, J. T. Wasson, J. W. Winchester,

F. Burns, paper presented at the symposium on Trace Gases and Artificial Radioactivity in the Atmosphere, Utrecht, Netherlands, August 1962.

8. Since Br^{79} is "shielded" in fission by the 7×10^4 -yr Se^{79} , perturbations in the natural Br^{79}/Br^{81} ratios in upper atmospheric bromine might also be anticipated.
9. This work is supported in part by the U.S. Atomic Energy Commission.
10. R. L. Bogner, R. C. Koch, I. Gruverman, unpublished work.
11. Since preparing this manuscript, I have seen a prepublication copy of a paper from Argonne National Laboratory presenting a detailed analytical procedure for I^{129} (M. H. Studier, C. Postmus, Jr., J. Mech, R. R. Walters, and E. N. Sloth, *J. Inorg. Nucl. Chem.*, in press).
12. I acknowledge with pleasure stimulating discussions on this subject with R. A. Brightsen and other colleagues at NSEC, and with Professors Truman P. Kohman of Carnegie Tech. and Paul K. Kuroda of the University of Arkansas.

2 August 1962