DOE/PC/89772-T7

4TH QTR 1989

APPLICATIONS OF MICELLAR ENZYMOLOGY TO CLEAN COAL TECHNOLOGY

First Quarterly Report

DOE/PC/89772--T7

DOE Grant No. DE-FG22-89PC89772

DE92 016123

Submitted to:

Sayeed Akhtar
Technical Project Officer
PM-20, MS 922-206
U.S. Department of Energy
Pittsburgh Energy Technology Center
P.O. Box 10940
Pittsburgh, PA 15236

Prepared by:

Carol T. Walsh, Ph.D.
Principal Investigator
Boston University School of Medicine
80 E. Concord Street, L-603
Boston, MA 02118
617-638-4326

January 20, 1990

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

<u>Section</u>		<u>Page</u>
1	INTRODUCTION	1
1.1	Program Overview	1
1.2	Summary of Results	3
2	MATERIALS AND METHODS	4
2.1	Materials	4
2.2	Methods	4
2.2.1	Production of Reverse Micelle Solutions	4
2.2.2	Assay of Horseradish Peroxidase (HRP) Activity in Aqueous Solution	5
2.2.3	Assay of HRP Activity in Reverse Micelle Solutions	6
2.2.4	Quantitative Analysis of DBT Metabolites: HPLC Procedure	7
3	RESULTS	8
3.1	Production of Reverse Micelle Solutions with AOT in Isooctane	8
3.2	HRP Activity in Aqueous and Reverse Micelle Solutions	8
3.2.1	Aqueous	8
3.2.2	Reverse Micelles	10
3.3	Development of Analytical Procedure for DBT Metabolites	14
3.3.1	Separation of DBT Metabolites from AOT by Extraction	14
3.3.2	Analysis of DBT Metabolites in 0.1 M AOT-Isooctane by HPLC	15
4	PLANS FOR THE SECOND QUARTER	17
4.1	Analytical Procedures	17
4.2	Studies with HRP	17
4.3	Studies with Laccase	18
4.4	Studies with Microbial Extracts	18
5	BIBLIOGRAPHY	19

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
Figure 1:	DBT and Metabolites in Sulfur Oxidation Pathway	2
Figure 2:	HRP Activity in Aqueous Solution	10
Figure 3:	HPLC Chromatogram of DBT and its Metabolites in 0.1 M AOT-Isooctane with 2.3% H ₂ O	16

LIST OF TABLES

<u>Table</u>		<u>Page</u>
Table 1:	Stability of HRP Activity in Reverse Micelles as a Function of Enzyme Concentration	12

Section 1

INTRODUCTION

1.1 Program Overview

Full implementation of coal fuel sources will require more effective methods of providing "clean coal" as a fuel source. Methods must be developed to reduce the coal content of sulfur which significantly contributes to environmental pollution. This project is designed to develop methods for pre-combustion coal remediation by implementing recent advances in enzyme biochemistry. The novel approach of this study is incorporation of hydrophilic oxidative enzymes in reverse micelles in an organic solvent. Enzymes from commercial sources or microbial extracts are being investigated for their capacity to remove organic sulfur from coal by oxidation of the sulfur groups, splitting of C-S bonds and loss of sulfur as sulfuric acid (Figure 1). Dibenzothiophene (DBT) and ethylphenylsulfide (EPS) are serving as models of organic sulfur-containing components of coal in initial studies.

A goal of this project is to define a reverse micelle system that optimizes the catalytic activity of enzymes toward desulfurization of model compounds and ultimately coal samples. Among the variables which will be examined are the surfactant, the solvent, the water:surfactant ratio and the pH and ionic strength of the aqueous phase. Studies by several groups (Martinek et al., 1981; Kabanov et al., 1988) have shown that the surfactant AOT over a broad concentration range in organic solvents produces micelles, comparatively uniform in diameter,

Figure 1

DBT AND ITS SULFUR OXIDATION PRODUCTS

which incorporate hydrophilic enzymes. The activity (kcat) of certain enzymes in this system is higher than in aqueous solution. This surfactant is therefore being examined first, although the potential disadvantages of an SO_3 containing molecule have been recognized. Other surfactants to be tested include Tritons, Tweens and Brij 35.

1.2 Summary of Results

Among the accomplishments of the first quarter of this project are: study of the literature on coal chemistry and reverse micelle enzymology by the new P.I., Carol T. Walsh; hiring and training of a graduate student at Boston University to serve as research assistant, Majid Yazdani; development of interaction between the P.I. and Dr. Kitchell and her staff at DynaGen; meeting of the P.I. with DOE personnel including David Boron, Pat Barnett and Ian Campbell.

Procedures for monitoring enzyme activity in reverse micelle solutions have been worked out for one system, horseradish peroxidase (HRP) in 0.1 M AOT-isooctane with 13:1 H_2O :AOT and 4-aminoantipyrine as substrate. Reaction velocity was shown to increase as a function of incubation time and enzyme concentration. The oxidizing activity of HRP towards 4-aminoantipyrine in this system appeared to be less than that in aqueous solution. In order to test for HRP-catalyzed sulfur oxidation in reversed micelles, an HPLC method is being developed for measuring DBT sulfoxide, sulfone and dihydroxybiphenyl. Initial studies indicate some interference of the surfactant,

AOT, with the sulfoxide peak so the method is undergoing modification.

Section 2

MATERIALS AND METHODS

2.1 Materials

Studies were carried out with HRP, Type I RZ=1.20, obtained from Sigma Chemical Co. A more purified form of the enzyme (Type VI, RZ=3.1) was purchased from Sigma as well; this form of the enzyme is comparable to that which has been used in structural studies of HRP in reverse micelles (Balny and Douzou, 1979). Substrates for HRP assays included 4-amino antipyrine (Sigma Chemical Co.), phenol (Baker) and hydrogen peroxide (H_2O_2 , 30%, Fisher Scientific). Buffers included Tris HCl (Sigma Chemical Co.) and sodium phosphate (Fisher Scientific).

For formation of reverse micelle solutions the surfactant AOT, di(2-ethyl-hexyl)sodium sulphosuccinate, was obtained from Sigma Chemical Co. (The water content of the AOT has not been determined and therefore reported water content of micelle solutions do not account for the possible small contribution from this source.) Isooctane (Optima, Fisher Scientific) was used as organic solvent.

2.2 Methods

2.2.1 Production of Reverse Micelle Solutions

AOT (MW 444) was dissolved in isooctane to produce a 0.1 M solution. Aliquots of 36-144 μ l of 0.025 M phosphate buffer were then added to 2 ml of the organic solution to produce molar

ratios of H₂O/AOT of 10-40. Mixtures were examined for turbidity and absorbance readings in the 200-300 nm range measured.

2.2.2 Assay of HRP Activity in Aqueous Solution

The method for assay of HRP activity described in The Worthington Manual was used in initial studies. This procedure utilizes H₂O₂, 4-aminoantipyrine and phenol as substrates and entails monitoring the increase in absorbance at 510 nm as the reaction proceeds. Details are as follows:

- 1) Dilute 30% H₂O₂ 1:95 in distilled water and then this solution 1:50 in 25 mM Tris buffer (pH 7.1) or 25 mM phosphate buffer (pH 7.0). Flush all H₂O₂-containing vessels with N₂.
- 2) Under a hood dissolve 810 mg phenol in 40 ml distilled water, add 25 mg 4-aminoantipyrine and bring to 50 ml.
- 3) Make up HRP 1 mg/ml in distilled water and dilute 0.1 ml to 40 ml with distilled water.
- 4) Combine 0.7 ml of 4-aminoantipyrine solution and 0.75 ml of H₂O₂ solution in a cuvette, shake vigorously and measure absorbance at 510 nm every 15 sec for 5 min to determine nonenzymatic reaction rate (blank). Add 0.05 ml of HRP solution to substrate solutions for determination of enzymatic activity.
- 5) Absorbance was determined with an LKB Model 4050 spectrophotometer linked to an IBM-compatible PC. Software ("Reaction Rate") was used to collect absorbance data and compute reaction velocities.
- 6) Final concentrations in the standard reaction mixture were 1.17 mM 4-aminoantipyrine, 85 mM phenol, 0.88 mM H₂O₂ and 2.1×10^{-9} M HRP.

2.2.3 Assay of HRP in Reverse Micelle Solutions

The activity of HRP (Type I) in a reverse micelle solution was determined using the 4-aminoantipyrine-phenol assay technique. Final concentrations of reagents, based on total volume of the solutions, were approximately equivalent to those used in the aqueous assay. Partition coefficients of substrates between the organic and aqueous phases have not been determined however, so that actual concentrations of substrates in the aqueous microenvironment of the enzyme are not known. The activity of HRP in micelles was determined as follows:

1) 4-aminoantipyrine and phenol were dissolved in the organic phase, 0.1 M AOT in isooctane. To avoid precipitation of AOT, phenol was first added to isooctane (47.05 mg/10 ml) and AOT then added (0.444 mg/10 ml). The solution was gently heated to aid dissolution. 4-aminoantipyrine (2.377 mg) was then added. Higher concentrations of substrate could not be readily used because of solubility limits.

2) Two solutions containing reverse micelles were then formed by adding 0.237 ml of aqueous phase to 10 ml of organic (molar ratio of $\text{H}_2\text{O}:\text{AOT} = 13$). One aqueous component contained 30% H_2O_2 diluted 5000 fold with 0.025 M phosphate buffer. The second aqueous component contained 0.1 mg HRP/13.9 ml 0.025 M phosphate buffer ($1.8 \times 10^{-7}\text{M}$). After addition of the aqueous component, the two solutions were vortexed vigorously for 1 min and allowed to equilibrate for 10 min. This procedure is based on the method of Martinek et al. (1981) and is designed to allow

equilibration of the micelles, with respect to degree of hydration, before enzyme activity is assessed.

3) One-ml aliquots of the HRP and the H_2O_2 -containing solutions were then combined, vortexed for 10 sec and transferred into a cuvette. The absorbance at 510 nm was determined every 8 or 15 sec for up to 5 min. The extinction coefficient of the reaction product was not determined in the aqueous or reverse micelle solution. Enzyme activity is reported therefore as absorbance units per minute. Since the extinction coefficient could theoretically differ in the two types of solutions, definitive comparison of HRP activity cannot yet be made. It should be noted, however, that with other reaction products of HRP, such as purpurogallin in the pyrogallol assay, the extinction coefficient in 0.1 M AOT-isooctane is the same as in aqueous buffer (Martinek et al., 1981).

4) Final concentrations in the micellar reaction mixture were 1.17 mM 4-aminoantipyrine, 85 mM phenol, 0.21 mM H_2O_2 , 2.1×10^{-9} M HRP and $\text{H}_2\text{O}:\text{AOT}=13$, unless indicated otherwise.

2.2.4 Quantitative Analysis of DBT Metabolites: HPLC Procedure

DBT (0.1 mM), DBT sulfoxide (0.05 mM), DBT sulfone (0.05 mM) and dihydroxybiphenyl (0.05 M) were added to 0.1 M AOT in isooctane. The mixture was gently heated. Not all metabolites were completely solubilized. To one 10-ml aliquot, 0.237 ml of 0.025 M phosphate buffer was added to produce a reverse micelle solution. Samples were filtered through disposable syringe filter units (Rainin Instrument Co., No. 38-159, 0.2 μ pore size, 3 mm diameter Nylon 66 membranes). Filtrates (4 μ l) were

analyzed at DynaGen, Inc. with a Waters HPLC system containing a C18 resolve (5 micron spherical) column. The mobile phase contained tetrahydrofuran:acetonitrile:water (23:20:57) at a flow rate of 1.5 ml/min. The detector was a Waters Lambda Max 481 LC spectrophotometer which was set at 242 nm. The method, modified from Wyza et al. (1989), has been previously used by Dr. Kitchell at DynaGen, Inc. for analysis of DBT metabolites in organic solvents.

Section 3

RESULTS

3.1 Production of Reverse Micelle Solutions with AOT-Isooctane

Addition of phosphate buffer to 0.1 M AOT in isooctane rapidly resulted in clear, homogeneous solutions for H₂O/AOT molar ratios of 10 and 13. Samples with ratios of 18 became clear after 10 min. At a higher ratio (40), the sample was turbid with a foamy surface and phase separation occurred.

A wavelength scan of 0.1 M AOT in isooctane, measured using the organic solvent as the reference blank, indicated a broad strong absorbance band from approximately 200-245 nm with a peak at 227 nm. Solutions containing reverse micelles produced an absorbance peak at 238 nm, when 0.1 M AOT isooctane was used as the reference sample. Turbidity of the solution did not obviously alter the characteristics of the scan.

3.2 HRP Activity in Aqueous and Reverse Micelle Solutions

3.2.1 Aqueous

Under standard conditions of the enzyme assay in aqueous solution, reference blanks consistently generated reaction

velocities of 0.000 absorbance units per minute. Under all conditions tested in aqueous solution, the product formation was linear over the 5 minutes tested. The relationship of the 4-aminoantipyrine-phenol concentrations to enzyme activity was characterized by analysis of rates for concentrations ranging up to a 15 fold dilution. The results examined according to the Eadie-Hofstee transformation (Figure 2) suggested a $V_{\max}$ under these conditions of 0.095 absorbance units/min, a K_m of 0.185 mM for 4-aminoantipyrine and k_{cat} of $(4.5 \times 10^7 \text{ abs units/min/M enzyme})$. A 50% decrease in the H_2O_2 concentration was found to decrease enzyme activity to about 74% of the original value. A threefold increase in enzyme concentration, $2.1 \times 10^{-9}\text{M}$ to $6.1 \times 10^{-9}\text{M}$, produced an approximately threefold increase in activity from 0.05 to 0.19 abs. units/min.

3.2.2 Reverse Micelles

In initial studies of HRP in reverse micelles, the concentration of H_2O_2 in the aqueous phase (prior to addition to the organic solvent) was equivalent to that used in the aqueous assay, since it was assumed the peroxide would remain in the aqueous phase within the micelle. Negligible activity of HRP was detected however. Ten-fold higher concentrations of H_2O_2 resulted in detectable activity of approximately 0.010 absorbance units/min. A further five-fold higher concentration of H_2O_2 (0.967 mM) did not result in a further increase in enzyme activity (0.009 abs/min) but did produce higher nonenzymatic product formation (0.003 abs/min).

Figure 2: HRP Activity in Aqueous Solution

The activity of HRP was examined as a function of enzyme concentration. It should be noted that the molarity of HRP is computed on the basis of the total volume of the reaction solution. The lowest concentration tested, 2.1×10^{-9} M, is equivalent to the concentration used in the aqueous medium. The reaction rate was found to increase as a function of enzyme concentration (Table 1). The relationship was not linear; rate relative to concentration declined above 1.05×10^{-8} M. With the lower enzyme concentration, the reaction rate was approximately linear for 2.5 min and declined slightly during the subsequent 2.5 min. The curvi-linearity became marked at higher enzyme concentrations. At the highest concentration tested the absorbance declined after 1 min of the reaction, probably reflecting degradation of the product.

The basis for the decline in reaction rate despite the relatively low activity of HRP in the micellar as compared to the aqueous solution has not been established. One possibility is the availability of the substrates in the micellar core where the enzyme presumably is functional. In one experiment with 2.1×10^{-9} M HRP, the substrates (1.17 mM 4-aminoantipyrine and 85 mM phenol) were included in the aqueous phase added to the organic solvent. The reaction rate was comparable during the first 2.5 min but no decline was observed up to 5 min. A second possibility which may be important at higher enzyme concentrations is product inhibition of the enzyme.

Table 1: Stability of HRP Activity in Reverse Micelles
as a Function of Enzyme Concentration^a

HRP Concentration (M)	HRP Activity (Abs. Units/Min)	
	<u>Initial</u>	<u>After 24 hr at 4°C</u>
0	0.004 ± 0.001 ^b	0.004 ± 0.001
2.1 × 10 ⁻⁹	0.018 ± 0.001	0.010 ± 0.000
1.05 × 10 ⁻⁸	0.091 ± 0.015	0.044 ± 0.002
2.1 × 10 ⁻³	0.123 ± 0.001	0.056 ± 0.006
1.05 × 10 ⁻⁷	0.268 ± 0.090	0.215 ± 0.007

^aReverse micelle solutions contained 0.1 M AOT in isooctane with 13:1 molar ratio of H₂O to AOT.

^bMean ± S.D. for n = 2.

The stability of HRP in a reverse micelle solution was examined by monitoring baseline activity and activity after 24 hr at 4°C. These experiments were first attempted with HRP-containing reverse micelle solutions without 4-aminoantipyrine and phenol. However, baseline activity was low (0.005 + 0.001 abs. units/min), probably because the substrate concentrations in the reaction mixture were half the standard value. In the subsequent study the substrates were included in the micellar enzyme solution. The micellar solution was refrigerated for 24 hr to minimize autooxidation of the substrates. HRP activity was significantly lower after 24 hr (Table 1) at all enzyme concentrations tested. The reaction rate after 24 hr for the lowest enzyme concentration was similar, however, to baseline assays for comparable samples. Further study would be required to establish that the decline after 24 hr was attributable to alteration in enzyme function in the reverse micelle and not day-to-day variability in the assay. It should be noted that HRP in AOT-isooctane has been reported to retain its activity at least a week when pyrogallol is used as substrate (Martinek et al., 1981). Data has not been found which addresses the question of the stability of AOT in the presence of HRP AND H₂O₂. (Higher numbered Brij surfactants are susceptible to oxidation and are marketed containing antioxidants; these compounds will be avoided.)

Apparent kinetic parameters of HRP for these substrates in the reverse micelle solution (V_{max} , K_M , k_{cat}) have not been determined. One difficulty mentioned previously is the

solubility problem of the 4-aminoantipyrine phenol which have a lower affinity for HRP in aqueous media than other substrates such as pyrogallol. If the substrates are in excess so that velocities measured are close to $V_{\max}$ as was true for the aqueous assays, then k_{cat} ($= V_{\max} [\text{HRP}]$) can be compared for the lowest enzyme concentration ($[\text{HRP}] = 2.1 \times 10^{-9} \text{M}$). The k_{cat} of HRP in the reverse micelle solution, 8.5×10^6 abs. units/min/M HRP, is about one-fifth the value in aqueous solution, 4.5×10^7 abs. units/min/M HRP. This result is unlike that for HRP activity towards pyrogallol in this reverse micelle solution where k_{cat} exceeded the value found in aqueous solution and is about maximum for $\text{H}_2\text{O}:\text{AOT}$ of 13. Definitive comparison will require determination of the extinction coefficient of product in the aqueous and micellar solutions.

3.3 Development of Analytical Procedure for DBT Metabolites

3.3.1 Separation of DBT Metabolites from AOT by Extraction

For GC/MS analysis of experimental samples, a method is needed for reducing the concentration of AOT which is expected to impair column performance. Removal of AOT from 0.1 M solution in isooctane was attempted using 0.1 M KCl, 0.1 N NaOH or 0.1 N HCl. Samples were vortexed after addition of 1 part aqueous to 5 parts organic phase and then microfuged. The procedure was repeated three times. The acid wash was most effective in causing precipitation of AOT into the aqueous lower phase. The efficiency of the extraction was not quantified but considerable AOT was observed even in the third wash.

Based on these preliminary observations, an experiment was designed to determine the retention of DBT metabolites in the organic phase, after extraction of 0.1 M AOT in isooctane with 0.9% NaCl - 0.1 N HCl. The experiment could not be completed as designed but the following observations were made:

1) The solubility of DBT sulfone and dihydroxybiphenyl in isooctane was less than 0.1 mM.

2) Dihydroxybiphenyl, 0.1 mM, appeared to solubilize in 0.9% NaCl - 0.1 N HCl more readily than in isooctane.

3) The solubility of DBT in isooctane and in 0.1 M AOT-isooctane exceeds 1 mM.

4) To monitor separation of metabolites from AOT with UV absorbance, the peak at the higher wavelength of the metabolites (e.g. 280 nm for sulfoxide) should be used to avoid interference by AOT absorbance (200-245 nm).

3.3.2 Analysis of DBT Metabolites in 0.1 M AOT-Isooctane by HPLC

Initial results (Figure 3) suggest that HPLC will be a feasible method for quantitation of DBT metabolites in 0.1 M AOT-isooctane. Retention times of DBT, DBT sulfoxide, DBT sulfone and dihydroxybiphenyl in 0.1 M AOT isooctane (alone or with 2.3% 0.025 M phosphate buffer) were similar to values for compounds prepared in acetonitrile. Peak areas were considerably higher than expected, which will be explored. A problem with the method is some overlap of the signal from the sulfoxide (RT=1.42) and AOT (RT=2.15). Several approaches will be tried to improve the procedure.

FIGURE 3

4µl Sample (2) Anal. Version

CHANNEL A INJECT (22/90 12:49:46

Section 4

PLANS FOR THE SECOND QUARTER

4.1 Analytical Procedures

A major effort in the second quarter will be further development of analytical procedures for quantitation of DBT metabolites in surfactant-containing solutions. The preliminary HPLC procedure will be varied in several ways to reduce interference from the AOT surfactant:

- 1) Possibly a preinjection column can be used which removes AOT without affecting recovery of metabolites. The acidic sulfur group of AOT may represent a site which can be utilized for selective retention of the molecule.

- 2). The UV absorbance band of 0.1 M AOT in isooctane (compensated for isooctane) ranges from 200 to about 245 nm with a peak at 227 nm. A higher wavelength for the HPLC detector may improve resolution of AOT and sulfoxide without compromising the sensitivity of the assay. The UV peak of the metabolites at 280 nm may be preferable.

- 3) Previous work has shown that a slight difference in the ratio of components of the solvent system increases retention times and alters the separations of the metabolites. Solvent modifications for improving sulfoxide-AOT separation will be tried.

4.2 Studies with HRP

Once a suitable HPLC procedure is developed, studies will be carried out to determine the activity of HRP in reverse micelles toward DBT as substrate. Concentrations of DBT will be used that

would permit HPLC detection of 100-fold lower concentrations of metabolites. Initial studies will be carried out using reverse micelle parameters described in this report. Other surfactant-solvent conditions will be tested as well, since there are concerns about the appropriateness of a sulfur-containing AOT as surfactant and the relatively low solubility of the DBT metabolites in isooctane.

4.3 Studies with Laccase

Permission has recently been obtained from Professor Bengt Reinhammar, an expert on laccase (Reinhammar, 1984), to receive 100 mg of the enzyme which he has purified from Polyporus versicolor. The enzyme from this fungal source has been shown by Dr. Kent Kirk of the USDA to degrade lignin analogues. This enzyme preparation has not been studied in reverse micelles, but there is literature on another laccase in reverse micelles (Martinek, 1987). Experiments will be carried out to assess its activity in reverse micelles against a standard substrate (syringaldazine) and then with DBT as the substrate.

4.4 Studies with Microbial Extracts

Progress of Dr. Kitchell at DynaGen with desulfurization of DBT using microbial fractions (GB-1) is being followed. Once activity has been confirmed with GC/MS, attempts at enhancing activity by incorporation of microbial extracts in reverse micelles will be attempted. Lee and Yen (1988) have demonstrated the effectiveness of this approach for removal of total sulfur from coal, presumably mostly the pyritic form, by extracts of Thiobacillus ferrooxidans in reverse micelles.

5. BIBLIOGRAPHY

Balny, C. and Douzou, P. New trends in cryoenzymology: II - Aqueous solutions of enzymes in apolar solvents. Biochimie 61: 445-452, 1979.

Kabanov, A.V., Levashov, A.V., Klyachko, N.L., Namyotkin, S.N., Pshezhetsky, A.V. and Martinek, K. Enzymes entrapped in reversed micelles of surfactants in organic solvents: a theoretical treatment of the catalytic activity regulation. J. Theor. Biol. 133: 327-343, 1988.

Lee, K.-I. and Yen, T.F. Coal desulfurization through reverse micelle biocatalysis process. Proceedings of ACS Meeting, San Francisco, CA, 1988, pp. 573-579.

Martinek, K., Levashov, A.V., Klyachko, N.L., Pantin, V.I. and Berezin, I.V. The principles of enzyme stabilization vs. catalysis by water-soluble enzymes entrapped into reversed micelles of surfactants in organic solvents. Biochim. Biophys. Acta 657: 277-294, 1981.

Martinek, K., Berezin, I.V., Khmel'nitski, Y.L., Klyachko, N.L. and Levashov, A.V. Coll. Czech. Chem. Commun. 52: 2589, 1987.

Reinhammar, B. Laccase. In: Copper Proteins and Copper Enzymes, Vol. III, Lontie, R., ed., CRC Press, Inc., Boca Raton, FL, 1984.

Wyza et al. Presented at Microbial Coal Desulfurization Workshop, DOE Contractors Review Meeting, August, 1989.

**DATE
FILMED**
8/27/92