NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

PARTICIPATION IN HIGH ENERGY PHYSICS

TASK D

DOE/ER/40560--3D

DE92 016291

PROGRESS REPORT

FOR PERIOD MAY 1, 1991 - APRIL 30, 1992

LEON M. LEDERMAN

THE UNIVERSITY OF CHICAGO
5801 SOUTH ELLIS AVENUE
CHICAGO, IL 60637

APRIL 1992

PREPARED FOR

THE U.S. DEPARTMENT OF ENERGY

AWARD NO. DOE DE F602 290ER40560 TASK D

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Table of Contents

Progress Report

A. Communication and Advancement of High Energy Physics

B. Experimental High Energy Physics

Part I B-Quarks

Part II Secondary Vertex Trigger

1. David Chen's Thesis

2. E789 Work

C Science Education

1) Fermilab

2) IMSA

3) TAMS

REPORT ON DOE RESEARCH GRANT

Leon M. Lederman

April 30, 1992

A. Communication and Advancement of High Energy Physics

1. I continue my service as a member of the Board of Trustees of URA, devoted to the oversight of SSCL and Fermilab.
2. I continue as member of SEAB with special emphasis on the Future of National Labs panel whose final report will be out shortly.
3. I have visited numerous universities: Northern Iowa, University of Illinois in Chicago, Princeton, ELION, Jamaica (University of West Indies), Florida State, Florida A & M, Miami, (Ohio), Northern Michigan University, Ohio Northern University, Northern Iowa University, Kent State University, University of Wisconsin, Notre Dame, NYU, National Institute of Science and Technology, Lawrence University (Wisconsin), Rosary College (Chicago). In all of these visits, I talked about high energy physics as well as the general importance of science education.
4. Membership on the Boards of Science Services (sponsors of the Westinghouse Science Talent Search), the Museum of Science and Technology in Chicago, the Scientists Institute for Public Information and the Council of American Science Writers all add coherently to my ability to communicate and advance science in general and High Energy Physics in particular. This also goes for my activities as President and as Chairman of the Board of AAAS.

In the next year, I expect all of these activities to continue at about the pace of the 1991-1992 year.

B. Experimental High Energy Physics

Part I   B-Quarks

The field is enlivened by three approaches: the B-factory and the Cornell runner-up; the CDF/D0 hadron collider and the fixed target spectrometers which have just completed a long run. At this writing, HEPAP et al. recommendations notwithstanding, it is not clear where the "good stuff" will come from -- the CDF (UA1) data represent some tens of thousands of $B \rightarrow e^+x$ events trailing down to (CDF) a hundred or so fully reconstructed B exclusive decays. The next run may raise these by ~ 100 both because of the added vertex detector and general improvements in machine luminosity and detector efficiency. The fixed target experiments are deep in analysis and the E789 status is given below. Two or so years from now one can expect improvements of factor of 10 solidly and perhaps factor of 100 over the probable ~ 10 's of events now on top. CDF in the 1997 era may have 10^{10} $B\bar{B}$ pairs but the clarity of these events must await current analysis. Cornell will play away with increasing luminosity and and e^+e^- factory probably won't appear until 1999 ± 2 .

What follows is the E789 progress report extracted from a longer report. The University of Chicago participants in this research are Leon M. Lederman and Martin Schubb.

E789-RELATED WORK

The E789 collaboration includes physicists from eight institutions: Abilene Christian University, Berkeley, Chicago, Fermilab, Los Alamos, NIU, South Carolina, and the Taiwan Users Group. A sensitive search for rare low-multiplicity decay modes of particles containing c or b quarks, E789 had a test run with partial instrumentation in the spring and summer of 1990, and its first run with all equipment installed has just ended. The apparatus is shown in Figure 5.

The start of E789's data-taking in the 1991 run was delayed relative to that of the other fixed-target experiments due to concerns about radiation safety in our counting rooms. After an extensive series of measurements and considerable shielding work, we received approval on August 24 to run our spectrometer at the charm setting. In order to approve us for the (higher-intensity) beauty running, the Fermilab Research Division first installed a temporary trailer in the parking area to the east of Meson Laboratory to house experiment personnel, so that we no longer needed continuous occupancy of the electronics rooms inside Meson Lab. Beauty running began October 7. The run was divided roughly equally between charm and beauty running (two months each). A total of $\approx 1.5 \times 10^9$ events, collected over a total of $\approx 8 \times 10^4$ beam spills, were recorded on ≈ 1300 8mm tapes.

Charm Running

The charm running was crucial for tuning up our newly installed stereo-view silicon-strip detectors and our vertex trigger processor. Figure 6 displays the hits on all 16 silicon planes for a typical event. The silicon tracks associated with the tracks found in the downstream spectrometer are also indicated. Using the information from the silicon detectors to require a decay vertex downstream of our target, we see a clear $D^0 \rightarrow K\pi$ peak (Figure 7) in a preliminary analysis of less than 1% of our charm data. The on-line vertex processor, which makes a looser vertex cut than that used off-line, worked successfully and reduced the background trigger rates by a factor of ≈ 10 . The data acquisition upgrade, which replaced the E605 4-megabyte buffer memory with a 64-megabyte buffer, allowed a ten-fold increase in our data acquisition rate. These two factors together permitted data-taking at interaction rates up to 4 MHz, yielding up to $\approx 10^3$ reconstructible $D^0 \rightarrow K\pi$ events per day. The ring-imaging Cherenkov (RICH) counter worked well during a significant fraction of the run. Figure 8 shows Cherenkov rings observed in a preliminary analysis of RICH data.

Our charm measurements are expected to provide some 45,000 D^0 or $\bar{D}^0 \rightarrow K\pi$ events on tape, of which $\approx 7,000$ should survive vertex cuts in the gold-target sample and ≈ 1500 in the beryllium sample, permitting a useful measurement of the D cross section and its nuclear dependence in the small-Feynman- x region at 800-GeV beam energy. In addition, the RICH-identified data should yield a limit on the $D^0/\bar{D}^0$ production asymmetry. (It is possible that a reliable $D^0/\bar{D}^0$ identification can also be made kinematically; if so, the statistics of the entire charm sample will be available to support the results from the RICH analysis.) The gold-target sample should also contain some 400 $D^0 \rightarrow K^+K^-$ events and, by virtue of an acceptance enhancement for $\pi^+\pi^-$, 500 $D^0 \rightarrow \pi^+\pi^-$ events, yielding interesting measurements of the ratios of the three branching ratios; since in E789

we can distinguish $\pi\pi$, KK , and $K\pi$ decays kinematically, our measurement should have reduced systematic errors (as well as superior statistics) compared to that of the E691 collaboration⁷. We can also improve on current knowledge of dileptonic D^0 decays. The 90%-c.l. upper limits for $D^0 \rightarrow$ dileptons should be $\approx 0.5 \times 10^{-5}$ each for e^+e^- , $\mu^+\mu^-$, and $e\mu$.

Beauty Running

Beauty data were collected at a spectrometer setting (1500A in the "SM12" analyzing magnet) which simultaneously optimized sensitivity for $B \rightarrow J/\psi$ and for $B^0 \rightarrow$ dihadrons. We ran at interaction rates up to ≈ 50 MHz, limited by radiation levels in our trailer. The beauty data correspond to a total of 2.5×10^{13} interactions. Using standard assumptions for the size, shape, and A -dependence of the $b\bar{b}$ cross section, this should provide some 100 reconstructible $B \rightarrow J/\psi$ events; over a plausible range of cross-section assumptions, the number of events could well be as small as 40 or as large as 250. Figure 9 shows the dimuon mass spectrum from a preliminary analysis of $\sim 3\%$ of the beauty data sample. The J/ψ is evident at about the expected level, confirming our estimated sensitivity.

Assuming no dihadronic decays are observed after all cuts, the 90%-c.l. upper limit for each dihadronic B^0 decay mode is estimated at 1.0×10^{-4} (and similar limits for $B \rightarrow e^+e^-$, $\mu^+\mu^-$, and $e\mu$).

Data Analysis

We are presently doing preliminary analysis on both the charm and beauty running and are refining our understanding of detector alignments and the magnetic fields of our analyzing magnets. We expect to begin full-scale production analysis soon on both data sets, analyzing the charm data largely at Berkeley and Los Alamos and the beauty data at Fermilab.

Fig 5

3

E789 SCHEMATIC (PLAN VIEW)

Plan view of E789 spectrometer

EVT	14									
ZVTX, B, YSI:	-0.050	-5.139	-221	-232	-402	-420	0.997	0.956		
ZVTX, B, YSI:	-0.050	-2.914	81	133	133	0	0.954	0.805		

9 by

~4 tapes of data Fig 7

$D \rightarrow K\pi$

2. 21

fig 9

Part II   Secondary Vertex Trigger

We report on several fronts.

(A) A senior thesis of David Chen carried out at The University of Chicago looked in great detail at silicon strips and a fast, on-line read-out that fires on tracks emitted away from a target.¹ The cover, abstract, introduction and conclusions of David Chen's senior thesis are appended.

(B) The optical trigger of Charpak, Lederman and Giomataris is under study at CERN. Much analysis has been done and an experimental test in a beam is scheduled for early summer.

(C) A test of the optical trigger was carried out by Dan Kaplan and his students behind E789 at FNAL. A report is appended.

---

¹ A Trigger for Beauty, G. Charpak, Y. Giomataris and L. Lederman, Nuclear Instruments and Methods, 1991.

Part II 1. David Chen's Thesis

**A Speculative Trigger for the Silicon Vertex Detector
in a Fixed-Target Experiment**

A Senior Thesis presented by

Dave Chen

and sponsored by

Professor Leon M. Lederman

for the review of

**The Physics Department of
The University of Chicago**

May 1991

ABSTRACT

The experiment E789 at Fermilab proposes to measure two-body, two-prong, fixed-target decays of neutral b-quark hadrons using the earlier E605/E772 spectrometer. The spectrometer has been upgraded with an array of 16 planes of silicon microstrip detectors (SMD) placed closely behind the target, whose primary function is to pinpoint the two-body decays of a particular B state by the average distance that it decays downstream due to its characteristic lifetime. My thesis will not only examine the present configuration and vertexing algorithms of the SMD, but will undertake to design a faster, zero-level trigger for an immediate vertex discrimination. Such a trigger provides a possible solution to the high background typical of fixed-target experiments and may enhance the rejection efficiency to allow more time for the downstream detectors to gather data on the mass, lifetime, and production dynamics of the daughter particles. Problems encountered during the design of the trigger will also be discussed.

I. Introduction

The main objective of E789 is to study the dihadron decays of neutral b-quark hadrons in a fixed-target proton machine, in which a large number of events (an estimated 10^5 BB_'s per hour) can be produced. Due to the high background, however, it is imperative to implement a fast ($< \mu\text{s}$) zero-, or first-, level trigger, in order not to jam the data acquisition system. Such a front-end trigger, along with the higher-level triggers coming from the downstream detectors in the multi-step trigger scheme, will be able to reduce the final trigger rate to less than 10^4 per spill. In the case of the B's, the zero-level trigger is best provided using the signatures from a vertex detector placed closely behind the target. Since the B's have relatively high- P_T charged daughter particles and a relatively long lifetime, it will generate a distinctive secondary vertex downstream of the target.

In this paper I will start out with a brief discussion on the objective of experiment E789 as a whole and the major components of the spectrometer. A detailed account of the combined trigger system from all the various parts of the spectrometer is beyond the scope of this thesis. Instead, I will focus on the description of the newly installed silicon microstrip detector (or SMD) on which my speculative trigger is to be implemented. Before discussing my own trigger, however, I will first discuss briefly the trigger algorithm that is currently underway at the experiment, in order to serve as a model for comparison. Following will be a critical analysis of the Monte Carlo method I used in my simulation and the results that are generated. This includes a more in-depth discussion of the physical assumptions that have been made and an assessment of each of their validity. Problems such as multiple scattering among the silicon planes will also be addressed.

II. Physics Motivations & Objectives of E789

Presently E789 is proposing to study the dihadron decays of a neutral B state, mainly the decay of a neutral B_d or B_s into pairs of charged pions, kaons, or both, or a pair of proton/antiproton. B_d and B_s are bound states of the b quark and the u_- and s_- respectively, and they carry all the

The following plot gives the BAcceptance over the z position of the vertex and includes error bars due to statistical error:

In order to simulate tracks from the target, I need the Pt and Y (rapidity) distribution of the pions generated from the target.

`PtPtDist[Pt_] := Pt*Exp[-7.1*Pt+0.95*Pt^2]`

Detector is blind to target-originated particles and is fully efficient at $d \approx 10$ cm

problem of accidental filling, runs with higher statistics are clearly necessary.

IX. Conclusion

The addition of a zero-level trigger after the current pre-trigger of E789 could be useful in reducing the enormous ^{trigger} background that is typically produced in a fixed-target experiment. My trigger was able to achieve a per-track rejection efficiency of 99% for tracks coming from the point target. At the present interaction rate, this translates to an per-event rejection of close to 90%. However, in rejecting the huge number of target tracks we now end up with a 16% acceptance (for the configuration with 3 25-micron-pitch planes). In hope observing any of the rare B events, the experiment may have to run at a much higher interaction rate, which would in turn degrade the rejection efficiency. A decision over the optimum running condition requires a look at both the rejection of the target tracks and the acceptance of the B decay tracks.

Our test on the number of accidental filling, however unreliable, indicate to us that the acceptance of B's will further be undermined by approximately $3/8 = 38\%$ (8 is a number gotten from a similar but separate run) to only $1 - (16\% * 38\%) = 10\%$. By the same token, however, we expect the same effect of accidental filling of accepted target tracks to help us improve the rejection by $1 - (10\% * 38\%) = 96\%$. Furthermore, 85% of the B tracks that were rejected due to the trigger may also recuperate from the help of 10% of the non-rejected target tracks, adding another approximate 8.5% to the 10% B acceptance. A look at a compromise between rejection of target tracks and acceptance of B tracks should thus also take into account the effect of accidental filling.

In the process of deriving the per-track rejection efficiency to the event rejection efficiency, recall that we assumed a total number of 20 to 30 hits per arm; these numbers may have been overestimated. The number of tracks that actually fall on the silicon planes depend strongly on the running condition. However, to see how an overestimation may have been made, we can compare our number to an actual measurement of 20 p+p- per interaction, made by an experiment by CERN at a comparable energy. Since our experiment is expected to run at a rate of

10 interactions/bucket, and I get in one of my TargRejEff runs 300 tracks on both planes among 2000 thrown, I should expect about $0.5 * (300/2000) * 20 / \text{interaction} * 10 \text{ interactions} = 15$. This would raise the rejection up to 98% !

On the other hand, there are also numerous other problems with the trigger that need to be considered. The main problems that were discussed in this paper are the large multiple scattering exhibited by soft target pions, and the possibility of faked tracks due to the trigger's reliance on the simple geometry of the silicon planes. The roads defined by the scattering pions may require further optimization for better acceptance of the B tracks. This may call for the use of a Monte Carlo method for the simulation of the pions' momenta. Furthermore, if compatibility with the experiment's current setup could be ignored, then perhaps smaller number of planes is better for triggering purposes alone.

Many technological difficulties have been overlooked in the design of this trigger, which will eventually need to be addressed. One of the biggest problems involve the large amount of noise that may be produced in such a cramped, and high-interaction environment as the SMD. Planes of 99% efficiency are absolutely essential for the working of the trigger. The electronics design of the trigger is glossed over by this paper and may need further refinement before the trigger could be implemented.

From: FNAL::KAPLAN 20-APR-1992 14:46:30.41
 To: MHS
 CC:
 Subj: Preliminary statement of results

E789 TESTS
 OF OPTICAL TRIGGER

From: FNAL::KAPLAN 29-JAN-1992 21:29:46.65
 To: JNET%ioa@cernvm
 CC: UCHEP::LEDERMAN,DFANDERSON,CHUCKB,SWALK,APOLINSKI,LUEBKE,KAPLAN
 Subj: Preliminary note on optical-B-trigger test

Yanis,

This is a quick note to describe the preliminary result of our test. We seem to observe a surprisingly slow turn-on of phototube pulse-height with incident track angle. We tested a MgF2 crystal of dimensions 2.5 x 3.2 x 0.2 cm**3, attached to a Hamamatsu R1828-01 phototube having a glass window (to cut off the light at 300 nm and stay in the region where $n < \sqrt{2}$). Muons were tracked by a telescope consisting of eight single-wire drift tubes in three views at 60-degree stereo, spread out over about 0.8 m, with four tubes before and four after the crystal. The fiducial area was determined by two 2.5 x 2.5 cm**2 trigger scintillation counters located about 0.8m apart, at either end of the drift tube telescope. The trigger also required the muons to penetrate 4.6 m of shielding concrete, thus muon momenta were $> \sim 2$ GeV. After alignment and calibration, the drift tubes give r.m.s. resolution of about 100 microns, implying sub-milliradian angular resolution. We recorded over 100,000 events, covering about 50 mrad in incident angle, in three overlapping settings. The main result is in the form of mean ADC pulse-height from the crystal as a function of vertical angle (θ_y), since the crystal was oriented vertically, hanging down from the phototube. We see a slow, roughly linear increase of pulse height with angle over a range of about 20 mrad. We have tried imposing cuts on horizontal angle and distance from phototube, but the qualitative behavior is unchanged. While it is premature to draw conclusions, if these results hold up on further analysis, it seems to me that the proposed beauty trigger may not be feasible.

} Fig 1-6

We have not ruled out that the crystal may be scintillating or fluorescing, and these mechanisms could be producing light whose intensity may depend on angle in subtle ways. For example, Dave Anderson has suggested that if fluorescence is being stimulated by Cherenkov photons of wavelength about 150 nm, its light may be decreasing at the same time as the light from totally- reflected Cherenkov photons is increasing, since the path length in the crystal for short-wavelength Cherenkov photons would decrease as the muon angle steepens (i.e. as the muon points more directly at the phototube) - this because the (wavelength-dependent) index traps all 150-nm Cherenkov photons at normal incidence, but not at steeper angles.

We intend to refine the data analysis, improve the drift-tube calibrations and alignment constants, and make various measurements on the phototube and the crystal to establish number of photoelectrons per ADC count and fluorescence and degree of scintillation of the crystal. The crystal was purchased from Optovac, who claim that it does fluoresce slightly, whereas a more expensive grade (?) of MgF2 does not. We may therefore want to order a "non-fluorescing" MgF2 crystal and make further tests, if a copious source of muons can be found now that the fixed-target run is over...

Particle Detector Group) and my students Mark Apolinski and Bill Luebke. Apolinski is analyzing the data and will write up this work as his Master's thesis, and we expect to submit a paper to NIM. We are also preparing a paper describing the drift-tube telescope and its performance. We will keep you informed about any new results. (I'm sending this also to the above-named people and to Leon Lederman, Dave Anderson, and Chuck Brown who are all interested.) If you have any suggestions or ideas, please let me know. I would like to come to CERN at some point to discuss this with you, but it may be premature at present - what do you think?

Dan Kaplan

Hi, Leon,

Here are the plots Dan made up, plus a couple of descriptions of the apparatus and results. A good diagram of the apparatus does not yet exist, but Dan tells me there are Fermilab photos; I could have an 8-by-10 made up if you like.

I've numbered the plots in pencil in the lower right-hand corner. The coordinate system is as follows: the crystal is in the xy plane, with the phototube sticking straight up at positive y. The dimensions of the crystal are 3.2 cm in y, 2.5 cm in x, and 2 mm in z.

1. Phototube pulse-height versus theta-y. In this plot and plots 2-6, a cut has been made on the theta-x so that variations in theta-x don't smear out the turn-on with angle. The zero of theta-y has some arbitrary offset, so zero on the plot doesn't necessarily correspond to a track that's perpendicular to the largest face of the crystal. The slow rise in pulse-height is attributed to dispersion in the crystal, which makes both the Cerenkov angle and the trapping angle depend on wavelength, so that the Cerenkov photons that are emitted aren't monochromatic and also don't get trapped with equal efficiency. At negative theta-y, there's a non-zero pulse-height, which is presumably due to delta rays.
- 2-6. Same as 1, except that each plot shows only tracks that passed through a particular strip in y of the crystal; in plot 2, the tracks selected pass furthest from the phototube, and in plot 6 they pass closest. Ordinarily one would expect the turn-on to be sharpest furthest from the phototube, where the number of internal bounces is highest, but this doesn't seem to be the case, and again, this is attributed to dispersion.
7. Phototube pulse height versus theta-x, where theta-x, like theta-y, has some unknown offset. The asymmetry in this plot is not understood.

I know the above isn't the clearest description in the world, so if you have questions, let me know.

Martin

Fig 1

Fig 6

C. Science Education

Aside from some consultation work at Fermilab, with the Education group, and as vice-chairman of the Board of the Illinois Math Science Academy, my major effort has been with TAMS: the Teachers' Academy for Math and Science. This new institution has been supported by DOE and by NSF and has as its goal the retraining of 17,000 teachers in the City of Chicago. I append an overview and status report. By the end of the fiscal year, we will have had almost 700 teachers through our intense training program and over 4000 teachers participating in the workshops. Assessment and evaluation of the work uses a variety of early information techniques. Extensive records of teachers and principal commentary, hours of science and math being taught and truancy rates are among these indicators. It's a very ambitious program.

1) Fermilab: The education group is an outgrowth of "Friends of Fermilab" and has extensive experience in the kind of outreach pioneered at Fermilab and now mandated for all Labs. Fermilab's program is especially wide ranging, touching third graders as well as high school teachers. Recently the FNAL group has collaborated with TAMS in summer programs in the city of Chicago. These cooperative programs and Fermilab's long experience are increasingly valuable to the TAMS effort. We see this is a continuing process.

2) IMSA: This school was really started at Fermilab as a unique three year residence public school for gifted math and science high school students. It opened in 1986 and all three of its graduating classes have achieved the highest ACT scores in the nation! Fermilab continues to provide mentors to the students and act as a resource for the IMSA faculty.

3. Teachers Academy for Mathematics and Science

10 West 35th Street, Chicago, Illinois 60616 On the campus of Illinois Institute of Technology. Tel 312 808 0100 Fax 312 808 0103

I. Background

In a major urban setting and in a wide-reaching format, the Teachers Academy addresses the well-documented concerns with national math and science education:

- ♦ inadequate teacher preparation in science and mathematics;
- ♦ the lack of adequate instructional materials, particularly those for hands-on, quantitative mathematics and science activities;
- ♦ the lack of coordination of mathematics and science curricula;
- ♦ limited emphasis on problem-solving and real world applications;
- ♦ poor general understanding of the critical thinking skills that provide the basis for modern technology.

These problems are particularly severe in large urban areas and in the Chicago Public Schools. If these problems can be solved in Chicago, they can be solved anywhere. Chicago is a microcosm of the nation, and its public school system is typical of the nation's large, dysfunctional urban school systems. Therefore, the Academy will focus initial intensive staff development efforts on the Chicago Public Schools and eventually serve the entire metropolitan Chicago area.

II. The Teachers Academy

The Teachers Academy was born of Chicago's reform movement which has given parents and community members a greater stake in their schools. Developed under the leadership of Nobel-prize-winning physicist, Leon Lederman, the Teachers Academy's goal is a restructuring of school learning environments through the vehicle of mathematics and science. With active support from area universities, museums, national laboratories and corporate research centers, the Academy will initially link Chicago's extensive educational and scientific resources with its inner city school system and, subsequently, with all metropolitan area schools. It is an experiment of unprecedented boldness and scale that can provide a means, both for the Chicago area and as a national model for other urban areas.

The Teachers Academy was founded in 1990 as a project of the Chicago Education Federation, which is governed by a Council of Presidents composed of the Presidents or Chancellors of 13 leading Chicago-area colleges and universities. In addition to appointing the Teachers Academy Board of Directors, the Council of Presidents has taken an active role in coordinating the objectives of the Teachers Academy and the various universities, in involving the Teachers Academy in the development and operation of pre-service programs, and in developing and coordinating collaborative projects. Meaningful activities have already occurred, including the donation of 18 months

of rent-free space to house the Academy at the Illinois Institute of Technology Research Institute, and the loan of curricular experts from the University of Illinois at Chicago and the University of Chicago to serve as initial Teachers Academy faculty and trainers and provide input into design and development of program content. The University of Chicago and the two research laboratories (Argonne and Fermilab) also donated crucially-needed staff and financial support during the Academy's start-up phase. Further, there are plans in progress for additional loaned faculty, dual appointments, the use of undergraduate volunteers in Academy programs, and the collaborative development of college-credit follow-up programs for Teachers Academy graduates. Funding for the Teachers Academy is being provided by the U.S. Department of Energy, the National Science Foundation, the U.S. Department of Education, the State of Illinois and private sources.

III. The Teachers Academy Mission

In partnership with school communities, the Teachers Academy's mission is to restructure the total learning environment of each school through the related vehicles of mathematics and science, to transform the instructional system, and to energize the faculty. To accomplish this, the Teachers Academy will:

1. Focus on teachers, kindergarten through twelfth grade, as the key to such improvements.
2. Further involve parents and community leaders in the educational process.
3. Enhance the capabilities of teachers to deliver state-of-the-art mathematics and science instruction to all students.
4. Marshall the vast resources of the City, its universities, museums, national laboratories, and private sector and link these resources to the needs of the schools.

IV. The Teachers Academy Programs

Intensive Staff Development Programs

The Teachers Academy Staff Development Program, now offered to elementary school teachers but soon to be expanded to include secondary school teachers, is designed to involve the entire school community in new approaches to academics, via:

- ♦ a needs assessment conducted at each school;
- ♦ a systematic staff development program which works with public school teachers to revitalize the current system from within;
- ♦ replacement teachers paid by the Teachers Academy who make it possible for teachers to confidently leave their classes so they may attend Academy Intensive classes for one semester.
- ♦ Intensive follow-up: to assist teachers who have completed their semester of classes at the Academy, an Academy Implementation Specialist is assigned to the school for an additional semester.

Teacher Resource Center

A Teacher Resource Center will provide a continuing opportunity for teachers to expand their mastery in areas related to math and science. Now in the advanced planning stages, the Resource Center is scheduled to open in mid 1992. This collegial setting will be open to all area teachers, from elementary, middle and high school, from

public, private and parochial schools. This center is designed to bring teachers in contact with effective science and mathematics teaching curricula and techniques, as well as with technology specialists and colleagues from other schools. A multi-media library, well-equipped workstations, classrooms, plus information, reception and teacher hospitality areas will all play an important role in making the Resource Center a "teacher-friendly" place.

Special Focus Workshops

From one-day to two weeks, special focus workshops will continue to introduce area teachers to innovative new curricular methods, and the individual resources of practicing mathematicians and scientists. Workshops are designed to increase use of hands-on multi-sensory recreational and imaginative programs in the classroom, and to encourage teachers who have suffered from math anxiety and science phobia.

V. Council of Presidents

Mr. Mirron Alexandroff	President Columbia College
Rev. Raymond C. Baumhart, S.J.	President Loyola University
Dr. Nelvia Brady	Chancellor City Colleges of Chicago
Dr. Ronald Champagne	President St. Xavier College
Dr. Lewis M. Collens	President Illinois Institute of Technology
Dr. Dolores E. Cross	President Chicago State University
Dr. Hanna H. Gray	President University of Chicago
Dr. Theodore L. Gross	President Roosevelt University
Dr. Orley R. Herron	President National-Louis University
Dr. Gordon H. Lamb, Chairman	President Northeastern Illinois University
Rev. John T. Richardson, C.M.	President DePaul University
Dr. James Stukel	Interim Chancellor University of Illinois at Chicago
Dr. Arnold Weber	President Northwestern University

Ex Officio

Mr. Ted D. Kimbrough	General Superintendent Chicago Public Schools

VI. The Teachers Academy Board of Trustees

Dr. Adrienne Y. Bailey
Deputy Superintendent
Chicago Public Schools

Hon. Arthur L. Berman
State Senator
State of Illinois

Mr. Bruce Berndt
President
Chicago Principals Association

Dr. Gordon Berry
President, Citizens Schools Committee
Physicist, Argonne Laboratory

Mr. Bernie Bradley
Teacher
Newberry Academy

Dr. Lewis M. Collens
President
Illinois Institute of Technology

Mr. James Compton
President & CEO
Chicago Urban League

Mr. Judd R. Cool
Vice President, Human Resources
Inland Steel Industries, Inc.

Mr. Mario DiPrizio
Local School Council Member
Engineer, Motorola Corporation

Mr. Leonard J. Dominguez
Deputy Mayor for Education
City of Chicago

Dr. Leonard Ellinger
Director of Summer Sessions
Chicago State University

Dr. Ahmed A. Fareed
Dean, College of Education
Northeastern University

Ms. Barbara Franke
Director
Golden Apple Science Program

Ms. Cellie French
Teacher
Collins High School

Dr. Henry Frisch
Physicist
Enrico Fermi Institute
University of Chicago

Mr. Christopher Galvin
Executive Vice President
Motorola Corporation

Ms. Iris Goldfein
Partner, Actuarial Benefits
Coopers & Lybrand, Chicago

Ms. Marilou Von Ferstel
Executive Vice President
General Manager, Public Affairs
Ogilvie & Mather

Dr. Paul Helme
Director
Chicago Academy of Sciences

Mr. William A. Hensley
Vice President
Corporation Communications
Illinois Bell

Sr. Peter Mary Hetling
Archdiocese of Chicago

Dr. James S. Kahn
President
Museum of Science and Industry

Ms. Edith King
Teacher
McDowell Elementary School

Hon. Bob Kustra
Lt. Governor
State of Illinois

Dr. Gordon H. Leab
President
Council of Presidents and
Northeastern Illinois University

Dr. Leon M. Loderman,
Co-chair, Board of Trustees
Fermi Institute
University of Chicago

Dr. Robert Leininger
Superintendent of Education
State of Illinois

Ms. Coretta McFarren
Coordinating Director, WESCO

Mr. Jeffrey C. Miller
Associate Dean/Administration
Northwestern University
Medical School

Mr. Sam Mitchell, President
Chicago Association of
Commerce & Industry

Ms. Lourdes Monteagudo
MacArthur Foundation
Board Member

Mr. Richard Morrow
Ret'd CEO
Amoco Oil Corporation

Earl L. Neal
Co-Chair, Board of Trustees
Principal, Earl L. Neal Assoc.

Dr. John Peoples
Director
Fermi National Laboratory

Mr. Donald S. Perkins
Co-Chair, Board of Trustees
Ret'd CEO, Jewel Companies

Mr. Joseph D. Reed
Past President
Leadership for Quality Education

Dr. William Sampson
President
Chicago United

Dr. Alan Schricsheim
Director/CEO
Argonne National Laboratory

Mr. Daniel Solis
Executive Director
United Neighborhood Assn.

Dr. Jon R. Thompson
Director, Teachers Academy
for Mathematics and Science

Mr. Alvin VandenBosch
Chairman, Illinois Advisory
Committee, Non-Public Schools

Ms. Jacqueline Vaughn
President
Chicago Teachers Union

Mr. Mark Wagner
Teacher
Kenwood Academy

**DATE
FILMED**

8/12/92