for Am (2)

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2793
UC-427

PPPL-2793

INVERSE PROBLEM FOR BREMSSTRAHLUNG RADIATION

BY

K.E. VOSS and N.J. FISCH

October 1991

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
703-487-4650

Inverse Problem for Bremsstrahlung Radiation

K. E. Voss and N. J. Fisch

Princeton Plasma Physics Lab

July 25, 1991

Abstract

For certain predominantly one-dimensional distribution functions, an analytic inversion has been found which yields the velocity distribution of superthermal electrons given their Bremsstrahlung radiation.

Paper

In general, it is not possible to write an analytic expression for the distribution function of electrons simply on the basis of observing their Bremsstrahlung radiation in a plasma. There is a set of circumstances, however, in which just such a data inversion is possible: Consider a homogeneous plasma of electrons and ions of charge Z_i . The Bremsstrahlung emission at energy $\hbar\omega$ is given by

$$R(\hbar\omega, \theta) \equiv \frac{dP}{d(\hbar\omega)d\theta} = \int_{v_{\min}}^{\infty} d^3\mathbf{v} f(\mathbf{v}) n_e n_i v \hbar\omega \frac{d\sigma}{d(\hbar\omega)d\theta}, \quad (1)$$

where $f(\mathbf{v})$ is the electron distribution function, and $d\sigma/d(\hbar\omega)d\theta$ is the Bremsstrahlung radiation cross-section. If the electrons are non-relativistic, yet are of large enough energy

for the Born approximation to be valid ($v \gg c \cdot 2\pi Z_i/137$), then the cross-section is given by Koch and Motz (1959) as:

$$\frac{d\sigma}{d(\hbar\omega)} = \frac{16Z_i^2 r_0^2}{137} \frac{1}{\hbar\omega} \frac{c^2}{v^2} \ln \left[\frac{(\sqrt{\frac{1}{2}mv^2} + \sqrt{\frac{1}{2}mv^2 - \hbar\omega})^2}{\hbar\omega} \right]. \quad (2)$$

In certain cases, the distribution of the most energetic electrons is primarily one dimensional, i.e. $f(\mathbf{v}) = f(v_{\parallel})$, where $v_{\parallel}$ might be the dimension along a strong imposed magnetic field. This is the case in tokamak plasmas for a slide-away discharge (Pieroni and Segre, 1975) or for discharges sustained by rf-driven currents (Fisch, 1987). Using Eq. (2) and $f(\mathbf{v}) = f(v_{\parallel})$ in Eq. (1), we have

$$R(s) = R_0 \int_s^\infty dx f(x) \frac{1}{x} \ln \left[\frac{x}{s} + \sqrt{\frac{x^2}{s^2} - 1} \right] = R_0 \int_s^\infty dx f(x) \frac{1}{x} \cosh^{-1} \left(\frac{x}{s} \right), \quad (3)$$

where energies are normalized such that: $x^2 \equiv \frac{1}{2}mv_{\parallel}^2$, $s^2 \equiv \hbar\omega$. Note that the lower bound takes into account the cutoff of electrons below the photon energy.

If we now take the derivative of Eq. (3), we get

$$\frac{dR}{ds} = R_0 \int_s^\infty dx f(x) \left[-\frac{1}{s} \frac{1}{(x^2 - s^2)^{1/2}} \right] = \frac{R_0}{s} \int_s^\infty dx \left[\frac{f(x)}{2x} \right] \frac{2x}{(x^2 - s^2)^{1/2}}, \quad (4)$$

which is the familiar form of an Abel invertible equation (Bronshtein and Semendyayev, 1985), so we can invert to yield:

$$f(x) = \frac{2x}{R_0\pi} \int_x^\infty \frac{d}{ds} \left(s \frac{dR}{ds} \right) \frac{ds}{(s^2 - x^2)^{1/2}}. \quad (5)$$

A similar inversion can be made for a distribution which is primarily in the perpendicular direction, e.g. a distribution of highly energetic electrons created by electron cyclotron resonant heating. In this case (where we label the energy as $y^2 \equiv \frac{1}{2}mv_{\perp}^2$), the resulting inversion is:

$$f(y) = \frac{2}{R_0\pi} \int_y^\infty \frac{d}{ds} \left(s \frac{dR}{ds} \right) \frac{ds}{(s^2 - y^2)^{1/2}}. \quad (6)$$

The restrictions on this successful inversion of Bremsstrahlung data are that of high energy (Born approximation), yet non-relativistic electrons, and a primarily one-dimensional distribution. All three of these conditions can be met by the electrons in an extended tail distribution of a typical tokamak discharge (1-10 keV), where the tail extends to about 5 thermal speeds. Thus it should be possible to apply experimentally this inversion technique, after first subtracting off the background emission from thermal electrons, to determine, for example, the very important features of rf-driven tokamak discharges.

Acknowledgements

The authors would like to thank George Vetoulis for his comments. This work was supported by the U.S. Department of Energy under contract DE-AC02-76-CHO-3073.

References

- [1] BEKEFI, G. (1966) *Radiation Processes in Plasmas*, John Wiley and Sons, Inc., New York.
- [2] BRONSHTEIN, I. N. AND SEMENDYAYEV, K. A. (1985) *Handbook of Mathematics*, Van Nostrand Reinhold Company, New York.
- [3] FISCH, N. J. (1987) *Reviews of Modern Physics* **59**, 175.
- [4] KOCH, H. W. and MOTZ, J. W. (1959) *Reviews of Modern Physics* **31**, 920.
- [5] PIERONI, L. and SEGRE, S. E. (1975) *Physical Review Letters* **34**, 928.

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Data and Planning Center, Nagoya Univ., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Prof. S. Tanaka, Kyoto Univ., JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Library, Kyoto Univ., JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	S. Mori, Technical Advisor, JAERI, JAPAN
Ecole Royale Militaire, Lab. de Phys. Plasmas, BELGIUM	O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	H. Jeong, Korea Advanced Energy Research Inst., KOREA
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	Prof. D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Instituto De Pesquisas Especiais-INPE, BRAZIL	Inst. of Plasma Physics, PEOPLE'S REPUBLIC OF CHINA
Documents Office, Atomic Energy of Canada Ltd., CANADA	Librarian, Inst. of Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Library, Tsinghua Univ., PEOPLE'S REPUBLIC OF CHINA
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Dr. O. Petrus, AL I CUZA Univ., ROMANIA
Prof. T.W. Johnston, INRS-Energie, CANADA	Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Dr. C.R. James., Univ. of Alberta, CANADA	C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. P. Lukac, Komenského Univerzita, CZECHOSLOVAKIA	Dr. L. Stenflo, Univ. of UMEA, SWEDEN
The Librarian, Culham Laboratory, ENGLAND	Library, Royal Inst. of Technology, SWEDEN
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
P. Mähönen, Univ. of Helsinki, FINLAND	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	M. Durgut, Vice Chairman, Middle East Tech. Univ., TURKEY
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Ms. C. Rinni, Univ. of Ioannina, GREECE	Dr. G.A. Eliseev, Kurchatov Inst., USSR
Dr. T. Mui, Academy Bibliographic Ser., HONG KONG	Librarian, The Ukr.SSR Academy of Sciences, USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. B. Das Gupta, Saha Inst. of Nuclear Physics, INDIA	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Librarian, International Center for Theo Physics, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA
Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN	
Prof. I. Kawakami, Atomic Energy Res.Inst., JAPAN	
Prof. K. Nishikawa, Hiroshima Univ., JAPAN	

END

**DATE
FILMED**

11 / 19 / 91