QUARTERLY REPORT # 6
March 30, 1991

Project No. DE-FG22-89PC89782

"SELENIUM IN OKLAHOMA GROUND WATER AND SOIL"

Submitted to

Donna J. Tamilia
U.S. Department of Energy
Pittsburgh Energy Technology Center
P.O. Box 10940
Pittsburgh, PA 15236-0940

by

Asmare Atalay and Dharam Vir Maggon

University of Oklahoma
Civil Engineering and Environmental Science
202 W. Boyd, Room 334
Norman, OK 73019

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DMR

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

INTRODUCTION

Selenium was discovered in 1817 by the Swedish chemist, Berzelius, in residue from Sulphuric acid production. It was named from the Greek word, Selene, meaning the moon, because of its resemblance to Tellurium, an element which had been discovered earlier and was named from the Latin word tellus, meaning the earth (1). Selenium is located between Sulfur and Tellurium in the periodic table. It is both beneficial and toxic to animals, plants and humans. The narrow gap between necessary and toxic concentration of Se makes it imperative to understand the processes controlling the distribution of this element in the environment. The processes responsible for its distribution include volcanic activity, the burning of fossil fuels, the weathering of rocks and soils, ground water transport, precipitation of minerals, adsorption, chemical or bacterial reduction and oxidation, and metabolic uptake and subsequent release of the element by plants and animals (2).

Although Se in high concentrations is toxic to animals, the element is an essential nutrient in low concentrations. Selenium deficiency became an agricultural problem because of changes in animal nutrition, which resulted in losses of livestock in the U.S. This prompted the Food and Drug Administration to approve the use of Se as Selenate or Selenite in concentrations upto 0.2 ppm as an additive in animal feeds. Most Se intake is normally associated with consumption of cereals and sea foods. The average human diet in the U.S. contains 1.8 mg of Se per month (3).

Selenium is widely distributed in nature but does not occur evenly in the lithosphere. Its compounds are toxic to animals feeding on highly seleniferous plants, causing them to develop "alkali disease". Symptoms of chronic toxicity in animals include lameness, hoof malformation, loss of hair and emaciation (4). Most evidences indicate that there is a great overall potential for Se deficiency than for toxicity at current levels of Se intake. The maximum contaminant level in drinking water is 10 micrograms per liter, a level that barely provides the minimum nutrient amount of

Selenium with a consumption of 2 liters per day (5).

The objectives of this study are :

- 1 To determine the concentrations of Se in Oklahoma ground water and soil samples.
- 2 To map the Geographical distribution of Se species in Oklahoma.
- 3 To relate groundwater depth, pH and geology with concentration of Se.

LITERATURE REVIEW

Discovery of Selenium

Selenium is one of the most widely distributed elements of the earth's crust and is rarely present in any materials in concentrations exceeding 500 ug/g. The presence or absence of Se in any soil is dependent on a series of geochemical and environmental factors. Figure 1 shows a geochemical cycle of Se. Selenium originating from igneous rocks may be released to sediments and finally reaching the soils. Plants may take up Se from the soil. Studies conducted on the occurrence of Se indicate that Se in soils comes mainly from geochemical processes and anthropogenic sources(14)

This diagram summarizes the behaviour of Se in the environment. It indicates that Se is involved in a variety of physical, chemical and biological processes. These processes include volcanic activity, combustion of fossil fuels, weathering of rocks and soils, soil leaching, groundwater transport, plant and animal uptake and releases, adsorption and desorption, chemical and biological redox reactions and mineral formation. The importance of a given process is largely determined by the particular speciation of Se. Therefore, the biogeochemistry of Se and environmental parameters, such as pH and redox conditions, strongly influence the concentration, mobility and distribution of Se in the environment(2).

Geochemical Processes

Much of the Se in the earth's crust is associated with sulfide minerals and it is here that Se is recovered during the electrolytic refining of sulfide minerals. Selenium is associated with sulfides of silver, copper, lead, mercury, nickel and other metals(14). During the early magmatic stage, sulfides of iron, cobalt and nickel tend to separate from magma and, at the same time, Se separates and gets associated with some of the sulfide

Figure 1 Possible cycle of Se under field conditions (Sharma, 1989)

minerals (14). Kolgonen observed that Se being a chalcophile element, does not become incorporated into silicates during the endogenic conditions and, therefore, most of it is found in sulfides. During exogenic processes, Se partly separates from sulfur (14).

During magma crystallization, volatile components including sulfur and Se are concentrated in the residual fluid phase, thus becoming separated from magma and often from sulfide ore bodies. However, the average sulfur and Se content in igneous rocks is considered to be acceptable values. Goldschmidt reported an average of 520 ppm sulfur and 0.09 ppm Se in igneous rocks, thereby giving a S : Se ratio of 1 : 6000 (14). Coleman and Delevaux reported the following maximum values for Se: Pyrite 5 %, marcasite 0.65 %, chalcocite 5 % (14). It must be noted, however, that there is a wide variation in the Se content of sulfide minerals with such factors as mineralogical composition, type of deposits and geological setting of ore body that play an important role in dictating the Se levels. The highest content of Se is found in Chalcopyrite - pentlandite - pyrrhotite deposits and quartz - pyrite - chalcopyrite formations. Goldschmidt suggested that the Se content is higher in high temperature sulfides than in low temperature ones (14).

Selenium is very mobile in conditions under which Granite rocks are crystallized and is depleted in silicic plutonic rocks and in pegmatites (14). The behavior of Se during weathering of minerals is fairly simple compared with most other elements. When sulfides weather in acid soils, Se is oxidized to the elemental form or to Selenite. It is adsorbed to colloids and migrates with difficulty. Under neutral to alkaline environments, such as, in calcareous soils, Se is oxidized to Selenate, in which form it easily dissolves (14).

Sources of Selenium in the Environment

Sulfides or native sulfur deposits often contain Se in significant amounts because of the chemical similarity between

Sulfur and Selenium. Thus sulfides of iron, mercury, silver, copper, lead and zinc have been found to contain Se, sometimes at levels over 20 % (6).

Selenium in Rocks

Selenium is not evenly distributed in geologic materials. Igneous rocks have been estimated to contain Se in the range of 0.03 to 0.8 ppm, with an average of 0.09 ppm (7). This value was accepted as the average crustal abundance of the element, but revision of the recent data showed an estimated abundance of 0.05 ppm (8). Selenium was found in volcanic gases, which suggested that volcanic activity played an important role in the development of seleniferous geologic beds. In addition to volcanism, weathering of igneous rocks and leaching by rain water were considered to be important sources of the element (9). In practice, Se tends to concentrate in sedimentary rather than in igneous rocks. Silicate rocks are formed from the magma, and volcanic eruptions result in losses of volatile selenium. Volatile Se is distributed over the earth's surface, eventually that adds to the Se content of the present day soils (10). Igneous rocks are generally deficient in Se because of volatilization of the sediment by volcanic activity (11). Rock weathering is the major natural source of Se in the environment. Weathering of the earth's crustal material results in an estimated 7.2×10^6 kg Se being mobilized annually and transported by rivers draining the land mass (12).

Selenium in Soils

The concentration of selenium in most soils lies within the range of 0.01 to 2.0 mg Se/kg. However, high concentrations of 1200 mg Se/kg as total and 38 mg Se/kg as water soluble selenate have been reported in seleniferous areas of the world (13). The concentrations of Se in soils varies with the origin, pH and composition of soils (14). Shales are the parent materials for about 40% of the earth's soil. Sandstones, limestones and igneous rocks are about equally abundant at weathering surfaces. Soils

derived from shales are most likely to be supplied with Selenium. The majority of seleniferous soils are weathered from shales. Soils derived from sandstones and limestones may have variable Se concentrations. Soils of igneous origin are most likely to be very low in Se. Selenium concentrations in soils range from less than 0.1 ppm in a Se deficient area of New Zealand to as high as 1200 ppm in an organic soil in Ireland (15). Availability of Se to plants is governed by the chemical form of the element in the soil and by the pH and Eh in the root-soil environment (15).

Surveys in most countries like Denmark, Iceland, Sweden, Finland etc. indicate that high levels of Se in crops could be attributed to the geology of the area. Various reports from most countries have indicated areas with low, medium and high Se concentrations (14). Allaway (15) has summarized the relationship among the concentrations of Se in rocks, soils and plants as follows:

1. Where rocks with a high content of Se decompose to form well-drained soils in subhumid areas (<20 in. of annual rainfall), selenides and other insoluble forms of Se will be converted to selenates and organic Se compounds. These compounds will be available to plants, and vegetation containing potentially toxic levels of Se will probably be produced in these soils.
2. Where rocks with a high content of Se weather to form soils in humid areas, slightly soluble complexes of ferric oxide or hydroxides and selenate ions will be formed. These soils will also be slightly-to-strongly acid, and plants produced on them will not contain toxic concentrations of Se, but they may contain sufficient Se to protect livestock consuming them from Se deficiency.
3. Where rocks with a high content of Se weather to form poorly drained soils, or where Se from higher lying areas

is deposited in poorly drained areas by alluvial action, and where the soils are alkaline, plants containing toxic levels of selenium are likely to be produced.

4. Where rocks with a low content of Se decompose to form soils under either humid or dry conditions, the plants produced are likely to contain insufficient Se to protect animals from Se deficiency (15).

Selenium in Plants

Plants are capable of taking up some Se if growing in soils where Se is available. The greatest interest in plant Se is due to the importance of this element in nutrition, particularly for domestic animals, since Se can act both as micro nutrient and a toxin (16). Plants capable of assimilating large quantities of Se have been divided into primary and secondary Se absorbers. Primary absorbers grow only on Se toxic soils and include about 24 species. Secondary absorbers are capable of accumulating moderately large quantities of Se when growing on soils containing high quantities of available Se (17).

In general, Se uptake by plants depends on climatic conditions, water regime of soil, oxidation-reduction potential and pH. About 8 Se containing organic compounds have been identified in plants. Amino acids, especially Cysteine, are likely to bind Se, which may have several metabolic implications. During growth and accumulation, Se is mined from the soil and rocks and is concentrated near the soil surface. Plants decompose at the soil surface and release organic and inorganic Se to the environment. Some of the Se upon decomposition in-situ or by animal digestive processes is present as organic compounds but most of it is in inorganic form as Selenate or Selenite.

Selenium in Coal mines

Selenium also occurs in coal deposits, probably due to the presence of Se enriched biomaterials, which decompose during coal

formation. The Se content of 138 samples of coal from U.S. deposits, in 22 states, ranged from 0.46 to 10.65 ppm and averaged 2.8 ppm which was over 50 times the crustal abundance of the element (4). Coal often contains 1 to 20 mg Se per kg. Upon combustion, coal releases large amount of Se into the ecosystem. An estimated 6×10^{11} and 28×10^{11} of coal (having about 11% of ash) are combusted annually in the USA and the world, respectively (18). Upon combustion, about 40% of the ash occurs as bottom ash or slag, 60% as fly ash, and less than 1% as aerosols, that escapes to the atmosphere (19). These proportions may differ for other coal boiler plants because of design and operating conditions (2). Most coal consumed is burned at power plants where Se and other elements entering the boiler are partitioned between the bottom ash stream (slag) and flue gas stream containing suspended fly ash and the vapors of volatile elements or compounds (13).

Human activities are significantly impacting the global Se cycle. Andren et al (2) concluded that 1.5 - 2.5 times as much Se is mobilized through coal burning as by natural weathering process. Most of the bottom ash from the precipitators are transported to landfills and solid residuals are flushed with water to ash ponds, where Se may be leached from the ash and enter the aquatic environment. Vapors are discharged as aerosols to the atmosphere, only to enter the terrestrial and aquatic environment by wet or dry deposition (13). According to Se mass balance study conducted at Allen Steel Plant in Memphis (2), of the 2.2 mg Se/kg coal entering the boiler, 0.3 % exited with the slag, 68% was with the precipitated fly ash particles and the remaining 32% was associated with the vapor phase (13). A survey of 86 samples of coal from USA indicate that these samples contain an average of 3.36 ppm of Se, which is over 34 times the crustal abundance of the element (20).

Anthropogenic Sources of Selenium

Almost all primary Se is produced from copper refinery slimes. The present production technology consists of processes designed primarily for effective recovery of precious metals. Selenium

recoveries have therefore received secondary importance, which is reflected in low recovery efficiencies (4). Se is recovered from slimes by volatilization during roasting or furnacing as the oxide which is scrubbed from exhaust gas or by leaching of roasted calcine or furnace slag. All processes use sulfur dioxide to precipitate Se metal from solutions of sodium selenite and selenious acid (4). The annual consumption of Se in the United States increased 50% from 1964 to 1973. The most significant use of Se has been use in the electronic industry, including use in rectifiers, xerographics copying machines and photoelectric cells. The chemical industry accounts for an estimated 1/8 of the Se consumed. The chief use of the element in agriculture is in prevention of Se deficiency in livestock and poultry(4).

Occurance of Selenium in the Atmosphere

Selenium is introduced to the atmosphere by several natural sources. It has been suggested that volcanoes may be major contributors of Se in the air (9). Volatile Se compounds are released by accumulator plants mainly as dimethyl selenide. There are no accurate estimates, however, of the quantities contributed to the air by each of the above sources. The estimated industrial emissions of Se seems to be a small quantity to cause a serious pollution problem in comparison with the amounts of other industrial pollutants emitted (4).

Occurance of Selenium in the Natural Waters

The occurrence of soluble Se compounds in some Se soils and other geological materials suggests that natural waters would be effective in leaching and transporting the element from drainage areas to flood plains and other large water bodies. While considering the recent available information, the overall speciation data shows SeO_3^{2-} to be the dominant form of Se, with Se (-2) and Se(0) having the lowest concentrations (22). Nevertheless, the data also indicated that every river is unique with respect to its speciation and concentration of Se, and broad generalizations

are therefore, inappropriate. Cutter (22) has obtained speciation data for largest river system of California, the San Joaquin and Sacramento rivers. The San Joaquin, which drains the southern portion of the agriculturally important central valley of California, has total Se concentration of an order of magnitude higher than those of the Sacramento draining the northern valley. In San Joaquin, selenate is the dominant form, while selenite and selenide or elemental Se are primary species in Sacramento (22).

Selenium in Ground Water and Drinking Water

The content and speciation of Se in ground water and drinking water is of the most direct concern to humans. Seleniums as a minor constituent in drinking water in a concentration range of less than 0.1 - 100 ug/m³/liter. Safe drinking water act of USEPA, has established an upper limit of 0.01 mg/liter which is rarely exceeded. However, water derived from some of the cretaceous zones may have high concentrations of both Se and total soluble salts (23). In a study by Donovan et al (1981), in central Montana, it was reported that water from wells drilled into any of the geologic formations, cretaceous Colorado group, contained as much as 1000 mg/liter Se (23). An important source of drinking water, ground water, has total Se concentration ranging from less than 0.6-342 nmoles/kg. From these a worldwide average of 30 nmoles/kg can be extrapolated. All the available data shows that Selenate is the major Se species in ground water. Humic substances in ground water interfere with ground water Se speciation (22).

Selenium concentrations in surface water averages 2 ug/m³/liter or less. Drainage from irrigation and soil leaching are principal sources of Se in surface water. High concentrations of Se also occur in subsurface drainage waters from irrigated soils in much of the San Joaquin valley. However, the occurrence of elevated Se levels in ground water within this region is not uniform. Some areas having relatively low concentrations of Se occur among the nearby surrounding areas with much higher Se levels (24).

Environmental Cycle of Selenium in Nature

The schematic diagram summarizing the environmental cycle of Se is shown in Figure 2. This element is 40th in abundance in the earth (15). Volcanic eruptions often result in losses of volatile Se from the silicate rocks formed from magma. The volatile Se is distributed over the earth's surface which eventually adds to the Se content of the present day soils. Soils derived from shales are most likely to be supplied with Se. Plants absorb and accumulate Se from soils. Plants decompose at the soil surface and release organic and inorganic Se to the environment. Some of the plants and plant products are consumed by the animals and human beings. These plants products may be transported elsewhere before they are consumed and waste fractions are returned to the soil, streams, lakes and oceans (15).

Uses of Selenium Compounds

Unique optical properties of this element are useful in the laboratory and industrial applications. The familiar photographic exposure meters incorporate a Se photocell and a galvanometer which measures the current generated upon exposure of the cell to the light. It is extensively used to decolorise the green tint in glass resulting from ferrous sillicate (15). Selenate and also a 30% solution of potassium ammonium selenosulfide (selocide) have been used in control of mites and red spider. With the recognition of the role of Se in animal health and the development of safe and effective means of prevention of treatment of Se-responsive diseases, the use of Se as a food additive for animals may soon become one of the most important uses of the element and its compounds (15). However, commercial uses of Se are listed in Table 1.

Selenium Chemistry in Aqueous Systems

Increased mobility of Se into the environment via anthropogenic activities, and the potential oxidation-reduction behaviour of the element has made it imperative to study the

aquatic chemistry of Se. For this purpose, Se species are divided into two different categories: dissolved Se (that passes through filters with 0.45 μm openings) and particulate Se (of particle size $> 0.45 \mu\text{m}$), typically suspended sediment and other suspended solids. Element and colloidal phase, not truly dissolved, but passing through the filter is deemed to consist of Selenium (-2,0). In dissolved state, Se may exist in three of its four oxidation states; Se (-2), Se (+4) and Se (+6).

Particulate Se may exist in the same oxidation states as dissolved Se and can be found in different phases of the particulate matter. In sediments, Se may be within the organic material, Iron and manganese oxides, carbonates or other mineral phases. The actual chemical forms of the Se may be adsorbed to or co-precipitated with these phases (primarily selenites, selenates). Selenide can be covalently bound in the organic portion of the sediment. In addition, Se may be found in anoxic sediments as insoluble metal selenides precipitates, an insoluble elemental solution or as ferroselite and selenium containing pyrite.

Selenium exists in four oxidation states in aqueous systems; -2, 0, +4, +6. The reduced state, -2, is represented by the selenide species and the elemental state, 0, exists in different allotropic forms, whereas the two higher oxidation states are represented by selenites, +4, and selenate, +6.

The aquatic chemistry of Se is complicated since it can exist in four different oxidation states and as a variety of inorganic and organic compounds (25). The concentration, speciation and association of Se in a given environment depends on the pH and redox conditions, the solubility of its salts, the biological interactions and reaction kinetics. Many reactions in aqueous media involve transfer of electrons and protons. Such reactions can be represented in an Eh-pH diagram. The activity of the electrons is related to the Eh in the same manner as pH is related to the activity of the protons in solutions. Redox potential is a very useful parameter in describing and predicting the behavior of many important elements that occur in more than one oxidation state (26).

Figure 3 and 4 show the stability (Eh-pH) diagram for the different species of selenium in aqueous systems. Several species of selenium exist in the aqueous environment, however, the most dominant ones are those that are involved in oxidation reduction reactions. The main species involved are:

- a) **Elemental Se, Se(0)** - It is present in colloidal form and is presumed to result from decomposition of seleniferous minerals by bacteria, fungi and algae. The Eh-pH diagram indicates that elemental Se is stable in reducing environment. All crystalline forms are insoluble in water. In a study by Sarathchandra (27) 1981, it is reported that elemental Se can be oxidized to selenites and trace amounts of selenates by certain micro-organisms.

It is clear from figure 4 that a large portion of the stability domain of elemental selenium covers that of water. Therefore, it is stable in water and aqueous solutions at all pH values free of oxidizing and reducing agents. The stability of elemental selenium is demonstrated by its occurrence in sandstone in dry alkaline environments (4). Selenium exists in three allotropic forms; gray rhombic, which is the most stable, red-monoclinic, and the black amorphous form. The red and black amorphous allotropes are the most likely forms of selenium in soils and the aquatic environment. The red allotrop is found when selenium is precipitated from aqueous solution. Elemental selenium burns in air to form selenium dioxide, which is a solid substance (mp 340 degrees centigrade). During the combustion of fossil fuels selenium dioxide is formed, but it is readily reduced to elemental selenium by sulfur dioxide which is also found during the combustion of these materials. Contamination of water by elemental selenium would probably pose a minimal toxicity hazard.

- b) **Selenide, Se (-2):** It may be found in the dissolved states as hydrogen selenide in anoxic waters or as

Figure 3 An Eh-pH diagram for Se-H₂O system (McNeal, 1989)

Figure 4

Relation of Oxidation-Reduction potentials of some Se compounds to pH (Samuel , 1981)

organic selenides, primarily seleno-amino acids bound in soluble peptides, in oxic and anoxic waters (22). Equilibrium constants and solubility products show that selenide must exist in reducing environment as hydrogen selenide. Selenide is a foul smelling, poisonous gas. It is a weak acid, and when dissolved in water, it is easily oxidizes to elemental Se (27). Selenium is reduced to hydrogen selenide or other selenides at relatively low redox potentials(Figure III). Therefore, these compounds are unstable in aqueous systems. Hydrogen selenide by itself is not expected to exist in the aquatic environment since the $\text{Se}/\text{H}_2\text{O}$ Se couple falls even below the H^+/H_2 . Aqueous solutions of H_2Se are actually unstable in air. due to its decomposition into elemental Se and water. Under moderately reducing conditions, heavy metals are precipitated as selenides, which have extremely low solubilities. The precipitation of Se as heavy metal selenides can be an important factor affecting the cycling of the element in soils and natural waters.

- c) Selenite (SeO_3^{2-}) - Which according to a study in fresh water by Turner and Dickson (1) exists as HSeO_3^- at pH 6 and 31% (HSeO_3^-) + 69% (Selenite) at pH 9. Other thermodynamic data show that Selenite is favored under mildly oxidizing conditions. Selenous acid is a weak acid and is protonated when pH values are acidic to neutral. Most Selenite salts are less soluble than the corresponding Selenate salts. Selenite is reduced to elemental Se by mildly reducing agents (27). Selenious acid species are predominant in solutions over most of the area of moderately oxidizing conditions encountered in oxygenated waters. Between pH 3.5 and 9.0 biselenite ion is predominant in water. At pH values below 7.0 selenites are rapidly reduced to elemental Se under mildly reducing conditions.

Table 1. Uses of some inorganic Se compounds (National Academy of Science, 1976).

Selenium Compound	Industrial Use
Aluminium selenide	in preparation of hydrogen selenide; semiconductor research.
Ammonium selenite	in manufacture of red glass; as reagent for alkaloids
Arsenic Hemiselenide	in manufacture of glass
Cadmium selenide	in photoconductors, semiconductor photoelectric cells; in phosphors
Calcium selenide	in electron emitters
Cupric selenate	in coloring copper alloys black.
Indium selenide	in semiconductor research
Potassium selenate	as reagent
Selenium hexafluoride	as gaseous electric insulator
Selenium dioxide	in the manufacture of selenium compounds; as reagent for alkaloids
Sodium selenate	as veterinary therapeutic agent

Most Selenite salts are less soluble than the corresponding selenates. The extremely low solubility of ferric selenite and basic ferric selenite, which is found in dilute solutions of ferric chloride and sodium selenite, are important to environmental cycling of se. Selenites also form stable adsorption complexes with ferric oxides of even lower solubility than the ferric selenites. Under certain conditions, selenite seems to be completely adsorbed in high amounts by ferric hydroxide and to a lesser extent, by aluminium hydroxide, while selenate is not.

The adsorption of the selenite was shown to follow a langmuirian isotherm, with the specific adsorption being higher in acidic solutions and decreasing with an increase of pH value(4). The low levels of Se in ocean waters have been attributed to adsorption of Selenite by the oxides of the metals, such as iron and manganese (4).

- d) **Selenate, Se (+4)** - The +4 oxidation state of Se is stable in well oxidized environments. Selenic acid is a strong acid and therefore, not protonated over pH range of natural waters (27). Dissolved Se is found as Selenate in fresh waters from pH 6-9. Selenate salts are very soluble and conversion of selenate to selenite is a very slow process.

Selenates occupy the upper area of the Eh-pH diagram and therefore, are stable under alkaline oxidizing conditions. Most selenate salts are similar in solubility to the sulfates of the same metals. Soluble selenates would be expected in alkaline soils or alkaline weathering rocks in dry areas. Selenate appears to be the most significant form of Se as far as environmental pollution is concerned because of its stability at alkaline pH values, its solubility and its ready availability to plants.

Figure 5 shows the Pe-pH diagram for Se in an aqueous system under standard conditions. The redox range is considered to be bounded by the H_2O/O_2 and H_2/H_2O couples (28).

Figure 5

Predominance area diagram for Se in an aqueous system under standard conditions (Lakin, 1973)

The boundary is represented by the line,

$$Pe = 14.75 - 1.00 \text{ pH} \quad (1)$$

This line corresponds to Pe/pH conditions for which

$$\log PO_2/H_2O_2 = 6 \quad (2)$$

It can be seen from the predominance area diagrams (Figure 5) that Selenate, hydrogen selenite, and selenite are each predominant for some Pe/pH range within the overall conditions set on the system. Selection of the predominant form in the system is also based on the thermodynamic data.

The standard free energies of formation of some selenium species are shown in Table 2, which were used to develop the redox relationship (4). To predict the behaviour of Se in aquatic environment, the system can be divided into four broad categories - bulk water, interstitial water, suspended solids and sediment solids. Various transformations occurring in the system can, thus be predicted by making use of the above categories.

As can be seen from the predominance area diagram (Figure 5), the boundary between the selenate ion, $HSeO_3^-$, and selenite ion falls below the boundary determined by H_2O/O_2 couple. However, the oxidation potential in the natural aqueous system does not reach the boundary defined by the couple. Sato (28) proposed that the maximum oxidation potential in most aqueous systems may fall on or below the line defined by H_2O/O_2 couple which is given by

$$\log PO_2/H_2O = 6 \quad (3)$$

If this boundary is accepted as the upper limit on the oxidation potential in most aqueous systems, selenate may be slightly predominant under neutral to alkaline conditions, if the oxidation potential reaches the maximum potential predicted. Under acidic conditions, or under alkaline conditions with the oxidation

Table 2. Standard free energies of formation of some Selenium species(Susan Gibson, 1981)

Mineral name	ΔG_f (Kcal/mol)
Elemental Selenium	0.0
Hydrogen selenide	5.3
Hydrogen selenide ion	10.5
Selenide ion	30.9
Selenious acid	-101.87
Selenite ion	-88.40
Selenic acid	-105.42
Selenium dioxide	-41.5

potentials falling below the maximum predicted, selenite will be predominant. While predicting the relative amounts of selenates and selenites in natural waters, the kinetics for the reduction of the selenate ion to the selenite ion or to the elemental Se is also to be considered.

In a study conducted by Deveral and Millard (29) on shallow ground waters in San Joaquin valley, it is reported that Se is much more enriched in the coastal range alluvial soil zone and is strongly related to the selenity of the soil. The mobility and distribution of Se has been attributed to the geologic origin of the alluvial deposits.

Sampling and analysis for Se in San Joaquin valley in California has provided very important facts. Volatile methylated Se compounds (methyl selenides) and dimethyl selenonium compounds (DMSF) were documented in a study by Cooke and Bruland (25).

Selenium Entry to Aquatic Systems

Selenium is continuously cycled by natural processes. A general model explaining the cyclic process is shown in Figure 6. Se is transported from the core to the surface of the earth through igneous extrusions and volcanic gases. Soils may obtain Se from rocks, volcanic activities, industrial waste, irrigation water and fertilizers. Se and other trace elements are released through coal combustion and enter aquatic systems either by direct discharge or indirect input. Irrigation drainage effluent may be discharged into water bodies or into wet lands that are important wild life habitats (24).

In aquatic system Se is predominantly associated with the sediments, which act as both sink and reservoir, or with plants and animals living there, very small amounts are found dissolved in water. Aquatic bacteria can concentrate Se to high levels and provide a link into the food chain. The ultimate fate of Se and other trace elements is their return to sediments or other translocation systems. Other routes of escape for Se from an aquatic system include volatilization (24).

Figure 6 The cycling of Se in nature (Ohlerdorf, 1989)

Selenium Toxicity and Deficiency in Plants and Animals

Toxicity of Se to plants growing under natural conditions has not been reported, but symptoms of Se toxicity to crop plants have been described. Also, a decrease in yield of several crops was affected by the application of about 2 ppm Se into the soil (32). Increased Se levels in plants suppresses concentrations of N, P & S as well as several amino acids. So far, there is no evidence that Se is an essential element for the plants. Selenium deficient crops seem to occur intensely farmed humid areas on strongly leached soils.

Food constitutes the major route of human exposure to environmental Se in the general population. Estimates of adult exposure to environmental Se, generally, fall in the range of 20 - 300 ug/day. There is a very wide range of natural intake of dietary Se in different parts of the world. The Se compounds that may be encountered are Se dust and Selenium dioxide. The Grasslands of western Meced County, California, contain more than 30,000 hectares of seasonal and permanent wetlands in an area of largely native Grassland vegetation. This area comprises the largest tract of waterfowl habitat in the San Joaquin Valley and is also an important wintering area for waterfowl of pacific flyway. About two thirds of grasslands are located west of the San Joaquin River. Of this area, 20600 hectares are in the Grassland water District, 90% of which consists of Wetlands operated as private hunting clubs (24).

Subsurface agricultural drainage has been the sole source of water for Kesterson National Wildlife refuge. During 1983 to 85, aquatic birds nesting at Kesterson experienced poor reproductive success. The cause of this problem is considered to be exceptionally high concentrations of Se present in water and food chain organisms (24). Figures 7 & 8 show areas of USA where Se concentration in plants is less than 0.1 mg/kg and where Se deficiency disorders are likely to occur in livestock and poultry. Distribution of seleniferous vegetation in western USA and Canada is illustrated in figure 7. Each open dot represents the place of

Figure 7 Distribution of seleniferous vegetation in the western USA
and Canada (Rosenfeld & Beath, 1964)

Figure 8
Location and Classification of counties where plants
were collected for Se analysis (Kubota, 1967)

collection of a plant specimen containing 50-500 mg Se/Kg. Vertical lines, x's and solid dots indicate counties where more than one half of the crops sampled contained <0.1 , >0.1 , or where Se accumulator plants contained >50 mg Se/kg, respectively.

It has been found that Se concentrations are higher in sedimentary rocks where it is associated with the clay fraction, and thus, larger concentrations are found in shales and in limestones or sandstones. Even in shales, the concentrations are seldom >500 mg Se/kg. Values range from <1 to nearly 300 mg Se/kg in cretaceous black shales in Wyoming (33). Soils developed from the cretaceous shales of South Dakota, Montana, Wyoming, Nebraska, Kansas, Utah, Colorado and New Mexico tend to have high Se values. It is in these areas that natural Se toxicosis has occurred (34). Fig. 9 shows parts of North America which contain insufficient Se to meet the minimal animal nutritional needs.

■ $< 0.03 \text{ mg Se kg}^{-1}$

▣ $0.08 \text{ mg Se kg}^{-1}$

⊠ $0.05 \text{ mg Se kg}^{-1}$

□ $0.26 \text{ mg Se kg}^{-1}$

Figure 9

Concentrations of Se in forage crops in USA
(Kubota et al., 1967)

EXPERIMENTAL PROCEDURES

Sampling - Sampling of water and soil is a difficult and complex problem. The choice of sampling sites is of utmost importance in order to avoid trivial results. The objective of sampling is to collect a portion of material small enough in volume to be transported conveniently and handled in the laboratory while still accurately representing the material being sampled. This implies that the relative proportions or concentrations of all pertinent components will be the same in the samples as in the material being sampled.

Every effort was taken to get samples that are representative of the actual field conditions. The basic strategy included reaching the centre of every county in order to distribute the sampling sites throughout the map of Oklahoma. Samples were taken from sites which were being used for agriculture. use. Cost effectiveness and economic viability was given due consideration.

Water Sampling

Borosilicate brown glass containers were used to accomplish this task. Sampling was achieved only after the wells were pumped sufficiently to insure that the sample represents the groundwater source. In order to avoid any kind of contamination, the containers were cleaned with a metal free non-ionic detergent solution, rinsed with tap water, soaked in (1:1) Nitric acid and then rinsed with

metal free water. Chromic acid was used to remove organic deposits from the containers. All containers were then rinsed several times with deionised water to remove any residual chromium. Water to be sampled was allowed to run for a while before samples were taken. Every effort was done to reach as close to the source of the water as possible in order to avoid any contamination during transport through hose pipes etc. Immediately after sampling, water samples were kept in ice chest and then transferred to the refrigerator. Bottles were completely filled with water in order to avoid oxidation.

Soil Sampling

Soil sampling is not a easy task. Sampling site was cleaned of grass etc. Sampling was accomplished by digging approximately six inches of earth's surface in order to avoid any undesirable material. Soil samples were immediately stored in the ziplock bags.

Problems Encountered During Sampling

Some of the wells were denied access by the farmers and the property owners while some were declared contaminated due to oil spills. At times, chosen wells were found unoperational due to mechanical failures. On few occasions, on reaching the site, it was realized that the wells had caved in and hence it was logistically impossible to sample on that site.

Digestion of Groundwater Samples

To reduce interference by organic matter and to convert metals associated with particulates to a form (usually a free metal) that can be determined by atomic absorption spectroscopy, we have to use some digestion technique. We used the rigorous digestion method required to provide complete and consistent recovery compatible with the analytical methods and the metal being analysed.

Water samples were mixed by inverting the brown bottles several times. Fifty milliliter samples were pipeted and placed in digestion tubes. Samples were acidified by adding 5 ml. of conc. nitric acid. Two glass chips were put in each digestion tube in order to facilitate even boiling. Water was allowed to boil slowly until it evaporated to 15 to 20 ml. Then 5 ml conc. nitric acid and 10 ml conc. sulfuric acid to it. Samples were then allowed to evaporate until dense white fumes of SO_3 just appear. When the solution did not clear, 10 ml. conc nitric acid was added and evaporation continued.

When the solution was clear, white fume appears and the nitric acid boils off easily. Digested samples were then diluted using deionized water to the 100 ml mark on the tube. These samples were transferred into sample and kept bottles until analysis. Selenium was analyzed using International Laboratories (IL 451) Atomic Absorption equipped with a graphite furnace (30).

Digestion of Soil Samples

Soil samples from every county were air dried for one week by

spreading the moist soil on the laboratory bench. Samples were then ground using a Wiley Mill. The fine soil particles that passed through a 2 mm sieve were used for selenium analysis. Unused soil samples were stored in zip-lock plastic bags.

A 0.5 gm. soil was put in a digestion tube along with two boiling glass rods and 4 ml. of conc. HNO_3 . A small funnel was placed on the mouth of the digestion tube and the mixture was allowed to stand for 2 hours. The digestion tube containing the sample was then placed in the digestion system and gradually heated to 150 °C and maintained for 1 hour. It was then cooled and 5 ml. sulfuric acid added through the funnel. The temperature was increased to 235°C and held for 2 hours. The sample was then cooled and diluted to 100 ml. with deionised water. The resulting solution was filtered using whatman number 2 filter paper and analysed for its elemental analysis.

RESULTS AND DISCUSSION

Ground Water Samples from Oklahoma Counties

Table 3 shows the estimated amounts of total Se in groundwater samples from the 77 counties in Oklahoma.

Speciation of Se in Oklahoma Rural Groundwaters

Table 4 lists levels of Selenite in groundwaters from each of the 77 counties in Oklahoma. The results of this investigation indicate that Oklahoma's rural groundwaters contain primarily Selenite with no detectable Selenate present. This is in contrast to what is being reported for ground water samples by most other investigators. Selenate is being considered as the predominant species of Se in groundwaters due to its suspected high mobility and less adsorptive characteristics. It is apparent from this data that most of the counties exceed the allowable drinking water limit set for Se.

Figure 12 is a sampling distribution map for the state of Oklahoma. All dotted markings are approximate locations in each county where samples for groundwater and soils have been collected. The objective was to establish a well distributed sampling scheme.

Soil Samples

Table 5 shows the amount of total Se in soil samples in 77 counties in Oklahoma.

Table 3 Amounts of total Se in ground water.

COUNTY	Total Selenium (ppm)
ADAIR	0.02
ALFALFA	0.1
ATOKA	0.04
BEAVER	0.02
BECKHAM	0.02
BLAINE	0.02
BRYAN	0.02
CADDO	0.02
CANADIAN	0.02
CARTER	0.02
CHEROKEE	0.02
CHOCTAW	0.02
CIMARRON	0.04
CLEVELAND	0.02
COAL	0.02
COMANCHE	0.04
COTTON	0.02
CRAIG	0.02
CREEK	0.04
CUSTER	0.06
DELAWARE	0.04
DEWEY	0.06

DEWEY	0.06
ELLIS	0.02
GARFIELD	0.04
GARVIN	0.04
GRADY	0.02
GRANT	0.04
GREER	0.02
HARMON	0.12
HARPER	0.02
HASKELL	0.02
HUGHES	0.02
JACKSON	0.02
JEFFERSON	0.02
JOHNSON	0.02
KAY	0.16
KINGFISHER	0.02
KIOWA	0.10
LATIMER	0.02
LEFLORE	0.02
LINCOLN	0.04
LOGAN	0.02
LOVE	0.02
MAJOR	0.02
MARSHALL	0.02
MAYES	0.02
MACLAIN	0.02

MCCURTAIN	0.02
MCINTOSH	0.02
MURRAY	0.02
MUSKOGEE	0.02
NOBLE	0.02
NOWATA	0.04
OKFUSKEE	0.02
OKLAHOMA	0.02
OKMULGEE	0.04
OSAGE	0.02
OTTAWA	0.02
PAWNEE	0.02
PAYNE	0.02
PITTSBURG	0.02
PONTOTOC	0.02
POTTAWATIMIE	0.02
PUSHMATAHA	0.02
ROGER MILLS	0.02
ROGERS	0.04
SEMINOLE	0.02
SEQUOYAH	0.02
STEPHANS	0.02
TEXAS	0.04
TILLMAN	0.06
TULSA	0.06
WAGONER	0.02

WASHINGTON	0.02
WASHITA	0.1
WOODS	0.02
WOODWARD	0.02

Table 4. Speciation of Se in groundwater as estimated from selenite measurement.

COUNTY	SELENITE (ppm)
ADAIR	0.225
ALFALFA	0.212
ATOKA	2.360
BEAVER	0.177
BECKHAM	0.318
BLAINE	0.590
BRYAN	1.597
CADDO	0.296
CANADIAN	2.160
CARTER	0.333
CHEROKEE	0.016
CHOCTAW	0.064
CIMARRON	1.720
CLEVELAND	1.717
COAL	0.889
COMANCHE	1.520
COTTON	0.856
CRAIG	0.162
CREEK	0.967
CUSTER	0.074
DELAWARE	1.061

DEWEY	0.460
ELLIS	0.250
GARFIELD	1.880
GARVIN	0.270
GRADY	0.799
GRANT	4.033
GREER	4.190
HARMON	11.280
HARPER	0.096
HASKELL	5.310
HUGHES	0.203
JACKSON	1.860
JEFFERSON	0.884
JOHNSON	0.225
KAY	3.00
KINGFISHER	1.228
KIOWA	10.730
LATIMER	0.688
LEFLORE	0.645
LINCOLN	0.285
LOGAN	2.040
LOVE	0.492
MAJOR	0.447
MARSHALL	0.243
MAYES	0.120
MACLAIN	0.093

McCURTAIN	0.092
McINTOSH	1.092
MURRAY	3.404
MUSKOGEE	0.048
NOBLE	0.373
NOWATA	0.403
OKFUSKEE	0.580
OKLAHOMA	2.260
OKMULGEE	0.850
OSAGE	0.230
OTTAWA	0.206
PAWNEE	1.940
PAYNE	1.116
PITTSBURG	0.426
PONTOTOC	0.152
POTTAWATIMI	0.426
PUSHMATAHA	1.141
ROGER MILLS	0.161
ROGERS	1.485
SEMINOLE	9.490
SEQUOYAH	1.151
STEPHANS	6.210
TEXAS	0.728
TILLMAN	3.363
TULSA	1.034
WAGONER	0.946

WASHINGTON	0.796
WASHITA	1.460
WOODS	0.192
WOODWARD	0.298

pH Values for Different Water Samples

Table 5 lists the pH values recorded in the laboratory for 77 counties.

Table 5. pH values and depth to groundwater of water samples collected from different counties in Oklahoma.

County	pH	Depth to Groundwater (ft.)
ADAIR	6.67	70.0
ALFALFA	7.87	90.0
ATOKA	7.15	165.0
BEAVER	8.01	100.0
BECKHAM	7.52	140.0
BLAINE	7.90	200.0
BRYAN	7.53	150.0
CADDO	7.00	UNKNOWN
CANADIAN	7.57	UNKNOWN
CARTER	7.25	30.0
CHEROKEE	7.42	30.0
CHOCTAW	5.78	50.0
CIMARRON	8.06	280.0
CLEVELAND	8.07	100.0
COAL	5.85	75.0
COMANCHE	7.60	100.0
COTTON	7.53	30.0

CRAIG	7.18	117.0
CREEK	7.18	UNKNOWN
CUSTER	7.66	160.0
DELAWARE	7.26	120.0
DEWEY	7.55	28.0
ELLIS	7.75	150.0
GARFIELD	7.35	50.0
GARVIN	8.06	120.0
GRADY	7.79	>100.0
GRANT	8.17	100.0
GREER	7.52	654.0
HARMON	7.78	200.0
HARPER	8.03	80.0
HASKELL	7.45	25.0
HUGHES	7.66	90.0
JACKSON	7.47	54.0
JEFFERSON	7.28	16.0
JOHNSON	6.94	125.0
KAY	6.86	25.0
KINGFISHER	7.46	48.0
KIOWA	7.60	18.0
LATIMER	7.22	UNKNOWN
LEFLORE	6.95	135.0
LINCOLN	7.14	UNKNOWN
LOGAN	7.18	75.0
LOVE	6.56	110.0

MAJOR	7.65	58.0
MARSHALL	6.23	100.0
MAYES	7.26	100.0
MACLAIN	7.75	48.0
MCCURTAIN	6.02	80.0
MCINTOSH	6.75	UNKNOWN
MURRAY	7.42	100.00
MUSKOGEE	6.73	10.00
NOBLE	7.28	90.00
NOWATA	7.83	40.00
OKFUSKEE	7.65	UNKNOWN
OKLAHOMA	7.90	57.0
OKMULGEE	7.41	UNKNOWN
OSAGE	6.93	80.00
OTTAWA	7.40	UNKNOWN
PAWNEE	7.39	UNKNOWN
PAYNE	7.34	30.00
PITTSBURG	7.07	70.00
PONTOTOC	7.00	150.00
POTTAWATIMIE	7.30	90.00
PUSHMATAHA	7.20	UNKNOWN
ROGER MILLS	7.75	150.0
ROGERS	1.485	84.0
SEMINOLE	6.76	180.0
SEQUOYAH	7.19	UNKNOWN
STEPHANS	7.48	95.00

TEXAS	8.09	400.0
TILLMAN	7.62	30.00
TULSA	7.02	4.00
WAGONER	7.12	23.00
WASHINGTON	6.81	25.00
WASHITA	7.72	120.0
WOODS	7.73	150.0
WOODWARD	8.08	46.0

Distribution of Total Se in Groundwater

Table 5 also shows depth to the ground water at locations where the water samples were collected. Figure 13 shows the distribution of the total amount of Se in Oklahoma soils, as indicated on the map. Although no particular trend can be established by distributing the data on the map, it appears isolated pockets of land near the coal mine districts contain intermediate to high levels of selenium. Therefore, it may be conceivable that coal mining contributed to the release of selenium to the surrounding environment.

The main factors affecting Se concentrations are the degree of groundwater salinity and the geologic source of the alluvial soil material. In a study in the western San Joaquin valley, Ca, it was found that high groundwater Se concentrations generally are in the areas where saline soils and groundwaters are common (29). A better understanding of the origin, distribution and mobility of Se and its relation to the chemical composition of the groundwater in the agricultural areas is essential for effective water management.

This part of the study attempted to assess the relationship between Se concentration and various parameters that affect its distribution and mobility. The approach taken in this study was to evaluate the effect of geologic source material on Se concentration by analyzing the interrelationship between pH and other parameters.

The data obtained in the laboratory experiment and during the course of field investigation will be discussed in the light of following parameters:

Figure 13 Distribution of total Se in groundwater

1. Principal groundwater resources, recharge areas and geology of aquifers in Oklahoma.
2. Relation of concentration of Se with pH.
3. Interrelationship of total amount with depth of water.
4. Comparison of coal mining areas with the other areas in context of the total amount of Se present.
5. Copper deposits and Se concentrations in Oklahoma.

Principal Ground Water Resources in Oklahoma

Figure 15 shows the distribution of the principal bedrock aquifers in Oklahoma. The term aquifer refers to those rocks and sediments that are saturated with good-to-fair quality water and are sufficiently permeable generally to yield significant volumes of water from wells. The term recharge areas refers to those portions of the land surface where surface water (precipitation, surface runoff, streams, rivers and lakes) enters the subsurface and eventually migrates downward to the zone of saturation in the aquifers.

Oklahoma's bedrock aquifers are wide spread in distribution and differ in composition. Most of the aquifers are sands or sandstone, some are mainly limestones or dolomites. The main aquifers in the state are described below:

Ogallala Formation - Loosely cemented layers of fine to medium grained sand, silt, clay and gravel in western Oklahoma and pan handle. Thickness generally ranges from 100 to 700 ft. Weels

commonly yield 25 to 1500 gpm of water that is of good quality.
(generally less than 500 mg per liter dissolved solids)

Antlers sandstone - Loosely cemented fine grained sand and sandstone with some layers of shale and clay. It is situated in southeastern Oklahoma and includes underlying holly creek formations of eastern Maccurtain county. Antlers aquifer typically ranges from 200 to 700 ft. thick. Wells commonly yield 10 to 400 gpm.

Elk City Sandstone - Fine grained sandstone with minor amounts of silt and clay. Maximum thickness is about 185 ft. Wells commonly yield 25 to 300 gpm of good quality water. Its geographical location is in southwestern Oklahoma.

Rush Springs Sandstones and Morrow Formations - Fine grained sandstone with some layers of gypsum, shale and dolomite in western Oklahoma. Thickness ranges from 400 ft in the south to 200 ft in the north. Wells yield 25 to 300 gpm of water that is of good quality.

Cedar Hills sandstone - Fine grained sandstone interbedded with layers of siltstone and shale in north western Oklahoma. Thickness typically is about 180 ft.

Garber Sandstone and Wellington Formation - It is situated in central Oklahoma. Fine grained sandstone irregularly interbedded

with shale and siltstone. Thickness of the aquifer ranges from about 300 ft in the south to about 800 ft in the north. Wells commonly yield 25 to 400 gpm of water that is of good quality.

Oscar Group - Area designated Os-a consists of fine grained sandstone interbedded with shale and siltstone in central Oklahoma. Contains some conglomerate in the south. The thickness of the aquifer ranges from about 150 ft in the south to 300 ft in the north.

Arbuckle and Timbered Hills Group - Limestone and Dolomite interbedded with some sandstone and shale in Wichita mountains - Lawton region of southwestern Oklahoma. Maximum thickness is about 5000 to 6000 ft. Wells commonly yield 25 to 600 gpm of water that is of good to fair quality.

Distribution of total Se in soil samples

Table 6 shows total selenium in soil samples obtained from all 77 counties in Oklahoma. This data is shown in a distribution map in Figure 14. This distribution map shows that except for a few locations in the western part of the state, most of the soil contains low levels of selenium. Soils around the abandoned coal mine districts contain intermediate levels of selenium. However, these levels are higher than the average selenium level in soils.

Figure 14 Distribution of total Se in soil samples

ELLIS	2.20
GARFIELD	4.20
GARVIN	3.00
GRADY	2.00
GRANT	0.60
GREER	5.00
HARMON	2.00
HARPER	0.30
HASKELL	1.80
HUGHES	5.40
JACKSON	0.80
JEFFERSON	0.80
JOHNSON	0.30
KAY	0.30
KINGFISHER	0.30
KIOWA	1.60
LATIMER	0.80
LEFLORE	0.30
LINCOLN	1.40
LOGAN	0.30
LOVE	0.30
MAJOR	0.30
MARSHALL	0.30
MAYES	1.20
MACLAIN	3.00
MCCURTAIN	4.00

MCINTOSH	1.20
MURRAY	1.20
MUSKOGEE	0.30
NOBLE	1.00
NOWATA	2.80
OKFUSKEE	1.60
OKLAHOMA	2.80
OKMULGEE	0.30
OSAGE	4.80
OTTAWA	2.60
PAWNEE	0.30
PAYNE	0.30
PITTSBURG	0.30
PONTOTOC	3.60
POTTAWATIMIE	1.60
PUSHMATAHA	3.00
ROGER MILLS	3.40
ROGERS	0.40
SEMINOLE	4.20
SEQUOYAH	0.30
STEPHANS	4.00
TEXAS	2.20
TILLMAN	0.40
TULSA	6.00
WAGONER	4.20
WASHINGTON	0.30

WASHITA	2.00
WOODS	0.30
WOODWARD	4.00

Selenium in Geologic and Aquifer Materials

It has been reported that sedimentary rocks usually have Se concentrations much higher than the earth's average of 0.09 ppm (15). Lakin and Davidson gave ranges of Se contents of shale, sandstones and carbonate rocks of 0.24 to 277, 0.05 to 1.12 and 0.0 to 30 ppm, respectively. Igneous rocks are generally impoverished in Se as they are in Sulfur because of volatilization of these elements by volcanism (15).

Se concentrations in magmatic and sedimentary rocks are given in table 7. Analyzing the amounts of Se on figure 13, we find that Se enriched groundwater is found in southwestern and northeastern parts of Oklahoma. On the southwestern parts the principal water sources blaine and dog creek formations and rush spring sandstone formations. These aquifers are underlain by fine grained sandstone with some layers of gypsum, shale and dolomite which corresponds with the values in table 7.

It has already been reported that samples derived from south Dakota, Nebraska, Missouri, New Mexico, Colorado and Arkansas tend to have high Se levels. On the eastern side, Oklahoma is surrounded by Missouri and Arkansas while on the western side we have New Mexico and Colorado.

Moreover, the groundwater formations in northeastern parts of Oklahoma consists of limestone beds that tend to have high amounts of Se (table 7). Above all the north eastern parts of Oklahoma are surrounded by Missouri and Arkansas (established high Se content

Table 7 - Se concentrations in Magmatic and sedimentary rocks
(adapted from Kabata Pendias and Pendias, 1984)

Rock type	Se concentrations mg/kg
Magmatic rocks	
Ultramafic rocks	0.02 - 0.05
Maic rocks	0.01 - 0.05
Intermediate rocks	0.02 - 0.05
Acid rocks	0.01 - 0.05
Acid rocks - volcanic	0.02 - 0.05
Sedimentary Rocks	
Argillaceous sediments	0.4 - 0.6
Shales	0.6
Sandstone	0.05 - 0.08
Phosphorites	1 - 100
Limestones, Dolomites	0.03 - 0.10

Reeds spring formations that runs across Oklahoma and parts of Missouri too. Rock weathering has been established as the major source of environmental Se.

Correlation Between Selenium Content in Groundwater and pH

Figure 16 shows the graphical representation of pH with concentration of Se in the groundwater. Most natural waters tend to have low concentrations of Se. Notable exceptions can occur if waters are alkaline or if they leach and drain Seleniferous rocks and soil. One example of the pH effect is in the data of Scott and Voegli(2). When the pH of various Colorado rivers falls in the range of 6.1 to 6.9, the river water Se concentration is generally less than 0.01 mg per liter. However, when the pH of the water is between 7.8 and 8.2, the Se concentrations range from 0.01 to 0.4 mg per liter. One recent example of waters leaching and draining Seleniferous rocks and soils is in the San Joaquin valley (2).

Our data of pH in table 5 shows that most of the water samples are in alkaline conditions and the graph shows a rising trend which correlates to the data presented by previous researchers.

Relationships Between Total Selenium and Depth to Groundwater

The information provided in table 5 is based upon personal communication. Although we don't see any trend in the concentration of Se with respect to depth to water, yet we would anticipate that the deeper wells would show high levels of Se as compared to shallow wells. This analogy is based on the fact that the major

Figure 16 Principal bedrock aquifers in Oklahoma
(OSGS, 1989)

source of Se in water is rock weathering which indicates that water nearer to the rock will show high levels of Se concentration. In this case, more accurate information regarding depth of water wells will result from the hydrology and hydrogeological labs. More information regarding permeability, hydraulic conductivity, flow conditions etc. will also be needed to establish a relationship.

Comparison of Selenium Content Between Coal mining and Other Areas

Figure 17 shows the outline of the coal fields and areas of surface mining for coal in the state of Oklahoma. It is mainly situated in the northeastern parts of Oklahoma. If we again look at figure 13 we find that this area has higher levels of Se as compared to the other areas in Oklahoma. It, thus, agrees with the established fact that Se may leach from the coal deposits and enter the aquatic environment. It has also been reported that most of the bottom ash and the ash from the precipitators are transported to the landfills as solid residues or flushed with water to ashponds where Se is leached from ash and seeps down to groundwater(13)

The development of large scale coal mining operations in the western USA has raised concerns regarding the potential for Se contamination of soils, surface water and ground waters in and adjacent to these mining operations. It is expected that the Se content of the Oklahoma coal may be in the same range as those in the neighbouring states.

Figure 17 Coal mining areas in Oklahoma

Copper Deposits and Selenium Content

Se was discovered by Berzelius in 1870 in the residual slime from the preparation of sulfur dioxide from copper pyrites. Se, as usually recovered from these ores as Se bearing minerals, is not in a sufficient quantity to be considered a source of Se. Electrolytic refining of Copper and sludge from the sulfuric acid production accounts for much of the unrefined Se emissions (14). Figure 20 shows the map of northeastern Oklahoma, and adjoining parts of Colorado and New Mexico where copper minerals were mined more than 80 years ago.

Other Factors Affecting Accumulation of Selenium

Biological activity plays a strong role in the distribution of Se in the environment. The accumulation of Se in the plants and animals is highly variable and as noted before, can favourably or adversely affect their growth, survival and reproduction.

The speciation and the form of Se can change as a result of biological and physical processes. Selenate and Selenite can be reduced to red amorphous Se by numerous bacteria and yeasts. This process significantly limits the availability of Se to plants. Inorganic Se compounds can be converted to volatile organic compounds such as dimethyle selenide or demethyle diselenide by plants, microorganisms and animals(2).

Several inorganic processes control the distribution of Se. One is the adsorption of Se by particles. Selenite has a stronger affinity for particles, particularly Fe oxyhydrides than selenate,

and that adsorption is strongly influenced by pH. Sulfide minerals such as Galena and Pyrite tend to act as carriers for Se. Some sulfide minerals have been reported to contain upto 20% Se.

Industrial activities are also high contributors to Se level in the environment. This element is widely used in electronics because of its photosensitive and semiconducting properties; in the glass industry as a decolorizer and pigment; in ceramic industry to introduce color glazes; in rubber industry as a sulfur supplement; and it is also used in photographic exposures; metal finishing operations etc. The effluent from these industries is discharged to landfills, streams and open land management which would contribute significantly to environmental Se(14).

Agricultural activities are also responsible for some of the selenium released into the environment. Fertilizers made from rock phosphate may contain as much as 200 mg. Se per kg. These concentrations are generally much higher than those measured in fertilizers derived from rock carbonates, synthetic nitrogen fertilizers and potassium sulfates. Se deficient soils are often sulfur deficient, especially for the production of legumes. Increasing the concentration of sulfates in the forage also may have detrimental effect on Se availability to the animals (2).

REFERENCES

1. Rosenfeld, I. and Orville B., **Chemistry of Selenium**, McGraw Hill, New York, 1964, pp 299-329.
2. McNeal, L. and Blisstrieri L.S., **Geochemistry and Occurence of Selenium-An Overview**; American society of agronomy inc. special publication no. 23, 1989 pp. 15-50.
3. Schroeder, H.A., et al, "Diseases due to Selenium", **Journal of Chronic diseases**, 23:227, 1970.
4. Faust Samuel and Osman Aly, "Trace Elements in Aquatic Environment", **Chemistry of Natural Waters**, John Wiley Publishers, 1981, pp 301-380.
5. USEPA, National Interim Primary Drinking Water Regulations, EPA - 570/9-76-003, Washington, D.C., 1976.
6. Luttrell, G.W., "Geology of selenium", US Geological Survey Bulletin No. 1019M, Washington, 1959.
8. Lakin H.W. In: Trace elements in the environment, Advances in Chemistry Serries No. 123, pp. 96, 1973.
9. Byers H.G. et al., "Commercial Uses of Selenium", **Industrial Engineering and Chemical Engineering**, Edition 28:821, 1936.
10. Connor J.J. and Shacklette, H.T. "Background Geochemistry of Some Rocks, Soils, Plants and Vegetables in the Counterminous US", US Geological Survey, 574-F, pp. 126, 1975.
11. National Academy of Sciences (NAS). **Selenium**, Washington, D.C., 1976.
12. Goldberg, E.D. and K.K. Turekian, "Marine Chemistry", **Radioactivity in the Marine Environment**. NAS-NRC, Washington, D.C., 1971.
13. Mayland, James and Panter, "Selenium in Seleniferous Environment" **Selenium in Agriculture and the Environment**, Soil Science Soc. Amer. Special Publication No. 23., 1989.
14. Sudhirender Sharma and Ramendra Singh, "Selenium in Soil, Plants, and Animal Systems" **CRC Vol. 13**, Issue 1, 1989.
15. Johnson, C.M., "Residue Reviews: Selenium in Environment, vol 62, pp 102-125, 1976.
16. Kobata-Pendias, A., "Element of group VI", **Trace Elements in Soils and Plants**, CRC press, 1984, pp. 185- 191.

17. Garrett Fleming, "Essential Micronutrients II: Iodine and Selenium, **Applied Trace Elements**, pp 199-234.
18. Andren, Talmi, "Se in coal fired steam plant emissions" **Environmental Science and Technology**, Vol. 9, pp. 856-858, 1975.
19. Eisenberg S.H., "Chemical Characteristics of Selected Flyash Leachates", **Journal of Environmental Science and Health**, pp. 383-402, part 21, 1986.
20. Pillay, Kaminski, "Neutron Activation Analysis of Selenium Content of Fossil Fuels", **Nuclear Applic. Technol.**, 7:478-483.
21. Gregory A cutter, "Selenium in Fresh Water Systems", **Occurrence and Distribution of Selenium**, Milan Ihnat, editor, CRC press Inc., 1989, pp. 243-260.
23. Mayland, James and Soudergger, "Selenium in Seleniferous Environment", **Soil Science Soc. Amer. Special Publication No.23**, 1989, pp. 15-44.
24. Ohlardorf, Harry, "Bioaccumulation and Effects of Selenium in Wildlife", **Soil Science Soc. Amer.**, Special Publication No. 23, 1989, pp. 133-177.
25. Cooke and Bouland, 'Aquatic Chemistry of Selenium", **Environmental Science and Technology**, No.21, 1987, pp. 1214-1219.
26. Owen Bricker, "Redox potential", **Water Analysis**, No.1, 1987, pp. 55-83.
28. Gibson Susan, "Use of Transport/Transformation Matrices for Selenium Assessment", a M.S. thesis, University of Oklahoma, 1981.
29. Millard S.D., "Distribution and Mobility of Selenium in Shallow Groundwater of the Western San Joaquin Valley", **Environmental Science and Technology**, No. 22, 1988, pp. 697-702.
30. **Standard Methods for Examination of Water and Waste Water**, "Preliminary treatment of samples", **American Water Works Association 17th Edition**, 1989. pp 3-5 to 3-9.
31. Kabata-Pendias, A., "Elements of group VI", **Trace Elements in Soils and Plants**, CRC Press, 1984, pp. 185-191.

END

**DATE
FILMED
9 / 2 / 92**