National Synchrotron Light Source Annual Report 1991 Volume II

(For the period of October 1, 1990 through September 30, 1991)

Editors:
S. L. Hulbert and N. M. Lazarz

April 1992

The National Synchrotron Light Source Department
is supported by the
Office of Basic Energy Sciences
United States Department of Energy
Washington, D.C.

Brookhaven National Laboratory
Associated Universities, Inc.
Upton, New York 11973
Under Contract No. DE-AC02-76CH00016 with the
United States Department of Energy

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency, contractor or subcontractor thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency, contractor or subcontractor thereof.

Printed in the United States of America
Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

NTIS price codes:
Printed Copy: A14; Microfiche Copy: A01

TABLE OF CONTENTS

VOLUME II

SECTION I

List of Abstracts: VUV Beamlines	3
Abstracts of VUV Beamlines	11

SECTION II

List of Abstracts: X-ray Beamlines	83
Abstracts of X-ray Beamlines	103

TABLE OF CONTENTS

VOLUME I

SECTION I

INTRODUCTION

Introduction by the Chairman	Denis McWhan	3
Users' Executive Committee	Mark Rivers	4
Scientific Program Support Committee	Randy Alkire/Roger Klaffky	6

SCIENTIFIC DISCIPLINES:

Atomic and Molecular Science	Mei-Ling Shek	11
Energy Dispersive Diffraction	Earl Skelton	17
Lithography, Microscopy and Tomography	Gwyn Williams	20
Nuclear Physics	Andrew Sandorfi	24
UV Photoemission and Surface Science	Chi-Chang Kao	26
X-ray Absorption Spectroscopy	Lars Furenlid/Chuck Bouldin	32
X-ray Scattering and Crystallography	Lonny Berman/Peter Siddons	36
X-ray Topography	Michael Dudley	43

SECTION II

SYMPOSIA, WORKSHOPS AND PROJECTS

NSLS Users' Meeting	Mark Rivers	47
Workshop on Surface Structure	Doon Gibbs	49
Workshop on Electronic and Chemical Phenomena at Surfaces	Kevin Smith/Neal Shinn	50
Workshop on Imaging	Harald Ade/Shawn Williams	51
UV FEL Machine Reviews	Sam Krinsky/Ilan Ben-Zvi	52
DOE High School Honors Program	William Lynch	54

SECTION III

FACILITY REPORT

VUV Machine Operations	Steve Kramer	59
VUV Beamline Operations	Roger Klaffky	63
VUV Storage Ring Parameters	Nick Gmür, et. al.	64
X-ray Machine Operations	Norman Fewell	65
X-ray Beamline Operations	Roger Klaffky	68
X-ray Storage Ring Parameters	Nick Gmür, et. al.	70
Users Facilities	William Thomlinson	71

SECTION IV

NSLS PROJECTS

Superconducting X-ray Lithography Source (SXLS)	Richard Heese	77
SXLS Storage Ring Parameters	Jim Murphy, et. al.	78
The Accelerator Test Facility	Ilan Ben-Zvi	79
Proposed UV-FEL User Facility at the NSLS	Ilan Ben-Zvi/Sam Krinsky/Li-Hua Yu	81
Global Orbit Feedback Systems	Om Singh/Li-Hua Yu/Sam Krinsky	84
NSLS Computer System	Ed Desmond	85

SECTION V

1991 NSLS OPERATIONAL HIGHLIGHTS

Beamline Operational Highlights	89
---------------------------------------	----

SECTION VI

INFORMATIONAL GUIDE

Informational Guide to the NSLS VUV Beamlines	User Administration	105
Informational Guide to the NSLS X-ray Beamlines	User Administration	109

SECTION VII

APPENDICES

List of Abstracts: VUV Beamlines (Abstracts Located in Volume II)	Lisa Lo Sordo	119
List of Abstracts: X-ray Beamlines (Abstracts Located in Volume II)	Lisa Lo Sordo	127
NSLS Users' Publications	Lisa Lo Sordo	147
NSLS Staff Publications	Lydia Lever	167
NSLS Staff Preprints	Lydia Lever	173
NSLS Staff BNL Formal and Informal Reports	Lydia Lever	175

FRONT COVER PHOTOGRAPHS

- a - The National Synchrotron Light Source, Building 725.
- b - Scanning electron microscope photos of 0.1 micron lines and spaces in a 60 nm thick layer of PMMA resist, patterned using 3 different fabrication techniques by the Advanced Lithography Research Department of AT&T Bell Labs at beamline U13UB. See Section I - Lithography, Microscopy and Tomography.
- c - Three $2 \times 2 \mu\text{m}^2$ images of an Au test object, as viewed in the X1A Scanning Transmission Electron Microscope (STEM). See Section I - Lithography, Microscopy and Tomography.
- d - Copper monolith body of the beam position monitor developed by E. Johnson and T. Oversluizen (NSLS Beamline R & D Section). See Rev. Sci. Instrum. 60, 1947 (1989).
- e - The crystal structure of the superconducting Buckminsterfullerence compound K_3C_{60} . The raspberries are C_{60} and the smaller isolated spheres are potassium cations. This structure was solved by Peter Stephens and coworkers at SUNY and UCLA, from powder diffraction data taken at beamline X3. See Section I - X-ray Scattering and Crystallography.
- f - Andy Warkentien (Technical Specialist in the NSLS Mechanical Group) and John Flannigan (Computer Analyst in the NSLS Computer Controls Group) adjust the liquid helium refrigerator to optimize the performance of the X17 Superconducting Wiggler Cryogenic System.

- g - Movable exit slit chamber for the U13UA beamline, sandwiched between long welded bellows and mounted on a 880 mm-travel linear slide. The distance from the sample chamber to the center of the U13 insertion device is 23 meters. S. L. Hulbert (NSLS, Beamline R & D Section).
- h - The Phase I Superconducting X-ray Lithography Source (SMLS) Storage Ring.
- i - VUV front-end fast valves during assembly.
- j - Tony Lenhard (NSLS Beamline R & D Section) inspecting an adjustable crystal holder for a double crystal monochromator. These monochromators are currently installed at X19A, X12B/C and X7.
- k - Crystal holders for a double crystal monochromator. T. Oversluizen (NSLS Beamline R & D Section).
- l - Molecular structure of Buckminsterfullerene, C_{60} . D. Cox (BNL Physics) and P. Heiney and coworkers (U. Penn) at beamline X7A have carried out high resolution powder diffraction experiments which revealed the existence of an orientational ordering transition in C_{60} at 250K and an analogous transition in C_{70} . See Section I - X-ray Scattering and Crystallography.
- m - Schematic of the water cooled copper monolith body of the NSLS developed beam position monitors. The tungsten blades which are used to sense the beam position are also shown. E. Johnson and T. Oversluizen (NSLS Beamline R & D Section). See Rev. Sci. Instrum. 60, 1947 (1989).

BACK COVER PHOTOGRAPHS

- n - The U4IR beamline on the VUV Storage Ring includes three Nicolet interferometers covering the wavelength region from 1 micron to 1 mm. See Section V - Beamline Operational Highlights.
- o - A 44 year old male sustained an acute myocardial infarction on Feb. 19, 1991. On Feb. 25, cardiac catheterization revealed an 80% stenosis of the Right Coronary Artery (RCA) - see left hand Cine (film) image. Balloon angioplasty was performed to re-open the RCA - see middle Cine image. Transvenous coronary angiography was performed on Aug. 22, 1991 at the Synchrotron Medical Research Facility (SMERF) showing that the RCA was still patent - see right hand SMERF image, Left Anterior Oblique 40° view. Beam parameters: X17 superconducting wiggler field was 3.0 Tesla; ring current was 206mA.
- p - Computed microtomography (CMT) image of the internal structure in a 1 cm fragment of the Allende carbonaceous meteorite. This 2-dimensional image is one of 40 slices in a 3-dimensional tomogram obtained in the superconducting wiggler beamline (X17B1). The high energy wiggler radiation is essential to penetrate these large, dense objects. CMT of meteorites is valuable because of the ability to determine the positions of the pseudo-circular objects (silicate-bearing chondrules that formed in the early solar nebula) prior to developing sampling strategies and allocation plans. The techniques being developed at X17B1 will be particularly useful in studying rare meteorites collected in Antarctica, some of which come from the Moon and probably Mars.

- q - Developers of the X1A SPEM (left to right) D. Kern, IBM, E. Anderson, LBL, E. Johnson (back), NSLS, J. Kirz, SUNY at Stony Brook, S. Hulbert, NSLS, and H. Ade, SUNY at Stony Brook. H. Ade and J. Kirz were the principal investigators of this collaborative project which received a 1991 R & D 100 Award.

- r - Using a soft x-ray undulator source and a Fresnel zone plate to produce a focused x-ray probe, researchers at X1A have obtained 50 - 75 nm resolution images of the ZnS-based P31 phosphor grains shown here. C. Jacobsen, S. Lindaas, S. Williams, and X. Zhang (SUNY at Stony Brook). See Section I - Lithography, Microscopy and Tomography.

- s - Laue crystal used in monochromators.

- t - A Couette Shear Cell at beamline X10A is designed for *in-situ* x-ray scattering of macromolecular fluids under flow. The purpose of this research program is to explore new non-equilibrium steady-state structures of complete fluids that are produced under shear flow, due to the continuous input and dissipation of energy. See Section V - Beamline Operational Highlights.

- u - Beamline U3A during re-assembly. The large chamber at right-center houses the new monochromator. Two of three calibration chambers are located left (downstream) of the monochromator. See Section V - Beamline Operational Highlights.

- v - X-ray scattering at beamline X22C. Upper figure: The diffraction pattern obtained for Pt(111) at high temperatures in the reconstructed phase. Lower figure: "ring scans" taken around the (1,0) reflection as a function of temperature. A. R. Sandy, S. G. J. Mochrie (MIT), D. M. Zehner (ORNL), G. Grübel, K. G. Huang, and D. Gibbs (BNL Physics). See Section I - X-ray Scattering and Crystallography.

- w - Two X1A-SPEM images of a microfabricated device, acquired simultaneously by monitoring the sample current (representative of the sum of all electrons liberated) together with the intensity of the electron energy analyzer. D. Kern, IBM, E. Anderson, LBL, E. Johnson, NSLS, J. Kirz, SUNY at Stony Brook, S. Hulbert, NSLS, and H. Ade, SUNY at Stony Brook. See Section I - Lithography, Microscopy and Tomography.

**Schwarzschild Camera Images in PMMA
Using Patterned Reflection Masks**

**Etched Ge
Absorber**

**Liftoff of
Ge Absorber**

**RIE
Multilayers**

Section I
VUV Abstracts

List of Abstracts: VUV Beamlines

Beamline U1

Characterization of Fluorocarbon Polymer Films and Adsorbed Protein by Ultra-Soft X-ray Absorption Spectroscopy D.G. Castner, K.B. Lewis, D.A. Fischer, B.D. Ratner, and J.L. Gland.....	11
A Fluorescence-Yield Near-Edge Spectroscopy (FYNES) Study of the Formation and Surface Reaction of NiO on Ni(100) J.G. Chen, J.H. Hardenbergh, R.B. Hall, and D.A. Fischer.....	11
Fluorescence Yield Oxygen K Edge Study of Reduced/Reoxidized $\text{YBa}_2(\text{Cu}_{1-x}\text{Fe}_x)_3\text{CuO}_{7-8}$ D.A. Fischer and A.R. Moodenbaugh.....	12
Observation and Identification of a New Resonance in the Graphite Carbon K Near Edge Spectrum D.A. Fischer, R.M. Wentzcovitch, R. Carr, A. Continenza, and A.J. Freeman.....	12
Distortions of Fluorescence Yield X-ray Absorption Spectra Due to Sample Thickness G. Meitzner and D.A. Fischer.....	13
EXAFS Studies of Fluorine in Solid-Acid Catalysts G. Meitzner, G.B. McVicker, and D.A. Fischer.....	13
Near-Edge and EXAFS Studies of Binary Alloys of 3d Metals G. Meitzner, J.H. Sinfelt, and D.A. Fischer.....	14
Near-Edge and EXAFS Studies of Carbon on VUV Beamline U-1 G. Meitzner, D. Cox, and D.A. Fischer.....	14
X-ray Absorption Spectroscopic Study of New Carbon Allotrope C_{60} , Buckminsterfullerene G. Meitzner and D. Fischer.....	15
Analysis of Engine Deposits by Carbon and Oxygen Near Edge Spectroscopy P.A. Stevens and A.M. Schilowitz.....	15
Near Edge Absorption Measurements of the Decomposition of $\text{C}_4\text{H}_4\text{S}$ on Pt(111) P. Stevens, J. Gland, T. Rufael, and A. Yeh.....	16
X-ray Absorption Study of Diamond-Like Nanocomposites X.Q. Yang, V. Dorfmann, T.A. Skotheim, D.A. Fischer, and B. Pypkin.....	16

Beamline U2

Chemically Amplified Negative Photoresist for X-ray Lithography W. Conley, J. Gelorme, G. Spinillo, K. Stewart, and R. Wood.....	17
---	----

Beamline U3C

Progress Report of Calibration Work Done on U3C T. Archuleta, A. Bennett, R. Balke, B. Coulter, J. Cobble, R. Hockaday, T. Ortiz, S. LaMarra, R. Montoya, E. Gullikson, S. Gallagos, J. Gonzales, P.P. Gong, and Leif Singman.....	17
Photoelectron Spectroscopy in Ce Heavy Fermions: Inconsistencies with the Predictions of the Prevailing Kondo Theory A.J. Arko, J.J. Joyce, J.M. Lawrence, R.J. Bartlett, P.C. Canfield, Z. Fisk, and J.D. Thompson.....	18
Photoemission in YbCu_2Si_2 : Problems with the Kondo Model A.J. Arko, J.J. Joyce, J.M. Lawrence, R.J. Bartlett, P.C. Canfield, Z. Fisk, and J.D. Thompson.....	18
Quasiparticle Lineshapes in $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ A.J. Arko, J.J. Joyce, R.J. Bartlett, R. Liu, B.W. Veal, and A.P. Paulikas.....	19
Search for the Kondo Resonance in Ce Compounds A.J. Arko, J.J. Joyce, R. Bartlett, P. Canfield, Z. Fisk, J.L. Smith, J.D. Thompson, and J. Lawrence.....	19
Double Photoionization Studies of Argon, Neon and Helium R.J. Bartlett, P.J. Walsh, and J. He.....	20
Double Photoionization Studies of He from Threshold to 560 eV R. Bartlett, P. Walsh, J. Samson, and Z. He.....	20

Partial Photoionization Cross Sections of Ne Across the K-Edge R.J. Bartlett, P.J. Walsh, and J. He	21
Be, BeO, and Mo Mirror Calibrations R. Blake	21
Photoemission Responses and Resonant Behavior in $(\text{Mg}_{1-x}\text{Fe}_x)_2\text{SiO}_4$ L.M. Hirsch and T.J. Shankland	22
Photoemission Study of Electronic Structure Near EF in $\text{YBa}_2\text{Cu}_3\text{O}_x$, for $6.3 < x < 6.9$ R. Liu, B.W. Veal, A.P. Paulikas, J.W. Downey, A.J. Arko, and J.J. Joyce	22
Calibration of a Soft X-ray Film Spectrograph T.J. Tanaka, N.D. Shinn, R.E. Ripple, A. Widman, and M.A. Palmer	23
ARPES Study of $\text{YBa}_2\text{Cu}_3\text{O}_{6.7}$ and $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$ B.W. Veal, A.P. Paulikas, R. Liu, A.J. Arko, and J.J. Joyce	23
ARPES Study of $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ B.W. Veal, A.P. Paulikas, R. Liu, A.J. Arko, and J.J. Joyce	24
Diffusion in Ti:Si Multilayers Probed by Anomalous Dispersion in the Soft X-ray Regime Z. Xu, Z. Tang, S.D. Kevan, T. Novet, and D.C. Johnson	24

Beamline U4A

ARP Study of the Clean and H-Covered Pt(111) Surface W. Di, K. Smith, and S.D. Kevan	25
Partial Densities of States of Cu-Pd Alloys S.-J. Oh, M. Han, J.W. Allen, and G.K. Wertheim	25
Surface Atom Properties of Ta(110) D.M. Riffe, G.K. Wertheim, and P.H. Citrin	26
Vibrational Broadening of Bulk and Surface Core Levels of Na(110) D.M. Riffe, G.K. Wertheim, and P.H. Citrin	26
Surface Bands on Mo(001) and Mo(001)-2H K.E. Smith and S.D. Kevan.	27
Electron Mean Free Path in the Alkali Metals G.K. Wertheim, D.M. Riffe, and P.H. Citrin	27
Thermal Shift of Core-Electron Binding Energies G.K. Wertheim, D.M. Riffe, and P.H. Citrin	28

Beamline U4B

$L_{2,3}$ Edges and Electronic Structure of Iron in Molecular Complexes C. Cartier, P. Rudolf, A. M. Flank, and C. T. Chen	28
Spin Transition Evidenced by Soft X-ray Absorption Spectroscopy C. Cartier, P. Rudolf, A. M. Flank, and C. T. Chen	29
Electronic States and Phases of K_xC_{60} C.T. Chen, L.H. Tjeng, P. Rudolf, G. Meigs, J.E. Rowe, J. Chen, J.P. McCauley Jr, A.B. Smith, W.J. Romanow, and E.W. Plummer	29
Studies of Y-Ba-Cu-O Single Crystals by X-ray Absorption Spectroscopy A. Krol, Z.H. Ming, Y.H. Kao, N. Nücker, G. Roth, J. Fink, G.C. Smith, A. Erband, G. Muller-Vogt, J. Karpinski, E. Kaldis, and K. Schönmann	30
Magnetic Moment in a Gd Iron Garnet Studied by Soft-X-ray Magnetic Circular Dichroism P. Rudolf, F. Sette, L.H. Tjeng, G. Meigs, and C.T. Chen	30
Giant Cu 2p Resonances in CuO Valence-Band Photoemission L.H. Tjeng, C.T. Chen, J. Ghijsen, P. Rudolf, and F. Sette	31
Soft X-ray Magnetic Circular Dichroism: A New Technique for Probing Magnetic Properties of Magnetic Surfaces and Ultrathin Films L.H. Tjeng, Y.U. Idzerda, P. Rudolf, F. Sette, and C.T. Chen	31
Soft X-ray Absorption Spectroscopy on Ni Model Compounds J. van Elp, S.J. George, J. Chen, and S.P. Cramer	32

Beamline U4-IR

Pulsed Infrared Photoresponse Studies of Superconducting YBCO Films G.L. Carr, D.B. Romero, D.B. Tanner, S. Etemad, F. DeRosa, and A. Inam.....	32
Time-Resolved IR Studies of Photo-Induced States in High- T_c Superconductors G.L. Carr, D.B. Romero, D.B. Tanner, S. Etemad, A. Inam, W. O'Brien, D. Ederer, C.J. Hirschmugl, and G.P. Williams.....	33
Potassium Enhanced Oxidation of Cu(100) F.M. Hoffman, K. Lin, R. Tobin, P. Dumas, G.P. Williams, and C.H. Hirschmugl.....	33
Infrared Transmission of Superconducting Niobium Films C. Kendziora, D. Mandrus, D. Koller, S. Gausepohl, L. Mihaly, C. Hirschmugl, and G. Williams	34
Far-IR Study of Adsorbates on Cu(100) K.C. Lin, R.G. Tobin, P. Dumas, C. Hirschmugl, and G.P. Williams	34
Far Infrared Measurements of the Absorption of Ammonia Using a Mach-Zender Interferometer K.D. Moeller, T. Sears, H.T. Liu, C.J. Hirschmugl, and G.P. Williams	35

Beamline U5

Spin Polarized Photoemission Studies of Surface and Interfacial Magnetism N.B. Brookes, Y. Chang, A. Clarke, P.D. Johnson, and M. Weinert.....	35
Studies of K on Fe(001) with Spin Polarized Photoemission Y. Chang, N.B. Brookes, and P.D. Johnson.....	36
Spin-Polarized Photoemission Study of the Pd/Fe(100) Interface T. Chang, N.B. Brookes, and P.D. Johnson.....	36
Local Spin Correlations in Ultra-Thin Fe/W(100) Films G.A. Mulhollan, R.L. Fink, J.L. Erskine, and G.K. Walters	37

Beamline U6

X-ray Mask Distortion Due to Radiation Damage R.E. Acosta.....	37
X-ray Radiation Damage and a Reliability Study on Bipolar Devices D. Boyne, L. Hsia, R. Wachnik, R. Decker, and C. Wasik	38
Chemically Amplified Negative Photoresist for X-ray Lithography W. Conley, J. Gelorme, G. Spinillo, K. Stewart, and R. Wood.....	17
Effect of Synchrotron X-ray Radiation on the Channel Hot-Carrier Reliability of Reoxidized Nitrided Silicon Dioxide G.J. Dunn	38
A Study of X-ray Damage Effects in Open-Bottom Trench Isolation for Bipolar Devices L.C. Hsia	39
A Search for Neutral Electron Traps in Synchrotron Irradiated Silicon/Silicon Dioxide Structures P.M. Lenahan, T. Weist, L.C. Hsia, and J. Maldonado.....	39
X-ray Mask Repair A. Wagner, J.P. Levin, J.L. Mauer, P.G. Blauner, S.J. Kirch, P. Longo, and J.P. Silverman	40

Beamline U7A

The Condensation and Photon Decomposition of Diborane M.W. Ruckman, J. Mowlem, and D. Strongin.....	40
Photon-Assisted Formation of Diamond-Like Carbon Films from Organics M.W. Ruckman, J. Mowlem, and D. Strongin.....	41
Reaction of Potassium with Ammonia at Liquid Nitrogen Temperatures M. Ruckman, E. Horache, and M. Strongin.....	41
Photon-Assisted Reaction of Diborane and Ammonia to Form Boron Nitride D. Strongin, J. Mowlem, M.W. Ruckman, and M. Strongin	42

Beamline U7B

Valence States in Thin Eu-Rh and Eu-Pd Films C. Berg, S. Raaen, and M.W. Ruckman	42
Photon Modification of Aluminum Oxide on 6063 Aluminum T.S. Chou and M.W. Ruckman	43
Photoelectron Spectroscopic Studies of the Electronic Structure of Ferrous Sulfide P.M. Jones, J.A. Guckert, J.A. May, and E.I. Solomon	43
Polarization Dependence of X-ray Absorption in O ₂ ; the Square Well Model P. Kuiper and M.W. Ruckman	44
Spin-Resolved XAS of O ₂ P. Kuiper and M.W. Ruckman	44
Valence Density of States of Hydrogen-Reduced NiO(100) M.A. Langell, K.W. Wulser, and B.P. Hearty	45
Electronic Properties of Body-Centered-Tetragonal Copper H. Li, S.C. Wu, J. Quinn, Y.S. Li, D. Tian, and F. Jona	45
Structural and Electronic Properties of Ultrathin Films of Gd, Tb, Dy, Ho and Er H. Li, S.C. Wu, J. Quinn, Y.S. Li, D. Tian, and F. Jona	46
Ultra-Thin Films of Rh on Au(001) and Rh on Ag(001): Growth Mode and Magnetism H. Li, S.C. Wu, J. Quinn, Y.S. Li, D. Tian, and F. Jona	46
Core Level Lineshapes in Fe-V Alloys S.L. Qiu, R.G. Jordan, M.W. Ruckman, and M. Strongin	47
Partial Densities of States in CuAuI S.L. Qiu, R.G. Jordan, M.W. Ruckman, and M. Strongin	47
Atomic and Electronic Structure of Thin Films of Mn on Pd{111} D. Tian, H. Li, S.C. Wu, F. Jona, and P.M. Marcus	48
Re-Examination of the Electron Band Structure of Tb Along $\Gamma\Delta A$ S.C. Wu, H. Li, Y.S. Li, D. Tian, J. Quinn, F. Jona, D. Fort, and N.E. Christensen	48
Relativistic Effects in the Electron Band Structure of Cu{111} S.C. Wu, H. Li, Y.S. Li, D. Tian, and F. Jona	49
Surface State on Clean Tb(0001) S.C. Wu, H. Li, Y.S. Li, D. Tian, J. Quinn, and F. Jona	49

Beamline U8A/B

Chlorine Adsorption on Room Temperature GaAs(110) D.K. Shuh and J.A. Yarmoff	50
Unfilled Orbitals of C ₆₀ and C ₇₀ from Carbon K-Shell X-ray Absorption Fine Structure L.J. Terminello, D.K. Shuh, F.J. Himpsel, D.A. Lapiano-Smith, J. Stöhr, D.S. Bethune, and G. Meijer	50
Photoelectron Spectroscopy Study of the Cr/MoS ₂ Interface T.D. Durbin, D.K. Shuh, J.A. Yarmoff, J.R. Lince, and S.V. Didziulis	51
Photoemission Studies of the Chemisorption of Dichlorosilane on the Si(111) and Si(100) Surfaces T.D. Durbin, D.K. Shuh, J.A. Yarmoff, D.A. Lapiano-Smith, F.J. Himpsel, and F.R. McFeely	51
The Decomposition of BCl ₃ on Si(111) and the Subsequent Incorporation of Boron into Silicon D.A. Lapiano-Smith and F.R. McFeely	52
A Mechanism for Diborane Induced Selectivity Loss in the CVD of Silicon from SiH ₂ Cl ₂ D.A. Lapiano-Smith and F.R. McFeely	52
Studies of the XeF ₂ Etching of Si(111) at 200°C C.W. Lo, D.K. Shuh, and J.A. Yarmoff	53
Construction and Launch of an Improved Soft X-ray Telescope E. Spiller, J. Wilczynski, L. Golub, and G. Nystrom	53

Beamline U9B

Structural Analysis of Phage T7 Gene 5.9 Product, Using Circular Dichroism Spectroscopy and Primary Sequence Analyses	
L.L. France, L. Lin, B. Studier, and J.C. Sutherland	54
Structural Analysis of Variable Membrane Protein 7 from <i>Borrelia HermsII</i> , Using Circular Dichroism Spectroscopy	
L.L. France, J.J. Dunn, and J.C. Sutherland	54
A Vacuum UV CD Study of Pyrophosphatase Wild Type and Mutants	
M. Gulotta, D.J. Goss, and B. Cooperman	55
The Vacuum UV CD Spectrum of A G · G · C Triplex	
K.H. Johnson, M.E. Hogan, and J.C. Sutherland	55
Circular Dichroism Study of an Oligodeoxyribonucleotide Triplex	
Y.-W. Park, G.E. Plum, K.J. Breslauer, and J.C. Sutherland	56

Beamline U10A

Investigation of the Al L Exciton in AlGaAs by Soft X-ray Reflection and Emission	
Q.-Y. Dong, W.L. O'Brien, J.J. Jia, T.A. Callcott, D.R. Mueller, and D.L. Ederer	56
Copper M _{II,III} X-ray Emission Spectra for CuO, Cu ₂ O and Bi ₂ Sr ₂ CaCu ₂ O ₈	
D.R. Mueller, D.L. Ederer, J.J. Jia, W.L. O'Brien, Q.Y. Dong, and T.A. Callcott	57
The N _{IV,V} → O _{II,III} Soft X-ray Emission Spectrum of La	
D.R. Mueller, C.W. Clark, D.L. Ederer, J.J. Jia, W.L. O'Brien, Q.Y. Dong, and T.A. Callcott	57
Electronic Structure of MgO, α-Al ₂ O ₃ and MgAl ₂ O ₄ , Occupied and Unoccupied Bulk DOS	
W.L. O'Brien, J. Jia, Q.-Y. Dong, T.A. Callcott, D.T. Mueller, and D.L. Ederer	58
Phonon Coupling to Core Excitations in Ionic Solids	
W.L. O'Brien, J. Jia, Q.-Y. Dong, T.A. Callcott, D.R. Mueller, and D.L. Ederer	58

Beamline U11

Photon-Induced Chemistry in Weakly-Bound Thiophene Clusters	
P. Cid-Aguero, E.A. Walters, and J.R. Grover	59
Photoionization Studies of Trifluoromethyl Bromide and a Cluster with Methanol	
J.T. Clay, E.A. Walters, M.V. Willcox, and J.R. Grover	59
Complexes of Oxygen with Benzene and Hexafluorobenzene	
J.R. Grover, G. Hagenow, and E.A. Walters	60
Kinetic Energy Release in the Dissociative Photoionization of Van Der Waals Clusters	
J.R. Grover, E.A. Walters, D.L. Arneberg, G. Hagenow, J.T. Clay, and M.V. Willcox	60
Photoionization Mass Spectrometric Study of N + C ₂ H ₅ Reaction Products	
R.B. Klemm, W. Tao, F.L. Nesbitt, and L.J. Stief	61
Photoionization Spectra and Ionization Thresholds for H ₂ CN and D ₂ CN	
F.L. Nesbitt, G. Marston, L.J. Stief, M.A. Wickramaarachchi, W. Tao, and R.B. Klemm	61
A Possible Role for Ions in Flame Extinguishment by Halons	
E.A. Walters, J.R. Grover, J.T. Clay, M.V. Willcox, D.L. Arneberg, and J.S. Ninitz	62

Beamline U12B

Coverage Dependence of the Li K-Edge Absorption for Li/Be(1120)	
P.A. Bruhwiler and E.W. Plummer	62
Angle-Resolved Photoemission Investigation of Bi Overlayers on InSb(110) Surfaces	
N.J. DiNardo, D. McIlroy, D. Heskett, A.B. McLean, R. Ludeke, and H. Munekata	63
Core-Level Spectroscopy From Low-Index Surfaces of HfC Exposed to Oxygen	
G.R. Gruzalski, S.-C. Lui, D. Heskett, X. Shi, and D.M. Zehner	63
The Experimental Band Structure of Be(1120)	
J.B. Hannor, G. Lee, and E.W. Plummer	64

Core-Level Spectroscopy from Low-Index Surfaces of TaC S.-C. Lui, G.R. Gruzalski, D. Heskett, X. Shi, and D.M. Zehner	64
Interaction of Hydrogen with Mg(0001) P.T. Sprunger, P.A. Bruhwiler, and E.W. Plummer	65
Bulk Band Structure of Lithium G.M. Watson, X.Y. Liu, and E.W. Plummer	65

Beamline U13UA

Core Electron Excitation of Simple Molecules to Rydberg States - A New Method to See Hidden Resonance Excitations D.Y. Kim, K. Lee, C.-I. Ma, M. Mahalingham, D.M. Hanson, and S.L. Hulbert	66
Selective Excitation of Carbon in Acetone C.-I. Ma, K. Lee, D.Y. Kim, D.M. Hanson, and S. Hulbert	66

Beamline U13UB

X-ray Imaging and Projection Lithography J.E. Bjorkholm, J. Bokor, R.M. D'Souza, L. Eichner, R.R. Freeman, T.E. Jewell, W.M. Mansfield, A.A. MacDowell, E.L. Raab, L.H. Szeto, D.M. Tennant, W.K. Waskiewicz, D.L. White, D.L. Windt, and O.R. Wood, II	67
Soft X-ray Projection Lithography R. Freeman, J. Bjorkholm, J. Bokor, K. Early, L. Eichner, M. Himel, T. Jewell, A. MacDowell, L. Szeto, D. Taylor, D. Tennant, W. Waskiewicz, D. White, D. Windt, O. Wood, II, P. Takacs, J. Bruning, M. Ruda, and K. Thompson	67

Beamline U14A

Line Shape Of The Si $L_{2,3}$ VV Auger Spectrum of Stoichiometric SiO ₂ Measured by APECS Bartynski, R. A., See, A. K., Kao, C.-C., and Hulbert, S.L.	68
Electronic Structure of the SiO ₂ /Si(111) Interface Probed by APECS See, A.K., Bartynski, R.A., Kao, C.-C., Hulbert, S.L.	68
Photoemission Investigation of the Core Levels of Na/Cu(111) X. Shi, D. Tang, D. Heskett, and K.-D. Tsuei	69

Beamline U15

Fragmentation of Gaseous Freons: Electron-Multiple-Ion Coincidence Measurements T.A. Ferrett, S. Hsieh, M. Bayne, D.Y. Kim, K. Lee, C. I. Ma, and D.M. Hanson	69
Electron-Ion Coincidence Mass Spectra of CO R.G. Hayes, K. Lee, C.I. Ma, and D.M. Hanson	70
Electron-Ion Coincidence Mass Spectra of NH ₃ R.G. Hayes, K. Lee, C.I. Ma, and D.M. Hanson	70
Electron-Ion Coincidence Studies of the Behavior of Ammonia Upon Core Electron Excitation R.G. Hayes, K. Lee, C.-I. Ma, and D.M. Hanson.	71
Electron-Ion Coincidence Studies of the Behavior of Methylamine Upon Core Electron Excitation R.G. Hayes, K. Lee, C.-I. Ma, and D.M. Hanson.	71
The Anisotropic Fragmentation of CO ₂ , N ₂ O, and H ₂ O Following Core Electron Excitation D.Y. Kim, K. Lee, C.I. Ma, and D.M. Hanson.	72
Local Environment Around the Oxygen Atoms in Y-Ba-Cu-O Studied by X-ray Absorption Fine Structure (XAFS) Spectroscopy A. Krol, Z.H. Ming, Y.H. Kao, N. Nücker, G. Roth, J. Fink, and G.C. Smith	72
Local Structure Around F in Nd ₂ CuO _{4-x} F _x A. Krol, Y.L. Soo, Z.H. Ming, S. Huang, Y.H. Kao, G.C. Smith, K. Lee, A.C.W.P. James, and D.W. Murphy.	73
Angular Dependence of Photofragments from K-Shell Excited Diatomics K. Lee, C.I. Ma, D.Y. Kim, and D.M. Hanson.	73

Determination of the Harmonic Composition of the Synchrotron Radiation at U15 K. Lee, D.M. Hanson, and Y.H. Kao	74
Electron-Multiple-Ion Coincidence Studies of Methanol C.-I. Ma, K. Lee, D.Y. Kim, and D.M. Hanson	74

Beamline U16A

Spin-Polarized Angle-Resolved Photoemission Studies of Ultra-Thin Fe/W(110) Films A.B. Andrews, G.A. Mulhollan, B.S. Kim, J.L. Erskine, and X. Zhang	75
ARUPS Study of Bulk Gd(0001) B. Kim, A.B. Andrews, J.L. Erskine, K. Kim, and B.N. Harmon	75
ARUPS Temperature Dependence of RKKY Induced Magnetic Splitting in Bulk Gd(0001) B. Kim, A.B. Andrews, J.L. Erskine, K. Kim, and B.N. Harmon	76
High-Resolution Core-Level Spectroscopy of Epitaxial Bi-Metallic Interfaces B. Kim, N.D. Shinn, A.B. Andrews, and J.L. Erskine	76
Angle-Resolved Photoemission Evidence for a Gd(0001) Surface State D. Li, C.W. Hutchings, P.A. Dowben, C. Hwang, R.-T. Wu, M. Onellion, B. Andrews, and J.L. Erskine	77
Temperature Dependence of Core-Level Binding Energies of W(110) and Ta(100) D.M. Riffe, W. Hale, B.S. Kim, and J.L. Erskine	77

Beamline U16B

The Influence of Preadsorbed Potassium on the Adsorption of PF_3 on Ru(0001) at LN_2 Temperatures V. Chakarian, D.K. Shuh, J.A. Yarmoff, H.-S. Tao, B. Maschhoff, and T.E. Madey	78
Chemical Bonding and Structure in Epitaxial Aluminas(III): Analysis of Alumina Valence Band Spectra R.P. Merrill, T.N. Rhodin, S. Woronick, P. O'Hagen, N.D. Shinn, R. Absil, and A. Chester	78
X-ray Spectroscopy of Catalytic Materials: Surface Acidity in Epitaxial Alumina and the Effect of Heat Treatment on Ethylene Adsorption R.P. Merrill, T.N. Rhodin, S. Woronick, P. O'Hagen, N.D. Shinn, G. Woolery, R. Absil, and A. Chester	79
Decay Channels of $\text{Al L}_{2,3}$ Excitons and Absence of O K Excitons in $\alpha\text{-Al}_2\text{O}_3$ W.L. O'Brien, J. Jia, Q.-Y. Dong, T.A. Callcott, D.L. Mueller, D.L. Ederer, N.D. Shinn, and S.C. Woronick	79
Oxidation of W(110): W(4f) Core-Level Spectroscopy C.H.F. Peden and N.D. Shinn	80
The Structure of Thermally Nitrided Si(100) C.H.F. Peden, J.W. Rogers, Jr., N.D. Shinn, K.B. Kidd, and K.L. Tsang	80

CHARACTERIZATION OF FLUOROCARBON POLYMER FILMS AND ADSORBED PROTEIN BY ULTRA-SOFT X-RAY ABSORPTION SPECTROSCOPY

Beam Line U1

D. G. Castner (U. of Washington), K. B. Lewis (U. of Washington), D. A. Fischer (Exxon), B. D. Ratner (U. of Washington), and J. L. Gland (U. of Michigan)

Films of both conventional and radio frequency glow discharge (RF-GD) prepared fluorocarbon (FC) polymers were examined by ultra-soft X-ray absorption spectroscopy. Both total electron yield (EY) and fluorescence yield (FY) measurements were made at the carbon and fluorine K edges. The conventional FC polymers examined were poly(tetrafluoroethylene) (PTFE) and poly(vinylidene fluoride) (PVDF). Several RF-GD FC films deposited on Au and poly(methyl methacrylate) (PMMA) from tetrafluoroethylene (TFE) and perfluoropropane (PFP) were examined.

The PTFE sample, prepared by skiving a thin film from a rotating cylinder of PTFE, had visible striations on the surface. The near edge X-ray absorption fine structure (NEXAFS) spectra shown below indicate that the CF_2 chains in PTFE are oriented parallel to the surface striations. The transition to the C-C σ^* orbital (peak near 295 eV in the C NEXAFS spectra) is enhanced when the X-ray E-vector is parallel to the striation direction. Correspondingly, the transitions to the C-F σ^* orbitals (peaks near 292, 299, and 689 eV in the C and F NEXAFS spectra) are enhanced when the E-vector is perpendicular to the striation direction.

Preparation of FC films by RF-GD deposition is generally assumed to produce randomly oriented films containing several chemical species (e.g., CF , CF_2 , CF_3 groups). This was the case for all but one of the RF-GD films analyzed. This film, prepared by RF-GD deposition of TFE onto Au, showed strong polarization dependence in the C and F NEXAFS spectra. It appears this film is comprised of CF_2 chains that are aligned vertically on the Au surface. This preferential orientation is retained after soaking the film overnight in buffer solution and then exposing it to a protein solution for two hours. The structure of this RF-GD film may be responsible for its unique protein adsorption behavior. For thin (50-200 Å) FC RF-GD films deposited onto PMMA the type of detection (EY or FY) selected determined which component was examined. EY spectra were characteristic of the FC overlayer, while FY spectra were characteristic of the PMMA substrate.

This research was funded by NIH grant No. RR-01296. Travel assistant for DGC and KBL was provided by the DOE through the NSLS Faculty/Student Support Program.

U1

A FLUORESCENCE-YIELD NEAR-EDGE SPECTROSCOPY (FYNES) STUDY OF THE FORMATION AND SURFACE REACTION OF NiO ON Ni(100)

J.G. Chen, J.H. Hardenbergh, R.B. Hall (Exxon CRSL) and D.A. Fischer (Exxon PRT)

We have investigated the oxidation of Ni(100) and the subsequent reduction of NiO on Ni(100) by using fluorescence-yield near-edge spectroscopy (FYNES). A detailed understanding concerning the oxidation mechanism of Ni(100) and of the reduction kinetics of NiO is of technological importance since NiO-related compounds have been recently proposed as possible catalysts for methane coupling. The technique of FYNES is particularly suited for these types of studies, since it has the capability of detecting oxygen species deep inside the bulk and of monitoring the reduction experiments at moderate pressures (up to 0.1 torr).

A typical FYNES spectrum of NiO/Ni(100) is shown in Figure 1. The major O K-edge features are related to the electronic transition to the 3eg, 3a1g and 4t1u orbitals of NiO, respectively. An analysis of the intensities of the O K-edge features as a function of oxygen exposures (spectra not shown) provides complementary information to our previous investigation of the oxidation mechanism of Ni(100) by using other UHV techniques: The oxidation process at 300 K occurs via the sequence of dissociative chemisorption followed by the formation of amorphous NiO of about 3 monolayers; NiO(100) islands can be obtained by heating the surface to 800 K; and a more ordered NiO(100) overlayer can be produced by repeated cycles of oxygen adsorption at 300 K followed by annealing the surface to 800 K.

An example of the FYNES investigation of the H₂-reduction of NiO is shown in Figure 2. This spectrum was obtained by monitoring the 3a1g O K-edge feature as a function of reaction time. Our results demonstrate that the reaction rate for the reduction of NiO is first-order with respect to hydrogen pressure and zero-order with respect to oxide coverage. Our results also indicate that the activation barrier for the reduction is about 1.8 kcal/mole with a pre-exponential factor of 9.4×10^{13} molecules $cm^{-2} sec^{-1}$.

Figure 1

Figure 2

Fluorescence Yield Oxygen K edge study of Reduced/Reoxidized $\text{YBa}_2\text{Cu}_{1-x}\text{Fe}_x\text{O}_{7-8}$ D.A. Fischer (Exxon) and A.R. Moodenbaugh (BNL)

The oxygen k edge XAS studies were made possible using a fluorescence yield system described in detail in reference¹. This system allows rapid sample exchange (<15 min.) and the bulk sensitivity of fluorescence yield insures that surface effects play a minor role in the data. This is in contrast to the rather surface sensitive electron yield probes used in earlier works. Normalized fluorescence yield spectra for a typical (.08%) iron doped 123 material are presented in the figure below. The figure also shows near edge spectra for this sample after reduction and reoxidation. The reduction in edge jump for the reduced sample compared with the standard represents about a 6.5% loss of oxygen, which compares favorably to the mass change data. The reduced sample near edge spectra shows a large decrease in the first peak which is attributed to the reduction in oxygen 2p states near the Fermi level. This peak has been identified as due to hole states near the Fermi level by several research groups. The loss of these hole states for the reduced sample results in no superconductivity. However, after reoxidation of this sample, we find that the oxygen near edge spectra resembles that of the original unreduced sample. This would indicate that the number of hole states has returned to its original value before reduction and we would expect that the TC should be unchanged. This conclusion is in agreement with TC measurements on these materials.

1. D.A. Fischer, J. Colbert, and J.L. Gland, Rev. Sci. Inst. **60**, 1596 (1989).

OBSERVATION AND IDENTIFICATION OF A NEW RESONANCE IN THE GRAPHITE CARBON K NEAR EDGE SPECTRUM.

D.A. FISCHER, Exxon; R.M. WENTZCOVITCH†, BNL; R. CARR, SSRL, A. CONTINENZA††, A.J. FREEMAN††, Northwestern U.

We show for the first time polarization dependent fluorescence, electron yield and transmission near edge spectra above the carbon K edge for graphite (HOPG and thin films). The spectra consistently display a new peak above the π peak and below the σ threshold, which has not been noticed in previous electron yield near edge spectra. A comparison of our carbon K near-edge spectra data with the one-electron DOS obtained from a full-potential LAPW method indicates that this new feature could be caused by the lowest lying interlayer conduction states which were presumably submerged among the other σ and π states.

† Supported by DMS of US-DOE grant #DE-AC02-76CH00016

†† Supported by NSF (NU MRC)

STORTIONS OF FLUORESCENCE YIELD X-RAY ABSORPTION SPECTRA DUE TO SAMPLE THICKNESS

G. Meitzner (Exxon) and D.A. Fischer (Exxon PRT, NSLS)

Fluorescence yield x-ray absorption spectra are subject to distortions, when regarded as direct measurements of the true absorption coefficient, if the absorbing samples are not thin compared to the absorption length of the element of interest for the incident radiation. The systematic divergence between the fluorescent yield and absorption coefficient is illustrated by figure 1, which shows normalized fluorescence yield versus thickness of carbon films. These measurements were made at an energy just above the carbon K edge. At the right (thick) end of the curve the carbon fluorescent yield will have become insensitive to changes in the value of the absorption coefficient. Since this type of error cannot be detected by inspection of the spectrum, it is necessary to calculate an absorption length for the absorber and to establish that the solution or solid under study provides a thickness at least this short.

Figure 1. Intensities of fluorescence are plotted against sample thickness for a series of self-supporting carbon films. Absorption spectra from such films would comprise a reasonable approximation to measurements of true absorption coefficients for films much thinner than about 1000 Angstroms.

EXAFS STUDIES OF FLUORINE IN SOLID-ACID CATALYSTS

G. Meitzner and G.B. McVicker (Exxon), and D.A. Fischer (Exxon PRT, NSLS)

When it is added to Al_2O_3 -based catalysts to enhance their solid acidity, fluorine can produce AlF_3 from some of the aluminum. Other fluorine is associated with aluminum still bound in the oxide matrix. Figure 1 shows how fluorine EXAFS is able to discriminate these two fluoride environments. These experiments have produced the first physical description of catalyst structures associated with solid acidity.

Figure 1. The field at left shows phase-corrected Fourier transforms of oxygen and of fluorine EXAFS from 1%F- Al_2O_3 . The peaks occur at distances corresponding to coordination spheres of oxygen or aluminum neighbors. The correspondence between peaks in oxygen and fluorine transforms indicates that the fluoride has substituted for Al_2O_3 hydroxyl groups.

The field at right shows superimposed fluorine Fourier transforms from AlF_3 and from 6%F- Al_2O_3 . In this case the similarities indicate that fluoride is mainly in an AlF_3 -like phase.

G. Meitzner and J.H. Sinfelt (Exxon), and D.A. Fischer (Exxon PRT, NSLS)

A study was continued of L-edges of copper and the 3d metals iron, cobalt, nickel in their binary alloys. The magnetic properties of alloys formed from pairs of these metals suggest that alloying is accompanied by electronic rearrangements. The L-edges of the metals are sensitive to d-band filling and can test the rigid band model for changes in magnetization. Work to date indicates that electronic shifts occur but not of a magnitude predicted by the rigid band model (figure 1). In future work on these metals we will use the sensitivity of their L-edges to their oxidation states, to monitor the metals' reducibility in catalysts of the sort used commercially for petroleum upgrading by hydrogen addition and sulfur removal. All this work takes advantage of the ability to maintain a useful pressure of hydrogen above the sample during measurement of spectra with the FYNES end-station on U-1.

Figure 1. The curve shows the change in the areas of nickel L_{II} and L_{III} edge resonances (shaded, in inset) as a function of atomic percent copper in NiCu binary alloys. The resonances result from electronic transitions into the 3d band, and weaken as a result of increased occupancy of the d band by copper 4s electrons.

G. Meitzner and D. Cox (Exxon), and D.A. Fischer (Exxon PRT, NSLS)

We have been using the EXAFS capability of beamline U-1 to study the allotropes of carbon. We first verified our methodology with diamond and graphite films that have well-understood structures. We also measured what we believe is the first EXAFS and the first direct physical structural data on the newly discovered C_{60} allotrope of carbon. The phase-corrected Fourier transformed EXAFS from C_{60} is shown in the figure, below the phase-corrected Fourier transformed carbon EXAFS from graphite. The vertical lines on the graphs indicate true positions of coordination spheres in the two structures. Some mismatches near 2 Angstroms are not yet understood. The EXAFS radial distribution function of C_{60} in the figure is consistent with the proposed truncated icosahedral structure for the C_{60} molecule. This work will be extended in the future to various diamond and graphite-like carbons. The physical information from carbon EXAFS may identify useful sub-classifications of these variable carbon structural types.

G. Meitzner (Exxon Research and Engineering Co.) and D.A. Fischer
(National Institute of Science and Technology)

The isolation of C₆₀ stimulated a flurry of studies directed at characterization of the material. We used soft x-ray extended x-ray absorption fine-structure spectroscopy (EXAFS) for what we believe was the first determination of the carbon-carbon interatomic distances by a physical method. These measurements were made with beamline U-1 at NSLS using total electron yield detection, from C₆₀ deposited on a gold substrate from toluene solution. The sample was free of C₇₀.

The phase-corrected Fourier Transformed EXAFS of C₆₀ and of a graphite film are shown in figure 1. The calculated distances to the first few coordination spheres in C₆₀ and in graphite are represented on the RSFs by vertical lines. The strong peak in the graphite Fourier Transform, at about 2.2 Angstroms, was surprising since no such distance was expected. The distance may have resulted from wrinkles in the graphite sheets that produce structures like fragments of the C₆₀ molecule. The 2.2 Angstrom distance occurs naturally in C₆₀ as a diagonal second neighbor distance.

The low resolution radial distribution function in figure 1 demonstrated a reasonable consistency between the proposed and true structures of the Buckminsterfullerene molecule.

1 D.M. Cox, S. Behal, K. Creegan, M. Disko, C.S. Hsu, E. Kollin, J. Millar, J. Robbins, W. Robbins, R.D. Sherwood, P. Tindall, D. Fischer, and G. Meitzner, *Mat. Res. Soc. Symp. Proc.*, 206, 651(1991).

Figure 1. Phase corrected Fourier Transformed EXAFS from Graphite and from C₆₀ are compared. The graphite data were measured in transmission mode. Vertical lines mark nominal positions of coordination spheres, determined by diffraction for the graphite.

ANALYSIS OF ENGINE DEPOSITS BY CARBON AND OXYGEN NEAR EDGE SPECTROSCOPY

P. A. Stevens (EXXON) and A. M. Schlowitz (EXXON)

Deposits produced in internal combustion engines can reduce engine life and inhibit performance. These deposits are typically produced by oxidative degradation of the lubricant and/or fuel. The first step towards preventing formation of these deposits is to understand their chemical composition. The figures below depict the C(1s) and O(1s) near edge photoabsorption spectra of a typical engine deposit collected in total electron yield mode. Features in C-K edge spectrum indicate the presence of both aromatic and aliphatic carbon species. Similarly, the O-K edge shows the presence of both C-O and C=O species.

Near Edge Absorption Measurements of the Decomposition of C_4H_4S on Pt(111)

P. Stevens (Exxon), J. Gland, T. Rufael, and A. Yeh (U. of Mich.)

We have performed near edge X-ray absorption studies at the C-K edge of the decomposition of C_4H_4S on Pt(111). Fluorescence yield detection was used to monitor photon absorption. This allowed temperature dependent measurements to be performed in vacuum and in a 10 mTorr atmosphere of H_2 .

The C-K edge spectrum of a multilayer coverage of C_4H_4S adsorbed on Pt(111) at 80 K is shown in figure 1. The resonances structures are due to dipole transitions from the C(1s) levels to unoccupied molecular orbitals. The labels in the figure indicate the final state orbital associated with the individual structures. The most intense resonance is the $\pi^*_{3b_1}$ peak at 285.4 eV. The final state orbital identified with this peak is a π^* orbital located predominantly on the S atom and the nearest-neighbor C atoms in the ring.

Figure 2 shows the intensity of $\pi^*_{3b_1}$ peak as a function of temperature ($\beta=0.5$ K/sec). As indicated in the figure one of the spectra was recorded in vacuum while for the other there was a 10 mTorr atmosphere of H_2 present. The experiments indicate that the temperature dependence of the peak is not dependent on the presence of H_2 until a temperature of 380 K. Complementary temperature programmed reduction spectroscopy (TPRS) experiments show that in vacuum some of the thiophene molecules desorb at 160 K while the other react to form a stable surface intermediates. These intermediates then decompose near 380 K to leave C and S atoms on the surface. The photoabsorption experiments indicate that the presence of H_2 does not change the reactive intermediates formed by thiophene on Pt(111), but it does affect their subsequent decomposition.

Figure 1

Figure 2

X-ray Absorption Study of Diamond-Like Nanocomposites

U1A

X. Q. Yang, (DAS/BNL), V. Dorfmann and T. A. Skotheim, Moltech Corporation, D. A. Fischer, Exxon, B. Pytkin, Moscow Institute of Controlling Machines, USSR.

We report the study of the near edge x-ray absorption fine structure (NEXAFS) above the carbon k -edge of Diamond-Like Nanocomposites (DLN) films grown by r.f. glow discharge chemical vapor deposition (CVD) and annealed at 430 and 450 C in air. Near Edge X-ray Absorption Fine Structure study was carried out at beam-line U1-A. Electron total yield method was used to collect the NEXAFS spectra. The spectra were normalized by the spectra of a gold grid. The NEXAFS spectra of untreated DLN film and the DLN film annealed at 450 C are plotted in Fig. 1 and Fig. 2. Feature 1 at 285 eV is attributed to the transition from 1s initial state to a state with π symmetry. While feature 3 at 294 eV is attributed to the transition from 1s state to a final state with σ symmetry. Both of these two features are related to sp^2 bonding. Feature 2 at 289 eV could be assigned to a transition related to the C-H bonding due to the residue of hydrocarbon in the DLN film. The sp^2 bonding is a clear evidence of graphitization. The spectrum of untreated DLN film in Fig. 1 is similar to those spectra of graphitized DLC films we studied before. However, the intensity of feature 1 at 285 eV is much weaker, indicating small percentage of the sp^2 bonding. The relative intensities of feature 1 and feature 3 are weaker. After annealing at 430 C and 450 C, the intensity of feature 1 and 3 are significantly reduced which can be seen in Fig. 2 where the spectrum of DLN film after 450 C annealing was plotted. This is a clear evidence that the ratio of sp^2/sp^3 have a sudden increase after annealing around 430 C. This is in a good agreement with the Raman study.

Figure 1. NEXAFS of untreated DLN film. Figure 2. NEXAFS of DLN film annealed at 450 C.

Work partially supported by U. S. DOE under Contract No. DE-AC02-76CH00016.

Chemically Amplified Negative Photoresist for X-Ray Lithography

Willard Conley, Jeffery Gelorme, Gary Spinillo,
Kevin Stewart, Robert Wood

IBM Corporation
East Fishkill Facility
Rt. 52
Hopewell Jct. N.Y. 12533

Ureca formaldehyde prepolymer, an alkaline soluble novolac resin has been formulated with a triplate compound that produces catalytic quantities of acid after x-ray exposure which creates crosslinking during a brief post-exposure bake step. The resultant latent image is developed in aqueous alkaline medium. The photoresist dubbed T-BXTM (Electron Beam, X-Ray) Photoresist, has demonstrated excellent lithographic properties in x-ray exposure work at IBM beamline U-6 and U-2.

A gold-on-silicon mask (0.8um and 2.5um, respectively) was imaged on wafers with approximately 40um proximity gap. Wafers were adhesion treated in vapor HMDS and the T-BX photoresist was spin coated on the wafers and post apply baked at 90C for 60 seconds. Wafers were exposed using a resolution test mask at doses ranging from 20-120mj and post expose baked on a hotplate at 105C for 2 minutes at Brookhaven, then transported back to IBM East Fishkill and developed in 0.32N TMAH developer for 150 seconds. Wafers were then SEM'd at the 90mj dose to evaluate the resist resolution and profile to determine the feasibility of the system for x-ray lithography. (fig. 1)

Progress Report of Calibration Work Done on U3C.

Tom Archuleta, Elbert Bennett, Richard Blake, Bill Coulter,
Jim Cobble, Robert Hockaday, Tom Ortiz, Steve LaMarra, and Rick Montoya
LANL

Eric Gullikson
LBL

Sonny Gallagos, and Joe Gonzales
EG&G LAO

Ping Po Gong and Leif Singman
EG&G LV

A calibration run was run from November 12, 1990 to November 21, 1990 on U3C to calibrate a transmission grating streak camera system, a gallium arsenide photodiode, a gallium phosphide photodiode and eight aluminum photocathode x-ray diodes.

The transmission grating streak camera system consisted of a gold free standing transmission grating, a LBL active detector (gold photocathode and electron lens to CdS fast phosphor), a 49 element fiber array, and a EG&G LAO optical streak camera with a cooled CCD array readout. This system was used to diagnose SATURN holbraum experiments at Sandia National Laboratory. The calibrations covered 60 eV to 1500 eV, the full range of the instrument. The calibrations improved the system response over the pre-calibration system response estimates considerably and improved the confidence in the spectrometer's inferred measured spectra. The largest discrepancy between the measured response and the system response estimates was a factor of 100.

The aluminum x-ray diodes were calibrated from 40 eV to 1500 eV. These were detectors used to diagnose the SATURN holbraum experiments and the x-ray conversion efficiency measurements at AURORA (KrF laser at LANL). The calibrations agreed to within 10-15% of the independently measured data points. Where detector responses previously had been interpolated and extrapolated there were differences of order a factor of two.

The gallium arsenide and gallium phosphide photodiodes were calibrated between 40 eV and 1500 eV to explore the possibility that these detectors may be good absolute flux transfer detectors in the soft x-ray regime. It is useful to note that these detectors would be expected not to have a significant response jump at carbon and oxygen edges, a common feature of x-ray diodes, hopefully making them less ambiguous flux detectors in these spectral regions.

A.J. Arko, J.J. Joyce, J.M. Lawrence**, R.J. Bartlett, P.C. Canfield, Z. Fisk, and J. D. Thompson (Los Alamos National Laboratory, Los Alamos, NM, 87545)

For the last several years the issue of whether photoemission can see a temperature dependent Kondo resonance (KR) in heavy fermion systems seemed to be resolved. High resolution PES data for CeSi₂ by Patthey et al. [PRL 58, 2810 (1987)] was interpreted within the Gunnarson-Schonhammer (GS) framework [PRB 28, 4315 (1987)] for the Kondo model and the fit was claimed to be exceptionally good. In particular, both the KR (at E_F) and the spin-orbit split sideband (at -0.3 eV) were observed in the near-E_F 4f spectrum, which renormalized with temperature to some degree. The very small amount of temperature dependence was explained as being due to additional split-off crystal field states which have a low degeneracy (1/N approximation). The data, however, to us also seemed to fail the prediction of the spectral width (= kT_K is the Kondo temperature), as well as the spectral weight (should vary as T_K).

We have re-examined a large body of Ce heavy fermion systems using photoelectron spectroscopy in high resolution (ΔE=90 meV at hν=120 eV), checking particularly the so-called KR for compliance with predictions. For CeSi₂ we additionally did a careful study of the temperature dependence at 20K, 80K, and 300K, to see how much of the variation can be explained by simple renormalizations like Fermi function broadening and phonon broadening. The latter was determined by studying the Si-2p core levels at the stated temperatures and fitting the results with Voigt functions to extract the thermal portion (90 meV between 20K and 300K). Inasmuch as all materials experience phonon broadening, it is necessary to account for this first before any additional temperature dependence can be considered.

The top frame in the figure at right is actually the 20K data of Patthey et al. with our Lorentzian fit through the data. Our fit is much better than theirs, indicating that the lineshape is Lorentzian (core-level like). Superimposed on the data is the expected spectrum based on instrument broadened crystal field states as determined from the 3 quasielastic peaks in neutron diffraction. Clearly the observed data is much broader than predictions even accounting for the fact that most of the predicted width comes from the separation of the KR and crystal field states and not their individual widths. Thus the natural lineshape is actually much broader than the entire cluster of crystal field states, let alone the KR itself. The temperature dependence in the bottom frame is from our present data. Applying the thermal broadening from the Si-2p core levels, together with Fermi function broadening, we are able to exactly account for all the weak temperature dependence of the data in the bottom frame. It is not necessary to resort to the Kondo model to explain any of the temperature dependence.

We are thus forced to the following conclusions that: i) the near-E_F 4f features are far too broad for predictions, ii) they are not temperature dependent, except for phonon broadening and Fermi function effects, iii) they do not scale with T_K, and iv) they are observed even in non-Kondo systems where they should be absent. A simple explanation in terms of the GS model does not seem appropriate for heavy fermions.

* Work supported by the US Dept. of Energy
** Permanent address, Univ. of Calif., Irvine

A.J. Arko, J.J. Joyce, J.M. Lawrence**, R.J. Bartlett, P.C. Canfield, Z. Fisk, and J.D. Thompson (Los Alamos National Laboratory, Los Alamos, NM, 87545)

The prevailing theory of photoemission in heavy fermion compounds is the Kondo impurity theory in the 1/N approximation. It predicts a Kondo resonance (KR) in the spectrum located a distance kT_K (T_K is the Kondo temperature) above E_F for the single 4f-electron Ce compounds, and a distance T_K below E_F for the single 4f-hole Yb heavy fermions. Its width is also predicted to be of the order kT_K, although there may be complications from crystal field split sidebands. The Yb compounds are clearly much more advantageous for study since the full weight of the resonance is occupied, unlike the former where only a tail extends below E_F. Additional important predictions from the Kondo model regarding these resonances are that i) their width and spectral weight should scale with T_K, and ii) the spectral weight and width strongly renormalizes with temperature, so that by ~5 T_K the central amplitude is nearly zero.

In this abstract we report results for YbCu₂Si₂, which has a T_K of order 50K as reduced from specific heat or quasielastic linewidth. Experiments were carried out as a function of temperature, at 20K, 80K, and 300K, with resolution varying from 60meV to 95meV. Single crystals were cleaved in-situ at a base pressure of 8x10⁻¹¹ Torr. Some additional crystal field lines are seen in neutron diffraction, but we account for these in our analysis below.

While the full spectrum is not shown in the figure at right, all data were normalized to the Cu 3d emission peak, which should be temperature independent. We show instead in the figure only the first of the 4f¹⁴ bulk components which in the Kondo model should be the KR (The 4f¹⁴ spectrum consists of bulk and surface doublets, separated by ~1eV, with the latter at higher binding energy). All other features have been modeled with Lorentzians and subtracted out.

The top frame consists of the 4f_{7/2} transition fitted with a Lorentzian with values as shown. Superimposed is the expected spectrum based on a resolution broadened KR and crystal field lines as obtained from neutron diffraction. Clearly the measured width is much too broad, having a Lorentzian vs. Gaussian lineshape. The lower frame shows the 4f_{7/2} transition as a function of temperature. Some trivial amount of temperature dependence is observed, which, just as in Ce compounds, can be completely explained by Fermi function and phonon broadening. Thus the Kondo model fails in all of its major predictions. Our results imply that i) there is no temperature dependence, ii) lineshapes and widths are simple core-level-like and much broader than predictions, iii) valence remains constant, and iv) spectral weight is independent of T_K (not shown in the fig.) It is clear that previous interpretations of data in terms of a KR need revision.

* Work supported by the US Dept. of Energy
** Permanent address, UC, Irvine

QUASIPARTICLE LINESHAPES IN $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$

A.J. Arko, J.J. Joyce, R. J. Bartlett (Los Alamos), R. Liu, B.W. Veal, A.P. Paulikas (Argonne)

Ever since the first high quality angle resolved photoemission data on high-Tc materials (Olson et al., on $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$) were obtained, a controversy has raged over the nature of the quasiparticle peaks (are they Landau Fermi liquid, RVB, Marginal Fermi liquid, etc.) The source of the controversy was the lineshape of the data which suggested highly asymmetric peaks, with tails extending to higher binding energy, as well as an energy dependence which seemed to scale better with E rather than E^2 as in the Fermi liquid. Although subsequent analyses cast some doubt on the certainty of linear energy dependence as well their absolute interpretation, the asymmetric lineshapes have never been totally resolved. Part of the problem is that the signal levels in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$ are so small that any background signal, be it due to residual DOS or secondaries, raises the background to intolerable levels.

In $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ the problem is solvable from several points of view. In the first place the bands are much narrower so that the finite aperture of the analyzer does cause a problem. Secondly, a signal can be observed in the very flat bottom of the band (unlike in $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_8$), thus reducing the aperture problem still further. Most important, however, by working in higher zones the important quasiparticle peak intensity near E_F is dramatically enhanced, thus enabling us to distinguish between a background due to DOS or secondaries. We have looked at photoelectron spectra of many samples of $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$, and concluded that indeed most of the background signal is due to residual DOS as can readily be discerned from the figure at right.

In the top frame of the figure we show an angle resolved spectrum of $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ (near Γ , $\Delta E=100$ meV for the top frame, $T=20$ K, $\Delta\theta=2^\circ$). One can see that the quasiparticle peak near E_F is riding on a strong background signal which smoothly blends into the main DOS structure and is clearly not due to secondaries. Thus background subtraction is still somewhat arbitrary, but given the size of the peak it leads to only small errors in the high-binding energy side.

We assumed a linear DOS between -6 eV and 0 eV and subtracted it from the data (dashed line). A high resolution (50 meV) scan of the data subtracted in this fashion is shown in the bottom frame, together with a Lorentzian lineshape superimposed on top of the data. It does not not take much imagination to see that the lineshape is clearly Lorentzian (FWHM ≈ 200 meV), and centered at -160 meV, near the bottom of the band. The Lorentzian is cut off by the Fermi function, as one can see by the deviation from Lorentzian behavior within 30 meV of E_F . Our conclusion is that we are dealing with broad Lorentzian lineshapes characteristic of a Landau Fermi liquid. Indeed, the high binding energy tail is an artifact generated by the broad Lorentzian cut off by a sharp Fermi function.

* Work supported by the US Dept. of Energy.

SEARCH FOR THE KONDO RESONANCE IN Ce COMPOUNDS

A. J. Arko, J. J. Joyce, R. Bartlett, P. Canfield, Z. Fisk, J. L. Smith, J. D. Thompson, (Los Alamos National Lab.), J. Lawrence, (U. C. Irvine)

We have made measurements on four Ce compounds - CeSi_2 , CeGe_2Ni_3 , CeBe_{13} , and CeSb_2 - in our continuing search to try to nail down the source of the two f-electron features in the cerium compounds. The generally accepted explanation is that the peak at ~ -2 eV is a screened f-hole while the peak at E_F is the so-called Kondo Resonance which may consist of spin $5/2$ and $7/2$ components separated by about 0.28 eV. The Kondo resonance explanation carries with it a requirement of a very specific temperature dependence of the peak at E_F depending on the T_K .

We have made these measurements at 80 K and at 300 K on compounds with widely varying T_K , as well as on a system which goes magnetic (CeSb_2) thus implying $T_K = 0$. The measurements were carried out at the Fano resonance of Ce (120 eV) in order to maximize the $5f$ signal relative to the ligand valence band signal. The 170 meV resolution of our apparatus at 120 eV, while not as spectacular as that of Y. Baer and co-workers using a helium resonance lamp, was nonetheless sufficient to clearly resolve the double feature at E_F . We find an astounding similarity among all the spectra and almost no observable temperature dependence for any compound. Indeed, even the CeSb_2 displayed a typical spectrum expected from heavy Fermion systems. In addition, the condition of the surface caused large variations in the relative peak intensities leading us to suspect that some of the features may be surface related.

We are converging on the notion that perhaps there is a more or less universal spectrum associated with Ce compounds which reflects the f-screened and d-screened photoholes.

DOUBLE PHOTOIONIZATION STUDIES OF ARGON NEON AND HELIUM USC

Roger J. Bartlett and Peter J. Walsh (Los Alamos)
John He (University of Nebraska)

The ratio of double photoionization to single photoionization was measured from threshold up to 800 eV for Ar, Ne and He. The measurements were made using the USC Beasline and a Time-of-Flight charge state spectrometer. Thin film filters of B, Be and C were used to reduce the amount of scattered and higher order light in the beam. The results have been compared to literature values over the common photon energy ranges, and agree well with those of Schmidt et. al.¹. The Ne data is in good agreement with the calculations of Chang and Poe.² The He data were compared to the Wannier threshold law with good agreement obtained for the first several volts above threshold.

1. V. Schmidt, N. Sander, H. Kuntsemüller, P. Dhez, F. Wulleumier, and E. Kalline, Phys. Rev. A **13**, 1748 (1976)
2. T. N. Chang and R. T. Poe, Phys. Rev. A **12**, 1432 (1975)

ACKNOWLEDGEMENTS: This work was supported by the U.S. Department of Energy through Los Alamos National Laboratory.

DOUBLE PHOTOIONIZATION STUDIES OF He FROM THRESHOLD TO 560 eV USC

R. Bartlett, P. Walsh (Los Alamos), J. Samson, Z. He (U. of Neb.)

Double photoionization of atoms by a single photon is of considerable interest because it exemplifies the effects of electron-electron correlation particularly when the ionization takes place in the outer shell of the atom. In He the double ionization process is totally due to correlation effects and because of its simplicity accurate calculations can be performed. Past measurements have been limited to only about 150 eV above the threshold except for a recent single data point at 2.8 keV. Also, Samson² has recently hypothesized that the double ionization or ionization with excitation processes can be considered as two step processes where the absorbed photon produces a single electron which in turn can cause a secondary ionization or excitation. Thus, to test these ideas and to provide data up to 600 eV we have measured the double to single ionization ratio for He. The normalized results are shown in Fig. 1, along with several other data sets^{1,3,4}, the theory of Amusia⁵, and the electron impact ionization data of He⁶. (Of particular interest is the good agreement between the electron impact data and the photoionization data, strongly supporting the two-step double ionization process.

Fig. 1 Branching ratio data for He²⁺.

1. J.C. Levin et. al. Phys. Rev. Lett. **67**, 968 (1991).
2. J.A.R. Samson, Phys. Rev. Lett. **66**, 1234 (1990).
3. J.M. Bizau et. al. Phys. Rev. A **24**, 170 (1981).
4. H. Kossmann et. al. Phys. Rev. Lett. **63**, 2040 (1989).
5. M. Ya Amusia et. al. J. Phys. B **8**, 1248 (1975).
6. B. Peart et. al. J. Phys. B **2**, 1347 (1969).

Work supported by the US Dept. of Energy.

U3C
PARTIAL PHOTOIONIZATION CROSS SECTIONS OF NE ACROSS THE K-EDGE
R.J. Bartlett, P.J. Walsh (Los Alamos) and J. He (U. of Nebraska)

Photon impact ionization of an inner shell electron changes the screening of the nuclear charge. This in turn causes core rearrangement of the remaining electrons which may lead to fluorescence and/or the further ejection of one or more electrons. One of the simpler systems in which to study such processes is Ne. Thus, we have measured the partial photoionization cross sections for the first three ionization states of Ne across the K-edge energy range.

The data were taken using a time-of-flight ion state spectrometer and the ERG monochromator on the U3C beam line. The energy resolution of the monochromator at the K-edge (~870 eV) with 10 μ m slits was approximately 1.5 eV. This was not adequate to resolve all of the structure near the edge but does give the general shape of the cross sections. A more serious problem was caused by energy shifts in the data sets taken on different electron fills of the storage ring. This may have been caused by beam position instabilities in the storage ring or by excess heating of the beam line components due to the large emittance of the storage ring. This problem will have to be resolved before satisfactory data can be obtained.

The relative cross sections are shown in Fig. 1. Structure in the first ionization state is due to a Rydberg series leading to the K-edge, while the background continuum below the resonance region is due to single ionization from the valance shell. The double and triple ionization cross sections remain low below the edge gradually rising to the edge. It can be seen that the Rydberg states also affect the formation of the multiply charged states.

* Work supported by the US Department of Energy.

Be, BeO, AND Mo MIRROR CALIBRATIONS

R. Blake, Los Alamos National Laboratory

Two sets of measurements were carried out. A recalibration was done on Be and BeO mirrors, used previously in a spectroscopy experiment by SAIC. Recalibration showed that the mirrors used in the experiment had been contaminated with carbon. Modeling of the measured reflectivities allows an estimate of the amount of carbon on the surface.

A separate research project by Richard Blake of LANL was carried out with the same hardware. Molybdenum mirrors were made by evaporation, sputtering, polishing polycrystalline Mo, and from polished single crystals. It was found that each of the samples had measured reflectivities that matched very well with calculated values over the range of angles and energies where the measurements were of high quality. These measurements are being combined with others in the 5 - 11 KeV range from X8C to derive the low energy optical constants of Mo in the 100 - 600 eV range.

Photoemission Responses and Resonant Behavior in $(\text{Mg}_{1-x}\text{Fe}_x)_2\text{SiO}_4$

L. M. Hirsch and T. J. Shankland (Geophysics Group and Institute of Geophysics and Planetary Physics, MS D447, Los Alamos National Laboratory, Los Alamos, NM 87545)

In this work we are studying the mineral olivine $(\text{Mg}_{1-x}\text{Fe}_x)_2\text{SiO}_4$ as a representative component of the Earth's mantle (the most abundant in the outer 30-400 km of the Earth). We measured valence band ultraviolet photoelectron spectra from Rockport fayalite (Fe_2SiO_4), San Carlos olivine (Mg_2SiO_4), and synthetic forsterite (Mg_2SiO_4) single crystal surfaces cleaved in vacuum. For fayalite at photon energies between 55 and 58 eV, the total photoelectron yield increases beyond that at nearby higher and lower photon energies. This resonant excitation of electrons from $\text{Fe}(3p)$ states, which lie about 57 eV below the Fermi level, to the $\text{Fe}(3d)$ states within the valence band enables the $\text{Fe}(3d)$ -derived valence states to be distinguished from the overlapping $\text{O}(2p)$ and $\text{Si}(3s \& 3p)$ states. The resonance is demonstrated by constant initial state (CIS) spectra as in Fig. 1. $\text{Fe}(3d)$ derived states are found to extend about 18 eV below the Fermi level in Fe_2SiO_4 . From differencing spectra at photon energies in the range 51-62 eV, resonance appears to be dominated by $\text{Fe}(3d)$ with peaks about 2, 6, 10, and 15 eV below the Fermi level E_F . Resonant photoemission results for fayalite share some similarities with results by Lad and Henrich (*J. Phys. Rev.*, B39, 1989) for FeO , Fe_2O_3 , and Fe_3O_4 . Covalent interaction of localized $\text{Fe}(3d)$ electrons with neighboring O^{2-} ligands participate across the spectral width of the valence band and broaden it with respect to the valence band of Mg_2SiO_4 .

Fig. 1. CIS spectra from a cleaved iron orthosilicate single crystal across the $\text{Fe } 3p$ excitation threshold. The $\text{Fe } 3p$ binding energy is near 57 eV.

The similarity of Fe-derived energy levels in Fe_2SiO_4 and in simple oxides such as FeO suggest that iron-bearing silicates can behave electrically like transition metal oxides. For forsterite and San Carlos olivine at photon energies in the range 49-59 eV, the valence bands are considerably narrower, are dominated by peaks near 4.5 and 10 eV below the Fermi level, and exhibit a lower photoelectron yield as compared with fayalite. The 8% fayalite content in the olivine appears to be too little to produce detectable $\text{Fe } 3p \rightarrow 3d$ resonance. At photon energies between 130-160 eV, the valence bands of the forsterite and olivine are dominated by a peak 2.5 eV below E_F ; however the valence band in fayalite remains broad and the 2 eV peak becomes more distinct. The separation of $\text{Fe}(3d)$ from the valence band is significant because the $3d$ levels are intimately involved in conduction processes to temperatures of about 1600K in the earth.

PHOTOEMISSION STUDY OF ELECTRONIC STRUCTURE NEAR E_F IN $\text{YBa}_2\text{Cu}_3\text{O}_x$, FOR $6.3 < x < 6.9$

R. Liu, B. W. Veal, A. P. Paulikas, J. W. Downey (Argonne National Laboratory), A. J. Arko, J. J. Joyce (Los Alamos National Laboratory)

We have performed high resolution angle resolved photoelectron spectroscopy (ARPES) studies on single crystals of $\text{YBa}_2\text{Cu}_3\text{O}_x$ with varied oxygen concentration. Since Tc monotonically decreases as the oxygen concentration x varies from 6.9 to 6.4 (becoming non-superconducting for $x < 6.4$), $\text{YBa}_2\text{Cu}_3\text{O}_x$ provides an excellent system for systematic exploration of normal state electronic structure and superconducting properties. ARPES data were acquired at the low energy range of U3C capability from samples with stoichiometries $x = 6.3$ and 6.5. A well defined Fermi function cutoff was observed for both crystals indicating normal metallic behavior (although at 6.35 the sample is non-superconducting). Band dispersions along the Γ -S and Γ -X(Y) symmetry directions were observed for both stoichiometries. The points in k-space where these dispersing bands were observed to cross the Fermi level (points on the Fermi surface) are nearly the same for the three stoichiometries $x = 6.35$, 6.65, and 6.9. This startling result is probably related to the ordering of oxygen vacancies in this system and may reflect local short range order.

CALIBRATION OF A SOFT X-RAY FILM SPECTROGRAPH*

T. J. Tanaka, N. D. Shinn, R. E. Ripple, A. Widman, and M. A. Palmer
Sandia National Laboratories, Albuquerque, NM 87185

A grazing-incidence, dispersive-grating film spectrograph used in the radiation effects testing program at Sandia National Laboratories has been quantitatively calibrated by combining tunable, monochromatic soft X-ray radiation from the LANL/Sandia beamline with laboratory measurements. The calibrated spectrograph was used in July to measure the output (50 eV < $h\nu$ < 1000 eV) from the SATURN X-ray source at Sandia.

The spectrograph calibration consisted of systematic exposures to monochromatic radiation at energetic intervals over the range of interest. The exposed film is then digitized using a microdensitometer for quantitative analysis. An example of the exposed film is shown below on the left with the corresponding digitized result on the right. For this film, exposures were made in 5 eV increments from 60 eV to 100 eV and also at 110 eV. An added benefit of these measurements is the quantification of the higher order harmonic output from the beamline itself, which appears as numerous weaker peaks at lower wavelengths in the digitized results. This information will be used to improve the beamline characterization for future experiments.

* Work supported by the United States Department of Energy under contract DE-AC04-76DP00789 with Sandia National Laboratories.

ARPES study of $\text{YBa}_2\text{Cu}_3\text{O}_{6.7}$ and $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$

B. W. Veal, A. P. Paulikas, R. Liu (ANL), A. J. Arko, J. J. Joyce (LANL)

Comparative studies were made on $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ single crystals with varied oxygen stoichiometries by using angle-resolved photoemission. Synchrotron radiation from the U3C beamline was used. Spectra were taken using the Los Alamos VSW chamber. For these measurements, the angular resolution was 2° and the energy resolution was approximately 100 meV. Samples were cleaved at about 20 K in a vacuum better than 2×10^{-10} Torr.

Shown in Fig. 1 are the valence band spectra measured on a $\text{YBa}_2\text{Cu}_3\text{O}_{6.7}$ (superconductor) and a $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$ (insulator) sample. Photons with $h\nu = 74$ eV were used. Data for the two stoichiometries are presented for three k values. We observe that the intense features at the Fermi level apparent in $\text{YBa}_2\text{Cu}_3\text{O}_{6.7}$ are lost in $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$. A sharp peak at about 1 eV binding energy near $\text{Y}(X)$ point, apparent in $\text{YBa}_2\text{Cu}_3\text{O}_{6.7}$, is severely broadened and attenuated in $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$. However, the remaining valence band features and their evolution in k-space for these two stoichiometries are very similar. It appears that, as oxygen atoms are removed from the material, the states near the Fermi energy are significantly altered so that the material converts from a metal to an insulator; but the underlying bands at higher binding energies remain relatively unchanged.

It appears that band theory nicely accounts for the ARPES spectra in the metallic region (i.e., for $7-\delta \geq 6.4$). Apparently, strong electron correlation effects responsible for the transition to insulating behavior most significantly affect those states near E_F without profoundly altering the underlying band structure.

Work at Argonne National Laboratory is supported by the U. S. DOE under contract #W-31-109-ENG-38 (BWV,APP) and by NSF, Science and Technology Center for Superconductivity, under contract #DMR 8809854 (RL).

Fig. 1

ARPES study of $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$

B. W. Veal, A. P. Paulikas, R. Liu (ANL), A. J. Arko, J. J. Joyce (LANL)

We performed angle-resolved photoemission studies on $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ single crystals using synchrotron radiation from the U3C beamline and the Los Alamos VSW chamber. Data were acquired with angular resolution of 2° and energy resolution of approximately 100 meV.

Spectral features in the vicinity of E_F were systematically studied as a function of photon energy $h\nu$ and electron emission angle to study electron energy bands in high-symmetry directions. Data were acquired in the first Brillouin zone and in higher zones. We find that the relative weight of different spectral features varies substantially under different measurement conditions. Fig. 1 shows spectra for $\text{YBa}_2\text{Cu}_3\text{O}_{6.9}$ taken along the Γ -X- Γ line at $h\nu = 74$ eV, with k outside the first zone. An intense peak, shown amplified in Fig. 2, is observed at E_F . The peak shows significant dispersive behavior that is consistent with band theory predictions and with measurements at lower photon energies. The strong features at E_F means that electron bands, possibly with a relatively high density of states, lie close to E_F along Γ -X- Γ line. Studies at $h\nu = 28$ eV also show an intense peak dispersing along Γ -X. However, at 28 eV, the intensity falls very dramatically as k approaches Γ . The varying sensitivity to different electron states afforded by the photon energy and k dependence permits selective study of specific bands for detailed comparison with theoretical predictions.

Work supported by the DOE under contract #W-31-109-ENG-38 (BWV,APP) and by NSF, Science and Technology Center for Superconductivity, under contract #DMR 8809854 (RL).

DIFFUSION IN Ti-Si MULTILAYERS PROBED BY ANOMALOUS DISPERSION IN THE SOFT X-RAY REGIME

Z. Xu, Z. Tang, S.D. Kevan, T. Novet, D.C. Johnson, University of Oregon

We have applied a soft x-ray variant of anomalous dispersion to probe interfacial diffusion and reaction in multilayered structures. We have made careful measurements of the reflectivity of a Ti-Si multilayered structure with a 100 Å period at energies in the vicinity of the Ti and Si L-edges. We observed significant peaks in first-order reflectivity at energies just below the absorption edges. A sampling of these reflectivity data is shown in the figure below.

Si-Ti Multilayer D=101Å, N=20

The increase in reflectivity just below the edges is attributed to significant Kramers-Kronig-related deviation of the real part of the refractive index from one, while the abrupt decrease at the edges is attributed to enhanced absorption and thus to sampling of a limited number of reflective interfaces in the multilayer. The peak at 98 eV (450 eV) is thus a useful probe of the spatial profile of silicon (titanium), since the optical constants of titanium (silicon) near the silicon (titanium) edge are nearly those of vacuum. We then performed temperature-dependent measurements of the reflectivity in real time, and found several interesting results. The reflectivity at the silicon edge decreases at approximately the same rate as that at the titanium edge, implying that the initial diffusion is accomplished by motion of both types of atoms. This is a difficult piece of information to attain by any other technique. Higher reflection orders were actually observed to increase upon heating. Satellite peaks in the reflectivity curves caused by the finite number of periods in our multilayer were observed to be relatively unchanged upon heating as compared to the Bragg peaks. Finally, there is evidence that the character of the diffusion is temperature-dependent. We thus plan to repeat our results at other temperatures.

Work supported by the National Science Foundation and by the American Chemical Society Petroleum Research Fund.

ARP STUDY OF THE CLEAN AND H-COVERED Pt(111) SURFACE

Wei Di, Kevin Smith, Stephen D. Kevan, Department of Physics, University of Oregon

The surface electronic structure of clean and the H-covered Pt(111) was probed at room temperature using the high resolution angle-resolved photoemission (ARP). In addition to the 2D Fermi contours of the clean surface which were reported previously, we also mapped out the surface band dispersion relations along the symmetry lines of the surface Brillouin zone (SBZ). These are plotted on a projection of the calculated bulk band structure onto the SBZ (shaded) in the figures.

All the intrinsic surface bands (upper figure) are surface resonances, with the exception of the one inside a gap about 3eV below the E_F around $\bar{K}$ point, which is a pure surface state. The band which crosses E_F at $k \sim 0.85 \text{ \AA}^{-1}$ along $\bar{\Gamma}$ and $\sim 1.00 \text{ \AA}^{-1}$ along $\bar{\Sigma}$ and the flat band just below E_F around $\bar{K}$ point are responsible, respectively, for the hexagon-like pocket around the $\bar{\Gamma}$ point and the triangular-shaped pocket at each $\bar{K}$ point of the 2D Fermi contours. Along $\bar{\Sigma}$, the clean surface bands all have even (or mixed) mirror symmetry. The extrinsic surface band (lower figure) is tentative. We did not see solid evidence for any other H-induced states or resonances.

Our results exhibit qualitative similarities to as well as some intriguing differences from previous studies. Contrary to one previous ARP study of clean Pt(111) at room temperature, our data show a large perturbation of the d-like intrinsic surface states and resonances upon hydrogen-adsorption. We thus cannot support the sub-surface chemisorption site for H at room temperature.² The scarcity of extrinsic surface features in our data might be interpreted as supporting the model which describes adsorbed hydrogen as delocalized quantum liquid.³

¹ Wei Di, Kevin E. Smith, and S.D. Kevan, Phys. Rev. B, **43**, 12062 (1991).

² W. Eberhardt, F. Greuter, and E.W. Plummer, Phys. Rev. Lett. **46**, 1085 (1981).

³ C.H. Hsu, M. El-Batanouny, K.M. Martin, J. Elect. Spect. **54/55**, 353 (1990).

Work supported by the U.S. Department of Energy, Division of Materials Science.

Fig.1 Valence band photoelectron spectra of $\text{Cu}_2\text{Pd}_{1-x}$ alloys at $h\nu = 60 \text{ eV}$

Fig.2 Valence band photoelectron spectra of $\text{Cu}_2\text{Pd}_{1-x}$ alloys at $h\nu = 120 \text{ eV}$

PARTIAL DENSITIES OF STATES OF Cu-Pd ALLOYS

S.-J. Oh, M. Han (Seoul National U., Korea), J.W. Allen, Jachoon Park (U. of Michigan) and G.K. Wertheim (AT&T Bell Laboratories)

The electronic structures of disordered alloys are of interest both technologically and from the fundamental point of view. In particular, Cu-Pd alloy systems have received wide attention lately because of the controversy as to the role of hybridization between Cu and Pd states and the effect of disorder on their electronic structure. In this work, we have measured the valence band photoelectron spectra of $\text{Cu}_2\text{Pd}_{1-x}$ alloys with various composition ($x=0.1, 0.25, 0.5, 0.75, 0.9, 0.95$) in the photon energy range $h\nu = 40$ to 250 eV . By taking advantage of the Cooper minimum in the Pd 4d photoelectron cross section around $h\nu = 120 \text{ eV}$, we are able to determine experimentally the partial densities of states from Cu and Pd atoms, and compare them with theoretical calculations.

Figures 1 and 2 show the valence band photoelectron spectra of pure Pd, $\text{Cu}_{10}\text{Pd}_{90}$, $\text{Cu}_{50}\text{Pd}_{50}$, $\text{Cu}_{75}\text{Pd}_{25}$, and $\text{Cu}_{95}\text{Pd}_5$ alloys at $h\nu = 60 \text{ eV}$ and $h\nu = 120 \text{ eV}$, respectively. At the photon energy $h\nu = 60 \text{ eV}$, the Pd 4d cross section is larger than or comparable to the Cu 3d cross section[1], whereas at $h\nu = 120 \text{ eV}$ it is smaller by a factor of 30 to 40. Hence, the spectra in Fig.2 closely represent the Cu 3d partial density of states. We can see from this figure that the position of the Cu 3d state does not move much depending on the composition, unlike the results of the coherent potential approximation calculation on these alloys[2]. On the other hand, the spectra in Fig.1 indicate that the Pd 4d state moves toward higher binding energy as the Cu concentration is increased.

[1] J.J. Yeh and I. Lindau, At. Data Nucl. Data Tables **32**, 1 (1985)

[2] H. Winter, P.J. Durham, W.M. Temmerman, and G.M. Stocks, Phys. Rev. B **33**, 2370 (1986)

VIBRATIONAL BROADENING OF BULK AND SURFACE CORE LEVELS OF Na(110)

D. M. Riffe, G. K. Wertheim, and P. H. Citrin (AT&T Bell Laboratories)

Although there have been many reports that core-level spectra of surface atoms are broader than those of the bulk, no conclusive determination of the origin of this broadening has been offered. The increased width associated with the surface atoms could be due to changes in the core-hole lifetime or the vibrational properties, or to an inhomogeneous environment. Recent work on surface atom core-level shifts of the alkali metals showed that well resolved bulk and surface spectra are readily obtained from the outermost filled p states of these metals. These states are especially attractive for detailed investigation of the vibrational contributions to the line shape, because the lifetime widths are relatively small and clean unreconstructed surfaces are readily prepared.

We have obtained such spectra for Na covering the range from 77K to the temperature where the vapor pressure reaches 10^{-10} Torr. The Na crystal was grown *in situ* on a 77K Ni(100) substrate and annealed to 200K in order to improve the LEED pattern. The data were taken as the sample warmed up to room temperature over a period of a few hours. The data were analyzed by fitting them with two spin-orbit doublets composed of Doniach-Sunjić lines convolved with Gaussians and constrained to have the same spin-orbit splitting, lifetime width, and singularity index.

The Gaussian widths obtained from this analysis, which include the instrumental contribution, exhibit a systematic increase with temperature, suggestive of phonon broadening. According to theory, in the high-temperature limit, i.e., above the Debye temperature, the phonon width increases as $T^{1/2}$. Since Gaussian widths add in quadrature it is advantageous to plot the square of the measured Gaussian width vs. temperature, as shown in the Figure. In this presentation, the intercept at $T=0$ corresponds to the temperature independent contributions, i.e., the instrumental and inhomogeneous widths, and the slope is inversely proportional to the Debye temperature. For the bulk data we find that the intercept agrees with the independently known instrumental width, indicating that inhomogeneous contributions are indeed small. For the surface data there appears to be a small inhomogeneous contribution, but the important result is the larger slope, which shows that the surface atoms have a smaller effective Debye temperature. The quantitative results obtained from this analysis are in good agreement with the analysis of surface vibrational properties due to Jackson [1]. In his model there is a soft mode perpendicular to the surface which results in a reduced effective surface Debye temperature.

[1] D. P. Jackson, Surf. Sci. 43, 431 (1974).

Work supported by AT&T Bell Laboratories.

SURFACE ATOM PROPERTIES OF Ta(110)

D. M. Riffe, G. K. Wertheim, and P. H. Citrin (AT&T Bell Laboratories)

As part of the continuing investigation of the modification of the electronic properties of metals in the surface layer, we have taken core-electron photoemission data from Ta(110). The surface was cleaned by sputtering and annealing very close to melting point, and gave no evidence of surface contamination. The data taken with 70 and 100 eV photons, and instrumental resolutions of 80 and 90 meV, respectively, are illustrated in the Figure. The solid lines, obtained by fitting the lines with the Doniach-Sunjić function, clearly show the many-body asymmetry. There are no indications of deviations from this line shape due to the modulation of the electron-hole pair density of states in this transition metal. The spin-orbit splitting is 1.912 ± 0.004 eV, and the surface-atom core-level shift appears to be about $1/3$ eV. It is certainly larger than the value of 0.28 eV obtained from less well resolved data [1], and in better agreement with the theoretical shift of 0.40 eV quoted there.

More detailed data encompassing only the $4f_{7/2}$ electrons, taken to investigate the details of the line shape, contain a number of surprises. The two salient observations are (1) that, counter to general experience, the bulk line is broader than surface line and (2) that the surface atom core-level shift is always smaller in the more surface-sensitive 70 eV spectra. Both observations are accounted for if the "bulk" line is broadened by unresolved subsurface shifts. Fitting the data with this additional degree of freedom, but with the peak intensities constrained by an escape depth, yields a surface-atom core-level shift of 357 ± 4 meV and a subsurface shift of 55 ± 10 meV. Since the near-neighbor and next near-neighbor coordination numbers of the second layer of a (110) surface are identical to those of the bulk, this result indicates that surface-atom shifts cannot be explained in terms of the first and second coordination spheres alone. However, a subsurface shift is not unexpected since layer-wise based structure calculations generally show the density of states of the second layer significantly perturbed relative to that of the bulk. The escape depth, measured in terms of the Ta(110) layer spacing, are 2.4 and 3.6 Å at kinetic energies of 48 and 78 eV above the Fermi level. These values are comparable to those of W.

The lifetime width of the bulk $7/2$ line is in good accord with an earlier analysis of the Ta(111) surface [2]. The bulk phonon widths (FWHM) at 78 and 300 K are 105 ± 5 and 163 ± 5 meV, respectively. The shape of the surface line is best represented by using a Lorentzian width much greater than that of the bulk. It is not clear whether this represents a decrease in core-hole lifetime or an anharmonicity of the surface vibrations.

[1] C. Guillou, et al., Surf. Sci. 162, 46 (1985).

[2] G. K. Wertheim, P. H. Citrin, and J. F. van der Veen, Phys. Rev. B 30, 4343 (1984).

Work supported by AT&T Bell Laboratories.

SURFACE BANDS ON Mo(001) AND Mo(001)-2H

Kevin E. Smith and S.D. Kevan, Department of Physics, University of Oregon

The (001) surfaces of tungsten and molybdenum have been the subject of numerous investigations over the past two decades due to the plethora of electronic and structural phenomena they display. Among these phenomena are temperature-induced reconstructions, adsorbate-induced reconstructions, and non-adiabatic damping of adsorbate vibrational modes. Frustrating the goal of finding universal explanations for these phenomena is the occurrence of significant differences between the behavior of Mo(001) and W(001), despite these metals being isoelectronic and having nearly identical lattice constants. We have recently reported a complete study using high-resolution angle-resolved photoemission of the surface electronic states and the corresponding surface Fermi contours for W(001).¹ The purpose of the present investigation was to produce similar results for Mo(001) in order to look for important similarities and differences between the two surfaces.

The above figure presents a summary of the surface bands for clean Mo(001) (filled circles and squares) and Mo(001)-2H (open squares). These are plotted on a projection of the even bulk bands onto the (001) surface Brillouin zone (shaded regions). Also shown are the results of a calculated surface band structure (dashed lines). Our measured bands for clean Mo(001) show similarities and differences from those for W(001). The surface bands along $\bar{\Gamma}$ and the Fermi level crossings near M are qualitatively similar, although the quantitative differences can explain the different unit cell sizes for the clean surface reconstructions. Along $\bar{\Delta}$, significant differences exist between the two clean surfaces. These are apparently not related to the reconstructions, since wave vectors along this direction are not integrally related to the reconstruction wave vector.

¹ K.E. Smith, G.S. Elliott, and S.D. Kevan, Phys. Rev. B **42**, 5385 (1990); G.S. Elliott, K.E. Smith, and S.D. Kevan, Phys. Rev. B, in press.

² G.P. Kerker, K.M. Ho, and M.L. Cohen, Phys. Rev. Lett. **40**, 1458 (1978).

Work supported by the U.S. Department of Energy, Division of Materials Science.

ELECTRON MEAN FREE PATH IN THE ALKALI METALS

G. K. Wertheim, D. M. Riffe, and P. H. Citrin (AT&T Bell Laboratories)

Photoemission spectra from clean metal surfaces in which the signal from the first atomic layer is well resolved from that of the bulk make it possible to determine the electron mean free path (escape depth), λ , in terms of the layer spacing, d , normal to the surface. This technique has major advantages compared to the conventional overlayer method, because it circumvents the problems associated with the various growth morphologies encountered in hetero-epitaxy. It also avoids the need to determine the overlayer thickness.

We have obtained core-electron photoemission spectra for the alkali metals which meet this criterion, with photon energies from close to threshold to 140 eV. The alkali films were grown on a Ni(100) substrate cooled to 78 K, using SAES Getters sources. The escape depths were determined from the areas of lines fitted to the data using a modification of Marquardt's non-linear least-squares algorithm. The lines were represented by a Gaussian convolution of a Doniach-Sunjic profile. It is readily shown that the fractional area associated with the second and deeper layers, which make up the bulk signal, is equal to $\exp(-d/\lambda)$.

In the analysis of the Li and Rb data, it was found that the surface singularity index α is larger than that of the bulk. This indicates smaller p phase shifts in the surface layer, in accord with simple expectations, but counter to the theoretical results of Kao [1]. This increased α , together with the extra phonon widths associated with the surface layer [2], makes the the fractional bulk area significantly smaller than suggested by the peak heights. The escape depths obtained for Li, Na, and Rb are compared with the theoretical values from Penn [3] in the Figure. [The oscillations in the Rb data are due to photoelectron diffraction.] The agreement with theory is entirely satisfactory for Li and Rb. The discrepancy in the case of Na is due in part to the use of a common singularity index for bulk and surface. It is interesting to note that the small escape depths obtained for kinetic energies less than 100 eV imply that angle-resolved, valence-band spectra of the alkali metals, in which bulk and surface are unresolved, are dominated by surface contributions.

[1] M. Kao, Phys. Rev. B **38**, 10915 (1988).

[2] D. M. Riffe, et al., Phys. Rev. Lett. **67**, 116 (1991).

[3] D. R. Penn, Phys. Rev. B **13**, 5248 (1976).

Work supported by AT&T Bell Laboratories.

THERMAL SHIFT OF CORE-ELECTRON BINDING ENERGIES

G. K. Wertheim, D. M. Riffe, and P. H. Citrin (AT&T Bell Laboratories)

The thermal shift of core-electron binding energies in metals was first observed in an XPS study of metallic Li and attributed to a combination of initial and final state effects [1]. In the initial state, thermal expansion reduces the band width, lowering the Fermi energy and decreasing the core-electron binding energy, but it also decreases the coulomb potential of the conduction electrons, counteracting the change in Fermi energy. In the final state, thermal expansion results in a decrease of the final state relaxation energy, increasing the core-electron binding energy. In the conventional experimental set-up there is an additional small contribution due to the thermal emf in the wire used to ground the sample. It amounts to only a few μeV per degree and can be ignored.

We have now extended this work to include four of the alkali metals, Li, Na, K, and Rb, making detailed measurements with resolution better than 100 meV over the range from 80 K to room temperature. The samples were prepared by depositing alkali vapor on a Ni(100) substrate cooled by liquid nitrogen. The spectra of the Li 1s, Na 2p, K 3p, and Rb 4p shells were recorded as the sample warmed up continuously over a period of a few hours. The line positions were determined by fitting the spectra with Gaussian broadened Doniach-Sunjić lines.

The thermally induced shift of the bulk component, measured from the binding energy at 80 K, are plotted in the Figure. The solid lines shown in the figure are the scaled linear thermal expansions of the corresponding alkali, obtained by integrating the thermal expansion coefficient from 80 K to T. For Rb the thermal expansion coefficient is known only at room temperature, but the Debye temperature of 56 K indicates that the thermal expansion will be nearly linear in the temperature range of the data. The fact that the thermal expansion curves fit the core-electron data so well, reproducing the different shapes of the data for the four alkalis, confirms that the observed effect is caused by the change in lattice constant. Quantitative evaluation of the three contributions described above, based on free electron theory, yields shift in good accord with measurements, except in the case of Li. This approach also accounts for the small shift of Mg [1] and the negative thermal shift in Al[2] as due to the larger Fermi energies of the di- and trivalent metals, which allows the negative shift due to band narrowing to overwhelm the two positive contributions. The failure in Li is tentatively attributed to self-energy effects which result in a bandwidth well below the free-electron value [3].

[1] P. H. Citrin, G. K. Wertheim, and Y. Baer, *Phys. Rev. B* **16**, 4256 (1977).[2] J. A. Tagle, E. T. Arakawa, and T. A. Callcott, *Phys. Rev. B* **21**, 4552 (1980).[3] J. E. Northrup, M. S. Hybertsen, and S. G. Louie, *Phys. Rev. Lett.* **59**, 819 (1987).

Work supported by AT&T Bell Laboratories.

L_{2,3} edges and electronic structure of iron in molecular complexesC. Cartier¹, P. Rudolf², A. M. Flank¹ and C. T. Chen²¹ L.U.R.E., Bâtiment 209d, Université Paris-Sud, F-91405 Orsay, France² AT&T Bell Laboratories, Murray Hill, NJ 07974, U.S.A.

In order to extract all the information contained in the soft x-ray absorption spectra of the Fe L_{2,3} edges of molecular compounds, one has to understand the influence of electronic and structural parameters. The fine structure of the spectra can be interpreted qualitatively in terms of a very simple orbital molecular scheme, or, more rigorously, following the "multiplet structure" model which involves the exchange interaction between the 2p hole and the 3d electrons. A way to proceed in this field is to perform a systematic investigation of different electronic and/or stereochemical parameters by measuring the Fe L_{2,3} edges of a series of well characterized model compounds synthesized by coordination chemistry methods. Such a study has been undertaken at L.U.R.E. (SA 32) and continued at the N.S.L.S., benefiting from the high resolution obtainable at the AT&T Dragon beamline.

We collected experimental evidence for the influence of the following parameters:

- symmetry: Fe(III)Cl₆, O_h compared to Fe(III)Cl₄, T_d, see figure below;
- oxidation state: Fe(III)(H₂O)₆ compared to Fe(II)(H₂O)₆, Fe(III)(CN)₆ compared to Fe(II)(CN)₆;
- ligand field: Fe(III)X₆ with X = F, Cl, O, Br (O_h symmetry), Fe(III)X₄ with X = Cl, Br (T_d symmetry);
- spin state: see preceding abstract.

Spin transition evidenced by soft X-ray absorption spectroscopy

U4B

C. Cartier¹, P. Rudolf², A. M. Flank¹ and C. T. Chen²

¹ L.U.R.E., Bâtiment 209d, Université Paris-Sud, 91405 Orsay
² AT&T Bell Labs, Murray Hill, NJ 07974

In a number of transition metal molecular compounds, the ligand field strength has an energy of the same order of magnitude as the average electron pairing energy. The metal ion is then capable of undergoing a spin state interconversion under the effect of an external perturbation such as a temperature change. Compounds that present this kind of reversible spin transition could be useful for electronic devices (switching, signal processing, data storage). The understanding of the transition mechanism is essential for the design of new compounds in the molecular electronics field.

We have performed a study of an "in situ" thermocontrolled and reversible spin transition. The $\text{Fe(II)(o-phenantroline)}_2(\text{NCS})_2$ presents a spin interconversion $12g^4e_g^2[S=2] \rightarrow 12g^6e_g^0[S=0]$ which is abrupt and complete at $T=170\text{K}$. XAS at the Fe L_{2,3} edges is a fundamental probe of the 3d empty states and is very sensitive to the spin state. As shown in the figure, we measured these absorption spectra at room temperature and at LN₂ temperature, taking care to cool down the sample very slowly in order to avoid a quench of the spin state. We checked the reversibility of the transition by recording also the spectrum after warming back up to room temperature.

Experimental L_{2,3} iron absorption spectra of $\text{Fe(II)(o-phenantroline)}_2(\text{NCS})_2$ at (A) 300K (high spin configuration) $S=2$; (B) 77K (low spin configuration) $S=0$.

BEAMLINE U4B

Electronic States and Phases of K_xC_{60}

C. T. Chen,¹ L. H. Tjeng,¹ P. Rudolf,¹ G. Meigs,¹ J. E. Rowe,¹
J. Chen,² J. P. McCauley Jr.,² A. B. Smith,² W. J. Romanow,² and E. W. Plummer²

¹AT&T Bell Laboratories, ²LRSB, U. of Pennsylvania

High resolution soft-x-ray photoemission and absorption measurements have been utilized to investigate the electronic states and phases of K_xC_{60} . [1] Well-annealed high quality K_xC_{60} thin films were grown under ultra-high vacuum conditions at the beamline. The valence band photoemission and C 1s absorption spectra of K_xC_{60} as a function of x are found to be a linear combination of data from just three phases, C_{60} , K_3C_{60} , and K_6C_{60} , giving the first conclusive spectroscopic evidence for phase separation into 3 stable phases, with K_3C_{60} being the metallic phase responsible for superconductivity. The figure on the left shows the photoemission spectra near the Fermi edge for the three phases of K_xC_{60} . A well-defined Fermi edge is clearly observed in the K_3C_{60} phase with a density of states $N(E_F)$ of 1.9 states/eV/spin/ K_3C_{60} and an occupied band width of 1.2 eV. The figure on the right shows the soft-x-ray absorption spectra near the C 1s ionization threshold for the three phases. The data show large non-rigid-band shifts between the three phases with half and complete filling, in the K_3C_{60} and K_6C_{60} phases, respectively, of the conduction band formed from the lowest unoccupied molecular orbital of C_{60} , demonstrating that the conduction band has dominantly C 2p character. The non-rigid-band shift coupled with the anomalous occupied-band width implies that there is significant mixing of the electronic states of K and C_{60} in the superconducting phase. Utilizing energies of possible mediating phonons, the directly measured $N(E_F)$ suggests that K_3C_{60} may be a BCS like superconductor. [1] C. T. Chen *et al.*, Nature, 352, 603 (1991).

BEAM LINE U4B STUDIES OF Y-Ba-Cu-O SINGLE CRYSTALS BY X-RAY ABSORPTION SPECTROSCOPY

A. Krol, Z. H. Ming, and Y. H. Kao (SUNY/Buffalo), N. Nücker, G. Roth, and J. Fink (Kernforschungszentrum Karlsruhe, Germany), G. C. Smith (BNL), A. Erhard G. Müller-Vogt (Universität Karlsruhe, Germany), J. Karpinski and E. Kaldis (ETH Zürich, Switzerland), K. Schönmann (Walter-Meißner-Institut, Germany)

The symmetry and density of unoccupied states of $\text{YBa}_2\text{Cu}_3\text{O}_{7-x}$ and $\text{YBa}_2\text{Cu}_4\text{O}_8$ have been investigated by orientation-dependent x-ray absorption spectroscopy on the O 1s edge using a bulk-sensitive fluorescence-yield-detection method. The obtained normalized spectra are shown in Fig. 1. It has been found that the O 2p holes are distributed equally between the CuO_2 planes and CuO chains and that the partial density of unoccupied O 2p states in the CuO_2 planes are identical in both systems investigated. The upper Hubbard band has been observed in the planes but not in the chains in both systems.

Fig. 1. O 1s x-ray absorption spectra for $\text{YBa}_2\text{Cu}_3\text{O}_7$ $E//a$: filled squares, $E//b$: filled triangles; difference spectrum ($E//b - E//a$): filled circles; and $\text{YBa}_2\text{Cu}_4\text{O}_8$ $E//a$: open squares, $E//b$: open triangles, difference spectrum ($E//b - E//a$): open circles.

BEAMLINE U4B Magnetic Moment in a Gd Iron Garnet Studied by Soft-X-ray Magnetic Circular Dichroism

P. Rudolf, F. Setz, L. H. Tjeng, G. Meigs, and C. T. Chen
AT&T Bell Laboratories

The magnetic moments of Gd and Fe in gadolinium iron garnet ($\text{Gd}_3\text{Fe}_5\text{O}_{12}$) were probed at 300K and 77K by soft-x-ray magnetic circular dichroism (SXMCD) measurements.[1] The figures show the Fe $L_{2,3}$ (on the left) and Gd $M_{4,5}$ (on the right) absorption and SXMCD spectra taken at these two temperatures. The Fe data contain contributions from two anti-ferromagnetically coupled Fe^{+3} components, 3 on tetrahedral and 2 on octahedral sites per formula units. The opposite sign of the Gd signal with respect to that of dominate Iron signal at the same temperature clearly indicates that Gd is anti-ferromagnetically coupled to the net Fe magnetic moment, as expected. When the sample temperature is

lowered from 300K to 77K the SXMCD signals of both elements are reversed in sign and increase by a factor of 1.5 and 4 for Fe and Gd, respectively. This reversal of the local magnetic moment is due to the crossing of the compensation point ($T_{\text{comp}}=288\text{K}$), where the Gd magnetic moment dominates over that of Fe. The data, being in quantitative agreement with an earlier calculation[2], demonstrate that SXMCD is a powerful tool for probing element and site specific magnetic moments of multi-component systems. [1] P. Rudolf *et al.*, J. Mag. Mag. Mat. (in press); L. H. Tjeng *et al.*, SPIE 1548, 160 (1991). [2] R. Pauthenet, J. Appl. Phys. 29, 253 (1958).

Giant Cu 2p Resonances in CuO Valence-Band Photoemission

L. H. Tjeng,¹ C. T. Chen,¹ J. Ghijsen,² P. Rudolf,¹ and F. Sette¹¹AT&T Bell Laboratories, ²FUNDP, LISE, Belgium

Resonant photoemission is an invaluable technique for studying the electronic structure of highly correlated electronic systems. All previous resonant photoemission studies have been carried out using photon energies close to the 3p absorption edges for 3d transition metal systems. In such measurements, the resonant enhancements are quite weak, and the 3p spin-orbit splitting is relatively small compared to the core-hole life time width, causing difficulties in data interpretation. To avoid such difficulties, we have extended this technique to 2p resonant photoemission using soft-x-rays.[1] The figure shows a giant resonant effect (up to ~100 fold increase in intensity) in the valence-band photoemission spectra of CuO with photon energies in the vicinity of the Cu 2p_{3/2} absorption edge. The data allow an unambiguous identification of the local Cu 3d⁸ and 3d⁹ L configurations (indicated in the figure) in the valence band photoemission spectrum, providing conclusive evidence for the charge-transfer nature of the insulating gap in the electronic structure of CuO. The current debate on the Mott-Hubbard-like ($U_d < \Delta$) or charge-transfer-like ($U_d > \Delta$) nature of the insulating gap is therefore successfully resolved. [1] L. H. Tjeng *et al.*, Phys. Rev. Lett. 67, 501 (1991).

Soft-X-ray Magnetic Circular Dichroism: a new technique for probing magnetic properties of magnetic surfaces and ultrathin films

L. H. Tjeng,¹ Y. U. Izderda,² P. Rudolf,¹ F. Sette,¹ and C. T. Chen¹¹AT&T Bell Laboratories, ²Navel Research Laboratory

Recent advances in synchrotron radiation instrumentation have made possible the production of high quality circularly polarized light in the soft-x-ray region (150-1500 eV).[1] Several soft-x-ray magnetic circular dichroism (SXMCD) experiments on ferromagnetic transition metals and rare earth compounds have been conducted and strong SXMCD effect were observed.[1] To extend the capability of the SXMCD technique to magnetic surfaces and ultrathin films, measurements have been carried out on MBE grown Ni ultrathin films.[2] The figure on the left shows the L_{2,3} absorption and the SXMCD spectra of a thick Ni film taken with the spin directions of the photons and that of the majority 3d electrons parallel and anti-parallel to each other. The fine structure observed in the spectra has been discussed previously.[3] The figure on the right shows the SXMCD signal at the L₃ white line from 3, 4, 5 and 6 monolayer thick Ni films on a Cu(100) substrate as a function of temperature. The data exhibit strong temperature and thickness dependence, giving film thickness dependent Curie temperature and suggesting that a single monolayer of Ni on a Cu(100) substrate may be non-magnetic. Considering the element and site specificities of core-level absorption processes, this work demonstrates that the SXMCD technique is a powerful tool for probing the magnetic properties of multi-component magnetic surfaces and ultrathin films.

[1] C. T. Chen *et al.*, Phys. Rev. B 42, 7262 (1990); F. Sette *et al.*, in *X-ray Absorption Fine Structure*, edited by S. S. Hasnain (Ellis Horwood, 1991). [2] L. H. Tjeng *et al.*, SPIE 1548, 160 (1991); L. H. Tjeng *et al.*, J. Mag. Mag. Mat. (in press). [3] C. T. Chen, N. V. Smith and F. Sette, Phys. Rev. B 43, 6785 (1991).

Soft X-Ray Absorption Spectroscopy on Ni model compounds.

J. van Elp, S. J. George, Jie Chen, and S. P. Cramer.

Lawrence Berkeley Labs and UC Davis

Soft X-ray absorption spectroscopy is a powerful new probe for studying the electronic structure of 3d transition metals in biomorganic complexes. Important Ni enzymes are CO dehydrogenase, responsible for the biological reduction of CO₂ to acetate, and Ni hydrogenases.

We have measured a number of Ni model compounds to study the influence of the local ligand-Ni structure and the significance of different ligands. In the bottom of the figure on the left we show NiF₂ in which Ni is in an O_h symmetry. This spectrum has been very well simulated using atomic multiplet structure calculations. A ligand field splitting of 10Dq=1.5 eV is found. Compared to NiF₂ the spectral shape of the Ni²⁺-(cyclam) and Ni²⁺-(mnt) is significantly different. The structure of both is planar with for mnt sulfur and for the cyclam nitrogen ligands. By descending in symmetry from O_h symmetry down to D_{4h} symmetry we hope to be able to simulate these spectra, and get values for the ligand field splitting parameters Dq, Ds and the reduction of the Slater integrals. It is however quite clear that the Ni³⁺-(mnt) and Ni³⁺-(cyclam) compounds cannot be described by using atomic multiplet calculations. The Ni³⁺ calculations show much broader structures then seen in the experimental data. In both compounds we find more spectral weight at higher absorption energy and also the entire spectrum shifts to higher absorption energy. A system in which the formal Ni valence changes gradually from Ni²⁺ to Ni³⁺ is Li_xNi_{1-x}O. Here one also finds in the Ni L edge spectra a shift of spectral weight to higher absorption energy. In this system it is however very clear that the introduced holes are mainly localized on the oxygen ligands, resulting in a mainly d⁸L (L stands for a ligand hole) ground state for the Ni³⁺ as compared to mainly d⁸ for Ni²⁺. This will also be the most likely explanation for the Ni³⁺-mnt and cyclam spectra.

Acknowledgement: We thank the profs M. Maroney and J. Kovacs for preparation of the samples. This research is supported by NIH-GM-44080-2.

PULSED INFRARED PHOTORESPONSE STUDIES OF SUPERCONDUCTING YBCO FILMS

G.L. Carr (*Grunman Corp. Res. Ctr.*), D.B. Romero and D.B. Tanner (*Univ. of Florida*), S. Elmadi, F. DeRosa, and A. Inam (*Bellcore*).

We have studied the transient voltage response of very thin YBa₂Cu₃O₇ films on LaAlO₃ substrates when exposed to pulsed infrared radiation from port U4-IR. Each film was lithographically patterned into a narrow bridge, typically 200 μm wide and 600 μm long. The bridges were cooled to about 90K, current biased, and exposed to the IR pulses. Figure 1 shows the resulting induced voltage response for a 15nm thick bridge at two slightly different temperatures. The dashed curve is the response near T_c when the bridge resistance is 70Ω. This fast signal is bolometric, similar to previous studies of films on MgO. At a slightly lower temperature (and higher bias current) the bridge resistance drops to 33Ω and the response becomes even faster (solid curve), perhaps due to an additional response mechanism. We have also measured the spectral sensitivity of the response using bandpass filters (Fig. 2). The response for R=70Ω is independent of wavelength, consistent with estimates of the film's absorption. For R=33Ω, there is enhanced response toward longer wavelengths (relative to the response in the visible). The mechanism for this apparently non-bolometric response component is unknown, but may be related to infrared-induced flux flow in the film.

Fig. 1

Fig. 2

Fig. 1 Transient voltage response of a thin YBCO film bridge to a 1ns duration, infrared pulse. The R=33Ω response (lower T, higher bias) has a steeper onset than the R=70Ω response (higher T, lower bias).

Fig. 2 Relative spectral sensitivity of the response for the same film as for Fig. 1 (normalized to visible light response = 1). The response is enhanced at longer wavelengths when R=33Ω.

Research at U. Florida is supported by DARPA through contract MDA972-88-J-1006
Research at NSLS is supported by DOE through contract DE-AC02-76CH00016

TIME-RESOLVED IR STUDIES OF PHOTO-INDUCED STATES IN HIGH- T_c SUPERCONDUCTORS

G.L. Carr (*Grainman Corp. Res. Ctr.*), D.B. Romero and D.B. Tanner (*Univ. of Florida*), S. Etemad and A. Imam (*Bellcore*), W. O'Brien (*U. Tenn.*), D. Ederer (*NIST*), C.J. Hirschmugl and G.P. Williams (*NSLS*)

We have developed a new infrared measurement technique for studying the evolution of photo-excitations on a nanosecond time scale. Excitations are produced (pumped) by a short pulse of laser light and their decay is then sensed (probed) by a pulse of far-infrared radiation from the NSLS-VUV ring. By varying the delay between the pump and probe pulses, one can map out the time dependence of the excitation. In addition, the infrared radiation is analyzed spectroscopically to simultaneously yield the frequency (energy) dependence of the excitation.

We use a mode-locked Nd:YAG laser, frequency doubled to produce 530 nm radiation, for the pump source. The mode-locking circuit uses the VUV ring's rf signal to produce 100 ps duration pulses synchronized to the pulsed VUV radiation. The laser is located at the U10 beamline, so a fiber optic cable is used to transport the light to the U4-IR platform.

One system we have chosen for study is the high- T_c superconductor $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ (YBCO). The far-infrared transmission through a superconducting film decreases toward longer wavelengths due to screening by the superfluid carriers (pairs). Photons of visible light break the pairs, reducing their density, and increasing the long wavelength transmission. The figure below shows the normalized, photo-induced change in transmission $(\Delta T/T)$ for an epitaxial YBCO film on LaAlO_3 at $T=80\text{K}$. In all cases the laser pulses cause the transmission to increase, especially at the longer wavelengths. The broken curves illustrate the effect of varying the pump-probe time delays. The (small) variation with delay time is due to the recombination of excitations into pairs, which takes place in a few nanoseconds, and is probably governed by the characteristic time for phonons to escape from the film into the substrate. The solid curve is the change in transmission for a 3K temperature rise and its similarity to the photo-induced data demonstrates that the much of the signal can be attributed to heating of the substrate.

Research at U. Florida supported by DARPA through contract MDA972-88-J-1006.

Research at NSLS is supported by DOE through contract DE-ACO2-CH00016

Potassium Enhanced Oxidation of $\text{Cu}(100)$ [†]

F. M. Hoffmann, Exxon Research and Engineering, Amundale NJ, 08801, K. Lin and R. Tobin, Michigan State University, East Lansing, MI 48824, P. Dumas, LASIR, Thiais, France, G.P. Williams and C.J. Hirschmugl, NSLS, Brookhaven National Laboratory, Upton, New York 11973.

Using Reflection-Absorption Infrared Spectroscopy (IRAS) the interaction of oxygen with a potassium multilayer adsorbed on a $\text{Cu}(100)$ surface has been studied. Vibrational spectra indicate K-O bond formation of potassium superoxide. Annealing of the layer to 600K results in the decomposition of the alkali compound. Simultaneously the formation of cuprous oxide Cu_2O , which exhibits a characteristic phonon spectrum is observed. Some understanding of these modes can be learned from the isotopic shift measurements shown below, in which ^{18}O has been substituted for ^{16}O .

The results demonstrate that potassium promotes the oxidation of Cu, increasing the rate by several (4-6) orders of magnitude.

[†]Supported by US DOE Contract #DE-ACO2-76CH00016

INFRARED TRANSMISSION OF SUPERCONDUCTING NIOBIUM FILMS

Chris Kendziora, Dave Mandrus, Dan Koller, Steve Gausepohl, and Laszlo Mihaly
Department of Physics, SUNY, Stony Brook NY 11794, Carol Hirschmugl, Gwyn
Williams NSLS, Brookhaven National Laboratory, Upton NY 11973

We investigated the infrared transmission of thin films of Niobium in the wavenumber range of 20 cm^{-1} to 100 cm^{-1} at temperatures above and below the superconducting transition. Superconductivity in Niobium is well understood, and the optical transmission is expected to follow the predictions of the BCS theory. Our study is motivated by the need for a comparison with similar experiments on the high temperature superconductors, where the interpretation of the optical measurements¹⁻³ is controversial, primarily because no firm theoretical framework is available.

The infrared transmission spectra reported here have been measured on an approximately $220 \text{ \AA}$ thick film, made by evaporation of Nb metal on zapphire substrate. The dc resistivity of the film was measured separately, and the transition temperature was found to be $T_c = 9\text{K}$, close to the bulk value, indicating the absence of significant surface scattering. Due to the high conductivity of these films, they tend to reflect nearly all radiation in the low frequency regime, necessitating the use of the high beam flux available at U4IR.

The Figure depicts the transmission, relative to the transmission at $T=15\text{K}$, at several temperatures above and below T_c . For $T > 9\text{K}$, the transmission ratio is expected to be 1, independent of frequency, since the conductivity changes slowly with temperature. To a great extent this is borne out, as evidenced by the 12K and 20K curves in the figure.

Below 9K the films begin to superconduct. In the BCS theory the transmission should decrease for frequencies within the "BCS" gap 2Δ , and exhibit a peak for frequencies above 2Δ . In the weak coupling limit $2\Delta = 3.5k_B T_c$, or approximately 22 cm^{-1} . Although the experimental data do not extend to low enough frequencies to test every aspect of the BCS predictions, the peak feature common to both the 4.5K and 7K ratios indicate the presence of superconducting gap in Niobium.

This work has been supported by the NSF grant DMR 9016456.

1. D.B. Romero, G.L. Carr, D.B. Tanner, L. Forro, D. Mandrus, L. Mihaly and G.P. Williams, *Phys Rev B* **44** 2818 (1991)
2. K. Kamaras et al., *Phys Rev Lett* **64** 84 (1991)
3. Z. Schlesinger et al., *Phys Rev B* **41** 237 (1991)

FAR-IR STUDY OF ADSORBATES ON Cu(100)

K.C. Lin and R.G. Tobin (Michigan State U.), P. Dumas (LASIR), C.J. Hirschmugl and G.P. Williams (NSLS)

We have used far-infrared reflection-absorption spectroscopy to study the low-frequency vibrations of oxygen, sulfur, formate, methanol and ethanol on Cu(100). The adsorbate-substrate vibrations of the atomic adsorbates oxygen and sulfur were not detectable at the present sensitivity limit of 0.03%, indicating that their oscillator strengths are at least a factor two smaller than estimated from electron energy loss spectra.

Oxygen induces a systematic decrease in the broadband reflectance of the sample, similar to that observed for CO on Cu(100).

This can be seen in the lower curve of fig. 1. The change can be conclusively distinguished from experimental drift by exposing the oxygen overlayer to formic acid, which decomposes, removing the oxygen and leaving a formate layer. The result is shown in the upper curve of fig. 1; the broadband reflectance change is reversed, and a band is visible at 390 cm^{-1} , which we assign to the Cu-COOH stretch vibration. In future work we plan a detailed study of this mode, including its line shape, intensity, variations with temperature and oxygen and formic acid exposure, and interactions with coadsorbed potassium. EELS measurements have suggested a reversible change in the formate orientation with temperature; we intend to investigate this issue as well.

Fig. 1. : On Cu(100) Surface :
(a) 50L O₂; (b) 50L HCOOH & 50L O₂

Condensed multilayers of methanol and ethanol at 90 K exhibited strong bands at 320 and 430 cm^{-1} , respectively. These features have not yet been identified.

1. C.J. Hirschmugl, G.P. Williams, F.M. Hoffmann and Y.J. Chabal, *Phys. Rev. Lett.* **65**, 480 (1990).

2. B.A. Sexton, *Surf. Sci.* **213**, 103 (1989).

Research supported in part by the National Science Foundation under Grant #DMR-8815616, and by the Center for Fundamental Materials Research of Michigan State University.

N. B. Brookes, Y. Chang, A. Clarke, P. D. Johnson and M. Weinert
 Physics Department, Brookhaven National Laboratory.

Spin polarized angle-resolved photoemission has been employed to study the magnetic and electronic properties of iron-silver interfaces. Epitaxial films have been grown both of iron on Ag(001) and silver on Fe(001). The latter case represents a particularly good growth system for the first few layers allowing us to observe spin-polarized interface states. These states are related to the clean iron surface electronic states¹, and exhibit distinct quantum size effects - a single new electronic state being observed for each overlayer (0-3 layers) - see below. There is also evidence for similar interface effects in the iron/Ag(001) system. The experimental results are compared with existing FLAPW calculations² and new tight binding model calculations and good general agreement is found. These results have bearing on theoretical approaches³ where the bulk iron is coupled to the overlayer through the iron surface states.

1. N. B. Brookes, A. Clarke, P. D. Johnson and M. Weinert Phys. Rev. B **41**, 2643 (1990).
2. S. Ohnishi, M. Weinert and A. J. Freeman Phys. Rev. B **30**, 36 (1984).
3. e.g. Y. Wang, P. M. Levy and J. L. Fry Phys. Rev. Lett. **65**, 2732 (1990).

This work was supported in part by the U. S. Department of Energy under Contract No. DE-AC02-76CH00016 and the National Science Foundation under Contract No. DMR-86-03304.

K.D. Moeller, Physics Department, Fairleigh Dickinson University, Teaneck New Jersey 07666, T. Sears, Chemistry Dept. BNL, H.T. Liu, USTC, Hefei, PRC, C. J. Hirschmugl, Yale University, New Haven, CT 06520, G.P. Williams, NLS, BNL, Upton, New York 11973. ^{*}Supported by US DoE Contract #DEAC02-76CH00016

A Mach-Zender interferometer was constructed using exploratory research funds. The instrument has a maximum resolution of $.0008\text{cm}^{-1}$.

The instrument is shown in schematic form in the above figure. Light is input and split on one side of the central column which moves to shorten one path while lengthening the other. The two beams recombine on the other side of the central column. A $50\text{ }\mu\text{m}$ mylar beamsplitter was used for the first work.

The figure shows the spectrum obtained from a gas cell containing ammonia, showing the far infrared rotational spectrum.

Studies of K on Fe(001) with Spin Polarized Photoemission

Y. Chang, N. B. Brookes, and P. D. Johnson
Physics Department, Brookhaven National Laboratory

As a prototypical system of alkali-metal chemisorption on magnetic substrates, potassium adsorption on Fe(001) has been studied with spin-resolved photoemission. The spin-resolved valence band spectra show significant modifications in both majority and minority electron density of states as the K coverage increases. The most interesting phenomenon is the evolution of a K-induced peak with minority character, shown in Fig. 1. At low submonolayer K coverage this minority peak grows near the Fermi level. After the K coverage reaches about 20% of the room temperature saturation coverage, it starts shifting toward higher binding energy with increasing K coverage. The work function, on the other hand, shows the general behavior of having a minimum around half a monolayer. These results may be explained by the hybridization of K 4s with the Fe 3d minority density of states highly localized at the surface.

Figure 1. Spin-resolved Photoemission spectra of a Fe(001) surface and the surface covered with 0.2ML K at room temperature. The spectra are taken at normal emission with 19.5eV photons.

SPIN-POLARIZED PHOTOEMISSION STUDY OF THE Pd/Fe(100) INTERFACE

Yeh Chang, N. B. Brookes and P. D. Johnson, Physics, BNL

Currently there is strong interest in the magnetism of thin films and superlattices. Magnetic properties of these structures can be different from those of bulk materials, and interesting magnetic coupling between thin layers of transition metals has been discovered. A good understanding of the electronic properties at interfaces is essential for an understanding of the magnetism. Palladium is of particular interest because it is similar to Ni but is nonmagnetic in bulk. A Pd film grown on Fe(100) has been reported to have an expanded lattice(5%) and to be magnetic at the interface. An interesting inverted magnetic exchange splitting have been reported for the Pd/Fe(110) interfaces.

We have performed a spin-polarized angle-resolved photoemission study of the Pd/Fe(100) interface. At thin Pd coverage several interface features are observed. Figure 1. shows the spin resolved valence band photoemission spectra of 1ML Pd on Fe(100), taken at normal emission. The spectra in the lower panel are taken with mainly p-polarized 58eV photons, and the spectra in the upper panel are taken with more s-polarized photons of the same energy. Features at around 0.4eV, 1.7eV, and 3.7eV binding energy(B.E.) are not from the Fe substrate, and their intensities attenuate as the Pd layer is grown thicker. It is interesting to note that the features at around 1.7eV B.E. seem to exhibit an inverted exchange splitting, as the majority peak has lower B.E. than the minority peak. However the photon polarization dependence shows that this is not the case. The minority peak shows strong enhancement with s-polarized light, while the majority peak does not. This difference indicates that they are derived from bands of different symmetry, and therefore can not be considered as an exchange-split pair. The spin-split partner of the minority peak is more likely to be the shoulder on the higher binding energy side of the majority peak. We have performed Tight Binding calculations for this interface. Good agreement between experimental results and calculations have been obtained, showing similar interface features and their energy dispersion. This work was supported in part by the U.S. Department of Energy Contract No. DE-AC02-76CH00016 and National Science Foundation Materials Research Group Contract No. DMR-86-03304.

Fig. 1

of different symmetry, and therefore can not be considered as an exchange-split pair. The spin-split partner of the minority peak is more likely to be the shoulder on the higher binding energy side of the majority peak. We have performed Tight Binding calculations for this interface. Good agreement between experimental results and calculations have been obtained, showing similar interface features and their energy dispersion. This work was supported in part by the U.S. Department of Energy Contract No. DE-AC02-76CH00016 and National Science Foundation Materials Research Group Contract No. DMR-86-03304.

1. Z. Celinski, B. Heinrich, J. F. Cochran, W. B. Muir, A. S. Arrott, J. Kirschner, Phys. Rev. Lett., **65**, 1156(1990).
2. W. Weber, D. A. Wesner, G. Guntherodt, and U. Linke, Phys. Rev. Lett. **66**, 942(1991).

G.A. Mulhollan, R.L. Fink, J.L. Erskine(u. of Texas) and G.K. Walters(Rice U.)

The temperature dependence of the valence band energy distribution curves of ultra-thin Fe₂W(100) films has been measured using spin- and angle-resolved photoemission. Majority and minority features at $\bar{\Gamma}$ are stationary with increasing temperature; apparent peak broadening is caused by the emergence of extraordinary peaks. Comparison with a cluster calculation suggests that short range ferromagnetic order $\sim 4\text{\AA}$ above T_c in the 1.5 monolayer film.

Figure 1. Spin resolved EDCs for (a) majority: $\blacksquare$ $T=0.35T_c$, $\square$ $T=0.9T_c$ and $\bullet$ $T=1.05T_c$ and (b) minority: $\blacksquare$ $T=0.35T_c$, $\square$ $T=0.9T_c$ and $\bullet$ $T=1.05T_c$ spins of 1.5ML at $\bar{\Gamma}$. The lines labelled A,B, and C mark the locations of the centroids of the Fe minority, W Δ_5 and the Fe majority peaks respectively.

Figure 2. EDC for $\bullet$ $T=1.05T_c$ and $\circ$ 50-50% mixture of 0.35Tc majority and minority spin-resolved EDCs.

This research was supported by DMR-8906935(MRG) and DMR-8922359.

X-RAY MASK DISTORTION DUE TO RADIATION DAMAGE

R. E. Acosta
IBM Research Division
T.J. Watson Research Center
Yorktown Heights, NY 10598 USA

During their lifetime X-ray masks will receive exceedingly large accumulated doses, on the order of 500,000 J/cm². Because the allowed mask distortion for 0.25 μm ground rules is of the order of 0.075 μm , the permissible radiation-induced distortion is $\approx 0.025 \mu\text{m}$, thus requiring substrate materials with negligible distortion when subjected to large radiation doses. The radiation hardness of various candidate membrane materials was determined by mapping the position of fiducials using a metrology tool (Nikon 2i and Nikon 3i) at approximately 5,000 J/cm² (incident dose) increments. Irradiations were performed in helium at 20 Torr pressure, with the membranes separated by a 40 μm gap from a backing aluminum plate, to insure that the membrane was well heat-sunked. To accentuate the effect of the distortion, only one half of the membrane was irradiated. Because this method gives a "worst case" estimate, for comparison some of the Si/Polyimide structures were irradiated full field. The membranes were irradiated at beam line U-6 of the NSLS at Brookhaven National Laboratory and beamlines ES-0 and ES-1 of the Aladdin ring at the University of Wisconsin. The results shown below are given, except where noted, in terms of the maximum distortion (at the boundary irradiated/non-irradiated).

ACKNOWLEDGMENTS

The nature of this work required the invaluable contributions of very many individuals. In particular, those of R. Fair, B. Hill, W. Waldo (Motorola Corp.), H. Voelker, J. Buccignano, M. Hansen and F. Cerrina (U. of Wisconsin), J. Silverman, R. Scott, and R. Viswanathan. For many useful discussions, I thank A. D. Wilson, J. R. Maldonado, A. Wagner, R. Fuentes, S.-L. Chiu, and W. A. Johnson (Motorola Corp.).

D. Boyne, L. Hsia, R. Wachnik, R. Decker, and C. Wasik
IBM Corporation-East Fishkill, Hopewell Junction, New York 12533

The radiation damage and reliability of bipolar transistors have been studied. The device parameters were measured before and after irradiation and were measured again after the devices were annealed. The study shows that X-ray radiation degrades the common emitter current gain. However, the degradation can be completely removed by annealing. No reliability problems associated with irradiation have been found. Fig. 1 shows the common emitter current gain before and after an irradiation of 1500mJ/cm², and after annealing in forming gas for 30 min. The gain drops sharply at lower collector current. At a collector current of 5 uA the current gain drops about 50% after irradiation. Upon annealing the gain recovers. To determine whether exposure to X ray reduces device reliability, the transistors were stressed at 120C for 1000 h. with the emitter-base junction reversely biased. As shown in Fig. 2, it is not possible to distinguish between the irradiated and non-irradiated samples. It can be concluded that exposure to X ray does not impose any additional reliability problems for the bipolar devices employed in this investigation.

FIG. 1. Common emitter current gain before and after irradiation with 1500 mJ/cm² and after annealing.

FIG. 2. Percent shift of common emitter current gain as a function of stress time for a collector current of 10 uA.

Gregory J. Dunn, (M.I.T. Lincoln Laboratory)

X-ray lithography is a candidate for fabrication of VLSI circuits with feature sizes substantially below 0.5 μm . However, x-ray doses needed for lithography produce radiation damage in the gate oxide of MOSFETs. Positive charge, neutral electron traps, and interface traps are generated in large numbers. The interface traps and positive charge can be eliminated by a low temperature anneal in a hydrogen ambient, but the neutral electron traps do not completely anneal out. Higher temperature anneals are not possible after metallization due to the problem of aluminum penetration into the silicon junctions. Therefore, radiation damage incurred during the later steps in MOSFET fabrication is of great concern. The neutral electron traps remaining after the post-metallization anneal have been shown to contribute to device degradation due to channel hot carrier stress, particularly for p-channel devices and for n-channel devices stressed at $V_g \geq V_d$.

Previously we have shown that reoxidized nitrided oxides (RNO) exhibit substantially reduced positive charge generation and virtually no interface trap generation after exposure to ionizing radiation, such as would be encountered in space or weapons applications. We have now completed a study of process-induced radiation damage, and in particular the creation of neutral electron traps, in these dielectrics.

Finished n-channel transistors were irradiated by L.C. Hsia of IBM Corporation using the vacuum ultraviolet storage ring at the National Synchrotron Light Source. The doses of 500 and 1500 mJ/cm² simulated typical exposures anticipated for lithography using medium to low sensitivity x-ray resists. After irradiation, the samples were annealed at 400°C for 30 min in forming gas, simulating the final post-metal anneal. Channel hot carrier stress tests were then performed under conditions which maximize electron injection. The resulting degradation in transconductance was substantially less for RNO than for oxide. The data show that the lifetime of the RNO devices without irradiation is greater than that of the oxide devices, and that x-ray exposure produces less degradation of lifetime in RNO than in oxide, especially at the lower dose.

In conclusion, nitridation and reoxidation produce a gate dielectric which is more resistant than conventional oxide to creation of residual electron traps by synchrotron x-ray lithography. This improved resistance will lead to longer integrated circuit lifetime in future, very short channel devices.

*This work was sponsored by the Department of the Air Force.

X-RAY MASK REPAIR

A. Wagner, J. P. Levin, J. L. Mauer, P. G. Blauner, S. J. Kirch,
P. Lougo, and J. P. Silverman

IBM

T. J. Watson Research Center
Yorktown Heights, NY 10598

Focused ion beams are currently used to repair defects in X-ray masks. An image of the defective region on the mask is obtained using the ion beam in a manner analogous to a scanning electron microscope. Opaque defects are removed by physical sputtering of extra absorber. Clear defects (missing absorber) are repaired by ion beam induced decomposition of an organometallic compound to form an opaque film on the substrate. The X-ray opacity of ion beam deposited gold was measured. A series of gold patches of varying thickness were deposited on an X-ray mask. The mask was then exposed to three different X-ray doses using the National Synchrotron Light Source VUV ring at Brookhaven National Laboratory. The ratio of the energy transmitted through each gold patch to the incident X-ray energy was measured using dosimeter film. Figure 1 shows the transmitted X-ray energy as a function of the thickness of the gold patch. For films less than approximately 2 μm thick, the transmitted energy decreases exponentially with thickness as expected. The measured X-ray absorption coefficient was approximately 18,000 to 20,000 per cm which is roughly half the value measured for pure, bulk gold. Thus, repairs made with ion beam deposited gold must be approximately twice as thick as the electroplated gold to achieve the same X-ray opacity. The reduced X-ray absorption coefficient of ion deposited gold can be attributed to the large carbon content (approx 50 at. %) and resultant low density of gold in the films.

FIG. 1. X-ray transmission vs thickness of ion-beam-deposited gold. The X-ray absorption coefficient of the ion-beam-deposited gold is approximately one half the measured value for bulk gold.

THE CONDENSATION AND PHOTON DECOMPOSITION OF DIBORANE

M. W. Ruckman (BNL), J. Mowlem and D. Strongin (SUNY-Stony Brook).

Boron forms electron deficient compounds which have important uses in chemical synthesis. Unlike carbon, boron does not have enough valence electrons to make all the pairwise covalent chemical bonds with the atoms needed to achieve a closed octet. This produces an interesting hydrogen mediated chemical bond in diborane (B_2H_6) which is shown in the accompanying sketch. There are a series of boron hydrides that form polyhedral structures of increasing size that resemble the fullerenes and these structures can be formed by pyrolysis of diborane and the condensation of the resulting vapor. There is a lack of electron spectroscopic and near edge data for these interesting compounds and this work is an attempt to study boron hydrides in the solid state.

We obtained near-edge data for B_2H_6 condensed on a silver surface from a 5 % B_2H_6 mixture in argon liquid nitrogen temperature. We are aware of only one gas phase EELS result for B_2H_6 [1] which shows that diborane has a large π^* resonance at 190 eV and smaller features at 194 and 196 eV. Our data differs from the published gas phase data and has a principle peak at 192 eV, unresolved features near 195 eV and a large shape resonance near 202 eV. The difference between our result for B_2H_6 is and the gas phase data is puzzling since the gas phase result is in better agreement with theory. A possible explanation is that B_2H_6 was spontaneously converted to another boron hydride on a silver surface. This might be similar in concept to the conversion of benzene into a phenol species on Pt, but a problem with this idea is that the process continues past completion coverage.

The condensed diborane was subjected to photon irradiation to decompose it and simulate the thermal process known to produce a series of increasingly more complex boron hydrides and boron. The strong near-edge features of diborane are reduced after a 5 min. irradiation in the white beam. The B k-edge develops a sharp excitonic feature near 195 eV which is also seen when pure boron is examined using EELS or NEXAFS.

[1] T. M. Zimkina and A. S. Vinogradov, Bull. Acad. Sci. USSR Phys. Ser. 36, 229 (1972).

Research sponsored by the Division of Materials Sciences of the Office of Basic Energy Sciences, U. S. Department of Energy, under Contract Nos. DE-AC02-76CH00016.

Banana-shaped charge-clouds of the three-center bonds in B_2H_6 .

PHOTON-ASSISTED FORMATION OF DIAMOND-LIKE CARBON FILMS FROM ORGANICS

M. W. Ruckman (BNL), J. Mowlem and D. Strongin (SUNY-Stony Brook).

In the late 1970s, it was discovered that diamond and diamond like carbon films could be deposited from glow discharges onto heated substrates. Subsequent work established that sp^3 bonded CH species generated in the low pressure discharges were responsible for diamond formation. Diamond and diamond-like carbon films have been deposited under a number of conditions and have been successfully deposited on room temperature substrates in plasmas containing halogens.

We have studied the use of photon beams to react molecular solids with surfaces (i.e. the nitrification of GaAs(100) using NH_3 [1]) and promote the reaction of molecular solid mixtures to make refractory thin film coatings. The success of the above work, led us to examine the possibility that soft x-rays or photoelectrons could create the chemical species needed to grow hard carbon films instead of graphite.

The effect of white synchrotron radiation on a selection of solid organics was studied using NEXAFS and photoemission. The compounds included an aromatic (p-xylene), a ketone (acetone), carboxylic acid (acetic acid), an alcohol (CH_3OH), and an aldehyde (CH_3COH).

Near-edge results are shown for acetal and p-xylene. In both cases, the photon treatment breaks the molecule and produces a non-volatile carbon film. For acetal and the other methyl containing organics, the carbon film formed lacks the sharp π and σ K-edge features that exist for diamond and graphite. There are pre-edge features corresponding to CH bonding. For p-xylene, the near-edge has a closer resemblance to graphite. We believe the carbon films produced by the photon irradiation of acetal and similar organics are diamond-like or hard carbon films [2]. The formation of graphitic carbon is more likely in the aromatics because of the initial electronic state of the carbon is not sp^3 . In a substance like acetal, atomic oxygen, hydrogen or CO radicals are generated and can play a role in removing hydrogen from the methyl species which also serves to promote diamond-like carbon formation.

[1] M. W. Ruckman et al., Appl. Phys. Lett., 59, 849 (1991)
[2] D. Wesner et al., Phys. Rev. B 28, 2152 (1983).

Research sponsored by the Division of Materials Sciences of the Office of Basic Energy Sciences, U. S. Department of Energy, under Contract No. DE-AC02-76CH00016.

Figure 1

Figure 2

REACTION OF POTASSIUM WITH AMMONIA AT LIQUID NITROGEN TEMPERATURES

M. W. Ruckman, E. Horache, and Myron Strongin (BNL)*

Near-edge and photoemission techniques were used to characterize the chemical behavior of potassium/ NH_3 systems formed either by the deposition of ammonia on a thick potassium film or by the deposition of potassium vapor into an ammonia film at liquid nitrogen temperature. Earlier studies of cesium in NH_3 suggest that a hexamine or related compound forms at liquid nitrogen temperature and warming this phase to about 100 K causes the cesium atom to become solvated. The cesium hexamine-like phase has distinctive valence band features. For K on NH_3 (Fig. 1) these valence features are absent and the near-edge peaks identified the NH_3 Rydberg states (Fig. 2) become less intense. In addition, the ammonia shape resonance almost disappears when K is added. It does not do so when Cs is added. For NH_3 on potassium (not shown), we see the same result at low NH_3 coverages but at large NH_3 coverages the valence band begins to resemble that seen during the earlier experiments with cesium at much lower NH_3 coverages. At this time, we attribute the results for K/ NH_3 as indicating that potassium does not form the hexamine phase at liquid nitrogen temperature except at low potassium concentrations when NH_3 is deposited on potassium.

* Work supported by the U. S. Dept. of Energy, Division of Materials Science, Contract No. DE-AC02-76CH00016.

PHOTON-ASSISTED REACTION OF DIBORANE AND AMMONIA TO FORM BORON NITRIDE

D. Strongin, J. Mowlem (SUNY-Stony Brook), M. W. Ruckman and Myron Strongin (BNL).

Many useful materials like BN can be produced by the decomposition of gases over hot surfaces. Chemical vapor deposition (CVD) is not a practical means of depositing materials like BN on compound semiconductors like GaAs because it is not possible to heat the substrate hot enough to decompose gases like B_2H_6 and NH_3 or a precursor like borazene. This limitation can be overcome, in part, by using photon or plasma-assisted decomposition of the gaseous precursors. Unfortunately, the energy of the incident species is still high enough to damage the substrate.

In earlier work by Gao et al. [1], solid O_2 was used as a reactant in a reactive deposition process to grow an aluminum oxide on GaAs(100). In this work, the feasibility of using solid molecular precursors in a photon-assisted chemical deposition process to produce a refractory BN film was examined. B_2H_6 and NH_3 were co-condensed on a silver film at 78 K and irradiated with a white synchrotron beam. After the film was reacted at liquid nitrogen temperature, it was heated to room temperature to remove volatile reaction products and to test its thermal stability. Near-edge and photoemission spectra were used to determine the composition of the resulting films.

The accompanying figure shows B K-edge data which is sensitive to the changing chemical composition of the thin film. When diborane and ammonia are co-condensed, diborane disproportionates and the complex $(NH_3)_2BH_2BH_4$ forms [2]. The photons cause this precursor to break up into volatile components like hydrogen and H_2BNH_2 . Molecular BN also forms and contributes to the strong π^* pre-edge feature at the B K-edge. Our conclusion that BN is formed is supported by the appearance of the valence and core level photoemission spectra. The N K-edge is not as sharp as that obtained by NEXAFS for hexagonal BN [3] and this suggests significant atomic level disorder. This disorder can be partially removed by further exposure to the white beam at 300 K.

[1] Y. Gao et al., J. Appl. Phys. **67**, 7148 (1990).

[2] K. Wade, Electron Deficient Compounds, (Thomas Nelson & Sons, London, 1971) p 74.

[3] B. M. Davies et al., Phys. Rev. B **24**, 3537 (1981).

Research sponsored by the Division of Materials Sciences of the Office of Basic Energy Sciences, U. S. Department of Energy, under Contract Nos. DE-AC02-76CH00016.

VALENCE STATES IN THIN Eu-Rh AND Eu-Pd FILMS

C. Berg and S. Raaen (U. of Trondheim, Norway), and M.W. Ruckman (BNL)

Europium 4f photoemission as well as CFS measurements in the vicinity of the Eu 4d absorption threshold, have been performed on Eu on Rh and Eu on Pd overlay systems. The Eu 4f emission was dramatically enhanced by tuning the photon energy to the giant resonance energy of ~141 eV. Eu was evaporated at room temperature onto vacuum deposited substrates of Rh and Pd. Mixed interface films that extended over many atomic lattice spacings were found to form at room temperature. Eu 4f emission showed that only divalent Eu ions were present at the surfaces, whereas less surface sensitive CFS measurements indicated the presence of trivalent Eu ions below the surfaces in the Pd case. Annealing of the samples at ~500 °C resulted in an increased fraction of Eu 3+ ions. The Eu ions at the surface were always divalent. Trivalent Eu ions are likely to occupy sites where the local environments are similar to those in $EuRh_2$ and $EuPd_3$ for the Rh and Pd systems, respectively. The mixed interfaces that were not annealed, are argued to be amorphous in nature, since they form at temperatures that are well below those where nucleation and crystalline growth are expected to occur.

Fig.1
CFS spectra of the Eu/Pd interface system. About 30 Å Eu was deposited on a Pd substrate. The solid line is the as deposited sample, and the dashed line is a sample that had been annealed at ~500 °C for one minute. The feature near 152 eV photon energy indicates trivalent Eu.

This work was supported by the Norwegian Research Council (NAVF)

PHOTON MODIFICATION OF ALUMINUM OXIDE ON 6063 ALUMINUM

T.S. Chou (NSLS), M.W. Ruckman (Physics Dept.)

Photon modification of surface oxides generated on 6063 aluminum by different surface treatments was studied using photoemission at the U7B beam line. The surface of 6063 aluminum with mild alkaline detergent (Oakite) is covered by a mostly amorphous oxide. A further in situ oxygen glow discharge at 20ma for 30 minutes at room temperature did not produce the crystalline oxide, but removed surface impurities like carbon. When the surface is exposed to white synchrotron radiation the surface is modified. For the aluminum cleaned with Oakite the aluminum 2p binding energy decreases from 76.19 eV down to 75.34 eV for 1 Amp-Hr white light radiation, and for the glow discharge cleaned sample white light radiation causes the aluminum 2p core level to move from -76 eV to -75.34 eV. The sample subjected to the oxygen glow discharge treatment and white light radiation was subsequently exposed to moist air and given a cumulative exposure of 2.3×10^{19} Langmuirs. The Al 2p binding energy shifts from 75.15 eV to 74.77 eV under the same photon dosage. Fig. 1(A) shows the results for Aluminum treated with Oakite only. Fig. 1(B) for Aluminum treated with in situ oxygen glow discharge. Fig. 1(C) shows the results of Aluminum exposed to air after the end of Fig. 1(B) experiment. In Fig. 1 the solid line is the UPS spectrum without photon radiation, the dotted line is with 0.8 Amp-Hr of synchrotron radiation, and the broken line is with 1 Amp-Hr photon dosage. The results are consistent with oxygen loss via photo-desorption and with increment of surface oxide disorder.

This work was supported by the U.S. Dept. of Energy under contract PE-AC01-CH00076.

PHOTOELECTRON SPECTROSCOPIC STUDIES OF THE ELECTRONIC STRUCTURE OF FERROUS SULFIDE

P. M. Jones, J. A. Guckert, J. A. May, and E. I. Solomon* (Stanford University)

Variable photon energy experiments have been performed on tetrahedral high spin d^6 Ba_2FeS_3 to probe the bonding and effects of ionization on their electronic structure. Due to photoionization cross section behavior, relative intensity changes of valence band photoemission peaks through a photon energy range of 25-120 eV allow definitive assignment of the low binding energy bands. Consistent with the normal bonding description of transition metal compounds, the valence band PE spectra indicate that the highest occupied antibonding levels (Peak 1, Figure 1a) consist primarily of metal character in the initial state, while the deeper binding energy bonding levels are predominantly ligand in character (Peak 3, Figure 1a). CIS resonance studies through the iron $3p \rightarrow 3d$ absorption edge, however, show dominant resonance enhancement in the deeper binding energy bonding peak (Figure 1b), indicating that in the ionized final state this bonding peak becomes primarily metal in character. Further evidence of electronic relaxation is found in the Fe3d shake-up satellite intensity observed at photon energies away from resonance. Thus significant changes in the electronic structure are observed upon ionization of high spin d^6 FeS_4^{6-} as a result of the large increase in the exchange interaction on going to a high spin d^5 final state. These orbital wavefunction changes are also being investigated using spin polarized SCF-X α -SW electronic structure calculations.

Figure 1. (a) Valence band PE spectra of Ba_2FeS_3 collected at photon energies of 30, 40, 60, 80, and 110 eV; (b) intensity-energy profiles of valence peaks compared to the iron $3p \rightarrow 3d$ CFS absorption edge.

Financial support for this work was provided by the NSF (Grant # NSF-891968704841) and the DOE (Grant # AC03 82ER-13000). We would like to thank NSLS and SSRL for providing beam time for these experiments.

Polarization dependence of X-ray absorption in O_2 : the square well model.

Pieter Kuiper and M.W. Ruckman, Physics Department, BNL

We have studied the oxygen K-edge of O_2 on graphite (HOPG) at 25 K. At high coverage the molecular (z) axes are perpendicular to the graphite (xy) plane¹, so that at normal incidence of the X-rays only transitions from $O\ 1s$ to $np\ x_{||}$ are possible. The $2p_{x_{||}}$ atomic orbitals form the partly occupied π^* (antibonding) molecular orbital, giving rise to the first absorption peak at 531 eV in the π spectrum shown in the figure below.

The $2p_z$ atomic orbitals form a σ^* molecular orbital. This orbital can be reached when at grazing incidence the electric field vector is parallel to z . Langhoff and coworkers² have proposed a simple model for estimating the energies of σ orbitals in diatomic molecules. They show that these orbitals are very similar to the wave functions of a particle in a cylindrical well three times as long as the internuclear distance. The energies of sigma orbitals are then given by the square well formula $E(n) = (\pi^2 \hbar^2 / 8m)(n^2 / l^2) - V_0$ where V_0 is the calculated binding energy of the lowest sigma valence level. Counting gives $n=4$ for the lowest unoccupied sigma level. This model explains nicely why in the solid this orbital is at higher energy, closer to the vacuum level than in the gas phase: the neighbouring molecules reduce the effective width of the well. The resonance due to $n=5$ can also be observed. This model shows why the standard EXAFS model fails for O_2 .³ The π spectrum has a strong feature at 565 eV, that may be due to a double ionization, and extended fine structure that is probably due to scattering by neighbouring molecules in hexagonal coordination.

¹ M.F. Toney and S.C. Fain, Phys. Rev. B 36, 1248 (1987), and references therein.

² J.A. Sheehy, T.J. Gil, C.L. Winstead, R.E. Farren, P.W. Langhoff, J. Chem. Phys. 91, 1796 (1989).

³ B.X. Yang, J. Kirz, and T.K. Sham, J. de Physique 47, C8-585 (1986).

Supported by US Department of Energy under contract DE-AC02-76CH00016.

Spin-resolved XAS of O_2 .

Pieter Kuiper and M.W. Ruckman, Physics Department, BNL

Oxygen is magnetic: droplets of liquid oxygen are attracted to a magnet and a flame burns higher in a strong magnetic field. The magnetic moment is due to two spin-aligned electrons in two degenerate antibonding π^* molecular orbitals, forming a triplet state ($S=1$). In the X-ray absorption spectrum peak A is due to transitions of a spin-down electron from $1s$ to the π^* orbital. But transitions to all other unoccupied levels occur for up- and down-spins. These final states will have different energies because of the exchange interaction. Recent calculations show that the exchange splitting of the σ^* peak is only 0.5 eV. The calculated σ^* -multiplet coincides with peak B.¹

We have used the Auger spectrum to differentiate between these two spin states. Let 'up' be the spin direction of the π^* electrons. If the excitation to the σ^* -orbital involves a spin up core-electron, the π^* electrons can take part in deexcitation, and the relaxation of these least-bound electrons will produce high energy Auger electrons. But if a spin-down core-electron is excited, the π^* -electrons cannot fall down into the core, and Auger-electrons with less kinetic energy will be emitted. We measured the Auger spectra around 540 eV, and saw indeed differences in the high-kinetic energy part. In the figure the total yield (line), the yield of high-energy electrons (up triangles), and the yield of lower energy Auger electrons (down triangles). Indeed we see that the spin-down σ^* excitation is about 0.5 eV higher than the spin-up state. This shows that Auger electrons can be used to measure spin-resolved XAS, even in anti-ferromagnets. The much larger difference at peak C is due the energy shift between the one-hole and the two-hole Auger-spectrum.²

¹ M.W. Ruckman, J. Chen, S.L. Qiu, P. Kuiper, M. Strongin, and B.I. Dumlup, PRL (in press).

² J. Chen, C.L. Lin, S.L. Qiu, M. Strongin, and M.L. denBoer, J. Vac. Sci. Techn. (A) 8, 2591 (1990).

Supported by US Department of Energy under contract DE-AC02-76CH00016.

VALENCE DENSITY OF STATES OF HYDROGEN-REDUCED NiO(100) M. A. Langell, K. W. Wulker and B. P. Hearty (Univ. of Nebraska-Lincoln).

Slow hydrogen reduction of NiO(100) has been studied by photoemission, LEED and Auger spectroscopy. Under 1.3×10^{-2} Pa H₂ and 625 K, NiO gradually loses oxygen in a three dimensional process that can be modeled kinetically by competitive surface reduction coupled to diffusion of subsurface oxygen to the surface reduction front [1]. The LEED retains good quality NiO(100) structure for a considerable time, although an eventual epitaxy of Ni(100) can be observed along with NiO(100) in the LEED.

Photoemission results were obtained for $h\nu = 40\text{--}120$ eV as a function of surface reduction, with Auger O 510/ Ni 848 ratios taken as a measure of oxygen content. The accompanying figure samples the data at 50 eV. Stoichiometric NiO(100) yields valence band spectra comparable to previously published results [2]. As the surface reduction proceeds, intensity from reduced nickel states gradually builds up in the band gap in approximate correlation with the decrease in Auger ratios. After 3 hr (O/Ni = 0.98), the reduction enters a near steady state, with O/Ni ratio and photoelectron spectra changing only very gradually. At ~ 11 hrs. (O/Ni = 0.88), Ni(100) epitaxy is observed and while Auger ratios continue their gradual decrease, the photoelectron spectra again change substantially. We are presently attempting to correlate changes in the PES data with reduced Ni cluster size.

- [1] J. J. Scholtz and M. A. Langell, Surf. Sci. 164, 543 (1985).
[2] J. M. McKay and V. E. Henrich, Phys. Rev. Lett. 53, 2343 (1984).

Research sponsored by the National Science Foundation (# CHEM- 9117069) and NLSL Faculty-Student Support Program (United States Dept. of Energy Contract No. DE-AC02-76CH00016).

Fig. 1. Experimental band structure (circles) of strained fcc body-centered-tetragonal Cu along the $\Gamma\Delta X$ line. The numbers on top of the two panels refer to final-state energies. Left: Cu film grown on Pd(001); Right: Cu film grown on Pt(001). The solid curves are calculated for bulk (unstrained) fcc Cu.

† Present address: The BCC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.
; On leave from: Department of Physics, Peking University, The People's Republic of China.

We are grateful to the National Science Foundation and to the Department of Energy for partial support of this work with Grants DMR-8709021 and DE-FG02-86ER45239, respectively. One of us (S.C.W.) is also indebted to the Science and Technology Commission of the People's Republic of China for partial support through Grant No. 19074004.

ELECTRONIC PROPERTIES OF BODY-CENTERED-TETRAAGONAL COPPER

H. Li, S.C. Wu, J. Quinn, Y.S. Li, D. Tian and F. Jona (SCUY Stony Brook)

Angle-resolved photoemission with synchrotron light is used to determine the band structure of 3- and 6-layer films of Cu grown epitaxially on Pd(001) and on Pt(001). The films have body-centered-tetragonal structure resulting from the large plane strains of 8 to 9% imposed by the substrates. The band structure is the same, within experimental error, for films grown either on Pd(001) or Pt(001), and is similar to that of stable fcc Cu.

Fig. 1. Experimental band structure (circles) of strained fcc body-centered-tetragonal Cu along the $\Gamma\Delta X$ line. The numbers on top of the two panels refer to final-state energies. Left: Cu film grown on Pd(001); Right: Cu film grown on Pt(001). The solid curves are calculated for bulk (unstrained) fcc Cu.

† Present address: The BCC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.
; On leave from: Department of Physics, Peking University, The People's Republic of China.

We are grateful to the National Science Foundation and to the Department of Energy for partial support of this work with Grants DMR-8709021 and DE-FG02-86ER45239, respectively. One of us (S.C.W.) is also indebted to the Science and Technology Commission of the People's Republic of China for partial support through Grant No. 19074004.

STRUCTURAL AND ELECTRONIC PROPERTIES OF ULTRATHIN FILMS OF Gd, Tb, Dy, Ho and Er

H. Li†, D. Tian, J. Quinn, Y.S. Li, S.C. Wu and F. Jona (SUNY Stony Brook)

Single-crystal ultra-thin films of Gd, Tb, Dy, Ho and Er were grown epitaxially on W(110) is ultra-high vacuum and studied by both LEED and UPS. Comparison with bulk single crystal data showed, at least in the case of Tb, that while the purity of the films may be higher, the crystallinity, i.e., the long-range order, is less well developed in the films than in bulk single-crystals. Electron distribution curves from Gd, Dy, Er and Tb are shown in the figure (the Ho data were not representative of pure material owing to the presence of a strong oxygen-derived peak at about -6 eV). All curves exhibit a peak at approximately -0.5 eV, which is associated with a surface state located in a band gap of the projected band structure at $\bar{\Gamma}$ (see also Barrett and Jordan, Z. Phys. B66, 375 (1987) for Y(0001); Wu et al., Phys. Rev. B in press for Tb(0001); and Li et al., J. Magn. Magn. Mater. in press for Gd(0001)).

Fig. 1. Normal-emission electron distribution curves from single-crystal thin films of Gd, Dy, Er and Tb measured with photon energies $h\nu$ and (75% - 25%)-polarized light. The zero of energy is taken at the Fermi energy.

†Present address: The BOC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.

‡On leave from the Department of Physics, Peking University, The People's Republic of China.

This work was supported by the Department of Energy with Grant No. DE-FC02-86ER45239.

ULTRA-THIN FILMS OF Rh ON Ag(001) AND Rh ON Ag(001): GROWTH MODE AND MAGNETISM

H. Li†, S.C. Wu†, D. Tian, Y.S. Li, J. Quinn and F. Jona (SUNY Stony Brook)

Deposition of minute amounts of Rh on a clean and reconstructed Au(001) surface destroys the reconstruction and yields a 1×1 structure. Prolonged deposition produces epitaxial but not pseudomorphic films of Rh with the bulk equilibrium fcc structure. Deposition of Rh on a clean Ag(001) surface produces overlayers which are partially covered by or intermixed with Ag, but which are pseudomorphic with the Ag(001) substrate. Photoemission experiments in the very early stages of overlayer growth reveal a 4.1 ± 0.3 eV splitting of the $4s$ levels of Rh, which is not observed in thick Rh films. The $4s$ splitting is explained by the presence of a magnetic moment on the Rh atoms and is related to recent theoretical predictions of ferromagnetism in a monolayer of Rh on either Ag(001) or Au(001) [Zhu et al., Phys. Rev. B 43, 4007 (1991); Eriksson et al., Phys. Rev. Lett. 66, 1350 (1991)].

Fig. 1. Top: Rh $4s$ levels for increasing Rh coverage, given on the left in layer-equivalents (LE). The experimental data (black dots, after background subtraction) were fitted with two Gaussians (dashed curves). The splitting is obvious at low Rh coverages (bottom three panels), but is not visible in thick Rh films, which produce curves indistinguishable from those of single crystal Rh (top panel). Photon energy 165 eV. Bottom: Rh $4s$ levels for increasing Rh (background not subtracted). Photon energy 110 eV.

†Present address: The BOC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.

‡On leave from: Department of Physics, Peking University, The People's Republic of China.

We are grateful to the National Science Foundation and to the Department of Energy for partial support of this work with Grants DMR-8921123 and DE-FC02-86ER45239, respectively. One of us (S.C.W.) is also indebted to the Science and Technology Commission of the People's Republic of China for partial support through Grant No. 19074004.

CORE LEVEL LINESHAPES IN FE-V ALLOYS

S.L. Qiu and R.G. Jordan (Florida Atlantic U.), M.W. Ruckman and M. Strongin (BNL)

Some while ago it was suggested that the spin-splitting of the s core levels in magnetic $3d$ transition metals might be used as a probe of the local moment in metals, alloys and compounds. The spin-splitting results from an exchange coupling between the two different spin orientations of the s core level relative to that of the $3d$ spin. It has been shown theoretically that, in principle, one can measure the local moment by investigating the lineshapes of photoelectron spectra and, apparently, there have been a number of successful applications of this approach in the case of the Fe $3s$ core-level splitting using XPS. However, a recent detailed study and compilation of results on a wide range of Fe-based alloys and inorganic compounds revealed that there appears to be a poor correlation between the measured Fe $3s$ spin-splitting and the local moments on the Fe-site. In order to help clarify this dilemma we carried out a preliminary investigation of the photoemission from the Fe $3s$ core levels from some Fe-V alloys using XPS. We chose this particular system because there are both experimental and theoretical data that indicate that the local moment on the Fe-site decreases in a regular way with V concentration, and disappearing at $\sim 77\%$ V. Our measurements indicated quite clearly that the profiles of the spectra change with V concentration and we concluded that there is increasingly less weight in the higher binding energy regions of the spectra as the local moment on the Fe-site decreases. However, due to an inferior count-rate and correspondingly poor resolution we were not able to identify whether the changes in profiles were due to a reduction in spin-splitting or the intensity of the higher binding energy component.

In an attempt to obtain data at improved resolution we carried out a series of photoemission measurements from three Fe-V alloy samples using the facilities on U7B. We investigated samples of pure Fe and V, Fe-20% V (disordered bcc), FeV (CsCl-structure) and Fe-70% V (disordered bcc) at room temperature. *In-situ* cleaning was carried out using Ar^+ bombardment and annealing at 320°C for 30 minutes. Previous work had shown that after cleaning there was a small excess of V in the surface region.

We collected data from several core levels of both Fe and V. Below we show the results for the Fe $3s$ core levels (left) and for the Fe $3p$ core levels (right) taken at a photon energy of 200 eV. (For convenience, we have removed the background, aligned the peak positions and normalized the heights. Both the Fe $3s$ and $3p$ core levels shift to increased binding energy with increasing V concentration.) It is clear that the lineshapes change as the V concentration increases. We also observe changes to the lineshapes of the V core-levels. One explanation is that the spin-splitting is reduced as the V concentration increases, i.e., as the local moment decreases. However, before we can be certain that this is the cause we will investigate the changes in more detail by fitting lineshapes. This analysis is currently underway.

This work was carried out as part of proposal #91-U-116. SLQ and RGJ are grateful to the NSLS faculty research support program for financial assistance and Florida State University for the allocation of supercomputer resources.

PARTIAL DENSITIES OF STATES IN CuAuI

S.L. Qiu and R.G. Jordan (Florida Atlantic U.)

M.W. Ruckman and M. Strongin (BNL)

We have carried out a series of photoemission measurements from a polycrystalline sample of (ordered) CuAuI over the photon energy range 20 – 160 eV. Our aim is to determine the Cu- and Au- contributions to the density of states by exploiting the strong photon energy dependence of the relative photoemission cross sections from the Cu $3d$ - and Au $5d$ -states. According to data tables, for example, the cross section of the Au $5d$ -states decreases by at least two orders of magnitude over the photon energy range of our measurements. In principle, therefore, it should be possible to extract the individual contributions to the density of states.

The sample was cleaned *in situ* by repeated Ar^+ bombardment and annealing cycles. Previous studies have shown that the ordered phase (CuAuI) can be readily obtained by annealing at 350°C for 1 hour. We show in figure 1 a selection of spectra taken at room temperature. Our analysis of the measurements is not complete but, using earlier work on Cu_3Au as a guide, we can make the following preliminary observations. At 160 eV the experimental photoemission spectrum is dominated by the contribution from the Cu-states but at about 40 eV the Au-states dominate. Thus, the measurements at 160 eV should reflect closely the Cu $3d$ partial density of states and those in the range 40 – 50 eV should provide a reasonable picture of the Au $5d$ partial density of states, with a small contribution from the Cu $3d$ states near a binding energy of 2.5 eV.

We have calculated the site decomposed densities of states in CuAuI (Li_0 structure) using the SCF-LMTO-ASA method. In figures 2(a) and (b) we show the results after they have been cut-off at E_F and smoothed using a Lorentzian function with an energy dependent FWHM (of 0.5 eV at E_F and 1 eV at the bottom of the Au $5d$ -bands). The latter function is taken to represent a combination of the instrumental resolution and lifetime effects.

A direct comparison between the data in figures 1 and 2 is not entirely appropriate. For instance, in the photon energy of our measurements the emission is predominantly from the *surface* region, (where previous work has shown there is an excess of Au), but the calculations were made for the (uniform) *bulk*; also the effect of the electron-photon matrix elements should be taken into account. Nevertheless, the development of the photon-energy dependent features and their general shape are in accord with the calculations. We are carrying out a more quantitative analysis, involving comparisons with similar measurements on Cu and Au and with angle-resolved XPS of CuAuI . Calculations will also be made of the Cu and Au contributions to the photocurrent that take into account the electron-photon matrix elements.

This work was carried out as part of proposal #91-U-116. SLQ and RGJ are grateful to the NSLS faculty research support program for financial assistance and Florida State University for the allocation of supercomputer resources.

ATOMIC AND ELECTRONIC STRUCTURE OF THIN FILMS OF Mn ON Pd(111)

D. Tian, H. Li, S.C. Wu, F. Jona (SUNY Stony Brook) and P.M. Marcus (IBM Yorktown)

Epitaxial films of Mn can be grown on Pd(111) to thicknesses of the order of 15 to 20 layers. Vacuum deposition on a room-temperature substrate produces a 3-fold symmetric 1x1 phase with in-plane lattice constant $a=2.75$ Å (pseudomorphic to Pd(111)) and bulk interlayer spacing $d_{111}=2.16$ Å. This structure probably corresponds to strained magnetic α -Mn. Vacuum deposition on a 150°C substrate produces a 6-fold symmetric $\sqrt{3} \times \sqrt{3}$ 30° strained magnetic α -Mn. This structure probably produces a bulk phase with in-plane lattice constant $a=4.76$ Å. Both the atomic and the electronic structure of the $\sqrt{3} \times \sqrt{3}$ 30° phase are different from those of the 1x1 phase, but the magnetic moments of the Mn atoms are essentially the same in the two structures (see Fig. 1) and larger than in α -Mn.

Fig. 3. Top: Angle-resolved normal-emission electron distribution curves measured with s-polarized 90-eV photons on a 2.1LE film of Pd(111)-1x1-Mn and on a 12-LE film of Pd(111)- $\sqrt{3} \times \sqrt{3}$ 30°-Mn. Bottom: 3s core levels for both films mentioned above.

† Present address: The BOC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.

‡ On leave from: Department of Physics, Peking University, The People's Republic of China.

We are grateful to the National Science Foundation and to the Department of Energy for partial support of this work with Grants DMR-892123 and DE-FG02-86ER45239, respectively. One of us (S.C.W.) is also indebted to the National Natural Science Foundation of the People's Republic of China for partial support through Grant No. 19074004.

RE-EXAMINATION OF THE ELECTRON BAND STRUCTURE OF Tb ALONG Γ -A

S.C. Wu, H. Li, Y.S. Li, D. Tian, J. Quinn, F. Jona (SUNY Stony Brook), D. Fort (C. Birmingham), and N.E. Christensen (Aarhus C.)

The electron band structure of Tb metal has been re-examined by means of angle-resolved photoemission experiments on Tb(0001) samples free of Fe impurities in the surface region. Comparison of the new data with those collected from an Fe-contaminated sample Wu *et al.*, Phys. Rev. B 41, 11913 (1990) shows that the presence of Fe in the surface region caused a shift of the top portion of the $6s$ $3d_{5/2}$ -type Δ_1 band toward larger binding energies by approximately 1 eV. The energy position of the top of the Δ_1 band (E_F) is -2.6 eV. The agreement with the result of the relativistic linear muffin-tin-orbital calculation reported earlier ($E_F = -2.3$ eV) is markedly improved.

Fig. 1. Electron band structure of Tb along the Γ -A line. Dashed curves: theory. Solid and dash-dotted curves: experiment with different choice of the position of the bottom of the band.

† On leave from: Department of Physics, Peking University, The People's Republic of China.

‡ Present address: The BOC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.

We acknowledge partial support of this work by the Department of Energy with Grant No. DE-FG02-86ER45239. One of us (S.C.W.) is also indebted to the Science and Technology Commission of the People's Republic of China for partial support through Grant No. 915702.

RELATIVISTIC EFFECTS IN THE ELECTRON BAND STRUCTURE OF Cu(111)

S.C. Wang, H. Li, Y.S. Li, D. Tian and F. Jona (SUNY Stony Brook)

Angle-resolved photoemission from Cu(111) with both s - and $25^\circ p$ -polarized light shows evidence for the lifting of degeneracies of the two occupied Λ_3 bands. Spin-orbit coupling leads to splitting of the non-relativistic upper Λ_3 band into Λ_3^6 and Λ_3^5 bands in the middle portion of the Γ -AL line, and of the non-relativistic lower Λ_3 band into Λ_3^3 and Λ_3^4 bands near the Γ point along the Γ -AL line. The critical points for the relativistic bands are measured to be $\Gamma_6 = -2.90 \pm 0.05$ eV, $\Gamma_5 = -3.45 \pm 0.10$ eV and $\Gamma_8 = -3.65 \pm 0.05$ eV, in good agreement with the results of self-consistent relativistic band-structure calculations done by Eickardt, Fritzsche and Noffke (EFN) with a Slater-type X_α potential and $\alpha = 0.70$.

Fig. 1. Band structure of Cu along the Γ -AL line. Dashed and solid curves represent non-relativistic and relativistic bands as calculated by Janak, Williams and Moruzzi [Phys. Rev. B 11, 1522 (1975)] and by Eickardt, Fritzsche and Noffke [J. Phys. F 14, 97 (1984)] respectively. Dots and crosses are experimental data obtained in this work as described in the text; the associated numbers refer to the photon energies used to measure the electron-distribution curves E_{em} which the experimental data were obtained. The final energy above the Fermi level E_F is given by the scale on top of the figure. The open circles are experimental results of Knapp, Himpel and Eastman [Phys. Rev. B 19, 4952 (1979)]. The arrows at the edges indicate critical points as calculated by EFN with a Slater-type X_α potential ($\alpha = 0.70$)—the solid curves were calculated by EFN with an Hedin-Lunqvist exchange-correlation potential. Note that the ordinate scale is different above and below the Fermi level E_F .

†On leave from: Department of Physics, Peking University, The People's Republic of China.
‡Present address: The BOC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.

We are grateful to the Department of Energy for partial support of this work with Grant DE-FG02-86ER45239, respectively. One of us (S.C.W.) is also indebted to the Science and Technology Commission of the People's Republic of China for partial support through Grant No. 9187002.

SURFACE STATE ON CLEAN Tb(0001)

S.C. Wang, H. Li, Y.S. Li, D. Tian, J. Quinn and F. Jona (SUNY Stony Brook)

Angle-resolved photoemission experiments on clean Tb(0001) reveal the existence of a state 0.5 eV below the Fermi level which has the properties of a surface state. It exhibits no dispersion with respect to the momentum component perpendicular to the surface in the photon-energy range between 15 and 55 eV; it is located in a band gap of the projected band structure at Γ ; and it is very sensitive to oxygen contamination of the surface. The wave function of this surface state has Λ_1 symmetry, as the intensity of the photoemission peak decreases when the p -component of the exciting radiation is reduced. Both the very weak energy dispersion with respect to the parallel component $k_{\parallel}$ of the momentum and the photon-energy dependence of the photo-ionization cross section indicate that the surface state has $5d_{3/2}$ orbital character. The existence of this surface state can be explained by the same theory of surface core-level shifts that is related to the narrowing of the surface density of states in transition metals.

Fig. 1. Angle-resolved electron distribution curves measured on clean Tb(0001) at normal emission with $(75^\circ \pm 25^\circ)$ p -polarized radiation.

†On leave from the Department of Physics, Peking University, The People's Republic of China.
‡Present address: The BOC Group, Technical Center, Murray Hill, New Jersey 07974, U.S.A.

We acknowledge partial support of this work by the Department of Energy with Grant No. DE-FG02-86ER45239. One of us (S.C.W.) is also indebted to the Science and Technology Commission of the People's Republic of China for partial support through Grant No. 9187002.

CHLORINE ADSORPTION ON ROOM TEMPERATURE GaAs(110)

David K. Shuh and Jory A. Yarnoff (UC Riverside Dept. of Physics)

The elucidation of the mechanisms responsible for the chemical etching of semiconductor is of considerable importance as technology is applied to the etching of III-V semiconductor materials. Of particular interest are the interactions of Cl_2 with GaAs, since chlorine is the primary etchant employed in GaAs device fabrication. One of the most useful techniques for investigating halogen chemisorption is soft x-ray photoemission spectroscopy (XPS). We have investigated the reaction at high chlorine exposure levels, and find that the chemistry involves not only both As-Cl and Ga-Cl bonding, but a rich variety of higher chloride formation as well.

GaAs(110) wafer substrates were subjected to large exposures of Cl_2 gas. The resulting interfaces were examined by surface sensitive synchrotron XPS of the Ga 3d and As 3d core levels. All of the work was performed at the NSLS VUV-8 beamline utilizing an angle-integrating ellipsoidal mirror analyzer. The measured Ga and As core levels were numerically fit to a sum of Gaussian-broadened Lorentzian lineshapes in order to deduce to the chemical states of Ga and As in the surface region.

The Ga and As core levels both show clear evidence of bonding to Cl. As the Cl_2 exposure is increased, we observe the formation of both As and Ga monochlorides as well as the higher chlorides, AsCl_2 and GaCl_2 . The distribution of arsenic chlorides favor the monochloride, while the distribution of gallium chlorides favor the formation of GaCl_2 . These species are the reaction intermediates which are formed on the surface by the etching process. At the highest exposures studied, a large build up of GaCl_3 is observed on the surface, which may inhibit further etching of the GaAs material. The conversion of GaCl_2 to the volatile Ga chloride species, GaCl_3 , is thus the limiting factor in the etching reaction at room temperature.

Figure 1. Surface sensitive photoemission from the As 3d and Ga 3d core levels to those measured after exposure to chlorine gas.

UNFILLED ORBITALS OF C_{60} AND C_{70} FROM CARBON K-SHELL X-RAY ABSORPTION FINE STRUCTURE

L. J. Terminello (Lawrence Livermore), D. K. Shuh (U. of CA, Riverside), F. J. Himpsel and D. A. Lapiano-Smith (IBM Res., Yorktown), J. Stöhr, D. S. Bethune, and G. Meijer (IBM Res., Almaden)

The K-shell near edge x-ray absorption fine structure spectra of solid C_{60} and C_{70} have been recorded with high spectral resolution. The spectra exhibit sharp resonances associated with transitions to unfilled π^* molecular orbitals and, at higher energy, broader resonances associated with σ^* -like orbitals. The observed energy splitting of the π^* resonances can be accounted for by published extended Hückel calculations. Our data clearly reveal that the fullerene films resemble molecular solids with considerably stronger intra-molecular than inter-molecular interactions.

Photoelectron Spectroscopy Study of the Cr/MoS₂ Interface

U-8B

T.D. Durbin, D.K. Shuh and J.A. Yarnoff
Department of Physics, University of California, Riverside, CA 92521
J.R. Lince and S.V. Didzulis
The Aerospace Corporation, P.O. Box 92957, Los Angeles CA 90009

The interface between metals and the MoS₂ (0001) basal plane surface exhibits interesting surface chemistry in that the highly anisotropic MoS₂ crystal structure has been found to be relatively inert to chemical reactions with metallic adsorbates when compared to metal adsorption on other semiconductors. We have studied the Cr/MoS₂ interface using core level and valence band soft x-ray photoelectron spectroscopy in order to better understand the chemistry at the metal/MoS₂ interface and to determine ways to maximize the adhesion of MoS₂ based solid lubricants to stainless (i.e. Cr containing) steel surfaces.

Our results show that vapor deposited Cr forms a relatively flat overlayer which is primarily composed of metallic Cr although some sulfur is incorporated in the film. Core level spectroscopy of the Mo 3d and S 2p, shown in the figures, reveals that Cr partially reacts with MoS₂ and forms metallic Mo and a chromium sulfide species at the metal/MoS₂ interface. Using variable photon energies to sample the Cr overlayer at different depths it was also found that the topmost portion of the Cr film is enriched with sulfur. The reactivity of Cr with respect to MoS₂ was found to be intermediate between that of Fe and Mn when these results are compared to the results of previous metal/MoS₂ interface studies.

Annealing the Cr covered sample to progressively higher temperatures results in additional chemical reaction between Cr and the MoS₂ substrate. The interaction caused by annealing the sample to 425°C involves the uncovering Mo in a metallic state and the formation of another S-containing species. Annealing the sample further results in the formation of several new S containing species.

Photoemission Studies of the Chemisorption of Dichlorosilane on the Si(111) and Si(100) Surfaces

U-8B

T.D. Durbin, D.K. Shuh and J.A. Yarnoff
Department of Physics, University of California, Riverside CA 92521
D.A. Lapiano-Smith, F.J. Himpel and F.R. McFeely
IBM T.J. Watson Research Center, Yorktown Heights, NY 10598

Dichlorosilane has been the subject of recent interest due to its applicability in atomic layer epitaxial growth of silicon films. Recently, Nishizawa *et al.* demonstrated the growth of epitaxial layers of silicon using alternating exposures of SiCl₂H₂ and H₂.¹ We have studied the chemisorption and subsequent reaction of SiCl₂H₂ on Si(111) and (100) surfaces at both the low pressures and temperatures typical of a surface science approach and the higher pressures and temperatures employed for film growth.

In this study, core level photoemission of the Si 2p level was used to characterize the silicon surface after exposure to SiCl₂H₂. We found that SiCl₂H₂ chemisorbs dissociatively resulting in the formation of adsorbed silicon monochloride. The coverage of monochloride increases as the temperature of the substrate during deposition is increased, reaching a maximum at ~700°C. The increased Cl coverage observed at higher temperatures correlates with the thermal desorption of H₂ from the surface, which leaves more sites available for Cl₂ adsorption.² We also found that the exposure to SiCl₂H₂ resulted in larger chlorine coverages on Si(100) as compared to Si(111) at all temperatures studied. The results of an 8 kL dose of SiCl₂H₂ taken with the sample at room temperature and 600°C are shown in Fig. 1 for both the Si(111) and Si(100) surfaces.

I.J. Nishizawa, A. Aoki and S. Suzuki, J. Crystal Growth 89, 502 (1990).
P.A. Coon, P. Gupta, M.L. Wise and S.M. George, to be published

The Decomposition of BCl_3 on $\text{Si}(111)$ and the Subsequent Incorporation of Boron into Silicon

D.A. Lapiano-Smith and F.R. McFeely, IBM Research Division, Thomas J. Watson Research Center, Yorktown Heights, N.Y. 10598

A series of photoemission experiments were conducted in order to probe the sequence of steps whereby boron atoms are incorporated into silicon via the reaction between BCl_3 and $\text{Si}(111)$. The reaction between BCl_3 and Si is complex due to the presence of Si, B and Cl in a number of chemically distinct environments simultaneously. Row A (rA) in the figure below shows the Cl 2p, Si 2p and B 1s core levels following an initial exposure to 1000 L BCl_3 which results in saturation coverage. The Cl 2p has been fit with two spin-orbit split doublets, separated by 1.1 eV which have been assigned to B-bound to Cl and Si-bound to Cl components. The Si 2p shows a shoulder arising from the Si-Cl component, which is shifted by 0.9 eV from the bulk peak (69.1 eV kinetic energy) and the filling up of the region at 68.3 eV is attributed to a Si-B component which is shifted 0.4 eV from the bulk. The B 1s data show a sharp peak at 106 eV kinetic energy assigned to elemental B bonding to Si and the broad band may be fit with the assumption that it arises from BCl_2 and BCl_3 moieties. An anneal of the sample to 570°C (row B) effectively desorbs all molecularly chemisorbed BCl_3 . Practically all of the Cl is desorbed from the surface following an anneal to 710°C (row C). Both the Si 2p and B 1s data indicates that the amount of Si-bound-B has increased. Annealing to 870°C (row D) completes the reaction process. Cl photoemission is reduced to a trace. The Si-B features continue to grow showing the gradual incorporation of boron into the silicon with successive high temperatures anneals.

A Mechanism for Diborane Induced Selectivity Loss in the CVD of Silicon from SiH_2Cl_2

D.A. Lapiano-Smith and F.R. McFeely, IBM Research Division, Thomas J. Watson Research Center, Yorktown Heights, N.Y. 10598

Soft x-ray photoemission measurements were acquired to illustrate a mechanism whereby the introduction of B_2H_6 into a SiH_2Cl_2 CVD reaction mixture for in situ doping can have a dramatic effect, namely, the loss of selectivity of SiH_2Cl_2 to SiO_2 . These selectivity loss results from the thermally activated, autocatalytic dissociative chemisorption of B_2H_6 on the SiO_2 . Due to the autocatalytic nature of the process, the oxide surfaces are, in contrast to clean SiO_2 , quite reactive to SiH_2Cl_2 .

Panels a and b in the figure below show the Si 2p and Cl 2p lines resulting from the exposure of a boron covered oxide surface to 300L SiH_2Cl_2 at room temperature. The Si 2p level is broad and shifted to higher binding energy by about 2 eV wrt the elemental position. This indicates that Si is bound to two Cl atoms. The Cl 2p spin orbit doublet is poorly resolved, indicative of both B and Si bound moieties. After annealing the surface to 550°C, not only does the Si peak shift to the elemental position (panel c) but its intensity also decreases by about a factor of 2. The Cl 2p peak (panel d) sharpens due to the decrease of the contribution from Si-bound Cl, however, the total intensity is conserved indicative of the quantitative transfer of chlorine atoms to the boron substrate.

STUDIES OF THE XeF_2 ETCHING OF Si (111) AT 200°C

C.W. Lo, D.K. Shuh and J.A. Yarnoff

Department of Physics, University of California, Riverside, CA 92521 USA

In order to understand the basic phenomena involved in the technological processing of silicon, there have been many fundamental investigations of the interaction of atomic fluorine with silicon. A large number of these investigations have used XeF_2 as a source of atomic fluorine, due to the ease with which XeF_2 can be employed. It has been shown, however, that XeF_2 has a different behavior than atomic fluorine, particularly at elevated temperature. There is a minimum in the rate at which XeF_2 etches Si at approximately 100°C, while atomic fluorine displays Arrhenius behavior.¹ A possible cause of the minimum in the XeF_2 etch rate is that there are different mechanisms involved at low and high temperatures.

In this study, the surface chemical and geometric structure of a silicon sample is investigated after the sample had been exposed to XeF_2 at room temperature (RT) and at 200°C. These two temperatures are on each side of the etching rate minimum, and have a similar etch rate. Also, mass spectrometry has shown that the gas phase etch products evolved from the surface are independent of the temperature in this regime.² In the present work, however, we show that there is a difference in structure of the reaction layer which forms on the surface as a result of etching.

Si (111) samples were exposed to sufficient amounts of XeF_2 to achieve steady state etching at both RT and 200°C. The sample which was exposed at RT was subsequently annealed to 200°C in order to induce a similar amount of thermal desorption. Soft x-ray photoemission spectroscopy (SXPS) and photon stimulated desorption (PSD) spectra were collected from each of these surfaces. The SXPS results, shown in the figure, and the PSD results both indicate that although the reaction layers contain similar concentrations of fluorine, they have quite different structures. This demonstrates that the difference in etching at RT and 200°C is due to a difference in the reaction mechanisms.

¹D.L. Flamm, D.E. Ibbotson, J.A. Mucha and V.E. Donnelly, Solid State Technol. 4 (April, 1983) 117.

²H.F. Winters and I.C. Plumb, J. Vac. Sci. Technol. B9 (1991) 197.

Photoemission Intensity (Arb. Units)

CONSTRUCTION AND LAUNCH OF AN IMPROVED SOFT X-RAY TELESCOPE

E. Spiller and J. Wilczynski (IBM Research), L. Golub and G. Nystrom (Ctr. for Astrophysics)

The main limitation on the resolution (about 0.8 arcsec) of the telescope launched in Sept. 89 was photon shot noise, due to the small photon flux available during the short rocket flight. A new telescope was built with the goal to improve the throughput, mainly by improving the mirror reflectivity and filter transmission. Polishing of the boundaries of the 140-layer Co-C coating with an ion beam was used to increase the mirror reflectivity. X-ray transmission of the light blocking filter was improved by replacing the parylene films in the previous design by thin films of phthalocyanine (replace the carbon-parylene structure with a carbon-phthalocyanine structure). The additional light absorption of the phthalocyanine films permits to use thinner films with enhanced x-ray transmission.

The new telescope was launched on Feb. 22, 1991 and produced spectacular x-ray images of the corona of the sun. The exposures indicate an increase of the photon flux by about a factor 3. Resolution in the images is around 1 arcsec; due to a break of some of the pre-filters during launch, the photos with the longest exposure times are useless (visible light leak) and we could not realize the expected improvement in resolution. A new launch date has been set for July 11, 1991.

Ion beam polished mirrors as used for the telescope where tested at beamline U8.

Fig. 1. The soft x-ray corona of the sun on Feb. 22, 1991. Photographed from a NASA sounding rocket with a normal incidence x-ray telescope.

U9B
STRUCTURAL ANALYSIS OF PHAGE T7 GENE 5.9 PRODUCT, USING CIRCULAR
DICHROISM SPECTROSCOPY AND PRIMARY SEQUENCE ANALYSES.
L.L. France, Li Lin, Bill Studier and J.C. Sutherland.
Biology Department, Brookhaven National Laboratory, Upton, NY.

Phage T7 gene 5.9 product has been identified as an inhibitor of the RecBCD protein in *E. coli*, and evidently acts to protect the T7 phage DNA from degradation by RecBCD. Circular dichroism (CD) spectra and primary sequence analysis of the protein are currently being conducted to predict the quantity and locations of secondary structures. The results indicate that this protein (which consists of 51 amino acid residues) contains predominantly α -helix ($\approx 73\%$), which occurs as two helical regions (residues 8 - 22, and 29 - 47), interrupted by a β -turn (at residues 25-29). Helical wheel projections support these conclusions, and further indicate that both helical regions are amphipathic.

The CD spectra of T7 gene 5.9 product in 10 mM phosphate buffer (pH 7.0) were measured in the presence (—) and absence (---) of 150 mM NaCl in order to determine the effect of ionic strength on this protein. In both cases the protein concentration was 10 mg/ml. The second purpose of this experiment was to compare the CD spectrum of this protein under conditions currently being used to measure its NMR spectrum (100 - 150 mM NaCl), with those generally used for CD (no NaCl). These spectra are compared in Fig.1, and indicate that the presence of 150 mM NaCl causes no significant structural changes in the protein.

Research supported by the Office of Health and Environmental Research, United States Department of Energy, and a grant from the National Institutes of Health (GM34662).

U9B
STRUCTURAL ANALYSIS OF VARIABLE MEMBRANE PROTEIN 7 FROM *BORRELIA HERMSII*, USING CIRCULAR DICHROISM SPECTROSCOPY.
L.L. France, J.J. Dunn and J.C. Sutherland.
Biology Department, Brookhaven National Laboratory, Upton, NY.

Borrelia hermsii is the etiological agent of relapsing fever, which is marked by periodic, alternating episodes of illness and remission of symptoms. One of the more abundant outer membrane proteins of *B. hermsii*, called "Variable membrane protein" (Vmp) undergoes antigenic variation within the host, allowing the spirochete to avoid the host's immune response and causing a resurgence of symptoms. Several serotypes of Vmp are being cloned and expressed in *E. coli*, by Dr. John Dunn (Biology Department, Brookhaven National Laboratory). We have measured the circular dichroism (CD) spectrum of Vmp serotype 7 in the UV range (Fig.1) in order to predict the secondary structural content of the protein (Table 1). The research goal for this project is to characterize and compare the structures and properties of the different Vmp serotypes and related outer surface proteins from the Lyme disease spirochete (*Borrelia burgdorferi*), in order to investigate the structural basis of inter- and intra-species antigenic variation.

Fig.1: CD spectrum of Vmp7 in 10 mM sodium phosphate buffer (pH 6.0).

Table 1: Secondary structure prediction of Vmp7 from *B. Hermsii* from analysis of the CD spectrum (260 nm - 178 nm).

	α -helix	β_s -sheet	β_p -sheet	β -Turn	Other	Total
fraction	0.63	0.02	0.00	0.27	0.10	1.02
st.dev.	± 0.01	± 0.02	± 0.00	± 0.01	± 0.02	

Research supported by the Office of Health and Environmental Research, United States Department of Energy, and a grant from the National Institutes of Health (GM34662).

A VACUUM UV CD STUDY OF PYROPHOSPHATASE WILD TYPE AND MUTANTS
M. Gulotta, D.J. Goss (Hunter College, CUNY), and B. Cooperman
(Univ. Pennsylvania)

Pyrophosphatase (PPase) is an essential enzyme for both prokaryotic and eucaryotic systems. Its function is to catalyze the high energy hydrolysis reaction:

Cooperman's laboratory has looked at the enzymatic activity and the hydrophobicity of the wild type enzyme. His group has done site specific mutagenesis on the enzyme and has found that single amino acid changes have resulted in drastic decreases in the enzymatic activity and sometimes changes in the hydrophobicity of the enzyme as well. To show that the mutated amino acid is part of the binding site, we need to show that the mutation hasn't altered the conformation of the protein.

E. coli PPase is a symmetrical enzyme with 6 identical lobes of approximately 100 amino acids which makes it a good candidate for vacuum UV CD studies.

We have used the vacuum UV circular dichroism spectrometer which uses the beam as its light source to monitor the conformation of E. coli pyrophosphatase in its wild type and mutant forms. This set up is preferable to what is available commercially because we can obtain clear spectra down to 176 nm on an aqueous sample and the thinness of the beam allows us to use very little sample. Any differences in the CD spectrum of the mutant versus that of the wild type indicate that the mutation has caused a structural change and therefore there is insufficient evidence to prove that this enzyme is part of the binding site.

The spectrum of the wild type PPase (solid line) and that of the DE97 mutant where an aspartic acid was replaced with a glutamic acid (dashed line) are shown below. No substantial difference in is seen between these spectra.

Grant Support:

NIH AM13212, 5T32GM08399 /M10⁴

THE VACUUM UV CD SPECTRUM OF A G-G-C TRIPLEX

Kenneth H. Johnson (Baylor C.O.M.), Michael E. Hogan (Baylor C.O.M.), John C. Sutherland (BNL)

Vacuum UV circular dichroism (CD) spectra were measured down to 175 nm for d(C)₁₀, d(G)₁₀, d(G)₁₀·d(C)₁₀, and d(G)₁₀·d(G)₁₀·d(C)₁₀. A CD difference spectrum was calculated for d(G)₁₀·d(G)₁₀·d(C)₁₀ (Figure). The difference spectrum was the CD induced by triplex formation from d(G)₁₀·d(C)₁₀ and d(G)₁₀. In the near-UV, the difference spectrum resembled the difference spectrum for d(G)₁₀·d(C)₁₀. We interpreted this as indicating that the bases of the purine strands of d(G)₁₀·d(C)₁₀ and d(G)₁₀·d(G)₁₀·d(C)₁₀ were stacked in a similar way. The d(G)₁₀·d(G)₁₀·d(C)₁₀ vacuum UV difference spectrum had a negative band at 195 nm and a positive band at 180 nm. Poly[d(G-C)·d(G-C)] in the Z conformation also has a negative band at 195 nm and a positive band at or below 180 nm (Proc. Natl. Acad. Sci. USA 78, 4801-4804). The similar bands could indicate that the third strand of d(G)₁₀·d(G)₁₀·d(C)₁₀ contains guanine in the syn conformation as in the Z conformation of poly[d(G-C)·d(G-C)].

Research sponsored by the NSLS Faculty-Student Support Program and by grants from the Office of Naval Research and Triplex Pharmaceutical Corporation.

CIRCULAR DICHROISM STUDY OF AN OLIGODEOXYRIBONUCLEOTIDE TRIPLEX

Y.-W. Park, G. E. Plum, K. J. Breslauer (Rutgers U.) and J. C. Sutherland (BNL)

We have collected simultaneously ultraviolet circular dichroism (CD) and pseudoabsorbance spectra of an oligodeoxyribonucleotide triplex and its constituent duplex and single strands.

Prior to use, the samples were desalted, lyophilized and dissolved in H₂O containing 10 mM Phosphate, 200 mM NaCl, and 1 mM EDTA at pH 6.5. Molar extinction coefficients at 260 nm were determined by phosphate analysis. Spectra were collected from 330 to 170 nm at 1 nm intervals with a 2 mm slit width and 10 second signal averaging. All spectra were collected at 20 °C in a dismountable quartz windowed "Gray" cell. The pathlength, as determined from the absorbance of oligonucleotide solutions of known concentration, was from 3 to 4 μm. CD spectra were corrected by subtraction of a solvent spectrum and normalized by multiplication by the known molar extinction coefficients and division by the measured absorbance at 260 nm.

Fig. 1 shows spectra of a 15mer, y15 [TTTTCTCTCTCTCT] and two 21mers, u21 [GCTAAAGAGAGAGAGATCG] and y21 [CGATCTCTCTCTCTTTTATGCT]. Each of the three spectra has a positive band at 192 nm. The two predominantly pyrimidine molecules, y15 and y21, exhibit very similar CD spectra. The purine rich oligonucleotide, u21, has a spectrum which contains an additional positive band at 216 nm.

Fig. 2 shows the CD spectrum of the Watson-Crick base paired duplex which is composed of the two complementary 21mer strands, y21 and u21. The 192 nm band is present but is of greater magnitude than the sum of the constituent single strand spectra.

Fig. 3 shows the CD spectrum of the triplex formed by the Hoogsteen pairing of y15 in the minor groove of the u21-y21 duplex. For comparison, the sum of the duplex and y15 spectra is also displayed in Fig. 3. The formation of triplex causes a shift of the major band from 192 to 184 nm. Also evident is the appearance of a negative band between 205 and 230 nm.

Research sponsored by NIH grants GM 23509, GM 34469, and CA 47795 (KJB)

Fig. 1 CD Spectra of Single Stranded DNA, y15 (—), y21 (---), u21 (....)

Fig. 2 CD Spectrum of the 21mer Duplex, u21-y21

Fig. 3 CD of Triplex, u21-y21-y15 (—) and Composite u21-y21 + y15 (---)

INVESTIGATION OF THE AL L EXCITON IN AlGaAs BY SOFT X-RAY REFLECTION AND EMISSION.

Q.-Y. Dong, W. L. O'Brien, J. J. Jia and T. A. Callcott, (U. of TN) D. R. Mueller and D. L. Ederer (NIST)

The Al L₂₃ soft-x-ray reflection and emission spectra of Al_xGa_{1-x}As have been measured as a function of x (0.17 < x < 1.0). Valence band maxima were determined from the emission spectra using a fit based on band structure parameters. The dominant features in the reflection spectra are produced by the L₂₃ core excitons. Since both the reflection spectra and emission spectra were obtained using the same soft x-ray spectrometer, the energy of the excitons above the valence band maximum are accurately obtained. The known values of the optical band gap, as a function of x, are then used to determine the core exciton binding energy. The core excitons are found to follow and lie above the conduction band minimum by 0.3 eV as the Al composition changes (see Fig.). Our results are compared to various core exciton theories and to previous experiments.

Measured position of exciton relative to VBM compared to local minima at X, L and Γ , vs. x. This work was supported by the NSF under grant number DMR-8715430 and Martin Marietta Labs-Baltimore and USDOE contract DE-AC05-84OR21400.

COPPER $M_{II,III}$ X-RAY EMISSION SPECTRA FOR CuO , Cu_2O AND $Bi_2Sr_2CaCu_2O_8$.

D. R. Mueller, D. L. Ederer NIST, Gaithersburg, MD. 20899
J. J. Jia, W. L. O'Brien, Q. Y. Dong, & T. A. Callcott U. Tenn. Knoxville 37996

Soft X-ray emission spectroscopy can provide a relatively bulk-sensitive measure of the electronic partial density of states around the atoms within a sample. Photon excited soft X-ray emission spectra for CuO , Cu_2O and several cuprate-based high temperature superconductors in the region of the $Cu\ 3d \rightarrow 3p$ transitions were measured with the aim of obtaining electronic structural information. For CuO and Cu_2O transitions from the valence levels into the M_{II} and M_{III} levels were clearly resolved due to the narrowness of the copper d contribution to the valence band in these compounds. The emission features were separated by the 2.3 eV spin-orbit splitting of the copper $3p_{1/2}$ and $3p_{3/2}$ states, and appeared with the statistical intensity ratio. The emission spectrum for Cu metal is shifted by several eV with respect to spectra from the simple oxides. For all three oxides the overall width and shape of the copper M_{III} emission band, extracted from the measurements, was found to agree fairly well with calculations^{1,2,3} for the ground state copper $3d$ partial density of states. [1] K. T. Park et al. J. Phys. Soc. Japan **57** (1988) 3445. [2] P. Marksteiner et al. Z. Phys. B **54** (1986) 119. [3] S. Massidda et al. Physica C **152** (1988) 251

Soft X-ray emission in the region of the Cu valence to $3p$ transitions. The spectra have been corrected for background and variations in spectrometer response, and have been offset for clarity. The oxide spectra were excited by 102 eV photons, the metal spectrum by 2 keV electrons. Near threshold excitation reduces the satellite emission intensity.

THE $N_{IV,V} \rightarrow O_{II,III}$ SOFT X-RAY EMISSION SPECTRUM OF La .

D. R. Mueller, C. W. Clark, D. L. Ederer NIST, Gaithersburg, MD. 20899
J. J. Jia, W. L. O'Brien, Q. Y. Dong, & T. A. Callcott U. Tenn. Knoxville 37996

Soft X-ray emission spectra for the $La\ N_{IV,V} \rightarrow O_{II,III}$ transition in lanthanum aluminate include contributions from several relatively intense multiplet lines near 80 eV. The appearance of this spectrum contrasts sharply with the three diagram lines (a weak $4f = 0$ and two $4f = 1$ transitions) observed for the $N_{IV,V} \rightarrow O_{II,III}$ emission spectrum of Ba in barium fluoride. An atomic model of the $La\ N_{IV,V} \rightarrow O_{II,III}$ transition assuming single occupancy of the $4f$ shell by a spectator electron provides a good approximation to the measured spectrum. Emission features due to the low energy $3p$ and $3d$ configurations of $4d^9 4f$ are also apparent in the lanthanum aluminate emission spectra but not in emission spectra from barium fluoride. Reflection measurements indicate that the $4f$ level is not occupied in the ground state in $LaAlO_3$. The data suggest that charge transfer from ligand orbitals following creation of a $4d$ vacancy prior to the emission process is the predominate mechanism giving rise to the required $4f$ occupancy.

Soft X-ray emission in the region of the $N_{IV,V} \rightarrow O_{II,III}$ ($4d \rightarrow 5p$) transitions for barium in barium fluoride (lower curve) and lanthanum in lanthanum aluminate (upper curve). The spectra have been offset for clarity.

ELECTRONIC STRUCTURE OF MgO , $\alpha\text{-Al}_2\text{O}_3$ and MgAl_2O_4 , OCCUPIED AND UNOCCUPIED BULK DOS.

W. L. O'Brien, J. Jia, Q.-Y. Dong and T. A. Callcott, (U. of TN) D. R. Mueller and D. L. Ederer (NIST)

The electronic structure of MgO , $\alpha\text{-Al}_2\text{O}_3$ and MgAl_2O_4 have recently been the subject of both experimental and theoretical studies. Taken as a group these three oxides cover a broad range of crystalline structures. MgO is highly ionic and has a simple NaCl-type structure. $\alpha\text{-Al}_2\text{O}_3$ has the more complex corundum structure with each Al atom in a distorted octahedral hole. The spinel oxide MgAl_2O_4 has an even more complicated structure with Al atoms in octahedral sites and Mg atoms in tetrahedral sites. In this work we investigate the electronic properties of MgO , $\alpha\text{-Al}_2\text{O}_3$ and MgAl_2O_4 by soft-x-ray emission (SXE) and soft-x-ray reflection (SXR) spectroscopy. With these two techniques we can measure both the occupied and unoccupied local partial-density of states with the same spectrometer (see Fig). The density of states measured are local since a core hole participates in the electronic transition, and the density of states are partial since dipole selection rules are obeyed. These new measurements are compared to optical, and theoretical results on the same three materials. It is shown that these new measurements allow a more quantitative comparison to theory than the more traditional optical measurements which yield a joint density of states. A discussion on the need for theories to include both local and symmetry projections will be included.

This work was supported by the National Science Foundation under grant number DMR-8715430, and the National Institute of Standards and Technology.

PHONON COUPLING TO CORE EXCITATIONS IN IONIC SOLIDS.

W. L. O'Brien, J. Jia, Q.-Y. Dong and T. A. Callcott, (U. of TN) D. R. Mueller and D. L. Ederer (NIST)

Phonon coupling to electronic transitions have been of interest for over thirty years. Ionic solids have played a significant role in this interest due to the simplifying assumptions which can be used to accurately describe the observed coupling. We present new temperature dependent soft x-ray emission (SXE) and soft x-ray absorption (SXA) measurements on MgO which show in detail the final states affect on phonon coupling for transitions involving the Mg 2P core hole. As the temperature is increased both the SXE and SXA spectrum are observed to change. In SXE the valence band edge is broadened and shifted towards higher energies while no energy shift or broadening is observed for the $2P \rightarrow 2S$ core-to-core transition. In SXA the core-exciton is broadened and shifted to lower energies while conduction band features are broadened but not shifted. The temperature dependent broadening of the conduction band features are similar to the broadening of the valence band edge but significantly different than that of the core-exciton, (see Fig). The total width of these different features, assuming Gaussian broadening, are discussed in terms of partial relaxation and the phonon self energies for the different transitions. Temperature dependent energy shifts of the valence band edge, core-exciton and valence band and conduction band features are discussed in terms of band structure and thermal expansion. Comparisons are made to similar measurements on $\alpha\text{-Al}_2\text{O}_3$ and $\alpha\text{-SiO}_2$.

This work was supported by the National Science Foundation under grant number DMR-8715430, and the National Institute of Standards and Technology.

PHOTON-INDUCED CHEMISTRY IN WEAKLY-BOUND THIOPHENE CLUSTERS

P. CIP-AGUERO, E.A. WALTERS, Dept. of Chemistry, University of New Mexico, Albuquerque, NM 87131
J.R. GROVER, Dept. of Chemistry, Brookhaven National Laboratory, Upton, NY 11973.

Van der Waals clusters of thiophene, C_4H_4S , were prepared by free-jet expansions of 1:9 mixtures of C_4H_4S :Ar and C_4H_4S :He and studied by photoionization of molecular beams containing the clusters with mass spectrometry for ion detection. Size distributions of neutral clusters in both gas mixtures were determined, Fig. 1a. A quantitative analysis was made of the efficient cooling properties of Ar compared to He as a carrier gas. For the clusters, we obtain $IP((C_4H_4S)_2) = 8.27 \pm 0.03$ eV and $IP((C_4H_4S)_3) = 8.17 \pm 0.04$ eV. The product $C_4H_4S^+$ (m/e 134) is observed in considerable abundance; we find $AP(C_4H_4S^+) = 15.50 \pm 0.07$ eV. Its photoion yield curve is shown in Fig. 1b. Analysis of the pressure dependence of m/e 134 yields shows that the ion signal tracks the abundance of $(C_4H_4S)_3$ instead of the dimer. Thus, the reaction leading to $C_4H_4S^+$ is $(C_4H_4S)_3 + h\nu \rightarrow C_4H_4S^+ + C_4H_4S$. The results are compatible with the ion $C_4H_4S^+$ being the complex $(C_4H_4S \cdot C_4H_2)^+$ with $C_4H_2^+$ formed in its ground electronic state.

Figure 1. (a) The number density distribution of $(C_4H_4S)_n$, $n = 1-3$, in expansion of a 1:9 Ar: C_4H_4S mixture.
(b) Photoionization yield curve for the production of $C_4H_4S^+$ from a 1:9 Ar: C_4H_4S mixture at the stagnation pressure of 149 torr.

Supported in part the U.S. Department of Energy and in part by the U.S. Air Force.

U11
PHOTOIONIZATION STUDIES OF TRIFLUOROMETHYL BROMIDE AND A CLUSTER WITH METHANOL

J.T. CLAY, E.A. WALTERS, M.V. WILLCOX, Dept. of Chemistry, University of New Mexico, Albuquerque, NM 87131
J.R. GROVER, Dept. of Chemistry, Brookhaven National Laboratory, Upton, NY 11973.

The production of CF_3Br^+ from jet-cooled CF_3Br in a molecular beam was determined as a function of wavelength by single photon ionization, Fig. 1a. Striking features of this curve are the lack of step-like structures on the rise from threshold and the abrupt decrease in ion production that gives rise to the maximum at 1030 Å. We deduce that the threshold for forming CF_3Br^+ ($2E_{1/2}$) is 11.50 ± 0.03 eV and that three vibrational states are weakly evident. At 11.85 eV the higher spin-orbit state is produced, CF_3Br^+ ($2E_{1/2}$), and one bound vibrational level is seen about 725 cm^{-1} higher. Immediately above this CF_3Br^+ efficiently dissociates into CF_3^+ and Br.

Further methodology was developed to measure neutral cluster distributions from systematic changes in photoion yield curves at energies well above the ionization threshold. This was used to separate the yields of m/e 180, $CH_3OH \cdot CF_3Br^+$, and m/e 298, $(CF_3Br)_2^+$, into the fraction arising from the corresponding neutral dimer. The balance comes from larger clusters; Fig. 1b. The source gas in both cases was a 9:0.3:100 mixture of CF_3Br : CH_3OH :Ar.

Figure 1. (a) Threshold photoion yield curve for CF_3Br^+ from neat CF_3Br at 499 torr, open circles, and the derivative of the cubic spline smoothed fit, line. (b) Fraction of observed $(CF_3Br)_2^+$ circles, and $(CF_3Br \cdot CH_3OH)^+$, diamonds, coming from corresponding neutral dimers as a function of pressure.

Supported in part the U.S. Department of Energy and in part by the U.S. Air Force.

COMPLEXES OF OXYGEN WITH BENZENE AND HEXAFLUOROBENZENE

J. R. Grover*, G. Hagenow* and E. A. Walters*

*Department of Chemistry, Brookhaven National Laboratory, Upton, NY 11973,

†Institut für Physikalische Chemie, Freie Universität Berlin, Berlin, Germany

#Department of Chemistry, U. of New Mexico, Albuquerque, NM 87131

Complexes of molecular oxygen with benzene and hexafluorobenzene were synthesized by jet expansion and studied by one-photon ionization. The neutral heterodimers $C_6H_6 \cdot O_2$ and $C_6F_6 \cdot O_2$ were found to have ionization potentials of 9.172 ± 0.004 eV and 9.856 ± 0.003 eV, respectively, where both thresholds are sharp because of autoionization. Final values (see NLS Ann. Rpt. 1989) of the heterodimers' dissociation energies (in kcal mol⁻¹) are $D_0(C_6H_6 \cdot O_2) = 1.65 \pm 0.32$, $D_0[(C_6H_6 \cdot O_2)^+] = 3.37 \pm 0.34$, $D_0(C_6F_6 \cdot O_2) = 2.1 \pm 0.4$, and $D_0[(C_6F_6 \cdot O_2)^+] = 3.2 \pm 0.4$. Empirical calculations from electrostatic and van der Waals terms give $D_0(C_6H_6 \cdot O_2) = 1.35$ and $D_0(C_6F_6 \cdot O_2) = 2.3$. Surprisingly, the ion $(C_6H_6 \cdot O_2)^+$ is formed only from $C_6H_6 \cdot O_2$ and not from larger clusters. On the other hand, intracuster reaction products $C_6H_6O^+$ and $C_6F_6O^+$ are not formed at all from the dimers, but only from larger clusters, and their onsets are far above their thermodynamic thresholds. These data are consistent with a model, supported by the known electronic states of oxygen, in which an excited organic ion is created intracuster, dissociates its O_2 neighbor, captures one of the recoiling O atoms, and the reaction product is then stabilized by the excitation/ejection of a third neutral member of the cluster. The above calculations give structures for the heterodimers, Fig. 1, that are consistent with this model.

Figure 1. Calculated structures for the trimers $(C_6H_6)_2O_2$. A, and $(C_6F_6)_2O_2$. B, that are supported by the data.

This research was carried out at Brookhaven National Laboratory under contract DE-AC02-76CH00016 with the U.S. Department of Energy and supported by its Division of Chemical Sciences, Office of Basic Energy Sciences.

KINETIC ENERGY RELEASE IN THE DISSOCIATIVE PHOTOIONIZATION OF VAN DER WAALS CLUSTERS

J. R. Grover*, E. A. Walters†, D. L. Arneberg†, G. Hagenow†, J. T. Clay†, and M. V. Willcox†, *Department of Chemistry, Brookhaven National Laboratory, Upton, NY 11973, †Department of Chemistry, University of New Mexico, Albuquerque, NM 87131, †Institut für Physikalische Chemie, Freie Universität Berlin, Berlin, Germany

Dissociative photoionization of van der Waals clusters occurs quite easily because the forces holding the constituent molecules together are very weak. This is especially true for clusters of mixed composition. When the kinetic energies of such ions were measured with a recently-developed instrument, we were surprised to find that they are produced with energies larger than van der Waals bond strengths, which are about 0.1 eV. Figure 1 shows a test of the instrumental response, using the reaction $CO_2 + h\nu \rightarrow CO^+ + e^-$, where two vibrational states of CO are populated, and the O^+ is produced at energies of 0.10 and 0.27 eV with respect to the center-of-mass. Approximate deconvolution of these data gives 0.09 and 0.28 eV. The spectrum for $(C_4H_6 \cdot SO_2)^+$ produced from the photoionization of $(1,3-C_4H_6)_n(SO_2)_m$, figure 2, shows a major component at about 0.08 eV, and a minor component at 0.16 eV. This corresponds to c.m. energies of 0.11 and 0.24 eV if the $(C_4H_6 \cdot SO_2)^+$ is produced from trimers, or 0.16 and 0.32 eV if from tetramers.

Fig. 1 Instrumental energy distribution of O^+ from dissociative photoionization of a molecular beam of CO_2 . The solid line gives the locus of O^+ born with zero energy.

Fig. 2 Instrumental energy distribution of $(C_4H_6 \cdot SO_2)^+$ from dissociative photoionization of a cluster beam of $(1,3-C_4H_6)_n(SO_2)_m$.

This research was carried out at Brookhaven National Laboratory under contract DE-AC02-76CH00016 with the U.S. Department of Energy and supported in part by its Division of Chemical Sciences, Office of Basic Energy Sciences and in part by the Engineering and Services Laboratory, Air Force Engineering and Services Center, Tyndall AFB, FL 32403.

PHOTOIONIZATION MASS SPECTROMETRIC STUDY OF $N + C_2H_5$ REACTION PRODUCTS*

R.B. Klemm,[†] W. Tao[‡], F.L. Nesbitt,[§] and L.J. Stief[§]

The products of the reaction of ground-state atomic nitrogen with the ethyl radical have been determined in a discharge flow system with collision-free sampling to a photoionization mass spectrometer (DF-PIMS) using dispersed synchrotron radiation as the ionizing source. The only products detected at ionization energies ≤ 11.5 eV were C_2H_4 and the free radicals H_2CN and CH_3 . Potential products not observed but detectable at these ionization energies are $H(CH_3)CN$ and adducts such as CH_3CHNH or $cyc-C_2H_4NH$. Identification of the C_2H_4 , H_2CN and CH_3 products is based on the unambiguous combination of product m/e values, the shift observed for the $N + C_2D_5$ system (based on mass scans) and the features of the photoionization spectra and photoionization threshold (based on wavelength scans). These results suggest that the major channels for the reaction are:

The figure shows a mass scan performed at $\lambda = 114$ nm that illustrates the observed products from the reaction of N with C_2D_5 : CD_3 ($m/e = 18$); D_2CN ($m/e = 30$); C_2D_4 ($m/e = 32$); and reactant C_2D_5 ($m/e = 34$).

*This work was supported by the Division of Chemical Sciences, U.S. Department of Energy, Washington, D.C., under contract No. DE-AC02-76CH00016. Also, we thank E. Gregory for his contributions to this research.

[†]Brookhaven National Laboratory, Department of Applied Science.

[‡]NASA/Goddard Space Flight Center, Greenbelt, MD.

PHOTOIONIZATION SPECTRA AND IONIZATION THRESHOLDS FOR H_2CN AND D_2CN *

F.L. Nesbitt,[§] G. Marston,[§] L.J. Stief,[§] M.A. Wickramaratne,[†] W. Tao,[‡] and R.B. Klemm,[†]

The photoionization spectra of the H_2CN and D_2CN radicals were obtained by photoionization mass spectrometry. The radicals were generated by the reaction of N atoms with CH_3 and CD_3 , respectively. For both H_2CN and D_2CN a prominent feature was observed near 118.6 nm (10.5 eV) and the ionization threshold was determined to be 9.4 ± 0.1 eV (see Figure 1). Using the value for ΔH_f (H_2CN) derived from a recent electron affinity measurement (Ellison et al.) and other available measured or calculated thermochemical quantities for H_2CN and $HCNH$ radicals and radical-ions, we derive a value of 10.8 ± 0.6 eV for the ionization energy of H_2CN . The much lower value derived for the ionization energy of $HCNH$ (6.8 eV and 7.0 eV for the cis and trans isomers, respectively) is consistent with the product of the $N + CH_3$ reaction being the H_2CN isomer and not $HCNH$. The ionization threshold observed at 9.4 eV is attributed to autoionization arising from high Rydberg states of H_2CN which couple into vibrationally excited states of the linear $HCNH^+$ ground state of the ion.

Figure 1. Photoionization spectrum of H_2CN . The photoion yield is the ion counts at m/e 28 divided by the light intensity in arbitrary units.

*This work was supported by the Division of Chemical Sciences, U.S. Department of Energy, Washington, D.C., under contract No. DE-AC02-76CH00016. Also, we thank E.J. Gregory for his contributions to this research.

[†]Brookhaven National Laboratory, Department of Applied Science.

[‡]NASA/Goddard Space Flight Center, Greenbelt, MD.

A POSSIBLE ROLE FOR IONS IN FLAME EXTINGUISHMENT BY HALONS

E.A. Walters (U. New Mexico), J.R. Grover (BNL), J.T. Clay, M.V. Willcox, D.L. Arnebergs, and J.S. Ninitz (U. New Mexico)

Halons, a class of halogenated hydrocarbons, are widely used as fire extinguishing agents because of their effectiveness, cleanliness, and low toxicity. They have long tropospheric lifetimes, so they eventually enter the stratosphere where photolysis produces atomic Br which catalyzes ozone decomposition. Consequently, replacements must be found that will not adversely impact stratospheric ozone. We have investigated some of the initial reactions between Halon 1301 (CF_3Br) and the primary flame free radicals O , OH , and H by dissociative photoionization of weakly bound molecular complexes of CF_3Br with free radical precursors, O_2 and CH_3OH , in a mass spectrometer. A striking result is that the only product of intra-cluster chemical reactions is HBr^+ . The absence of oxygenated reaction products is highly unusual. Insight into why this occurs comes from the photoion yield curve for CF_3Br , Fig. 1, and its fragment CF_3^+ , Fig. 2. In Fig. 1, the ionization threshold of CF_3Br is at 11.48 ± 0.04 eV and the ion yield increases to an abrupt peak near 11.90 eV, after which the yield remains approximately constant to 20 eV before decreasing. The abrupt peak and low ion yield are due to the onset of fragmentation to CF_3^+ beginning at 11.82 ± 0.04 eV, Fig. 2. We deduce the C-Br bond energy in CF_3Br^+ to be 8 kcal mol^{-1} , too low to be a chemical bond. This suggests a special role for this family of ions. One possibility is proton transfer reaction of CF_3Br with the dominant ion in flames CHO^+ :

$\text{CHO}^+ + \text{CF}_3\text{Br} \rightarrow \text{CO} + \text{HBr} + \text{CF}_3^+$. This reaction is exothermic by 94 kcal mol^{-1} and is driven in large measure by the stability of CF_3^+ . The HBr product is important because it participates catalytically in the chain-terminating recombination of atomic hydrogen.

Fig. 1. CF_3Br^+ photoion yield curve; the threshold is 1078.39 ± 0.04 eV. (Supported by the U.S. Dept. of Energy and the U.S. Air Force)

Fig. 2. CF_3^+ photoion yield curve; the appearance potential is 1046 ± 3 eV 11.82 ± 0.04 eV

COVERAGE DEPENDENCE OF THE LI K-EDGE ABSORPTION FOR $\text{Li}/\text{Be}(1120)$

P.A. Bruhwyler, E.W. Plummer, University of Pennsylvania

We have measured the Li K absorption spectrum using total electron yield for many coverages Θ between 0 and the monolayer saturation of $\text{Li}/\text{Be}(1120)$, as well as for all n-layer coverages up to $n=4$. Fig. 1 summarizes the data. Θ was monitored by work function and by the Li 1s core level photoemission spectrum, which for Θ below saturation was a single line, and for Θ above saturation was a multiplet corresponding to interface, surface, and "in-between" layers of the films for $n \geq 3$. We estimate that the coverages are accurate to within 10%. Judging from the core line intensities, the deposition was largely epitaxial. A metal-like step function threshold occurs at all values of Θ , indicating that the Li atoms are significantly metallicity bonded even at the lowest coverages.

Li/Be(110) Yields vs Coverage

ANGLE-RESOLVED PHOTOEMISSION INVESTIGATION OF BI OVERLAYERS ON InSb(110) SURFACES

N. J. DiNardo (Drexel U.); D. McLroy and D. Heskett (U. of Rhode Island); A. B. McLean (Queen's U.); R. Ludeke and H. Munkaza (IBM)

The electronic structure of epitaxial Bi-InSb(110) interfaces has been studied using angle-resolved photoelectron spectroscopy. This study is part of a comprehensive effort to systematically determine the relationship between geometric structure, electronic structure, and overlayer strain at semimetal/III-V(110) interfaces. This work may be related to recent studies of the Bi/GaAs(110)¹ and Bi/InAs(110)² systems. In contrast to those systems, however, two epitaxial Bi structures can be grown on InSb(110) surfaces³ - a 1x1 structure for Bi coverages up to one monolayer and a 1x2 structure above one monolayer. For these studies, a clean surface was obtained by sputtering and annealing a thick MBE-grown film of InSb on a GaAs(110) template. Concentrating on the 1x1 Bi overlayer for comparison to the previous work,^{1,2} we illustrate a series of photoemission spectra to track the dispersion of surface states associated with the Bi overlayer from $\bar{X}$ to $\bar{\Gamma}_2$. We identify the surface states S' , S'' , and S''' , which can be assigned⁴ to originate from bonding surface state bands (S' , S'') and from intrachain bonding p states (S'''). The dispersion throughout the surface Brillouin zone and the polarization-dependence of these features have also been measured. For the (1x1) overlayer at $\bar{X}$, it was found that S' is polarized perpendicular to the surface while S''' is polarized parallel to the surface, supporting the predicted origin of the bands.⁴

References

1. A. B. McLean, R. Ludeke, M. Prietsch, D. Heskett, D. Tang, and T. Maeda Wong, *Phys. Rev. B* **43**, 7243 (1991).
2. D. McLroy, D. Heskett, D. Swanson, A. B. McLean, J. Chen, N. J. DiNardo, M. Prietsch, R. Ludeke and H. Munkaza, to be published.
3. W. K. Ford, T. Guo, S. L. Lantz, K. Wan, S.-L. Chang, C. B. Duke, and D. L. Lessor, *J. Vac. Sci. Technol.* **B8**, 940 (1990).
4. C. Mailhior, C. B. Duke, and D. Chadi, *Phys. Rev. Lett.* **53**, 2114 (1984).

WORK CONDUCTED ON BEAM LINE

BEAM LINE U12B

CORE-LEVEL SPECTROSCOPY FROM LOW-INDEX SURFACES OF HfC EXPOSED TO OXYGEN*

G. R. Gruzalski,¹ S.-C. Lui,¹ D. Heskett,² X. Shi,² and D. M. Zehner¹
¹ORNL, ²Univ. of Rhode Island

The effect of oxygen upon the Hf-4f core levels of the (111), (100), and (110) surfaces of HfC was investigated using angle-resolved ultraviolet photoemission spectroscopy with synchrotron radiation.

The surfaces were cleaned by a series of 2-sec anneals to ~2500°C by bombarding their back sides with electrons (2 kV, 100 mA). Exposures were made by filling the chamber with oxygen of 5N purity; to reduce contamination from exchange reactions, the vacuum system was continuously ion pumped during all exposures. An energy resolution of ~220 meV was attained from the combination of a toroidal-grating monochromator and angle-resolved analyzer.

Photoemission spectra from the Hf-4f region of the (111) surface of HfC are shown in Fig. 1. The photon energy of 60 eV was selected in an attempt to optimize conditions for meeting two criteria: a large sensitivity to the valence-band oxygen signal, which was used to monitor surface cleanliness, and a small escape depth of the photoemitted electrons, so as to maximize surface sensitivity. A shoulder to higher binding energies (BEs) can be seen in the spectrum from the clean (111) surface. Upon exposure to oxygen, a shoulder reduced in intensity appears (see the spectrum following 0.1 L exposure), which seems to break away and shift to even higher BEs with additional exposure. Similar effects are seen in the spectra from the (100) and (110) surfaces with two differences: The surface core-level shift is much less obvious in the spectra from the clean (100) and (110) surfaces, and a larger oxygen exposure is required before the extra set of peaks emerges. Significant here is the fact that the high-BE shoulder first reduces in intensity; this observation supports our interpretation that the shoulder is due to photoemission from Hf atoms on the surface.

Fig. 1. Photoemission spectra from the Hf 4f region of clean HfC(111) and HfC(111) exposed to oxygen.

*This research was sponsored by the U.S. Department of Energy under Contract No. DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc., and by the URI Council for Research.

THE EXPERIMENTAL BAND STRUCTURE OF Be(1120)

J.B. Hannon, Geunseop Lee, and E.W. Plummer
Department of Physics, University of Pennsylvania

We have performed angle-resolved photoemission from the (1120) surface of Beryllium. One goal of this experiment was to see if a proposed surface reconstruction would alter the surface electronic structure via unklapp processes that would be forbidden for the bulk-terminated surface. Two non-dispersing features were observed at 1.5 and 2.5 eV binding energy, which may imply localized surface states, or may be due to poor surface quality, and is still under investigation. Our measurement of the bandwidth (11.1 eV) agrees with previous measurements from the Be(0001) surface, as does the FWHM of the lowest occupied bulk band state (3.5 eV).

Figure Caption : The lowest occupied bulk band of Be at Γ . The fit is to a lorentzian, with a standard inelastic background.

WORK CONDUCTED ON BEAMLINE

BEAMLINE U12P

CORE-LEVEL SPECTROSCOPY FROM LOW-INDEX SURFACES OF TaC*

S.-C. Lui,¹ G. R. Gruzalski,¹ D. Heskett,² X. Shi,² and D. M. Zehner¹
¹ORNL, ²Univ. of Rhode Island

We have employed the technique of angle-resolved ultraviolet photoemission spectroscopy with synchrotron radiation to study the 4f core levels of the (111), (100), (110), and (130) faces of TaC.

The surfaces were cleaned by a series of 2-sec anneals to ~2500°C by bombarding their back sides with electrons (2 kV, 100 mA). An energy resolution of ~220 meV was attained from the combination of a toroidal-grating monochromator and angle-resolved analyzer.

In Fig. 1 are shown normal emission spectra of the Ta 4f region for the (100) and (111) surfaces of TaC. The top curve shows emission from the surface and deeper layers of TaC(100); similar spectra were obtained from the (110) and (130) faces, which we believe microfacet to expose regions of (100) surface. The bottom curve shows emission from the 4f levels of TaC(111). Note that for this surface there is a large surface core level shift (~0.58 eV) to lower binding energies (BEs) and that the FWHMs of the surface peaks in the spectrum are larger than those of the corresponding bulk peaks.

As the carbon concentration decreases in bulk TaC_x, the electron density increases in the vicinity of the Ta ions and decreases in the vicinity of the C ions, causing the 4f BE to decrease.³ A metal atom on a Ta-terminated (111) surface of TaC has three of six nearest-neighbor carbon atoms missing. If the resulting charge redistribution were similar to what occurs in the bulk owing to the formation of carbon vacancies, one might expect the surface core-level BE to be smaller than that of the bulk, which is exactly what is seen in Fig. 1. Although other factors contribute to surface core-level shifts, we contend that charge transfer plays the major role for TaC(111), and we therefore interpret these results as suggesting that the (111) surface is Ta terminated, at least in part. Similarly, the small surface core level shifts on the (100), (110), and (130) faces suggest that the charge redistribution owing to their creation is small.

*G. R. Gruzalski and D. M. Zehner, *Phys. Rev. B* 34, 3841 (1986).

*This research was sponsored by the U.S. Department of Energy under Contract No. DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

Fig. 1. Photoemission spectra of Ta 4f region for TaC(100) and TaC(111).

Interaction of Hydrogen with Mg(0001)

P.T. Sprunger, P.A. Bruhwiler, and E.W. Plummer
Department of Physics and LRSM, University of Pennsylvania

We have studied the electronic structure of the H/Mg(0001) system by angle-resolved photoemission and core level spectroscopy. The results of the core level spectroscopy has shown that at 150K the adsorbed atomic hydrogen forms a "surface hydride"; but upon annealing the system to 425K (before the hydrogen desorbs) the hydrogen is absorbed into the bulk. This is consistent with earlier studies of this system with other spectroscopies (LEED, EELS, $\Delta\Phi$) which has shown that the interaction of atomic hydrogen on Mg(0001) at 150K has vibrational losses characteristic of an adsorbed "hydride" system; but annealing the system to 425K the vibrational spectra has shown that the hydrogen might be absorbed into the subsurface or bulk region. Using core level spectroscopy, this system has been investigated as a function of coverage and temperature. At 150K (fig.1), the Mg(2p) core level (49.6 eV) spectra shows a large shifted core peak (-1.65 eV) which increases as a function of exposure indicating a strongly chemisorbed system ("surface hydride") is forming in which there is a large amount of charge transfer in the surface region. The attenuation of the unshifted core peak is attributed to the subsurface Mg atoms which are unperturbed by the surface hydrogen. Fig. 2 shows that upon annealing the system to 425K, the shifted core peak is absent and the unshifted core is broader indicating that the hydrogen has been absorbed into the subsurface region in which the electronic structure is reminiscent of a soluble hydrogen system.

Figure 1. Mg(2p) core level spectra as a function of atomic hydrogen coverage (150K)

Figure 2. Mg(2p) core level spectra as a function of annealing temperature.

BULK BAND STRUCTURE OF LITHIUM

G. M. Watson, X. Y. Liu, and E. W. Plummer
Physics Dept., University of Pennsylvania, Philadelphia, PA 19104

We have measured the bulk band structure of BCC Li along the $\Gamma \rightarrow N$ direction of the bulk Brillouin Zone. Figure one shows a plot of the measured initial state energy as a function of photon energy. The solid line in figure one is the expected dispersion assuming Li is a free electron metal. Clearly both the occupied bandwidth and the shape of the dispersion are not described by the free electron model. The LDA value for the bottom of the band is 3.45 eV¹, in better agreement with our results. The difference between our results and the LDA value is attributed to the quasiparticle self-energy. The quasiparticle bandwidth in the GW approximation is 2.84 eV¹ in nearly exact agreement with our measurement of 2.85 eV.

Figure 1.

1 J. E. Northrup, M. S. Hybertsen, and S. G. Louie, Physical Review B 39, 8198 (1989).

CORE ELECTRON EXCITATION OF SIMPLE MOLECULES TO RYDBERG STATES - A NEW METHOD TO SEE HIDDEN RESONANCE EXCITATIONS

D. Y. Kim, K. Lee, C.-I. Ma, M. Mahalingam, and D. M. Hanson (SUNY at Stony Brook)
S. L. Hulbert (NSLS, BNL)

High resolution and high flux makes it possible to determine the angular distribution of molecular fragmentation following core electron excitation. Two detectors at 90° and 0°, which are installed with high pass electrostatic filters, measured high kinetic energy ion yields. Relative collection efficiencies of the two detectors are calibrated by CH₄ and the degree of polarization of the synchrotron radiation is found from N₂ 1s→π* excitation. This angle-resolved measurement is directly correlated with photoabsorption anisotropy.

Fig. 1 shows ion yields of two detectors and the anisotropy parameter, β. The β spectrum is constructed from ion yields at the two detectors and the degree of light polarization. Excitations to π symmetry electronic states appear only at 90°, and excitations to σ symmetry electronic states appear only at 0°. Resonance peaks that have not been resolved by conventional high resolution spectroscopy can now be revealed in the two different detectors. The excitation resonance peak of 1s→3pσ is clearly shown as an isolated peak in the 0° detector. The number 6 peak in Fig. 2 of Ref. 1 is not a single σ state but consist of both π and σ symmetry states. Such polarized spectra provide much information that high resolution absorption spectroscopy can not provide. N₂, CO, O₂ and H₂O have been studied.

Fig. 1 Ion yields from N₂ at 90° and 0° and β spectra.

[1] C. T. Chen, Y. Ma and F. Sette, Phys. Rev. A **40**, 6737 (1989).

This research was supported by the National Science Foundation (CHE-8921729).

SELECTIVE EXCITATION OF CARBON IN ACETONE

C.-I. Ma, K. Lee, D. Y. Kim, and D. M. Hanson (SUNY at Stony Brook)
S. Hulbert (NSLS, BNL)

The effect of site and state selective core excitation on the electronic decay of molecules is demonstrated here by the carbon K-edge resonances in acetone. Two major resonances from methyl carbon 1s→3p and carbonyl carbon 1s→π* were observed in the total ion yield spectrum in Fig. 1 and the electron spectra at these two resonances are shown in Fig. 2 and 3. The spectral shape in Fig. 3 resembles the Auger spectrum of methane which implies that the site of the core hole is the methyl carbon. This similarity also indicates the spectator nature of the 3p electron. The features in the high energy region of Fig. 2 correspond to the one-hole states appearing in photoelectron spectra, implying participant decay processes in this region. These two spectra, together with the associated electron-ion coincidence data, exemplify the effect of the site and state specificity in molecular core hole decay. This is contrary to an earlier result¹ where site specificity was claimed based upon their assignment of unresolved resonances to carbonyl carbon 1s→π* transition.

I. W. Eberhardt, T. K. Sham, R. Carr, S. Krummacher, M. Strongin, S. L. Weng, and D. Wesner, Phys. Rev. Lett. **14**, 1038 (1983).

This research was supported by the National Science Foundation (CHE-8703340).

SOFT-X-RAY PROJECTION LITHOGRAPHY

R. Freeman, J. Bjorkholm, J. Bokor, K. Early, L. Eichner, M. Himel, T. Jewell, A. MacDowell, L. Szeto, D. Taylor, D. Tennant, W. Waskiewicz, D. White, D. Windt, O. Wood, II (AT&T Bell Labs.), P. Takacs (BNL), J. Bruning (GCA/Tropel), M. Ruda (Talandic Res.) and K. Thompson (Opt. Res. Assoc.).

The goal of this project is to demonstrate a reduction lithographic camera capable of resolution better than 0.1 microns. Our most recent progress includes the following:

1. We have compared the performance of Mo/Si multilayer coated reflection masks on rugged, stable substrates to that of a transmission mask in a 20:1 reduction Schwarzschild optic at 14 nm. There was no observable difference among reflection masks fabricated 3 different ways! (See Figure) or between the reflection and transmission mask results. This supports the contention that the ability to use a reflecting mask is a potential advantage of soft-x-ray projection printing over x-ray proximity printing.

Figure: SEM photo of 0.1 μm lines and spaces in a 60 nm thick layer of the resist PMMA produced with 3 reflection mask types.

2. We have dramatically improved the image quality in a 1:1 Offner ring-field imaging system² by illuminating each point on the mask with radiation converging to that point with a cone of angles and directing all the light from each point into the entrance pupil of the imaging system. This eliminated all evidence of edge ringing, speckle and other unwanted interferences and has led to a significantly better illumination system for use in the experiments planned for the U9A beamline.

J.D. M. Tennant, J. E. Bjorkholm, R. D'Souza, L. Eichner et al. J. Vac. Sci. Technol. B, Nov/Dec. 1991, to be published.
 2A. A. MacDowell, J. E. Bjorkholm, L. Eichner, R. R. Freeman et al., J. Vac. Sci. Technol. B, Nov/Dec. 1991, to be published.
 Supported by AT&T & NIST under Coop. Agreement No. 70NANB1H115.

X-RAY IMAGING AND PROJECTION LITHOGRAPHY

J. E. Bjorkholm, J. Bokor, R. M. D'Souza, L. Eichner, R. R. Freeman, T. E. Jewell, W. M. Mansfield, A. A. MacDowell, E. L. Raab, L. H. Szeto, D. M. Tennant, W. K. Waskiewicz, D. L. White, D. L. Windt, and O. R. Wood, II (AT&T BELL LABORATORIES)

Ring-Field Camera

We have carried out a number of experiments using an Offner ring-field camera system which provides 1:1 imaging over a slit-shaped field several cm long and 100 μm wide. Current experiments are carried out with 46 nm radiation using Ir-coated optics; in the future we will use 14 nm radiation with multilayer-coated optics to achieve even better resolution. First, we used an open-stencil transmission mask to demonstrate the printing of features as small as 0.2 μm . This resolution is at the limit set by the Rayleigh criterion and it attests to the high quality of the spherical mirrors used in this objective. In fact, we can infer that the surface figure accuracy is better than can be measured using state of the art interferometry. Next, we dramatically improved the fidelity of the imaging by taking pains to correctly illuminate the mask in order to properly fill the entrance pupil of the system. Finally, we carried out our first imaging using a reflection mask; the mask was composed of a fine pattern of Au on a silicon substrate. Imaging at the diffraction limit was obtained. In the near future we will use the ring field system at 14 nm; to do this we will coat another set of mirrors with appropriate multilayers. In this configuration the 1:1 imaging will be very useful for the inspection of multilayer reflection masks intended for use with reduction systems having sub-0.1 μm resolution.

Multilayer Reflection Masks

We have carried out imaging at 14 nm using the 20:1-reduction Schwarzschild optical system described previously and multilayer reflection masks. These masks were fabricated by etching a pattern into a reflective multilayer coating, thereby yielding a finely-patterned reflector. This demonstration is important since reflection masks of this type promise to be far superior to transmission masks which are, of necessity, very thin and fragile in this part of the spectrum.

Precision Metrology of Surface Figures

The US optical fabrication industry does not currently possess the technology to test and/or fabricate mirrors with the precision required for practical soft x-ray projection lithography. We are initiating a major program to address this problem and we have received a 3 year grant from the Advanced Technology Program administered by the National Institute of Standards and Technology to partially fund this work. The undulator beamline U13UB will play a central role in our plans to develop precision tests of the surface figures of soft x-ray optics.

Work supported by AT&T Bell Laboratories

LINE SHAPE OF THE Si L₂₃VV AUGER SPECTRUM OF STOICHIOMETRIC SiO₂ MEASURED BY APECS

R.A. Bartynski and A.K. See (Ruigers U.), C.-C. Kao and S.L. Hulbert (BNL)

Employing the technique of Auger-photoelectron coincidence spectroscopy, we have studied thin (about 7 Å thick) films of SiO₂ grown on the Si(111) surface by thermal oxidation. For such thin oxides, there is a significant contribution to the Si L₂₃VV Auger spectrum from substoichiometric interface oxides. We have obtained the Si L₂₃VV Auger spectrum in coincidence with the Si 2p core level associated with stoichiometric SiO₂, thereby eliminating from the Auger spectrum any contribution from substoichiometric regions of the sample.

The Si 2p core level spectrum obtained from a 7 Å film of SiO₂ on Si(111) is shown in Fig. 1. The open circles are the background subtracted data. The solid line is a fit to the data assuming a series chemically shifted Si 2p core levels associated with substoichiometric oxides as proposed by Himpsel et al.¹ We find that our core level spectrum is well reproduced by this analysis.

Coincidence Auger spectra presented below are obtained in coincidence with photoelectrons with a kinetic energy of 41.85 ± 0.5 eV. As seen from this figure, these photoelectrons come exclusively from Si sites in stoichiometric SiO₂.

The background subtracted spectrum is presented in Fig. 2. The dashed curve shows the results of a calculated spectrum by Ramaker.² The calculation reproduces the major peaks in the data at 66 eV and 78 eV quite well. However, the spectral weight is severely underestimated near 68 eV and at threshold. The additional spectral weight at threshold has been observed previously in conventional Auger studies and alternatively attributed to substoichiometric oxide generated by beam induced damage, or from Si orbitals of d-character participating in the Auger transition. This latter interpretation is favored by our results since we have eliminated any contribution from suboxide species through the coincidence experiment. The additional intensity near 68 eV has been attributed either to correlation effects in the 2-hole final state or to shake-up generated in the creation of the core hole. Having specified the kinetic energy of the outgoing electron in the coincidence measurement, the possibility of shake-up participation is eliminated. Therefore we assign the origin of this feature to a correlation satellite, in agreement with a CI calculation performed by Ramaker.²

1. F.J. Himpsel, F.R. McFeely, A. Taleb-Ibrahimi, J.A. Yarnoff, and G. Hollinger, Phys. Rev. B **38**, 6084 (1988).
2. D.E. Ramaker, Phys. Rev. B **21**, 4608 (1980).

This work was supported by NSF grant No. DMR89-07553

ELECTRONIC STRUCTURE OF THE SiO₂/Si(111) INTERFACE PROBED BY APECS

A.K. See and R.A. Bartynski (Ruigers U.), C.-C. Kao and S.L. Hulbert (NSLS)

We have obtained Auger-photoelectron coincidence spectra from thin films of SiO₂ grown on the Si(111) surface by thermal oxidation. The oxide was about 7 Å thick as determined by the Si 2p core level photoemission spectrum. The Si L₂₃VV Auger spectrum was measured in coincidence with oxide shifted core levels associated with the substoichiometric SiO_x at the semiconductor/oxide interface. The core level spectrum in this region (~2 eV greater binding energy than the substrate Si 2p_{3/2} level) is composed of overlapping signals associated with Si¹⁺, Si²⁺, and Si³⁺ oxidation states, but is dominated by the Si²⁺ contribution.¹

The figure below compares the noncoincidence, or singles spectrum (dashed line) to the Auger spectrum obtained in coincidence with Si 2p photoemission from the substoichiometric oxide. Several important observations can be made from this spectrum. The broad emission feature peaked near 85 eV accounts for the similar feature observed near that energy in the singles spectrum. However, a previous APECS study¹ of the Si L₂₃VV Auger spectrum from stoichiometric SiO₂ shows that emission in this energy region is not exclusively from suboxide species. In addition, the emission rapidly decreases above 90 eV, however, significant emission is observed up to an energy of 95 eV. From energetic considerations, emission from the top of the SiO₂ valence band occurs at 92 eV. The emission above this energy range is associated with states in the oxide band gap generated by the substoichiometric regions of the oxide. However, these states overlap the Si valence band and therefore do not generate traps in the Si band gap. Finally, there is a small emission feature observed between 95 and 98 eV which is accounted for by emission from states at the Fermi level. These observations are consistent with theoretical calculations of the electronic structure of defects in SiO₂ and the SiO₂/Si interface performed by Barrio et al.²

1. A.K. See, R.A. Bartynski, C.-C. Kao and S.L. Hulbert, previous abstract.
2. R.A. Barrio, R.J. Elliott and A.S. Carro, Phys. Rev. B **34**, 879 (1986).

Kinetic Energy (eV)

*Supported by SEMATECH

PHOTOEMISSION INVESTIGATION OF THE CORE LEVELS OF Na/Cu(111)

X. Shi, D. Tang, and D. Heskett (Univ. of R.I.); and K.-D. Tsuei (BNL)

We have performed angle-integrated photoemission measurements of the core levels of the system Na/Cu(111). We have measured the coverage-dependence of the Na 2p and 2s cores from submonolayer to multilayer coverages using photon energies of 70 eV and 120 eV for the 2p and 2s cores, respectively. In Figure 1 is plotted our results for the binding energy of the Na 2p core level as a function of coverage. The Na coverage was determined by a correlation of evaporation times, LEED, and work function measurements from a previous investigation of this system¹. For both the 2p and 2s cores, we observed a monotonic decrease in binding energy as a function of increasing coverage in the first layer. The magnitude of this decrease is approximately 0.85 eV and 0.66 eV for the 2p and 2s cores, respectively.

For coverages greater than ~0.44 (the completion of the first Na layer), a second 2p core level is detected which is associated with the second Na layer. As depicted in Figure 1, this peak shifts slightly to higher binding energy for increasing coverage in the second layer, while the first layer peak continues to shift to lower binding energy. For the Na 2s core level, only a single peak was detected for multilayer coverages, probably due to the poorer resolution used in these measurements. We interpret our results as supporting a covalent model of alkali adsorption, as, for example, recently suggested by the calculations of Ishida². In addition, we believe that the binding energy shifts of alkali core levels as determined here and in similar studies of other systems is most likely caused by final state, screening effects.

1. D. Tang, D. McIlroy, X. Shi, C. Shu, and D. Heskett Surf. Sci. 255, L497 (1991).
2. H. Ishida, Surf. Sci. 242, 341 (1991).

U15

FRAGMENTATION OF GASEOUS FREONS: ELECTRON-MULTIPLE-ION COINCIDENCE MEASUREMENTS*

T.A. Ferrett, S. Hsieh, M. Bayne (Carleton College) and D.Y. Kim, K. Lee, C.I. Ma, and D.M. Hanson (SUNY, Stony Brook)

Synchrotron radiation from U15 was used to selectively excite/ionize the F 1s core electrons in the gaseous freon series CF₃Cl, CF₂Cl₂, and CFCI₃. The wavelength dependence of subsequent fragmentation patterns was investigated below the F 1s threshold, on F 1s core-excited resonances, and in the F 1s continua. Time-of-flight ion mass spectra and ion coincidence maps were collected by triggering off electrons detected with a retarding potential analyzer.^{1,2} The resulting ion spectra represent fragmentation events which follow emission of both high and low energy electrons. The electron analyzer is particularly sensitive to low-energy electrons, thus permitting experiments which probe fragmentation events induced by multiple-electron emission.

Preliminary examination of the ion spectra and coincidence maps reveals several trends: 1) the production of many atomic ion fragments (>60% are atomic), with no detectable parent ions surviving on the microsecond timescale, 2) the predominance of F-containing molecular fragments in all three freons despite the variation in relative numbers of Cl and F in the parent molecules, 3) the presence of very sharp peaks in some mass spectra, indicating fragmentation with neutrals, and 4) subtle but clearly evident changes in the mass spectra with wavelength. Further analysis of peak intensities and shapes will provide information about the kinetic energy of the ion fragments and the relative importance of particular types of fragmentation channels versus incident photon energy.

¹ D.M. Hanson, C.I. Ma, K. Lee, D. Laipano-Smith, and D.Y. Kim, J. Chem. Phys. **93**, 9200 (1990).

² J. Murakami, M.C. Nelson, S.L. Anderson, and D.M. Hanson, J. Chem. Phys. **85**, 5755 (1986).

* This work was supported by Carleton College and a William and Flora Hewlett Foundation Award of Research Corporation.

ELECTRON-ION COINCIDENCE MASS SPECTRA OF CO

R. G. Hayes (U. Notre Dame), K. Lee, C. I. Ma, D. M. Hanson (SUNY at Stony Brook)

We have studied the behavior of carbon monoxide molecules upon O^{1s} electron excitation by electron-ion coincidence experiments. The Auger electron-ion coincidence spectra were particularly informative. One of these spectra appears in the figure. This mass spectrum is in coincidence with Auger electrons in the kinetic energy range 465-475 eV. It represents primarily the behavior two states of CO²⁺, which lie 37.2 eV and 39.3 eV above the energy threshold for fragmentation to C⁺ + O⁺.

The spectrum shows that the kinetic energies of C⁺ and of O⁺ are strongly peaked. The maximum kinetic energy in the C⁺/O⁺ pair is 21. eV. Thus, the atomic ions produced by fragmentation must be strongly excited electronically, since there is 16-18 eV to be accounted for.

The kinetic energies of the dipositive ions are much more broadly distributed. O²⁺ has a maximum kinetic energy of 11. eV. The threshold for production of C + O²⁺ is 59.8 eV. Thus, the ground states of O²⁺ and C may be produced. O²⁺ is also produced with less kinetic energy, though, which requires that at least one of the fragments be excited.

C²⁺ is produced with a maximum kinetic energy of 8.4 eV. The threshold for C²⁺ + O is at 46.8 eV, so the products must be highly excited if this is the process which produces C²⁺.

ELECTRON-ION COINCIDENCE MASS SPECTRA OF NH₃

R. G. Hayes (U. Notre Dame), K. Lee, C. I. Ma, D. M. Hanson (SUNY at Stony Brook)

We have studied the behavior of ammonia molecules upon N 1s electron excitation by electron-ion coincidence experiments. The Auger electron-ion coincidence spectra were quite informative, partly because the Auger spectrum of ammonia is sparse, so each spectrum arose primarily from only two states of the NH₃²⁺ ion which Auger decay produces. One of these spectra appears in the figure. This mass spectrum is in coincidence with Auger electrons in the kinetic energy range 369-379 eV. It represents the behavior of the ground singlet state of NH₃²⁺, ¹A₁, and of the first excited state, ¹E. ¹A₁ of NH₃²⁺ lies 31.6 eV (vertical energy) above the ammonia ground state and ¹E lies 4.6 eV higher. The threshold for fragmentation to NH₂⁺(¹A₁) + H⁺ lies at 30.6 eV. The threshold for fragmentation to NH⁺ + H⁺ + H lies at 35.6 eV.

The figure shows NH₃²⁺ and NH₂⁺ in intensity ratio of 2:1 and little NH⁺. The H⁺ feature has a component with large kinetic energy, a maximum of 6.4 eV, and a component with smaller kinetic energy, a maximum of 0.9 eV.

NH₃²⁺ comes mostly, perhaps exclusively, from the ¹A₁ state. NH₂⁺ has a maximum kinetic energy which associates it with the large kinetic energy component of H⁺, thus it comes from ¹E, and essentially all the electronic excitation energy of shows up as fragment kinetic energy. A number of pathways for production of NH⁺ and the small kinetic energy component of H⁺ can be imagined, all of which originate from ¹E of NH₃²⁺. The small kinetic energy component of H⁺ can not come from fragmentation of some of NH₃²⁺(¹A₁) because this would produce NH₂⁺ with small kinetic energy, which is not observed.

ELECTRON-ION COINCIDENCE STUDIES OF THE BEHAVIOR OF AMMONIA UPON CORE ELECTRON EXCITATION

R. G. Hayes (Norre Dame); K. Lee, C.-I. Ma and D. M. Hanson (SUNY Stony Brook)

We have observed electron-ion coincidence mass spectra from individual Auger features of ammonia upon N 1s electron ionization and upon excitation of a N 1s electron to a 3p Rydberg state. The figure, which shows the mass spectrum from the first excited singlet electronic state (1E) of NH_3^{2+} , is representative of our data. This state lies 8.6 eV above the threshold for fragmentation of NH_3^{2+} into $\text{H}^+ + \text{NH}_2^+$. The presence of NH_3^{2+} in the spectrum, thus, argues against rapid equilibration of the electronic excitation of the dication. The fragmentation which is observed releases 5.7 eV as kinetic energy of $\text{H}^+ + \text{NH}_2^+$, so NH_2^+ is left with 2.9 eV of internal energy. This is enough to populate any of the three lowest singlets of NH_2^+ . All three of these states connect adiabatically with 1E of NH_3^{2+} .

Work supported, in part, by Research Corp. (Grant R-60 to RGH)

ELECTRON-ION COINCIDENCE STUDIES OF THE BEHAVIOR OF METHYLAMINE UPON CORE ELECTRON EXCITATION

R. G. Hayes (Norre Dame); K. Lee, C.-I. Ma and D. M. Hanson (SUNY Stony Brook)

We have observed electron-ion coincidence mass spectra from individual Auger features of methylamine upon N 1s electron ionization. The figure shows the mass spectrum from the Auger transition of methylamine which produces electrons with the greatest kinetic energy. This transition leaves the methylamine dication in an electronic state (or states) which lies 28.0 eV above the ground electronic state of methylamine. This is 4.2 eV above the energy threshold for the fragmentation of $\text{CH}_3\text{NH}_2^{2+}$ to $\text{CH}_4\text{N}^+ + \text{H}^+$. The observed kinetic energy release of the fragmentation is near this energy, as may be obtained from the maximum kinetic energy of H^+ . One notes, though, that a significant amount of the dication is metastable. Also, the fragmentation to $\text{H}_2\text{CN}^+ + \text{H}_3^+$ is prominent. This fragmentation also releases 4.2 eV.

Work supported, in part, by Research Corp. (Grant R-60 to RGH)

THE ANISOTROPIC FRAGMENTATION OF CO₂, N₂O, AND H₂O FOLLOWING CORE ELECTRON EXCITATION.

D. Y. Kim, K. Lee, C. I. Ma, and D. M. Hanson (SUNY at Stony Brook).

It is demonstrated that the fragmentation of the triatomic molecules following core electron excitation is strongly anisotropic. Molecules of different symmetries, CO₂ which is in the D_{∞h} group, N₂O which is in the C_{∞v} group, and H₂O which is in the C_{2v} group, have been studied. Experimentally, two detectors positioned 0 degrees and 90 degrees with respect to the light polarization measure ion yields simultaneously. Application of a retarding potential, energy selects the ions. The polarization of the synchrotron radiation is determined from the π resonance of N₂, and the collection efficiencies of the two detectors as well as geometric factors are calibrated from CH₄. The ion yields and the anisotropy parameters at the oxygen edge of CO₂, N₂O, and H₂O are shown in the figure below. Results at different retarding potentials, at the other atomic edges for these molecules, and for acetone and CS₂ have been obtained but are not reported here.

Figure. The top spectra are the normalized ion yield spectra measured with +2.0 V retarding potential. The bottom spectra are the anisotropic parameters, β , spectra determined from the ion yield data.

Acknowledgement

This research was supported by the National Science Foundation (CHE-8921729).

LOCAL ENVIRONMENT AROUND THE OXYGEN ATOMS IN Y-Ba-Cu-O STUDIED BY X-RAY ABSORPTION FINE STRUCTURE (XAFS) SPECTROSCOPY

BEAM LINE U15

A. Krol, Z. H. Ming, and Y. H. Kao (SUNY/Bufalo), N. Nücker, G. Roth, and J. Fink (Kernforschungszentrum Karlsruhe, Germany), G. C. Smith (BNL)

The local environment about oxygen atoms in the CuO₂ plane in YBa₂Cu₃O₇ twin free single crystals has been investigated by orientation-dependent x-ray absorption fine structure (XAFS) spectroscopy at the O 1s edge using bulk-sensitive fluorescence-yield-detection method at 90 K. By choosing the orientation of the x-ray electric E -vector parallel to the a -crystallographic axis of the single crystal one can selectively probe bonds formed by O(2) atoms with its Cu(2) neighbor in the CuO₂ plane. The dominant peak in the magnitude of Fourier transform, between 0.8 and 2.4 Å is due to XAFS interference from O(2) - Cu(2), O(3) - Y and O(2,3) - O(3,2) pairs. In order to spectrally isolate these first three shells the XAFS spectrum was backtransformed using window shown in Fig. 1. The resulting $\chi(k)$ function is shown in Fig. 2 as open circles. A theoretical XAFS was then fitted and the result of this procedure is represented by solid line. The obtained bond length is practically unchanged within error limits ($\sim \pm 0.03$ Å) as compared to crystallographic data at room temperature. The amplitude reduction factor S_0^2 for small k ($3.5 \leq k \leq 8 \text{ Å}^{-1}$) due to oxygen shake-up and shake-off effects can be estimated as equal to 0.87 ± 0.15 . The obtained Debye-Waller factors σ^2 (in 10^{-3} Å^2 units) are 6.5, 2.0 and 27.0 for first three shells, respectively, which are close to the results obtained for polycrystals. *Within experimental accuracy no structural change in local ordering about oxygen atoms in the CuO₂ plane at 90 K (i.e. in superconducting state vs. normal state) has been detected.*

Fig. 1. Magnitude of Fourier transform and the window used to backtransform the contribution of the first 3 shells.

Fig. 2. Isolated XAFS contribution of the first 3 shells (open circles) vs. fit (solid line). The contribution from O(2)-Cu, O(3)-Y and O(2,3)-O(3,2) shells is also shown.

ANGULAR DEPENDENCE OF PHOTOFRAGMENTS FROM K-SHELL EXCITED DIATOMICS

K. Lee, C. I. Ma, D. Y. Kim, D. M. Hanson (SUNY at Stony Brook)

Ion yield spectra for each of the molecules N_2 , O_2 , NO, and CO around their K-edges were recorded simultaneously by two retarding field analyzers with their axes oriented perpendicular (90°) and parallel (0°) to the electric vector of the synchrotron radiation. In the figures of relative ion yields shown below, the solid curves represent the 90° spectra whereas the dashed curves denote the 0° spectra. The spectra of the calculated photoabsorption anisotropy parameter β based on these ion yields display a general pattern in that they all exhibit a pre-edge π^* resonance ($\beta = -1$) and a σ^* resonance ($\beta > 0$). In addition, the qualitative features of the β spectrum of NO at the N K-edge are similar to those of N_2 , whereas those of NO at the O K-edge resemble those of O_2 .

This research was supported by the National Science Foundation (CHE-8921729)

LOCAL STRUCTURE AROUND F IN $Nd_{2-x}Cu_{1+x}F_2$ SUPERCONDUCTORS

A. Krol, Y. L. Soo, Z. H. Ming, S. Huang and Y. H. Kao, (SUNY Buffalo); G. C. Smith, (Brookhaven National Laboratory); K. Lee, (SUNY Stony Brook); A. C. W. P. James, and D. W. Murphy, (AT&T Bell Laboratories, Murray Hill).

Soft x-ray absorption fine structure (XAFS) measurements were performed on $Nd_{2-x}Cu_{1+x}F_2$ electron superconductors with $x = 0.15$ and 0.30 , respectively. The magnitude of the Fourier transforms of the XAFS data multiplied by k^2 (denoted $FT[k^2\chi(k)]$) obtained for the NdF_3 and Cu_2O model compounds, the $Nd_2Cu_{3.85}F_{7.15}$ and the $Nd_2Cu_{3.7}F_{7.0}$ superconductors, and NdOF are shown in Fig. 1 (curves a, b, c and d, respectively). The analysis consisted of the spectral isolation of the contribution from the first shell and backtransformation into the k space. Backscattering amplitudes of the $k\chi_2(k)$ obtained for the investigated superconductors resembled very closely that obtained for the NdF_3 compound at the F K edge and disagreed with the backscattering amplitude obtained for a Cu_2O sample at the O K edge (see Fig. 2 with the same notation as in Fig. 1). This finding indicated that the first-neighbor shell consisted predominantly of the Nd atoms. The single shell XAFS interference function $k^2\chi_2(k)$ was subsequently compared with the NdF_3 model compound single shell XAFS function $k^2\chi_{Nd}(k)$ by means of the ratio method. It is found that F atoms substitute exclusively on the O(2) sites in the NdO layers. The $F(2)$ site is slightly distorted with longer $F(2) - Nd$ bond length ($2.34 \text{ \AA}$) as compared to the $O(2) - Nd$ bond ($2.32 \text{ \AA}$). Observed coordination number is close to 4, as expected for the O(2) site. A symmetry of anionic (F) and cationic (Ce and Th) substitution in NdO layers is observed.

FIG. 1. This research at the State University of New York at Buffalo is supported by DOE under Grant No. DE-FG02-87ER45283 and at Brookhaven by DOE under Grant No. DE-AC02-76CH00016.

FIG. 2.

DETERMINATION OF THE HARMONIC COMPOSITION OF THE SYNCHROTRON RADIATION AT U15

K. Lee, D. M. Hanson (SUNY at Stony Brook), Y.H. Kao (SUNY at Buffalo)

To evaluate the degree of chromatic purity of the synchrotron radiation reaching the end station at U15, studies were made to resolve the photon flux into its harmonic components. Photons emanating from a toroidal grating monochromator (TGM) were reflected by one of a set of interchangeable mirrors and then detected by a parallel plate avalanche chamber, filled with a mixture of argon/C₂H₆, with a Mylar window. The amplified signal was pulse-height analyzed by a MCA. For a given mirror at each fundamental photon energy setting of the TGM, an energy resolved spectrum is thus obtained. Shown below are spectra for three mirrors (quartz, nickel, and gold), each at three fundamental photon energies (300, 400, and 500 eV), with peaks and shoulders due to different orders of light as labeled. After deconvolution and account for the detector window transmission and gas absorption, the harmonic composition for each spectrum can be obtained.

This research was supported by the National Science Foundation (CHE-8921729), by AFSOR (AFS0R-88-0095), and by the Department of Energy (DE-FG02-87ER45283).

ELECTRON-MULTIPLE-ION COINCIDENCE STUDIES OF METHANOL

C.-I. Ma, K. Lee, D. Y. Kim, and D. M. Hanson (SUNY at Stony Brook)

The fragmentation of methanol following carbon 1s to continuum excitation was studied by electron-multiple-ion coincidence mass spectroscopy. Photoelectrons, Auger electrons and secondary electrons from Auger electrons were collected to trigger the TOF spectrometer. The figure shows a coincidence map of events containing two detected ions and the total mass spectra were aligned with the map to show the respective coincident ion pairs. Those ions with flight times shorter than atomic hydrogen ion were spurious. H⁺ was collected in coincidence with C⁺, CH_n⁺, n = 1, 2, 3, O⁺, OH⁺, CO⁺, and CHO⁺. Both O⁺ and OH⁺ were in coincidence with C⁺ and CH_n⁺, n = 1, 2, 3. Other observed species include H₂⁺, H₃⁺, C²⁺, O²⁺, CH₃O⁺, CH₃OH⁺, and CH₃OH₂⁺.

This research was supported by the National Science Foundation (CHE-8921729).

U16A SPIN-POLARIZED ANGLE-RESOLVED PHOTOEMISSION STUDIES OF ULTRA-THIN Fe/W(110) FILMS

A.B. Andrews, G.A. Mulholland, B.S. Kim, J.L. Erskine (U. of Texas at Austin), and X. Zhang (Rice U.).

The prediction of novel magnetic properties at surfaces and interfaces has stimulated broad efforts to understand the physics of thin-film magnetism. Noble metal substrates have been adopted because of their small hybridization effects despite evidence that they provide non-ideal templates for 3d transition metals due to poor growth characteristics. The (110) face of bcc W affords excellent epitaxy as well as an opportunity to test the accuracy of LSDA calculations¹ for a strongly hybridized system.

The spin-resolved photoemission spectra were taken on the U5 beamline. Monolayer films showed only a very weak remnant polarization ~5% at $T = 100K = 5 T_C(1)$. Even symmetry exchange split bands mapped along the Σ line for 2 ml thick films showed polarizations of 30%. Spin-integrated spectra at 1 ml coverage were taken on the U16A beamline in both even and odd geometries.

Our normal emission spin-resolved spectra show similarities to Fe(110), but the dispersion of the $T_{25}^{\uparrow}$ and $T_{12}^{\uparrow}$ precursors is smaller than that reported for the bulk states². The exchange splitting is an average of 15% larger than found in the bulk³, indicating an enhanced moment in the thin film. One must be cautious in comparing the k dispersion seen in the thin film with the bulk bands since the lattice constant is stretched by 10%.

To our knowledge, this is the first example of enhanced exchange splitting in a strongly coupled 2D ferromagnetic system. The interesting result of a small in-plane polarization for the monolayer is in need of further investigation.

1. S.C. Hong, A.J. Freeman, and C.L. Fu, *Phys. Rev. B* **38**, 12156 (1988).
2. A.M. Turner, A.W. Donoho, and J.L. Erskine, *Phys. Rev. B* **29**, 2986 (1984).
3. K. Schroder, G.A. Prinz, K.H. Walker, and E. Kocker, *J. Appl. Phys.* **57**, 3660 (1985).

This work is supported by the NSF DMR89-22359 and the NSF DMR89-06935.

ARUPS STUDY OF BULK Gd(0001)

Bongsoo Kim, A.B. Andrews, J.L. Erskine (U. of Texas at Austin), Kwangjoo Kim and B.N. Harmon (Iowa State U.)

We have carried out angle-resolved photoemission studies of the electronic properties of bulk Gd(0001) along the Γ -A direction of the three-dimensional Brillouin zone (corresponding to normal emission). An electropolishing technique was used to clean (and provide a passivated chloride layer on) the surface which was removed by *in situ* sputter-annealing cycles in the UHV chamber.

Figure 1 displays normal emission angle-resolved photoemission spectra vs. photon energy at 80K. The Γ_2' point was inferred from a maxima in the initial state energy (~1.2 eV) at $h\nu=35$ eV and $k_{\parallel}$ was determined by applying the direct transition model assuming free electron final state bands and an inner potential of 10.8 eV.

Figure 2 displays the $k_{\parallel}$ dependence of binding energies of the two bulk band features in the EDCs plotted over solid lines representing our calculated bands¹. These states are interpreted as the exchange split states of predominantly 6s character, which manifest increasing 5d_{3/2} contributions near Γ . The intense peak at ~0.25 eV binding energy lies in a gap of the projected bulk bands and does not disperse with $k_{\parallel}$. This feature is produced by a surface state which has been predicted theoretically², and which has also been observed by photoemission using Gd films grown on W(110)³.

Previous attempts⁴ to measure band dispersion on bulk rare-earth metals have met with limited success. While the agreement between our experimental results and calculations can only be regarded as fair, it is clear that photoemission experiments will be capable of testing LSDA calculations in the rare-earth metals where spin-orbit coupling, magnetic ordering, and strong correlations are expected to play an important role.

References

1. K. Kim and B.N. Harmon, unpublished.
2. R. Wu, C. Li, A.J. Freeman, and C.L. Fu, to appear in *Phys. Rev. B* **44** (1991).
3. D. Li, C.W. Hutchings, P.A. Dowben, C. Hwang, R.T. Wu, M. Onellion, A.B. Andrews, and J.L. Erskine, to appear in *J. Mag. Mag. Mat.*
4. F.J. Himpsel and B. Reihl, *Phys. Rev. B* **29** (1983) 574.

Research sponsored by NSF/DMR-8922359 and by the DoE (NSLS).

U16A

ARUPS TEMPERATURE DEPENDENCE OF RKKY INDUCED MAGNETIC SPLITTING IN BULK Gd(0001)

Bongsoo Kim, A.B. Andrews, J.L. Erskine, (U. of Texas at Austin), Kwangjoo Kim and B.N. Harmon (Iowa State U.)

Gd is a prototype Heisenberg spin system in which the magnetic ordering is achieved by RKKY indirect exchange between non-overlapping 4f states via conduction electrons. Early non-self-consistent calculations^{1,2} concluded that the 4f coupling to the 5d electrons alone dominated the exchange mechanism; more recent self-consistent relativistic calculations³ suggest the s and d electrons contribute on a more equal footing. The most recent⁴⁻⁷ calculational efforts include spin-orbit coupling and treat 4f electrons in narrow-band states.

Figure 1 displays normal emission EDCs for Gd(0001) at $h\nu=38$ eV as a function of temperature. $T=295$ corresponds to a temperature just above the Curie point of bulk Gd ($T_c=293$ K), $T=80$ K corresponds to $T/T_c=0.35$. In the accompanying abstract, the two peaks which exhibit temperature dependence were shown to agree reasonably well with LSDA calculations of the exchange split s band along $\Gamma-A$.

Figure 2 displays the temperature dependence of the exchange splitting obtained from curve fitting the spectra in Fig. 1. The experimental data has been fit using a simple power law yielding a critical exponent $\beta=0.378$ which agrees very well with the result obtained for bulk Gd based on measurements of the magnetization. This behavior is a strong indication that the splitting is of magnetic origin and is consistent with the RKKY mechanism for magnetic ordering.

References

1. J.O. Dimmock and A.J. Freeman, *Phys. Rev. Lett.* **13** (1964) 750.
2. B.N. Harmon and A.J. Freeman, *Phys. Rev. B* **10** (1974) 1979.
3. B.N. Harmon, *J. Physique Coll.* **40** (1979) C5 65.
4. J. Sticht and J. Kubler, *Solid State Comm.* **53** (1985) 529.
5. B.C.H. Kuitzen and F. Springelkamp, *J. Phys. Cond. Matt.* **1** (1990) 8369.
6. W.M. Temmerman and P.A. Sterne, *J. Phys. Cond. Matt.* **2** (1990) 5529.
7. K. Kim and B.N. Harmon, unpublished.

Supported by NSF/DMR-8922359 and DoE Light Source.

HIGH-RESOLUTION CORE-LEVEL SPECTROSCOPY OF EPITAXIAL BI-METALLIC INTERFACES

Bongsoo Kim, N.D. Shinn (Sandia National Laboratories), A.B. Andrews, and J.L. Erskine (U. of Texas at Austin)

High-resolution core-level spectroscopy is a powerful probe of chemical bonding at surface and interfaces. For epitaxial bi-metallic interfaces, previous studies¹ have shown that the surface W_{4f} peaks exhibit overlayer-specific chemical shifts to higher binding energies, yet remain distinct from the bulk W_{4f} peaks. We have exploited these findings to study the nucleation, growth, and thermal stability of epitaxial Fe overlayers on $W(110)$.

Figure 1 shows a series of $W_{4f} 7/2$ spectra as a function of Fe coverage in monolayers. One monolayer is defined as the atomic density in the $W(110)$ plane. The well-resolved surface peak at $\Delta E=320$ meV is seen to broaden and shift to higher binding energies with increasing Fe coverage at 300K. The spectrum for the complete monolayer, and subsequent thicker layers, shows a single shifted component at $\Delta E=156$ meV. This corresponds to the W atoms at the (1×1) Fe/W(110) interface.

These data and other data have been modelled by monte carlo simulations of 2-D Fe overlayer growth. A discrete binding energy is chosen for the interface/surface W atoms depending upon their coordination number. The results of one model which reproduces the experimental data well are shown in Figure 2. This experimental technique and model are being used in a complete systematic study of (1×1) bi-metallic interfaces on W surfaces to better understand nucleation and growth kinetics as well as chemical bonding at interfaces.

This work is supported by the NSF DMR89-06935 and by the U.S. Department of Energy, BES, Division of Materials Sciences under contract DE-AC04-76DP00789 with Sandia National Laboratories.

- 1 N.D. Shinn, C.H.F. Peden, K.L. Tsang, and P.J. Berkowitz, *Physica Scripta* **41**, 607 (1990).

ANGLE-RESOLVED PHOTOEMISSION EVIDENCE FOR A Gd(0001) SURFACE STATE

D. Li (Syracuse U.), C.W. Hutchings (U. Heidelberg), P.A. Dowben (Syracuse U.), C. Hwang, R.-T. Wu, M. Onellion (U. of Wisconsin), B. Andrews and J.L. Erskine (U. of Texas)

We have used angle-resolved photoemission to study the electronic properties of Gd(0001) thin film crystals grown on W(110). We observe a prominent surface state at $\bar{\Gamma}$ having A_1 symmetry and a binding energy of approximately 0.2 eV, values in agreement with recent first principles calculations. We also study the emergence of 3d band structure with film thickness, the dispersion of 2d bands of the thin films including the surface state, and examine lifetime effects.

Figure 1. Normal emission EDCs for Gd(0001) vs film thickness. The large peak near E_F is a surface state. Peaks around 8 eV are 4f levels.

Figure 2. 2D band structure and energy half width for Gd(0001) bands along $\bar{\Gamma} - \bar{M}$.

This work was supported by NSF DMR89-22359.

TEMPERATURE DEPENDENCE OF CORE-LEVEL BINDING ENERGIES OF W(110) AND Ta(100)

D. M. Riffe, W. Hale, B. S. Kim, and J. L. Erskine (*The University of Texas at Austin*)
Recent photoemission data from the core-levels of In and four alkali metals (Li, Na, K, and Rb) show a temperature dependent shift in the binding energy (BE) of up to 63 meV between 77 and 295 K [1]. In principle, there are contributions to the shifts in both the initial and final states. In the initial state both the core-level eigen-value and the Fermi level decrease due to the expansion of the valence charge as the lattice expands with temperature. In the final state the relaxation energy decreases with increasing temperature, also due to the expansion of the valence charge. For the alkali metals the dominant effect appears to be a reduction in the core-level eigen-energy relative to the Fermi level which results in the positive BE shift. Interestingly, the surface and bulk BEs shift by essentially the same amount.

In order to further elucidate the mechanisms responsible for these thermal shifts in metals we have undertaken an investigation of the thermal shifts of surface and bulk core-levels from transition metals. The figure illustrates the changes in the 4f_{7/2} spectrum of the Ta(100) surface at 80 and 295 K. In contrast to the alkali metals, the surface core-level shift (SCS) is clearly seen to decrease over this temperature range. Surprising are the observations that the shift is dominated by motion of the bulk (second layer and deeper) line relative to the Fermi level and that the bulk and surface lines shift in opposite directions. From lineshape analysis, the shift of the bulk level BE is determined to be ~15 meV while that of the surface is ~6 meV so that the change in SCS is ~21 meV. On W(110) much smaller shifts are observed. Like Ta(100) the bulk line is observed to also move more than the surface line relative to the Fermi level. In this case the bulk line decreases in BE, but since the SCS of W in the opposite direction to Ta, the magnitude of the SCS is again observed to decrease as the temperature increases.

Qualitatively, the motion of the core levels for these transition metals with increasing temperature can be simply understood. As the lattice expands both the surface and bulk core level eigen-energies must approach the core-level position of the free atom. Hence, the binding energies of the bulk and surface must come together as the temperature is raised. Additionally, since the surface atoms are already more atomic-like than the bulk, their motion should be slower than that of the bulk. The unmeasurable difference in the surface and bulk BE shifts of the alkalis is also understandable since from work-function calculations one can deduce that the surface atoms of a simple metal have a valence charge density which more closely resembles their bulk counterparts more than do the surface atoms of a transition metal.

[1] D.M. Riffe, G.K. Wertheim, D.N.E. Buchanan, and P.H. Citrin (submitted to Phys. Rev. B).

V. Chakarian, D.K. Shuh and J.A. Yarnoff, *Dept. of Physics, U. of California, Riverside, CA*
H.-S. Tao, B. Maschhoff and T.E. Madey, *Dept. of Physics, Rutgers, The State U. of New Jersey, Piscataway, NJ*
N.D. Shinn, *Sandia National Laboratories, Albuquerque, NM*

The preadsorption of alkalis on metal surfaces is known to promote chemical reactions. It has been observed that molecules that normally adsorb molecularly, when coadsorbed with alkalis, either dissociate or experience bond weakening. Many of the properties of these coadsorbed systems are similar for numerous metal substrates, alkali metals, and molecules, such as CO, NO, and PF₃, suggesting that the processes involved are rather fundamental in nature.

The adsorption of PF₃ alone on Ru(0001) has been previously studied by employing a variety of surface spectroscopic techniques.¹ The chemisorption of the PF₃ on Ru(0001) pre-exposed to a submonolayer of potassium at room temperature has been reported previously.² In this report the results of soft x-ray photoemission experiments aimed at determining the chemical state of the adsorbates for various PF₃ coverage regimes at liquid nitrogen (LN₂) temperatures are presented.

In the absence of potassium PF₃ adsorbs molecularly at room temperature as well as at LN₂ temperature. The preadsorption of potassium induces dissociative chemisorption at both room temperature and LN₂ temperature but the details of the reaction are different at different temperatures. At the initial stages of exposures PF₃ completely dissociates in both cases. However, differences are seen between the chemisorption at room temperature and the LN₂ temperature for higher PF₃ exposures. The P 2p core level photoemission at room temperature and at LN₂ temperature are shown in the Figure below where the PF₃ exposures indicated correspond to the saturation coverage in each case. At room temperature and at LN₂ temperature alike, changes in the K 3p core level lineshape (not shown) are seen, indicative of the direct involvement of potassium in the dissociation process.

- 1 S.A. Joyce, A.L. Johnson, J.A. Yarnoff, and T.E. Madey, to be published.
- 2 V. Chakarian, D.K. Shuh, J.A. Yarnoff, H.-S. Tao, B. Maschhoff and T.E. Madey, *J. Chem. Phys.*, to be published.

R. P. Merrill,* T. N. Rhodin,* S. Woronick,* P. O'Hagen,* N. D. Shinn,** R. Absi*** and A. Chester****
*Cornell Materials Science Center, Cornell University, **Sandia National Laboratories, ***Mobil Research and Development

Analysis of valence band spectra of epitaxial alumina provides significant information on the effect of heat treatment and ethylene adsorption on changes in chemical bonding and structure resulting from heat treatment. A typical valence band spectrum for the water-grown oxide is shown in Fig. 1a. The valence band consists of two broad peaks. The upper valence band is identified with O_{2p} states and the lower valence band with O_{2s} states. States of low binding energy (BE) for each peak consist of flat nonbonding oxygen orbitals while the deeper bands are O_{2p}/Al_{2p} and O_{2s}/Al_{2s} hybrid bands, respectively. The large dispersion in these hybridized states and the level of charge transfer suggest that the Al-O interaction has significant covalent character. The difference spectra for the high temperature heat treatments up to 450°C minus the low temperature valence band spectra are shown in Fig. 1b. The loss and gain shown at 3 and 7 eV, respectively, indicate a shift in the density of states (DOS) associated with stabilization of the Al-O hybridized states accompanying the change from amorphous to crystalline alumina.

Additional changes are observed when ethylene is adsorbed on the oxides pretreated at various temperatures (Fig. 1c). During this chemisorption much of the hydroxyl-derived DOS in the upper valence band, lost during the high temperature treatment, is restored because of reversibly adsorbed water from the background gas. The DOS region around 22 eV of the lower valence band also increases. The features in the upper valence band at 5 and 6 eV are attributed to ethylene-derived states. While the observation of the 2p_{3/2} molecular orbital of ethylene is hindered by the co-adsorbed hydroxyl, both the 1b_{1g} and the 1b_{3g} molecular orbitals of the ethylene are clearly shown.

The losses and gains observed in these valence level spectra are completely consistent with the previously reported O_{1s} core-level spectra and support our previous interpretations which indicate that the maximum surface acidity develops from heat treatment at about 375°C. In addition, the features of the C_{1s} core level spectra (not shown) are also consistent with those derived from the alumina valence level spectra. It appears that changes in valence band spectra provide an excellent indirect indicator of modifications in crystallinity and local structure as well as identifying the electronic structure of the reactive adsorbate molecule. The conclusions on bonding and coordination remain to be substantiated by using direct structural probes such as Al_{2p} NEXAFS and EXAFS.

R. P. Merrill, T. N. Rhodin, S. Woronick, P. O'Hagen (Cornell U.), N. D. Shinn (Sandia), G. Woolery, R. Absil, and A. Chester (Mohr)

Aluminum oxides are important as catalysts and catalyst supports; hence understanding the role of composition, structure and chemical activity on catalytic effectiveness are of much interest. It is particularly important to have detailed information on the factors of local chemical bonding, surface acidity and crystallinity obtained in a correlated manner. Photoemission and absorption of soft x-rays are ideally suited for this purpose. It is also of interest to explore how UVV techniques on planar materials could be applied to commercial catalysts. The specific objective of this study was to investigate the influence of thermal treatment on the development of surface acidity in alumina overlayers grown epitaxially on single crystal aluminum. Dehydroxylation of the hydrated aluminas produced a combination of Brønsted and Lewis acid sites. An optimum heat treatment temperature for the development of surface acidity on our samples was observed between 350-400°C. The states in the O 1s spectra induced by the C₂H₄ adsorption also showed a maximum at 370°C. A typical O 1s photoemission spectrum for a sample heat treated at 300°C after ethylene adsorption at 120K is shown in Fig. 1 where the photoemission intensity is plotted as a function of binding energy. The majority state at 532.0 eV is consistent with the O 1s binding state reported for aluminum oxides. The next minority state at 533.7 eV has been assigned to hydroxyl species in aluminum oxide and the remaining minority state at 534.6 eV was assigned to a C₂H₄-derived state. The effect of heat treatment on surface acid sites for a range of temperatures from 300 to 450°C is summarized in Fig. 2. The surface compositions of oxide-, hydroxyl-, and C₂H₄-derived states are plotted as a function of treatment temperature. It is observed that the oxide composition goes through a maximum at 375°C, and, as expected, the hydroxyl content decreases monotonically with treatment temperature. The C₂H₄-derived species pass through a maximum of about 40% at 370°C.

1. Deconvoluted typical O 1s core level spectra after C₂H₄ exposure. (Sample heat treated at 300°C)

2. The density of states for the oxygen 1s photoemission vs. treatment temperature. Oxide states, O; C₂H₄ derived states, Δ; hydroxyl states prior to C₂H₄ adsorption, •; hydroxyl states after C₂H₄ adsorption at 120K, ▲; hydroxyl state after warming to 210K.

DECAY CHANNELS OF Al L_{2,3} EXCITONS AND ABSENCE OF O K EXCITONS IN α-Al₂O₃

W. L. O'Brien, J. Jia, Q.-Y. Dong and T. A. Callcott (U.Tenn), D. L. Mueller and D. L. Ederer (NIST), N.D. Shinn (SANDIA), S.C. Woronick (Cornell U.)

The Al L_{2,3} and O K thresholds for single crystal α-Al₂O₃ have been studied by photoemission. Sample charging was compensated for by use of an electron flood gun. Energy distribution curves (EDCs), constant initial state (CIS) and constant final state (CFS) spectra were obtained and compared to the absorption spectrum reported previously¹. An exciton appears as a doublet at threshold in the Al L_{2,3} CFS, CIS and absorption spectra. These are the first reported absorption measurements, using photoelectron techniques, which have resolved this doublet. The details of the Al L_{2,3} CFS spectrum and absorption spectrum are similar while the exciton is the only feature present in the CIS spectrum, Fig.1. The CIS spectrum (using the valence band as the initial state) shows a strong Fano resonance. Comparison of the various Al L_{2,3} spectra allow the probabilities of different exciton decay channels to be determined. The probability for nonradiative direct recombination of the exciton is found to be 8±1% and the probability for Auger decay of the exciton is found to be 72±20%. Comparison of the O K CIS and CFS spectra suggest that no O K exciton is formed.

Fig.1 CIS and CFS compared to absorption at Al L threshold

W.L. O'Brien, J.J. Jia, T.A. Callcott, J.-E. Ruebenisson, D.L. Mueller and D.L. Ederer, Phys. Rev. B, 44, 1013 (1991).

This work was supported by NSF-DMR-87-13430, and the NIST

OXIDATION OF W(110): W(4f) CORE-LEVEL SPECTROSCOPY

C. H. F. Peden and N. D. Shinn
Sandia National Laboratories, Albuquerque, NM 87185-5800

Oxygen chemisorption onto, and subsequent oxidation of tungsten (W) surfaces have been studied extensively. Because oxygen adsorption can lead to significant substrate structural changes, the specific O(1s) adsorbate geometries, coverages and electronic structure are very sensitive to the single crystal surface studied [1]. The initial stages of W(110) oxidation are quite complex owing, in part, to its open structure. In contrast, O₂ adsorption on the close-packed W(110) surface up to coverages of 1 monolayer (ML) results in, at most, minor changes in the positions of the surface W atoms.

We have characterized the stages of oxygen adsorption on W(110) surfaces using surface-sensitive W(4f) core-level photoemission, analogous to a recent study of the W(100) surface [1]. Shown below are representative W(4f)_{7/2} and W(4f)_{5/2} spectra for clean W(110) (1a), a saturation oxygen coverage at 300K (1b), and after high-temperature (1550K) O₂ reaction (1c). The clean surface spectrum (1a) exhibits the well-known features due to bulk W, and coordinatively unsaturated surface atoms (at 0.3 eV lower binding energy). At room temperature, oxygen adsorbs in islands of 1/2 ML local coverage (p(2x1) LEED structure). When fully developed ($\theta = 0.5$ ML), this structure is characterized by a surface W core level feature shifted by 0.3-0.4 eV to higher binding energy from the bulk W peak as shown in Fig. 1c. O(1s) coverages above 1/2 ML are conveniently obtained by exposures at higher substrate temperatures. This gives rise to a new W(4f) feature at 0.6-0.7 eV higher binding energy from the bulk peak, which saturates at $\theta_0 = 1.0$ ML (Fig. 1c).

Figure 1: W(4f)_{7/2} and W(4f)_{5/2} spectra for clean W(110) (1a), a saturation oxygen coverage at 300K (1b), and after high-temperature (1550K) O₂ reaction (1c). The features due to bulk W atoms are labelled in the figure (B).

[1] P. Alnot, D.J. Auerbach, J. Behm, C.R. Brundle and J. Viescas, Surf. Sci. 213 (1989) 1.

This work was supported by the U. S. Dept. of Energy, Office of B.E.S., Division of Materials Science, under contract No. DE-AC04-76DP00789 with Sandia National Laboratories.

THE STRUCTURE OF THERMALLY NITRIDED Si(100)

C. H. F. Peden, J. W. Rogers, Jr., N. D. Shinn, K. B. Kidd and K. L. Tsang
Sandia National Laboratories, Albuquerque, NM 87185-5800

Silicon nitride (Si₃N₄) ceramics and thin films are used extensively for commercial applications in the ceramics and microelectronics industries owing, in part, to their chemical stability, and desirable electrical and mechanical properties. Traditional methods for the formation of Si₃N₄ ceramic precursor powders and thin films include the direct thermal nitridation of silicon powders and single crystals.

Synchrotron photoemission measurements of the Si(2p) and N(1s) core-levels have been made on Si₃N₄ thin films grown in-situ by high temperature (> 800K) reaction of Si(100) with NH₃. The tunability of the excitation photons allowed the mean-free-path of the emitted photoelectrons to be varied, thereby providing the capability to determine not only the types of chemical species present, but their location within the film as well. Photoemission spectra of the Si(2p) region near 100 eV binding energy are shown in Fig. 1 for Si₃N₄ films thermally grown for various lengths of time in NH₃. In the figure, the raw data and the calculated best-fit to the data are illustrated together with the various Si(2p) spectral features that make up the fit. The four 2p_{3/2}/2p_{1/2} doublets, labeled A-D in the figure can be assigned to elemental Si (A), Si in stoichiometric Si₃N₄ (D), and Si in intermediate oxidation states with less than 4 bonds to N atoms in the structure (B,C). Fig. 2 is a plot of the integrated intensities of the various Si(2p) doublets as a function of nitride thickness. These results can be rationalized with a film structure as follows: the nitride film consists of an outer monolayer of Si atoms (C) on top of the growing, stoichiometric nitride film (D). There is also a significant quantity (1/2 ML) of interfacial Si atoms (B) between the nitride film (D) and bulk Si (A). Consistent with this film structure are the binding energies of the outer layer- and interfacial-Si atoms, both being intermediate between elemental (bulk) Si and Si₃N₄.

Figure 1: Si(2p) spectra ($h\nu = 130$ eV) as a function of NH₃ exposure for the direct thermal nitridation of Si(100) at substrate temperatures above 850 K.

Figure 2: Integrated intensities of the four Si(2p) doublets (A-D) shown in Fig. 1. The solid squares (—) are due to bulk Si (A), triangles (---) are interfacial-Si atoms (B), diamonds (---) are outer layer Si (C), and circles (---) are for the spectral features due to Si atoms in the stoichiometric nitride Si₃N₄ (D).

This work was supported by the U. S. Dept. of Energy, Office of B.E.S., Division of Materials Science, under contract No. DE-AC04-76DP00789 with Sandia National Laboratories.

**Synchrotron Image
Venous Catheter**

**6 months
Post-Angioplasty**

**Section II
X-ray Abstracts**

List of Abstracts: X-ray Beamlines

Beamline X1A

Improved Images with the Scanning Photoemission Microscope H. Ade, C.-H. Ko, and E. Anderson	103
Photon Beam Mediated Oxidation on Aluminum Surfaces H. Ade, C.-H. Ko, E. Jonahon, Janos Kirz, S. Hulbert.	103
Investigation of Calcium Deposits in Human Cartilage Using STXM C.J. Buckley, G.F. Foster, R.E. Y.S. Ali, C. Scotchford, and J. Kirz	104
Imaging of Wet, Unfixed and Unstained Muscle at 50nm Resolution G.F. Foster, C.J. Buckley, and R.E. Burge	104
Quantitative Measurements of Protein Flux Across the Zymogen Granule Membrane Using STXM K.K. Goncz and S.S. Rothman	105
Diffraction-Limited Imaging in a Scanning Transmission X-ray Microscope C. Jacobsen, S. Williams, J. Kirz, X. Zhang, E. Anderson, M.T. Browne, C.J. Buckley, D. Kern, and M. Rivers	105
Scanning Luminescence X-ray Microscopy C. Jacobsen, S. Lindaas, S. Williams, and X. Zhang	106
X-ray Holography Using Photoresists C. Jacobsen, S. Lindaas, and M. Howells	106
Differential Phase Contrast Imaging with the STXM G.R. Morrison, J.R. Palmer, C. Jacobsen, and S. Williams	107

Beamline X1B

Screening Power of Spectator Electrons: Observation of the Electronic Decay of Core Hole-Rydberg Electron States in N ₂ W. Eberhardt, J.-E. Rubensson, K.J. Randall, J. Feldhaus, A.L.D. Kilcoyne, A.M. Bradshaw, Z. Xu, P. Johnson and Y. Ma	107
Soft X-ray Spectroscopy at the X1 Undulator K.J. Randall, J. Feldhaus, W. Erlebach, A.M. Bradshaw, W. Eberhardt, Z. Xu, P.D. Johnson, and Y. Ma	108
Core Level Photoelectron Spectroscopy of Gas Phase Molecules at Vibrational Resolution K.J. Randall, J. Feldhaus, A.L.D. Kilcoyne, A.M. Bradshaw, W. Eberhardt, J.-E. Rubensson, Y. Ma, Z. Xu and P.D. Johnson	108
Decay of Vibrationally Selected Core Hole States in Small Molecules J.-E. Rubensson, W. Eberhardt, K.J. Randall, J. Feldhaus, A.L.D. Kilcoyne, A.M. Bradshaw, Z. Xu, P. Johnson and Y. Ma	109

Beamline X2

Microtomography of Calcium Deposits in Cartilage Tissue K.L. D'Amico, J.H. Dunsmuir, S.R. Ferguson and C.J. Buckley	109
Determination of the Spatial Distribution of Metal Atoms in a Supported Catalyst System K.L. D'Amico, J.H. Dunsmuir, S.R. Ferguson and G.B. Mcvicker	110
Observation of the Settling of Particulates in a Heavy Oil J.H. Dunsmuir, K.L. D'Amico and S.R. Ferguson	110

Beamline X3A2

Phase Determination for Macromolecular Crystals Using X-ray Multiple Diffraction S.L. Chang, H. King, M.T. Huang and Y. Gao	158
Phase Separation Kinetics of a Segmented Polyurethane B. Chu, Y. Li, and T. Gao.	111

Synchrotron SAXS Study of Mean-Field and Ising-like Critical Behavior of Poly(2-chloro-styrene)/Polystyrene Blends B. Chu, Q. Ying, K. Linliu, P. Xie, T. Gao, Y. Li, T. Nose, M. Okada, A. Darovsky, and W. Lenhert	111
Structural Changes Upon Electrochemical Insertion of Lithium in the 2212 BiSrCuO Superconducting Phase P. Coppens, P. Lee, Y. Gao, N.A. Fleischer, and J. Manassen	112
Lattice Instabilities and the Effect of Copper-Oxygen Sheet Distortions on Superconductivity in La_2CuO_4 -Superconductors: $\text{La}_{1.60-x}\text{Nd}_{0.40}\text{Sr}_x\text{CuO}_4$ M. Crawford, R. Harlow, E. McCarron, and W. Farneth	112
Morphology Studies of CHDI Urethane Elastomers by X-ray Diffraction Using Synchrotron Radiation C.R. Desper, B. Chu, and Y. Li	113
SAXS and WAXS Study of Polyurethanes Containing Polyethylene Soft Segments C.R. Desper, B. Chu, T. Gao, Y. Li, and C.A. Byrne	113
Single Crystal Anomalous Scattering Studies of Mixed-Valence Complexes A. Frost-Jensen, M. Pressprich, Y. Gao, and P. Coppens	114
Lattice Instabilities in Nd-Doped La_2CuO_4 : Presence of a PCCN Phase? R. Harlow, M. Crawford, E. McCarron, and W. Farneth	125
Lattice Instabilities and the Effect of Copper-Oxygen Sheet Distortions on Superconductivity in La_2CuO_4 -Superconductors: $\text{La}_{1.88-x}\text{Nd}_x\text{Sr}_{0.12}\text{CuO}_4$ R. Harlow, M. Crawford, E. McCarron, and W. Farneth	114
Crystallization and Melting Study of Poly(Aryl Ether Ketones) B. Hsiao, K. Gardner, D. Wu, and B. Chu	115
Time-Resolved Synchrotron Study of Poly(Aryl Ether Ketones): Crystallization and Melting B. Hsiao, K. Gardner, D. Wu, and B. Chu	115
Thermally-Induced Structural Changes in a Segmented Polyurethane Y. Li, J. Liu, B. Chu, A. Darovsky, and W. Lenhert	116
Protein Crystals for Physical Estimation of X-ray Triplet Phases F. Mo, B.C. Hauback, E. Husstad, M. Teeter, L.C. Sieker, and M.S. Lehmann	116
Structures of Alkali-Doped Fullerenes by Powder Diffraction P.W. Stephens, L. Mihaly, F. Diederich, K. Holczer, S.-M. Huang, R.B. Kaner, R.L. Whetten, J.B. Wiley, and P.L. Lee	117
Time-Resolved X-ray Scattering Studies of the Crystallization of Polyethylene Blends K. Tashiro, R.S. Stein, M.M. Satkowski, B. Chu, and Y. Li	117
Determination of the Structural Distortions Corresponding to the q_1 -Type and q_2 -Type Modulations in Niobium Triselenide S. van Smaalen, J.L. deBoer, A. Meetsma, H. Graafsma, H.-S. Sheu A. Darovskikh, and P. Coppens	118
Temperature Effect on the Small Angle X-ray Scattering of an Ionomer J. Wang, Y. Li, D.G. Peiffer, R.D. Lundberg, and B. Chu	118
Small-Angle X-ray Scattering from Microvoids in Amorphous Silicon-Based Semiconductors D.L. Williamson and Y. Chen	119

Beamline X3B1

Investigation of Microstructure of Interfaces in $\text{Si}/\text{Ge}_{0.2}\text{Si}_{0.8}$ Superlattices by Means of X-ray Reflectivity and X-ray Standing Wave Z.H. Ming, A. Krol, Y.L. Soo, Y.H. Kao, and K.L. Wang	119
--	-----

Beamline X5

The Tensor Force Between Quarks and the Deformation of Baryons G. Blanpied, M. Blecher, A. Caracappa, C. Djalali, M.-A. Duval, G. Giordano, S. Hoblit, M. Khandaker, C. Kistner, G. Matone, L. Miceli, W. K. Mize, B.M. Preedom, A.M. Sandorfi, C. Schaerf, R.M. Sealock, C.E. Thorn, S.T. Thornton, K. Vaziri, C.S. Whisnant, X. Zhao, and M.A. Moinester	120
---	-----

Beamline X6B

Anomalous X-ray Reflectivity Measurements on Ni/Fe Alloy Thin Films J.M. Bai, P.A. Montano, and M. Tomkiewicz	121
Glancing Angle X-ray Fluorescence Study of Ni/W Multilayers J.M. Bai, K. Huang, P.A. Montano, M. Grimsditch, E. Fullerton, D. Kelly, and I.K. Schuller	122
<i>In Situ</i> X-ray Diffraction of the Early Stages of the Crystallization of Fe ₈₀ B ₂₀ C. Brizard, B. Rodricks, E. Alp, and R. Macchiarie	122
X-ray Reflectivity Studies of Diblock Copolymer Films Z. Cai, G.P. Felcher, K. Huang, P.A. Montano, J.M. Bai, T.P. Russell, and G.W. Zajac	123
<i>In-Situ</i> X-ray Reflectivity Study of Sputter Deposition R.P. Chiarello, H.K. Kim, and H. You	123
Interface Structure of Epitaxial Si/CoSi ₂ /Si(001): an X-ray Reflectivity and Fluorescence Study K.G. Huang, J.M. Bai, Q.F. Xiao, and L.J. Schowalter	124

Beamline X7A

Structural Investigations of Some Synthetic Zeolite Phases K.J. Andires, J.V. Smith, and S. Han	124
Lattice Instabilities and the Effect of Copper-Oxygen Sheet Distortions on Superconductivity in La ₂ CuO ₄ -Superconductors: La _{1.60-x} Nd _{0.40} Sr _x CuO ₄ M. Crawford, R. Harlow, E. McCarron, and W. Farneth	112
<i>In Situ</i> Powder Diffraction Studies of the Hydriding Behavior of Nanocrystalline Pd J.A. Eastman, M.R. Fitzsimmons, and L.J. Thompson	125
Lattice Instabilities in Nd-Doped La ₂ CuO ₄ : Presence of a PCCN Phase? R. Harlow, M. Crawford, E. McCarron, and W. Farneth	125
Lattice Instabilities and the Effect of Copper-Oxygen Sheet Distortions on Superconductivity in La ₂ CuO ₄ -Superconductors: La _{1.88-x} Nd _x Sr _{0.12} CuO ₄ R. Harlow, M. Crawford, E. McCarron, and W. Farneth	114
Orientalional Ordering Phase Transition in C ₆₀ P.A. Heiney, J.E. Fischer, A.R. McGhie, W.J. Romanow A.N. Denenstein, J.P. McCauley, Jr., A.B. Smith, and D.E. Cox	126
The Crystal Structure Determination of BiMo ₂ O ₇ (OH) · 2H ₂ O from Powder Diffraction Data J.A. Hriljac and C.C. Torardi	126
The Crystal Structure Determination of CuPt ₃ O ₆ from Powder Diffraction Data J.A. Hriljac, J.B. Parise, G.H. Kwei, and K.B. Schwartz	127
The Effects of High-Pressure Annealing on the Phase Transitions in La _{1.88} Ba _{0.12} CuO ₄ J.A. Hriljac, D.H. Chen, D.E. Cox, and A.R. Moodenbaugh	127
Atomic Structure of Al ₆₅ Ru ₁₅ Cu ₂₀ R. Hu, T. Egami, A.P. Tsai, A. Inoue, and T. Masumoto	128
Compression of Solid Krypton to 130 GPa A.P. Jephcoat, L.W. Finger, and D.E. Cox	128
High-Resolution, High-Pressure Powder Diffraction A.P. Jephcoat, L.W. Finger, and D.E. Cox	129
Microstructural Aspects of the LTO-LTT Transition in La _{1.875} Ba _{0.125} CuO ₄ M. Marezio, H. Takagi, B. Batlogg, P. Bordet, D.E. Cox, and Uchida	129
Preliminary Structural Investigations of the (sic) Some Univalent Cation Exchanged Forms of the RHO Framework J.B. Parise, D.R. Corbin, L. Abrams, T.M. Nenoff, and G.D. Stucky	130
Anomalous Diffraction Study of Pb(Mg _{1/3} , Nb _{2/3})O ₃ H.D. Rosenfeld and T. Egami	130
The Local Atomic Structure of Pb(Mg _{1/3} , Nb _{2/3})O ₃ H.D. Rosenfeld and T. Egami	131

High Resolution Synchrotron X-ray Diffraction Study of Phase Separations in $\text{Pr}_2\text{NiO}_{4+\delta}$ J.D. Sullivan, D.J. Buttrey, D.E. Cox, and J. Hriljak	131
Re-Examination of the Phase Diagram in Heavily-Doped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ H. Takagi, M. Marezio, R.J. Cava, B. Batlogg, P. Bordet, J. Hodeau, and D.E. Cox	132
Orientalional Disorder in Solvent-Free Solid C_{70} G.B. Vaughan, P.A. Heiney, J.E. Fischer, D.E. Luzzi, D.A. Ricketts-Foot, A.R. McGhie, Y.W. Hui, A.L. Smith, W.J. Romanow, B.H. Allen, N. Coustel, J.P. McCauley Jr., A.B. Smith, and D.E. Cox	132
Isothermal Compression of Fe_2SiO_4 (SPINEL) at Pressures Up to 24 GPa at Room Temperature L.-J. Wang, J.-A. Xu, L.C. Ming, J. Balogh, and H. Manghanani	133
High Temperature Behavior of $(\text{Mg,Fe})\text{SiO}_3$ Perovskite at 1 Bar Y. Wang, Y. Zhao, D.J. Weidner, R.C. Liebermann, J.B. Parise, and D.E. Cox	133
Comparative Study of Elemental Contrast by Neutron and Resonant X-ray Powder Diffraction J.K. Warner, A.P. Wilkinson, A.K. Cheetham, and D.E. Cox	134
$\text{Fe}^{\text{II}}/\text{Fe}^{\text{III}}$ Valence Contrast by Resonant X-ray Powder Diffraction J.K. Warner, A.K. Cheetham, and D.E. Cox	134
Reproducibility of Site Occupancies Obtained from Resonant X-ray Powder Diffraction J.K. Warner, A.P. Wilkinson, A.K. Cheetham, and D.E. Cox	135
Atomic In-Plane Disorder in Co/Pt Superlattices X. Yan, T. Egami, E.E. Marinero, R.F.C Farrow, and C.H. Lee	135
Atomic Interdiffusion in Co/Pt Superlattices X. Yan, T. Egami, E.E. Marinero, R.F.C. Farrow, and C.H. Lee	136
Perovskite Structure as Function of Temperature & Pressure: The Results from NaMgF_3 Perovskite Y. Zhao, D.J. Weidner, J.B. Parise, and D.E. Cox	136

Beamline X7B

High Pressure Single Crystal X-ray Diffraction Investigations of CaF_2 : Data Collection at 6.7 GPa and at Normal Pressure S. Åsbrink, J.C. Hanson, M. Malinowski, and A. Waskowska	137
Single Crystals with Intercalated Proteins from Biomineralized Materials A. Berman, L. Addadi, L. Leiserowiz, S. Weiner, J. Hanson, and T. Koetzle	137
Study of the Low Temperature Structure of the 2212 Tl-Ba-Ca-Cu-O Superconductor M. Blomberg, I. Mutikainen, and J. Hanson	138
Diffraction Linewidth of Charge-Density Wave Satellite Spots S.M. DeLand, G. Mozurkewich, and L.D. Chapman	138
Single Crystal Study of Hydrated Aluminophosphate at 20 K T. Ericksson, Å. Kvik, Y. Long, and J.C. Hanson	139
EXAFS Investigation of Amorphous Platinum Uridine Clusters in H_2O -Solution V. Eteläniemi, R. Serimaa, T. Laitalainen, T. Paakkari, and I. Mutikainen	139

Beamline X9A

EXAFS Studies of Copper-Containing Monooxygenases N.J. Blackburn and R.S. Strange	140
X-ray Diffraction Studies of Tethered Protein Monolayers and Ultrathin Lipid Multilayer Films on Ge/Si Multilayer Substrates J.K. Blasie, J. Chupa, S. Xu, M. Murphy and R. Fischetti	140
Time-Resolved, Resonance X-ray Diffraction Studies of Sarcoplasmic Reticulum Membrane Multilayers J.K. Blaise, F. Asturias, L. DeLong and R. Fischette	140
XAFS Studies of the Iron Containing Subunit of Ribonucleotide Reductase G. Bunker, K. Zhang, B. Chance, M. Sahlin, and B.-M. Sjöberg	141
The Structure of the Transient State Intermediate of the H^+ -ATPASE from Chloroplasts C. Carmeli, O. Tadmor, R. Ophir, and S. Carmeli	141

<i>In-Situ</i> XANES Study of the Chemistry of Passive Oxides on AlV and AlMn Thin Films Under Electrochemical Control	
A.J. Davenport, H.S. Isaacs, G.S. Frankel, A.G. Schrott, C.V. Jahnes, and M.A. Russak	142
Purification, EXAFS, and Edge Data of Vitamin-B12 Binding Proteins with Cyano-and Aquo-Cobalamin	
S.M. Frisbie and M.R. Chance	142
EXAFS Studies of Ni Tetraazamacrocycles and Their Adducts as Models of F430	
L.R. Furenlid, M.W. Renner, and E. Fujita	143
<i>In-Situ</i> Characterization of a Copper Monolayer Electrochemically Adsorbed on a Platinum Electrode	
T.E. Furtak and T.M. Hayes	143
The Structure of the Zinc Sites of <i>E. coli</i> DNA-Dependent RNA Polymerase	
W.-J. Huang, F.Y.-H. Wu, R.B. Sinclair, and L. Powers	144
Drug Structures In Biological Membranes	
L. G. Herbetete and L. A. Nakhimovsky	144
Discotic Liquid Crystal Strands	
S.H.J. Idziak, N.C. Maliszewskyj, G.B.M. Vaughan, P.A. Heiney, J.P. McCauley, Jr. A.B. Smith, III, C. Meresdorf, and H. Ringsdorf	144
X-Ray Absorption Spectroscopy of Manganese in Chloroplasts	
M.P. Klein, V.K. Yachandra, V.J. DeRose, M.J. Latimer, I. Mukerji, and K. Sauer	145
EXAFS Experiments on Iron, Copper, and Zinc Proteins	
R. Korszun	145
Protein Damage and Possible Protection: EXAFS Studies of Metallo Enzymes	
C. Kumar, K.S. Reddy, C. Lee, and B. Chance	146
X-ray Induced Damage to Proteins During EXAFS Studies	
C. Kumar, B. Chance, and K.S. Reddy	146
The Vertical X-ray Beam Stability at Beam Lines X-9A & B at NSLS	
H.-C. Lee, S. Khalid, M.-Z. Zhang, and B. Chance	147
Glancing-Angle XAFS Studies of Premelting at Cu/Al ₂ O ₃ Interfacial Regions	
R.A. Mayanovic, B.A. Bunker, and N. Jisrawi	147
Polarized X-ray Absorption Spectroscopy	
J.E. Penner-Hahn, G.S. Waldo, K. Clark, and T. Stemmler	148
Hemeprotein Catalysis: Structure ↔ Function Relationship	
L. Powers, R.B. Sinclair, C.-S. Chang, S. Hallam, and I. Yamazaki	148
EXAFS Studies of [(HB(3,5- ¹ Pr ₂ pz) ₃)Fe ^{II} (OBz)(MeCN)] (1)	
L. Que, Jr., C.R. Randall, A.E. True, R.C. Holz, and T.E. Elgren	148
EXAFS Investigations of E7 Mutant Myoglobins	
K.S. Reddy, L. Powers, H. Tang, and B. Chance	149
Photomultiplier/Phosphor X-ray Detector for EXAFS Measurements	
K.S. Reddy, S. Khalid, C. Kumar, and B. Chance	149
Prototype Stopped Flow Apparatus for EXAFS Study of Enzyme Intermediates	
K.S. Reddy, C. Kumar, and B. Chance	150
Improved Sagittal Focus of Scanning Monochromator	
G. Rosenbaum, M. Sullivan, and L. Rock	150
Small Angle Diffraction Pattern from Smooth Muscle	
G. Rosenbaum and A. Somlyo	151
EXAFS Structure of Retroviral Gag Zinc Finger Protein-RNA Complex in Intact Retroviral Particles	
I. Sagi, L.O. Arthur, L.E. Henderson, M.F. Summers, J.W. Bess, and M.R. Chance	151
EXAFS Studies of Zinc Finger Binding Proteins	
I. Sagi, M.F. Summers, M. Wirt, E. Chen, and M.R. Chance	152
SIT Detector Characterization	
V. Skita, R. Kelly	152
Ligninase Active Site Structure	
R. Sinclair, I. Yamazaki, J. Bumpus, B. Brock, C.-S. Chang, A. Albo, and L. Powers	152

EXAFS of Fe^{3+} in Active Soybean Lipoxygenase R.C. Scarrow, G. Grove, M.G. Trimitsis, M. Nelson, and R. Cowling	153
Solution Scattering Studies of the Polyphenolic Protein Isolated from the Blue Mussel <i>Mytilus Edulis</i> M.W. Trumbore, V. Skita, and R. Kelley	153
XAS Investigations Into the Structural Conformations of the Redox States of Hydrogenase from <i>Thiocapsa Roseopersicina</i> J.P. Whitehead, Cs. Bagyinka, G.J. Colpas, and M.J. Maroney	154
Ge Detector Dead Time Loss and Its Effect on XAFS K. Zhang, G. Rosenbaum, and G. Bunker	154
XAFS Studies of Carboxypeptidase and its Intermediate State K. Zhang, D.S. Auld, K.S. Larson, and B. Chance	155
An X-ray Absorption Study on Pro-carboxypeptidase K. Zhang, D.S. Auld, and K.S. Larson	155
An XAFS Study on HPV-16 E7 Protein K. Zhang and D. Patrick	156

Beamline X10A,B

Transverse Structure Function of the Liquid-Vapor Interface of Ga A. Acero, Z.-H. Cai, E. Flom, B.-H. Lin, N. Maskil and S.A. Rice	156
Direct Phase Determination of Large Macromolecule Crystals Using Three-Beam X-Ray Interference S.-L. Chang, H. E. King, Jr., M.-T. Huang, and Y. Gao	157
High Resolution Small Angle X-ray Scattering Study of Styrene-Butadiene Diblock Copolymer S.-L. Chang, K.S. Liang, Z. Fu, R.J. Rose, and K.J. Joen	157
Phase Determination for Macromolecular Crystals Using X-ray Multiple Diffraction S.-L. Chang, H. King, M.T. Huang, and Y. Gao	158
GIXD Study of Self-Assembled Monolayers P. Fenter, J. Li, P. Eisenberger, T.A. Ramanarayanan, and K.S. Liang	158
Misfit Dislocations in the Interface of Epitaxial Nb Films on Sapphire C.H. Lee, and K.S. Liang	159
The Structure of Dilute Lamellar L_α Phases of Surfactant Membranes N. Lei and C.R. Safinya	159
Surface-Induced Order on a $\text{Cu}_3\text{Au}(001)$ Surface K.S. Liang, H.H. Hung, Z. Fu, and S.-L. Chang	160
High Resolution Powder Diffraction Studies of Disorder in Lithium Nickel Oxides I.J. Pickering, J.T. Lewandowski, and A.J. Jacobson	160
Location of Iron in Framework-Substituted Zeolite L Using High Resolution Powder Diffraction I.J. Pickering, D.E.W. Vaughan, and K.G. Strohmaier	161
Effect of Excess Gold on the Surface Ordering Kinetics of the $\text{Cu}_3\text{Au}(001)$ Surface S.B. Rivers, W.N. Unertl, H.H. Hung, K.L. D'Amico, and K.S. Liang	161
Melting and Reassembling Transition of the 2-D Protein Lattice of Bacteriorhodopsin Y. Shen, C.R. Safinya, K.J. Rothschild, and B. Thomas	162
X-ray Diffraction Studies of Phospholipid-Based-Tubules B. Thomas, C.R. Safinya, B. Ratna, R. Shashidhar, and N.A. Clark	162
Specular and Off-Specular X-ray Reflectivity from Sputtered Fe/Cr Multilayers M.F. Toney and S.S.P. Parkin	163
Observation of Ordered CoPt_3 in Single Crystal Magnetic Multilayers M.F. Toney, R. Farrow, and R. Marks	163
X-ray Reflectivity Study of Oxidation Induced Roughening of the $\text{Pt}(111)$ Single Crystal Surface Under In-Situ Electrochemical Potential Control H. You, D.J. Zurawski, Z. Nagy, R.P. Chiarello, H.K. Kim, and R.M. Yonco	164
Pressure-Induced Staging Transition in TiS_2 Intercalation Compounds O. Zhou, J.E. Fischer, and K.S. Liang	164

Structure of M_6C_{60} ($M = K, Rb, \text{ and } Cs$) O. Zhou, J.E. Fischer, N. Coustel, and Q. Zhu	165
X-Ray Diffraction Study of Rb_xC_{60} Phase Diagram Q. Zhu, O. Zhou, N. Coustel, G. Vauhan, and J.E. Fischer	165

Beamline X10C

XANES of Molybdenum Compounds and Catalysts at the Molybdenum $L_{2,3}$ -Edges S.R. Bare, G.E. Mitchell, and J.L. Gland	166
XAFS of Two Novel Tungsten-Containing Enzymes from the Hyperthermophilic Bacteria <i>Pyrococcus Furiosus</i> and <i>Thermococcus Litoralis</i> Y. Gea, G.N. George, R.C. Prince, S. Mukund, and M.W.W. Adams	166
Polarized Single Crystal Sulfur K-Edge X-ray Absorption Spectroscopy of Substituted Thiophenes G.N. George, M. Sansone, and M.J. Gorbaty	167
X-ray Absorption Spectroscopy of Zinc Dialkyl Dithiophosphates G.N. George, M. Sansone, and D. Martella	167
X-ray Fluorescence XAFS Microscopy at $50\mu m$ Resolution G.N. George and M. Sansone	168
Polarized EXAFS Studies of Nitrogenase Crystals S.L. George, J. Chen, J. Christiansen, S.P. Cramer, N. Campobasso, J. Bolin, M. Sansone, and G.N. George	168
XAS Study of Metal Partitioning at Water/Oxide Interfaces K.F. Hayes, L.E. Katz, and J.E. Penner-Hahn	169
<i>In-Situ</i> XANES of an Iron Porphyrin Irreversibly Adsorbed on an Electrode Surface S. Kim, I.T. Bae, M. Sandifer, P.N. Ross, R. Carr, J. Woicik, M.R. Antonio, and D. Scherson	169
Investigation of the Chemistry of Chromium in Anti-Corrosion Finishes on Aluminum F.W. Lytle and R.B. Gregor	170
Near-Edge vs EXAFS Studies of Oxides: Understanding Without the Tears G. Meitzner and E. Hayden	170
DIFFRAXAFS: A Novel Technique for Probing Local Atomic Environment I.J. Pickering, M. Sansone, J. Marsch, and G.N. George	171
X-ray Absorption Spectroscopy of Transition Metal Substituted Zeolites I.J. Pickering, G.N. George, G.H. Via, D. Sondericker, K.G. Strohmaier, and D.E. Vaughan	171
Sulfur K-Edge X-ray Absorption Studies of Photographic Materials T. Smith, J. DeWitt, K. Hodgson, and B. Hedman	172

Beamline X11A

Titanium Germanide Formation D.B. Aldrich, C.L. Jahncke, R.J. Nemanich, and D.E. Sayers	172
Annular Detector for Obtaining Accurate X-ray Absorption Spectra in Fluorescence from Thick Concentrated Samples D.L. Brews, D.M. Pease, P. Chartier, J.I. Budnick, C.E. Bouldin, and Z. Tan	173
Morphology of Supported Platinum Catalysts Prepared from Platinum Carbonyls on Magnesium Oxide J.-R. Chang and B.C. Gates	173
Genesis of Pt-Re Bimetallic Interactions in Supported Pt-Re Catalysts J.-R. Chang, S.K. Purnell, and B.C. Gates	174
Electron-Yield and Fluorescence XAFS Study of E-Beam Deposited Co/Pd Multilayers M. Choi, J.H. Joo, S.K. Kim, J.S. Kang, Y.-P. Lee, S.-C. Shin, and S.M. Heald	174
Electron-Yield EXAFS Study of Multilayers for Magneto-Optical Device Materials M. Choi, J.H. Joo, J.G. Lee, C.S. Yang, K. Kim, D. Whang, and S.M. Heald	175
XAFS of Strained Discontinuous Au Films G.A. DeRose and R.W. Hoffman	175

XAFS of Chemically Synthesized II-VI Nanoclusters R.H. Felton, A.J. Bard, and C. Hoener	176
Glancing Angle X-ray Study of Crystallization of Amorphous Ge at the Ge-Al Interface S.M. Heald and J.K.D. Jayanetti	176
Operation of a Dynamically-Bent Sagittally-Focussing Double Crystal Monochromator for XAFS Studies S.M. Heald and G.M. Lambie	177
Interfacial Reactions of Ni-Cr Alloys and Aluminum S.M. Heald, Z. Tan, and J. Jayanetti	177
X-ray Reflectivity Study of Thin Film Density for Al, Cu, Ni, Co, Cr, Mn, and Pd S.M. Heald, Z. Tan, and J.K.D. Jayanetti	178
Ligand Effects in Supported Metal Carbonyls: X-ray Absorption Spectroscopy of Rhenium Subcarbonyls on Magnesium Oxide A. Honji, L.U. Gron, J.-R. Chang, and B.C. Gates	178
EXAFS Studies of the Interfacial Phase of Highly Milled Sn/Ge Composites J.K.D.S. Jayanetti, S.M. Heald, and Z. Tan	179
σ^2 -Derivative Method for Disorder in PbTiO_3 M. Joo, A.M. Edwards, and D.E. Sayers	179
Slope Analysis of Disorder in SrTiO_3 M. Joo, A.M. Edwards, and D.E. Sayers	180
Temperature-Dependent EXAFS Study of $\text{Sr}_{0.8}\text{Pb}_{0.2}\text{TiO}_3$ M. Joo, A. Edwards, and D. Sayers	180
Measurement of Internal Strains Using Embedded Micromarkers E.H. Jordan, S.C.U. Ochi, D.M. Pease, and J.I. Budnick	181
XANES of Conversion Coatings on Aluminum Alloys M. Kendig, A. Davenport, and H. Isaacs	181
A Time-Resolved Diffraction Study of the Ta-C Combustion System E.M. Larson, J.B. Holt, J. Wong, P.A. Waide, B. Rupp, and L. Terminello	182
Comparison Between Thorium and Cerium Doping and Study of the Role of Reduction for η -Type Superconductor Related Systems G. Liang, Y. Guo, D. Badresingh, M. Croft, J. Chen, J. Peng, R.F. Jardim, E.A. Early, M.B. Maple, O. Beom-hoan, J.T. Markert, M. Nagoshi, and T. Suzuki	182
Electronic Structure Study of $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$ by Polarization X-ray Absorption Near-Edge Structure G. Liang, W. Xu, M. Croft, J. Chen, J. Peng, O. Beom-hoan, and J.T. Markert	183
X-ray Absorption Study of Electron-Doped High- T_c Superconductor-Related Compounds $\text{R}_{2-x}\text{Ce}_x\text{CuO}_{4-\delta}$ G. Liang, Y. Guo, D. Badresingh, W. Xu, M. Croft, J. Chen, J. Peng, O. Beom-hoan, and J.T. Markert	183
Thermal Vibrations and Melting from a Local Perspective P. Livins, E.A. Stern, and Z. Zhang	184
XANES and EXAFS Study of Mn Doped CaF_2 Dosimetric Phosphor A.N. Mansour and J.B. Barkyoumb	184
Characterization of a $\text{Cu}/\text{Al}_2\text{O}_3$ Buried Interface Using Glancing-Angle X-ray Reflectivity and XAFS Studies R.A. Mayanovic, B.A. Bunker, C. Zhong, and A. Anagnostopoulos	185
Glancing-Angle XAFS Studies of Premelting at $\text{Cu}/\text{Al}_2\text{O}_3$ Interfacial Regions R.A. Mayanovic, B.A. Bunker, and N. Jisrawi	185
EXAFS Investigations of $\text{Pt-Au}/\gamma\text{-Al}_2\text{O}_3$ Catalysts K.I. Pandya, S.M. Heald, J. Fraissard, and L. Petrakis	186
$L_{3,2}$ Core Hole Lifetime Widths of 3d Transition Metals J.M. Pease	186
The Structural Characterization of Metal Oxide Clusters in Zeolite-Y and Zeolite-L Hosts, Produced by the Photo-oxidation of Group(VI) Metal Hexacarbonyls, Followed by Various Degrees of Subsequent Reduction I.A. Prokopowicz, A. Malek, M. Steele, A. Stein, and G.A. Ozin	187
Spontaneous Ordering EXAFS Study of $\text{Ga}_{0.5}\text{In}_{0.5}\text{P}$ Thin Films Deposited on GaAs A. Tabor-Morris, K. Kemner, R. Mayanovic, and B. Bunker	187

Rare Earth Valence and Doping in T', T'- and T*-Phase R_2CuO_4 (R=Rare Earths) Z. Tan, S.M. Heald, S.-W. Cheong, A.S. Cooper, and J.I. Budnick	188
X-ray-Absorption Studies of Local Structure of $LaR_2SrCu_2O_{6+y}$ (R=La, Pr, Nd, Sm and Gd) Z. Tan, S.M. Heald, S.-W. Cheong, H. Hwang, A.S. Cooper, and L.W. Rupp, Jr.	188
EXAFS Study of Fe Core Structure in Reconstituted Ferritins Q. Wang, A. Edwards, D. Sayers, and E. Theil	189
EXAFS Study of Local Structure of Ferroelectric $Zn_xCd_{1-x}Te$ Alloys Z. Wang, R. Mayanovic, B. Bunker, U. Debska, and J. Furdyna	189
EXAFS and X-ray Edge Studies of Cobalt Substituted Bacteriorhodopsin: Irreversible Light Induced Cation Uptake M.D. Wirt, M.R. Chance, and M. Engelhard	190
X-ray Absorption Near-Edge Study of Several Bi- and Tl-Based High- T_c Superconductor Systems W. Xu, G. Liang, M. Croft, J. Chen, J. Peng, R.S. Liu, B. Jayaram, and J.M. Tarascon	190
Trends in Protein Structure Changes Around the Transition Metal Site in Concanavalin A When the Crystallized Protein is Dissolved Y. Zhang, E.A. Stern, S.L. Lin, and A.J. Kalb	191
Structure and Photo-Induced Structural Changes in Non-Stoichiometric $a-As_xS_{1-x}$ W. Zhou, M.A. Paesler, and D.E. Sayers	191

Beamline X12B

X-ray Diffraction Studies of Liquid Crystalline Polymers P. Cebe, Y. Aihara, P. Huo, C. You, H. Gibson, A. Bluhm, and W. Yeomans	192
SAXS Study of the Folding of a Globular Protein in the Presence and Absence of Chaperonins J. Flanagan, M. Kataoka, M. Capel, and D. Engelman	192
The Role of Histone H1 in Chromatin Higher-Order Structure V. Graziano, M. Chapel, and V. Ramakrishnan	193
X-ray Diffraction Studies of Blends of Poly(Butylene Terephthalate) with Polyarylate P. Huo, C. You, Y. Aihara, and P. Cebe	193
X-ray Diffraction from Mixtures of Collagen Types I and III V. James, J. McConnell, and M. Capel	194
Phase Study of AOT/Water/Decane Dense Microemulsion E.Y. Sheu, M.M. De Tar, and M. Capel	194
Properties of Vacuum Residue Fractions in Organic Solvents E.Y. Sheu, M.M. De Tar, and D.A. Storm	195
SAXS Studies of Porous Carbons J. Wegrzyn, A.N. Golland, M. Capel, and J. Schwartz	195

Beamline X12C

Structure Determination and Refinement of Bovine Lens Leucine Aminopeptidase and Its Complex with Bestatin S.K. Burley, P.R. David, R.M. Sweet, A. Taylor, and W.N. Lipscomb	196
Three-Dimensional Structure of T7 Lysozyme X. Cheng, J.W. Pflugrath, X. Zhang, and F.W. Studier	196
Structural Studies of Tryptophanyl-tRNA Synthetase Using Multiwavelength Anomalous Dispersion from Selenomethionine-Substituted Protein Crystals S. Doublié, R.M. Sweet, and C.W. Carter, Jr.	197
Structural Studies of the Malic Enzyme from the Parasitic Nematode <i>ASCARIS SUUM</i> and the Interleukin-1 Receptor Antagonist Protein H.M. Einspahr and B.C. Finzel	197
Probable Amylase/Protease Inhibitor-B from Rice Seeds K.Y. Hwang, K.K. Kim, and S.W. Suh	198
Thermostable α -Amylase from <i>B. Licheniformis</i> K.Y. Hwang, K.K. Kim, and S.W. Suh	198

X-ray Crystallography of Arylesterase K.K. Kim, K.Y. Hwang, and S.W. Suh	199
X-ray Crystallography of Lipase K.K. Kim, K.Y. Hwang, and S.W. Suh	199
Structural Studies on Flavo-Protein Reductables J. Kuriyan, T.S.R. Krishna, B. Guenther, X.P. Kong, and R.M. Sweet	200
The Structures of DNA and GM-CSF J. LaLonde, K. Swaminathan, R.M. Sweet, and D. Voet	200
Structural Analysis of the <i>E. Coli</i> TRP Repressor L-Tryptophan Binding Pocket: Why 5-Methyltryptophan is a Better Corepressor C.L. Lawson and R.M. Sweet	201
Structural Basis of Latency in Plasminogen Activator Inhibitor-1 J. Mottenen, A. Strand, J. Symersky, R.M. Sweet, D.E. Danley, K.F. Geoghegan, D. Gerard, and E.J. Goldsmith,	201
Three-Dimensional Structure of cAMP-Dependent Protein Kinase J.W. Pflugrath, J. Kuret, X. Cheng, and R.M. Sweet	202
Crystal Structure Determination of Ribosomal Protein S5 V. Ramakrishnan, R.M. Sweet, and S. White	202
Structure of the Globular Domain of Histone H5 V. Ramakrishnan, V. Graziano, and R.M. Sweet	203
Crystallographic Studies of Complexes of Bovine Somatotropin with a Soluble Fragment of its Receptor, Renin and EPSP Synthase H-S. Shieh, R.A. Stegeman, A.M. Stevens, W.C. Stallings, and R.M. Sweet	203
A Comparison of High-Resolution Trypsin Crystal Structures Obtained by White-Beam Laue and Monochromatic X-ray Diffraction Using Synchrotron Radiation P.T. Singer, R.P. Carty, L. Berman, and R.M. Sweet	204
Multiwavelength Data Collection on BamHI T. Strzelecka, L. Dorner, I. Schildkraut, and A. Aggarwal	204
Structure Determination of <i>Staphylococcal Enterotoxin B</i> S. Swaminathan, W. Furey, J. Pletcher, and M. Sax	205

Beamline X14

Diffuse X-ray Scattering from the Alloy Fe-25% Ni-1%C B.D. Butler, P.R. Zschack, H. Yalcinkaya, and J.B. Cohen	205
Studies of the Structure of Amorphous Melanins J. Cheng, M. Eisner, and S.C. Moss	206
Isothermal Thickening and Thinning Processes in Low MW PEO Fractions Observed From Synchrotron SAXS S.Z.D. Cheng, A.-Q. Zhang, J.S. Barley, J.-H. Chen, A. Hubenschuss, and P.R. Zschack	206
Powder Diffraction Study of Magnetoelastic Strain Broadening in Nanocrystalline Dysprosium M.R. Fitzsimmons, J.A. Eastman, and L. Thompson	207
X-ray Scattering Study of Ag/Si(111) Buried Interface Structures H. Hong, R.D. Arburano, D.-S. Lin, H. Chen, T.-C. Chiang, P. Zschack, and E.D. Specht	207
Inelastic X-ray Scattering Spectroscopy in Lithium Hydride E.D. Isaacs, G.E. Ice, and P. Zschack	208
Crystallographic Study of a Single Quasicrystal of Al-Cu-Fe X B. Kan, S.C. Moss, J.L. Robertson, T. Ishimasa, M. Mori, D. Gratias, V. Elser, and P. Zschack	208
X-ray Determination of Site Occupation Parameters in Ordered Ternaries Cu(Au _x M _{1-x})M=Ni,Pd: Anomalous Scattering and Rietveld Refinement R. Kumar, C.J Sparks, E.D. Specht, P. Zschack, G.E. Ice, T. Shiraishi, and K. Hisatsune	209
Direct Determination of Precipitate Strain and Sizes Using High-Order X-ray Diffuse Scattering B.C. Larson, M.D. Galloway, G.E. Ice, and P. Zschack	209
Magnetic X-ray Scattering in Dy/Lu Superlattices M.B. Salamon, R.S. Beach, B.A. Everitt, K. Pettit, A. Matheny, and C.P. Flynn	210

X-ray Studies of Dysprosium/Lutetium Superlattices M.B. Salamon, R.S. Beach, C.P. Flynn, A. Metheny, and J. Borchers	210
Anomalous Behavior of Diffraction Peak Widths in Lipid Monolayers M.C. Shih, T.M. Bohanon, J.M. Mikrut, P. Dutta, and P. Zschack.	211
Effect of pH on a Fatty Acid Monolayer with Calcium Ions in the Subphase M.C. Shih, T.M. Bohanon, J.M. Mikrut, P. Dutta, and P. Zschack.	211
High-Pressure Phases of a Lipid Monolayer and Their Relationship to Isotherm Anomalies M.C. Shih, T.M. Bohanon, J.M. Mikrut, and P. Dutta	212
Time-Domain Spectrum of Resonantly Filtered X-rays from a Nuclear Bragg Monochromator J.C. Tischler, B.C. Larson, and P. Zschack	212
Thermal Effects on Solid Neon C.T. Venkataraman, R.O. Simmons, M.A. Fradkin, and I. Fujita.	213
Study of Alkoxide-Derived Amorphous TiO ₂ Powder Q.J. Wang, S.C. Moss, M. Shalz, A.M. Glaeser, H.W. Zandbergen, and P. Zschack.	213

Beamline X15A

Structural Study of Si-Alkali Metal Interface by XSW S. Lagomarsino, F. Scarinci, P. Castrucci, C. Giannini, J.R. Patel, and E. Fontes	214
X-ray Standing Waves: YBa ₂ Cu ₃ O ₇ on SrTiO ₃ (100) and Si(111) J. Zegenhagen, H.-U. Habermeyer, T. Siegrist, J.R. Patel, and E. Fontes	214

Beamline X16A

<i>In-Situ</i> X-ray Diffraction Study of Silicide Growth P.A. Bennett, B. Devries, P. Eng, and I.K. Robinson	215
Monolayers of Pb on Si(111) K. Evans-Lutterodt, E. Fontes, B.J. Hinch, L.H. Dubois, and J. Patel	215
Adsorption Geometry of Antimony on Si(001) 2 × 1 R. Felici, C. Ottaviani, I.K. Robinson, and P. Eng	216
The Structure of K Monolayers on Cu(100) H.L. Meyerheim, V. Jahns, W. Mortiz, H. Schulz, I.K. Robinson, and P.J. Eng.	216
Oxygen and Sulfur Structures on the Cu(111) Surface I.K. Robinson, R. Feidenhans'l, F. Besenbacher, and F. Jensen	217
X-ray Diffraction Under Grazing Incidence on Si ⁺ Implanted Silicon After Thermal Annealing S. Rugle, H. Metzger, and I. Robinson.	217

Beamline X16B

Investigation of MBE-Grown Iron Silicide Films D.-M. Smilgies, I.K. Robinson, N. Onda, and H. von Känel.	218
Hydrocarbon Films on Silicon I: The Structure of Wetting Films I.M. Tidswell, T.A. Rabedeau, S.D. Kosowsky, and P.S. Pershan.	218
Hydrocarbon Films on Silicon II: The Structure of Chemisorbed C ₁₈ Alkylsiloxane Monolayers I.M. Tidswell, T.A. Rabedeau, S.D. Kosowsky, P.S. Pershan, J.P. Folkers, and G.M. Whitesides	219

Beamline X17B1

DPA Phantom Studies with a Preliminary MECT System F.A. Dilmanian, E. Nachaliel, R.F. Garrett, W.C. Thomlinson, L.D. Chapman, N.F. Gmür, N. Lazarz, H.R. Moulin, T. Oversluizen, H. Raback, M. Rivers, D.N. Slatkin, P. Spanne, P.M. Stefan, A.C. Thompson, N.D. Volkow, and H.D. Zeman	219
Biological Efficacy of the Auger Effect B.H. Laster, W.C. Thomlinson, C. Gordon, N. Gmür, N. Lazarz, V. Benary, L. Warkentien, and R.G. Fairchild	220

In-Situ X-ray Diffraction Study on Modified Spinel (β) and Spinel Phases (γ) of Mg_2SiO_4 at High Pressure and High Temperature	
Y. Meng, D.J. Weidner, G.D. Gwanmesia, R.C. Liebermann, M.T. Vaughan, R.E. Pacalo, A. Yeganeh-Haeri, K. Leinenweber, Y. Wang, and Y. Zhao	220
Performance of the MECT/Materials Science Monochromator in the Laue-Laue Diffraction Mode	
E. Nachaliel, F.A. Dilmanian, R.F. Garrett, T. Oversluizen, L.E. Berman, C. Brite, L.D. Chapman, J.B. Hastings, N. Lazarz, A. Lenhard, H. Rarback, M. Shleifer, D.P. Siddons, P.M. Stefan, and W.C. Thomlinson	221
Preliminary Studies of the Potential for Structure Refinement at High Pressures Using the DIA Apparatus: SiO_2 , Stishovite, at High Pressure	
J.B. Parise, K. Leinenweber, D.J. Weidner, J. Rakovan, M. Vaughan, G. Gwanmesia, and R.C. Liebermann	221
Computed Microtomography of Meteorites using a Linear Array Detector	
S.R. Sutton, P. Spanne, M.L. Rivers, and K.W. Jones	222
More on the High-Pressure, High-Temperature Equation of State of MgSiO_3 Perovskite	
D.J. Weidner, Y. Wang, R.C. Liebermann, M.T. Vaughan, K. Leinenweber, Y. Meng, R.E.G. Pacalo, A. Remsberg, A. Yeganeh-Haeri, and Y. Zhao	222
Perovskite Structure as Function of Temperature & Pressure: - The Results from NaMgF_3 Perovskite	
Y. Zhao, D.J. Weidner, J.B. Parise, and D.E. Cox	136
The Pressure Dependence of Transition Temperature for Structural Phase Transition in NaMgF_3 Perovskite	
Y. Zhao, D.J. Weidner, R.C. Liebermann, X. Liu, J. Ko, Y. Meng, R.E.G. Pacalo, J.B. Parise, M.T. Vaughan, Y. Wang, and A. Yeganeh-Haeri	223

Beamline X17B2

Monochromator Harmonic Content Measurements and Calculations at Energies Above 20 keV	
D. Chapman, H. Moulin, and R. Garrett	223
First Human Transvenous Coronary Angiography Images at the NSLS	
W. Thomlinson, N. Gmür, D. Chapman, N. Lazarz, R. Garrett, J. Morrison, P. Reiser, V. Padmanabhan, L. Ong, S. Green, A. Thompson, H. Zeman, G. Brown, R. Hofstadter, E. Rubenstein, J. Giacomini, and H. Gordon	224
Imaging an Occluded Artery Using Transvenous Coronary Angiography	
W. Thomlinson, N. Gmür, D. Chapman, N. Lazarz, R. Garrett, J. Morrison, P. Reiser, V. Padmanabhan, L. Ong, S. Green, A. Thompson, H. Zeman, G. Brown, R. Hofstadter, E. Rubenstein, J. Giacomini, and H. Gordon	224
A Kinesthetic Charge Detector for Intravenous Coronary Angiography Using Synchrotron Radiation X-rays	
H.D. Zeman, F.A. DiBianca, and W.C. Thomlinson	225

Beamline X17C

Diffraction From Single Crystals of Sub-Attoliter Volume and a New Phase of Bismuth	
E.F. Skelton, L.W. Finger, S.B. Qadri, J.Z. Hu, J.D. Ayers, and H.K. Mao	225
A Method For Probing Variations in δ in $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ Single Crystals Using Energy Dispersive Diffraction	
E.F. Skelton, S.B. Qadri, J.Z. Hu, M.S. Osofsky, J.L. Cohn, and T.A. Vanderah	226

Beamline X18A

Early Stages of Precipitation of δ' in Al - 3.2 wt% Li	
V. Mahadev, K. Mahalingam, and G.L. Liedl	226
A Study of Substrate-Induced Alignment of Smectic LC Phases: Presence of Surface of Substrate Stabilized Phases Near the LC-Air Interface	
L.J. Martinez-Miranda and E. Smela	227
Investigation of the Local Atomic Structure in HgCdTe	
D.Q. Na, J.B. Cohen, and P. Georgopoulos	227
X-ray Characterization of 10 nm Sn Film on Si(111)	
K.F. Peters, J.B. Cohen and P. Georgopoulos	228

X-ray Study of Substrate-Induced Alignment on a Smectic-A Liquid Crystal S.E. Snella and L.J. Martinez-Miranda	228
Multilayer Structure and Growth of Xenon on Ag(111) S.-K. Wang, P. Dai, S.N. Ehrlich, T. Angot, J.Z. Larese, and H. Taub.	229
<i>In-Situ</i> X-ray Diffraction Study of YBCO Thin Films J.Q. Zheng, S. Williams, X.K. Wang, T. Rippert, S. Maglic, D. Segel, R.P.H. Chang, J.B. Ketterson, T. Roberts, Y. Lin, R.T. Kampwirth, and K. Gray	229

Beamline X18B

X-ray Reflectivity Study of Si/C Multilayers J.M. Bai, Y.L. Chan, H. Homma, and P.A. Montano	230
X-ray Reflectivity Study of Ti/C and W/C Multilayers Y.L. Chan, J.M. Bai, P.A. Montano, E. Ziegler, and I.K. Schuller	230
Characterization by EXAFS Spectroscopy of Silver, Sodium Halo-Sodalites and Silver Oxalato-Sodalites R.A. Prokopowicz, A. Stein, and G.A. Ozin.	231
Experimental Characterization of Circular Fresnel Zone Plates W. Yun, J. Chrzas, B. Lai, D. Legnini, J. Viccaro, R. Bionta, and K. Skulina	231

Beamline X19A

<i>In-Situ</i> Investigation of Passive Films on Fe-Cr Alloys J.A. Bardwell, A.J. Davenport, H.S. Isaacs, G.I. Sproule, B. MacDougall and M.J. Graham.	232
Fe K-Edge EXAFS Studies of Nitrogenase J. Chen, S.J. George, J. Christiansen, S.P. Cramer, R. Tittsworth and B. Hales	232
X-ray Absorption Studies of 4d Transition Metal Elements J. Chen, Y. Jeon, F. Lu, and M. Croft.	233
<i>In-Situ</i> XANES Study of Cr Valency Changes in Passive Oxides on AlCr Thin Films Under Electrochemical Control A.J. Davenport, H.S. Isaacs, G.S. Frankel, A.G. Schrott, C.V. Jahnes, and M.A. Russak	233
Microbial Transformations of Uranium: Speciation Studies A.J. Francis, C.J. Dodge, J.B. Gillow, and F. Lu	234
X-ray Absorption Measurements on RT ₂ Si ₂ Compounds Y. Jeon, J. Chen, F. Lu, M. Croft, S.A. Shaheen, and P.H. Ansari	234
Speciation of Uranium Compounds by XANES F. Lu, C.J. Dodge, and A.J. Francis	235
Multi-Shell Fitting of EXAFS Data in Complex Vanadium Compounds F. Lytle, M. Thompson, R. Friedman, and J. Ebner	235
EXAFS of Fe ³⁺ in Nitrile Hydratase M. Nelson, H. Jin, R. Scarrow, G. Grove and C. Buck	236
X-ray Absorption Studies of Organo-Disulfide Redox Polymer Electrodes X.Q. Yang, K.H. Xue, H.S. Lee, J. McBreen, T.A. Skotheim, and F. Lu	236

Beamline X19C

X-ray Topographic Studies of the Phase Transformation Occuring at -143°C in <i>p</i> -Terphenyl Single Crystals R. DiSalvo, M. Dudley, J. Wu, and D. Gordon-Smith	237
Characterization of Chemomechanically Polished CdTe and CdZnTe Single Crystal Wafers by Synchrotron X-ray White Beam Topography T. Fanning, M. Dudley, D. Di Marzio, and L. Casagrande.	237
Modification of Precipitate Strain Fields Accompanying RTP of High Carbon Content Silicon T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith, and D.T. Hodul.	238
Observation of Combined Climb and Glide Induced by RTP of High Carbon Content Silicon T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith, and D.T. Hodul.	238
Observation of RTP Induced Dislocation Nucleation at Precipitates in High Carbon Content Silicon T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith, and D.T. Hodul.	239

Studies of RTP Induced Damage in High Carbon Doped Silicon T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith, and D.T. Hodul	239
Characterization of Defect Structures Close to Grain Boundaries in Ice Crystals F. Liu, I. Baker, G.-D. Yao, and M. Dudley	240
Observation of Faults Produced by Vacancy Condensation in Ice Single Crystals F. Liu, I. Baker, G.-D. Yao, and M. Dudley	240
Synchrotron Topographic Investigation of Damage Distributions Resulting from Electromagnetic Breakdown of Epitaxial Silicon p-n Junctions G. Tolis, M. Dudley, C. Fazi, and D. Gordon-Smith	241
Assessment of Damage Induced by Electromagnetic Breakdown in Silicon Bi-Polar Diodes by Synchrotron White Beam X-ray Topography S. Wang, M. Dudley, D. Gordon-Smith, and C. Fazi	241
Grazing Bragg-Laue Synchrotron X-ray Topographic Imaging of Near-Surface Structural Damage in Silicon Bi-Polar Diodes After Failure with Microwave Pulses S. Wang, M. Dudley, D. Gordon-Smith, and C. Fazi	242
Characterization of Dislocation Structures Around Indents on {111} Indium Antimonide Surfaces J. Wu and M. Dudley	242
Characterization of Gel-Grown Crystals by Synchrotron White Beam Topography J. Wu, S.-P. Wang, and M. Dudley	243
Analysis of Defect Structures in NdGaO ₃ Single Crystals G.-D. Yao, S.-Y. Hou, M. Dudley, and J.M. Phillips	243
Analysis of Twin Structures in LaAlO ₃ Single Crystals G.-D. Yao, S.-Y. Hou, M. Dudley, and J.M. Phillips	244
Comparison Between Observation and Calculation of Surface Relaxation Effects on Dislocation Strain Fields in Thin Films G.-D. Yao, J. Wu, M. Dudley, V. Shastri, and P. Anderson	244
Imaging of Defect Structures in CdZnTe Single Crystals G.-D. Yao, M. Dudley, T. Fanning, and D. Larson	245
Synchrotron Topography Observations of the Second Order Phase Transition in Lanthanum Aluminate Single Crystals G.-D. Yao, S.-Y. Hsu, M. Dudley, and J.M. Phillips	245

Beamline X20A

Observation of Enhanced Ordering in Polymer Films Near Air and Substrate Interfaces Using Grazing Incidence X-ray Diffraction B.J. Factor, T.P. Russell, and M.F. Toney	246
Magnetic X-ray Scattering Study of the 3D Random Field Ising Model J.P. Hill, P. Westbrook, M. Greven, R.J. Birgeneau, and T.R. Thurston	246
X-ray Scattering Studies of Strain at the SiO _x /Si(001) Interface S.D. Kosowsky, T.A. Rabedeau, P.S. Pershan, J. Bevk, and B.S. Freer	247
Nitrogen Induced (2 × 3) Reconstruction of Cu(110) A.R. Sandy, C. You, S.G.J. Mochrie, A. Baddorf, D.M. Zehner, and L.D. Gibbs	247
Electrochemically Adsorbed Bismuth Monolayers on Silver (111): Surface Crystallography of the Uniaxially Commensurate Structure M.F. Toney, J.G. Gordon, M.G. Samant, G.L. Borges, D.G. Wiesler, D. Yee, and L.B. Sorensen	248
Structural and Magnetic X-ray Scattering Measurements of Epitaxial Dy Thin Films M.F. Toney, D.G. Wiesler, B.D. Hermsmeier, and R.F. Farrow	248
Underpotentially Deposited Thallium on Silver (111): Two-Dimensional Compressibility M.F. Toney, J.G. Gordon, M.G. Samant, G.L. Borges, O.R. Melroy, D. Yee, and L.B. Sorensen	249
Structure and Epitaxy of Anodic TiO ₂ /Ti(11 $\bar{2}$ 0) D.G. Wiesler, M.F. Toney, M.G. Samant, O.R. Melroy, W.H. Smyrl, and C.S. McMillan	249

Beamline X20B

<i>In-Situ</i> X-ray Scattering During Electrodeposition Under Controlled Hydrodynamics M.J. Armstrong, G.M. Whitney, and M.F. Toney	250
The Commensurate and Incommensurate Phases of Bilayer Xenon on Single Crystal Graphite W.J. Nuttall, K.P. Fahey, M.J. Young, and R.J. Birgeneau	250

Beamline X20C

Rapid Crystallization of Amorphous Co_2B Studied by Time Resolved X-ray Scattering S. Brauer, H.E. Fischer, J.O. Ström-Olsen, M. Sutton and G.B. Stephenson	251
Polarization Measurements on a 3d Ising System J.P. Hill, P. Westbrook, R.J. Birgeneau, T.R. Thurston, and D. Gibbs	251
Evidence for Two-Step Disordering of the Au(110) 1×2 Reconstruction D.T. Keane, P.A. Bancel, J.L. Jordan-Sweet, G.A. Held, A. Mak, and R.J. Birgeneau	252

Beamline X22B

X-Ray Resonance Magnetic Scattering in $\text{Eu}(\text{As}_{0.02}\text{P}_{0.98})_3$ T. Chattopadhyay, G. Grübel, and L. Rebersky	252
X-ray Resonance Magnetic Scattering in EuTe T. Chattopadhyay and G. Grübel	253
X-ray Reflectivity Study of Interface Growth Dynamics: In-Situ Sputter Deposition of Au on Silicon R.P. Chiarello, H.K. Kim, and H. You	253
Resonant Magnetic Scattering In Europium Compounds G. Grübel, J.D. Axe, L. Rebersky, D. Gibbs and T. Chattopadhyay	254
Potential Induced (1×3) Reconstruction of the Au(110) Surface B. M. Ocko, G. Helgesen, B. Schardt, J. Wang, and A. Hamelin	254
X-ray Crystal Truncation Rod Analysis of Untwinned $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ Single Crystals: Termination Finger Print H. You, U. Welp, G.W. Crabtree, Y. Fang, S.K. Sinha, J.D. Axe, X. Jiang, and S.C. Moss	255
X-ray Diffuse Scattering from Polybromostyrene PBrS/Toluene Solutions W. Zhao, X. Zhao, M. Rafailovich, J. Sokolov, M. Sanyal, and S.K. Sinha	255

Beamline X22C

Capillary Modes on a Close-Packed Metal Surface: Pt(001) D.L. Abernathy, S.G.J. Mochrie, D.M. Zehner, G. Grübel, and D. Gibbs	256
Order, Disorder and Structure of Crystals of $\text{C}_{60}/\text{C}_{70}$ J. Bohr, D. Gibbs, S.K. Sinha, W. Krätschmer, G. Van Tendeloo, E. Larsen, H. Egsgaard, and L.E. Berman	256
Magnetic X-ray Scattering Study of Tb P.M. Gehring, L. Rebersky, D. Gibbs, and G. Shirane	257
Reconstruction of the Pt(111) Surface G. Grübel, K.G. Huang, D. Gibbs, D.M. Zehner, A.R. Sandy, and S.G.J. Mochrie	257
In-Plane Structural Modulation of the CaF_2 Thin Film on Si(111) K.G. Huang, J. Zegenhagen, J.R. Patel and J.M. Phillips	258
Specular and Diffuse Scattering from ErAs Buried Epitaxial Layers P.F. Miceli, M. Sanyal, S.K. Sinha, and C.J. Palmstrøm	258
Au(113) Surface Studies A.R. Sandy, D.L. Abernathy, S.G.J. Mochrie, D.M. Zehner, and L.D. Gibbs	259
Anomalous Reflectivity Measurement of Electron Density Profiles in Thin Films M.K. Sanyal, S.K. Sinha, A. Gibaud, K.G. Huang, M. Rafailovich, J. Sokolov, X. Zhao, W. Zhao, and S.A. Schwarz	259
Determination of Electron Density Profile of a Thin Film Using Anomalous Reflectivity M.K. Sanyal, S.K. Sinha, A. Gibaud, K.G. Huang, B.L. Carvalho, M. Rafailovich, J. Sokolov, X. Zhao, and W. Zhao	260

Resonance X-ray Magnetic Scattering Study of Erbium M.K. Sanyal, D. Gibbs, J. Bohr, and M. Wulff	260
Magnetic Structure in $U_{0.85}Th_{0.15}Sb$ Using Resonant X-ray Scattering C.C. Tang, W. Stirling, G.H. Lander, D. Gibbs, and O. Vogt	261
Magnetic Critical Scattering in Holmium T.R. Thurston, D. Gibbs, J.P. Hill, and G. Shirane	261
X-ray Crystal Truncation Rod Analysis of Untwinned $YBa_2Cu_3O_{7-\delta}$ Single Crystals: Termination Finger Print H. You, U. Welp, G.W. Crabtree, Y. Fang, S.K. Sinha, J.D. Axe, X. Jiang, and S.C. Moss	255
X-ray Diffuse Scattering Study of Liquid Polymer Films W. Zhao, X. Zhao, M. Rafailovich, J. Sokolov, M. Sanyal, and S.K. Sinha	262

Beamline X23A2

Spatially Selective Local Structural Information for InGaAs Multilayers Using Diffraction Anomalous Fine Structure C.E. Bouldin, J. Woicik, H. Stragier, D. Yee, C. Bitz, L.B. Sorensen, and K. Kuhn	262
Local Structural Determinations Using Diffraction Anomalous Fine Structure (DAFS) Measurements J.O. Cross, H. Stragier, L.B. Sorensen, C.E. Bouldin, and J. Woicik	263
Atomic Exchange at Semiconductor Superlattice Interfaces K.M. Kemner, B.A. Bunker, N. Sathya, H. Luo, and J.K. Furdyna	263
Total Electron Yield EXAFS at Low Temperatures K.M. Kemner, Z. Wang, R. Mayanovic, and B.A. Bunker	264
Study of Phase Transitions in Potassium Niobate K.H. Kim, W.T. Elam, and M.I. Bell	264
Characterization of a Cu/Al_2O_3 Buried Interface Using Glancing-Angle X-ray Reflectivity and XAFS Studies R.A. Mayanovic, B.A. Bunker, C. Zhong, and A. Anagnostopoulos	265
Glancing-Angle XAFS Studies of Premelting at Cu/Al_2O_3 Interfacial Regions R.A. Mayanovic, B.A. Bunker, and N. Jisrawi	265
Intermediate-Range-Order Around Copper in Crystalline and Amorphous Silicates Using EXAFS D.A. McKeown	266
EXAFS Study of Structural Changes of the Ni_3Al Ordered Intermetallic by Mechanical Milling T. Nasu, A.M. Edwards, D.E. Sayers, and C.C. Koch	266
Orientation-Dependent XAFS Studies of High- T_c Superconducting Pb-Bi-Sr-Ca-Cu-O System W.F. Pong, P.K. Tseng, J.B. Shi, H.C. Ku, R.A. Mayanovic, and B.A. Bunker	267
X-ray Absorption Investigation of Amorphous Platinum Pyrimidine Derivatives R. Serimaa, V. Eteläniemi, T. Laitalainen, T. Paakkari, and I. Mutikainen	267
Site Selective Local Structure Information for $YBaCuO$ Using Diffraction Anomalous Fine Structure H. Stragier, C. Bitz, D. Yee, L.B. Sorensen, C.E. Bouldin, J. Woicik, J.M. Tranquada, and K. Ludwig	268
EXAFS Study of Spontaneous Ordering in $Ga_{0.5}In_{0.5}P$ Thin Films Deposited on GaAs A. Tabor-Morris, K. Kemner, R. Mayanovic, and B. Bunker	268
EXAFS Study of Gold-Induced Germanium Crystallization Z. Tan, S.M. Heald, C.E. Bouldin, and J.C. Woicik	269
EXAFS Study of Ferroelectric PbS_xTe_{1-x} Alloys Z. Wang and B.A. Bunker	269
EXAFS Study of the Ferroelectric $Zn_xCd_{1-x}Te$ Alloys Z. Wang, R. Mayanovic, B. Bunker, U. Debska, and J. Furdyna	270

Beamline X23A3

In-Situ Observation of Defect Formation in CdTe Crystals at High Temperatures Due to Imposed Mechanical Stresses R. Balasubramanian and W.R. Wilcox	270
--	-----

Diffraction Imaging of a Homoepitaxial Diamond Film D.R. Black and H.E. Burdette	271
Diffraction Imaging of Isotopically Pure Diamond D.R. Black and H.E. Burdette	271
Diffraction Imaging of Isotopically Pure Man-Made Diamond D.R. Black and H.E. Burdette	272
Absolute Calibration of Small-Angle X-ray Scattering R.R. Jemian, G.G. Long, S. Krueger, and D.R. Black	272
Morphology of L-Arginine-Phosphate Crystals for Nonlinear Optical Devices G.G. Long, D.R. Black, and H.E. Burdette	273
X-ray Diffraction Imaging of L-Arginine Phosphate (LAP) G.G. Long, D.R. Black, H.E. Burdette, S.M. Rao, and R.B. Lal	273
Structure of Microporous Foams G.G. Long, S. Krueger, B.J. Olivier, R. Lagasse and D.W. Schaefer	274
High-Resolution SAXS Study of the Structure of Microporous Foams B.J. Olivier, D.W. Schaefer, S. Krueger, and G. Long	274
Diffraction Topography and High Resolution Diffraction Microscopy of Lattice Imperfections, Impurity Inclusions and Fracture in Amethystine α -Quartz H.H. Schloessin, R.D. Spal, and R.A. Secco	275
Phase Contrast Fresnel Diffraction of Hard X-rays by a Thin Film Edge R.D. Spal	275
The Effect of Microgravity on the Mesoscopic Structure of Lead Tin Telluride: High Resolution Monochromatic Diffraction Imaging of Crystals Grown in Microgravity on Space Shuttle STS 61A and on the Ground B. Steiner, R.C. Dobbyn, D. Black, H. Burdette, M. Kuriyama, R. Spal, R. Simchick, and A. Fripp	276
The Mesoscopic Structural Transition from Czochralski to Bridgman Growth of Gallium Arsenide: High Resolution Monochromatic Diffraction Imaging B. Steiner, R.C. Dobbyn, D. Black, H. Burdette, M. Kuriyama, R. Spal, D. Matthiesen, and B. Ditchek	276
The Mesoscopic Structure of Mercuric Iodide Crystals Grown in Spacelab III and on the Ground: High Resolution Monochromatic Diffraction Imaging B. Steiner, R.C. Dobbyn, D. Black, H. Burdette, M. Kuriyama, R. Spal, and L. van den Berg	277
Structural Continuity in Triglycine Sulfate Crystals Grown in Spacelab III: High Resolution Monochromatic Diffraction Imaging B. Steiner, R.C. Dobbyn, D. Black, H. Burdette, M. Kuriyama, R. Spal, R.B. Lal, and A. Batra	277

Beamline X23B

Ag $L_{2,3}$ Edge XANES Study of Charge Redistribution at the Ag Site in Au-Ag Alloys A. Bzowski, Y.M. Yiu, and T.K. Sham	278
Structure in Multilayer Systems C.L. Foiles, J.A. Cowen, D. Koppel, H.-K. Sung, and T.I. Morrison	278
EXAFS Studies of Structural Anisotropy in Fe-Tb Amorphous Films V.G. Harris, K.D. Aylesworth, N.C. Koon, and W.T. Elam	279
EXAFS Studies of Amorphous-To-Crystalline Transition in $Co_{74}Fe_6B_{15}Si_5$ Thin Film Specimens V.G. Harris, S.A. Oliver, P. Dorsey, C. Vittoria, K.H. Kim, and W.T. Elam	279
EXAFS Studies of IBS Fe-Tb Alloy Films V.G. Harris, K.H. Kim, and W.T. Elam	280
Fe Thickness Effect on Structure and Magnetism in Low-Dimensional Fe/Tb Multilayered Structures V.G. Harris, K.H. Kim, N.C. Koon, W.T. Elam, R. Coehoorn, and W. Hoving	280

Beamline X24A

X-24A Experimental Techniques: Flux Measurements from Grating-Multilayer Monochromator Combinations for Low Energy Studies T. Jach, S. Southworth, P.L. Cowan, R.C.C. Perera, and B.A. Karlin	281
---	-----

X-ray Adsorption Isotherms J.L. Jordan-Sweet, B.D. Swanson, and L.B. Sorensen	281
X-ray, Soft X-ray, and VUV Beam Position Monitor B.A. Karlin, P.L. Cowan, and J.C. Woicik	282
X-ray Standing-Wave Study of Sb on InP(110), GaP(110) and GaAs(110) T. Kendelewicz, J.C. Woicik, K.E. Miyano, P.L. Cowan, B.A. Karlin, C.E. Bouldin, P. Pianetta, and W.E. Spicer	282
Double Photoionization of Helium at 2.8 keV J.C. Levin, D.W. Lindle, N. Keller, R.D. Miller, Y. Azuma, N.B. Mansour, H.G. Berry, and I.A. Sellin	283
X-ray Beam Position Monitor Using A Quadrant Pin Diode S.H. Southworth and P.L. Cowan	283
X-ray Standing-Wave Study of Rare Gas Atoms Physisorbed on Single Crystal Graphite B.D. Swanson, C. Bitz, D. Yee, L.B. Sorensen, J. Jordan-Sweet, P. Cowan, and B. Karlin	284
Determination of the Sb/Si(111) Interfacial Structure by Back Reflection X-ray Standing Waves and Surface Extended X-ray Absorption Fine Structure J.C. Woicik, T. Kendelewicz, K.E. Miyano, P.L. Cowan, C.E. Bouldin, B.A. Karlin, P. Pianetta, and W.E. Spicer	284
X-ray Standing Wave Determination of the Clean InP(110) Surface Reconstruction J.C. Woicik, T. Kendelewicz, K.E. Miyano, P.L. Cowan, C.E. Bouldin, B.A. Karlin, P. Pianetta, and W.E. Spicer	285

Beamline X24C

Efficiency of Soft X-ray Detector Phosphors E.L. Benitez, D.E. Husk, and S.E. Schnatterly	285
Photoemission Study of Hydrogen Adsorption on Vanadium Dioxide Near the Semiconductor - Metal Phase Transition V.M. Bermudez, R.T. Williams, J.P. Long, R.K. Reed, and P.H. Klein	286
Combined Pulsed Laser and Synchrotron Radiation Measurements of Photochemical Decomposition of GaAs(110) J.P. Long, S.S. Goldenberg, and M.N. Kabler	286
Photoelectron Spectroscopy of Semiconductors Exhibiting Surface-Potential Inhomogeneities J.P. Long, S.S. Goldenberg, and M.N. Kabler	287
C-1s NEXAFS and EXAFS of Graphite, Natural Diamond, and Synthetic Diamond Thin Films J. Nithianandam and J.C. Rife	287
Process-Dependent Spectral Reflectance Studies of Thermal SiO ₂ on Si H.F. Wei, A.K. Henning, T. Hochwitz, J. Slinkman, J. Rife, and W.R. Hunter	288

Beamline X25

Performance of an Adaptive Jet Cooled Silicon Monochromator in a High Power Density X-ray Beam L.E. Berman, M. Hart, and S. Sharma	288
Order, Disorder and Structure of Crystals of C ₆₀ /C ₇₀ J. Bohr, D. Gibbs, S.K. Sinha, W. Krätschmer, G. Van Tendeloo, E. Larsen, H. Egsgaard, and L.E. Berman	286
Performance of Be Crystals in a High Power Density X-ray Beam A.K. Freund, S. Joksche, K. Kawata, G. Marot, E. Ziegler, L.E. Berman, and J.B. Hastings	289
Performance of Very Thin Si Crystals Under High Heat Load A.K. Freund, G. Marot, K. Kawata, S. Joksche, E. Ziegler, L.E. Berman, and J.B. Hastings	289
Inelastic X-ray Scattering Spectroscopy in Aluminum E.D. Isaacs, H. Williams, P.M. Platzman, P. Zschack, and G.E. Ice	290
High Resolution X-ray Absorption Spectroscopy K. Hämäläinen, D.P. Siddons, J.B. Hastings, L.E. Berman, C.-C. Kao, V. Stojanoff, and S.P. Cramer	290
Performance of a Gallium Cooled 85° Inclined Silicon Monochromator for a High Power Density X-ray Beam W.K. Lee, A.T. Macrander, D.M. Mills, C.S. Rogers, R.K. Smither, and L.E. Berman	291

High Resolution Inelastic X-ray Scattering Study of the Boron K-Absorption Edge S. Manninen and K. Hämäläinen	291
Cryogenic Cooling of Si Monochromators G. Marot, A.K. Freund, M. Rossat, S. Joksche, K. Kawata, E. Ziegler, L.E. Berman, D. Chapman, J.B. Hastings, M. Iarocci, G. van der Laske, and N. Lazarz	292
Kinetics of the Electrochemical Induced Reconstruction of the Au(111) Surface B. Ocko, J. Wang, H. Isaacs, A. Davenport, and L.E. Berman.....	292
Potential Induced (1 × 3) Reconstruction of the Au(110) Surface B.M. Ocko, G. Helgesen, B. Schardt, J. Wang, and A. Hamelin.....	254
A Comparison of High-Resolution Trypsin Crystal Structures Obtained by White-Beam Laue and Monochromatic X-ray Diffraction Using Synchrotron Radiation P.T. Singer, R.P. Carty, L. Berman, and R.M. Sweet.....	204
Intensity Fluctuation Spectroscopy Using Coherent X-rays M. Sutton, E. Dufresne, S.G.J. Mochrie, L.E. Berman, G. Held, and G.B. Stephenson	293
X-ray Search for CDW Satellites in Ba _{1-x} K _x BiO ₃ P. Wochner, Q.J. Wang, S.C. Moss, S.K. Sinha, G. Grübel, H. Chou, L.E. Berman, J.D. Axe, C.-K. Loong, and J.Z. Liu.....	293
High Heat Load Study of Multilayer Monochromators at Room and Low Temperatures E. Ziegler, A.K. Freund, G. Marot, S. Joksche, K. Kawata, M. Rossat, L.E. Berman, J.B. Hastings, M. Iarocci, G. van der Laske, and N. Lazarz	294

Beamline X26A

Lunar Volcanic Glasses: Trace Element Abundances in Individual Spheres J.W. Delano, S.R. Sutton, and J.V. Smith.....	294
Detection and Quantitation by SRXRF of Some Trace Metals in Corn Bran and Corn Bran Passed Through the Pig Gastrointestinal Tract F. Dintiz and K. Jones	295
Cosmic Class Stratospheric Particles: Trace Elements and Zn Depletions G.J. Flynn and S.R. Sutton.....	295
Identification and Quantification of Toxic Trace Elements in Mid-Continental Atmospheric Aerosols R.H. Grant, D.G. Schulze, S.R. Sutton, and M.L. Rivers.....	296
Chromium Oxidation State in Individual Lunar Olivine Grains Using MicroXANES S.R. Sutton, K.W. Jones, B. Gordon, M.L. Rivers, and J.V. Smith	296
Experimental Determination of the X26 Bending Magnet X-ray Source Size S.R. Sutton and M.L. Rivers	297
Analysis of Oceanic Fluid Inclusions by Synchrotron X-ray Fluorescence D.A. Vanko, S.R. Sutton, and M.L. Rivers	297
Analysis of Synthetic Fluid Inclusions in Quartz by SXRF Microbe D.A. Vanko, S.R. Sutton, and M.L. Rivers	298

Beamline X26C

Analysis of Titanium in Boron Nitride Coating on a Beryllium Window N. Gmür and M. Rivers	298
Measurements of Compton-Ejected Electrons for Ne Gas Using Filtered "White Light" of 10-15 keV N.A. Guardala, J.L. Price, M.F. Stumborg, D.G. Simons, G.A. Glass, Jr., B.M. Johnson, and K.W. Jones.....	299
Development of Wire Scanner to Determine Vertical Profiles of Intersecting Photon and Ion Beams B.M. Johnson, D.H. Lee, M. Franklin, and R. Hedeman	299
Synchrotron X-ray Fluorescence Microanalysis of Synthetic Calcite Single Crystals J. Paquette	300

Dynamic Studies of High-Resolution Trypsin Crystal Structures During Catalysis Obtained by White-Beam Laue X-ray Diffraction Using Synchrotron Radiation	
P.T. Singer, R.P. Carty, K. Jones, W.F. Mangel, and R.M. Sweet	300
Laue Diffraction Studies at X26C	
E. Westbrook, M. Westbrook, I. Schlichting, D. Ringe, K.W. Jones, P. Singer, P. Spanne, and R. Sweet	301
Experimental Characterization of Circular Fresnel Zone Plates	
W. Yun, J. Chrzas, B. Lai, D. Legnini, J. Viccaro, R. Bionta, and K. Skulina	231

Improved Images with the Scanning Photoemission Microscope

Harald Ade¹, Cheng-Hao Ko¹, and Erik Anderson²

¹ Department of Physics, SUNY @ Stony Brook, Stony Brook, NY 11794

² Center for X-Ray Optics, Lawrence Berkeley Laboratory, Berkeley, CA 94720

The SPEM is a zone plate based instrument whose imaging characteristics, i.e. its Modulation Transfer Function (MTF), is determined by the size and profile of the zone plate produced microprobe. The sample is then mechanically scanned through this probe. Details regarding the instrumentation have been reported elsewhere [1],[2].

The major improvement in the recent past has been the elimination of astigmatism introduced by the beamline transport optics, with a correspondingly higher spatial resolution. The improvement was achieved through the incorporation of an elliptical zone plate [3], which has different focal lengths in orthogonal directions corresponding to its major and minor radii. In order to assess the performance for both configurations a gold test pattern was imaged in transmission. It consists of 32 radial spokes and constitutes a sample with continuously varying spatial frequencies at all angles. Micrographs of this pattern acquired with the round zone and elliptical plate are shown in Fig. A (image at plane of least confusion) and B (image at focal plane). Artifacts due to contrast reversal are present in image Fig. A, taken with the astigmatic configuration. These artifacts were eliminated in the astigmatism corrected configuration (Fig. B). Furthermore the contrast for all spatial frequencies got markedly improved in Fig. B, and the smallest feature size resolved corresponds to 0.12 μm . However, even at this level of performance, the SPEM is not diffraction limited. The resolution is still dominated by chromatic aberrations and figure errors in the beamline transport optics [3]. The theoretical diffraction limited resolution of the present generation of zone plates used is about 60 nm, and work continues to approach more closely to this limit.

Fig. A

Fig. B

[1] H. Ade, Ph.D. thesis, Stony Brook (unpublished, 1990).

[2] H. Ade, J. Kirz, S. Hulbert, E. Johnson, E. Anderson and D. Kern, "Scanning Photoelectron Microscope with a Zone Plate Generated Microprobe", *Nucl. Instr. Meth. A* **291**, 126-131 (1990).

[3] H. Ade, C.H. Ko, E. Anderson, submitted to *Appl. Phys. Lett.*

This work is supported in part by the NSF under grant # DMR-8815043. The Center for X-Ray Optics is supported by the DOE under contract DE-AC-02-76SF00098.

Photon Beam Mediated Oxidation on Aluminum Surfaces

Harald Ade¹, Cheng-Hao Ko¹, Erik Johnson², Janos Kirz¹, Steve Hulbert²

¹ Department of Physics, SUNY @ Stony Brook, Stony Brook, NY 11794

² National Synchrotron Light Source, Brookhaven National Laboratory, Upton, NY 11973

We have observed enhanced oxidation of oxygen pre-dosed aluminum surfaces in regions exposed to the intense x-ray beam of the Scanning Photoemission Microscope at beamline X1A [1]. The sample was repeatedly Ar-sputtered until photoelectron spectra showed that there were no contaminants left on large areas of the surface. We typically dosed the aluminum surface with 100L to 250L of oxygen, and subsequently exposed the sample to either the focused or defocused photon beam at a total base pressure of 1.5×10^{-9} torr. After several hours of continuous irradiation, the Total Electron Yield and Oxygen KVV Auger images showed a repeatable effect as represented in figure 1. Spectra taken on and off the bright central spot in figure 1 confirmed the significantly increased oxygen concentrations inside the bright spot. The spot size corresponds to the outermost limits of the photon exposed area. However, due to a central stop in the zone plate with about half the zone plate radius, the defocused beam used in figure 1 is ring-shaped with the same proportion as the open to apodized area of the zone plate. For a simple photon induced reaction mechanism, we would therefore not expect to see any oxidation in the center, where no photons are hitting the sample. The defocused beam in this example had an outer and inner radius of about 3.8 μm and 1.6 μm , respectively. With a total photon flux of about 10^9 photons/s the photon flux density was therefore 2.7×10^{17} photons/s/ μm^2 .

Figure 1. TY (right) and O KVV (left) images of a 150L oxygen pre-dosed aluminum sample after the central area was exposed to the defocused photon beam. Data array is 90x90 pixels with a pixel size of 0.35 μm , and a dwell time of 400 msec. Photon energy was 646 eV.

This work is supported in part by the NSF under grant # DMR-8815043

[1] H. Ade et al., to be published *Surf. Interf. Analysis* (April 1992)

Investigation of Calcium Deposits in Human Cartilage using STXM

C.J. Buckley¹, G.F. Foster¹, R.E. Y.S. Ali², C. Scotchford², and J. Kirz³

¹ Department of Physics, Kings College, London WC2R 2LS, UK.

² Institute for Orthopaedics, Brockley Hill, Stanmore, MIDDLEX, UK.

³ Department of Physics, SUNY @ Stony Brook, Stony Brook, NY 11973, USA.

A previously unknown form of mineral deposit has been found near the surface of human femoral head articular cartilage¹. On inspection in the TEM, these deposits (which range in size from 50 to 650nm) appear cubic in shape. It is thought that this form of deposit may be involved in the onset of arthritis. The STXM was used to image the deposits on the micro scale because of its ability to perform high dynamic range elemental imaging in a time scale which is very much shorter than is possible using x-ray microanalysis imaging in the TEM. Also, the STXM does not cause substantial crystal sublimation during imaging which is a problem using TEM.

The Articular cartilage is composed of chondrocyte cells in a collagen matrix. The question arises, are the deposits generated by cells, or are they matrix borne. If they are generated by cells, then it is likely that they would be found in or close to the chondrocytes. Several cells from a number of cartilage sections were imaged at 350.6 eV, and 351.6 eV from which calcium maps were made to investigate the distribution of the deposits. A sample result is shown in fig. 1. The transmission image on the left shows what could be a deposit in side the chondron housing cell, however when the map is taken (right) it is clear that the deposit or artefact is NOT a calcium crystal. This is a typical result from much data, and supports the theory that the cuboid crystals are matrix born.

Fig. 1. STXM generated transmission image at 350.6 eV (left), and calcium map (right) of a 0.1 micron thick fixed, but unstained section of human femoral head cartilage. The scale bar is 1 micron, and the calcium concentration (right) is depicted by a grey scale.

Ref: 1 J.A. Rees et al "Scanning Microscopy and Microanalysis of 'cuboid' crystals in human articular cartilage". Cell mediated calcification and matrix vesicles. ICS 705 1986.

Imaging of Wet, Unfixed and Unstained Muscle at 50nm Resolution

G.F. Foster, C.J. Buckley and R.E. Burge.

Department of Physics, Kings College, London WC2R 2LS.

Images of wet, unfixed and unstained myofibrils have been acquired using 350eV x-rays with the Scanning Transmission X-ray Microscope (STXM) on the X1A beamline of the NSLS. An example image is shown below. The high brightness of the NSLS results in image acquisition times of some minutes with a high signal to noise ratio. The current resolution of this technique is limited by the optics to 50 nm. While this is not yet comparable to the resolution that can be obtained in the electron microscope, the specimen is examined in conditions close to its natural state. The fibres are neither chemically fixed, or stained, allowing transmission images to be obtained - from which the protein distribution along the myofibril can be deduced. We expect this data to be helpful in refining the current model of muscle structure. This data is also beneficial in the calculation of dose imparted to muscle fibres during irradiation¹.

A wet, unfixed, unstained muscle fibre imaged at 350 eV using the scanning transmission x-ray microscope at the X1A beam line. The repeat is 2.5µm.

References

1. G.F. Foster, C.J. Buckley, P.M. Bennett, R.E. Burge "Investigation of radiation damage to biological specimens at 'water window' wavelengths", accepted for publication in Rev. Sci. Instrum. (to be published in 1992)

X1A **QUANTITATIVE MEASUREMENTS OF PROTEIN FLUX ACROSS THE ZYMOGEN GRANULE MEMBRANE USING STXM**

K.K. Goncz (U.C. Berkeley & LBL), Center for X-Ray Optics and S.S. Rothman (U.C. San Francisco)

The zymogen granule is a sub-cellular organelle (~1 µm diameter) found in the acinar cells of the mammalian pancreas. These cells manufacture and secrete, into the small intestine, the enzymes that are used to digest the food we eat. Zymogen granules are storage vesicles for these enzymes inside the cell.

With the scanning transmission X-ray microscope (STXM)¹, we have imaged individual granules, isolated from the pancreas and suspended in 0.3M sucrose, at a resolution of ~50 nm, and calculated the protein content of each one using the numerical absorption data from the pixels in the image. For the population of granules examined thusfar (~100), the protein concentration ranged quite broadly from about 50 mg/ml to 700 mg/ml.

The ability to measure such small amounts of protein (femtograms), has made detecting minute changes in the quantity of protein in individual granules possible and therefore, the flux of protein out of a granule, as induced in various ways, can be quantified. This is exciting since historically^{2,3}, flux measurements could only be performed on populations of such small objects and so the results were the average responses of a population. With STXM we can calculate quantitatively the protein release from a single granule over a period of time and observe, at high resolution, any structural changes.

We have collected protein release curves for different suspending solutions; water, 0.3M sucrose at pH 6-8 and 50 & 250 mg/ml chymotrypsinogen (an endogenous protein) in 0.3M sucrose, over the past three years (Figure). However, it is only in the past year that all of the elements of the experiment have come to the point where we have been able to collect a statistically significant amount of information. This new combination of quantitative and visual data from individual granules releasing their contents will help in understanding how these storage vesicles accumulate and release their proteins in the process of secretion.

A protein release curve for 4 granules from the same sample preparation, suspended in 0.3M sucrose. Granule size does not change significantly over time while protein concentration decreases, on average, about 50%. Pixel size, 32x32 nm; dwell time, 2 msec/pix., λ=34 Å.

¹H. Rarback, et. al., *Nucl. Instr. Meth.* A291, 54-59 (1990).

²C. Liebow & S.S. Rothman, *Nature*, 249, 176-178 (1972).

³S.S. Rothman, J. Underwood, K. McQuaid & J. Grendell, *Biochim. Biophys. Acta* 1074, 85-94 (1991).

This work is supported by the U.S. Department of Energy (O.E.R., O.B.E.S., O.H.E.R.) under Contract No. DE-AC03-76SF00098 and by the U.S. Department of Defense (A.F.O.S.R.) under Contract No. F49620-87-K-0001

X1A

DIFFRACTION-LIMITED IMAGING IN A SCANNING TRANSMISSION X-RAY MICROSCOPE

C. Jacobsen, S. Williams, J. Kirz, X. Zhang (Stony Brook), E. Anderson (LBL), M.I. Browne, C.J. Buckley (King's College, London), D. Kern (IBM), and M. Rivers (Chicago)

The Scanning transmission X-ray microscope (STXM) has been upgraded, using a 45 nm finest linewidth phase zone plate, and a new scanning stage. In the new stage, the motion is generated by piezoelectric pushers with built-in capacitance position sensors providing closed loop feedback arrangement. The electron beam microfabricated zone plate is 90 microns in diameter, and is made of 100 nm thick nickel zones supported on a 100 nm thick silicon nitride membrane. A central stop and order sorting aperture are used to produce a "pure" focussed beam. The thicker central stop, along with a separately mounted order sorting aperture, assures that only the radiation thrown into the first order focus reaches the specimen plane.

To characterize the optical performance of the microscope, a high resolution test pattern was made, utilizing the same techniques as were used to make the zone plate. The test specimen was first imaged in an electron microscope, to determine its "true" form (Fig. "STEM"). The image acquired using the STXM shows the sharp features somewhat blurred, as is expected (Fig. "STXM"). The modulation transfer function (MTF) determined from these images agrees remarkably well with that predicted from the optics of the beamline and the microscope. This agreement provides justification for calculating the point spread function, which can in turn be used to improve the image by deconvolution. Although the result of the deconvolution involves some increase in the noise content of the image, the visibility of the high resolution features is considerably improved (Fig. "Deconv").

Research supported in part by the NSF under grant DIR-9005893

SCANNING LUMINESCENCE X-RAY MICROSCOPY

C. Jacobsen, S. Lindaas, S. Williams, and X. Zhang (Physics Dept., SUNY at Stony Brook)

Scanning luminescence x-ray microscopy is based on the use of the very small focused probe of a scanning x-ray microscope to stimulate visible light emission from dyes usually used in a Fresnel zone plate to produce a focussed x-ray probe, we have obtained 50-75 nm resolution images of P31 phosphor grains, and we have imaged luminescence from polystyrene spheres loaded with 50-100 $\mu\text{mol/g}$ of fluorescent dye. The resolution was limited not by the focussed x-ray probe (we have resolved 36 nm structures in the STXM) but by dark noise and low net efficiency (0.7%) in the luminescence detection system used for this first investigation. This technique should make possible the imaging of dye-tagged sites of biochemical activity at 20-60 nm resolution in wet, unsectioned, and perhaps even initially living cells.

Figure 1: Scanning x-ray microscope images of polystyrene spheres (1.0 μm mean diameter) loaded with 50 $\mu\text{mol/g}$ of a fluorescein-like proprietary dye. The x-ray transmission image is shown at left, and the visible light luminescence image is shown at right.

We are grateful for the crucial contribution of zone plates fabricated by Erik Anderson of the Center for X-ray Optics (LBL) in Dieter Kern's lab at IBM Yorktown Heights. Operation of the X-1A STXM is supported by the NSF under grant DIR-9005893, and OHER, DoE, under contract DE-FG02-89ER60858.

X-ray holography using photoresists

C. Jacobsen and S. Lindaas, Department of Physics, SUNY at Stony Brook, Stony Brook, NY 11794.

M. Howells, Advanced Light Source, Lawrence Berkeley Laboratory, Berkeley, CA 94720

We have recorded Gabor holograms of biological specimens using photoresist detectors and 1.8-3.6 nm x-rays from the X-1A undulator beamline. Figure 1 shows that we are able to image thick biological specimens, and we are presently working to extend our capabilities to deal with wet biological samples. While our presently attained image resolution is 56 nm, there is evidence that the holograms may contain image information at the 16 nm level; we believe that field distortions in the transmission electron microscopes used to enlarge the hologram are imposing the 56 nm limit. As a result, we are building a scanning force microscope with a linear field scanning stage for hologram readout. Finally, Gabor holograms are known to suffer from twin-image noise, and we have developed a technique for reducing twin-image noise. By iteratively transforming between the specimen plane and the hologram plane, we are able to use the constraints of known object boundaries in the specimen plane and measured x-ray intensity in the hologram plane to more fully recover the specimen wavefield.

Fig. 1: With Dr. Stephen Rothman of UCSF, we have studied the structure of rat pancreatic zymogen granules. At left is shown a 1.5 MeV electron micrograph of granules in which enzyme release was stimulated prior to fixation and drying. Similar non-uniformities of granule mass are also shown in the reconstructed hologram at right of an identically prepared sample.

We gratefully acknowledge the support of the NSF under grant DIR-9006893.

DIFFERENTIAL PHASE CONTRAST IMAGING WITH THE STXM

G.R. Morrison and J.R. Palmer (King's College London, UK),
C. Jacobsen and S. Williams (SUNY at Stony Brook, USA)

The usual form of image produced in the scanning transmission x-ray microscope (STXM) is a map of the local x-ray absorption by the specimen, and is determined by the imaginary part of the specimen's complex refractive index. It is, however, possible to obtain information on the real part of the refractive index by sensing changes in the beam direction after it has traversed the specimen, and this can be achieved by replacing the standard single-element transmitted flux detector with a multi-element detector, and then forming appropriate difference signal combinations from the various detector elements. Image signals formed in this way are said to be in differential phase contrast (DPC). The practical implementation of a DPC imaging system on the STXM requires the use of multiple detectors that are both physically very close together, and have a closely matched response to soft x-rays. This is difficult to achieve with gas filled detectors, so the first series of experiments has concentrated on assessing the relative performance of several different avalanche photodiodes (APDs) as alternative photon-counting detectors for use with soft x-rays. The aim is to replace the existing high performance (but single element) gas flow proportional counter (GFPC) installed on the STXM with a four-element APD detector, since modern processing methods mean that it is relatively straightforward to manufacture multi-element devices on a single substrate, and indeed quadrant APD arrays are available commercially for use in visible-light tracking applications.

Single element 'reach-through' APDs from both RCA and Hamamatsu were tested. The usual glass front window was removed and replaced by a 100 nm thick Si_3N_4 membrane coated with ~ 100 nm of aluminium that allowed soft x-rays from beamline X1A (in the range 1.5 to 5.0 nm) to reach the APD, but which largely excluded any visible light. The devices also have an anti-reflection coating on their surface, typically about 110 nm thick as they were designed for use with visible light, but this 'dead' layer does not appear to have an unduly deleterious effect on their use with soft x-rays. To permit photon counting, the APDs were operated in the so-called 'Geiger' mode, so that any photo-induced carriers initiated an avalanche breakdown of the device that was subsequently quenched by a simple passive circuit. Such a mode of operation does not allow any form of energy discrimination to be performed, but with a monochromatic incident beam this is not a requirement at present. Below

All the devices successfully detected soft x-rays; the best were found to be comparable in speed and efficiency to the existing GFPC. All were capable of handling count rates ~ 1 MHz, a limit set by the use of a passive quenching circuit since the devices themselves can be operated at much higher rates. As the APDs were operated at room temperature it was found that the dark count in some cases was quite significantly higher than for the GFPC and increased rapidly as the reverse bias overvoltage was increased. The dark count can be reduced by cooling the APD, but, by careful selection of the device and the operating conditions, this has not so far been necessary.

The first transmission x-ray micrographs using a single-element APD detector have been successfully recorded, and the first tests of a four-channel system to permit DPC imaging have been made. To accommodate the extra signal data produced by the quadrant APD used in this latter case has required some modification to the existing STXM hardware and software, and there remain a few problems to be overcome, principally concerned with the sensitivity of the quadrant APD, which has been poorer than that of an ostensibly similar single-element device. Some crosstalk exists between adjacent detector elements, and this is presently being characterised.

The APD is more compact, robust, and in many ways simpler to operate than a GFPC, and our initial experience of their use suggests that they may become the general-purpose detector of choice in future years, particularly in circumstances where the ability to perform soft x-ray photon counting at high count rates is more important than any form of energy discrimination.

We gratefully acknowledge the support of NATO grant CRG910276 for this work, and the support of the UK Science and Engineering Research Council (Palmer).

Screening Power of Spectator Electrons: Observation of the Electronic Decay of Core, Hole-Rydberg Electron States in N_2

W. Eberhardt, J.-E. Rubensson (KFA Jülich), K. J. Randall, J. Feldhaus, A. L. D. Kilcoyne, A. M. Bradshaw (Fritz-Haber-Institut, Berlin), Z. Xu, P. Johnson (BNL) and Y. Ma (Pacific Northwest Lab.)

A core excited neutral molecule can decay via Auger-type processes where the excited electron either leaves the system (participator decay) or stays as spectator during the process (spectator decay). The former process leads to single hole final states, identical to those reached in direct photoemission, and the latter leads to states with two valence holes and an extra electron in an orbital unoccupied in the ground state. These two-hole-one electron states can be compared to the two-hole final states of the normal Auger process, and information about the screening properties of the extra electron can be extracted.

For intensity reasons, the only core hole excited state in a small molecule, available to decay spectroscopy, was the state corresponding to the intense π^* resonance, where the core electron is promoted to the first empty molecular orbital of the system. The π^* electron strongly interacts with the rest of the electron system, but higher excited states where the excited spectator electron is in an atomic-like Rydberg orbital, the Coulomb interaction is totally dominating and the interpretation of the spectra is simplified, highlighting the screening properties of the spectator electron.

To study the decay spectra of these Rydberg-type states two orders of magnitude higher excitation intensity is needed, compared to the π^* decay spectra. The X1B beamline provides this exciting new possibility, and below we present the first preliminary study of the decay of various excited states of the N_2 molecule.

In Fig. 1 the decay spectra of the two first Rydberg excited states, $1s^{-1}3p$ (c) and the $1s3p$ (b) are compared to the normal Auger spectrum (a) and the decay of the $1s\pi^*$ state (d). All the structures with kinetic energy less than 380 eV are due to spectator decay processes. It is immediately clear that the shape of curves (a) to (c) is very similar. This observation agrees with the prediction that the spectator Rydberg electron couples to the rest of the system, mainly via the Coulomb interaction. This interaction energy determines the shift of the spectral structure. From the shifts one can deduce the screening efficiency of the spectator electron in the initial and the final configurations.

The shape of the $1s\pi^*$ decay spectrum, here recorded with unprecedented statistics, also shows similar structures as curves (a)-(c) so that the screening efficiency can be deduced. However the shape of curve (d) also shows the effects of the spin-orbit coupling to the two hole final state.

Fig. 1 Electronic decay of various core excited states of the N_2 molecule.

Soft X-ray spectroscopy at the X1B undulator

K.J. Randall¹, J. Feldhaus¹, W. Erlebach¹, A.M. Bradshaw¹,
W. Eberhardt², Z. Xu³, Y. Ma³ and P.D. Johnson³

- 1) Fritz-Haber-Institut der Max-Planck-Gesellschaft, 1000 Berlin 33, FRG
- 2) EXXON, USA / Institut für Festkörperforschung, KFA Jülich, FRG
- 3) Brookhaven National Laboratory, Upton, USA

The NSLS X1B undulator is probably the worlds brightest source of soft x-ray photons currently available. On the recently completed spectroscopy beamline X1B (1) we have attempted to use the intrinsically narrow beam size and divergence to maximise the intensity throughput in a spherical grating monochromator similar to 'The Dragon', NSLS U4B (2).

The high power density of the undulator radiation requires the use of water-cooled mirrors, entrance slit and grating. Cooling is necessary such that thermal deformation of figure error is negligible in the SGM design. This has been confirmed by attainment of resolution superior to the Dragon, the SSRL-LBL SGM and the BESSY SX700-II monochromators over the energy range 300-540eV ($C1s^{-1}$, $N1s^{-1}$ and $O1s^{-1}$). As an example we present the oxygen $O 1s^{-1}$ photoabsorption spectrum taken in the gas phase. Figure a) is the overall near edge region taken at the Dragon, the spectrum is dominated by the $1s-\pi^*$ transition at approximately 531eV. The region enclosed by the dashed lines is expanded in figure b). Two overlapping Rydberg series of transitions approaching the spin split ionic levels at 543.1 and 544.2eV, together with vibrational sidebands and broad underlying resonances, result in a complicated series peaks. The same spectrum, taken under similar conditions at X1B is shown in figure c). The improvement in resolution is immediately apparent.

1. K.J. Randall, J. Feldhaus, W. Erlebach, A.M. Bradshaw, W. Eberhardt, Z. Xu, Y. Ma and P.D. Johnson. Rev. Sci. Instr., in press (proc SRI '91 Chester).
2. C.T. Chen Nucl. Instr. Meth **A256** 595 (1987)

CORE LEVEL PHOTOELECTRON SPECTROSCOPY OF GAS PHASE MOLECULES AT VIBRATIONAL RESOLUTION

K.J. Randall¹, J. Feldhaus¹, A.L.D. Kilcoyne¹, A.M. Bradshaw¹, W. Eberhardt²,
J.-E. Rubensson², Y. Ma³, Z. Xu⁴, P.D. Johnson⁴

¹Fritz-Haber-Institut, ²KFA Jülich, ³Pacific Northwest Laboratories, ⁴BNL-Physics

High resolution photoelectron spectroscopy of molecules can provide detailed information on the dynamics of the photoionization process where both the N-electron system and the inter-nuclear geometry respond to the photo-excitation in a strongly correlated way. In particular, the vibrational fine structure depends on the equilibrium geometry of the molecular ion and can thus give qualitative insight into the electronic relaxation. So far photoelectron spectroscopy at vibrational resolution has mainly concentrated on valence electrons. For core electrons with binding energies >200 eV such measurements have only been possible with monochromatized Al K α radiation¹ (1487 eV) and, very recently, with zero electron kinetic energy (ZEKE) photoemission using synchrotron radiation². However, at 1487 eV the accuracy is limited by the source linewidth and Doppler broadening, and at zero kinetic energy by post collision interaction (PCI). The high photon flux at X1B now makes such work possible at photon energies between ~200 eV and ~700 eV.

As a first example we have investigated the carbon $1s$ ionization of methane. In the Figure we show the $C 1s$ photoelectron spectrum taken at a photon energy of ~304 eV. The vibrational peaks are clearly much better resolved than in the corresponding Al K α and ZEKE spectra. The results are consistent with those published in References 1 and 2. Preliminary work on CO indicates changes in the vibrational fine structure in the region of the shape resonance.

1. U. Gelius, S. Svensson, H. Siegbahn, E. Basilier, A. Faxälv and K. Siegbahn, Chem. Phys. Letters **28**, 1 (1974); L. Asplund et al., J. Phys. B **18**, 1569 (1985)
2. P. A. Heimann, L. J. Medhurst, M. R. F. Siggel, D. A. Shirley, C. T. Chen, Y. Ma and F. Sette, Chem. Phys. Lett. **183**, 234 (1991)

Decay of Vibrationally Selected Core Hole States in Small Molecules

X1B

J.-E. Rubensson, W. Eberhardt (KFA Jülich), K. J. Randall, J. Feldhaus, A. L. D. Kilcoyne, A. M. Bradshaw (Fritz-Haber-Institut, Berlin), Z. Xu, P. Johnson (BNL) and Y. Ma (Pacific Northwest Lab.)

The electronic decay of a core excited neutral molecule can leave the system in various single valence hole states, identical to the final states of the direct photoemission process. The photoemission process is a direct transition between the ground state and the single valence hole states. The deexcitation process involves an intermediate core excited state with a different equilibrium geometry, and thus the vibrational fine-structure in the spectra differ. The vibrational structure is of particular interest for the small 2nd-row-element-containing molecules, since here the vibronic level spacing is of the same order of magnitude as the lifetime width of the excited states. The vibrational fine-structure in decay spectra thus contain information about the coupling between the electronic states and the nuclear motion, and, in general, information about the time evolution of the system on a time scale comparable to the core hole "life" time (typically 10^{-14} s).

For a systematic investigation of these effects it is desirable to excite the core hole states with vibrational resolution and also record the decay spectra with comparable resolution. Until recently the intensity and resolving power needed for such studies were not at hand.

The performance of the X1B beamline now opens up this field of research, and here we present the first observation of the effect of vibrational selection on core hole decay spectra. In Fig. 1a we show the decay of the first vibronic level of the $1s\pi^*$ state of gas-phase N_2 .

The sharp peak at 16.7 eV corresponds to the first vibrational level ($v=0$) of the $1\pi_u$ final state, and the weaker progression starting at 15.6 eV corresponds to the first vibrations of the $3\sigma_g$ final state. The spectrum can be compared to the previously recorded spectrum, recorded without vibrational resolution in the excitation (Fig. 1b). The peak at 16.7 eV looks very similar in the two spectra, due to the fact that the $\Delta v=0$ transitions are favored when the initial and final states have the same equilibrium geometry. The peak in Fig. 1b corresponds to several accidentally overlapping transitions.

In the energy region corresponding to the $3\sigma_g$ final state there is a clear difference between the two spectra. The integrated spectrum shows a pronounced shoulder at 15.4 eV, which is missing in the selectively excited spectrum. This shoulder corresponds to a $\Delta v=-1$ transition, i. e. the electronic deexcitation is accompanied by a vibronic deexcitation. This type of transition is obviously blocked when only the $v=0$ state is excited in the intermediate state (Fig. 1a).

For a quantitative analysis, better statistics is needed, and the $v=0$ spectrum should be compared to the decay of the other vibrational levels. Work along these lines is currently under way.

Fig. 1. Participant decay spectra of the $1s\pi^*$ state in N_2 , with (a) and without (b) vibrational resolution in the excitation

Microtomography of calcium deposits in cartilage tissue

X2

K.L. D'Amico, J.H. Dunsmuir, and S.R. Ferguson (Exxon); and C.J. Buckley (King's College, London, UK)

We have used the X2B Microtomography Beamline to investigate the spatial distribution of Ca deposits in cartilage tissue. The sample was mounted in an epoxy substrate and cut to a cube approximately 1.5 mm on an edge, and was imaged at about 15 keV. Despite the low optical depth of the epoxy and tissue material, the relatively large $\mu\text{-contrast}$ required the high energy in order for them to be penetrated with sufficient signal-to-noise to permit reconstructions. The relative size and orientations of the deposits was sought. It was clear from the data that the Ca deposits varied up to sizes in excess of 150 microns. In addition, the roughly spherical particles appeared to be arranged in a well defined pattern, much like beads on a string. Further experiments are planned to investigate the location of deposits relative to physical features such as the neighboring bone tissue. The figure is a volume rendering of the cartilage sample showing the tissue matrix (lighter color) and the Ca deposits, the dark nodules which appear to be strung together.

X2 Determination of the spatial distribution of metal atoms in a supported catalyst system

K.L. D'Amico, J.H. Dunsmuir, S.R. Ferguson, and G.B. Mcvicker (Exxon)

We have used the facility at X2B to map the spatial distribution of a metal atom in a supported metal catalyst. The technique involves taking radiographs of a material at X-ray energies above and below the Pt L-absorption edge. Projection radiographs, maps of the spatial distribution of In (104), are subtracted to yield the metal distribution on the support. The sample consisted of a cylindrical-shaped zeolite extrudate support with a Pt metal loading. The extrudate was cut normal to the cylinder axis to yield a sample about 1 mm thick, and the beam was incident on the material parallel to the cylinder axis. The total loading of the Pt on the extrudate was about 1 wt. %. Typical data acquisition times for a given sample were about 1 hour. The integrated signal over the entire sample can be analyzed quantitatively to yield the absolute Pt concentration for the sample. Alternatively, the relative concentration of the Pt as a function of position in the extrudate can be determined. The figure shows the net Pt distribution in a pellet as a function of position across the material. The plot shows the calculated weight fraction along the line shown.

Platinum Distribution in a Rim-Loaded Catalyst

X2 Observation of the settling of particulates in a heavy oil

J.H. Dunsmuir, K.L. D'Amico, and S.R. Ferguson (Exxon)

We have used radiography to observe the settling of particulates in a sample of a heavy oil. The hydrocarbon was obtained from the bottoms of a Cat Cracker reactor. This material is extremely viscous and typically contains a significant quantity of "fines", which are particles of the Cat Cracker catalyst that remain entrained in the oil after processing. This material must be removed for the hydrocarbon to be used in further processes. Typical removal involves placing the hydrocarbon into a tank and raising the temperature to about 100°C, thus permitting the material to settle out as a result of gravity. Solvents may be added to mediate the settling process. The mechanism of their action is not known, with speculation centering on the possibility that they cause the agglomeration of the particles. Thus we are interested in directly observing the settling process. We have obtained a time-lapse X-ray movie of the settling at room temperature in the absence of a solvent. The data were acquired over the course of about 3 hours, and consisted of a series of radiographs. Particles observed ranged in size from 10-100 microns and settled at the rate of about 100-500 microns per hour. The advantage of observing the particles in this manner is that the entire 1 mm thick sample could be seen by the X-rays, which readily pass through the hydrocarbon. Future experiments will focus on the effects of temperature and solvent additions in influencing the settling process. The figure shows a 3 x 2 mm² field of view of the above sample showing the suspended particles.

PHASE SEPARATION KINETICS OF A SEGMENTED POLYURETHANE

B. Chu, Y. Li, and T. Gao (SUNY)

Segmented polyurethanes (SPUs) usually form microphase separated structures which give rise to unique properties of the materials. Therefore how the microphase separated structures develop is a topic of interest both academically and industrially. The kinetics of phase separation could be studied in a decisive way probably only by using the synchrotron X-ray source. The sample used in this study consists of 4',4'-diphenylmethane diisocyanate (MDI) and 1,4'-butanediol (BD) as the hard segment and polytetramethylene oxide end-capped with polypropylene as the soft segment. The hard segment content is 47%.

Figure 1 (left) shows the small-angle X-ray scattering patterns as a function of elapsed time after the melt was quenched to 140°C. The experimental integrated scattering intensity (Q) normalized by the integrated scattering intensity (Q_0) from ideal phase separated system could be used as an approximate indication of degree of phase separation. The interdomain spacing (d) could be obtained by using the Bragg equation. The interdomain spacing remained unchanged throughout the whole process. The phase separation process could be fitted by an equation of relaxation

$$[(Q/Q_0) - (Q/Q_0)_{\infty}] / [(Q/Q_0)_{\infty} - (Q/Q_0)] = \exp(-t/\tau)$$

where τ is the relaxation time and ∞ , t , and 0 denotes elapsed time at infinity, t and 0 , respectively. The τ value is 3.7×10^3 seconds, which means that the phase separation process is very slow in the SPU system. The good fitting shown in Figure 2 (right) suggests that mobility of the hard segment and viscosity of the system are key factors controlling the structural changes in the SPU system. This is in good agreement with our annealing experiments.

This work is supported by the US Dept of Energy (DEFG0286ER45237A004).

Synchrotron SAXS Study of Mean-Field and Ising-like Critical Behavior of Poly(2-chloro-styrene)/Polystyrene Blends

B. Chu, Q. Ying, K. Linliu, P. Xie, T. Gao, Y. Li (SUNY, Stony Brook), T. Nose, M. Okada (Tokyo Inst. Tech., Japan), A. Darovsky, W. Lehnert (NSLS, BNL)

Critical fluctuations of a polymer blend poly(2-chloro-styrene)/polystyrene (P2CIS/PS) with 22.6% weight fraction dibutyl phthalate (DBP) has been studied by synchrotron SAXS. The experiments have been performed on five samples of P2CIS/PS blends with different weight fractions of PS, listed in Table 1. The molecular weight of PS and P2CIS are 37K and 233K, respectively. The sample holder was set in a thermal block which can be controlled to 0.02°C over a temperature range from 60°C to 215°C. Each run was collected for a 1000 s period.

Based on the Ornstein-Zernike relationship,

$$I(q, T)^{-1} = I(0)^{-1} (1 + q^2 \xi^2)$$

where $I(q, T)$ is the excess scattered intensity at q and T ($q = 4\pi \lambda^{-1} \sin(\theta/2)$, θ is scattering angle, T is temperature) due to critical fluctuations. Both $I(0)$, intensity scattered at zero angle, and ξ , the correlation length, were extracted from SAXS results in a plot of the reciprocal of the excess scattered intensity $I(q, T)^{-1}$ versus q^2 . For the mean-field behavior of sample E, $I(0) \propto \epsilon^{-2}$, $\gamma = 1.00$ (reduced temperature $\epsilon = (T - T_c)/T_c$), as is shown clearly in Figure 1. However, a deviation from the linear behavior was also observed between 131.7°C and the spinodal temperature T_s . In a log-log plot of reciprocal susceptibility versus temperature, a critical exponent γ of 1.23 was observed. Similarly, a critical exponent $\nu = 0.63$ in a log-log plot reciprocal correlation length versus temperature, $\xi \propto \epsilon^{-\nu}$, further strengthens the Ising-like behavior.

In this study, SAXS results clearly reveal a crossover behavior in the static susceptibility $I(0)$ from the mean-field to the Ising-like behavior at 2.8 K below the critical temperature. The critical temperature $T_c = 134.30^\circ\text{C}$ is consistent between SAXS and independent phase separation measurement.

Table 1.

sample	Binodal W _{P2CIS} /W _{PS}	SAXS T _p /°C	T _s /°C	T _c /°C
E	48/52	134.30*	134.30	134.30
A	40/60	136.92	137.20	
B	51/49	134.66	154.60	
C	31/69	157.68	180.50	
D	61/39	162.80	200.00	

samples contain 22.6wt% of DBP

M_w PS : 3.7×10^4

M_w P2CIS : 2.33×10^5

* critical temperature T_c

T_p : binodal temperature

Research supported by the Department of Energy (DEFG0286ER45237A004 and DEFG0589ER75515), the SUNY beamline (DEFG0286ER45321A004) at the NSLS, and the National Science Foundation (INT 8815115).

STRUCTURAL CHANGES UPON ELECTROCHEMICAL INSERTION OF LITHIUM IN THE 2212 BiSrCuO SUPERCONDUCTING PHASE

P. Coppens, P. Lee, Y. Gao
Chemistry Department, SUNY at Buffalo, Buffalo, NY 14214

N. A. Fleischer and J. Manassen
Weizmann Institute of Science, Rehovoth, ISRAEL

The product obtained upon electrochemical insertion of Li into $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8-y}$ pellets was analyzed by powder diffraction methods. Up to a Li content of 2, the reaction proceeds with retention of the host crystal structure and without the appearance of new phases. Bulk superconductivity persists. The c-axis expands by 0.17 Å for $x_{\text{Li}} = 1.85$, while the a- and b-axes undergo a smaller expansion of about 0.02 Å. Structure analysis including the satellite reflections shows the expansion to be due to an increase in interlayer spacing of the Bi-O planes, with a possible small contraction of the remainder of the lattice (see figure). The lithium atoms are assumed to be located in tetrahedral sites between the Bi-O layers. The small expansion of the a- and b-axes is evidence for a decrease of the antibonding hole concentration in the Cu-O plane, due to the addition of electrons. However the decrease is insufficient to account for all the electrons that must be incorporated in the lattice upon Li^+ insertion. This conclusion is supported by the retention of superconductivity which is evidence for incomplete filling of the holes in the Cu-O plane.

Research supported by the Division of Basic Energy Sciences of the Department of Energy (DEFG0291ER45231) and the National Science Foundation (CHE9021069).

LATTICE INSTABILITIES AND THE EFFECT OF COPPER-OXYGEN SHEET DISTORTIONS ON SUPERCONDUCTIVITY IN $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ SUPERCONDUCTORS:

$\text{La}_{1.60-x}\text{Nd}_{0.40}\text{Sr}_x\text{CuO}_4$

M. Crawford, R. Harlow, E. McCarron and W. Farneth (Du Pont)

Measurement of T_c as a function of Sr doping in the $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ series shows an irregularity at $x=0.12$ where the T_c is anomalously low (see figure 1). Furthermore, the ^{160}Nd isotope effect on T_c reaches its maximum at $x=0.12$ even though the maximum T_c occurs at $x=0.15$. These anomalies have suggested that there is something electronically special about $x=0.12$ (one hole per eight Cu atoms). Although the series of Sr doped compounds with $x<0.2$ remains in the LTO, Bmab structural form at all temperatures where these samples superconduct, it is clear from the previous work that a sample doped with 0.4 Nd undergoes two further phase transitions, first to the orthorhombic, Pccn form and finally to a tetragonal, P4₂/mnm form as the temperature of the sample is lowered. We were now curious to see what effect 0.4 Nd doping would have on the superconductivity and the phase transitions in the series $\text{La}_{1.6-x}\text{Nd}_{0.4}\text{Sr}_x\text{CuO}_4$. In particular, would the $\text{Sr}=0.12$ doping remain electronically special even if the tilts of the copper-oxygen octahedra were altered by a phase change from Bmab to P4₂/mnm. The results of the powder diffraction studies and the T_c measurements are given in figure 1. Clearly, the 0.12 Sr concentration remains anomalous, particularly with respect to T_c . However, the diffraction data suggest that the phase transitions are also anomalous at this concentration of Sr leading to the suggestion that the electronic and structural instabilities are correlated to some degree.

La_{1.6-x}Nd_{0.4}Sr_xCuO₄

Figure 1.

Morphology Studies of CHDI Urethane Elastomers by X-ray Diffraction using Synchrotron Radiation.

C. RICHARD DESPER, US Army Materials Technology Laboratory, BENJAMIN CHU and YINGJIE LI, SUNY Stony Brook

Urethane elastomers based on *t*-(1,4)-cyclohexane diisocyanate (CHDI) have reported¹ DSC and TMA softening points well in excess of 200 C. The morphology of a series of CHDI elastomers has been studied by SAXS and WAXS at the National Synchrotron Light Source² at varying temperature and compositions. The WAXS results show a broad amorphous peak with a single weaker and narrower peak superimposed. The narrower peak, which disappears above 130-140 C upon heating and reappears at that temperature upon cooling [see below], is indicative of hard segment ordering falling short of crystallinity. SAXS results indicate lamellar order arising from microphase separation with periodicity in the 16 to 21 nm, depending upon composition. The persistence at elevated temperature of hard segment microstructure, in the absence of crystallinity, is attributed to strong hydrogen bonding interactions associated with the CHDI structure.

1. Janice W. Dieter and Catherine A. Byrne, *Polym. Eng. & Sci.*, **27**, 673-683 (1987).
2. The authors acknowledge the assistance of Jian Liu and Tong Gao, SUNY Stony Brook, in the X-ray experiments.

2.5 CHDI, 1 PTMO 2033, 1.5 BD

TEMPERATURES, LOW PLOT FIRST: 21 61 98 127 159 185 212 177 136 104 70 36
PROCESSED: 28-JAN-91 09:31:40 E:CS000001.00

This research was carried out under the sponsorship of the United States Department of the Army and Department of Energy.

SAXS AND WAXS STUDY OF POLYURETHANES CONTAINING POLYETHYLENE SOFT SEGMENTS

C. R. Desper,* B. Chu,* T. Gao,* Y. Li,* and C. A. Byrne*

* Polymer Research Branch, U. S. Army Materials Technology Laboratory, Watertown, MA 02172.
Chemistry Department, State University of New York at Stony Brook, NY 11794

Small- and Wide-Angle X-ray Scattering have been used to study the morphology of a series of polyurethane elastomers as a function of temperature, the polyurethanes having a common soft segment: "polytail" hydroxy-terminated polyethylene of $M_n = 3336$ (VPO) and functionality 2.6. Each polymer was prepared by condensation with a diisocyanate at a molar ratio 1.1 diisocyanate / soft segment without chain extender. Thermal mechanical analyzer data for each of the elastomers showed two softening points: the first in the 34-48°C range, and the final in the 210-254°C range.

TEMPERATURES, LOW PLOT FIRST: 30 40 50 60 70 80 100 120 150 180

The diisocyanates studied were: toluene diisocyanate (TDI), hexamethylene diisocyanate (HMDI), 4,4'-diphenyl methane diisocyanate (MDI), and 4,4'-dicyclohexyl methane diisocyanate (H12MDI). The above figure shows SAXS patterns from a typical heating experiment using the TDI polymer at the indicated temperatures. The SAXS data show a lamellar structure, evidenced by the intensity maximum, which shifts to smaller angle (higher repeat period) as temperature is raised, finally disappearing in the 60-70°C range. Complementary WAXS studies reveal a highly disordered monoclinic polyethylene structure in this range. For the polymers studied, the lamellar repeat period is 13 nm at 30°C.

Shifting to 14 nm at 40°C and 18 nm at 50°C, independent of the diisocyanate. The lower softening transition at 34-48°C is attributed to the onset of melting in the polyethylene structure, but presumably such structure is easily disrupted by deformation to account for elastomeric properties at these temperatures. With the breakup of lamellar structure above 70°C, elasticity at higher temperatures is attributed to chemical crosslinking arising from the functionality greater than 2 of the soft segment species.

This research was sponsored by U.S. Department of the Army and Department of Energy.
(#DEFG0286ER45237A005).

X3A2

SINGLE CRYSTAL ANOMALOUS SCATTERING STUDIES OF MIXED-VALENCE COMPLEXES

A. Frost-Jensen, M. Pressprich, Y. Gao and P. Coppens
Chemistry Department, SUNY at Buffalo, Buffalo, NY 14214

At specific wavelengths near an atomic absorption edge, atoms of the same element located in a different chemical environment can have different atomic scattering factors. We are investigating a number of *mixed-valence* organometallic complexes with octahedral coordination of the metal atoms (i.e. low anisotropy) to obtain site-specific information on the shift in the f minimum, to be correlated with the electrostatic potential in the region of the resonating-electron orbital. The latter is to be calculated from experimental electron densities on the same compounds.

Data have been collected at seven wavelengths near the K edge on single crystals of $(\text{Mn}(\text{bipy})_2)(\text{BF}_4)_3 \cdot 3\text{H}_2\text{O}$. The wavelengths were chosen from an EXAFS spectrum of the compound around the Mn K edge at points separated by 3–4 eV, selected so as to resolve interesting fine structure. Helium pipes were employed to reduce air absorption. At each wavelength, 25 strong reflections with large positive or negative Mn contributions were collected and corrected for decay, absorption, $L\beta$ effects, and beam fluctuations. The corrected intensities vary significantly and smoothly in the edge region, reflecting the varying f values. Interestingly, the data suggest an approximate 3 electron difference in f values for the two Mn atoms in the edge region, with the more oxidized atom having the more negative values. Further experiments are planned to verify this result.

A trinuclear valence-trapped complex with the oxo-centered Fe_3O core has been synthesized and will be the subject of following studies.

Research supported by the Division of Basic Energy Sciences of the Department of Energy (DEFG0291ER45231) and the National Science Foundation (CHE9021069).

LATTICE INSTABILITIES AND THE EFFECT OF COPPER-OXYGEN SHEET DISTORTIONS ON SUPERCONDUCTIVITY IN $\text{La}_{2-x}\text{CuO}_4$ -SUPERCONDUCTORS: $\text{La}_{1.88-x}\text{Nd}_x\text{Sr}_{0.12}\text{CuO}_4$

X3A7A

R. Harlow, M. Crawford, E. McCarron and W. Farneth (Du Pont)

Compounds of the type $\text{La}_{2-y}\text{Sr}_y\text{CuO}_4$, $0.0 < y < 0.2$, exist only in the high-temperature HTT, 14/mmm, and low-temperature LTO, Bmab, structural forms. Although the highest T_c in this series occurs at $y=0.15$, the $y=0.12$ compound shows the greatest ^{16}O - ^{18}O isotope effect and is presumably the compound in which the T_c would be most sensitive to any type of lattice distortion or transformation. Since it was established that Nd doping could be used to transform La_2CuO_4 from the Bmab to a Pccn structure, we were curious to see what effect Nd doping would have on the structures and T_c 's of $\text{La}_{1.88}\text{Sr}_{0.12}\text{CuO}_4$. Figure 1 plots the diffraction profiles of the (200) and (020) reflections as a function of temperature for the Nd=0.4 sample. As the temperature is lowered, the Bmab structure showed increasing orthorhombicity (as expected) until the temperature reached 70K where the pattern clearly showed the growth of the Pccn phase (two inner peaks). On further cooling, the two peaks unexpectedly coalesced into a single peak of a $P4_2/nm$ phase, previously known to exist only in Ba doped samples. Figure 2 is the complete phase diagram for the $\text{Nd}_x\text{Sr}_{1-x}\text{CuO}_4$ series. T_c values have also been plotted for these samples. Although the lattice instabilities reduce T_c , the samples remain bulk superconductors with sharp Meissner transitions through the Pccn phase and even when the structure is in the tetragonal, $P4_2/nm$, form. Substitution of Nd for La appears to be one way in which these lattice instabilities can be tuned, and this should allow us to study in detail the interplay between these instabilities (which reflect specific tilts of the copper-oxygen octahedra), doping and superconductivity.

Figure 1.

Figure 2.

CRYSTALLIZATION AND MELTING STUDY OF POLY(ARYL ETHER KETONES)

B. Hsiao, K. Gardner, D. Wu (DUPONT), and B. Chu (SUNY)

In recent years, high performance semi-crystalline poly(aryl ether ketones) have generated a lot of interest in the polymer science community for their excellent environmental and mechanical properties. These polymers can be schematically represented as (-P-*ti*-) where P is a 1,4-linked phenyl group and *ti* signifies either an ether or a keto linkage. The addition of keto linkages increases the stiffness of the molecule, as reflected by the increase of melting and glass temperatures. The crystallization and morphology of melting and glass temperatures directly affects the mechanical property. However, the crystallization kinetics of these materials is fast such that traditional characterization techniques are not capable of following the rapid structural change. In this case, the high intensity of a synchrotron is required to study the fast crystallization behavior.

We have constructed a high temperature (to 400C) dual-chamber jump cell device which is suitable for studying the rapid isothermal crystallization in real time using the simultaneous synchrotron SAXS/WAXS techniques. These experiments have been carried out at the SUNY X3A2 Beamline for polyetherylene and its blends. We propose to adapt similar experiments to study the crystallization of poly(aryl ether ketones). Some preliminary measurements have been made to demonstrate the feasibility of this work, which is shown in the following figure.

Synchrotron SAXS run for PEEK isothermal crystallization at 270 degree C

A unique member of the poly(aryl ether ketone) family is poly(aryl ether ketone ketone) or PEEK. PEEK differs from conventional poly(aryl ether ketones) in two aspects: it consists of different meta/para isomer ratios (poly(aryl ether ketones) are all para linked); it shows two stable crystalline polymorphs (poly(aryl ether ketones) have only one). We have prepared PEEK samples having different meta/para isomer ratios. It is also our objective to study the effect of meta isomer on the structure and crystallization behavior of these copolymers using the synchrotron x-ray scattering techniques.

This work is sponsored by DUPONT Fibers and Central Research and Development.

TIME-RESOLVED SYNCHROTRON STUDY OF POLY(ARYL ETHER KETONES) CRYSTALLIZATION AND MELTING

B. Hsiao, K. Gardner, D. Wu (Du Pont), and B. Chu (SUNY)

Time-resolved synchrotron SAXS experiments were carried out at the SUNY X3A2 beamline to study the crystallization and melting behavior of poly(aryl ether ketones). These experiments were accomplished in a high temperature dual-chamber jump cell, capable of reaching 450 degree C.

A typical SAXS measurement of a poly(aryl ether ether ketone) (PEEK) specimen at 240 degree C crystallization temperature is shown in Figure 1. The signal/noise ratio of the data was enhanced by a filtering subroutine. In this run, the minimum time resolution is 3 seconds. The maximum peak position shifts to a higher *q* value initially then remains constant at the later stage, which indicates that the average long period decreases only at the initial stage. In contrast, the invariant (*Q*, area under the curve) increases continuously with time, suggesting that the lamellar thickening persists throughout the entire crystallization process.

The crystallization rate of PEEK is usually fast such that conventional techniques are not capable of following the rapid structural changes. The high intensity of a synchrotron is proven to be suitable for studying the fast crystallization and melting behavior in polymers.

Figure 1. SAXS study of PEEK crystallization at 240 degree C.

This work is sponsored by Du Pont Fibers and Central Research and Development.

X3A2 THERMALLY-INDUCED STRUCTURAL CHANGES IN A SEGMENTED POLYURETHANE

Y. Li, J. Liu, B. Chu (SUNY), A. Darovsky, and W. Lehnert (NSLS)

Segmented polyurethanes (SPUs) are a class of short-block copolymers with unique structures and properties which have drawn a great deal of academic and industrial attentions. However, there have been few studies on the thermal effects of the structure-property relationship in SPUs starting from the melt state. The thermal effects were investigated in the present small-angle X-ray scattering study. The sample consists of 4,4'-diphenylmethane diisocyanate (MDI) and 1,4'-butanediol (BD) as the hard segment and polytetramethylene oxide end-capped with polypropylene as the soft segment. The hard segment content is 47%.

Figure 1 (left) shows the effects of annealing on the structure of the sample after the melt was quenched to (a) 25°C; (b) 80°C; (c) 107°C; (d) 135°C; and (e) 167°C. The experimental integrated scattering intensity (Q) normalized by the integrated scattering intensity (Q_i) from an ideal phase separated system could be used as an approximate indication of degree of phase separation. The interdomain spacing (d) could be obtained by using the Bragg equation. The results are shown in Figure 2 (right). The degree of phase separation increased first with increasing annealing temperature, reached a maximum at $\sim 107^\circ\text{C}$, and then decreased with further increase in the annealing temperature. The interdomain spacing showed a monotonic increase with increasing annealing temperature.

An increase in the mobility of the hard segment and the decrease in the system viscosity resulted in a maximum of degree of phase separation at $\sim 107^\circ\text{C}$. With further increase in the annealing temperature, short hard segments were dissolved in the soft segment matrix (phase mixing). Therefore a decrease in the degree of phase separation was observed. The increase in d with increasing annealing temperature could be explained by extending the Koberstein-Stein coiled/folded hard segment chain model. The hard segment chain extended gradually with increasing annealing temperature.

This work is supported by the US Dept of Energy (DEFG0286ER45237A004).

X-3A2 PROTEIN CRYSTALS FOR PHYSICAL ESTIMATION OF X-RAY TRIPLET PHASES

F. Mo, B.C. Hauback, E. Husstad (U. of Trondheim-NTH, Norway), M. Teeter (Boston College), L.C. Sieker (U. of Washington) and M.S. Lehmann (ILL, France).

Several crystals of each of three small proteins have been examined to explore the possibility for extracting phase information under multiple diffraction conditions. The proteins were crambin in space group $P2_1$, unit-cell volume $V \sim 17,500 \text{ \AA}^3$, rubredoxin (from *Desulfovibrio vulgaris*) in $P2_1$, $V \sim 19,400 \text{ \AA}^3$, and soybean hydrophobic protein in $P2_12_12_1$, $V \sim 65,200 \text{ \AA}^3$. An unfocused beam was used for this work, both with and without an analyzer crystal, and wavelengths $\lambda = 1.75, 1.5$ and $1.0 \text{ \AA}$.

Previous work has indicated that the single most critical parameter in physical phase estimation is the quality of the crystal, *viz.* splitting and broadening of the reflections. This property varied strongly in the specimens studied, from crystals where all reflection profiles as examined by ω scans had FWHM-values $> 0.5^\circ$, to one highly anisotropic case, featuring a wide range of peak characteristics, from sharp single maxima with FWHM $\sim 0.025^\circ$, to a string of split maxima appearing over a scan range of $\sim 0.7^\circ$.

Significant "Urnweg" diffraction was observed in the primary diffracted intensity for a crystal of crambin. Attempts to map out the phase signal failed, however, due to hardware/software problems. FWHM values for the three reflections involved in the "best" triplet were $0.028, 0.041$ and 0.091° , respectively.

Successful phase estimation has been reported recently for single crystals of myoglobin (1) and hemoglobin (2). In both these cases FWHM-values for the intensity maxima were in the range $0.017 - 0.025^\circ$. Our experiments confirm the criticality of crystal mosaicity in this type of experiment, but also indicate that phase information may be extracted if the intensities involved have profiles with FWHM $\lesssim 0.05^\circ$.

1. K. Hümmer, W. Schweigle and E. Weckert. *Acta Cryst. A* **47**, 60-62 (1991).
2. S.L. Chang. Personal communication (1991).

Research supported by the Norwegian Research Council (NAVF), Grants 424.90/064 and 424.91/098.

Structures of Alkali-Doped Fullerenes by Powder Diffraction

P. W. Stephens and L. Mihaly (SUNY Stony Brook), F. Diederich, K. Holzer, S.-M. Huang, R. B. Kaner, R. L. Whetten, and J. B. Wiley (UCLA), and P. L. Lee (SUNY Buffalo)

We have used powder x-ray diffraction and Rietveld analysis to determine the structure of several phases of alkali-doped fullerenes. These materials have attracted particular interest since the recent discovery of superconductivity in K_3C_{60} .

Based on powder diffraction profiles such as that shown below, we found that K_3C_{60} has a face-centered-cubic structure with cubic lattice constant 14.24 Å, nearly equal to undoped C_{60} . [1] The K ions fit into interstitial sites, where they constrain the orientation of the C_{60} into the two possible states having eight hexagonal rings facing {111} directions. The orientation of C_{60} throughout the lattice is random, so that the structure has space group Fm3m. (The group Fm3, for identically oriented C_{60} , is specifically excluded.) We find that this two-fold orientational disorder persists to temperatures as low as 11 K.

We have also studied a series of samples of Rb_3C_{60} . [2] We found three structures: superconducting Rb_3C_{60} is fcc, with the same binary orientational disorder as K_3C_{60} ; bcc Rb_3C_{60} has a structure analogous to K_3C_{60} and Cs_3C_{60} described earlier; [3] and a body-centered tetragonal phase Rb_4C_{60} , which was also described by Fleming et al. [4]

- [1] P. W. Stephens et al., Nature **351**, 632 (1991).
- [2] P. W. Stephens et al., submitted to Phys. Rev. B.
- [3] O. Zhou et al., Nature **351**, 462 (1991).
- [4] R. Fleming et al., Nature **352**, 701 (1991).

Work supported by NSF, D&L Packard Foundation, and DOE grant DEFG0286ER45231

TIME-RESOLVED X-RAY SCATTERING STUDIES OF THE CRYSTALLIZATION OF POLYETHYLENE BLENDS

K. Tashiro, R.S. Stein (U. Mass.), M. M. Saitowski (Procter& Gamble), B. Chu, Y. Li. (SUNY Stony Brook)

Simultaneous wide angle x-ray diffraction (WAXD) and small angle x-ray scattering (SAXS) were used to study the crystallization behavior of blends of linear polyethylene (High Density PE) and short chain branched polyethylene (Linear Low Density PE). While miscible in the melt, differently branched PEs have different crystallization temperatures and rates. Consequently, blends may phase separate to some extent during the crystallization process. WAXD and SAXS results show that LLDPE with 18 short chain branches (SCB) per 1000 backbone carbons blended with HDPE does indeed phase separate when crystallized. However when deuterated linear PE (dHDPE) is used instead of normal hydrogenous HDPE, cocrystallization occurs based on WAXD and independent infrared spectroscopy data. Since the melting point of dHDPE is about 6°C lower than HDPE, this indicates that the degree of supercooling of the two components is the important parameter. The lamellar size of the linear PE is controlled by the degree of supercooling. Shown below are the SAXS patterns for HDPE/LLDPE and dHDPE/LLDPE at 80°C during a slow cooling (1°C/minute) from the melt. SAXS results indicate that the lamellar size of the neat dHDPE is much closer to that of the branched LLDPE than the lamellar size of HDPE homopolymer. Blends of dHDPE and LLDPE have one intermediate lamellar periodicity, while HDPE and LLDPE blends have a SAXS peak and a shoulder indicating two lamellae sizes. The shoulder is associated with phase separated LLDPE domains and the peak associated with primarily HDPE domains. Given the apparent superposition of HDPE and LLDPE scattering, the size scale of the phase separation must be greater than the lamellar size scale. For the dHDPE blends with LLDPE SCB of greater than 20 SCB/1000, cocrystallization is not evident.

Fig. 1. SAXS of HDPE/LLDPE at 80°C

Fig. 2. SAXS of dHDPE/LLDPE at 80°C (Peaks scaled to same height)

DETERMINATION OF THE STRUCTURAL DISTORTIONS CORRESPONDING TO THE q_1 -TYPE AND q_2 -TYPE MODULATIONS IN NIOBIUM TRISELENIDE

Sander van Smalen, Jan L. de Boer and Auke Meetsma
(Department of Inorganic Chemistry, University of Groningen, The Netherlands);
Hans Graafma, Hwo-Shuen Sherr†, Alex Darovskikh and Philip Coppens
(Chemistry Department, SUNY at Buffalo)

A complete structure determination of the modulations corresponding to both the q_1 type and q_2 type Charge Density Waves in NbSe_3 has been completed, based on 972 unique main and satellite reflections, collected at 20K. The structure of NbSe_3 consists of columns of face sharing triangles: prisms of selenium atoms, parallel to the unique monoclinic b-axis, with a niobium atom at the center of each prism. There are three crystallographically independent columns, labelled I, II and III, respectively. The analysis shows that the q_1 -type modulation involves displacements of Nb(III) as well as the selenium atoms on column III and the Se atoms on column II adjacent to Nb(III) . The q_2 type modulation is analogous, but resides on column type I, and also involves the displacements of selenium atoms on column II. The modulation amplitudes are very small: the largest is 0.05 Å and is found on one atom (Nb(III)) only.

The results provide a better understanding of the dimensionality of NbSe_3 . The bond distances imply that the interaction between the modulations on neighboring columns III is very small along c, thus explaining the small correlation length parallel to c^* above T_c . The interaction along a involves a path of only two Nb-Se bonds and a single non-bonded interaction. The latter is between Se atoms which have a relative large modulation amplitude, and thus can be expected to interact relatively strongly. This provides an explanation for the 2-dimensional character of the CDW in terms of elastic interactions between the atoms.

ACKNOWLEDGMENTS

The research of SvS has been made possible by financial support from the Royal Netherlands Academy of Arts and Sciences (KNAW). Partial support of this work by the National Science Foundation (CHE8711736) and support of the SUNY X3 beamline operated by the Division of Basic Energy Sciences of the U.S. Department of Energy (DEFG0286ER45231) are gratefully acknowledged.

† Permanent address: Synchrotron Radiation Research Center, 8th floor, No.6, Roosevelt Road, Sec. 1, Taipei 10757, Taiwan, Republic of China.

J. Wang(SUNY/SB), Y. Li(SUNY/SB), D. G. Peiffer(Exxon), R. D. Lundberg(Exxon) and B. Chu(SUNY/SB)

The ionic peak in small angle X-ray scattering(SAXS) patterns has generally been attributed to the existence of ion-rich microphases. However, the reasons for the absence of an ionic maximum in SAXS patterns with different thermal history and for the irreversible morphology change with temperature are still obscure. In the present study, an attempt was directed toward a better understanding on the microstructure changes with temperature and on the kinetics of phase-diffusion at certain temperatures for a model sulfonated polystyrene(SPS) neutralized with a zinc salt.

Fig.1 shows the temperature-induced SAXS patterns of a 4.5% ZnSPS. The sample film of 1 mm thickness was heated up continuously from room temperature to 300°C and held at each temperature for 15 min before being measured. All the curves were accumulated for 15 min. It is clearly seen that the intensity of the ionic peak associated with the ion-rich microphase increases with increasing temperature from glass transition temperature($\sim 100^\circ\text{C}$) to $\sim 150^\circ\text{C}$, and decreases with increasing temperature from $\sim 150^\circ\text{C}$ to 300°C. A semi-quantitative analysis of these curves was carried out by using the Porod-Debye two-phase model. Both the diffuse-boundary thickness and Porod inhomogeneity length at elevated temperatures suggest that the ion-rich microphase could be disintegrated at higher temperatures. From the scattering curves, we could consider the interphase between the ion-rich microphase and the polymer backbone background becomes sharper from $\sim 100^\circ$ to 150°C and more diffused from $\sim 150^\circ$ to 300°C .

BC acknowledges the U. S. Army Research Office(DAALO391G0040) for the support of this research.

Table I. Data analysis results for 4.5 mol% ZnSPS at varying temperatures

Temp. °C	I^a 10^3	σ nm	E nm	l_p nm
25	3.34	0.060	0.208	3.96
68	3.49	0.062	0.217	4.05
90	3.56	0.063	0.218	4.07
120	3.45	0.057	0.196	4.40
150	3.41	0.056	0.196	4.58
180	3.45	0.055	0.192	4.50
210	3.54	0.087	0.300	4.45
240	3.33	0.092	0.319	4.70
260	2.88	0.093	0.324	4.41
280	2.26	0.180	0.624	4.18

SMALL-ANGLE X-RAY SCATTERING FROM MICROVOIDS IN AMORPHOUS SILICON-BASED SEMICONDUCTORS

D.L. Williamson and Yan Chen (Colorado School of Mines)

Recent small-angle x-ray scattering (SAXS) experiments [1-3] have demonstrated the existence of very small microvoids about 1 nm in diameter in amorphous(a)-Si-based solar-cell materials. The role of these microvoids in the opto-electronic properties of such material is not yet established but the significant internal surface area (about $15 \text{ m}^2/\text{g}$) and number density (about $10^{10}/\text{cm}^3$) associated with such defects are likely to be detrimental. Our research at NSLS is directed toward obtaining more detailed information from current state-of-the-art, device-quality a-Si:H that exhibits relatively weak SAXS due to relatively small volume fractions (about 1%) of the very small microvoids [3].

Figure 1 displays some of our initial results in the form of a Guinier plot ($h^2 = 4\pi/\lambda$, $\lambda = 0.150 \text{ nm}$). The intensity has been normalized to correct for sample thickness, mass absorption, and incident intensity [1] so that a direct comparison of relative SAXS intensity for various samples is presented. The SAXS from a crystalline(c)-Si wafer and the high purity Al foil often used as the substrate for the preparation of suitable SAXS samples is clearly smaller than the two samples of a-Si:H. The latter are device-quality materials prepared by two techniques, glow-discharge (GD) and sputtering (SP). The variation in slope in the plot will be modeled with a distribution of microvoid sizes. Qualitatively, the SP sample has a larger void fraction of slightly larger void sizes than the GD sample.

Figure 2 shows data from the GD sample in two orientations relative to the incident x-ray beam: film surface normal parallel (non-tilted) and at an angle (30° tilt) with respect to the beam. The reduced intensity seen for the tilted orientation (corrected for the increased thickness) is consistent with non-spherical microvoids that can be modeled as ellipsoids with their longer axis preferentially aligned parallel to the film surface normal [2].

1. D.L. Williamson, A.H. Mahan, B.P. Nelson and R.S. Crandall, Appl. Phys. Lett. 55 (1989) 783.
2. A.H. Mahan, D.L. Williamson, B.P. Nelson and R.S. Crandall, Solar Cells, 27 (1989) 465.
3. A.H. Mahan, D.L. Williamson, B.P. Nelson and R.S. Crandall, Phys. Rev. B40 (1989) 12024.

Acknowledgement: This work was supported by the Solar Energy Research Institute (Contract No. XG-1-10063-3) and by the Electric Power Research Institute (Technical Agreement No. RP2824-05).

INVESTIGATION OF MICROSTRUCTURE OF INTERFACES IN $\text{Si}/\text{Ge}_{0.2}\text{Si}_{0.8}$ SUPERLATTICES BY MEANS OF X-RAY REFLECTIVITY AND X-RAY STANDING WAVE

Z. H. Ming, A. Krol, Y.L. Soo, and Y. H. Kao (SUNY/Buffalo), and K. L. Wang (UCLA)

$\text{Si}/\text{Ge}_{0.2}\text{Si}_{0.8}$ superlattice obtained by MBE was investigated by means of x-ray specular reflectivity and x-ray standing waves. Fluorescence yield was monitored at $\text{Ge K}\alpha$ emission line detected by $\text{Si}(\text{Li})$ detector. Fig. 1 shows log of specular reflectivity vs. grazing angle of incidence obtained at $h\nu = 12 \text{ keV}$ for $\text{Si}/\text{Ge}_{0.2}\text{Si}_{0.8}$ superlattice which consists of 40 bilayers with $2 \times 50 \text{ \AA}$ thickness on the Si substrate. The superlattice peaks including the first and third Bragg are clearly observed. Fig. 2 shows the x-ray standing wave pattern at the Bragg peaks obtained for $h\nu = 12 \text{ keV}$. A model assuming uncorrelated interfacial roughness is used to fit the obtained data.

Fig. 1 Log of specular reflectivity

at 12 keV; data: open circles, fit: solid line.

Fig. 2 Fluorescence yield for

incoming photons with 12 keV

The Tensor Force between quarks and the Deformation of Baryons

The LEGS collaboration

G. Blanpied⁴, M. Blecher⁶, A. Caracappa¹, C. Djalali⁴, M.-A. Duval⁴, G. Giordano², S. Hoblit⁵, M. Khandaker⁶, O.C. Kistner¹, G. Matone², L. Miceli¹, W.K. Mize⁴, B.M. Freedom⁴, A.M. Sandorff¹, C. Schaerf³, R.M. Sealock⁵, C.E. Thorn¹, S.T. Thornton⁵, K. Vaziri⁷, C.S. Whisnant^{1,4}, X. Zhao⁶ and M.A. Moinester⁸

¹Physics Department, Brookhaven National Laboratory, Upton N.Y., 11973;

²INFN-Laboratori Nazionali di Frascati, Frascati Italy;

³Universita di Roma and INFN-Sezione di Roma, Rome Italy;

⁴Department of Physics, University of South Carolina, Columbia S.C., 29208;

⁵Department of Physics, University of Virginia, Charlottesville Va., 22903;

⁶Physics Department, Virginia Polytechnic Institute and State University, Blacksburg Va., 24061.

⁷Rensselaer Polytechnic Institute, Troy, N.Y., 12180-3590.

⁸School of Physics and Astronomy, Tel Aviv Univ., 69978 Tel Aviv, Israel

Essentially all constituent quark models invoke a tensor interaction between the quarks in a proton which comes about through one-gluon exchange. This tensor force between quarks mixes in a D state into what would otherwise be a purely S wave proton. The D wave component breaks spherical symmetry, resulting in a non-vanishing $\langle r^2 Y_2 \rangle$ matrix element for the nucleon and a static quadrupole moment and deformation for its first excited state, the delta. The magnitude and sign of this D state component are quite sensitive to how the internal structure of the proton is treated in quark models and have been of great interest in recent years.

The experimental signature of a D wave component would lie in the excitation of the nucleon to the delta resonance, at about 300 MeV. The delta is excited mainly by M1 photons. If there is a D wave component in the delta then this transition can also be excited by an E2 photon. The problem is in understanding the relative magnitude of this E2 admixture compared to the dominant M1 transition. A variety of models predict this mixing to be quite small, anywhere from -0.9 to -4%, so that a high degree of precision is demanded of experiment.

The delta is most readily observed in photo-pion production. During the last 6 or 7 years there has been a flood of attempts to extract this mixing from existing data on pion production with unpolarized photons, and this has led to numbers for the mixing ratio that are anywhere from +4 to -6%. The high level of activity underscores the importance of this question. However, these attempts are hampered by the low degree of sensitivity in reactions with unpolarized beams, and by the presence of large non-resonant backgrounds.

A new set of pion production experiments is underway at LEGS, which have been specifically designed to maximize sensitivity to the E2 amplitude. The observable most

sensitive to an E2 component is the ratio $d\sigma_{\parallel}/d\sigma_{\perp}$ of $p(\gamma,\pi^0)$ cross sections measured using photons with linear polarization parallel and perpendicular to the reaction plane, respectively. Such ratios of cross sections have the advantage of canceling many systematic errors. To test the sensitivity to systematic uncertainties that may survive this ratio, three independent experiments have been conducted at LEGS with different detectors, different methods of defining the gamma ray energy, different polarizations, and using two targets of liquid hydrogen having different cell configurations. The final results from the first two experiments, along with the preliminary results of the third, are in excellent agreement with one another and have been merged into the single data set shown in the figure.

Plotted with the data in the figure are the results of two recent model calculations, both of which are rather sophisticated but very different in approach. The first is the work of Nozawa, Blankleider and Lee [1]. They calculate the contributions of the various diagrams for photo-pion production in a model that is both unitary and gauge invariant. The second model calculation is the work of Davidson, Mukhopadhyay and Wittman [2]. Their approach differs in that they impose Watson's Theorem, in which the photo-pion amplitude is written as a real part multiplied by an exponential containing the pion-nucleon scattering phase shift. The critical difference between

Anomalous X-Ray Reflectivity Measurements on Ni/Fe Alloy Thin Films*

J.M. Bai (Brooklyn College of CUNY), P.A. Montano (Argonne National Laboratory) and M. Tomkowiak (Brooklyn College of CUNY).

We have measured glancing angle x-ray reflectivity from Ni/Fe alloy (0.8 Ni/0.2 Fe) which was sputtered on 5 cm by 5 cm SiO₂ glass. The anomalous reflectivity measurements were carried out by the tuning the x-ray energy below and above the respective K-edges of Fe and Ni. The x-ray energies were chosen to be 7.0 Kev (below the Fe edge), 7.5 Kev (above the Fe edge but below the Ni edge) and 9.0 Kev (above the Ni edge). The phase information obtained from the absorption component of the index of refraction allows a more accurate determination of the electron density profiles of the Ni/Fe alloy.

We use in our analysis of the data the matrix formalism developed by B. Vidal and P. Vincent [1] for the x-ray specular reflectivity. A 3-layer model calculation is fitted to the experimental data in order to determine the parameters. This analysis of the data gives us information on the electron density profile and roughness at the interfaces. The energy-independent parameters (layer thickness and interface roughness) were obtained from the fittings of the three measurements (Fig. 1). The energy dependence of the effective electron density and absorption are in good agreement with the theoretical calculation[2].

REFERENCE

- [1] B. Vidal and P. Vincent, *Applied Optics*, Vol. 23, No. 11, June 1984.
- [2] Satoshi Sasaki, *IEK Report* 90-16, November 1990.

these models is in how they decompose the E2 reaction amplitude into resonant (having to do with the excitation of the delta) and non-resonant parts. This difference can be tested by comparing with measurements both on- (~320 MeV) and off-resonance (~50 MeV lower). These two theories differ significantly in the energy dependence of the parallel/perpendicular cross section ratio. The top curves in the figure are from Nozawa et al. The long-dashed line is their calculation for ~3.1% E2 admixture. The dotted curve is obtained by turning off the resonant part of the Isospin-3/2 E2 amplitude. Near the peak of the delta resonance, about 320 MeV, the long-dashed curve is quite close to the data. However, this calculation clearly has the wrong energy dependence, which implies that the backgrounds have not been treated adequately. The lower curves in the figure are from Davidson et al. The solid line is their result for a mixing ratio of ~1.4% and the dashed curve is obtained by turning off the resonant part of the E2 amplitude. These calculations exhibit the correct energy dependence, but they appear low. The same general trends are repeated at other angles.

If the fact that the Davidson calculations exhibit the correct energy dependence is taken as a confirmation that the backgrounds are properly treated in this model, then a mixing ratio of about 4% would put the full calculation in reasonable agreement with these new polarization data. That would imply a larger deformation for the delta than expected in most models. However, there are outstanding problems. Data at far backward angles (150°), which are determined by only the dominant Magnetic dipole excitation, are not reproduced by either the Nozawa or the Davidson calculations, despite the fact that this multipole is generally regarded as being very well known. The extraction of a small E2 component through its interference with the dominant multipole will be affected by uncertainties in the M1 amplitude, and these are evidently not as small as one would hope. New experiments are underway at LEGS that will provide a significantly improved decomposition of the pion multipoles in the region of the delta resonance, as well as a further reduction in the uncertainties associated with polarization observables such as shown in the above figure.

This work was supported by the US Department of Energy under contract Nos. DE-AC02-76-CH00016 and DE-FG0589-ER4050, and in part by the Istituto Nazionale di Fisica Nucleare (Italy), the US National Science Foundation, and the US-Israel Binational Science Foundation.

[1] S. Nozawa, B. Blankleider, and T.-S. H. Lee, private comm.;

Nucl. Phys. A513, 459 (1990).

[2] R. M. Davidson, N.C. Mukhopadhyay, and R.S. Wittman, private comm.; *Phys. Rev.* D43, 71 (1991).

*This work is supported by the U. S. Department of Energy, BES-Materials Sciences, under contract No. #W-31-109-ENG-38.

Glancing Angle X-Ray Fluorescence Study of Ni/W Multilayers*

J. M. Bai (Brooklyn College of CUNY), K. Huang, P. A. Montano, M. Grimsditch, and E. Fullerton (Argonne National Laboratory), D. Kelly and I.K. Schuller (University of California San Diego).

Ni/W multilayers have been found to exhibit a dramatic softening of the elastic constants as a function of modulation wavelength. Since it has been proposed that this elastic anomaly is due to interfacial effects we have investigated them using surface-sensitive X-ray fluorescence which is a powerful nondestructive technique for surface and interface characterization. In this study we simultaneously measured the angular dependence of the X-ray fluorescence for Ni and W atoms in Ni/W multilayers and the reflectivity at grazing incidence angles. The X-ray energy was set to 11.0 Kev which is high enough to excite both the Ni and W fluorescence. The double-layer-thicknesses of the samples investigated range from 22 to 300 Å, with Ni and W having the same thicknesses within each bi-layer. The figure shows the Ni and W fluorescence together with the reflectivity for one of the samples (with a period of 120 Å). The electric field intensity distribution inside the multilayers depends on the incident angle, the depth-concentration profile can be obtained by analyzing the angular dependence of the fluorescence signals. The electric field intensity inside the multilayer can be calculated according to Fresnel's recursion formula for different incident angles. A full analysis of the data is in progress.

*This work is supported by the U. S. Department of Energy, BES-Materials Sciences, under contract No. #W-31-109-ENG-38.

In Situ X-Ray Diffraction of the Early Stages of the Crystallization of Fe₈₀B₂₀*

C. Brizard, B. Rodriche, E. Alp (Argonne National Laboratory), and R. Macchirri (AT&T Bell Laboratories).

A new technique has been used for the early stages of crystallization of amorphous materials, like metallic alloys. *In situ* x-ray diffractions have been performed during the early stages of crystallization of Fe₈₀B₂₀. The samples are resistively heated to 600° C in a customized vacuum chamber. A programmable Charge Coupled Device (CCD) detector records simultaneously the evolution of the 3 phases: α-Fe, Fe₃B and Fe₂B in the minute scale. This is the first *in situ* x-ray diffraction study of this system in these temperature and time scales. Interesting behaviors have been seen: appearance and disappearance of phases, α-Fe supersaturation solution in boron (B) (found for the first time in this compound), and migration of B out of the α-Fe matrix. The 2-dimensional diffraction pictures show topography irregularities indicating crystallite inhomogeneities.

X-ray diffraction peaks obtained by summing the CCD chip rows. From low to high angle: Fe₃B, α-Fe, and Fe₂B.

*This work is supported by the U. S. Department of Energy, BES-Materials Sciences, under contract No. #W-31-109-ENG-38.

IN SITU X-RAY REFLECTIVITY STUDY OF SPUTTER DEPOSITION

R.P. Chiarello, H.K. Kim, and H. You (ANL)

Sputter deposition is one of the most extensively used techniques of film growth in modern materials science. Despite the extensive usage, understanding the effects of various environmental conditions on the nature of the film growth remains primitive due to limited experimental techniques with which one can directly probe during the deposition. Because of weak scattering and absorption of x-rays through the gaseous environment of sputter deposition, x-ray reflectivity technique is ideal for *in situ* study of sputtered film growth.

The chamber, previously used for Cu-O phase diagram studies [1] and modified for sputtering at cryogenic temperatures, consists of face-magnetron sputter guns [2], a 4 mm wide, 180° open miller window, and sample holder. A low-current power supply was used to control the deposition rate to ~ 1 Å per second.

Reflectivity measurements were made at various thickness of Au film on a polished silicon substrate. The figure shows selected samples of reflectivity scans for 2, 8, 24, 44, 60, and 120 minute deposition. The scans were sequentially offset by a factor 100 for display. It is clear from the scans that there are three stages of growth. At the top, the critical angle still remains that of silicon. The oscillation, however, shows thin layers (~ 50 Å) of Au indicating island formation. The Au critical angle is evident in subsequent scans. From the second the frequency of the oscillation increases indicating the thickening of the layer. At last the oscillation of the intensity is no longer visible because of roughness of the Au surface.

[1] D.J. Miller, et al., submitted to Applied Phys. Lett. (1991).

[2] J.Q. Zheng, et al., J. Vac. Sci. Technol. A9, 128 (1991).

*Supported by DOE under Contract # W-31-109-Eng-38

Symmetric diblock copolymers are macromolecules composed of two unlike polymers A and B of equal molecular weight linked by a chemical bond. The two blocks in a copolymer are incompatible at temperatures below the microphase separation temperature. As a result of specific interactions of A and B with interfaces the thin film of the symmetric diblock copolymers exhibits a multilayered lamellar morphology oriented parallel to the external surface. The diblock copolymer films can be initially prepared in a disordered state with an arbitrary thickness and ordered by annealing. The total film thickness at any point, after ordering, is given by $t(n+1)L$ or $t(n+1/2)L$, where the lamellar period L corresponds to the thickness of the -ABBA- sequence and n is an integer. We wanted to show that, for a model PS/PMMA copolymer system, the annealed film consists of a layer of uniform thickness t_n or t_n+1 , with the unbalance from the full layer appearing in form of islands or holes.

X-ray reflectivity measurements were made on two PSD/PMMA copolymer films on polished silicon substrates. As shown by optical microscopy, one of the specimens had islands on the top of the film while the other was smooth. The figure shows the corresponding reflectivities (the scan for the film with islands is offset by a factor of 100 for the sake of clarity). The high frequency oscillations show that both films have a total thickness of 1482 Å. However, the bottom curve shows also a low-frequency oscillation which corresponds to islands on the surface with a height of 423 Å. This experiment confirms that the layered structure takes place in quantized steps (no tilt). We are now examining the lateral size, correlation and dynamics of the copolymer islands.

* This work is supported by the U.S. Department of Energy, BES-Materials Sciences, under contract No. #W-31-109-ENG-38

Reflectivity

0.1 0.2 0.3 0.4 0.5
 $q [\text{\AA}^{-1}]$

X6B
Interface Structure of Epitaxial Si/CoSi₂/Si(001): an X-ray Reflectivity and Fluorescence Study

K.G. Huang (MSD, ANL*); J.M. Bai (Brooklyn College); Q.F. Xiao (SUNY/Albany); L.J. Schowalter (RPI).

CoSi₂ has a cubic, CaF₂ structure and a lattice mismatch of -1.2% to silicon at room temperature. Due to this small lattice mismatch, CoSi₂ can be grown epitaxially on the low index surfaces of a silicon substrate. Si/CoSi₂/Si(001) system is a promising candidate for applications such as buried epitaxial metal layers and novel devices like the metal-base transistor because of the high efficiency of the electron transmission across the CoSi₂-Si(001) interfaces.

To investigate the structural properties of this system with a buried CoSi₂ layer of various thicknesses, we have carried out an x-ray reflectivity - fluorescence measurement with x-rays of 11 keV, by monitoring the reflectivity and Co K fluorescence simultaneously. Fig.(a) shows the reflectivity of a 400ÅSi/70ÅCoSi₂/Si(001) sample, which gives interference beatings from the two interfaces of the sandwich structure. The Co-fluorescence yield recorded during the reflectivity measurement is shown in Fig.(b). A characteristic feature was observed near the critical angle of the total external reflection, which may yield additional structure information on the density profile of the buried CoSi₂ in the surface normal direction. A detail data analysis is underway.

* Work supported by US DOE, BES-MS, under contract W-31-109-ENG-38.

X7A
STRUCTURAL INVESTIGATION OF SOME SYNTHETIC ZEOLITE PHASES

Andries, K.J., Smith, J.V. and Han S.
 Department of Geophysical Sciences, University of Chicago, Illinois 60637.

Powder synchrotron X-ray data have been collected for some synthetic zeolite phases in order to determine unit cell dimensions and possible space group symmetries which will help to resolve their three-dimensional framework structures. Techniques used for developing theoretical framework structures matching the experimentally derived properties are model building and direct method strategies. Several samples are currently under investigation:

1. A synthetic AlPO₄ framework structure has been indexed on orthorhombic axes with $a = 13.915$ Å, $b = 18.015$ Å, $c = 9.654$ Å. The highest possible space group as deduced from the systematic absences is Pcab. No suitable topological model has been determined so far.
2. An AlPO₄ framework structure, obtained after calculation of the previous one, has been indexed as monoclinic with $a = 13.483$ Å, $b = 5.106$ Å, $c = 10.468$ Å, $\beta = 51.711^\circ$. The systematic absences indicate that Cm, C2 or C2/m or possible space groups. Several low-intensity impurity peaks are present. Purer samples are being prepared. Model building and direct method techniques were unsuccessful so far.
3. The as synthesised (K-) form of a Linde zeolite sample is orthorhombic with $a = 9.555$ Å, $b = 9.938$ Å, $c = 9.455$ Å. The experimentally derived space group is P2₁2₁2₁ with evidence for a pseudo 4₁ or 4₃ axis parallel to b. In the NH₄⁺ form, the structure is tetragonal with $a = 9.632$ Å, $c = 10.019$ Å. The experimental data are consistent with the framework topology of a known synthetic material, but refinement has been unsuccessful so far. We are currently applying direct methods to try and resolve the structural details.
4. The crystal structure of the synthetic zeolite Linde Q⁽¹⁾ (hexagonal, $a = 13.502$ Å, $c = 13.390$ Å) has been refined using the XRS-82 Rietveld refinement program⁽²⁾ to a final R_f = 6.3 %. Results so far indicate that there is no significant deviation from the highest possible topological symmetry (P-62m) and that one K-site is located in the eight-ring between neighbouring hexagonal sheets. Refinement using the GSAS Rietveld refinement program⁽³⁾ is being carried out.

Acknowledgements: We thank D.E. Cox for assistance during data collection.

- (1) Andries, K.J., Bosmans, H.J. & Grobet, P.J. *Zeolites* **11**, 124-131 (1991).
- (2) Baerlocher, Ch. The X-Ray Rietveld System, Version of September 1982, Institut für Kristallographie und Petrographie, E.T.H., Zürich, Switzerland.
- (3) Larson, A.C. & Von Dreele, R.B. General Structure Analysis System, LANSCE, MS-H805, Los Alamos National Laboratory, Los Alamos NM 87545 (1990).

IN SITU POWDER DIFFRACTION STUDIES OF THE HYDRIDING BEHAVIOR OF NANOCRYSTALLINE Pd

J. A. Easman (ANL), M. R. Fitzsimmons (LANL) and L. J. Thompson (ANL)

The small grain sizes (typically 5-50 nm) attainable in metal or ceramic samples formed from gas-condensed powders [1] are responsible for the occurrence of a number of special properties in these "nanocrystalline" materials [2]. One interesting property that has recently been studied is the hydriding behavior of Pd. Müttschle and Kirchheim [3] charged nanocrystalline Pd samples electrolytically with H₂ and observed a significant narrowing of the miscibility gap compared with that typically observed in conventional coarser-grained materials. We have carried out powder diffraction experiments while hydriding nanocrystalline and coarse-grained samples in situ in order to characterize the microstructural changes that occur during hydriding and dehydriding cycles. Diffraction data from nanocrystalline and coarse-grained samples were obtained prior to hydriding at temperatures ranging from ambient to 16K in order to characterize the thermal expansion properties and obtain the temperature dependence of the Debye-Waller parameter. The hydriding transformation was observed at ambient temperature using a position-sensitive detector, which allowed several complete scans to be acquired prior to complete transformation to the PdH phase. The figure below shows the angular region in the vicinity of the (220) reflection at several times during the transformation. While the nanocrystalline sample was observed to transform completely to PdH within several hours, the coarse-grained reference foil did not transform at ambient temperature, indicating a fundamental grain-size dependent difference in properties. Following transformation of the nanocrystalline sample, the temperature experiments were repeated to allow comparison of the properties of nanocrystalline Pd and PdH.

PdH (220) Pd(220)

- [1] K. Kimoto, Y. Kamiya, M. Nonoyama and R. Uyeda, Jpn. J. Appl. Phys. 2, 702 (1963).
- [2] H. Gleiter, in Deformation of Polycrystals: Mechanisms and Microstructures, N. Hansen et al., eds., Risø National Laboratory, Roskilde, (1981).
- [3] T. Müttschle and R. Kirchheim, Scripta Met., 21, 1101 (1987).

Work supported by the U. S. Department of Energy, BES-Materials Sciences, under contract W-31-109-Eng-38.

LATTICE INSTABILITIES IN Nd-DOPED La_2CuO_4 : PRESENCE OF A PCNN PHASE?

R. Harlow, M. Crawford, E. McCarron and W. Farneth (Du Pont)

La_2CuO_4 exists in a high-temperature tetragonal form (HTT, $a/b \neq c$) in which the copper-oxygen sheets are, on average, perfectly planar and in a low-temperature orthorhombic form (LTO, $a \neq b$) in which the copper-oxygen octahedra tilt around one diagonal of the HTT cell by a few degrees. Partial substitution of La with Ba leads to yet another phase transition from the LTO form to a low-temperature tetragonal form (LT_T, $a/b \neq c$) which results from equal rotations about both diagonals. These tilt structures can be described with the aid of figure 1: for HTT, $Q_1=Q_2=0$; for LTO, $Q_1 \neq 0, Q_2=0$; for LT_T, $Q_1=Q_2 \neq 0$. The amount of the tilt in the LTO phase is related to the orthorhombicity of the unit cell which is defined as $[(a-b)/b]$. Figure 2 plots orthorhombicity vs temperature for a series of $\text{La}_{2-x}\text{Nd}_x\text{CuO}_4$ samples with $0.0 < x < 0.4$. Our measurements on undoped La_2CuO_4 have confirmed previous findings that the orthorhombicity increases in a smooth manner as the temperature is decreased. Doping with Nd is seen to have two effects. First, at moderate temperatures, the orthorhombicity (presumably) the tilts, increase significantly with Nd concentration. Second, while the general trend of increasing orthorhombicity with decreasing temperature is observed for moderate temperatures, an abrupt departure from this expected behavior is seen below ca. 100K where the orthorhombicity actual decreases. While this reduction could represent a simple reversal of the octahedral tilts, we would like to speculate instead that the doped samples are undergoing a phase change from a Bmab structure to a lower symmetry orthorhombic structure in space group Pccn. The resulting structure would be intermediate between the LTO and LT_T structures with tilts described as $Q_1 \neq 0, Q_2 \neq 0, Q_1 \neq Q_2$. Superlattice reflections which would confirm our speculation are notably absent in the x-ray diffraction patterns. Neutron diffraction patterns (NIST, A. Santoro and Q. Huang) are presently being analyzed.

Figure 1.

Figure 2.

 $\text{La}_{2-x}\text{Nd}_x\text{CuO}_4$

ORIENTATIONAL ORDERING PHASE TRANSITION IN C₆₀

P. A. Heiney, J. E. Fischer, A. R. McGhie, W. J. Romanow, A. N. Denenstein, J. P. McCauley Jr., A. B. Smith (U. Penn), D. E. Cox (BNL)

High resolution synchrotron x-ray powder diffraction has revealed the occurrence of a crystallographic phase transition at about 250K in C₆₀ [1]. At 300K the diffraction pattern shows peaks characteristic of a face-centered cubic cell with an edge of 14.16 Å (Fig. 1, top). Below 250K additional peaks are observed which can be indexed on the basis of a simple cubic cell, as illustrated in Fig. 1 (bottom). At 15K the cell edge is 14.04 Å.

The 300K structure can be interpreted in terms of a close-packed fcc lattice of C₆₀ molecules, which have the form of truncated icosahedra (Fig. 2). The unit cell contains four such molecules and the intensities are consistent with a model with Fm3m symmetry in which the molecules are completely orientationally disordered. Below 250K, orientational order sets in such that the molecules are rotated about local <111> axes, the rotation angle being about 25° at 15K. The resulting space group is Pa3.

[1] P. A. Heiney, J. E. Fischer, A. R. McGhie, W. J. Romanow, A. N. Denenstein, J. P. McCauley Jr., A. B. Smith and D. E. Cox, Phys. Rev. Lett. 66, 2911 (1991)

Fig. 2. Molecular structure of C₆₀ projected on a cubic (100) plane.

Fig. 1. Powder diffraction patterns from C₆₀ at 300K (top) and 15K (bottom).

Work at U. Penn supported by NSF MRL Program Grant No. DMR88-19885, NSF Grant No. DMR89-01219 (PAH, JPM and ABS) and DOE Grant No. DE-FC02-86ER45254 (JEF). Work at BNL supported by DOE, DMS, Contract No. DE-AC02-76CH00016.

THE CRYSTAL STRUCTURE DETERMINATION OF BiMo₂O₇(OH)·2H₂O FROM POWDER DIFFRACTION DATA.

J. A. Hriljac (BNL*) and C. C. Torardi (Du Pont).

Synchrotron x-ray powder diffraction data were collected on a flat plate sample of a material of nominal composition BiMo₂O₇(OH)·2H₂O using a wavelength of 1.1988 Å. The reflection positions were consistent with the monoclinic unit cell obtained via auto-indexing laboratory diffraction data, and least-squares refinement gave a = 6.347(1), b = 11.594(1), c = 5.794(1) Å, and β = 113.34° in space group P2₁/m. The intensities of 387 reflections in the range 10<2θ<70° were extracted via an automated procedure and these were used as the input to a direct methods routine¹ which successfully located the Bi and Mo atoms. Alternation of Rietveld refinements and difference Fourier techniques led to the placement of all of the non-hydrogen atoms. In the final model, the severe anisotropic broadening along h0l was included as a least-squares variable and the current agreement factors are: R_{wp} = 16.92%, R_s = 7.27%, and R_{et} = 5.99% for 453 reflections.

The structure is built from distorted octahedral molybdenyl (MoO₂²⁺) units and irregular 6-coordinate Bi atoms. Bond length - bond valence sums (see Table) indicate one of the bismuth bound oxygen atoms, O(1), is part of a hydroxyl group and one of the molybdenum bound ones, O(4), is from a water molecule. This leads to the correct formulation as BiMo₂O₇(OH)·2H₂O. The MoO₆ units form edge-shared dimers which are corner linked to form infinite 1-dimensional chains which run along [010]. These chains are held together via weak (2.515 Å) Bi-O bonds to form sheets, as shown in the figure. The hydroxyl and water molecules project between these sheets, and presumably they are held together via hydrogen bonding.

Atom	Valence
Mo	+5.66
Bi	+3.27
O(1)	-1.11
O(2)	-1.99
O(3)	-1.91
O(4)	-0.25
O(5)	-1.75
O(6)	-1.85

1. G. M. Sheldrick in "Crystallographic Computing 3", eds. G. M. Sheldrick, C. Kruger and R. Goddard, pp 175-189, Oxford University Press, 1985.

* Work supported by the Div. of Materials Sciences, US Dept. of Energy under contract DE-AC02-76CH00016.

THE CRYSTAL STRUCTURE DETERMINATION OF CuPt_3O_6 FROM POWDER DIFFRACTION DATA.

J. A. Hrijiac (BNL*), J. B. Parise (SUNY, Stony Brook), G. H. Kwei (LANL), and K. B. Schwartz (Raychem).

In a continuing effort to elucidate the structures of multi-component copper oxides, the previously reported title material¹ was synthesized and powder diffraction data were collected on a flat plate sample at a wavelength of 1.20030 Å. The observed peak positions were consistent with the reported unit cell (orthorhombic, $a = 6.1154(1)$, $b = 11.560(1)$, $c = 3.1012(1)$ Å), but the observation of a weak 100 reflection revealed the true space group is $Pn2_1m$ rather than $Pnmm$. The integrated intensities of 84 reflections were extracted and used to generate a Patterson map, and manual interpretation led to the initial positions of the Pt atoms. Continued alternation of Rietveld analysis and difference Fourier techniques led to a final structural model. This was then refined against neutron diffraction data collected at HIPD at LANSCE to give more accurate positional parameters for the light atoms.

In common with other members of the MPt_3O_6 ($M = \text{Mn}, \text{Co}, \text{Ni}, \text{Zn}, \text{Cd}, \text{Mg}$) family of materials,² the structure contains two octahedrally coordinated Pt^{4+} and one square planar Pt^{2+} per asymmetric unit. In contrast, the coordination around the copper atom is found to be a Jahn-Teller distorted square pyramid rather than an octahedron (Ni) or distorted square ($\text{Mn}, \text{Co}, \text{Zn}, \text{Cd}, \text{Mg}$). The 3-dimensional structure can be built from the PtO_6 octahedra which are alternately *cis* and *trans* corner-linked to form buckled slabs approximately in (100). The slabs are held together by alternating infinite stacks of the PtO_4 squares and edge-shared CuO_3 groups along [001].

1. O. Muller and R. Roy, *J. Less-Common Metals* **19**, 209 (1969).
2. C. T. Prewitt, K. B. Schwartz, and R. D. Shannon, *Acta Crystallogr.* **B39**, 519 (1983); K. B. Schwartz, J. B. Parise, C. T. Prewitt, and R. D. Shannon, *Acta Crystallogr.* **B39**, 217 (1983).

* Work supported by the Div. of Materials Sciences, US Dept. of Energy under contract DE-AC02-76CH00016.

THE EFFECTS OF HIGH-PRESSURE ANNEALING ON THE PHASE TRANSITIONS IN $\text{La}_{1.88}\text{Ba}_{0.12}\text{CuO}_4$.

J. A. Hrijiac, D. H. Chen, D. E. Cox, and A. R. Moodenbaugh (BNL*)

The high-temperature superconductors of formula $\text{La}_{2-x}\text{Ba}_x\text{CuO}_4$ ($0.05 < x < 0.20$) are known to undergo a series of phase transitions upon cooling. The high-temperature tetragonal structure (HTT, $I4/mmm$) becomes orthorhombic (LTO, Bmm), and then converts to a different low-temperature tetragonal form (LTT, $P4_2/nm$). The appearance of the LTT phase correlates with an anomalous depression of T_c , and this effect is maximized for materials with x ca. 0.125.

In order to determine the effects of high-pressure annealing on these transitions, we have subjected a phase-pure sample of $\text{La}_{1.88}\text{Ba}_{0.12}\text{CuO}_4$ to a pressure of 100 kbar and a temperature of 1000 °C for 1 hour. Magnetization data indicates that the amount of superconducting material increases substantially, with a T_c onset of ca. 28K. High resolution x-ray diffraction data were collected at several temperatures on beam-line X7A at a wavelength of 0.7002 Å, and plots of the 200 reflection for the original (left) and treated (right) samples are shown below. The severe line broadening observed in the treated sample hampers the interpretation of the results, but it appears that the same series of transitions occurs for both. Attempts to modify the anneal conditions in order to prepare a more crystalline sample which will allow quantification of the fractions of LTO and LTT present are currently being performed.

* Work supported by the Div. of Materials Sciences, US Dept. of Energy under contract DE-AC02-76CH00016.

Ruizhong Hu, T. Egami (U. of Pennsylvania) and A. P. Tsai, A. Inoue T. Masumoto (Toboku University)

The atomic structure of $\text{Al}_{48}\text{Ru}_{12}\text{Cu}_{12}$ quasicrystal has been studied by the differential atomic pair distribution function (PDF) analysis of anomalous x-ray diffraction measurements carried out at the Ru K-edge. The goal is to understand the local atomic structure and chemical ordering of the material.

The PDF is obtained from a Fourier transform of the partial structure factor of Ru. This can be obtained by taking the difference between two diffraction measurements taken with incident energies at the Ru edge and below it. These anomalous x-ray diffraction measurements were carried out at 12.6 eV and 300 eV respectively below the Ru K absorption edge which is at E-22.1193 KeV. The energy resolution of the channel-cut Si [111] monochromator was 6.0 eV FWHM. An intrinsic Ge detector was used to separate fluorescent radiation.

The analysis of the total pair distribution function (PDF) and the DDF of Ru illustrates several atomic pair correlations with strong Ru-Ru and Al/Cu-Al/Cu character respectively. The results also show that there are no Ru-Ru Ru-Cu correlations located at the nearest neighbor distances.

A structure model obtained by a direct projection of a 6-dimensional hypercubic lattice with the vertex decoration using a non-Penrose window is shown to account for the results quite well. Compositional ordering is specified by subdividing the window for each composition. The structure of the short range atomic order of this phasless quasicrystal is fundamentally different from the structures of other quasicrystals with phasons.

* Work supported by the National Science Foundation through DMR-8617950.

Figure 1

Figure 1 documents the pressure-load curve for the cell up to diamond failure at 150 GPa: a 1 MPa increment in applied load to the membrane results in a 26 GPa increase in sample pressure. Starting sample size was 0.040 mm; diamond culet 0.1 mm; internal pressure standard: gold; a 0.05 mm collimator was located in the convergent beam before the cell.

Up to the maximum pressure of the study, there was no evidence for a structural transition from the fcc phase. Diffraction from the sample was weak necessitating long counting times (0.7 Å is close to the K absorption edge for Kr). Kr also appears a strong solid with peak widths on the order of $0.15^\circ 2\theta$ at 20 GPa. Figure 2 illustrates the compression data reduced to 0 K with a Mie-Grüneisen thermal model and fitted with a 3rd-order Birch-Murnaghan EOS. The apparent stability of the fcc phase to above 100 GPa should motivate first-principles theoretical calculations of the metallization pressure in Kr, similar to those performed for the other heavy rare-gas solids, where the hcp phase occurs as a precursor to band gap closure.

Figure 2

Research supported under NATO CRG 890944 and the Royal Society, London (APJ); the National Science Foundation grant EAR 8817263 (LWF); and the Division of Materials Sciences, US Department of Energy, under contract DE-AC02-76CH00016.

HIGH-RESOLUTION, HIGH-PRESSURE POWDER DIFFRACTION

A.P. Jephcoat (Oxford U.), L.W. Finger (CIW) and D.E. Cox (BNL)

We have performed compression experiments in a diamond-anvil cell in a monochromatic, high-resolution scattering geometry with a linear, position-sensitive detector (PSD). A compact, diamond-anvil cell (DAC) of the Mao-Bell type was designed to mount at the second axis position of the diffractometer. Pressures were generated with a stainless-steel membrane (LeToullec et al., 1988) coupled to a 150 bar N_2 cylinder external to the hutch. Sufficient loads could be applied to reach pressures in the 100 GPa range without disturbing the sample alignment. The combination of an asymmetrically-cut, bent silicon crystal monochromator and linear PSD provided a moderately high flux at the sample and the advantage of parallel data collection over a maximum window of $10^\circ 2\theta$. At high resolution, the central portion of the order $\pm 1.0^\circ$ can be used. A 10mm horizontal portion of the incident white beam is focused down to about 0.25mm at the second axis position giving a $40:1$ gain in flux at, or slightly below, $0.7^\circ 2\theta$ with zero $\Delta\lambda/\lambda$. A pinhole collimator of 0.05 mm was placed just before the cell. The detector was operated with krypton gas at 4 bar and at a distance of 45 cm. The experiment is designed to explore low-symmetry transitions and the effects, if any, of strain-induced broadening, depending on the use of a pressure medium, in the 100 GPa range. The method contrasts with standard energy-dispersive diffraction techniques where the resolution can be detector-limited at low pressures.

The figure panel shows results for ferroelectric $LiNbO_3$ at 12 GPa: the importance of a pressure medium is shown by the order of magnitude increase in line width obtained with and without solid helium as a medium. Use of a smaller collimator (0.020 mm pinhole) does not result in narrower peak profiles, indicating that pressure gradients are not the dominant factor in lineshape at this pressure.

Research supported under NATO CRG 890944 and the Royal Society; London (APJ); the National Science Foundation grant EAR881726(LWF); and the Division of Materials Sciences, US Department of Energy; under contract DE-AC02-76CH00016.

MICROSTRUCTURAL ASPECTS OF THE LTO-LTT TRANSITION IN $La_{1.875}Ba_{0.125}CuO_4$

M. Marezio (A. T. & T. Bell Labs and CNRS), H. Takagi (A. T. & T. Bell Labs), B. Baolgg (A. T. & T. Bell Labs), P. Bordet (CNRS), D.E. Cox (Brookhaven), and Uchida (U. Tokyo)

Moodenbaugh et al. (Phys. Rev. B38 (1989) 4596) discovered that in $La_{2-x}Ba_xCuO_4$ for $x=0.125$ T_C has a narrow minimum. Axe et al. (Phys. Rev. Letters 62 (1989) 2751, and IBM J. Res. Develop. 33 (1989) 382) found that for this composition, besides the tetragonal to orthorhombic transition (HTT-LTO) at about 210 K, an additional phase transition to a tetragonal phase (LTT) takes place at about 60 K and pointed out that the most important crystallographic feature of the transition is a 45° rotation of the octahedral tilting.

We investigated the LTO-LTT transition by using two differently prepared samples (A spray drying and D solid state) of $La_{1.875}Ba_{0.125}CuO_4$ and high resolution synchrotron x-ray powder diffraction at X-7A (Si(111) monochromator/Ge (220) crystal analyser and 1.29914 as a wavelength). The reflections of sample D are all asymmetric and the best fit was obtained with two (one sharp and one diffuse) peaks. The sharp peak would correspond to domains with a homogeneous composition $x=0.125$ while the domains giving rise to the diffuse peak would have an average composition $x=0.11$. The domains are large enough to yield two separate peaks. On the contrary, all reflections of sample A are symmetric and can only be fitted with one peak. Fig. 1 shows the percentage of the LTT phase as a function of temperature for samples A and D. In both cases the LTO-LTT transformation is complete, however, the transition takes place over a much larger temperature range for sample A than for sample D.

A microstructural model for samples A and D implying the existence of microdomains can be envisaged to explain the different behavior. Because all reflections are larger than the instrumental FWHM, both samples must contain a certain spread in composition. We assume that the broadening due to the crystallite size is negligible with respect to that due to the spread in composition. The sharp and symmetrical reflections of sample A indicate that in this sample the spread is rather small. Instead, sample D must contain two types of domains, one homogeneous and the other less so, and each with a different average composition and both larger than the x-ray diffraction correlation length (about 200 Å).

If the geometrical component is an important factor for the LTO-LTT transition then an appreciable strain release must occur at the transition. In sample D the domains are small and the homogeneous ones are likely to be surrounded by inhomogeneous domains where the transition does not take place. These domains could act as strain absorbers and the transition occurring in the homogeneous domains becomes sharper. There is no strain release in sample A because the entire sample undergoes the phase transition. These results are corroborated by electron microscopy results (Chen et al. Physica C 175, 301, 1991).

Figure 1. The percentage of the LTT phase vs. T

PRELIMINARY STRUCTURAL INVESTIGATIONS OF THE SOME UNIVALENT CATION EXCHANGED FORMS OF THE RHO FRAMEWORK

John B. Parise, D. R. Corbin, L. Abrams (DuPont), T. M. Nenoff and G. D. Stucky (Santa Barbara).

The structures of the dehydrated, Ti^{4+} -exchanged forms of $SiAlO$, $BeAsO$ and $BePO$ -RHO have been refined using either neutron or synchrotron x-ray powder diffraction data coupled with interatomic distance constraints; the structure of Rb^{+} -exchanged $BePO$ -RHO has also been refined. All materials, except Ti - $SiAl$ -RHO, crystallizes in the space group $I\bar{2}3$, with $a = 13.4791(5)$ Å for Rb - $BePO$ -RHO, $a = 13.6465(6)$ Å for Ti - $BePO$ -RHO and $a = 13.998(1)$ Å for (Rb,Ti) - $BeAsO$ -RHO; for Ti - $SiAl$ -RHO ($I\bar{4}3m$) $a = 14.4636(7)$ Å. The extra framework cations are distributed over the six double 8-ring sites (D8R) and two single 6-ring sites (S6R). Once normalized, by taking account of differences in ionic radius sums, all materials with the RHO framework studied thus far, fall on the same universal curve describing the distortion of elliptical double 8-ring pore opening as a function of unit cell parameter.

Figure 1. Scatter plot of cubic unit cell edge (a) versus Δ/a for dehydrated materials with the framework of RHO. Data taken from references 1 and 12.

ANOMALOUS DIFFRACTION STUDY OF $Pb(Mg_{1/3}Nb_{2/3})O_3$

H. D. Rosenfeld and T. Egami (U. of Pennsylvania)

We are investigating the local atomic structure of the relaxor ferroelectric $Pb(Mg_{1/3}Nb_{2/3})O_3$ (PMN). The diffuse nature of the ferroelectric phase transition in the perovskite PMN has been attributed to local variations in the distribution of niobium atoms on the B site sublattice¹. In order to determine this local distribution, we have obtained the niobium differential distribution function in PMN at room temperature by the method of anomalous diffraction². Diffraction data for a powder sample of PMN were obtained at 30 eV and 300 eV below the niobium absorption edge at 18.989 keV on beam line X-7A of the NSLS. Structure factors were obtained for these two energies, and the differential of $S(q)$ for niobium was determined. Fourier transformation of this differential $S(q)$ yielded the Nb differential distribution function (Nb-DDF), which describes the spatial distribution of atoms surrounding niobium.

While we are still in the process of a detailed interpretation of this data, we can at this time make two important observations. The difference of the structure factors for the two energies reveals a bcc selection rule (Fig. 1). Positive peaks satisfy the bcc selection rule $h+k+l$ odd. Since only the scattering amplitude of Nb varies strongly in this energy region, this suggests that some fraction of the niobium atoms must be arranged in a $1r$ bcc configuration. We note that this is not a superlattice configuration, but rather has a lattice constant close to that determined for PMN by standard diffraction methods.

Inspection of the Nb-DDF reveals an unexpected feature (Fig. 2). There is a split peak centered at 2.68 Å. This peak does not correspond to any interatomic distance involving niobium, based on niobium occupying the B sublattice. It is, however, close to what one would expect for the A-O nearest neighbor distance in a cubic perovskite with a lattice constant of 3.8 Å. Assuming a model based on niobium partially occupying the A and B sites of a cubic perovskite of lattice constant 3.8 Å, additional features in the Nb-DDF at larger r , inconsistent with the crystallographically determined structure of PMN, may be explained in addition to explaining the bcc like selection rule observed in the difference $S(q)$. Detailed modelling of the local distribution of niobium in PMN based on the Nb-DDF will be carried out in the near future.

¹ G. A. Smolensky, J. Phys. Soc. Jap. 28, 26 (1970)

² D. D. Kofalt, Ph. D. Dissertation, U. Pennsylvania, 1988

Fig. 1

Fig. 2

Work supported by NSF contract DMR88-19885 and ONR contract N00014-91-J-1036.

THE LOCAL ATOMIC STRUCTURE OF $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})\text{O}_3$

H. D. Rosenfeld and T. Egami (U. of Pennsylvania)

We are investigating the relationship between local atomic structure and the diffusive ferroelectric phase transition in the perovskite $\text{Pb}(\text{Mg}_{1/3}\text{Nb}_{2/3})\text{O}_3$ (PMN) by the method of pair-distribution function (PDF) analysis. The PDF is the Fourier transform of the square of the crystallographic structure factor. No assumption of crystallographic periodicity is required in analyzing the PDF, and consequently short-range order in aperiodic structures may be treated exactly, rather than as an approximation.¹ Since the Fourier transform includes diffuse as well as Bragg scattering, information regarding local atomic pair correlations is included in the PDF. The Fourier transform need not be carried out to q equals infinity, because significant contributions to coherent scattering are contained within a Debye envelope, due to atomic motions and static displacements of atoms from their ideal crystallographic sites. The range of q space accessible using the high energy x-rays available at the NSLS is sufficient for an accurate Fourier transform.

We have shown¹ that the local atomic structure suggested by the PDF obtained by neutron powder diffraction is inconsistent with the crystallographically determined structure of PMN. Initial modelling of the local atomic structure suggests that lead is shifted approximately 0.5 Å from its crystallographic site, and that these shifts, while lacking long range periodicity, are locally correlated resulting in short-range order. We have studied the temperature dependence of the PDF in the vicinity of the Curie temperature (-7°C) using x-ray powder diffraction at beam line X-7A of the NSLS. Data were collected at 250K, 270K, 300K and 323K. Significant temperature dependence is observed over the range 2-5 Å (See figure). All peaks in this region are contributed to by atomic pairs involving lead (Pb-O, Pb-Pb, Mg/Nb, and Pb-O 1st and 2nd nearest neighbors). This suggests that changes in short-range order involving lead atoms are related to the bulk ferroelectric phase transition in PMN. We plan to develop a detailed model of these correlations and their temperature dependence, as well as studying the local distributions of oxygen, magnesium, and niobium using both x-ray and neutron powder diffraction data. This effort will be aided by the contrast provided by the differences in x-ray scattering amplitudes of the species comprising this compound, as compared to the relatively uniform neutron scattering lengths.

1 H. D. Rosenfeld, T. Egami, A. Shalla, IEEE Trans. UFFC, Nov. 1991 (in press)

Work supported by NSF contract DMR88-19885 and ONR contract N00014-91-J-1036.

HIGH RESOLUTION SYNCHROTRON X-RAY DIFFRACTION STUDY OF PHASE SEPARATIONS IN $\text{Pr}_2\text{NiO}_{4-\delta}$

J. D. Sullivan, D. J. Buttery (U. of Delaware), D. E. Cox, and J. Hriljak (BNL)

The influence of oxygen nonstoichiometry on the structure of $\text{Pr}_2\text{NiO}_{4-\delta}$ has been investigated with high resolution X-ray diffraction at several compositions and temperatures. Similar to other oxides with the K_2NiF_4 -type structure, including the high-Tc superconducting copper oxides, $\text{Pr}_2\text{NiO}_{4-\delta}$ shows multiple phases of distinct oxygen composition with a complex temperature dependence. In this investigation, several samples of varying oxygen content were examined with synchrotron radiation at temperatures between 15 and 1000K. These samples had been previously annealed under various oxygen pressures at temperatures above 1200K for at least six hours to establish the degree of oxygen nonstoichiometry. The samples were rapidly quenched to room temperature to minimize additional oxygen exchange with the environment, and were subsequently isodometrically fired to determine the amount of excess oxygen.

The room temperature oxygen concentration phase diagram for $\text{Pr}_2\text{NiO}_{4-\delta}$ developed from these results shows four distinct phases, separated by regions of phase coexistence. On the basis of observed superlattice reflections and previous neutron diffraction studies (1), the most stoichiometric phase ($\delta < 0.01$) has been assigned orthorhombic Pccn symmetry. This Pccn phase undergoes a low temperature phase transition to tetragonal $\text{P4}/\text{mnm}$ symmetry (118K), and can coexist with a $\delta = 0.06$ phase of tetragonal $\text{P4}/\text{mnm}$ symmetry. The orthorhombic structure and the absence of superlattice reflections necessitates the assignment of Fmmm symmetry to the two most nonstoichiometric phases, containing $\delta = 0.17$ and $\delta = 0.21$. No low temperature phase transitions were observed down to 15K for either of the Fmmm phases, or the $\delta = 0.06$ tetragonal phase. The room and low temperature phase diagram data are published in the Journal of Solid State Chemistry (2). High temperature (300-1000K) synchrotron diffraction experiments were also done isothermally on several samples that were sealed in evacuated quartz capillaries. The preliminary results show first and second order phase transitions occurring in samples upon heating, culminating in a thermally randomized (no coherent octahedral distortions) tetragonal structure (sp. gr. $\text{I4}/\text{mmm}$). The transition temperatures appear to be a function of the oxygen nonstoichiometry, but are not always reversible on cooling. Further investigation of the irreversibility and the nature of the high temperature phases is in progress in our ongoing efforts to elucidate the nature of phase separation phenomena in layered perovskites.

1. D. J. Buttery, J. D. Sullivan, G. Shirane, and K. Yamada, Phys. Rev. B 42, 3944 (1990).
2. J. D. Sullivan, D. J. Buttery, D. E. Cox, and J. Hriljak, J. Solid St. Chem. 94(2), 337 (1991).

RE-EXAMINATION OF THE PHASE DIAGRAM IN HEAVILY-DOPED $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$

H. Takagi, M. Marezio, R. J. Cava, B. Ballogg (AT&T Bell Labs), P. Bordet, J. Hodeau (CNRS) and D. E. Cox (BNL)

The reason for the disappearance of superconductivity in heavily doped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ is still not known, in spite of its importance to the understanding of high- T_c superconductivity in general. A close connection between delicate structural distortions and high- T_c has been suggested by the observation of the suppression of T_c associated with the low temperature orthorhombic (LTO)/low temperature tetragonal (LTT) structural transformation in $\text{La}_{1.875}\text{Ba}_{0.125}\text{CuO}_4$. This structural transformation is reminiscent of the presence of the composition-induced high temperature tetragonal (HTT)/low temperature orthorhombic (LTO) phase boundary in heavily doped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$.

Although most earlier work has claimed that the HTT/LTO boundary has nothing to do with the disappearance of superconductivity, these studies were limited by insufficient diffraction resolution and/or uncontrolled sample quality. We have therefore re-examined the structural and electronic phase diagram of heavily doped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ on carefully prepared samples by combining high resolution synchrotron x-ray diffraction and sensitive magnetic measurements.

A series of $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ samples ($0.15 \leq x \leq 0.3$) was carefully prepared by conventional ceramic processing techniques. In order to examine the possibility of phase separation, as is frequently suspected to occur in this composition range, the samples were further annealed for one month at 850°C and 1000°C. A subsequent high resolution x-ray diffraction study revealed that the diffraction peaks shifted systematically upon doping, and that their width remained essentially unchanged up to $x \approx 0.3$. We therefore concluded that the La-Sr solid-solution exists at least up to $x \approx 0.3$.

In these high quality samples, we determined the phase boundary between HTT and LTO, on the x - T plane, using high resolution x-ray diffraction at temperatures between 10K and 300K. This is the first detailed study of this kind. As is seen in Fig. 1, which shows the FWHM of the tetragonal (200) reflection, we observed evidence for the transition to LTO up to $x \approx 0.2$. In the magnetic measurements, we found a significant reduction of the Meissner signal near $x \approx 0.2$. This combined study thus revealed (1) a much narrower composition range for superconductivity in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$, and (2) coincidence of a structural phase boundary with the disappearance of superconductivity. We are finally beginning to get insight into this longstanding and complex problem.

Fig. 1 Temperature dependence of the FWHM of the tetragonal (200) in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$

ORIENTATIONAL DISORDER IN SOLVENT-FREE SOLID C_{70}

G. B. Vaughan, P. A. Heiney, J. E. Fischer, D. E. Luzzi, D. A. Rickes-Foot, A. R. McGhie, Y. W. Hui, A. L. Smith, W. J. Romanow, B. H. Allen, N. Coostel, J. P. McCauley Jr., A. B. Smith (U. Penn), D. E. Cox (BNL)

High resolution synchrotron x-ray powder diffraction data were collected on C_{70} [1]. A sample crystallized from toluene which contained an appreciable amount of solvent gave a diffraction pattern that could be indexed on a hexagonal cell with $a = 10.63$, $c = 17.39$ Å (Fig. 1a), suggestive of a hcp structure containing two C_{70} molecules per cell. A sample sublimed at 810 K showed a qualitatively similar pattern with significantly sharper peaks (Fig. 1b) consistent with a mixture of fcc ($a = 15.01$ Å) and hcp ($a = 10.62$, $c = 17.31$ Å) phases. Since all the strong peaks from the two phases coincide, this indexing was based on uniquely indexed weak peaks such as the fcc (200) at 0.837 Å $^{-1}$ and the hcp (102) at 0.936 Å $^{-1}$. A sample prepared by sublimation at 810 K and annealed at 575 K for 200 hours was still two-phase but with a much lower proportion of the hcp component (Fig. 1c).

Fits based on a model of complete orientational disorder of the C_{70} molecule (shown in the inset at the bottom of Fig. 1) gave good agreement with the observed pattern shown in Fig. 1c. The calculated pattern is shown in Fig. 1d, the individual contributions from the hcp and fcc phases in Figs. 1e and 1f respectively, and diffuse scattering due to stacking disorder and the glass sample capillary in Fig. 1g. The logarithmic plots shown in Figs. 1c and 1d are reproduced on a linear scale in Fig. 1h.

Fig. 1. Observed and calculated powder diffraction patterns from C_{70} as described in text.

[1] G. B. Vaughan, P. A. Heiney, J. E. Fischer, D. E. Luzzi, D. A. Rickes-Foot, A. R. McGhie, Y. W. Hui, A. L. Smith, D. E. Cox, W. J. Romanow, B. H. Allen, N. Coostel, J. P. McCauley Jr., A. B. Smith, Science (submitted).

Work at U. Penn supported by NSF MRL Program Grant No. DMR88-19885, NSF Grant No. DMR89-01219, and DOE Grants No. DE-FC02-86ER45254 and DE-FG05-90ER75596. Work at BNL supported by DOE, Division of Materials Sciences, Contract No. DE-AC02-76CH00016.

ISOTHERMAL COMPRESSION OF Fe_2SiO_4 (SPINEL) AT PRESSURES UP TO 24 GPa AND AT ROOM TEMPERATURE

L.-J. Wang, J.-A. Xu, L. C. Ming, J. Balogh and M. H. Manghnani (University of Hawaii at Manoa)

In view of a suggestion that Fe_2SiO_4 (spinel) would change its equation of state at pressures above 12 GPa, we have remeasured these material at pressures up to 24 GPa and at room temperature using a diamond-anvil cell and the energy dispersive x-ray diffraction method. A mixture of 4 methanol and 1 ethanol was used as pressure medium in the sample chamber, and Au was used as an internal pressure calibrant. Experiment was carried out with beam line 7A at NLSL. Examining the data indicated that those above 18 GPa appear to be slightly stiffer (Figure 1). Assuming the pressure derivative of the zero-pressure bulk modulus to be 4 in the Birch-Murnaghan equation, the zero-pressure bulk modulus derived from those data below 18 GPa is 191 ± 2 GPa, which is in excellent agreement with those reported previously (Wilburn et al., 1976; Sato, 1977) and those above 18 GPa is 215 ± 2 GPa, which is much smaller than the value of 265 GPa reported by Bi et al., 1991. It seems to us that those stiffer data at higher pressures are probably due to the nonhydrostatic condition rather than to the different equation of state as proposed by Bi et al., 1991.

Figure 1

1. Wilburn, D.R. and W.A. Bassett, *High Temp. High Pressure*, 8, 343-348, 1976.
2. Sato Y., Yagi T., Ida Y. and Akimoto S., *High Temp. High Pressures*, 7, 315-323, 1975.
3. Bi, Q., J.M. Brown and Sato S., 1991., preprint

We wish to thank D. E. Cox for his assistance on instrumentation and data acquisition during our experiment. This work was supported by NSF Grant EAR-8708482.

HIGH TEMPERATURE BEHAVIOR OF $(\text{Mg}, \text{Fe})\text{SiO}_3$ PEROVSKITE AT 1 BAR

Y. Wang, Y. Zhao, D. J. Weidner, R. C. Liebermann, J. B. Parise (CHPR and ESS, Stony Brook) and D. E. Cox* (Physics, BNL)

$(\text{Mg}, \text{Fe})\text{SiO}_3$ -perovskite dominates the properties of the Earth's lower mantle and constitutes as much as half of the earth's volume. We have carried out a high-temperature synchrotron x-ray diffraction study on MgSiO_3 and $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{SiO}_3$ perovskite at X7 A beamline of Brookhaven National Light Source with a position sensitive detector. Perovskite specimens were synthesized in the 2000-ton Uniaxial Split-Sphere Apparatus (USSA-2000) at 26 GPa; and 2173 K. MgO was used as an internal standard and diffraction patterns were collected on both heating and cooling cycles. Rietveld refinements were performed simultaneously on both the perovskite and MgO . Both MgSiO_3 and $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{SiO}_3$ perovskites retained orthorhombic symmetry (space group $Pbnm$) throughout the temperature range in our study.

Because of the metastability of the silicate perovskite under ambient conditions, changes in defect concentration and/or partial conversion to an amorphous phase might be expected to occur which could cause abnormal behavior at high temperatures.

A slight decrease ($< 0.1\%$) in the room-temperature unit-cell volume was observed for MgSiO_3 after being heated to 404 K and diffraction became significantly weaker above this temperature. In subsequent heating cycles, however, the cell parameters were reversible within the error. Overall, thermal expansivities were reversible up to 717 K. The volumetric thermal expansion (α_v) can be satisfactorily fitted by a Suzuki-type formula, with $\alpha_v = 1.4 (10^{-5} \text{ K}^{-1})$ near 300 K and 2.4 near 710 K. Crystals converted completely to an amorphous phase near 740 K.

For $\text{Mg}_{0.9}\text{Fe}_{0.1}\text{SiO}_3$, the room-temperature unit-cell volume was reduced by 0.25% after being heated to 472 K. Below 472 K α_v was reversible within the error, with values similar to those of MgSiO_3 . Above 472 K, α_v became significantly higher than that of MgSiO_3 ; complete amorphization occurred near 660 K.

*DOE contract No. DE-AC02-76CH00016

COMPARATIVE STUDY OF ELEMENTAL CONTRAST BY NEUTRON AND RESONANT X-RAY POWDER DIFFRACTION.*

J.K. Warner, A.P. Wilkinson, A.K. Chetani (Oxford U.) and D.E. Cox (BNL)

There is a long standing tradition of using powder neutron diffraction methods to distinguish between elements that are adjacent in the periodic table and therefore difficult to distinguish with X-rays. Recent studies have shown that anomalous scattering can also provide contrast between elements of similar atomic number, and the method can be applied to any element that has an absorption edge at a wavelength that is suitable for diffraction and is accessible with synchrotron X-rays. In this site ordering study of transition metal phosphates, occupancies obtained using anomalous scattering at the Fe K-edge are compared to those from high resolution neutron powder diffraction.

Farringtonite, $M_3(PO_4)_2$ contains two distinct metal sites, which are 5- and 6- coordinate with multiplicities of 4 and 2 respectively. Of the range of possible solid solutions, $Co_2Fe(PO_4)_2$ and $Zn_2Fe(PO_4)_2$ were selected to measure the distribution of iron over these two sites, in each case offering excellent elemental contrast for neutron scattering ($b(Fe)=0.954 \times 10^{-14}m$, $b(Co)=0.253 \times 10^{-14}m$ and $b(Zn)=0.568 \times 10^{-14}m$). $Fe_3(PO_4)_2$ *graffonite* was chosen as an X-ray standard to measure the value of f for Fe in a chemically similar environment. An incident energy of around 9 eV below the absorption edge, measured for these materials using XANES, was selected to provide a large, negative value of f (-7.81(9) electrons) whilst keeping f'' and hence absorption low. Table 1 gives the results of the Rietveld refinements for Fe occupancies over the two sites, with the overall composition constrained to the chemical formula. The values obtained using X-rays for Fe/Zn ordering are not as good as the Fe/Co case as the absorption edge measured by XANES was 0.7 eV higher in energy than that of the f standard $Fe_3(PO_4)_2$, compared to 0.3 eV for the Fe/Co case. This demonstrates that the use of a standard material to determine f is limited to materials which have edges very close in energy. At energies more than 15 eV below the Fe K absorption edge, the theoretical free atom values of f seem adequate and it is only in the near-edge region where coordination environment effects change the shape of the f' curve that the use of a chemically similar standard is recommended.

The good agreement and comparable relative precision found using neutrons and anomalous scattering indicates the value of using resonant X-ray powder diffraction in cases where the use of neutrons is not feasible.

Table 1. Occupancy of Fe over metal sites in *Farringtonite*.

Compound	Method	M1 Occupancy	M2 Occupancy
$Co_2Fe(PO_4)_2$	Neutron TOF	0.251(4)	0.498(8)
	X-ray Fe K-edge	0.243(5)	0.514(10)
$Zn_2Fe(PO_4)_2$	Neutron TOF	0.073(5)	0.854(9)
	X-ray Fe K-edge	0.109(5)	0.783(10)

Work at Brookhaven supported by the Div. Mater. Sci., US Dept. of Energy, under Contract No. DE-AC02-76CH00016.

*Accepted for publication in *J. Phys. Chem. Solids* 33, 1991.

FeII/FeIII VALENCE CONTRAST BY RESONANT X-RAY POWDER DIFFRACTION

J.K. Warner, A.K. Chetani (Oxford U.) and D.E. Cox (BNL)

One of the exciting opportunities available with anomalous scattering is the possibility of distinguishing between different valence states of the same element. The value of f' obtained for an element at an energy close to its absorption edge is very sensitive to its valence state and the immediate chemical environment. The studies of Sarode et al¹ on the x-ray absorption spectra of transition metal compounds indicate a difference of around 3 eV between the absorption edges of FeII and FeIII. This is in agreement with a series of XANES spectra of divalent and trivalent iron phosphates we have collected at SRS Daresbury Lab. (UK) which show an average difference of 2.6(4)eV. X-ray powder diffraction data were collected on α - Fe_2PO_5 at wavelengths below and in the proximity of its K-edge for Fe (7120.9(5)eV) on Station 8.3 at Daresbury in an attempt to distinguish between the FeII and FeIII sites. Although a significant difference was detected between the scattering power of the two Fe sites, the bandpass of the incident energy was comparable to the expected difference between the edge positions. Another on-edge powder diffraction pattern was collected using a different instrumental configuration and smaller energy band pass on the diffractometer X7A (NSLS, Brookhaven) to confirm the findings of the previous experiments.

α - Fe_2PO_5 is a mixed valence compound (S.G. = $Pnma$, $a=7.378\text{\AA}$, $b=6.445\text{\AA}$, $c=7.471\text{\AA}$, $Z=4$) in which FeII and FeIII are located in two different crystallographic sites. The octahedron surrounding FeII form chains parallel to the b-axis and share edges. The octahedra containing FeIII are connected on both sides of these chains, alternating with the PO_4 tetrahedra. A multiple data set Rietveld refinement (GSAS) was performed using a common structural model, but allowing instrumental parameters and values of f' to vary with each experiment. The table summarises the experimental conditions, and gives details of the values obtained for f' on the FeII and FeIII sites. The most significant finding is that the two sites can clearly be distinguished as having different scattering powers close to the absorption edge, and that this effect is observable even when the energy spread of the incident beam is comparable to the difference in the edge positions.

Diffractometer	8.3	8.3	X7A
Wavelength (Å)	1.78316(4)	1.74179(4)	1.74149(1)
Estimated Band Pass (eV)	3.1(2)	3.1(2)	1.6(1)
Monochromator (Ch. cut)	Ge(111)	Ge(111)	Si(111)
Slits	Soller	Soller	single
Detector	scint.	scint.	solid state
ΔE from Fe^{2+} edge (eV)	-1.67(8)	-2.6(5)	-1.4(3)
f' (FeII) (electrons)	-3.57(4)	-9.74(5)	-9.33(6)
f' (FeIII) (electrons)	-3.56(5)	-8.04(7)	-7.81(7)
$\Delta f'$ (FeII-FeIII)	+0.01(9)	-1.70(12)	-1.52(15)

Work at Brookhaven supported by the Div. Mater. Sci., US Dept. of Energy, under Contract No. DE-AC02-76CH00016.

1. P.R. Sarode, S. Ramasesha, W.H. Madhusudan and C.N.R. Rao, *J. Phys. C: Solid State Phys.*, 12, 2439 (1979).

REPRODUCIBILITY OF SITE OCCUPANCIES OBTAINED FROM RESONANT X-RAY POWDER DIFFRACTION.

J.K. Warner, A.P. Wilkinson, A.K. Cheetham (Oxford U.) and D.E. Cox (BNL).

Several site ordering studies have been published which make use of anomalous scattering to enhance the x-ray contrast between elements of similar atomic number. Neutron powder diffraction can often be used as an independent method for confirming these results, generally providing good scattering contrast, particularly between members of the first transition series. One exception is nickel and iron, where the similarity in neutron scattering lengths ($b(\text{Ni})=1.030 \times 10^{-14} \text{m}$ and $b(\text{Fe})=0.954 \times 10^{-14} \text{m}$) makes them virtually indistinguishable. Anomalous scattering is the only diffraction technique capable of distinguishing between these atoms, with an appropriate choice of wavelength providing more than 6 electrons difference in scattering power. To test the reproducibility of this technique, experiments were performed using the powder diffractometers X7A (BNL) and 8.3 (SRS, Daresbury Lab, UK) at energies close to the Fe K-edge. Differing instrumental configurations and proximity to the absorption edge were used to provide a wide range of experimental conditions.

The sample, $\text{NiFe}_2(\text{PO}_4)_2$, is isostructural with the mineral *sarcopside*, consisting of chains of edge shared octahedra, similar to those of *olivine*, but with every fourth octahedra missing. The iron and nickel distribute themselves over the two metal sites, with the central M1 cation possessing a fairly regular (-1) symmetry and smaller volume compared with the distorted sites of the two terminal M2 octahedra. A standard, $\text{Fe}_3(\text{PO}_4)_2 \cdot \text{graffiotite}$ was chosen to measure the value of f for Fe to be used in the subsequent refinement. All other values of f and r were calculated from free details of the values obtained for f and Fe occupancies from the two sets of experiments. The K absorption edge of Fe in $\text{NiFe}_2(\text{PO}_4)_2$ was measured using XANES, and the overall composition was constrained to the chemical formula. The excellent agreement of the cation occupancies obtained by this technique with different experimental conditions and energies highlights the versatility and reproducibility of resonant x-ray powder diffraction for site ordering studies of this kind.

Table 1. Summary of instrumental conditions and occupancies obtained for Fe in $\text{NiFe}_2(\text{PO}_4)_2$.

Diffractometer	X7A	8.3
Wavelength (Å)	1.7437	1.7433
Band Pass (eV)	1.6	3.0
Monochromator	Si(111)	Ge(111)
Slits	single	single
ΔE (Fe ²⁺ edge)(eV)	-9(1)	-8(1)
$f(\text{Fe})$ (electrons)	-7.81(9)	-10.7(3)
$f(\text{Ni})$ (electrons)	-1.741	-1.742
Occ. Fe [M1]	0.363(5)	0.376(3)
Occ. Fe [M2]	0.818(3)	0.812(2)

Work at Brookhaven supported by the Div. Mater. Sci., US Dept. of Energy, under Contract No. DE-AC02-76CH00016.

ATOMIC IN-PLANE DISORDER IN CO/PT SUPERLATTICES

X. Yan and T. Egami (U. of Pennsylvania), E. E. Marinero, R. F. C. Farrow, and C. H. Lee (IBM ARC)

Description of multilayered films grown by molecular beam epitaxy (MBE) frequently ignores the effect of structural defects. However, uniaxial structural defects are observed in magnetic Co/Pt superlattices which could be important in producing anisotropic physical properties such as magnetic anisotropy.

X-ray diffraction experiments on both films were performed with the scattering vector Q covering a plane having both components perpendicular ($Q_\perp$) and parallel ($Q_\parallel$) to the film plane. In the MBE grown $\text{Co}(3\text{\AA})/\text{Pt}(1\text{\AA})$, the diffraction peaks corresponding to the atomic arrangement are shown in Fig. 1 for the $\langle 111 \rangle$ oriented sample. It is interesting to note that the peak profile for the scattering vector parallel to the film plane ($Q_\parallel$) has a broad tail. The nature of the broadening was found to be orientation-dependent, characterized mostly by a Gaussian peak shape for the $\langle 001 \rangle$ oriented sample, and by a Lorentzian shape for the case of the $\langle 111 \rangle$. The Lorentzian shape can be best explained as an exponentially decayed $1/r$ strain field created by uniaxial structural defects. As a result, there are small domains among which an atomic orientational correlation is maintained within the film plane but a positional correlation is not. This is believed to be the first three-dimensional analogy of the hexatic phase which exists during a two-dimensional melting.

Fig. 1 High angle diffraction intensity distribution in the Q - $Q_\parallel$ plane for the $\langle 111 \rangle$ oriented $\text{Co}(3\text{\AA})/\text{Pt}(18\text{\AA})$ superlattice.

This work is supported by IBM through the IBM shared University Research program, 810312.

ATOMIC INTERDIFFUSION IN CO/PT SUPERLATTICES

X. Yan and T. Egami (U. of Pennsylvania), E. E. Marinero, R. F. C. Farrow, and C. H. Lee (IBM ARC)

On Co(3Å)/Pt(18Å) superlattices grown by molecular beam epitaxy from the GaAs single crystal substrate, synchrotron x-ray diffraction experiments were performed using a monochromatic beam (15.038 keV) in a symmetric θ - 2θ reflection geometry. A position-sensitive detector was used to measure the diffraction intensity as a function of the scattering vector Q having both components perpendicular (Q_z) and parallel (Q_y) to the film plane.

As shown for the $\langle 111 \rangle$ films in Fig. 1, the superlattice diffraction peaks were found to have a large width parallel to the plane of the film, indicating strong in-plane disorder. To evaluate more accurately the amount of interdiffusion at the interface in the presence of interfacial roughness, the broad peaks were integrated for both $\langle 111 \rangle$ and $\langle 001 \rangle$ oriented films. Contrary to the conventional low angle diffraction, it was found that the extend of Co between the Pt layers is similar (6.4 Å for the $\langle 001 \rangle$ and 6.9 Å for the $\langle 111 \rangle$), indicating that the previously proposed CoPt₂ alloys at the interface can not be extensive. Thus the CoPt₂ should not be the origin for the magnetic anisotropy observed only in the $\langle 111 \rangle$ oriented films.

Fig. 1 Low angle diffraction intensity profiles in the Q_z - Q_y plane for the $\langle 111 \rangle$ oriented Co(3Å)/Pt(18Å) superlattice.

This work is supported by IBM through the IBM shared University Research program. 810312.

PEROVSKITE STRUCTURE AS FUNCTION OF TEMPERATURE & PRESSURE: THE RESULTS FROM NAMGF₃ PEROVSKITE

Y. Zhao¹, D.J. Weidner¹, J.B. Parise¹, D.E. Cox² (1. Mineral Physics Institute, Dept. of ESS, SUNY, Stony Brook, NY 11794; 2. Physics Department, BNL, Upton, NY 11973)

We report the results of an *in-situ* high temperature and high pressure X-ray powder diffraction study conducted on NaMgF₃ perovskite using synchrotron radiation (X7A position sensitive detector and X17B superconducting wiggler port, NSLS, Brookhaven National Laboratory). Thermal expansion and compression data of the NaMgF₃ perovskite agree well with the proposed model that the temperature and pressure induced volume change of the orthorhombically distorted ABX₃ perovskite can be empirically expressed as a combination of the change of the [B-X] bond length and the change of tilting of the BX₆ octahedral framework. We find that NaMgF₃ perovskite transforms directly from orthorhombic $Pbnm$ to cubic $Pm\bar{3}m$ at a temperature of $T_c = 765^\circ\text{C}$ at atmospheric pressure. The tilting angle of the MgF₆ octahedral framework rapidly decreases, in the manner expected for ferroelastic phase transition, toward zero as the temperature approaches T_c . More interestingly, the [Mg-F] bond lengths of the MgF₆ octahedra apparently shrink throughout a temperature interval of about 100°C before the transition. At $P = 6.3$ GPa, the apparent [Mg-F] bond length (as derived from the unit cell assuming regular octahedra) decreases gradually with increasing temperature and contributes a *negative* thermal expansion to the overall volumetric thermal expansion. For centrosymmetrically distorted perovskites, the excess thermal expansion of the orthorhombic phase with respect to its prototype cubic phase may be separated into two contributions: octahedral tilting and octahedral expansion/shrinking. Applying this concept to silicate perovskite, the density and hence the iron content of the Earth's lower mantle as deduced from the seismically determined density is very sensitive to the actual state of distortion of perovskite in the lower mantle.

Acknowledgement: Work at X-7A beam line of NSLS supported under DOE contract DE-AC02-76CH 00016

HIGH PRESSURE SINGLE CRYSTAL X-RAY DIFFRACTION INVESTIGATIONS OF CaF_2 : DATA COLLECTION AT 6.7 GPa AND AT NORMAL PRESSURE.

S. Åsbrink*, J. C. Hanson†, M. Malinowski‡ and A. Waskowska*,

*Department of Inorganic Chemistry, Arrhenius Laboratory, Stockholm University, S-10691 Stockholm, Sweden

†Department of Chemistry, Brookhaven National Laboratory, Upton, NY 11973, USA

‡Department of Crystallography, Institute of Low Temperature and Structure Research, Polish Academy of Sciences, PL 50950 Wrocław, Poland

X-ray diffraction data were collected from a single crystal of CaF_2 at NSLS, station X7B, using a wavelength of 0.92432 Å. The sample was a thin plate of approximate dimensions 90x60x5 µm. It was placed in a cylindrical hole of diameter 110 µm in an inconel gasket in a diamond anvil cell. The cell, constructed by M. Malinowski, is a larger and improved version of the one previously described (Malinowski, 1987)¹ allowing pressures up to 500 kbar. As a pressure fluid a mixture of methanol and ethanol in the ratio 4:1 was used. No pressure standard was needed as the unit cell dimension of CaF_2 had formerly been determined as a function of pressure up to the phase transition point at 9.5 GPa (Gerward et al. 1991; Malinowski et al. 1990).^{2,3}

Data were collected at 6.7 ± 0.1 GPa and, as a control, at normal pressure, the temperature being kept at 25°C. 30 reflections, with $2\theta_{\text{max}}=66^\circ$, were investigated, many at several different azimuthal angles (ψ), giving a total number of data of 71. For each registration both ω - and ω -2 θ -scans were used. The measurements at several ψ -values were undertaken to facilitate a proper absorption correction of the reflection intensities. For each pressure the transmission function was registered as a function of χ - and ω -angles (cf Malinowski, 1987).¹ The computer treatment of the data is in progress.

References

1. M. Malinowski, J. Appl. Cryst. 20, 379 (1987).
2. L. Gerward, J. Staun Olsen, S. Steenstrup, M. Malinowski, S. Åsbrink and A. Waskowska, in preparation.
3. M. Malinowski, S. Åsbrink and A. Waskowska, NSLS, BNL, Annual Report 1990.

Supported in part by the Swedish Natural Science Research Council under contract R-RA2877-307, and by U.S. Department of Energy, Division of Chemical Sciences, under contract number DE-AC02-CH00016.

SINGLE CRYSTALS WITH INTERCALATED PROTEINS FROM BIOMINERALIZED MATERIALS.

A. Berman, I. Addadi, L. I. Eisenowitz, S. Weiner (Weizmann Inst. Rehovot, Israel), J. Hanson, T. Koetzle (NSLS).

In biomineralization, composite materials with unique properties are built by organisms out of crystals and biological macromolecules. Sea urchins form macroscopic spines and plates out of single calcite crystals. Acidic proteins, preferentially adsorbed on (110) planes during crystal growth in amounts of 0.02%, appear to mechanically reinforce the single crystals by interfering with their cleavage. The prismatic layer of the mollusk shells of *Altrina serrata* is composed of single calcite crystals (~300x40x40 µm) enveloped by organic matrix. Each crystal contains intracrystalline acidic proteins (0.1%). The proteins do not appear however to be preferentially adsorbed on specific crystal planes, and do not reinforce the crystal against fracture. In previous studies (1) we measured the coherence length and angular spread of sea urchin spines, and compared them to synthetic crystals of calcite with or without occluded sea urchin proteins. As a continuation of these studies, we measured single calcite prisms from *Altrina* shells and synthetic crystals with and without occluded acidic proteins from *Altrina*. Two sets of profiles were collected for a number of reflections in the ω and $\omega/2\theta$ modes, with a Geranium (220) analyzer crystal. The coherence length values, derived from the FWHM of the $\omega/2\theta$ scans, are shown in Fig. 1a,b. Fig. 1c-e show the FWHM of the ω scans for pure calcite, synthetic calcite with occluded protein and for the biogenic elements respectively. The data are superimposed on those obtained in the previous studies for the sea urchin samples. The mosaic spread for the *Altrina* affected crystal is smaller than for the sea urchin affected crystal, while the coherence length is slightly less for that crystal. The biogenic crystals are significantly more perfect in *Altrina*, although the amount of occluded protein is larger. These results are consistent with non-specific adsorption and occlusion of the *Altrina* proteins inside calcite, in contrast with the controlled intercalation of proteins from the sea urchin. Preliminary data were also collected on crystals of calcium malonate, grown in the presence and absence of the same acidic proteins. Morphological evidence indicates that the proteins are preferentially adsorbed at the crystal's {101} planes during growth. $\omega/2\theta$ scans were collected for a limited number of hkl and $hkl0$ reflections, with the same set-up as above. A preferential broadening of hkl reflections was observed, seemingly in agreement with the anisotropy of protein adsorption. These are the first crystallographic data on the effect of anisotropic adsorption of specific crystal growth inhibitors on crystal texture.

Reference:
(1) Berman et al. Science 250, 664-667, (1990).

This research was supported by a grant from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel.

STUDY OF THE LOW TEMPERATURE STRUCTURE OF THE 2212 Tl-Ba-Ca-Cu-O SUPERCONDUCTOR

M. Blomberg (U. of Helsinki, Finland), I. Mutikainen (BNL, Chemistry), and J. Hanson (BNL, Chemistry)

Measurements for the determination of the structure of 2212 Tl-Ba-Ca-Cu-O were performed using the Huber six-circle diffractometer of the X7B beamline at the National Synchrotron Light Source. The single crystal samples were prepared in the Institute of Crystallography in Moscow, U.S.S.R., and a crystal ground to a sphere of radius 0.113 mm was selected for this study. At room temperature the structure is tetragonal (14/mmm), $a = 3.8545(3)$ Å and $c = 29.317(2)$ Å. Transition to the superconducting phase occurs at about 110 K.

The sample was cooled to 20 K with a closed-cycle two-stage helium cryostat (Displex), and a set of intensity data for about 800 reflections was collected using radiation of wavelength $\lambda = 0.9389$ Å. Because the cooling system was in operation the very first time, the experiment can be considered as a performance test of the instrument. Due to instrumental instabilities, especially in the refrigerator and sample mount, but probably also in the primary beam, there are some difficulties in the interpretation of the results, and the analysis of the data is at a preliminary stage.

To study the possible modulations of the structure, a search for satellite reflections or diffuse intensity occurring about the space-group forbidden reflections was performed by numerous scans in reciprocal space. The results indicate that the modulation, if any, is very weak in comparison to the modulations in the structurally related 2212 Bi-Sr-Ca-Cu-O compound. Furthermore, there is some evidence that the sample is an intergrowth of 2212 and 2201 phases.

DIFFRACTION LINEWIDTH OF CHARGE-DENSITY WAVE SATELLITE SPOTS

S. M. DeLand and G. Mozurkewich (U. Illinois), and L. D. Chapman (NSLS)

High resolution X-ray diffraction was used to measure the charge-density-wave phase-coherence length L in electron-irradiated blue bronze ($K_{0.3}MoO_3$) at 14 K. The anisotropy of L and its dependence on the pinning-site density n was determined by measuring the excess width of the superlattice peaks in three perpendicular directions for various values of n . In the nominally pure specimen, an excess linewidth was observed only along $2a^*-c^*$ direction, corresponding to a coherence length of 1400 Å. In the other two directions, a^*2c and b^* , the lines were resolution limited, corresponding to coherence lengths of at least 2100 Å. For large values of n , L was resolved in all three directions. At $n = 0.034$ at.%, the fully-resolved coherence volume was 3.7×10^8 Å³, containing approximately 2000 pinning sites. The resolved coherence lengths decreased with pin density roughly as $n^{-0.4}$.

Our principle results are (1) that a single coherence volume contains many pinning sites, even in the nominally pure sample, and (2) that L depends on n much more weakly than $1/n$. Taken together, these results are inconsistent both with the weak pinning model of Fukuyama, Lee, and Rice and with the conventional model of strong pinning. These results may be consistent with a modified theory of strong pinning presented by Abe and further analyzed by Tucker et al., but further theoretical analysis is required before a convincing comparison can be made.

Supported by NSF DMR84-51935 and DE-AC02-76CH00016

SINGLE CRYSTAL STUDY OF HYDRATED ALUMINOPHOSPHATE AT 20 K

T. Ericksson*, Å. Kvick*, Y. Long†, J. C. Hanson‡

*Experiments Division, ESRF, BP 220, F-38043 Grenoble Cedex, France

†Fudan University, Shanghai, China

‡Department of Chemistry, Brookhaven National Laboratory, Upton, NY 11973

Aluminophosphates form a vast group of materials with structure types ranging from dense phases like berillite, an analogue of quartz, through hydrates and semi-dense phases to microporous, zeolite like, materials. In contrast to the zeolites the AlPO_4 moiety is electrostatically neutral, hence there are no charge compensating cations present, and the procedure of varying the characteristics with ion-exchange does not exist within aluminophosphates. The networks formed are based on strictly alternating phosphate tetrahedrons and four, five or six oxygen coordinate aluminium. The five and six aluminium coordination give rise to a large complexity of network structures and indeed is one of the main factors behind this work.

The compound used in this study is the synthetic aluminium phosphate hydrate, $\text{AlPO}_4 \cdot 1.5 \text{H}_2\text{O}$. The structure has previously been solved from room temperature data by Pluth & Smith (ref. 1) and it has a three dimensional network, H3, based on up and down linkages between 6.6 and 4.8 Å two-dimensional nets. The aluminium has two positions, one in which it is linked to four network oxygens and a second where it has two additional water oxygens. There is a third water molecule in the structure, which apparently is not within bonding distance to any other atom. This water molecule resides in an 8-ring channel. The aim of this low temperature measurement was to study the bonding between the third water molecule and the framework structure and also examine the electron densities in the channel structure.

A rod shaped crystal of dimensions $30 \times 30 \times 80 \mu\text{m}$, grown by hydrothermal methods, was used for the data collection at the NSLS X-7B crystallography station using a $\lambda = 0.9235$. The crystal was cooled in a quartz capillary to prevent dehydration in the refrigerator at 20°K . The crystal was enclosed in a DISPLEX-202 closed cycle refrigerator down to 20°K . The space group at 20°K was the orthorhombic $Pbca$ with the unit cell; $a = 9.7450(8) \text{ Å}$, $b = 9.7564(5) \text{ Å}$ and $c = 19.382(2) \text{ Å}$. A total of 2500 intensities out to $\sin(\theta)/\lambda = 0.7 \text{ Å}^{-1}$ were collected, out of these 1500 intensities were unique. The collection was made using ω -scan with a step of $\Delta\omega = 0.006^\circ$ and a scan width of 0.8° . Absorption correction was performed by the method of Gaussian grid with $\mu = 15.3 \text{ cm}^{-1}$.

The structure was refined using the coordinates from ref. 1 as a starting set down to $R = 0.07$. Water oxygen and hydrogen atoms were located in difference Fourier maps. Two of the water molecules are bonded to one of the aluminium atoms with Al-O distances of 1.975(4) and 1.965(5) Å. The third water molecule is hydrogen bonded to one of the water oxygens, and to a framework oxygen in a disordered way with O-O distances of 2.924(6) and 3.015(6) Å. There are four oxygen atoms within hydrogen bonding distances to the water molecule. The large and elongated thermal ellipsoid on the water oxygen atom suggests several alternative water orientations. The P-O distances range from 1.509(4) to 1.543(4) Å, and the Al-O distances for the four fold and six fold aluminium range from 1.724(4) to 1.750(4) Å and from 1.836(4) to 1.975(4) Å respectively.

References: 1. J. J. Pluth and J. V. Smith, Acta Cryst. (1986), C42, 1118-1120.

This research was carried out at Brookhaven National Laboratory under contract DE-AC02-76CH00016 with the U.S. Department of Energy Office of Basic Energy Sciences.

EXAFS Investigation of Amorphous Platinum Uridine Clusters in H_2O -solution

V. Eteläniemi, R. Serimaa, T. Laitalainen*, T. Paakkari and I. Mutikainen*

University of Helsinki, Department of Physics, Helsinki, Finland

* University of Helsinki, Department of Chemistry, Helsinki, Finland

We have carried out an EXAFS investigation of water solutions of platinum uridine complex to determined the local structure of Pt atoms. The purpose of these investigations was to compare the local structure of Pt atoms in solid state (see X23A2 in bibb) with the structure in H_2O -solution. Fourier transform of the EXAFS signals of liquid and solid samples (k -range $3\text{--}11 \text{ Å}^{-1}$) are shown in figure. Small differences are visible but these may be also due to distortions in the data from the liquid samples. Further analysis is in progress to clarify the differences.

A purpose of the work was to study the beamline for EXAFS investigations. A double crystal monochromator, the Si(311) reflection, and a focusing mirror were used. Some of the samples were simultaneously investigated at the beamline X23A2. Comparison of the data shows similar signals up to point $k = 10 \text{ Å}^{-1}$ but unfortunately at larger k -values the statistical precision of the X23A2 data was too low. Also in the X7B data distortions were visible at larger k range due to fluctuations of the incident flux (order of $\pm 1\%$). Further work is needed to stabilize the incident flux.

Research supported by Neste Foundation, by the Technology Development Centre and by the Academy of Finland, Finland.

EXAFS STUDIES OF COPPER-CONTAINING MONOOXYGENASES

Ninian J. Blackburn and Richard S. Strange (Oregon Graduate Institute).

The structure of the reduced form of the Cu containing monooxygenase dopamine- β -hydroxylase (DBH) has been the subject of controversy concerning the presence or absence of a heavy atom scatterer coordinated to the Cu(I) center. As part of our continuing studies on DBH and related Cu-containing enzymes we have been investigating the EXAFS spectroscopy of various states of the enzyme, and have used this and previous beam time allocations to settle the question of the identity of the ligands to Cu(I). We have now measured 5 different preparations of the reduced enzyme as follows (i) and (ii) ascorbate reduced enzyme from different batches (same method of preparation) (iii) reduced enzyme prepared under rigorous exclusion of Cl^- ion and (iv) sample (iii) plus 2 mM Br^- (v) native enzyme fully oxidized with $\text{Fe}(\text{CN})_6^{3-}$ then reduced with ascorbate. All of these samples have given essentially identical spectra which can be simulated by 2-3 His ligands at 1.94 Å and a second row scatterer (S or Cl) at 2.24 Å. The insensitivity of the spectra to rigorous exclusion of chloride ion and to the presence of millimolar concentrations of bromide would argue strongly that the heavy atom scatterer is S not Cl. In addition we have measured the Cl concentration in the ultrafiltrate of fully oxidized native enzyme which is known not to exhibit any second row scatterer, and found it to be less than 200 μM . The EXAFS of the reduced form of this fully oxidized native sample was identical to all the other spectra. Thus we believe that we have established beyond reasonable doubt the presence of a S ligand to Cu(I) in reduced DBH.

We have also continued our EXAFS studies on the CO derivatives of DBH and related system such as CO-hemocyanin and inorganic Cu(I)-carbonyls. A combination of EXAFS, FTIR and chemical analysis of bound CO has provided evidence that DBH binds CO to form a simple carbonyl adduct at one of the metal centers, with little perturbation of the reduced enzyme structure. Since CO is an analogue of the dioxygen substrate, this indicates that the enzyme is designed to form a simple dioxygen adduct at one of the copper centers.

We have initiated EXAFS studies on a second Cu-dependent monooxygenase, phenylalanine hydroxylase, in collaboration with Professor Steve Benkovic, Pennsylvania State University. So far we have only succeeded in obtaining data on the oxidized Cu(II) enzyme. Analysis using restrained refinement has strongly suggested the presence of tyrosine as a ligand. Attempts to measure the reduced enzyme have so far proved unsuccessful.

We have also continued our long and fruitful collaboration with Professor K. D. Karlin on EXAFS of Cu(I)-dioxygen inorganic model complexes. A new complex, $\text{Cu}(\text{O})\text{-tris}(2,3\text{-dimethylimidazole})\text{O}_2$ has been measured. This work complements previous work on X9A on Cu(II)-peroxo complexes prepared by Dr. Nobumasa Kitajima and oxy hemocyanin.

Supported by Grant R01-NS 27583 from National Institute of Health.

X-RAY DIFFRACTION STUDIES OF TETHERED PROTEIN MONOLAYERS AND ULTRATHIN LIPID MULTILAYER FILMS ON Ge/Si MULTILAYER SUBSTRATES

J.K. Blasie (U. Penn), J. Chupa (U. Penn), S. Xu (U. Penn), M. Murphy (U. Penn) and R. Fischetti (Univ. City Sci. Center, Phila.)

Meridional x-ray diffraction $\Pi(Q_{xy} = 0 \text{ Å}^{-1}, Q_z)$ data were collected from a cytochrome c protein monolayer covalently tethered to the surface of a Ge/Si multilayer substrate, a self-assembled monolayer of octadecylsilane on a Ge/Si multilayer substrate, and both a Langmuir-Blodgett monolayer and head-to-head bilayer of cadmium arachidate on bare vs. alkylated Ge/Si multilayer substrates, respectively. Doubly focusing x-ray optics (cylindrically bent Ni-mirror for vertical focus along Q_z and a cylindrical-ly bent ribbed Si-crystal for horizontal focus in the layer (xy) plane), specimen oscillation and a linear position-sensitive detector were employed for data collection. The Ge/Si multilayer substrates were fabricated by molecular beam epitaxy-MBE (J. Bean, AT&T Bell Labs.) to provide strong, but continuous x-ray diffraction to higher Q_z than can be achieved with inorganic multilayers fabricated via magnetron sputtering. X-ray interferometry utilizing the Ge/Si finite multilayer as the "known structure" is being used to uniquely derive the electron-density profile structures of these organic/biorganic overlayers.

Reference: Blasie, J.K., Xu, S., Murphy, M., Chupa, J., McCauley, J.P., Smith, A.B., III, Bean, J.C., and Petricolas, L.J., 1991. "Profile Structures of Macromolecular Monolayers on Solid Substrates by X-ray Holography", *Mat. Res. Soc. Symp. Proc.* (in press).

TIME-RESOLVED, RESONANCE X-RAY DIFFRACTION STUDIES OF SARCOPLASMIC RETICULUM MEMBRANE MULTILAYERS

J.K. Blasie, F. Asturias, L. DeLong (U. Penn), and R. Fischetti (Biostructures Institute)

The resonance x-ray scattering effects from lanthanum ions, bound to the two high-affinity metal-binding sites of the Ca^{2+} -ATPase, on the meridional x-ray diffraction $I(Q_{xy} = 0 \text{ Å}^{-1}, Q_z)$ from oriented multilayers of isolated sarcoplasmic reticulum membranes were investigated for the $(\text{La})_2\text{E}_1$ and $(\text{La})_2\text{E}_1\text{-P}$ forms of the enzyme. The meridional diffraction was collected in serial time-frames at three energies about the La L_{III} absorption edge (i.e., edge $\pm 100 \text{ eV}$) for the $(\text{La})_2\text{E}_1$ form of the Ca^{2+} -ATPase, and then for the $(\text{La})_2\text{E}_1\text{-P}$ form following the flash-photolysis of caged -ATP substrate in the oriented multilayer, using a linear position-sensitive detector. These data are being analyzed to provide the profile structure $\Delta\rho(z)$ for the Ca^{2+} -ATPase for the $(\text{La})_2\text{E}_1$ vs. the $(\text{La})_2\text{E}_1\text{-P}$ forms of the enzyme, via the time-resolved data as constant x-ray energy and for the position(s) of the lanthanum ions in each of these two profiles via the resonance data prior to, vs. following, the flash-photolysis of caged-ATP. The conversion of the $(\text{La})_2\text{E}_1$ to the $(\text{La})_2\text{E}_1\text{-P}$ form of the Ca^{2+} -ATPase results in the occlusion of these two metal binding sites involved in the active (energy-dependent) transport of calcium across the sarcoplasmic reticulum membrane.

Supported by the NIH grants HL18708 and RR01633.

THE STRUCTURE OF THE TRANSIENT STATE INTERMEDIATE OF THE H⁺-ATPASE FROM CHLOROPLASTS.

C. Carmeli[#], O. Tadmor[#], R. Ophir[#] and S. Carmeli*, Departments of Biochemistry[#] and Chemistry*, Tel Aviv U., Tel Aviv, Israel.

The H⁺-ATPase in the chloroplast membranes utilizes the electrochemical potential of protons for the synthesis of ATP from ADP and Pi. The process requires cooperative interaction among the three β subunits in which the catalytic sites of this multisubunit enzyme are located. We have shown earlier a positive cooperative interaction among three Mn²⁺ binding sites on the enzyme which affects the catalytic activity. There are several indications that these cooperatively interacting metals are bound at the active sites of the enzyme. Among these are EXAFS determination of the Mn²⁺ environment which indicated the appearance of three phosphorous atoms near each of the metals on addition of ATP or ATP analogues to the enzyme. Further support comes from the fact that activation of the enzyme shortened the Mn-P distance from 4.85 Å to 3.35 Å. Such an effect might be required to reduce the energy of activation by inducing a more positive phosphorous which becomes more susceptible to nucleophilic attack by water. Recently we have shown that such an attack is likely to proceed through a formation of a pentacoordinate phosphorous intermediate since vanadate which readily forms a pentacoordinate analogue to phosphate inhibited enzyme activity. Binding of the analogues to the enzyme was determined with the aid of nuclear magnetic resonance techniques. The nature of the transient state intermediate formed during the catalytic process was studied by determination of the environment bound Mn²⁺ in the presence of vanadate and of bound vanadate in the presence of Mn²⁺. An appearance of vanadium was observed on addition of vanadate to an enzyme prebound with Mn-ADP at the three interacting sites. An apparent Mn-V distance of approximately 3.3 Å was observed. This distance indicated that the vanadate is placed at the same distance from Mn²⁺ as the two phosphates of ADP when bound to the Mn²⁺ on the enzyme. Such a structure could be obtained on a formation of an oxygen bridge between the β phosphate of ADP and the vanadate both of which contribute oxyanions which were ligated to the Mn²⁺.

Based on these results it is suggested that a pentacoordinate phosphorous intermediate which is formed at the γ phosphate of ATP serves as the transient state in the process of hydrolysis and possibly in the synthesis of ATP. The formation of the intermediate might be facilitated by an exchangeable water molecule in the Mn²⁺ hydration shell committed for the displacement process. The close proximity of the phosphorous to the positively charged metal seem to be an essential element in affecting the nucleophilic attack of the water molecule during catalysis.

X9-A

XAFS studies of the iron containing subunit of Ribonucleotide Reductase

Grant Bunker
Physics Dept, Illinois Institute of Technology
Ke Zhang, B. Chance
Biostructures Institute, University City Science Center
Margareta Sahlin & Britt-Marie Sjöberg
Department of Molecular Biology, Stockholm University

XAFS spectra were collected on a recombinant mutant form (very similar to the wild type) of the iron containing subunit of Ribonucleotide Reductase from *E. coli*. The oxidized ferric form was compared to a form savagely reduced with sodium dithionite because more civilized reduction procedures failed. The situation should be greatly improved when the PRT's anaerobic glove box becomes operational. Two individual scans are shown below. Analysis and comparison with previously obtained data is underway.

This work is supported by NIH GM39612 and RR01633

X9A
IN SITU XANES STUDY OF THE CHEMISTRY OF PASSIVE OXIDES ON AlV AND AlMn THIN FILMS UNDER ELECTROCHEMICAL CONTROL

A. J. Davenport (BNL), H. S. Isaacs (BNL), G. S. Frankel (IBM), A. G. Schrott (IBM), C. V. Jahnes (IBM), and M. A. Russak (IBM)

The chemistry of passive oxide films on sputtered AlV and AlMn alloys in aqueous environments has been compared with that of AlCr alloys¹. XANES measurements were made on films in an electrochemical cell under potential control. The alloy was present as a thin film ~40Å in order to minimize signal from the underlying metal. Measurements were made in fluorescence geometry using a 13-channel solid state detector (Canberra). It had previously been determined that slow increase in the potential caused Cr to dissolve from AlCr alloys through formation of the 6-valent state on the surface. In contrast, polarization of AlV alloys results in formation of the high valent vanadate (Fig.1, presence of characteristic pre-edge peak) without any dissolution. For AlMn alloys, the highest valence state is +4, which then dissolves without formation of 7-valent permanganate (Fig.2).

Fig.1 V K edge for Al-21%V polarized in a borate buffer solution, potentials quoted with respect to a mercurous sulfate electrode.

1 A. J. Davenport, H. S. Isaacs, G. S. Frankel, A. G. Schrott, C. V. Jahnes, and M. A. Russak, *J. Electrochem. Soc.*, **138** 337 (1991).

This work was performed in part under the auspices of the U.S. Department of Energy, Division of Materials Sciences, Office of Basic Energy Science under Contract No. DE-AC02-76CH00016.

X9A

PURIFICATION, EXAFS, AND EDGE DATA OF VITAMIN-B12 BINDING PROTEINS WITH CYANO- AND AQUO-COBALAMIN
S.M. Frisbie, M.R. Chance, Chem. Dept., Georgetown U., Washington, D.C., 20057

Cobalophyllin and a mixture of TCII-IF have been purified from human plasma and bound to either cyanocobalamin or aquocobalamin. EXAFS and edge data have been obtained on the protein/cobalamin at NSLS. The edge data show a 0.5 eV shift to lower energy for cyanocobalamin bound to either protein fraction. The aquocobalamin/cobalophyllin does not show an edge shift. The shift to lower energy provides strong evidence that the σ -donor power of the cobalamin axial ligands has changed on binding to protein. This is due to either exchange of the 5,6-dimethylbenzimidazole (DMS) ligand with a protein ligand, or changes in the cobalt-ligand distances without exchange. Changes in the cobalt-ligand distances were seen for both protein fractions. The cobalt-DMS distance for aquocobalamin changes from 2.06 to 2.28 Å when cobalophyllin is bound. See Table I and Table II below.

BOND	CNCEL	AQCEL	CN/CP	AQ/CP	CN/TCII-IF
Co-Neq	1.89	1.90	1.89	1.89	1.88
Co-DMS	2.15	2.06	2.13	2.28	2.12
Co-Åx	1.93	1.93	1.91	1.95	1.95

Table I. EXAFS data of cyano- (CNCEL) and aquocobalamin (AQCEL) for cobalophyllin (CP) and TCII-IF (transcobalamin II and intrinsic factor). Co-Neq = ave. distance of 4 corrin ring nitrogens to cobalt, Co-DMS = dimethylbenzimidazole, and Co-Åx = upper axial ligand.

COMPOUND	CHARGE	POSITION	COMMENTS
Aquocobalamin	+3	0.0	standard
CN-cobalamin	+3	0.0	standard
Aquo/CP	+3	0.0	no edge shift
CN/CP	+3	-0.5	large shift
CN/TC-IF	+3	-0.5	large shift

Table II. Edge data of cobalamins and protein complexes.

This research is supported by the CSRS/USDA Dept. of Agriculture Grant # 90-37200-5357 of the Program in Human Nutrition. Beam line X9A is supported by NSF-RSC1633 and NSF-DMS910959

EXAFS STUDIES OF Ni TETRAAZAMACROCYCLES AND THEIR ADDUCTS AS MODELS OF F430

L. R. Furenlid (NSLS, BNL), M. W. Renner (DAS, BNL), and E. Fujita (Chem. BNL)

F430 is a nickel(II) hydrocorphin and the prosthetic group of methyl coenzyme M reductase. It catalyzes the reductive cleavage of S-methyl coenzyme M to coenzyme M and methane in the final step of the reduction of carbon dioxide to methane. As models of F430, we studied the EXAFS of Ni(II) complexes, the one electron reduced species and their adducts (L = 2,3,9,10 - tetramethyl - 1,4,8,11 - tetraazacyclotetradeca - 1,3,8,10-tetraene) in CH_3CN and other solvents to characterize the structural differences as a function of oxidation state and axial ligation of the metal in solution at 300 K and 173 K.

The near-edge structure of Ni(II)L_2^{2+} , shown in Figure 1a was solvent dependent: six coordinate in CH_3CN (solid line), four coordinate in CH_3NO_2 (broken line). The reduced species, Ni(II)L^+ , is clearly four coordinate in CH_3CN , Figure 1b. The edge of NiLCO^+ (Figure 1c) in $\text{CH}_3\text{CN}/\text{C}_3\text{H}_7\text{CN}$ ($v/v = 1/5$) at 300°K (solid line) is complicated, consistent with the existence of an equilibrium mixture of monomer Ni(II)L^+ , its dimer and Ni(II)LCO^+ . At 173 °K (broken line), the pre-edge clearly has the structure associated with five coordinate NiLCO^+ . This observation is supported by the optical spectra of NiLCO^+ in $\text{C}_3\text{H}_7\text{CN}$ under 720 torr of CO at various temperatures.

Fourier transforms of k^3 weighted EXAFS of the Ni(II) parent complex in CH_3CN (a) and in CH_3NO_2 (b), the one-electron reduction product (c) and the CO adduct at 173 K (d) are displayed in Figures 2. The EXAFS results reveal that the Ni(II) to Ni(II) anion radical reduction leaves the geometry around the nickel atom unchanged. The π anion radical, Ni(II)L^+ , undergoes a monomer-dimer equilibrium ($K_1 = 5.5 \times 10^4 \text{ M}^{-1}$) in CH_3CN as evident from the optical spectrum. However the x-ray absorption edge features show no signs of an axial interaction. Furthermore, the EXAFS showed no obvious evidence for a Ni-Ni interaction, suggesting that the dimer has a long Ni-Ni distances ($> 2.5 \text{ \AA}$), and/or is characterized by a large Debye-Waller factor at room temperature. EXAFS data on CO adducts of Ni(II)L , clearly indicate that these are five-coordinate complexes with a short Ni-C bond. Ni-N distance in the CO adducts of the Ni(I) complexes increase quite dramatically compared to that in the parent Ni(II) complexes.

This research was carried out at Brookhaven National Laboratory under contract DE-AC02-76CH00016 with the U.S. Department of Energy and supported by its Division of Chemical Sciences, Office of Basic Energy Sciences.

IN SITU CHARACTERIZATION OF A COPPER MONOLAYER ELECTROCHEMICALLY ADSORBED ON A PLATINUM ELECTRODE

T. E. Furtak, Physics, Colo. Sch. Mines, Golden, CO 80401

T. M. Hayes, Physics, Rensselaer Poly. Inst., Troy, NY 12181

We have continued our program under which a thin-layer electrochemical sample chamber is used to acquire grazing incidence fluorescence detection EXAFS from an adsorbed monolayer of Cu on a large (55cm^2) polycrystalline Pt electrode, while the sample is under electrochemical control. The deposition is performed with the Kaptan window expanded with excess electrolyte (0.5 M sulfuric acid with 0.2 mM Cu^{++}) above the sample. The sample voltage is controlled with respect to a Cu-wire reference to define the "underpotential" condition where only the first monolayer is deposited (0.1 eV on this scale). By using stripping voltammetry as well as x-ray edge calibration we have verified that the coverage corresponds to one Cu atom per Pt surface atom. Under the conditions of the experiment the water layer is approximately 10 microns thick. We average 15 scans at the Cu K-edge with the signal being detected by a large aperture ion chamber equipped with Soller slits and Ni filters. The Fourier transformed data appear below.

R (ANGSTROMS)

We have improved our sample preparation to avoid Pt restructuring which has been observed with this system. The result is a much cleaner signal. Analysis of these data seem to indicate that the near neighbor is Cu, not oxygen as was originally thought. If this is true, then the distance is 1.6 Å, much closer than would be predicted. Apparently the Pt induces Cu dimerization, or a stable condition under which Cu-Cu bonds are very short.

Sponsor: ONR N0014-90-J-1326 and N0014-90-J-1332.

THE STRUCTURE OF THE ZINC SITES OF E. coli DNA-DEPENDENT RNA POLYMERASE

W.-J. Huang¹, F.Y.-H. Wu^{1,2}, R.B. Sinclair³, and L.Powers³

¹ Department of Pharmacological Sciences, State University of New York at Stony Brook, Stony Brook, New York 11794

² Institute of Biomedical Sciences, Academia Sinica, Taipei 11529, Taiwan, R.O.C.

³ National Center for the Design of Molecular Function, Department of Chemistry and Biochemistry, Utah State University, Logan, Utah 84322-4630

X-ray absorption spectroscopy is ideally suited for the investigation of the electronic structure and the local environment ($\leq 5\text{\AA}$) of specific atoms in biomolecules. While the edge region provides information about the valence state of the absorbing atom, the chemical identity of neighboring atoms, and the coordination geometry, the EXAFS region contains information about the number and average distance of neighboring atoms and their relative disorder. The development of sensitive detection methods has allowed studies using near physiological concentrations [as low as $\sim 100\mu\text{M}$]. RNA polymerase from *E. coli* contains two zinc atoms: one tightly bound in the β' subunit, the subunit which participates in template binding, and the other loosely bound in the β subunit, the subunit which participates in substrate binding. X-ray absorption studies of these zinc sites in the native protein and of the zinc site in the β' subunit after removal of the zinc in the β subunit site by p -(hydroxymethyl)benzenesulfonate [D.P. Giedroc and J.E. Coleman, Biochem. 25, 4969-4978 (1986)] indicate that both Zn sites have octahedral coordination. The zinc in the β' subunit zinc site has four sulfur ligands at an average distance of $2.36\pm 0.015\text{\AA}$ and two oxygen (or nitrogen) ligands at an average distance of $2.23\pm 0.02\text{\AA}$. The β subunit zinc site has five sulfur ligands at an average distance of $2.38\pm 0.15\text{\AA}$ and one histidine nitrogen ligand at $2.14\pm 0.15\text{\AA}$. These results are in general agreement with earlier biochemical and spectroscopic studies.

This work was supported by State of Connecticut Department of Higher Education, High Technology Grants and was completed at beamline X9A which is supported by NIH/RR01633.

X9-A

DRUG STRUCTURES IN BIOLOGICAL MEMBRANES

Leo G. Herbette and Lina A. Nakhimovsky (Dept. of Radiology, Biomolecular Structure Analysis Center, University of Connecticut Health Center)

These were the first preliminary experiments to obtain information on the structure of drug molecules partitioned in membrane bilayers. Dioleoyl phosphatidylcholine (DOPC) dispersions in excess water and partially dehydrated oriented multilayers of DOPC were prepared in a 0.8 mm (OD) glass capillary and placed in a 2 Tesla magnetic field during x-ray exposure on Beamline X9A. In these experiments, the width of the lamellar reflections was being monitored to indicate an effect of the magnetic field on the order/disorder of the lipid lamellae. Analysis of data is currently in progress. Preliminary data do not indicate a strong effect of the magnetic field on the width of the lamellar reflections. Specially designed lipids with magnetic particles were recently synthesized and will be used in the next set of experiments. The ultimate goal is to obtain sufficiently high resolution of drug-lipid complexes in fully hydrated membrane bilayer.

Dioleoyl Liquid Crystal Strands

S. H. J. Mziak, N. C. Maliszewski, G. B. M. Vaughan, P. A. Henney, J. P. McCauley, Jr., and A. B. Smith, III, U. Pennsylvania; C. Moraschoff and H. Ringsdorf, U. Mainz, Germany.

The liquid-crystalline structure of hexacyclic derivatives has been extensively studied, and the structure was previously suggested to be hexagonal-tubular. However higher-order characteristic hexagonal peaks have never been observed in any hexacyclic derivative, even in high resolution powder XRD experiments. Moreover, computer simulations revealed that the cores of the molecules tend to collapse and that they prefer to take on a rod-like conformation when in the gas phase. These results suggested a lamellar smectic phase, although a hexagonal columnar structure could not be completely ruled out [Mziak et al. to be published in JACS]. We have now performed high resolution powder XRD on two new compounds: 1b and 1c (shown below).

To ensure consistency, we repeated the measurements previously [Mziak et al.] made on 1a with this experimental setup. For 1a only the (100) diffraction peak at $0.1883\text{\AA}^{-1}$ can be seen. Both the (200) and the (110) are reduced in intensity by a factor of at least 5000. The existence of the latter (i.e., a $\sqrt{3}:1$ ratio of the peak positions) is crucial for the identification of hexagonal order. By contrast, the diffraction pattern for 1b shows a clear (100) peak at $0.190\text{\AA}^{-1}$, and also a weaker (110) peak at $0.328\text{\AA}^{-1}$ is visible, demonstrating the existence of hexagonal order. Even clearer results can be seen for 1c: strong (100), (110) and (200) peaks, at $0.178\text{\AA}^{-1}$, $0.3085\text{\AA}^{-1}$, and $0.3575\text{\AA}^{-1}$ respectively, are present.

This is the first conclusive experimental evidence of hexagonal columnar order in any hexacyclic mesophase. It is still an open question whether 1a and other hexacyclic derivatives necessarily are hexagonal, or whether the structural order depends on details of the molecular architecture.

Supported by the National Science Foundation under Grants DMR 89-01219 and DMR-8819885.

X-RAY ABSORPTION SPECTROSCOPY OF MANGANESE IN CHLOROPLASTS. X9A

Melvin P. Klein, Vital K. Yachandra, Victoria J. DeRose, Matthew J. Latimer, Ishita Mukerji and Kenneth Sauer.

Melvin Calvin Laboratory, Chemical Biodynamics Division, Lawrence Berkeley Laboratory, Berkeley, CA 94720.

Our Mn X-ray absorption spectroscopy experiments have provided a structural model for the Mn complex in Photosystem II of the oxygen evolving complex in plants and cyanobacteria. The complex is made up of two di- μ -oxo bridged Mn clusters where the Mn-Mn distance is about 2.7 Å, which are linked to each other by at least one μ -oxo bridge with a Mn-Mn separation of about 3.3 Å. It is also known that chloride (or bromide) and Ca (or Sr) are required cofactors for maintaining activity. We have been interested in determining if the halide and Ca are in the proximity of the Mn complex. We have pursued Mn EXAFS studies of samples where chloride has been replaced by bromide and Ca by Sr. Since Br and Sr have greater X-ray backscattering amplitudes we expected to be able to deconvolute the backscattering more easily. We have grown cyanobacteria in medium containing only bromide and Mn X-ray results show that one halide could be a ligand to Mn. We have also shown that the Mn EXAFS spectra change dramatically when Ca is replaced by Sr. However these experiments are hampered by the presence of maybe only one halide and one Ca (or Sr) per four Mn atoms. So the backscattering from the other ligand atoms to four Mn atoms is convolved with backscattering from one Br or Sr making the deconvolution difficult. We have recently managed to prepare samples containing only one Br or one Sr per four Mn atoms, which makes the 'reverse' experiment of looking for Mn backscattering from Sr or Br feasible, and we have collected preliminary data with these samples. The data are being analyzed.

This research was supported, in part, by the Director, Office of Energy Research, Office of Basic Energy Sciences, Division of Energy Biosciences of the U.S. Department of Energy, under contract DE-AC03-76SF00998, and by a grant from the National Science Foundation (DMB83-04526).

EXAFS Experiments on Iron, Copper, and Zinc Proteins

Richard Kucszyn

Three EXAFS experiments were successfully performed during the last reporting period. These include the determination of the stereochemistry of

- 1) a Zn site in a mammalian transcription factor protein;
- 2) the Cu site in a nitrite reductase;
- 3) the Fe and Ni sites in a carbon monoxide dehydrogenase (CODH).

In the first experiment, the Zn EXAFS spectrum of a mammalian transcription factor was measured and indicated that the metal site was co-ordinated by 4 S atoms at an average distance of 2.33 Å from the Zn atom. The EXAFS results suggested that the protein was indeed a member of the class of so-called Zn-finger proteins which regulate the expression of DNA activity. Subsequent to the EXAFS study, several chemical and biological studies have confirmed the EXAFS result and firmly established that this factor is indeed that this factor is indeed a member of the Zn-finger class of proteins.

In the second experiment, the EXAFS spectra of *Achromobacter* nitrite reductase were determined in its oxidized resting state and with nitrite and azide bound. In addition, the EXAFS spectrum of its resting reduced state was determined. This enzyme contains one Type I and one Type II copper atom per monomer and x-ray crystallography has shown that the Cu atoms are approximately 12 Å apart and share a common histidine ligand. The EXAFS spectrum of the resting oxidized enzyme indicated that the Type I Cu atom was coordinated by two N atoms at 1.98 Å, one S atom at 2.15 Å and one S atom at 2.60 Å. The Type II Cu atom was shown to be coordinated by three N atoms at 1.98 Å and a water molecule at 2.25 Å. The ligands to each Cu atom were unchanged in the reduced molecule except that the average bond lengths increased by approximately 0.15 Å. These observations are consistent with the crystallographic observations. The EXAFS spectra of the oxidized enzyme in the presence of either nitrite or azide were identical to within experimental error and indicated that the Type I Cu environment was similar to that observed in the resting enzyme but the Type II Cu could best be fit by one average Cu-N distance of 1.97 Å. This has been interpreted in terms of a tetrahedral Type II Cu environment where the azide is bound by a terminal N atom and nitrite is bound by its central N atom.

In the third experiment, the Fe EXAFS spectrum of oxidized CODH was similar to the spectra obtained from typical Fe-S cluster proteins while the spectrum of the reduced protein showed significant differences compared to other Fe-S proteins. The Ni EXAFS indicated that the Ni atom was coordinated by 4 S atoms at 2.20 Å. Neither the Ni nor the Fe EXAFS showed evidence for the other metal but the reduced Fe XANES showed profound changes when compared to other Fe-S proteins suggesting that the Ni atom is intimately associated with the Fe-S clusters in CODH. Model building suggests that the Ni may share coordination with the clusters through the cysteine ligand.

Protein Damage and Possible Protection: EXAFS Studies of Metallo Enzymes.
 † C. Kumar, * K. S. Reddy † Chang Lee, and * B. Chance; † Bio-Structures Institute, 3401 Market Street; and * Dept. Of Biophys. & Biochem.; U. of Penn.; Philadelphia PA19104

X-ray Absorption Spectroscopy with Synchrotron Radiation is a powerful tool and only the choice for determining the distances of active sites in metallo-enzymes. Insertion devices and focussing mirrors in near future will produce very intense fluxes of x-rays (Advanced Photon Source) and obviously are the key to the opening of considerable extension to present EXAFS applications. For example in near future, it will be possible, to determine the structures of intermediates of metallo-enzymes in solution state. In the present study we show that protein damage is a potential problem for such EXAFS studies and one has to characterize them during and after EXAFS data collection

We have studied the radiation induced changes on number of proteins and enzymes starting with myoglobin, cytochrome-C, HRP, bovine serum albumin and superoxide dismutase. These experiments were performed on beam line X-9A (focussed beam) which is normally used for EXFAS studies. We have used U.V-Visible, Fluorescence & EPR spectral bands and by on-line optical spectroscopic studies to monitor the changes in the active site. The ligand binding studies were used for assaying the activity of the proteins. It has been observed that all the heme proteins studied were partially reduced and the intensity of the solet band decreases during irradiation indicating the heme is partially destroyed during the X-ray exposure.

We have also investigated various protectants (apart from Cr(NTA) complex) specially N-Oxide radical scavengers for protein protection. EPR technique was used to quantify the radical production due to radiation. At present it has been concluded that the N-Oxides were not effective protectors. The detailed study of the mechanism for the protective action of various Cr complexes is in progress.

The results indicate that the proteins studied in this report do lose their activity (30% in 1/2hr exposure) and we conclude that it is important to know the state of the sample under investigation and check for the X-ray induced damage by on-line monitoring of the sample during the EXAFS data collection.

This work was supported by RR01633

X-RAY INDUCED DAMAGE TO PROTEINS DURING EXAFS STUDIES X9A
 C. Kumar, B. Chance and K. S. Reddy, Dept. of Biochem. & Biophys., University of Pennsylvania; Philadelphia PA 19104.

EXAFS spectroscopy with synchrotron radiation has become a powerful tool for determining the active site structures of metalloenzymes in solution. Radiation damage to the protein and changes in the oxidation state of the metal can occur during the EXAFS measurements, compromising the results. Insertion devices and focussing mirrors producing very intense fluxes of X-rays are the key to the extending EXAFS applications but, are also likely to increase the radiation damage to biochemical samples. In the present study we show that the heme proteins like myoglobin, cytochrome-C, peroxidase are reduced and damaged at room temperature on exposure to the X-ray flux obtained in the X-9A beam line operating in the focussed mode.

Optical spectral alteration of myoglobin during 90 min. exposure in X-9A beam line. X-ray beam was horizontally focussed to 4mmX 2mm. The initial decrease in the 635 peak is an indication of protein reduction.

Reduction of CytoC in X-9A beam line (Focussed beam) at room temperature. The protein was reduced completely within 12 mts. This was followed by further spectral changes over 60 minutes.

These studies indicate that proteins at room temperature are damaged by X-Ray exposure. So, during EXAFS investigations, we consider it important, to test the activity of the proteins before and after X-ray irradiation, collect EXAFS and transmission optical spectroscopy data together to monitor potential sample damage. Combination of these methods will make biological EXAFS investigations more reliable

This work was supported by NIH grant RR 01633 and a BRSG from UCSC.

The Vertical X-ray Beam Stability at Beam Lines X-9A & B at NSLS X-9
H.-C. Lee, S. Khalid, M.-Z. Zhang, and B. Chance; Bio-Structure Institute, 3401
Market Street, Philadelphia, PA 19104

Upon the analysis of the vertical X-ray beam stability at NSLS of Brookhaven National Laboratory, both performance of the vibrating reed instrumentation and X-ray beam stability were tested. The phosphor coated reed chopped the X-ray beam at 200 Hz and the emitted photons were picked up by a PMT. The PMT had a high voltage closed loop feedback system to maintain a constant signal with the decaying beam. The phase shift in the pulses caused by the beam movement produced a proportional voltage variation and was recorded. Two reed detectors were used inside the X-9A hutch.

The beam movements were observed in three different situations by mounting the reed 6 m from the monochromator and vertically focusing the beam to 0.4, 2.0, and 4.5 mm FWHM. A Ni coated mirror was used to focus the beam. The Si(111) monochromator was set at 7.011 KeV. In each case the monochromator was tuned by using the piezo voltage change from a lower to a higher voltage, changing the incident flux from 10 % of the peak to the peak and back to 10 % of the peak at the higher voltage end. There was significant vertical beam movement in all three conditions. It was expected from the control of the tilting of the second monochromator crystal. By comparison, the maximum to minimum beam movements were observed in the order of fully unfocused, fully focused and half focused beam (Fig. 1).

Long time survey of the vertical X-ray beam stability suggested a random bidirectional drift of X-ray beam. Average approximate ± 560 microns vertical displacement was observed prior to the installation of global feedback system, which reduced the beam drift to ± 23 microns during a fill and greatly reduced the occurrence of sudden beam movements. In the past beam jumps of 20 to 300 microns were observed.

Fig. 1 ΔVT (Units 1&2) vs. ΔP° with beam half-focused

Fig. 2 X-Ray Beam Stability: X-9A vs. X-9B

Two detectors were also setup at monochromatic beam line X-9A and white beam line X-9B. A simultaneous record of the beam movement exhibits some common features and, in a way, testifies the integrity of the performance of the two detectors. (Fig. 2)

Work Sponsored by the NIH grant RR-01633.

X9A
GLANCING-ANGLE XAFS STUDIES OF PRE-MELTING AT Cu/Al₂O₃ INTERFACIAL REGIONS

R. A. Mayanovic, B. A. Bunker, and N. Jisrawi, Physics Dept., U. of Notre Dame.

Studies on metals have shown that premelting at the surface occurs considerably below the bulk melting temperature, indicating that the wet surface of a solid may act as a nucleation source during melting. Our aim in this work is to study the atomic thermal-motion disorder and vacancy formation associated with the precursors to melting at metal-insulator interfaces using XAFS. Cu K-edge XAFS spectra were taken on Cu/Al₂O₃ multilayers, grown using RF sputtering on optically flat Si wafers having 20 layers each of $\sim 100\text{\AA}$ of Cu and $\sim 100\text{\AA}$ of Al₂O₃, using the glancing-angle of incidence technique in the fluorescence mode. Measurements were taken at several angles of incidence, ranging from ~ 3 to 13 mrad, at room temperature and $420 \pm 2^\circ\text{C}$. The multilayers were studied under two orientations: the x-ray polarization vector parallel (P orientation) and perpendicular (S orientation) to the Cu/Al₂O₃ interface. Analysis of the polarization-dependent EXAFS data reveals a directional anisotropy in the EXAFS Debye-Waller factor, occurring both at room temperature and in a much more pronounced fashion at $420 \pm 2^\circ\text{C}$, indicating that the amplitude of relative Cu atom in-plane motion is larger than for out-of-plane motion at the interface (see Fig. 1). As shown in Fig. 2, we have obtained preliminary results for the incident angle dependence of the relative in-plane EXAFS Debye-Waller factor, measured at $420 \pm 2^\circ\text{C}$, indicating a steady decrease of vibrational amplitudes with depth. This, along with an estimate for the average in-plane vibrational amplitude at that temperature of approximately 3-4% of the nearest-neighbor distance for Cu atoms down to a depth of about 30Å from the interface, indicates that we are close to the premelting regime. The implication of this study is that the interface very likely undergoes premelting via a shearing mechanism rather than the recoil mechanism used to explain premelting at a free surface, for which the directional anisotropy in atomic vibrational amplitudes is reversed. The shearing model may ultimately help to explain superheating of encapsulated solids.

Fig. 1 Normalized EXAFS oscillations for the Cu K-edge, measured at a Cu/Al₂O₃ interface, at 3.9 mrad incidence angle and $420 \pm 2^\circ\text{C}$, in S (solid line) and P (dashed line) orientations.

Fig. 2 The Cu atom in-plane EXAFS Debye-Waller factor (P orient.), as a function of incident angle, measured relative to the out-of-plane component (S orient.) obtained at 3.9 mrad.

L. J. W. M. Frenken and J. F. van der Veen, Phys. Rev. Lett. 54, 134 (1985); P. H. Fuoss, L. J. Norton, and S. Brennan, Phys. Rev. Lett. 60, 2046 (1988).

Supported by Office of Naval Research under contract N00014-89-J-1108.

POLARIZED X-RAY ABSORPTION SPECTROSCOPY

James E. Penner-Hahn, Geoffrey S. Waldo, Kimber Clark, and Timothy Stemmler (University of Michigan)

In preparation for polarized studies of Mn catalase single crystals, we have completed a study of the isotropic XAFS of this enzyme. We determined the oxidation state composition of Mn catalase solutions using a combination of isotropic XANES and EPR. Quantitative curve fitting analyses of the XANES spectra are consistent with the presence of Mn(II) in the reduced protein, a mixture of Mn(III)/Mn(IV) in the oxidized (inactive) protein, and ca. 50% Mn(II) and 50% Mn(III) for the as-isolated protein. The combined EPR and XANES data provide convincing evidence that Mn catalase cycles between Mn(II)/Mn(III) and Mn(III)/Mn(IV) oxidation states (1). Isotropic EXAFS data show that the Mn-Mn interaction is detectable (at 2.7 Å) for the Mn(III)/Mn(IV) derivative, but is not detectable for the Mn(II)/Mn(III) derivative (2). We have examined several binuclear Mn models and find that (at least in the models) Mn-Mn interactions are detectable for structures that two single atom bridges (i.e., bis-oxo; oxo-hydroxo, or bis-hydroxo). This information will now be used in conjunction with polarized XAFS on Mn catalase to identify the bridges in the Mn(II)/Mn(III) derivative of the enzyme.

1. G.S. Waldo, R.M. Fronko, & J.E. Penner-Hahn, "Characterization of the Mn oxidation States in Manganese Enzymes. Evidence for a Mn(II)/Mn(III)/Mn(IV) catalytic Cycle in L. Plantarum Mn Catalase", Biochemistry, in press.

2. G.S. Waldo, S. Yu, & J.E. Penner-Hahn, "Structural Characterization of the Mn site in Mn catalase. Evidence from x-ray absorption spectroscopy", submitted to the J. Am. Chem. Soc.

Supported in part by the NIH (GM-38047 and GM-45205).

HEMEPROTEIN CATALYSIS: STRUCTURE- $\leftrightarrow$ FUNCTION RELATIONSHIP

L. Powers; R.B. Sinclair; C.-S. Chang; S. Hallam; and I. Yamazaki; National Center for the Design of Molecular Function, Department of Chemistry and Biochemistry, Utah State University, Logan, UT 84322-4630

The peroxidases enzymes and the oxygen transport proteins both contain an active site consisting of a single iron-heme with a proximal histidine ligand. Unlike the oxygen transport proteins, however, the peroxidases are capable of catalyzing the oxidation of substrates having very high redox potentials. How can this same structure perform these two radically different functions? X-ray absorption spectroscopy has been used to compare the iron-heme active site structures of the peroxidases [lignin peroxidases, niteroxidase, cytochrome c peroxidase, and microperoxidases (lignin peroxidase, catalase, *Aerthromyces ramosus* peroxidase, and microperoxidases MP8 and MP11)] with those of the oxygen transport proteins [myoglobin and hemoglobin]. The distinguishing feature in the active site structures is the iron to proximal histidine distance which is 0.2 Å shorter for the peroxidases. This results in a longer iron to pyrrole nitrogen average distance [Fe-Np] in the heme of the peroxidases. Significant differences are also found in the Fe-Np distances among the various peroxidases. Do these structural differences relate to their respective reactivity? Preliminary data suggest that the reactivity of the peroxidases with various substrates is correlated to the Fe-Np distances in just the opposite way as that found for the oxygen transport proteins. In the oxygen transport proteins, those states having short Fe-Np distances [R-state] are more reactive with oxygen, while for the peroxidases, those having longer Fe-Np distances are most reactive with substrates.

EXAFS STUDIES OF $[(HB(3,5\text{-}iPr_2pz)_3)Fe^{II}(OBz)(MeCN)]$ (1)

L. Que, Jr., C.R. Randall, A.E. True, R.C. Holz, T.E. Elgren (U. of Minnesota)

$LFel^{II}(OBz)(MeCN)$ (1, L = $HB(3,5\text{-}iPr_2pz)_3$) is the first example¹ of a "nonnuclear iron(II) complex with a symmetrically bound, chelating carboxylate ligand. From its x-ray crystal structure, the three pyrazole nitrogen atoms are bound at an average distance of 2.14 Å, and the two carboxylate oxygen atoms are bound at distances of 2.14 Å and 2.25 Å. The nitrogen atom of the solvated acetonitrile is at a longer distance than that of the pyrazole nitrogens, 2.28 Å. As a well-characterized iron(II)-carboxylate complex, it will be useful as a model in the study of non-heme iron-containing proteins. However, what is most interesting to us is that the unsolvated form of 1 binds dioxygen reversibly below -20° in toluene solution¹. Work is proceeding on the analysis of the x-ray absorption spectrum of the dioxygen adduct of $LFel^{II}(OBz)(MeCN)$. What is reported here is the analysis of the EXAFS data taken of 1, and its comparison with the data obtained by x-ray crystallography.

The energy of the K-edge absorption and magnitude of the 1s-3d transition in the K-edge XANES data are consistent with a six-coordinate ferrous site. The EXAFS simulations which are most consistent with the crystallographic data have 3N/O at 2.08 Å and 3 N/O at 2.20 Å. These distances are shorter than those of the crystal structure by 0.04 Å. The EXAFS can also be modelled with 4-5 N/O at 2.14 Å and 1 N/O at 2.40 Å. The reason for such a long interaction is as yet unknown. The remaining features in the Fourier transform of the data can be modelled by 3 pyrazole nitrogen atoms at 3.05 Å and 6 carbons, probably from the trispyrazolylborate ligand, at 4.11 Å.

Figure 1: A) χk^3 Fourier-filtered (back transform: 0.8 - 4.2 Å) and B) Fourier-transformed χk^3 EXAFS data (x) and simulations (—) of 1.

1. Kitajima, N., et al., *J. Am. Chem. Soc.*, 1990, 112, 6402-3.

Acknowledgements. We would like to thank Dr. Nobumasa Kitajima for kindly supplying us with $LFel^{II}(OBz)(MeCN)$. This work was supported by the N.I.H. (GM 33162). CRR is grateful for a Pre-doctoral Fellowship from the U.S. Public Health Service (T32 GM 07323-16).

EXAFS INVESTIGATIONS OF E7 MUTANT MYOGLOBINS

K. S. Reddy, L. Powers#, H. Tang and B. Chance : Dept. of Biochem&Biophys. Univ. of Penn; and #Utah State Univ.

The E7 histidine is a highly conserved amino acid in heme proteins and has been considered to be involved in the ligand binding process. Here we present the EXAFS results of sperm whale and horse myoglobin mutants in which the E7 histidine is replaced by phenylalanine, valine and E7 tyrosine mutant of horse Mb. The study showed that the replacement of distal histidine with different amino acids directly affects the heme structure of metmyoglobins. Whereas the deoxyMb and MbCO structures were not affected. The EXAFS data also indicated that the metmyoglobins are six coordinated with an 'O' atom being the sixth ligand. Significant changes were found in XANES region of metMb (E7 tyr), possibly due to second co-ordination of tyrosine mutant. This may be due to direct bonding of tyrosine 'O' atom to iron. Detailed modelling of different myoglobin structures is in progress. It is possible that E7 amino acid apart from its steric role for ligand entry, it electronically interacts with the heme either through the sixth ligand or directly.

X-ray absorption spectra of E7His, E7Val, and E7Phe sperm whale myoglobin (1mM sample) after conversion in to k space, multiplication by k^2 and background removal

First shell, filtered data of E7 tyrosine and E7 histidine horse metmyoglobins.

This work was supported by NIH grants HL18708 and RR01633

Photomultiplier/Phosphor X-ray Detector For EXAFS Measurements

K. S. Reddy, S. Khalid, C. Kumar and B. Chance ; Bio-Structures Institute, 3401 Market Street; and Dept. Of Biophys. & Biochem.; U. of Penn.; Philadelphia PA19104.

The success of an EXAFS experiment with biological samples (often <1mM), where fluorescence detection is the only way depends on the detector performance. The detector noise and nonlinearities introduce distortions in the amplitude function, which contains very important information such as coordination number. Our past comparative studies on fluorescent ion chamber (Stern-Heald type) and Photomultiplier/Scintillator-ZnS(PMS) indicated that the PMS detector is linear within the dynamic range of typical EXAFS fluorescence intensities and also that the S/N ratio of EXAFS spectra with a properly designed PMS detector is in fact found to be superior to that of Ion chamber (1).

The PMS detector performance critically depends on the X-ray to visible light conversion efficiency. We have so far used the commercially available Ag doped Zinc Sulfide as phosphor. Though it is efficient in converting X-rays to visible light, It is difficult to couple ZnS(Ag) powder to the large area photomultiplier tube. Hence we studied the the performance of ZnS(Ag) phosphor films prepared in different ways.

The large area (6" dia) phosphor films were made by three different methods. a). By spin coating of ZnS(Ag) onto a glass with polymer support (2). Spray Pyrolysis of $Zn(NO_3)_2$, $Ag(NO_3)$ with thioacetamide, (3). Chemical deposition of ZnS onto a glass substrate using thiourea-zinc nitrate reaction. We have found all the above methods give phosphor quality films and were efficient as X-ray converters. The comparative performance of PMS-Phosphor and Photodiode-Phosphor combinations as an EXAFS detector is currently under way.

This work was supported by RR 01633

X9-A

**Prototype Stopped Flow Apparatus for EXAFS Study of
Enzyme Intermediates**

K. S. Reddy, C. Kumar and B. Chance ; Bio-Structures Institute, 3401 Market Street, and Dept. Of Biophys. & Biochem.; U. of Penn.; Philadelphia PA19104.

With the availability of high intensity x-rays from the wigglers and focussing mirrors, now it is possible to determine the structures of enzyme intermediates using time resolved X-ray absorption spectroscopy. Of the many approaches to time resolved EXAFS, Stopped Flow-EXAFS is an attractive technique. With the expectation that the NSLS X-9A beam line could be focussed to a 1 mm diameter, we have developed appropriate stopped flow device for interfacing with EXAFS technique.

Fig (1) illustrates the design of the stopped flow apparatus which has been under development and to be tested at the beam line in the current year. This system is capable of studying the reactions with an half time of 5msec. The temperature of the apparatus can controlled to -20 to +30C° so that the slow as well as fast enzymes intermediate could be studied by EXAFS. The entire operation of the stopped flow apparatus is being automated so that it can be controlled from the outside of the hut. Evaluation of different window materials for EXAFS measurements is underway

This work was supported by RR 01633

X9-A

IMPROVED SAGITTAL FOCUS OF SCANNING MONOCHROMATOR

G. Rosenbaum, M. Sullivan (Biostructures Institute, Univ. City Science Center), L. Rock (L. Rock Autom. Assoc.)

The sagittally focusing stage of the double crystal scanning monochromator of beam line X9-A has been routinely and successfully used in a series of diffraction experiments. The focusing stage replaces the second stage with a flat crystal which is normally used for X-ray spectroscopy experiments.

These experiments have exploited the flexibility of the focusing optics. For a small angle diffraction study aimed at detecting the meridional reflections of a 800 Å superstructure from a 0.1 mm diameter and 1 mm long specimen we have concentrated a large horizontal divergence onto the specimen with the sagittal focus while retaining resolution in the meridional direction by focusing on the detector with the vertically focusing mirror. In the case of diffraction from a 0.2 mm diameter and 4 mm long bundle of muscle fibers we have focused vertically on the fiber and horizontally on the detector. In both of these cases the specimen to detector distance had to be around 1 m. In cases of small samples and short specimen - detector distances we have focused onto the specimen in both directions.

The focus position has ranged from the front to the rear of the hut corresponding to a focal length of 3.8 m to 5.4 m for the sagittal focus and 2.4 m to 3.8 m for the mirror respectively. The magnifications were from 0.32 to 0.45 for the sagittal focus and from 0.18 to 0.28 for the mirror focus. The width of the sagittal focus was from 0.38 mm to 0.55 mm (FWHM). In all cases the horizontal source size calculated from the measured width of the sagittal focus was between 4s = 1.62 mm and 1.93 mm. The design number is 4s = 1.8 mm. The measured width of the sagittal focus is also almost independent of the horizontal divergence / convergence used.

Vertical foci produced by the mirror of 800 mm length have been measured ranging from 0.14 mm to 0.35 mm (FWHM). The size of the mirror focus is very dependent on the vertical divergence used indicating a substantial spherical aberration. This is to be expected for a mirror of that length. The vertical source size calculated from the smallest height measured is 4s = 1.0 mm whereas the design figure is 4s = 0.5 mm.

At maximum divergence collected by the sagittal crystal and the mirror the flux into the focus is 5×10^{11} photons/sec at a photon energy of 7.4 keV and 200 mA beam current. The average flux density in the center of the focus (FWHM area) is 5×10^{11} photons/sec/mm².

This work was supported by Grant RR-01633, National Institutes of Health, Division of Research Resources.

SMALL ANGLE DIFFRACTION PATTERN FROM SMOOTH MUSCLE

G. Rosenbaum (Biostructures Institute, Univ. City Science Center),
A. Somlyo (University of Virginia)

Small angle X-ray diffraction pattern have been collected from thin sheets of smooth muscle using the beam line in point focusing mode, i.e. with the sagittally focusing crystal installed.

The focal size on the detector was $0.45 \text{ mm} \times 0.14 \text{ mm}$ (horizontal $\times$ vertical, FWHM) with the illuminated size of the specimen of $3.2 \text{ mm} \times 0.5 \text{ mm}$. The specimen to detector distance was 965 mm , the photon energy was 6.00 keV . The flux into the focus was 2.2×10^{10} photons/sec at 200 mA beam current giving a flux density of 3.5×10^{11} photons/sec/mm² in the focus.

The recorded diffraction pattern from *Taenia coli* of a Guinea pig was of good quality showing well resolved peaks at $1/146\text{\AA}$, $1/130\text{\AA}$ and $1/73\text{\AA}$ in meridional direction and at $1/108\text{\AA}$ in equatorial direction. The $1/130$ and $1/108\text{\AA}$ -reflections arise from periods in the thin filament (actin). The $1/146$ and $1/73\text{\AA}$ -reflections correspond to periodicities in the thick filament (myosin). Additional reflections are due to collagen.

The goal of the experiment was to determine the strength of the "second actin layer line" at about $2/385\text{\AA} = 1/193\text{\AA}$. The measured intensity at this diffraction angle was not different from the background.

Since consecutive exposures show changes in the lattice parameters we plan to repeat these measurements with better control of the state of the specimen.

Work supported by NIH grant RR-01633

EXAFS STRUCTURE OF RETROVIRAL GAG ZINC FINGER PROTEIN-RNA COMPLEX IN INTACT RETROVIRAL PARTICLES. I. Sagi¹, L.G. Arthur², L.E. Henderson³, M.F. Summers³, J.W. Bess² and M.R. Chance¹. Georgetown Univ. Washington D.C. 20057¹, NCI-Fredrick Cancer Research and Development Center, Maryland 21701², Maryland Univ. Baltimore, MD 21218³.

Retroviral-type zinc fingers have been suggested to function as zinc binding domains (Berg et al., 1986) that are involved in genomic RNA recognition and packaging. It was shown recently that zinc is tightly bound in the gag protein; however, minimal direct evidence exists on the zinc environment in these structures. We collected EXAFS data at the zinc edge on concentrated samples of Equine Infection Anemia Virus (EIAV). EXAFS structural analysis shows that the zinc present in EIAV is bound to (CYS)₃(HIS)₁ as proposed by Berg et al., where the averaged Zn-S distances to Cys are at $2.30 \pm 0.01 \text{\AA}$ and the Zn-N distance to His is at $2.00 \pm 0.02 \text{\AA}$. EXAFS studies were also performed on an eighteen residue metalloprotein with the sequence of the NH₂-terminal r.t. zinc finger domain from the HIV-1 NC protein. The metalloprotein, which is known to have a (CYS)₃(HIS)₁ coordination, shows longer distances from the zinc site to the four ligands with a Zn-N distance to histidine of $2.05 \pm 0.03 \text{\AA}$ and the average of the three Zn-S sulfur distances to the cysteines at $2.35 \pm 0.02 \text{\AA}$. Our results show unambiguously that zinc is coordinated to multiple sulfurs and at least one nitrogen or oxygen ligand in EIAV. The differences in bond distances between the metalloprotein and EIAV suggest direct nucleic acid interaction; however, more work is needed to confirm this hypothesis. This work is supported by Petroleum Research Fund grant 23257-G3 (M.R.C.), a grant from the Upjohn Company (M.R.C.), and National Institute of Health grant GM 42561 (M.F.S.)

EXAFS STUDIES OF ZINC FINGER BINDING PROTEINS. I. Sagi¹, M. F. Summers², M. Witt³, E. Chén⁴ and M. R. Chance⁵, Dept. of Chemistry, Georgetown U., and Dept. of Chemistry and Biochemistry, U. of Maryland Baltimore County.

The retroviral *gag* polypeptide identifies and forms a complex with viral RNA during assembly and growth stages of the retroviral life cycle¹. The retroviral *gag* proteins are designated as non - classical zinc finger proteins which play an important role in protein - nucleic acid interactions. All retroviral *gag* proteins contain conserved and repeated sequence C-X₂-C-X₄-C-H-X₄-C² which is believed to be the zinc binding domain in these proteins. Direct mutagenesis experiments led to the conclusion that the repeated domains are intimately involved in retroviral RNA recognition³. The nature of the protein - nucleic acid interaction is not completely concluded and still has to be determined. An extensive structural investigation and analysis of these proteins and their nucleic acid complexes can contribute some insight to the nature of the protein - nucleic acid interaction and also might determine the function of the zinc metal site in the zinc finger like domains.

2D high resolution NMR structures of zinc finger like retroviral *gag* proteins were proposed recently by Summers et al.⁴. The *gag* peptide exhibits a unique folding pattern and is stabilized by extensive internal hydrogen bonding and by coordination of three Cys and one His residues to zinc. The backbone folding of the *gag* retroviral peptide is very different from that of a "classical" DNA binding zinc finger. Yet, missing from the picture is the exact structural information around the metal site. This information is necessary in order to derive a direct answer to whether the zinc site in these proteins actually has a role in the retroviral protein - nucleic acid interactions.

EXAFS and X-ray edge data of an eighteen amino acid sequence *gag* peptide, containing one zinc atom per peptide, were collected. EXAFS data of ZnPP and ZnS were also collected and will be used as model compounds for the EXAFS analysis. EXAFS and X-ray edge data will be analyzed at the zinc metal site in order to provide direct structural information at the active finger like domains in the *gag* retroviral proteins. Since these X-ray measurements are directed to the metal site, accurate structural information around the zinc atom in the retroviral *gag* proteins is obtained. This information can be used to determine the function of the zinc metal in *gag* protein - nucleic acid interaction.

REFERENCES

1. Dickson, C., Eisenman, R., Fan, J., Hunter, E., and Teich, N. (1985) In *RNA Tumor Viruses*, 2nd Ed., Vol 1 and 2.
 2. Copeland, T. D., Morgan, M. A., and Orozlan, S. (1984) *Virology* **133** p. 137.
 3. Cotterick, R. J., Henderson, L. E., Hanser, J. P., and Rein, A. (1988) *Proc Natl. Acad. Sci.* **85** p. 8420.
 4. Summers, M. F., Sout, T. L., Kim, E., and Hare, D. R. (1990) *Biochemistry* **29** p. 729.
- * Dept. of Chemistry Georgetown University.
 * Dept. of Chemistry and Biochemistry University of Maryland Baltimore.

SIT DETECTOR CHARACTERIZATION

V. Skita, R. Kelley (U. of Connecticut Health Center) and R. F. Fischetti (Biostructures Institute)

The purpose of these experiments was to characterize the operation of the SIT detector with the high flux available in the point focused x-ray beam on X9A (~5 x 10¹¹ photons/sec/200mA). Standard specimens (fixed and embedded insect flight muscle and polycrystalline DPPC) were used to provide diffraction patterns for testing the linearity of response with flux and the dynamic range of the SIT detector. In a low gain mode, the diffraction patterns recorded with the SIT agreed well the expected patterns from the standard specimens. However, with higher gains a significant number of pixels in the diffraction pattern had reached the maximum number of counts per pixel for even the minimum exposure time (500 msec using the electronic internal gate in the SIT detector). A series of diffraction patterns were recorded with an increasingly attenuated direct beam until a diffraction pattern was recorded for which no pixels were saturated. Based on these measurements exposure times of less than 100 msec are required to avoid saturating any portion of the diffraction pattern when using the full available flux. The dynamic range was determined to be only approximately 400:1 and future efforts will concentrate on improving that range.

This work was supported by grant RR 01633 from NIH.

X9-A

LIGNINASE ACTIVE SITE STRUCTURE

R. Sindair¹, I. Yamazaki², J. Bumpus³, B. Brock⁴, C.-S. Chang⁵, A. Albo⁶, and L. Powers⁷
¹ Center for Biocatalysis Science and Technology, Utah State University, Logan, Utah 84322-4630
² University of Notre Dame, College of Science, Notre Dame, IN 46556

³ Oregon Graduate Institute of Science and Technology, 19600 Von Neumann Drive, Beaverton, OR 97006

⁴ Biotechnology Center, Utah State University, Logan, Utah 84322-4430

The wood degrading fungus *Phanerochaete chrysosporium* secretes a number of extracellular enzymes called lignin peroxidases which are involved in the degradation of both lignin and a number of persistent environmental pollutants. Lignin peroxidase isozyme H₂, a glycosylated protein of approximately 40 Kd, contains a single heme. X-ray absorption spectroscopy (XAS) has been used to probe the local environment of the iron in the active site resting enzyme, reduced enzyme and compound III. For the native, compound III and reduced forms respectively, the average Fe-pyrrrole distances are 2.075 Å, 2.045 Å and 2.03 Å (±0.015 Å); the Fe-proximal distance is 1.92 Å, 1.90 Å and 1.90 Å (±0.02 Å) while the Fe-distal distance is 2.14 Å, 1.70 Å and 2.10 Å (±0.03 Å). The heme iron pyrrrole nitrogen distance is more expanded in ligninase H₂ than in other peroxidases. The possible significance of this is discussed in relation to other heme proteins.

Supported in part by NIEHS Superfund Grant ES 04922 and by the Savannah River Laboratory.

EXAFS OF Fe^{3+} IN ACTIVE SOYBEAN LIPXYGENASE

Robert C. Scarow (Haverford College; spokesperson), Geoffrey Grove (H. C.), Milton G. Trimitsis (H. C.), Mark Nelson (duPont), Rebecca Cowling (duPont)

In September 1990 we spent three days of general user time at beam line X9-A collecting EXAFS spectra of frozen solutions of soybean lipxygenase-1 and of powdered solid samples of appropriate model complexes. The Si 111 monochromator was used for its higher intensity.

Two detectors and a sample of $[\text{NEt}_3][\text{FeCl}_4]$ were placed behind the sample for energy calibration employing the 1s-3d pre-edge peak (left). This relatively sharp peak gives a convenient measure of monochromator resolution. The gaussian half-width of the peaks shown are 1.7 and 1.3 eV with Si111 and Si220, respectively.

The XANES and EXAFS data from this and previous periods of data collection (at beam lines X9a, X11a and X19a) have been analyzed. We have initially focused on forms of lipxygenase in which the non-heme iron is Fe^{3+} , because the EXAFS of the native Fe^{2+} form has been previously published and because oxidation of the iron to Fe^{3+} has been shown to be a prerequisite for enzyme activity. The enzyme was oxidized by addition of 13-hydroperoxy-9,11-octadecadienoic acid followed by dialysis to pH 7 or 9 and concentration. The inhibitor 3,4-dihydroxybenzotriole (DHBN) was added to one of the pH 7 samples.

Fourier transforms ($k^3\chi$, 2-12 Å) of the EXAFS data show only a slight difference in coordination of the ferric enzyme between pH 7 and 9. Analyses of both spectra indicate the presence of one short Fe-O distance of 1.88 Å, probably a bound hydroxide, in addition to five Fe-O/N distances of about 2.13 Å, which are reasonable for high-spin iron(III) bound by several histidine ligands. Binding of DHBN to ferric lipxygenase causes major changes in the EXAFS Fourier transform. Analysis reveals two or three Fe-O distances averaging 1.96 Å, three or four Fe-N distances averaging ca. 2.12 Å, and a short Fe-C shell which confirms bidentate coordination of the DHBN to the iron.

Work at Haverford College is supported by the Petroleum Research Fund of the American Chemical Society (grant 21967-GB3).

SOLUTION SCATTERING STUDIES OF THE POLYPHENOLIC PROTEIN ISOLATED FROM THE BLUE MUSSEL MYTILUS EDULIS.

M. W. Trumble, V. Skita, and R. Kelley, Biomolecular Structure Analysis Center, UConn Health Center, Farmington CT 06030.

Adhesion is a way of life in the marine environment. The Blue Mussel is one of the many organisms that practices it. The mussel lives in the intertidal zone maintaining its position by means of an extracorporeal tendon known as the byssus. The byssus consists of a number of collagenous threads emanating from the base of the mussels foot and terminating at the substrate surface in adhesive plaques. One of the most interesting questions about this system is what is the exact nature of the adhesive contained in the plaques and how does its structure relate to its adhesive properties? The adhesive appears to be a composite consisting of multiple proteins and possibly lipid components. There are two classes of 3,4-dihydroxyphenylalanine (dopa) containing proteins which are believed to play a part in the adhesive composite. These are the 130 kDa dopa rich (= 30 mol%) polyphenolic protein now referred to as MeA, and the ≈ 50 kDa cysteine rich (7-8 mol%) dopa poor (2-3 mol%) MeB. For our initial work, we have been concentrating on MeA, attempting to relate its solution structure to its adhesive function. MeA consists of 70-75 repeats of the 10 amino acid consensus sequence: ala-lys-pro-ser-tyr-hyp-thr-dopa-lys. Because of its large size and repeated nature, MeA is not amenable to NMR studies, therefore we embarked on an x-ray solution scattering study of the protein. Previous circular-dichroism work by Williams et. al. (Arch. Biochem. Biophys. 269:415-422, 1989) indicated that MeA had a random coil conformation in solution. However, modeling studies done by us and by Nagy et. al. (Int. J. Peptides Protein Res. 38:84-89, 1991) indicated the possibility of stable non-random coil secondary structures in the protein. Preliminary solution scattering experiments conducted in our laboratory indicated that MeA was not random coil but assumed a fairly rigid linearly extended conformation with a Radius of Gyration of ≈ 71 Å. Our goal was to refine these experiments by collecting more accurate data. The initial challenge that we faced was adapting our scattering geometry to the beamline. Dr. Gerd Rosenbaum of the Biostructures PRT was instrumental in this endeavor. One of the most worthwhile suggestions made by Dr. Rosenbaum was to introduce limiting slits just before the detector face. This modification of our camera allowed us to use a much less attenuated beam than we had at the beginning of our run. After the initial set up we were able to collect data with a significantly better signal to noise ratio than was previously achieved. However, subsequent analysis of the data and the MeA used in the experiment indicated that the material had degraded to give a species with an Rg of ≈ 45 Å. We are planning to return to NSLS in October to repeat these experiments using freshly purified MeA. This work was supported by a grant from the State of Connecticut Department of Higher Education Cooperative High Technology Program in collaboration with BioPolymers Inc. of Farmington Connecticut.

OF HYDROGENASE FROM *THIOCAPSA ROSEOPERSICINA*
J.P. Whitehead, Cs. Bagyinka, G.J. Colpas, and M.J. Maroney
Dept. of Chemistry U. Massachusetts, Amherst, MA 01003

Hydrogenases (H₂ases) are enzymes that catalyze the reversible oxidation of H₂. The H₂ase isolated from *Thiocapsa roseopersicina*, a purple photosynthetic bacterium, is a Ni₂FeS enzyme. This H₂ase is isolated in an oxidized and inactive state, but may be reactivated by incubation under H₂. During this process several epr signals associated with the Ni site(s) have been observed and reflect the redox state of the enzyme. We report a study of structural changes in the Ni site as a function of redox state of the enzyme using x-ray absorption spectroscopy (XAS).

The energy of the Ni K-edges for Forms A (oxidized, aerobic) and B (oxidized, anaerobic) are identical within the resolution of the experiment. Similarly, the energies obtained for the edges of Form C (reduced, activated) and SR (super reduced) are identical. The difference in energy between Forms B and C is -0.8 eV, and refutes the hypothesis of a large change in electron density at the Ni site upon activation. Nickel K-edge data reveal five coordinate geometries for all of the forms as compared to various model compounds, except for form B which displays a well defined 1s-→3d peak, indicating a six coordinate environment.

The EXAFS spectra have also been analyzed. The first coordination sphere (backtransform window = 1.1-2.3 Å) of Ni for each form of the enzyme is clearly distinct, indicating that the coordination environment of Ni is not static. All four forms exhibit scattering from a mix of N/O and S donor ligands, at 2.05 Å and 2.25 Å, respectively. The fits of Forms A and B are improved by the inclusion of a long S(Cl) at 2.4-2.5 Å. Evidence for this long scatter is completely lacking in Form C.

Fourier-filtered EXAFS spectra arising from outer-sphere scattering atoms in Forms B and C can only be fit by Fe scattering atoms and constitute the first direct evidence of a novel Ni₂FeS cluster in H₂ase. The closest Ni-Fe distances are too long for Ni to be incorporated into an Fe₂S cluster, and suggest a bridged structure similar to that found involving a siroheme and an Fe-S cluster in sulfite reductase.

This work is supported by the NIH.

Ge Detector Dead Time Loss and Its Effects on XAFS

K. Zhang, G. Rosenbaum, and G. Bunker, Biostructures Inst. and Illinois Inst. of Tech.

The 13 element Ge detector (Canberra Model GL0110) has been used extensively at X9-A for XAFS data collection on dilute biological systems, as well as on concentrated model compounds. The detector is superior to the ionization chamber or the ZnS scintillation detector in terms of collecting effective signal counts on a sample with the metal concentration less than a couple of mM or so. However, the detector has its limitations, particularly its dead time loss associated with the electronics.

We have studied the detector dead time loss on some zinc and iron complexes. Figure 1 shows the relationship between a single channel Incident Count Rate (ICR) vs. the fluorescence signal count rate (SC). When ICR is small, the detected events SC can be approximated by a linear function of ICR. With the increase of ICR, SC levels off, then decreases.

Two types of non-linear problems can be introduced into the XAFS data collection. First, the I₀ noise cannot be fully divided out, which will result in random noise, glitches, and background change in XAFS spectra. Second, XAFS spectra for a concentrated sample can be substantially reduced. Figure 2 shows the XAFS $k^2\chi(k)$ data of ZnS collected with the Ge detector (about 90×10^3 ICR counts for the edge step) alone with the spectrum collected using a fluorescence ion chamber (solid line). The XAFS amplitude collected with the Ge detector is seriously reduced due to the dead time loss.

We are currently characterizing each channel of the detector in order to develop an algorithm for the dead time correction.

Figure 1. Plot of the fluorescence signal counts vs. the ICR counts on channel 3. A 2 mM solution Zn complex was used for the calibration.

Figure 2. XAFS $k^2\chi(k)$ data of ZnO solid collected using the Ge detector (solid) and an ionization chamber (dash). The ICR for the edge step was about 90×10^3 per second.

This work is supported by NIH grant No. RR01633.

XAFS STUDIES OF CARBOXYPEPTIDASE AND ITS INTERMEDIATE STATE

K. Zhang¹, D. S. Auld², K. S. Larson², and B. Chance³ (1 Biostructures Institute, 2 Harvard Univ., and 3 Univ. of Penn.)

Despite the extensive study of carboxypeptidase A (CPA) in the last half century or so, the mechanism of the function of the enzyme in hydrolyzing the carboxy-terminal peptide bond has not been fully understood.

We have measured Co(II)-CPA and Zn(II)-CPA ES₂ intermediate state in binding with peptide substrate, using freeze trapping technique. Both Si(111) and (220) double crystal monochromators were used for EXAFS and XANES data collection. The fluorescence signals from Co and Zn K-edge absorption were detected by a 13-element Ge detector.

Figure 1 shows the χ data of Zn(II)-CPA in binding with the peptide compared with the free enzyme. Significant differences of the spectra are observed. Data analysis is underway to reveal the active site structure differences between the free enzyme and the intermediate state.

Figure 1. Plot of χ data of Zn(II)-CPA free enzyme (solid) and its intermediate state binding with peptide (dash).

This work is supported by NIH grant No. RR01633.

An X-ray Absorption Study on Pro-carboxypeptidase

K. Zhang¹, D. S. Auld², and K. S. Larson², 1 Biostructures Inst./UCSC, 2 Harvard Medical School

Bovine pancreatic procarboxypeptidase A (PCP A S₅) is composed of two subunits (I and II). Subunit I is the immediate precursors of carboxypeptidase A (CPD A), while subunit II is the pro-ATEase. It has been proposed that¹ procarboxypeptidase can be activated by 3 steps: (1) activation of subunit II by trypsin; (2) dissociation of subunit I with subunit II; (3) activation of CPD A by cutting down terminal residues by trypsin or activated ATEase.

We used the XAFS technique to examine the structural changes around the zinc ion between the zymogen and activated CPD A. The experiments were carried out at Beamline X9-A of National Synchrotron Light Source. Using a Si(220) double crystal monochromator and a harmonic rejection mirror, the zymogen of 0.7 mM were measured at 300K and 150K.

The initial data analysis shows that the first shell structure of zinc in the zymogen is essentially the same between 300K and 150K, except for increased disorder in the former. Comparison of the XAFS $\chi(k)$ data between the zymogen and activated CPD A (Figure 1) revealed substantial differences. The structural origin of the differences is being investigated.

Figure 1. Plot of χ data of the zymogen (solid) and activated CPD A. (dash).

1. J. R. Urry and H. Neurath (1972) Biochemistry, 11, 4483.

This work is supported by NIH grant No. RR01633.

An XAFS Study on HPV-16 E7 Protein

K. Zhang, and D. Patrick, Biostructures Inst./UCSC, and Univ. of Pennsylvania

Human papillomaviruses (HPV's) are contributing factors in the pathogenesis of human cervical cancer. The HPV E7 gene is transcriptionally active in the disease and has been shown to transform mammalian cells *in vitro*. The HPV gene encodes a protein of 98 amino acids, which contains a cysteine-rich region in its C-terminal half similar to that of adenovirus E1A. It has been shown that the chemically synthesized HPV-16 E7 protein can bind zinc in a 1:1 molar ratio like E1A.

We have performed XAFS experiment on HPV-16 E7 protein produced in *E. coli* at Beamline X9-A. A Si (220) double crystal monochromator and a harmonic rejection mirror were used for the measurements. A 13 element Ge detector was used to collect Zn K-edge fluorescence from the absorption of the X-ray beam of the sample. The protein of 1.3 mM concentration was measured at 150K using a Displex refrigerator.

Figure 1 shows the XAFS $\chi(k)$ data of the HPV E7 protein compared with that of adenovirus E1A protein. The data are quite similar, especially in the XAFS phase, indicating that the similarity of the zinc environment.

Figure 1. XAFS $\chi(k)$ data of the HPV E7 protein (solid) compared with that of E1A protein (dash).

This work is supported by NIH grant No. RR01633.

TRANSVERSE STRUCTURE FUNCTION OF THE LIQUID-VAPOR

INTERFACE OF Ga

A. Acero*, Z.-h. Cai*, E. Flom*, B.-h. Lin*, N. Maskil** and S. A. Rice*

*U of Chicago

**Bar-Ilan U

We have determined the transverse (in-plane) structure function in the liquid-vapor interface of Ga at 10, 15, 20 and 30°C, using grazing incidence x-ray diffraction (GIXD) coupled with out-of-plane rod scans. The in-plane momentum transfer range studied was $\sim 1.5 - 5.5 \text{ \AA}^{-1}$; the out-of-plane momentum transfer range studied was $\sim 0 - 0.5 \text{ \AA}^{-1}$. These experiments were undertaken with two goals in mind:

(i) To learn how stratification of the density distribution along the normal to the interface (longitudinal density distribution) influences the transverse structure function in the interface;

(ii) To determine if the unusual structure of crystalline Ga, (in which there are pseudo-diatom molecules, hence anisotropic interactions) coupled with the increase in density on melting, leads to any unusual structural features in the liquid-vapor interface.

The results of the GIXD measurements at 30°C are shown in Fig. 1, along with the bulk liquid structure function determined by Narten (our measured bulk liquid structure function is identical with Narten's). These data, coupled with the data from the rod scans, lead to the conclusion that the transverse structure function in the liquid-vapor interface of Ga is, within the current experimental precision, the same as the bulk structure function, despite the stratification of the longitudinal density distribution in that interface. The data also show that the anisotropy in the interactions in crystalline Ga is either decreased to an ineffective level or not strong enough at the bulk level to induce unusual ordering in the liquid-vapor interface, such as is seen in the case of extreme anisotropy, e.g. in liquid crystals.

The lack of difference between the transverse structure function of the liquid-vapor interface and the bulk liquid structure function implies that the conventional local density approximation for the properties of an inhomogeneous fluid fails, and that the effective force that supports a density inhomogeneity must be included in any description, at least to lowest order, as in the Fisher-Metfess approximation.

This research was supported by a grant from the National Science Foundation.

High Resolution Small Angle X-ray Scattering Study of Styrene-Butadiene Diblock Copolymers

S.L. Chang * K.S. Liang, and Z. Fu (Exxon)
R.J. Rose and K.J. Jeon (University of Cincinnati)

The structure and morphology of diblock and triblock copolymers in relation to the variation of molecular weight and concentration of the constituent blocks have been extensively studied. Using a high-resolution Bonse-Hart type SAXS set-up at X10A, we have studied mixtures of polystyrene-polybutadiene diblock copolymers of 50K molecular weight with 26K homopolystyrene in various proportion ranging from 0% to 100%. The results are shown in the Figure. We have observed the transitions from lamellar to hexagonal (~40% wt) and to disordered (50% wt up) structures of copolymer blends as the concentration of homo-polystyrene increases.

The lamellar structures are identified for samples 1-6, 100%-60% wt. of copolymer, from the observed diffraction peaks. From these curves, the changes of the layer thicknesses and the lamellar spacing are determined as a function of the weight % of homopolystyrene. Asymmetric swelling of the polystyrene blocks is found for the wt% of homo-polystyrene ranging from 0 to 35% in the binary blends.

The hexagonal structure of cylinders are observed for samples 7 and 8. The radii of cylinders are about 150Å. Forth wt% about 15-35%, the structures of disordered cylinders with the radii about 185-210 Å are identified. For sample #12 (3 wt%), the structure is probably of disordered spherical micelle.

* Permanent address: National Tsing Hua University, Hsinchu, Taiwan

Beamline X10A

DIRECT PHASE DETERMINATION OF LARGE MACROMOLECULE CRYSTALS USING THREE-BEAM X-RAY INTERFERENCE

Shi-Lin Chang¹, H.E. King, Jr.², Mau-Tzai Huang¹, and Yen Gao³
(¹National Tsing Hua U, Taiwan, ²Exxon Research, 3 SUNY-Buffalo)
(³National Tsing Hua U, Taiwan, ²Exxon Research, 3 SUNY-Buffalo)

Phase information is lost in a typical diffraction intensity measurement because the intensity is only related to the product of the structure factor and its complex conjugate. This fact constitutes the well known x-ray phase problem in diffraction physics and crystallography. One can show theoretically that the relative phases can be determined through the use of three-beam diffraction, where two Bragg peaks of a single crystal are simultaneously brought into diffracting condition. From the resulting modulation of the diffracted intensity one can extract the phase. Using x-ray radiation ($\lambda=2.09\text{\AA}$) from Beamline X10A in addition to some experiments on X3A2, we have shown for the first time that such phase information can be obtained on a large-unit cell volume, $350,000\text{\AA}^3$, macromolecule crystal. The scattered intensity from such a structure is quite small requiring the use of a high-brightness synchrotron x-ray source. A single-crystal sample of hemoglobin (S.G. C2, $a=108.24\text{\AA}$, $b=63.13\text{\AA}$, $c=54.54\text{\AA}$, $\beta=110.85^\circ$) was mounted on a 4-circle diffractometer, oriented, then rotated about selected scattering-vectors. These were selected such that during the rotation a second Bragg peak would pass through the Ewald sphere. An example of the results of intensity modulation is given in Fig. 1, where the results of modelling the intensity using the known structure are indicated by the dashed curves. Good agreement between experiment and theory is found indicating the possibility of applying this technique to other such large-volume macromolecule samples.

X-ray multiple diffraction, utilizing the coherent interaction among diffracted waves, has long been considered as a proper means to solve the x-ray phase problem. However, difficulties associated with large crystal unit-cell hindered the applicability of this diffraction technique to large macromolecular crystals. To overcome those difficulties, we have carried out three-beam diffraction experiments with synchrotron radiation and demonstrated that direct phase determination of large macromolecular crystals, about 350,000 Å³ in unit-cell volume, can be achieved.

The three-beam experiments were performed on beam-lines X10A and X3A2 at NLSL, Brookhaven National Laboratory. Hemoglobin crystals were investigated. ω - and γ -scans were employed to track the peak positions of two centrosymmetrically related three-beam interactions. The crystal polarity is determined unambiguously from the peak intensity measurements. Quantitative phase analysis with a newly developed scaling scheme for intensities leads to the accuracy of about 60° in phase angle (Figure 1 shows a three-beam 512/622 diffraction profile of 0° phase).

Fig. 1

GIXD STUDY OF SELF-ASSEMBLED MONOLAYERS

We have studied the ex-situ structural properties of the self-assembled monolayers of $\text{CH}_3(\text{CH}_2)_{17}\text{SH}$ (referred to as C18) chemisorbed on Au(111) and Ag(111) using grazing incidence x-ray diffraction. Because of the competing length and energy scales that are present in these monolayer systems, complex structural behavior is expected. These monolayers form spontaneously from a dilute solution, resulting in uniform passivating layers which are only ~25Å thick. These experiments were performed on the Exxon beamline X10A with a wavelength of $\lambda = 1.09\text{\AA}$ and a (fixed) incident angle of 0.4° using a z-axis diffractometer.

In figure 1, we show a radial scan through the first order diffraction feature due to the ordering of C18/Au(111). The peak position ($1.45\text{\AA}^{-1}$) as well as its azimuthal orientation (inset) demonstrate that the monolayer forms a $\sqrt{3}\times\sqrt{3}\times\sqrt{3}$ structure with a spacing of $5.00\text{\AA}$. In addition, we find that the peak width is only $0.06\text{\AA}^{-1}$, which clearly shows that the monolayer exhibits long range order ($\sim 120\text{\AA}$).

In figure 2 we show similar data for C18/Ag(111) which indicate a more complex structure. The peak position in this radial scan clearly occurs at $1.57\text{\AA}^{-1}$ implying a spacing of only $4.77\text{\AA}$, which is 5% smaller than the $\sqrt{3}\times\sqrt{3}\times\sqrt{3}$ structure that is found on Au; that is, the monolayer is incommensurate. In addition, the azimuthal scan (inset) shows that the monolayer forms 2 domains which are rotated by $\pm 18^\circ$ from the expected $\sqrt{3}$ position. Like C18/Au, this monolayer also exhibits long range order with a peak width of $0.06\text{\AA}^{-1}$.

These data clearly demonstrate that by changing *only* the head group/substrate interaction, a drastic change in the static structural properties can be affected. Since the hydrocarbon tails tend to tilt in order to optimize the 3-d density, it is expected that there should be very large differences in the 3-d structure in these two systems. Indeed, we have measured the intensity distribution of these monolayer peaks perpendicular to the surface (i.e., a rodscan) and have found that the C18 rods exhibit drastic differences which can be attributed to the molecular tilt. On Au we find that the C18 molecules tilt by $\sim 30^\circ$ towards the molecule's next nearest neighbor, while on Ag the molecule is tilted by $< 8^\circ$. These differences in the 2-d spacing and the molecular tilt preserve the 3-d density of the film.

This work was supported by the Division of Material Science and Chemical Science under U. S. Department of Energy Contract No. DE-AC02-76CH-00016

MISFIT DISLOCATIONS IN THE INTERFACE OF EPITAXIAL Nb FILMS ON SAPPHIRE

C. H. Lee (Synchrotron Radiation Research Center, Taiwan)
K. S. Liang (Exxon)

Niobium films epitaxially grown on sapphire substrates are of special interests for studies of model metal-ceramic interfaces [1]. The interface structure of MBE grown Nb films on sapphire substrates was studied using grazing incidence x-ray diffraction and x-ray reflectivity measurements [2]. Satellite reflections accompanying the Nb Bragg peaks and an amplitude-modulated x-ray reflectivity curve were observed in these measurements [2]. Based on these results, we have proposed an atomistic model for the interface, which contains a soliton-type deformed lattice due to the presence misfit dislocations in the interface.

The structural model proposed here is an empirical model lattice with deformation near the dislocation core. We shall call our interface structural model as a soliton-type model. The final model employed in the calculation is shown in Fig. 1. The model has three soliton-type deformed layers of Nb in the interface region. Based on this model, we find that the calculated scattering profiles of this soliton-type model agree quite well with the experimental data of in-plane (110), (220), and (330) reflections of Nb grown on sapphire (0001) surface (Fig. 2).

1. J. Mayer, C. P. Flynn, and M. Ruhle, Ultramicroscopy 33, 51 (1990).
2. C. H. Lee, K. S. Liang, F. S. Shieu, S. L. Sass, and C. P. Flynn, in "Defects in Materials" Mat. Res. Soc. Symp. Proc. Vol. 209, p.679 (1991).

Figure 1

Figure 2

The Structure of Dilute Lamellar L_α Phases of Surfactant Membranes, Ning Lei, C. R. Safinya Exxon Research & Engineering Co., Annandale, NJ 08801

We report on synchrotron scattering studies of oriented dilute stacked-fluid membranes with the interlayer spacing ranging from 40 Å to larger than 350 Å. This dilute L_α phase is stabilized by entropically induced repulsive undulation forces [1], the existence of which was theoretically anticipated by Helfrich [2]. The structure factor $S(q_z)$ of this phase exhibits power-law singular behavior asymptotically close to the first Bragg peak position $q_0 = 2\pi/d$, with the exponent $\eta(q_z) = \text{kgT}q_z^2/8\pi k_B)^{1/2}$ [1]. (K and B are the bulk moduli for curvature and compression, and q_z is the component of the wave-vector normal to the layers.) We have carried out a detailed experimental study of $S(q_z)$ over a large q_z -range. Fig. 1, below shows the first harmonic of the structure factor of an intermediate dilution sample with $d=88\text{Å}$ which indicates that $S(q_z)$ is symmetric about q_0 . For dilute samples with large $\eta(q_0) = 1$ as shown in Fig. 2, with $d=230\text{Å}$, the explicit q_z -dependence of η , which varies only slightly asymptotically close to q_0 , leads to (i) an extremely asymmetric $S(q_z)$ in the non-asymptotic q_z -regimes around q_0 , and (ii) the onset of large small-angle-x-ray-scattering (SAXS) arising from the tails of the central peak, and overwhelming the (SAXS) arising from concentration fluctuations of the crumpled fluid membrane. The solid lines [3] are the fits to the discrete Caillé model [4] describing $S(q)$ of lyotropic smectic-A liquid crystals.

1. C. R. Safinya et al., Phys. Rev. Lett. 57, 2718 (1986).
2. W. Helfrich, Z. Naturforsch. 33a, 305 (1978).
3. N. Lei, C. R. Safinya and R. F. Bruinsma, in preparation.
4. A. Caille, C. R. Acad. Sci. Ser. B 274, 891 (1972)

Figure 1

Figure 2

χ : oil weight percentage

L. J. Pickering, J. T. Lewandowski and A. J. Jacobson (Exxon)

Lithium nickel oxide has been studied as a catalyst for the oxidative coupling of methane in both conventional and electrochemical reactors. *In situ* X-ray diffraction studies have shown that significant structural and compositional changes occur as reaction conditions are varied. Some changes are associated with the transformation of the ordered LiNiO_2 structure to the disordered $(\text{Li,Ni})\text{O}$ solid solution. However, the cation distributions and the nature of the disorder at intermediate compositions have not been studied in detail.

Lithium nickel oxides, $\text{Li}_x\text{Ni}_{1-x}\text{O}$ can be synthesized for values of x from 0 to close to $x=0.5$. For $x < 0.25$ the lithium atoms substitute randomly in the nickel sites of the NiO rocksalt structure. For values of x close to 0.5 an ordered, LiNiO_2 -type phase appears, in which the lithium and nickel atoms order on alternate close-packed (111) planes (fig. 1), giving rise to sharp superlattice peaks in the powder diffraction pattern. These superlattice peaks persist down to values of x close to 0.25, but are increasingly broadened with decreasing x .

As part of an on-going program to study the detailed structure of these compounds, high resolution powder diffraction patterns were recorded for different values of x in $\text{Li}_x\text{Ni}_{1-x}\text{O}$. Figure 2 shows a selected area of the diffraction patterns. Several changes can be seen as x is varied. The rhombohedral distortion (e.g. the splitting of the (006) and (012) peaks, coincident for ideal cubic) decreases, until for $x=0.29$ the structure is cubic within our resolution. The superlattice peaks (e.g. (003) decrease in intensity as x decreases, and also broaden, especially (003) for $x=0.34$. This selective broadening behaviour suggests islands of ordered LiNiO_2 -type material growing coherently within a matrix of random $(\text{Li,Ni})\text{O}$, the mean domain size decreasing with decreasing x . The domain size is estimated to be ca. 700 Å for $x=0.34$ and ca. 80 Å for $x=0.29$.

Fig. 1. Structure of LiNiO_2

Fig. 2. Portion of powder diffraction pattern for various x in $\text{Li}_x\text{Ni}_{1-x}\text{O}$

X10A
Surface-Induced Order on a Cu_3Au (001) Surface

K.S. Liang, H.H. Hung, Z. Fu and S.L. Chang (Exxon)

The possibility of surface effects on the first-order transition was first proposed by Lipowsky [1]. At temperatures away from the bulk transition, the surface can exhibit disordering (if approached from below the transition) or ordering (if approached from above the transition) phenomena. Cu_3Au surfaces have been investigated as a model system for the study of this newly proposed surface phenomenon [2]. Using grazing incidence x-ray scattering we have measured the surface order parameter on a Cu_3Au (001) crystal. The strong effect of a surface-induced order has been revealed [3].

The stoichiometry of the sample was determined by measuring the lattice constant ($a_0 = 3.752 \text{ Å}$) and the bulk transition temperature ($T_c = 390 \pm 2^\circ \text{C}$). Figure 1 shows typical x-ray scans taken around the (101) Bragg peak at temperatures above the bulk transition temperature. Scattering profiles shown in the figure were scans taken in the [100] direction. The broad diffuse background shown is associated with the short-range order from the bulk. The sharp component (coherent length $\sim 350 \text{ Å}$) shown on top of the diffuse background is a direct evidence that an ordered surface layer is present on the (001) surface of the disordered bulk Cu_3Au .

Our x-ray scans taken around the (100) Bragg peak showed similar results; i.e. an ordered surface layer was revealed by a sharp in-plane (100) surface peak was at temperatures above the bulk transition. In Figure 2, the measured peak intensity of the (100) reflection is plotted as a function of temperature, together with that of the (001) bulk peak. While the intensity of the (100) surface peak dropped by a factor of about 50 at bulk T_c , the peak remains to be very sharp (coherent length $\sim 300 \text{ Å}$) at temperatures above T_c . Eventually, the peak merged with the diffuse short-range-order background at 20°C above T_c . Empirically, the observed profiles can be fitted with a Gaussian plus Lorentzian lineshape. We attribute the Gaussian component to the presence of an ordered surface layer.

1. R. Lipowsky, Phys. Rev. Lett. 49, 1575 (1982); J. Appl. Phys. 55, 2485 (1984).
2. H. Deech, L. Mailänder, A. Lied, J. Peisl, F. Grey, R. L. Johnson, and S. Krummacher, Phys. Rev. Lett. 60, 2382 (1988).
3. K.S. Liang, et al. in "Surface X-Ray and Neutron Scattering", ed. H. Zabel and I.K. Robinson (Springer-Verlag).

Figure 1

Figure 2

LOCATION OF IRON IN FRAMEWORK-SUBSTITUTED ZEOLITE L USING HIGH RESOLUTION POWDER DIFFRACTION.

I. J. Pickering, D. E. W. Vaughan and K. G. Strohmaier (Exxon)

We have continued our structural investigations of framework-substituted zeolites by studying iron-substituted zeolite L. Previous EXAFS studies (from X10C) have shown the iron to be in the framework of this zeolite, evidenced by 4-coordination with respect to oxygen, and the presence of an outer shell of silicon. A high resolution powder diffraction pattern was taken at a wavelength of 0.9706 Å, and then a Rietveld profile refinement was carried out using the GSAS software. The starting structural model was taken from the known structure of zeolite L, and the fraction of Fe substituting for Si or Al on the two crystallographically distinct tetrahedral (T) sites was refined, with the constraint that the overall occupancy of each site is unity. Extra-framework cations and water molecules were also included in the model. The best fit between observed and calculated diffraction data was obtained with near equal substitution of Fe onto the two sites, with 9.9% atomic substitution on T1, the 12-ring site and 9.0% on T2, the 6-ring site. Figure 1 shows the results of the fit.

Figure shows the results of Rietveld profile analysis for Fe zeolite L. The observed data points (+), the calculated diffraction pattern and the difference (bent line) are shown. The inset shows a single channel of zeolite L; Fe substitution occurs randomly on the T sites (black).

Effect of Excess Gold On The Surface Ordering Kinetics of the Cu₃Au(001) Surface

S.B. Rivers and W.N. Unertl (University of Maine)*
H.H. Hung, K.L. D'Amico and K.S. Liang, (Exxon Research & Engineering)

We have discovered 1,2 that a thin layer at the surface of a 2% Au-rich Cu₃Au(001) crystal remains ordered at temperatures up to about 15-20 K above the bulk order-disorder temperature T_B = 648 K (Fig.1). We studied this transition and its kinetics using grazing-incidence x-ray scattering. The (001) beam was used to monitor the bulk order and the in-plane (100) beam was used to monitor order in the surface layer. In quasi-equilibrium measurements, we find that the order in the surface layer increases strongly just above T_B and then disorders smoothly as the temperature is raised; i.e. the disordering of the surface layer does not appear to be first order (Fig.1). Kinetics were studied by quenching from the fully disordered state. When the quench is into the temperature range where only the surface layer can order, growth follows a power law in time but with an exponent of -0.04 (Fig.2), about one order magnitude smaller than predicted for either diffusion or curvature driven growth mechanisms. For quenches to temperatures below T_B at which both the surface and bulk can order, growth is also slow and power law growth is not observed even for the longest times studied. The excess Au appears to act as an impurity that inhibits the rate at which long range order is achieved. Auger spectroscopy shows no evidence for changes in the surface composition in the transition region.

*Supported by NSF Grant DMR-8803529.

1. H.H. Hung, S.B. Rivers, K.S. Liang, K.L. D'Amico, and W.N. Unertl, to be published.
2. S.B. Rivers, W.N. Unertl, H.H. Hung, and K.S. Liang, in "Kinetics of Phase Transformation", MRS Symp. Proc. vol. 205 (1991).

Figure 1

Figure 2

Melting And Reassembling Transition Of The 2-D Protein Lattice of Bacteriorhodopsin
 Yi Shen (1), C. R. Safinya (1), K. J. Rothschild (2), B. Thomas (1) Exxon Research & Engineering Co., Annandale NJ 08801, (2) Boston Univ., Boston MA 02215

X10A

The photosynthetic bacteriorhodopsin (bR) protein self-assembles in the plasma bilayer lipid plane of the purple membrane of halobium halobacterium. The dimensions of the membrane sheets are one bR-molecule thick (49 Å), and ~ 0.5 mm by 0.5 mm wide. At low temperatures, a 2-D hexagonal lattice is formed within each membrane sheet. The unit cell consists of a triplet of bR-molecules, with internal degrees of freedom. This protein system is a prototype model of a 2-D lattice embedded in 3-D space [2].

We report on synchrotron x-ray scattering studies of the melting and reassembling phase transitions of oriented stacked purple membranes. The phase diagram is explored as a function of the chemical potential of water (relative humidity) and temperature. By increasing the relative humidity at constant temperature we increase both the inter-layer spacing d and the 2-D lattice constant, which directly affects the intra-layer lattice melting transition temperatures. At these high humidities, with large inter-layer spacings, we find a regime of completely decoupled ordered membranes, which exhibit a reversible melting transition. Figure 1 is a plot on semi-log scale of a $(0,0,q_z)$ scan (along the layer normal) through the first six harmonics of the structure factor resulting from the multilayer stacking with an inter-layer spacing of 200 Å at $T=30^\circ\text{C}$ and 100% relative humidity. Fig. 2 is a plot of a q_z scan in the layer plane through the (10) , (11) and (20) peaks of the 2D hexagonal lattice. The 2D rods in q -space are cylinders due to the random in-plane orientation of the sheets. The asymmetry of the line shape is a characteristic of diffractions from the stack of two dimensional lattices with finite out-of-plane mosaic. Rod-scans (not shown) through the (11) peak indicate a lack of correlation between the layers.

As shown in Fig. 2, the 2D lattice exhibits a melting transition, which we find to be reversible, $T_c=69^\circ\text{C}$. The scattering is expected to have power-law behavior, due to the lack of on-range order in 2D systems with continuous symmetry, with a characteristic exponent $\eta \approx \kappa B T_0 q_z^{-2/\mu}$. Here, μ is the Lamé shear modulus which is straightforwardly related to the inter-particle interactions. In contrast to atomic-based systems, the many measurable diffraction peaks will provide, (i) and accurate measurement of $\mu(T,RH)$ through the melting transition and (ii) a stringent test of melting models regarding the T-dependent nature of higher harmonics.

- [1] Walther Stockenius, Scientific American, June 1976.
 [2] D. R. Nelson and L. Feit, J. Phys. France 48, 1085 (1987); J. A. Aronovitz and T. C. Lubensky, Phys. Rev. Lett., 60, 2634 (1988). J. Toner, Phys. Rev. Lett., 64, 1741 (1990)

B. THOMAS and C. R. SAFINYA, Exxon Research & Eng. Co., Annandale, NJ 08801;
 B. RATNA and R. SHASHIDHAR, N. R. L., Washington D. C. 20375; N. A. CLARK,
 U. Colorado, Boulder, CO. 80309

We report on x-ray diffraction studies of phospholipid-water-ethanol systems (DC8,9PC, with diacylenic fatty acyl chains) that are known to spontaneously produce hollow cylindrical structures known as tubules at low temperatures $< 40^\circ\text{C}$. At higher temperatures, the system gives rise to spherical vesicles which is the natural shape of dilute lipid-solvent mixtures. We have studied these materials (1) in the dehydrated state, and (2) in their fully hydrated state. We present recent results regarding the nature of both the in-plane and inter-layer correlations associated with the wrapped multilayer structure of the tubules in these two states as a function of temperature. Fig. 1 below is of the tubules in their dehydrated state, which indexes to a monoclinic lattice with "in-plane" (e.g., $(h, k, 0)$) peaks, "lamellar" (e.g., $(0, 0, l)$) peaks and "combinational" (e.g., (h, k, l)) diffraction peaks evident. Figure 2 is of the tubules in their hydrated state; the absence of the "combinational" peaks indicate that no interlayer interactions exist within the hydrated tubule assemblies.

SPECULAR AND OFF-SPECULAR X-RAY REFLECTIVITY FROM SPUTTERED Fe/Cr MULTILAYERS

M.F. Toney and S.S.P. Parkin, (IBM Research Division, Almaden Research Center)

Recently, it was discovered that multilayer structures formed from alternating layers of Fe and Cr exhibit an unexpectedly large saturation magnetoresistance (MR).^{1, 2} The origin of this giant MR is not understood at present, but is related to the details of the interface structure (e.g., intermixing, strain, roughness). The goal of these experiments is to understand the X-ray reflectivity from these multilayers, to use the reflectivity to determine the multilayer structure, and to relate this to the magnetic properties.

Reflectivity measurements were conducted at 5980 eV, 11 eV below the Cr K edge, since this results in a reasonably large difference between Fe and Cr scattering factors. Slits defined the detector resolution with the result $\Delta\omega = 0.027^\circ$.

The figure shows the 'peak' ($\omega = 0$) and off-specular ($\omega = \pm 0.05^\circ$) reflectivity from a Fe/Cr multilayer. For large Q , the off-specular reflectivity (or diffuse scattering) significantly contributes to the 'peak' intensity; in fact for $Q > 0.62 \text{ \AA}^{-1}$, rocking curves show that the measured 'peak' intensity is almost completely diffuse scattering. To obtain the true specular X-ray reflectivity, it is essential to subtract this diffuse scattering; failure to do so results in an underestimate of the interfacial roughness and intermixing.

The diffuse scattering is caused by interfacial roughness. Since this scattering peaks at the same Q as the specular reflectivity (see figure), the roughness must be *conformal*: the interfacial meandering of each layer is replicated by all the layers in the multilayer stack. Preliminary analysis of these data shows that the interfacial roughness is $\approx 13 \text{ \AA}$ and that the correlation length parallel to the interfaces is $\approx 150 \text{ \AA}$. We are currently analyzing these data to separate the interfacial intermixing and the conformal and non-conformal roughness.

1. M.N. Baibich et al., Phys. Rev. Lett. **61**, 2472 (1988).
2. S.S.P. Parkin, N. More, and K.P. Roche, Phys. Rev. Lett. **64**, 2304 (1990).

OBSERVATION OF ORDERED CoPt_3 IN SINGLE CRYSTAL Co/Pt MAGNETIC MULTILAYERS

M.F. Toney, R. Farrow, and R. Marks (IBM Research Division, Almaden Research Center)

Multilayers of Co and Pt are candidates for future magneto-optical recording media, since they have large perpendicular magnetic anisotropy and a sizeable Kerr effect, particularly at short wavelengths. The origin of the anisotropy is not understood, but is related to the atomic structure of the interfaces. Several models have suggested that the anisotropy is due to the formation of a CoPt alloy near the interface,^{1, 2} although this is controversial.^{3, 4} The aim of our experiments was to investigate the presence of alloy formation in Co/Pt multilayers.

We have investigated the interface structure of (111)-oriented $\text{Co(3\AA)/Pt(18\AA)}$ multilayers grown by molecular beam epitaxy (MBE) onto c-axis sapphire. Two specimens were studied: one grown at 100°C and the other at 200°C . Data were obtained with an X-ray energy of 8019 eV; a solid state detector was used to eliminate Co fluorescence.

We find presence of an *ordered* CoPt_3 phase at the interfaces in these multilayers. The figure shows rocking scans through the (110) diffraction peaks. These peaks are forbidden in the fct structure of the multilayer, but occur in CoPt_3 , which has an sc lattice. From the integrated intensities of the (110) peaks, we find a substantial fraction of the Co is tied up in the ordered alloy (initial estimates are $60 \pm 30\%$ and $20 \pm 20\%$ for 100° and 200° growth, respectively). The figure shows that the azimuthal widths are broad $\Delta\phi \approx 15^\circ$; the radial widths are also large $\Delta Q \approx 0.3 \text{ \AA}^{-1}$. Thus, the CoPt_3 phase is not well-ordered. Q_z scans have a half-width of $\approx 0.3 \text{ \AA}^{-1}$, indicating the width of the interfacial CoPt_3 alloy is $\sim 10 \text{ \AA}$.

It is remarkable that an ordered CoPt_3 alloy forms in these multilayers at temperatures as low as 100°C , since experiments on bulk samples required heating at $\sim 600^\circ\text{C}$ for several days.⁵

1. R.F. Garcia, J. Appl. Phys. **63**, 3066 (1988).
2. C.J. Chien et al., Proc. Mater. Res. Soc. **231**, xxx (1991).
3. K. Spödl and D. Weller, J. Magn. Mag. Mat. **93**, 379 (1991).
4. B.D. Hermseier et al., J. Appl. Phys. **69**, 5646 (1991).
5. C. Leroux et al., J. Phys.: Condens. Matter **2**, 3479 (1990).

Azimuthal scan for the multilayer grown at 200°C . Data were taken at $Q_z = 0.1 \text{ \AA}^{-1}$; $\phi = 0$ corresponds to the (220) direction. The sharp peaks are caused by long tails of the multilayer diffraction rod that runs from the $\{002\}$ to $\{111\}$ peaks.

X10B

X-RAY REFLECTIVITY STUDY OF OXIDATION INDUCED ROUGHENING OF THE Pt(111) SINGLE CRYSTAL SURFACE UNDER IN SITU ELECTROCHEMICAL POTENTIAL CONTROL*

H. You, D.J. Zurawski, Z. Nagy, R.P. Chiarello, H.K. Kim, and R.M. Yonco (ANL)

Oxidation of the precious metals is one of the most extensively studied subjects in electrochemistry due to its scientific interest. The precious metals, such as platinum, do not form more than a monolayer of oxide in the ambient environment. However, it is known from charge transfer measurements that a much thicker layer of oxide can form in a solution. In this *in situ* x-ray reflectivity investigation we show that the Pt(111) single crystal surface in contact with a solution undergoes an irreversible roughening transition for the oxygen coverage beyond a monolayer.

A series of reflectivity scans of Pt(111) surface shown in Fig. 1 were made in 0.1 M CsF solution at 0 mV vs. Ag/AgCl (3 M KCl) reference electrode, after oxidation for 60 seconds at the potential indicated and reduction at -800 mV. A detailed description of the cell design and experimental procedure are given elsewhere [1]. The scans show little change up to the 500 mV cycle, then the anti-Bragg intensity progressively weakens following the cyclic potential control beyond 500 mV. This indicates that the initially flat surface completely recovers its original surface after reduction of the oxide film formed at potentials up to 500 mV; but the surface irreversibly roughens upon a cycle of the potential beyond 500 mV.

Cyclic voltammetry measurements of the oxygen coverage of the platinum surface indicated that a full monolayer of oxide coverage is not achieved at a potential negative to -600 mV, suggesting that oxidation/reduction of less than a monolayer of oxide film will not roughen the originally flat surface, but the surface will roughen if more than a monolayer of oxide is formed/reduced.

[1] H. You, et al., submitted to the Proceedings of Electrochemical Society Symposium, X-Ray Methods in Corrosion and Interfacial Electrochemistry, Oct. 1991, Phoenix, AZ.

*Supported by DOE under Contract # W-31-109-Eng-38

PRESSURE-INDUCED STAGING TRANSITION IN TiS_2 INTERCALATION COMPOUNDS

Otto Zhou and John E. Fischer (U. of Pennsylvania), and Keng S. Liang (Exxon)

A10B

Single crystal diamond anvil x-ray diffraction reveals a reversible stage-1 to stage-2 transition in $Ag_{0.35}TiS_2$, with an onset pressure < 1 kbar and coexisting stages up to 25 kbar¹. In contrast, no stages higher than stage-1 are observed in $Li_{0.5}TiS_2$ up to 55 kbar. The peak positions were consistent at all pressures with the 1-T stage-1 structure. No in-plane ordering of the Ag ions were observed under pressure. We argue that the different behavior of Li and Ag intercalates cannot be understood in terms of a continuum elastic model, but depends instead on the different interatomic distances and atomic radii. C-axis compressibility of $Li_{0.5}TiS_2$ is 90 % of that of pristine TiS_2 , indicating that occupying half of the octahedral sites in the Van der Waals gap has only minor effect in the stiffness. In both material, K_c is essentially independent of P in the pressure range of this experiment. Figure 1 shows stage-1 (001) and stage-2 (002) peaks of $Ag_{0.35}TiS_2$ at five pressures, along with fits to two Gaussians plus a quadratic background. Each profile is normalized to a constant total integrated Bragg intensity. Figure 2 represents data and fits in the Q region of the $Ag_{0.35}TiS_2$ (100) reflection for four pressures.

1 O. Zhou, J.E. Fischer, and K.S. Liang, *Phys. Rev. B*, Oct. 1 (1991)

This work is supported by DOE Grant DEFG02-87ER45254

X-RAY DIFFRACTION STUDY OF Rb_2C_{60} PHASE DIAGRAM
Qing Zhu, Otto Zhou, Nicole Coustel, Gavin Vaughan, and John E.
Fischer (U. of Pennsylvania) x10B

We have used high resolution x-ray powder diffraction to study the structure and bonding of Cs_60 doped to saturation with alkali metal^{1,2}. The powder Cs_60 was vacuum-doped with alkali metal ($M = \text{K}$, Rb , and Cs) in evacuated sealed tube at temperature range of 200 - 250°C for 12 to 7 days. Figure 1 shows the powder diffraction pattern of K and Cs doped Cs_60 in 0.5 mm capillaries, measured at 0.9617 Å wavelength. The saturation doped $M_x\text{Cs}_{60}$ phase is identified as body-centered cubic structure. We have carried out Rietfeld refinement with chosen space group of $\text{Im}\bar{3}$, from which integrated intensity R factor of $< 7\%$ is achieved. The refinement shows that the saturation doped bcc phase has six alkali metal per Cs_60 molecule in the bcc structure, consistent with weight uptake and atomic absorption results. In the $M_6\text{Cs}_{60}$ phase, all the Cs_60 molecules are orientationally ordered. Figure 2 shows a schematic view of a cube face, from which the essential features of the composite structure may be appreciated. Two equivalent Cs_60 molecules per cell are centered at (0,0,0) and $(1/2,1/2,1/2)$ (the latter is omitted from Fig. 2 for clarity). The C atom positions are localized to within a mean thermal displacement of 0.13 Å, therefore the orientational disorder which is known to exist in pure Cs_60 at 300K³ has been "frozen out". The molecules are oriented with two-fold axes along cube edges, thus the cube faces normal to x, y and z are not equivalent. Twelve M atoms per cell are located at (0,0.5,0.28) and allowed permutations in space group $\text{Im}\bar{3}$. These can be visualized as four-atom "cross" motifs centered on $(1/2,1/2,0)$ and equivalent positions. This motif also has a two-fold axis parallel to the cube edges. A typical motif, as shown in Fig. 2, lies in the $\{001\}$ plane with atoms displaced ± 0.28 along x and ± 0.29 along y from the face center.

FIG. 1. X-ray powder diffraction profiles of K-doped (a) and Cs-doped (b) BaTiO_3 . The intensity is plotted on a linear scale as a function of $Q = 2\pi/d = 2\pi \sin \theta / \lambda$ (Å⁻¹); the data were recorded with 2 θ steps of 0.01° or 0.015° for 2–4 s per point, which gave about 100,000 total counts at the strongest peak. These figures show raw data, including scattering from air and the glass capillary.

FIG. 2. Schematic of the cubic unit-cell face normal to the z direction, derived from Rietveld refinement of the Cs_2C_{60} profile in Fig. 1. Large circles represent Cs^+ in scale with the cube edge. Small circles are C atoms, not representing Cs^+ . An equivalent C_{60} molecule is centred at $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ (not shown). C_1 to C_{12} are equivalent (lookdown from top) are $(0.5, 0.72, 0.1)$, $(0.78, 0.5, 0.1)$, $(0.5, 0.28, 0.1)$ and $(0.22, 0.5, 0.1)$ with respect to an origin at the bottom left corner. The face normal to the x or y directions may be visualized by rotating the diagram $\sim 90^\circ$. This is a consequence of the molecular orientation of C_{60} molecules along cube edges.

This work is supported by DOE and NSF MRL program.

We carried out powder x-ray diffraction at 300K on equilibrated samples of several nominal compositions x in Rb_xC_{60} [1]. In addition to the superconducting fcc ($x=3$), bct ($x=4$) and bcc ($x=6$) stoichiometric phases [2], we find direct evidence for a dilute fcc doped phase $0 < x \leq 1$, and for a substoichiometric bcc phase $x \sim 5$. In contrast, $x = 3$ and $x = 4$ appear to be stoichiometric compounds with $\sim$ zero solubility of Rb vacancies and interstitials at 300K. We conclude that the miscibility gap between undoped and superconducting fcc structures does not extend entirely from $x = 0$ to $x = 3$, and that the saturated bcc phase can support a significant concentration of random vacancies. These results are summarized in a provisional binary phase diagram shown below.

[1]. Q. Zhu, O. Zhou, N. Coustel, G. Vaughan, J.P. McCauley Jr. W.J. Romanow, J.E. Fischer and A.R. Smith III. to appear in *Science*.

[2]. O. Zhou, Q. Zhu, N. Coustel, S. Kycia, Q. Zhu, A.R. McGhie, W.J. Romanow, J.P. McCannell, Jr. A R Smith III and D.F. Cox. *Nature* **351**, 462-464 (1991).

This work is supported by NSF MRL program, DMR88-19885, and by DOE grant, DE-FG02-86FR45254 and DE-FG05-90ER75596

XAFS of two novel tungsten-containing enzymes from the hyperthermophilic bacteria *Pyrococcus furiosus* and *Thermococcus litoralis*.

Y. Gea, G. N. George, R. C. Prince (Exxon Research), S. Mukund and M. W. W. Adams (U. of Georgia).

Tungsten and molybdenum possess quite similar chemistries, yet, while molybdenum enzymes catalyze a variety of different reactions in diverse organisms, until recently, only a single tungsten enzyme had been characterized (1). The marine hyperthermophiles *Pyrococcus furiosus* and *Thermococcus litoralis* both produce tungsten-containing aldehyde-ferredoxin oxidoreductases, which catalyze the conversion of aldehydes to carboxylic acids. Both enzymes contain a single tungsten, but with different amounts of iron and acid-labile sulfide. We have studied the tungsten environment of the *P. furiosus* and the *T. litoralis* enzymes by X-ray absorption spectroscopy at the tungsten L-III-edge. The tungsten EXAFS shows that both enzymes possess very similar coordination environments, and that the tungsten is bound by two terminal oxo ligands in the oxidized (assumed to be W(VI), form plus 3-4 thiolate ligands, and by a single terminal oxo plus 3-4 thiolates in the reduced (assumed to be W(V)) form. This oxo-thiolate coordination is analogous to the most common structural type among the more familiar molybdenum enzymes, such as sulfite oxidase and (desulfo) xanthine oxidase. The EXAFS-derived active site structures for the *T. litoralis* enzyme are shown in figure 1.

Figure 1. Tungsten active site structures of *Thermococcus litoralis* aldehyde-ferredoxin oxidoreductase.

1. Cramer et al. (1985) *J. Inorg. Biochem.* 23, 119-124.

XANES OF MOLYBDENUM COMPOUNDS AND CATALYSTS AT THE MOLYBDENUM L_{2,3}-EDGES

Simon R. Bare (Central Research Catalysis Laboratory, Dow Chemical, Midland, MI), Gary E. Mitchell (Analytical Sciences, Dow Chemical, Midland, MI), and J.L. Gland (Department of Chemistry, University of Michigan, Ann Arbor, MI)

Fluorescent XANES experiments have been performed at the Mo L-edges of selected molybdenum (VI) compounds and on some supported molybdenum oxide catalysts. Figure 1(a) shows the molybdenum L₃-edge of a series of molybdenum reference compounds. The spectra were obtained from powdered samples held in custom designed sample holders to prevent the absorption of water vapor from the ambient. Figure 1(b) shows the second derivatives of the edge region to highlight the differences between the spectra. The white lines are assigned to the dipole allowed 2p→4d transition. Splitting of the white lines is clearly observed, reflecting the ligand field splitting of the final state d-orbitals [1,2]. The splitting is smaller for tetrahedral coordination (2p→e and 2p→t_{2g} transitions) in Na₂MoO₄, CaMoO₄ and MgMoO₄ than for octahedral coordination (2p→eg and 2p→t_{2g} transitions) in MoO₃, consistent with predictions from ligand field theory. The compound (NH₄)₂Mo₂O₇ has molybdenum atoms in both tetrahedral and octahedral coordination in its unit cell; the XANES reflects the presence of both. The relative intensities of the features are related to the number of available orbitals. We have also studied a series of MoO₃/MgO catalysts. We have observed clear differences in the L-edge region as a function of weight loading of the catalyst and as a function of calcination temperature. Correlations between the microscopic local structure and catalytic performance have been established. Further work is underway to put these measurements on a quantitative basis, and to study some secondary effects in the XANES that are not understood.

1. B. Hedman, J.E. Penner-Hahn, K.O. Hodgson in EXAFS and Near Edge Structure,

Springer-Verlag, Berlin, 1984, p.64.

2. G.N. George, W.E. Cleland, J.H. Enemark, B.E. Smith, C.A. Kipke, S.A. Roberts, S.P.

Cramer, *J. Am. Chem. Soc.*, 112 (1990) 2541.

X-ray absorption spectroscopy of Zinc dialkyl dithiophosphates.

Graham N. George, Michael Sansone (Exxon Research and Engineering Co.) and David Martella (Exxon Chemical Co.)

Zinc dialkyl dithiophosphates (ZDDPs) are very widely used lubricating oil additives. They combine anti-oxidant properties with anti-corrosion properties to function as generalized antiwear agents in motor oils, which typically contain about 1 wt. % of ZDDP. Despite their wide use and importance their mechanism is of action remains essentially unknown. Two different forms are used industrially, the neutral (or normal) ZDDP, and the basic ZDDP. We have used XAFS spectroscopy to structurally characterize both types of commercial ZDDPs and model (in which the alkyl *i*-butyl) ZDDPs both in solution in oil and in the solid state. Figure 1 shows the Zn K-edge EXAFS Fourier transforms of basic and neutral ZDDP in solution. A curve fitting analysis of the raw EXAFS data indicates that the structure of the basic form changes on solvation in oil, from being a tetranuclear Zn species with a μ_4 bridging oxo with three Zn-Zn interactions in the EXAFS, to a species in which only one Zn-Zn interaction is detected in the EXAFS. This can be seen in the Fourier transforms shown in figure 1 as a decrease in the Zn-Zn peak amplitude at about 3.1 Å. In contrast no detectable change was observed for the neutral ZDDP on dissolving in oil. Similar results were obtained with the commercial product in solution (no well defined solid phase exists), and no major structural changes were observed on freezing the sample.

Earlier EXAFS work on ZDDPs by Burn *et al.* (1) reportedly confirmed the then currently accepted core structures for basic and neutral ZDDP, which are similar to the model compound structures shown in figure 1. Although our ZDDP structure for both the basic and neutral solids is essentially in agreement with the conclusions of Burn *et al.* it is important to note that their studies were of solutions, for which we detect quite major structural changes for the basic ZDDP. We believe that the most probable reason for this discrepancy is that Burn *et al.* neglected Zn-P interactions in their analysis, and thereby arrived at erroneous conclusions.

Figure 1. Zn K-edge EXAFS Fourier transforms of neutral and basic ZnDPPs, data were collected at a temperature of 4K, and the Fourier transforms were calculated over a *k*-range of $1-18.2 \text{ \AA}^{-1}$, with phase correction for sulfur. The insets show the structures of the model compound ZDDPs.

Burn, A.J., Joyner, R.A., Meehan, P. & Parker (1986) M.A. J. Chem. Soc.: Chem. Commun. 12, 982-984.

Polarized single crystal Sulfur K-edge X-ray absorption spectroscopy of substituted thiophenes.

Graham N. George, Michael Sansone and Martin L. Gorbaty (Exxon Research and Engineering Co.)

In an attempt to determine the factors which govern the variability of the structure observed in sulfur X-ray absorption K-edge spectra we have investigated the effects of chemical substituents upon the spectra a large number of 2- and 3- substituted thiophenes. It was found that the spectra depended very strongly upon the electron-withdrawing tendency of the substituent, the white-line feature showing a splitting which depended quantitatively upon the well-known Taft-Ingold resonance parameter σ_r , and was essentially independent of the inductive component σ_i . In order to better understand these observed changes and trends we have performed polarized single crystal XAFS on substituted thiophene derivatives. Figure 1 shows the sulfur K-edge spectra of a single crystal of thiophene 3-carboxylic acid. The molecules are arranged so that the plane of the thiophene rings are inclined at about 30° to the C-axis of the crystal (see inset on figure 1). Thus, for crystal orientations such that the X-ray *e*-vector is parallel to the C-axis, σ^* resonances are expected to be maximal, and for orientations such that the *e*-vector is perpendicular to the C-axis, σ^* resonances are expected to be minimal. It can be clearly seen in figure 1 that the lower-energy feature of the white line is maximal when *e* is parallel to C and that the higher-energy feature is maximal when *e* is perpendicular to C. We conclude that these represent $1s-\pi^*$ and $1s-\sigma^*$ resonances, respectively. This assignment agrees with the earlier calculations of Hitchcock and co-workers (1) on (unsubstituted) thiophene, who concluded that the white line was comprised of a $1s-\pi^*$ and a $1s-\sigma^*$ (C-S) resonance.

Figure 1. Single crystal polarized X-ray absorption sulfur K-edge spectra of thiophene 3-carboxylic acid the upper set of traces shows the spectra for crystal orientations with the C-axis parallel, perpendicular, and at 45° to the X-ray *e*-vector (dashed line). The lower trace shows the spectrum of a non-oriented powder, a the inset shows the crystal structure (2) of thiophene 3-carboxylic acid.

(1) Hitchcock, A.P., Horsely, J.A. and Stohr, J. (1986) J. Chem. Phys. 85, 4835.

(2) Hudson, P. and Robertson, J.H. (1964) Acta Crystallogr. 17, 1497.

X-ray fluorescence XAFS microscopy at 50 μ m resolution.

Graham N. George and Michael Sansone (Exxon Research and Engineering Co.)

We have used the focussed X-ray beam on X10C, together with our 13 element energy dispersive germanium detector array, to perform spatially resolved X-ray absorption spectroscopy at a resolution of 50 μ m with high sensitivity in the concentration range 10-100ppm. In these and in previous lower resolution experiments we investigated the calcium, iron, copper, zinc and arsenic contents of thin sections of Green-River oil shale. The experiment basically proceeds by positioning the sample with a high-precision sample stage so that the beam interrogated different parts of the sample. The dimensions of the beam were defined by using two sets of high-precision tantalum slits. The sample could either be "raster-scanned" while monitoring the X-ray fluorescence of particular elements (figure 1) and an "image" of

Figure 1 Typical MCA spectrum of a 50 $\times$ 50 μ m region of a section of Green-river oil shale.

Figure 1. Orientation dependent Fourier transformed data of nitrogenase crystals.

Here, we report the first Fe K-edge polarized EXAFS study of MoFe protein crystals. For this experiment, we employed the X10C focusing mirror with its newly installed position sensitive feed-back system, which allowed us to measure high quality EXAFS data from our small (0.5mm x 0.25mm) samples. The comparison between this result and our previous data shows that the position sensitive feed-back system is essential for our small samples.

Figure 1 shows a series Fourier transformed data (transform range: $k = 2-14 \text{ \AA}^{-1}$) of single crystal MoFe protein taken at different orientations. The beam E vector is in the crystal bc plane, and β is the angle between E and b axis. For all orientations, like in the solution sample, the spectra are dominated by Fe-S and Fe-Fe interactions at 2.29 Å and 2.61 Å, as well as the long Fe-Fe distance 3.7 Å. The spectra also show orientation dependence. The relative intensity of Fe-S and Fe-Fe peaks depends very much on orientations. This ratio changes by approximately a factor of 2 when β changes from 0° to 90°. The averaged spectrum over all orientations is very similar to previously measured solution data. Preliminary analysis suggests that there are some longer interactions at 4.2 Å and 4.7 Å.

This work was supported by DOA under grants no. DOA-91-37305-6514.

(1) Flank, A.M.; Weininger, M.; Mortenson, L.E.; and Cramer, S. P.; *J. Am. Chem. Soc.* **1986**, *108*, 1049-1055.

Figure 2 Arsenic K_{α} image of a piece of Green River oil shale together with XANES spectra showing chemical variations at specific regions on the sample.

XAS STUDY OF METAL PARTITIONING AT WATER/OXIDE INTERFACES

K. F. Hayes (U. of Michigan), L. E. Karz (U. of Maine), and J. E. Penner-Hahn (U. of Michigan)

Sorption processes often control the fate and transport of metal ions in aquatic environments. The specific structural configuration of sorbed metal species will, in turn, control the extent of sorption in a particular system. Over the past several decades, surface complexation models (SCMs) have been developed to describe the sorption process. These models assume that metal ions form complexes with surface functional groups in a manner similar to metal-ligand association in solution phase. SCMs vary depending on the assumptions regarding the structure and composition of the sorbed species and on the physical description of the interfacial potential. In some cases, the metal ion forms a simple complex consisting of one metal ion or complex reacting with one surface site whereas in other cases multinuclear surface complexes have been identified. In this study, XAS was used as a tool to provide structural information required for accurately incorporating mass law expressions for the formation of multinuclear species and surface precipitates into SCMs. In addition, the results of the study were used to assess the impact of particle-particle interactions on the formation of multinuclear species.

α -Al₂O₃, Co(II) and NaNO₃ were used in all experiments. In previous XAS studies using γ -Al₂O₃ and Co(II), Chisholm-Brause et al.¹ used XAS to identify changes in the predominant sorption process as a function of surface coverage. In these studies, best patterns in the EXAFS and the size of the second shell Co-Co peaks in the radial structure functions (RSFs) at about 3 Å were used to conclude that multinuclear species formed at surface coverages greater than 0.25 $\mu\text{moles/m}^2$, and the size and/or number of these complexes increased with increasing coverage. It was also suggested that the presence and increasing size of a fourth shell Co-Co peak at about 6 Å, unclear at what phase shifts, was indicative of medium range order in the surface complexes formed. It is still unclear at what coverages the predominant surface species changes from a multinuclear sorbed complex to a surface precipitate.

Batch sorption experiments were performed by adding appropriate quantities of solid, Co(NO₃)₂·6H₂O, and acid or base to polypropylene centrifuge tubes and mixing the solutions at 25°C for two to several days. The samples were then centrifuged and the pH and cobalt concentration in the supernatant were determined. Samples of the wet paste were placed in aluminum sample cell holders and immersed in liquid nitrogen. The samples were maintained in a frozen condition until analysis. The solid concentrations in these batch experiments were varied from 0.2 $\mu\text{mol/L}$ to 200 $\mu\text{mol/L}$ and cobalt concentrations ranged from 2x10⁻⁶ M to 1x10⁻⁵ M. Surface coverages ranged from 0.24 $\mu\text{moles/m}^2$ to 3 $\mu\text{moles/m}^2$. All of the sorption samples were collected in fluorescence mode using a 13 element germanium detector at 4°K. Comparison of the EXAFS of sorbed systems to those of model compounds, Co(OH)₂ and aqueous Co, and to data obtained by Chisholm-Brause et al.¹ indicates that multinuclear species formed at coverages greater than 0.24 $\mu\text{moles/m}^2$. Further, comparison of RSFs and XANES collected at 2 and 100 $\mu\text{mol/L}$ indicates that multinuclear species formation occurs at nearly equivalent surface coverages. The size of the fourth shell Co-Co peak in the RSFs increased significantly at coverages above 0.64 $\mu\text{moles/m}^2$ indicating that the order of the structure becomes more defined with increasing surface coverage. Based on these spectroscopic observations, three multinuclear species were postulated and formation reactions for these complexes were incorporated into the SCM. The model constants for these reactions were determined from data collected at 0.2 $\mu\text{mol/L}$, 2x10⁻⁶ M. Whereas reasonable predictions were obtained for the 2 and 20 $\mu\text{mol/L}$ data, the model significantly overpredicted the extent of multinuclear species formation for the 100 $\mu\text{mol/L}$ samples. The reduction in multinuclear species formation at 100 $\mu\text{mol/L}$ may be due to particle-particle interactions.

In order to evaluate the effect of particle-particle interactions, a recirculating batch reactor (RBR) was used to generate sorption samples. In this reactor, the particles are packed in a column at concentrations in excess of 500 $\mu\text{mol/L}$. The particle concentration in the column and external recirculating solution was maintained at 2 $\mu\text{mol/L}$. The results showed that there was a significant reduction in sorption in the RBR presumably due to electrostatic interactions between the particles. To determine whether the reduction in sorption capacity was accompanied by a concomitant change in the type of surface complex formed, the EXAFS of the RBR samples and the batch reactor samples were compared. Based on the XANES and EXAFS information obtained, it was concluded that the RBR samples consisted primarily of mononuclear sorbed species. The lack of multinuclear species formation was due to the large reduction in surface coverage caused by particle-particle interactions.

¹ Chisholm-Brause et al. (1996) Nature 384, 528-531.

This research was sponsored by the National Science Foundation grant BCS-8958407. Travel and subsistence funds to perform experiments at NSLS were sponsored through DOE/NSLS faculty research support group.

IN SITU XANES OF AN IRON PORPHYRIN IRREVERSIBLY ADSORBED ON AN ELECTRODE SURFACE

S. Kim (CMRU), I. T. Bae (BP), M. Sandifer (CMRU), P. N. Ross (LBL), R. Carr (SSRL), J. Wolcik, (NIST), M. R. Antonio (BP) and D. Scherson (CMRU)

In situ iron K-edge X-ray Absorption Near Edge Structure (XANES), has been employed to examine the axial coordination of μ -oxobis[iron meso-tetrakis(methoxyphenyl) porphyrin] (FeTMP)20, irreversibly adsorbed on a high area carbon substrate, Black Pearl (BP), as a function of applied potential. The cyclic voltammetry of 40% (FeTMP)20/BP in 0.1 M NaOH, shown in Fig. 1, was found to display a characteristic redox peak centered at about -0.72 V vs SCE. The in situ fluorescence XANES spectra of the same material, shown in Fig. 2, provided conclusive evidence that the coordination about Fe(3⁺) in the supported, fully oxidized macrocycle (middle curve) is remarkably different from that about Fe(2⁺) in the corresponding fully-reduced macrocycle (bottom curve). In the adsorbed, oxidized state, (FeTMP)20 retains its μ -oxo character and it undergoes a two-electron reduction to yield predominantly four-coordinate square planar spectra of supported (FeTMP)20 compared to that of the same material in FeTMP without axial ligation. The differences between the in situ XANES spectra of supported (FeTMP)20 and the same material in FeTMP without axial ligation, a process that could generate a multiplicity of molecular relaxation induced by the removal of the macrocycle from its original crystal lattice, a process that could generate a multiplicity of closely related iron environments (...Fe(3⁺)-O-Fe(3⁺)...), and thus lead to a global/broadening alteration of the XANES.

Figure 1. Cyclic voltammetry of 40%w/(FeTMP)20/BP in 0.1 M NaOH aqueous solution. Scan Rate: 5 mV.s⁻¹.

Figure 2. Normalized iron K-edge fluorescence XANES data for microcrystalline (FeTMP)20 (top curve) and 40% (FeTMP)20/BP in 0.1 M NaOH at (-0.43 V vs SCE) (middle curve) and NaOH at (-0.90 V vs SCE) (bottom curve). The vertical scale is offset for clarity.

This work supported by the Gas Research Institute

INVESTIGATION OF THE CHEMISTRY OF CHROMIUM IN ANTI-CORROSION FINISHES ON ALUMINUM

F. W. Lytle and R. B. Greeger, The Boeing Co., Seattle, WA 98124

We have obtained data comparing the role of Cr in three different anti-corrosion surface finishes used on Al alloys. The different surfaces were prepared by 1) the Alodine process, which is the industry standard, in which the cleaned part is dipped into a solution containing chromic acid which forms a hydrated $\text{Cr}^{3+}/\text{Cr}^{6+}$ oxide on the surface, 2) anodized in dilute chromic acid and 3) anodized in dilute boric/sulfuric acid followed by sealing in dilute chromic acid. The standard by which all aluminum anti-corrosion surfaces are measured is exposure for a week in a salt spray chamber. Satisfactory performance is the formation of no corrosion pits. Our samples were measured before and after salt spray (SS) in order to determine the changes if any in the amount and chemistry of the Cr.

The data were obtained with a detector in which the electron yield and fluorescent x-ray signals were measured, simultaneously, which allowed us to sample approximately the top 1000 Å and the entire surface layer, respectively. The respective edge jumps were used to estimate the amount of Cr present and were calibrated using Ar ion sputtering and Auger spectroscopy. Treatments 1, 2 and 3 resulted in relative retention of total Cr in the surface layer of 100, 35, and 25 (after SS it was 80, 10 and 10), respectively. The important entity in corrosion prevention appears to be Cr^{6+} which could be accurately determined by monitoring the intense 3d pre-edge peak which is characteristic of Cr^{6+} . Examples of the Cr K edge spectra from a failed Alodine panel, before and after SS, where all of the Cr^{6+} was reduced to Cr^{3+} by the salt spray is shown in the figures. In contrast more of the Cr was Cr^{6+} in the anodized panels both before and after SS. Treatments 1, 2, and 3 resulted in 25%, 90% and 80% (after SS occasionally to 0% but typically to 7%, 30% and 75%), respectively, of the total Cr as Cr^{6+} .

Near-Edge vs EXAFS Studies of Oxides: Understanding Without the Tears

G. Meitzner (Exxon Research and Engineering Co.) and E. Hayden (Department of Pharmacology, Rutgers University)

Structural studies of oxides, with EXAFS, are deterred by the complexity of the materials. The radial distribution functions, especially of mixtures of oxides, often show multiple unresolved coordination spheres. Moreover, the signal contributions from disordered phases may be deceptively weak.

For these reasons, we have concentrated on the use of K near-edge spectra to characterize oxides. In one set of experiments, a series of spectra were measured during the in situ reduction of an oxide of gallium. Reduction was indicated by three effects:

- 1) Diminished K-edge shift relative to gallium metal;
- 2) Gradual qualitative change in near-edge fine-structure;
- 3) Changes in the oxide structure as reflected by EXAFS.

Edge shifts were measured relative to a germanium standard. Fractions of reduced and oxidized gallium were calculated from fits to the near-edge spectra, and from fits to EXAFS spectra. These EXAFS fits used the entire manifolds of peaks below 5 Å.

We found that there was a good correlation between degrees of reduction as indicated by any pair of these three sets of results. For example, we fitted the series of Ga K edge spectra with standard spectra from tetrahedral and octahedral Ga(III) samples, and from a reduced Ga sample. The mole fraction of (tetrahedral plus octahedral) Ga(III) standards, from EXAFS fits, is plotted against Ga K edge shift in figure 1. This study indicates that near-edge spectra alone can provide useful chemical information, even though the EXAFS may be intractable.

Figure 1. X-ray absorption spectra were measured from a gallium oxide as the sample was reduced in H_2 at 500°C. EXAFS fits vs edge shifts are correlated, suggesting that guidance in EXAFS analysis can be drawn from edge analyses.

DIFFRAXAFS: a novel technique for probing local atomic environment.

Ingrid J. Pickering, Michael Sansone, James Marsch and Graham N. George (Exxon Research & Engineering Co.)

Powder X-ray diffraction (PXD) and X-ray absorption fine structure (XAFS) spectroscopy are both well-established techniques for structural investigations of solid-state samples. Both techniques use X-rays over similar energy ranges, and both give structural information, but they depend upon rather different physical phenomena, and are generally practiced by quite separate communities. The technique of anomalous X-ray diffraction depends upon changes in the X-ray scattering factor at X-ray energies (E) close to an X-ray absorption edge. The atomic scattering factor f is given by $f(E) = f_0 + f'(E) + if''(E)$, where f_0 is the non-anomalous part of the scattering factor, and f' and f'' are the real and the imaginary parts of the anomalous scattering, respectively. The function f'' is simply proportional to the atomic absorption coefficient multiplied by the X-ray energy (E), and f' and f'' are interrelated by Kramers-Kronig relationships. Both f' and f'' are therefore expected to contain structure analogous to XAFS. By monitoring the intensity of a particular diffraction peak, while simultaneously correcting for sample absorbance, we have observed this structure, or DIFFRAXAFS, for a number of different compounds, in experiments including measurements at the K-edges of manganese, cobalt and nickel. Our experiments have shown that DIFFRAXAFS is capable of discriminating individual components in mixtures, and, by judicious choice of diffraction peak, of probing the individual sites in compounds that contain more than one type of site (i.e. differing oxidation states and/or coordinations of a metal site). We expect DIFFRAXAFS to be a useful tool in future studies probing the structure of a variety of solid-state crystalline materials.

Figure 1. A single sweep of the XAFS region of the XAFS together with a single sweep of the (211) peak DIFFRAXAFS of a 5wt.% sample of KMnO_4 . The $1s-3d$ peak and other features of the XAFS can be clearly seen reflected in the DIFFRAXAFS.

X-ray Absorption Spectroscopy of transition Metal Substituted Zeolites.

Ingrid J. Pickering, Graham N. George, Grayson H. Via, Dale Sondericker, Karl G. Strohmaier and David E. W. Vaughan (Exxon Research and Engineering Co.)

The application of zeolites as ion exchangers, sorbents and in hydrocarbon conversion catalysis and separations is of importance to Exxon. These materials are typically aluminosilicates whose framework structures have pores of molecular dimensions. A large number of different framework geometries are known, and these give rise to a variety of shape-selective properties. In addition to geometry, the properties of zeolites can be modified by controlling other parameters such as the Si/Al ratio, or by framework cation substitution. There is considerable interest in the incorporation of different elements into the framework, as their presence may favorably modify the properties of the zeolite to provide more effective materials. XAFS is a valuable technique for establishing the degree of substitution into the tetrahedral framework positions in synthetic zeolites. Structural information derived using techniques such as XAFS is (ultimately) required to correlate the structural properties with the catalytic or sorbent behavior of the zeolite, and thus to tailor the material to improve its performance.

In studies using X10C we have used XAFS spectroscopy to investigate the local structures of first-row transition metal ions substituted into a number of different zeolites, namely Cr, Mn, Fe, Co, Ni and Zn substituted into a range of different synthetic zeolites. The EXAFS Fourier transform of FeLTL, together with the Fe K-edge XANES spectrum, is shown in figure 1. The XANES spectrum shows an intense (relative to other FeLTL XANES) $1s-3d$ transition, which is suggestive of a tetrahedral geometry, and none of the features typically associated with other geometries. In addition the EXAFS curve-fitting analysis indicates a coordination of 4 oxygens with a characteristically tetrahedral bond-length (1.86Å), additionally there are indications of outer-shell Fe-Si interactions, although some disorder is obviously present. Overall we conclude that for FeLTL there is good evidence for transition metal substitution into the framework. Other metal substituted zeolites, such as CoLTL (not shown), show evidence that the bulk of the framework metal is *not* incorporated into the framework (i.e. 6 coordinate to oxygen with pronounced Co-Co interactions in the EXAFS).

Figure 1 Fe K-edge EXAFS Fourier transform and XANES spectrum (inset) of FeLTL. The solid line shows the experimental data, and the broken line the results of a curve fitting analysis. Data were recorded at 4K using a Si(111) monochromator.

SULFUR K-EDGE X-RAY ABSORPTION STUDIES OF PHOTOGRAPHIC MATERIALS

T. Smith (Kodak); J. DeWitt (Stanford U.); K. Hodgson (Stanford U.); B. Hedman (SSRL)

Sulfur K-edge x-ray absorption spectra have been measured for a series of model compounds relevant to the study of sulfur-containing photographic surface species. Included in this series of compounds are cyanine and merocyanine sensitizing dye nuclei, silver and gold sulfides and complexes of other sulfur-containing ligands such as thiocyanate, mercaptotriazoles, and thioureas. Data were measured at room temperature in excitation fluorescence mode using a Stern/Heald/Lyle detector and Si (111) double-crystal monochromators.

The spectra of sensitizing dye nuclei indicate a sensitivity to the local sulfur environment (Fig. 1). Both the heterocyclic sulfur of the cyanine dyes and the terminal sulfur of the merocyanines reveal a high-intensity transition at approx. 2474 eV. The terminal sulfur also exhibits a sharp, low-energy feature between 2470 and 2472 eV which has no counterpart in the cyclic sulfur environment. Oriented single-crystal polarized measurements reveal that this low-energy feature is polarized normal to the molecular plane and is attributed to a 1s-to- π^* transition. Complexation of terminal-sulfur ligands by silver and gold atoms results in an increased intensity of the predominant edge feature at 2474 eV and a loss of the 1s-to- π^* feature. (Fig. 2) indicating a bonding interaction between the π orbitals of the terminal sulfur and the filled d orbitals of the metal ion.

We recently have measured low levels (1-2 monolayers) of sulfur on surfaces at grazing-incidence angle (10-20 mrad). A comparison of the surface spectra of a sulfided silver bromide sheet crystal (produced by chemical treatment with aqueous thiosulfate) and that of silver sulfide evaporated onto quartz with the spectrum of silver sulfide powder provides evidence for silver sulfide being the form of sulfur present on the chemically-treated silver bromide surface and indicate our ability to detect low levels of sulfur on these surface samples (Fig. 3).

Research was supported by funds from Eastman Kodak Company. Data were measured at SSRL and NLSL, which are supported by the NIH and DOE.

TITANIUM GERMANIDE FORMATION

X-11A

D.B. Aldrich, C.L. Jahncke, R.J. Nemanich, and D.E. Sayers
North Carolina State University, Dept. of Physics, Raleigh, NC 27695-8202

The evolution of titanium germanide was studied, from initial metal deposition through the formation of the stable TiGe₂ phase (isomorphous with the C54 phase of TiSi₂) [1]. Samples for this study were prepared by depositing Ti on cleaned Ge substrates and subsequently annealing them at temperatures from 300°C to 700°C in UHV. Various Ti layer thicknesses and annealing times were used to study the effects of these parameters on the Ti-Ge reaction. Samples with 400 Å Ti layers and annealed in 50°C steps were used to characterize the Ti-Ge reaction.

EXAFS and XANES data were obtained from the Ti K-edge using a total electron yield detection configuration. After annealing at 460°C there is no significant Ti-Ge reaction detected. At 500°C there are no distinct XANES features indicating the titanium has completely interdiffused with germanium and the structure formed is not well ordered. At 700°C distinct features appear in the XANES data signaling the formation of a well ordered structure. The XANES from the samples annealed at temperatures between 500°C and 700°C contain, to varying degree, features from both the 500°C XANES and the 700°C XANES indicating these samples are composed of a mixture of the two phases.

The 550°C-650°C XANES were fitted with a linear combination of the 500°C and 700°C XANES to determine the degree of intermixing of phases in these samples. No significant change in composition was observed from 500°C to 550°C. Between 550°C and 600°C the TiGe₂ nucleates and begins to grow. At 600°C the sampled area is composed of ~17 percent TiGe₂. A visual inspection of the 650°C sample reveals two distinguishable regions. The composition of region 1 is ~34 percent TiGe₂ suggesting nucleation and/or growth rate is temperature dependent. The composition of region 2 is ~92 percent TiGe₂ which indicates the nucleation sites were not evenly distributed in this sample.

Normalized XANES of UHV prepared samples and a metallic titanium standard. Samples were annealed for 10 minutes at the specified temperature. XANES were taken in regions R1 and R2 of the 650°C sample.

Relative X-ray Energy (eV)

[1] D.B. Aldrich, C.L. Jahncke, R.J. Nemanich, and D.E. Sayers, to appear in Mat. Res. Soc. Symp. Proc. 221

Annular Detector for Obtaining Accurate X-Ray Absorption Spectra in fluorescence from Thick Concentrated Samples

D.L. Brews, D.M. Pease, P. Chartier, J.I. Budnick., Physics Department and Institute of Materials Science, University of Connecticut.

C.E. Bouldin, National Institute of Science and Technology.

Z. Tan, Brookhaven National Laboratory

Previously, we have demonstrated that accurate X-ray absorption spectra (XAS) may be obtained in fluorescence mode from concentrated bulk absorbers by using a normal x-ray incidence, small exit angle x-ray detection geometry (1). As previously applied, this method yields counting rates sufficiently inferior to rates obtained by the transmission or electron yield techniques, that excessively noisy data may result. We have now increased the count rates obtainable in the small exit angle fluorescence mode by utilizing an annular detector made with silicon photo diodes. The annular detector is used in conjunction with a modified dispersive refrigerator so that temperature dependent, accurate, low noise XAS may be obtained from concentrated bulk specimens, including insulators. The method is less surface sensitive than the electron yield technique. The figure below shows a copper XAFS spectrum obtained by small exit angle fluorescence using a conventional ion chamber detector and by an annular detector using photodiodes. The annular detector is seen to give a spectrum with significantly lower noise.

(1) D.M. Pease, D.L. Brews, Z. Tan, J.I. Budnick, and C.C. Law, Phys. Lett. A138 (1989) 230.

XAFS spectrum from bulk copper by the small exit angle fluorescence method, using a conventional ion chamber and using an annular detector of silicon photodiodes.

This work is supported by the U.S. Department of Energy.

MORPHOLOGY OF SUPPORTED PLATINUM CATALYSTS X-11A PREPARED FROM PLATINUM CARBONYLS ON MAGNESIUM OXIDE

J.-R. Chang and B. C. Gates (University of Delaware, Newark, DE 19716)

A structurally unique platinum carbonyl cluster, $[\text{Pt}_{15}(\text{CO})_{30}]^{2-}$, first reported by Chini et al., was prepared for the first time on the surface of a solid. The solid was MgO powder. The preparation involved a surface-mediated synthesis from Na_2PtCl_6 in the presence of CO under mild conditions. The goal of the research was to prepare a supported Pt catalyst with a unique structure; this was made by decarbonylation of the $[\text{Pt}_{15}(\text{CO})_{30}]^{2-}$ at 120°C under vacuum. The formation and decarbonylation of the $[\text{Pt}_{15}(\text{CO})_{30}]^{2-}$ were characterized by EXAFS spectroscopy. The decarbonylated cluster was found to have an average Pt-Pt bond distance of $2.76 \text{ \AA}$ and an average coordination number of 3.70 . These results suggest that the trigonal prismatic structure of the Pt carbonyl precursor framework had been retained after the decarbonylation; thus the Pt_{15} on MgO is modelled as a rod-like structure. The sample was recarbonylated, and the process was followed by infrared spectroscopy; the spectra of the resulting species indicated the presence of Pt particles on the support; the Chini cluster was not reformed. EXAFS spectra of the recarbonylated species indicated an increase in the average Pt-Pt bond distance to $2.78 \text{ \AA}$ and an increase in the coordination number to 6.0 . The increase in coordination number indicates aggregation of the metal and a change in the cluster morphology to one resembling hemispheres. The lengthening of the Pt-Pt bond resulting from the CO -induced change in the morphology is consistent with a change from trigonal prismatic clusters to clusters that more closely resemble bulk Pt. The effects of CO adsorption on the electron density of Pt clusters were evident in the increase of white line intensity. The result is consistent with the IR spectra; the absorption band of the adsorbed CO is shifted to lower energy than that of the free CO molecule.

• CO treatment caused increase in Pt-Pt coordination number and bond distance.
• Increased amplitude of 2nd and 3rd Pt-Pt shells indicates more 3-D character.

We gratefully acknowledge the support of the U. S. Department of Energy, Division of Materials Sciences, under contract No. DE-FG02-87ER13790 for its role in the operation and development of Beamline X-11 at the National Synchrotron Light Source. The NSLS is supported by the Department of Energy, Division of Materials Science and Division of Chemical Sciences under contract number.

J. R. Chang, S. K. K. Purnell, B. C. Gates (University of Delaware, Newark, DE 19716)

Pt-Re γ -Al₂O₃ catalysts were first introduced for naphtha reforming in 1968. Despite their wide use and superior performance, the role of the Re in enhancing the stability of Pt-Re γ -Al₂O₃ catalysts remains unclear. To help understand the role of Re, model catalysts were made from structurally well-defined organometallic clusters, Re₂Pt(CO)₁₂ and Re₂(CO)₁₀, as presursors. The surface species present during the reduction process have been characterized by EXAFS, DRIFTS (Diffuse Reflectance Infrared Fourier-Transform Spectroscopy), and TPDE (Temperature-Programmed Decomposition). These complementary characterization techniques provide evidence of Pt-Re bimetallic interactions.

After the reduction at 150°C under H₂ of Re₂Pt(CO)₁₂ γ -Al₂O₃ and Re₂(CO)₁₀ γ -Al₂O₃, Re subcarbonyls were formed. The Pt was reduced to form three-dimensional metallic crystallites. As the reduction temperature was increased to 300°C, the coordination number of Re-CO decreased concomitantly with the formation of Pt-Re interactions. The formation of a Pt-Re bimetallic interaction is indicated by the following evidence: (1) high metal-metal shell contributions (2nd, 3rd, 4th) have been observed for Re-subcarbonyl (150°C) in the presence of Pt; the bond distances of these high-shell contributions are very close to those of Pt foil; (2) the intensity of the white line (Pt edge) decreased as the reduction temperature increased from 150°C to 300°C, although the coordination number of Pt did not increase significantly; (3) the IR absorption peak characteristic of Re subcarbonyl was shifted to higher wave numbers upon treatment at 300°C, when Pt was present; (4) Re-subcarbonyls were catalytically reduced only in the presence of Pt.

Evidence of Pt-Re Interaction

- existence of peaks at 150°C
- shift of peaks to shorter distances

We gratefully acknowledge the support of the U. S. Department of Energy, Division of Materials Sciences, under contract No. DE-FG02-87ER13790 for its role in the operation and development of Beamline X-11 at the National Synchrotron Light Source. The NSLS is supported by the Department of Energy, Division of Materials Science and Division of Chemical Sciences under contract number.

ELECTRON-YIELD AND FLUORESCENCE XAFS STUDY OF E-BEAM DEPOSITED Co/Pd MULTILAYERS*

M. Choi(CUMC), J.H. Joo(KIST), S.K. Kim, J.S. Kang, Y.-P. Lee(RIST), S.-C. Shin(KAIST), S.M. Heald (BNL)

Transition metal-transition metal(TM-TM) multilayers such as Co/Pd or Co/Pt are promising materials for magneto-optical recording. The magneto-optical and magnetic properties of these multilayers have recently been studied extensively, since these TM-TM multilayers are suitable for the use of shorter wavelength laser beams which will be one of future development trends. The sublayer thickness or the total film thickness dependence of these useful properties has been studied with various techniques, and the results have been of much interest. In the present study, some Co/Pd and Co/Pt multilayers with varying sublayer thicknesses have been prepared for XAFS measurements. Electron-yield and fluorescence XAFS data have been obtained for the e-beam evaporated and sputtered Co/Pd multilayers, and some preliminary analysis results of e-beam deposited samples are presented here. Some samples were deposited on quartz substrates and had the total thickness of about 1300Å (2.1Å/13.5Å & 2.2Å/4.5Å). The rest were deposited on float glass substrates and had the total thickness of about 100 or 200Å with various sublayer thicknesses. The Fourier transforms of the Co K-edge EXAFS data for Co/Pd multilayers show a splitting of major peak, and the magnitude ratio of these split peaks varies systematically with the ratio of the Pd sublayer thickness to the Co sublayer thickness. The Fourier transforms of the Pd K-edge EXAFS data for the same samples do not show as significant a variation with the Co to Pd sublayer thickness ratio. Further analyses of the data including the fluorescence correction are in progress.

FIG.1. Fourier transforms of the Co K EXAFS data (The 2nd y-scale is for the dashed line.)

FIG.2. Fourier transforms of the Pd K EXAFS data (The 3A/4A sample had a Pd predeposition.)

* Supported by KOSEF, PLS, and USDOE under contract no. DE-AC02-76CH00016

XAFS OF STRAINED DISCONTINUOUS AU FILMS

G. A. DeRose and R. W. Hoffman, Case Western Reserve University

Thin films of Au were physically evaporated at a pressure of 10^{-8} Torr in the thickness range 25 to 200 Å and characterized by electron yield and fluorescence X-ray Absorption Fine Structure (XAFS) spectroscopy. Substrate materials used include polyethylene, vinylidene chloride, and Kapton™ polyimide films. Scans about the Au L_{II} absorption edge (11919 eV) were obtained as a function of substrate strain between 0 and 18% for polyethylene and 0 and 5% for vinylidene chloride and Kapton™. As indicated in Figure 1 below, initial results show little difference between 1% and 5% strain, and further analysis is required to explain this result. This work was supported by the Office of Naval Research under contract number N0014-89-5-1555.

Figure 1: k^1 -weighted Fourier transform magnitudes of electron yield EXAFS data for 20 Å of Au on Kapton™ polyimide. (a) 1% strain, (b) 5% strain in polyimide.

ELECTRON-YIELD EXAFS STUDY OF MULTILAYERS FOR MAGNETO-OPTICAL DEVICE MATERIALS*

M. Choi (CUMC), J. H. Joo (KIST), J. G. Lee, C. S. Yang (SKC), K. Kim, D. Whang (POSTECH), S. M. Heald (BNL)

By using a helium-flow electron detector, electron-yield EXAFS measurements were made on Tb/FeCo and Tb/Fe multilayers as well as on a TbFeCo alloy and some reference thin films. These multilayers were prepared by sputtering onto glass substrates, and had the total thickness of 1000 Å with various sublayer thickness values. Rare earth-transition metal (RE-TM) amorphous alloy films are useful as magneto-optical recording device materials, and those of TbFe and TbFeCo are especially promising. Tb/FeCo multilayers exhibit superior characteristics compared with a homogeneous alloy film of TbFeCo, and the perpendicular magnetic anisotropy of the multilayer is also much greater than that of the alloy. As the repeat distance decreases and the number of interfaces increases, the crystalline structure of each layer in the bilayer case changes into an amorphous structure. The structure of the Tb/FeCo 100 layer sample is almost like that of the amorphous TbFeCo alloy. But there still is a slight difference between the two. Also to probe any structural difference between perpendicular and parallel directions, the EXAFS for the Tb/FeCo 100 layer sample was first measured with the polarization of x-ray parallel to the sample plane, and then measured again after the sample was tilted by 50° . When the EXAFS data and Fourier transforms of the two cases are compared, there is a very small difference which may need to be studied more carefully in the future with an electron detector operating at low temperatures.

Co/Al multilayers are useful as magnetic recording media. Alternately deposited Co-Al thin film layers were prepared by sputtering onto glass substrates and the electron-yield EXAFS measurements were also made on these samples. The individual layer thickness was close to that of a monolayer, and the Ar pressure was varied for different samples. An Al rich side sample (Co₁Al₂) made with the Ar pressure of 7 mTorr showed a splitting of major peak in the Fourier transform, indicating the possible existence of CoAl phase and/or more than a single Co phase environment. Whereas a diffractometer scan could not lead to any phase information, since the major peaks of Co, CoAl, and Al overlap. Further analyses of the EXAFS data are in progress.

FIG. 1. Raw data of electron-yield EXAFS measurements

FIG. 2. Fourier transforms of the Co K-edge EXAFS data

* Supported by KOSEF, PLS, and USDOE under contract no. DE-AC02-76CH00016

XAFS CF CHEMICALLY SYNTHESIZED II-VI NANOCCLUSERS

R. H. Felton (Ga. Tech), A. J. Bard (U. Texas), and C. Hoener (U. Texas)

Nanometer size clusters of II-VI semiconductors are synthesized by performing particle growth within mesoscopic cavities. Structural information on the clusters is accessible by X-ray diffraction, optical spectroscopy, and XAFS, *inter alia*. We have analyzed XAFS data for a series of clusters, composed of Zn^{2+} , Cd^{2+} , and Se^{-2} and grown in inverted micelles, present within the polymer, Nafion, or the AOT-H₂O-hexane ternary component system. Selected EXAFS results are shown in the table, where the first element listed corresponds to the K-edge measured. Samples in Nafion and denoted D are synthesized under anhydrous conditions, while W samples are the D samples grown out in deoxygenated and hydrated Nafion films.¹ Phenyl-capped clusters, annealed in pyridine, have different nominal diameters indicated by a number (Å).

A marked loss or decrease of second shell coordination number N_2 is seen in all samples (Fig.). In the $SeZnW$ and $SeZnW$ cases, the Debye-Waller factor is essentially identical to that of bulk standards. We can exclude the small particle size or a decreased mean-free path as causes and conclude that the nanoclusters in the W samples are composed of approximately equal amounts of crystalline and amorphous phases. For samples with a first-shell coordination number N_1 of ~ 3 , the average diameter is ~ 20 Å, while for a value of 4, the diameter is ≥ 50 Å.

Heterojunction structures were prepared, using a nanocluster as the inner seed for growth of the heterostructure. The EXAFS analysis indicated extensive surface reconstruction in both samples. Etching of Zn is found in $(ZnSe)_n(CdS)_{out}$, in agreement with EDS results and an X-ray diffraction study of a similar capped structure, $(ZnS)_n(CdSe)_{out}$.² For this species, N_1 is 3.9 (Se-Cd) and 3 (Cd-Se). In $(CdSe)_n(ZnS)_{out}$, N_1 is 1.9 ± 0.1 (Se-Cd), 0.75 (Se-Zn), 1.8 (Cd-Se), and 2.5 (Cd-S).

Table. EXAFS Results

Nafion				
Sample	N_1	N_2	$\Delta\sigma^2_{2nd\ shell}$ (Å ²)	
$SeZnD$	2.9 ± 0.1	<1	-	
$SeZnW$	3.8	6 ± 1.5	0.002	
$SeCdD$	2.9	<1	-	
$SeCdW$	4	6	0.001	
$CdSeW$	4	-	-	
Phenyl capped particles				
$SeZn16$	3.1	<1	-	
$SeZn32$	3.1	<1	-	
$SeCd32$	3.1	<1	-	
Sample	$N_{1,Se}$	$N_{1,O}$	N_2	
$ZnSe12$	2.6 ± 0.2	1.2	<4	
$CdSe15$	3	2	-

Fig. SeZnW in Nafion (—), bulk SeZn (---).

1. Sirokin, E.S., Brown, R.M., Rabenberg, L.K., Salomon, K., Bard, A.J., Campion, A., Fox, M.A., Mallouk, J.E., Webber, S.E., White, J.M., J. Phys. Chem., 1990, 94, 7543.
2. Kertun, A.R., Hull, R., Opila, R.L., Bawendi, M.G., Steigerwald, M.L., Carroll, P.J., Brus, L.E., J. Am. Chem. Soc., 1990, 112, 1327.

X-11A,B

GLANCING ANGLE X-RAY STUDY OF CRYSTALLIZATION OF AMORPHOUS GE AT THE GE-AL INTERFACE

S. M. Heald and J. K. D. Jayanetti, Brookhaven National Laboratory

The crystallization temperature of amorphous Ge and Si is substantially reduced for thin films in contact with various metals. The mechanism is generally thought to be due to enhanced diffusion caused by the interaction of the semiconductor and metal. To study this process in more detail we have carried out glancing angle EXAFS and reflectivity measurements at beamline X-11A and diffraction measurements at beamline X-11B on samples consisting of 2000 Å of a-Ge covered with 1000 Å of Al. Crystallization was observed to begin at the interface at temperatures of 120-150° C, with substantial disruption of the interface. Time dependent diffraction measurements which are shown below allowed the kinetics of the crystallization to be determined. After an initial nucleation period the Ge diffraction intensity follows a $t^{1/2}$ time dependence. This indicates that after the reaction moves away from the interface it is diffusion limited, likely due to the need for the metal atoms to diffuse to the crystallization front.

The x-ray diffraction data obtained from the Ge-Al thin film while annealing at 152° C. On the left is the raw data, and on the right the intensity of the Ge(111) reflection is plotted versus the square root of the annealing time.

This work supported by the U.S. Dept. of Energy, Div. of Mat. Science under contracts Nos. DE-AC02-76CH00016 and DE-AS02-80-ER10742. We also thank Dr. C. K. Saw of Hoechst-Celanese for loan of the x-ray detector.

S.M. Heald and G.M. Lammie (BNL)

The X-11A beamline at the NSLS has been upgraded to include a dynamic sagittally focussing second crystal in the monochromator. The focussing element is a triangular shaped Si(111) crystal which achieves an approximate cylindrical bend by applying a force at a single point on the apex, which is adjusted to maintain a constant focus at the sample during the XAFS scan. The crystal is positioned such that the optic axis is along the fixed base of the triangle, so that no vertical translation is required as its radius is changed. The approach used to minimise antilastic bending was to cut an array of slots in the diffracting face, such that the bending is permitted by the resulting weak links[1]. The monochromator accepts 3.3 mrad of horizontal radiation and a spot size of $\sim 0.5 \text{ mm}^2$ is achieved at the sample. Between 6 and 10 keV, where the system performs most favourably, total intensity gains over flat crystal running (with a 1 cm horizontal slit width) are between a factor of 2 and 4, without any observable noise introduced from the bending process. Bending errors hinder the performance at higher energies and also preclude effective harmonic rejection, by piezo crystal de-tuning, throughout the operational energy range. The bending behavior can be improved with better specifications on the crystal cutting. Thus, with the focussing system as it currently functions, the most useful application is for fluorescence measurements of dilute samples. Background subtracted XAFS data illustrating the improvement in statistics are shown below.

[1] B.W. Batterman and L.F. Berman, Nucl. Instr. and Meth. **208** (1983) 327.

This work was supported by the U.S. Department of Energy under contract Nos. DE-FG05-89ER45384 and DE-AC02-76CH00016.

INTERFACIAL REACTIONS OF Ni-Cr ALLOYS AND ALUMINUM* S. M. Heald, Zhenguan Tan and J. Janyaneti (Brookhaven National Laboratory).

We have studied the interface reactions of Ni-Cr alloy and Al thin-film bilayers using glancing-angle x-ray reflectivity and extended x-ray-absorption fine structure (EXAFS) techniques. The diffusion behavior and concentration profile (Fig. 1) were inferred from the reflectivity data, while the reaction products were identified by the depth-selective EXAFS spectra. The Al/Ni-Cr bilayers were prepared by evaporation in UHV with a clean interface (Clean) or an interface exposed to 600 Langmuir oxygen (O-exposed). No significant interdiffusion was observed at room temperature in either the Clean or the O-exposed samples. Upon annealing for 10 minutes, reaction starts at about 250 °C in the Clean sample and 310 °C in the O-exposed sample. The oxygen impurity at the interface retards the initial reaction and also modifies the diffusion behavior (see Fig. 1). In both Clean and O-exposed systems, the reaction starts between Ni and Al forming NiAl₃; only at higher temperatures Cr and Al react and form CrAl₃. As a consequence of the different diffusion rates of Ni and Cr, the original face-centered-cubic Ni-Cr alloy near the interface transforms to a thin Cr-rich layer, in which body-centered-cubic chromium was identified at sufficiently high temperatures.

Figure 1. Ni and Cr density profile for Al/Ni-Cr on float glass obtained from fitting the x-ray reflectivity at 8633 and 8033 eV. (a) clean Al-(Ni-Cr) interface and (b) interface with 600 Langmuir O₂ exposure.

*We acknowledge the support from the U. S. Department of Energy under Contract Nos. DE-AC02-76CH00016 and DE-AC05-80-ER10742.

A. Honji, L. U. Gron, J. -R. Chang, B. C. Gates (University of Delaware,
 Newark, DE 19716)

Supported catalysts with well defined and nearly molecular structures were prepared by the reaction of $[\text{HRe}(\text{CO})_5]$ with the surfaces of MgO powders, one being partially dehydroxylated (about 55%) and the other being almost fully dehydroxylated (about 93%). The surface species were characterized by infrared and EXAFS spectroscopies. The initial surface species were molecularly adsorbed $[\text{HRe}(\text{CO})_5]$, which, upon heating to 80°C in H_2 or under vacuum, gave rhenium subcarbonyls with three CO ligands and three oxygen-containing ligands provided by the MgO surface. Infrared spectra indicate C_{3v} symmetry in both structures, with bands at 2011 (vs), 1895(vs), and 1862(sh) cm^{-1} for rhenium subcarbonyl on the partially dehydroxylated MgO and at 2017(vs), 1908(vs), and 1867(sh) cm^{-1} for the rhenium subcarbonyl on the dehydroxylated MgO. The average bond distances were determined by EXAFS spectroscopy: on the partially hydroxylated MgO, the Re-C, Re-O* (O* is a carbonyl oxygen), and Re-O_s (O_s is a surface oxygen) bond lengths were 1.87, 3.11 and 2.13 Å, respectively; and on the dehydroxylated MgO, the Re-C, Re-O*, and Re-O_s bond lengths were 1.91, 3.12, and 2.13 Å, respectively. These distances indicate greater electronic backbonding donation from the Re to the CO on the partially hydroxylated MgO than from the Re to the CO on the fully dehydroxylated MgO surface and suggest electron transfer from the MgO to the Re. The MgO surface can thus be modelled as a source of rigid oxide ligands like those in molecular analogues. The X-ray absorption near edge data confirm a higher electron density on the Re atoms bonded to the oxygens of partially dehydroxylated MgO than on Re atoms bonded to dehydroxylated MgO. The family of rhenium subcarbonyls on MgO constitutes some of the best-defined metal-containing structures on surfaces.

A structural model of the surface species

We gratefully acknowledge the support of the U. S. Department of Energy, Division of Materials Sciences, under contract No. DE-FG02-87ER13790 for its role in the operation and development of Beamline X-11 at the National Synchrotron Light Source. The NSLS is supported by the Department of Energy, Division of Materials Science and Division of Chemical Sciences under contract number .

S. M. Heald, Z. Tan, and J. K. D. Jayanetti, Brookhaven National Laboratory

The properties of vacuum deposited metals are often taken to be the same as the bulk metals. However, in many cases the deposition process can introduce defects which substantially reduce the density of the film. In this work x-ray reflectivity was used to measure the film density of a number of metal-Al bilayer structures. Substantial reductions from bulk density were found for a number of metals as are shown in table 1. All of these films were deposited in UHV conditions at room temperature with rates of about 1 Å/sec. The samples consisted of approximately 1000 Å of metal covered by about 500 Å of Al to prevent oxidation. Fits to the resulting reflectivity curves were used to extract the Al and metal density. In all cases the Al density was close to the bulk value. The samples were also annealed at temperatures ranging from 150° to 400° C. In most cases the films reacted, but for Co-Al there was a clear density increase after annealing to 380° C. This is shown in the figure which compares the reflectivity before and after annealing. Clearly the Co critical angle near 0.45 degrees increases. The corresponding density change is from 0.92 to 0.97 with annealing.

Table 1

Metal	Density
Al	0.28
Mn	0.87
Cr	0.93
Co	0.92
Ni	0.97
Cu	0.96
Pd	0.93

Figure 1
 Comparison of the reflectivity from a unannealed (dashed) and annealed Co Al bilayer.

This work supported by the U.S. Dept. of Energy, Div. of Mat. Science under contracts Nos. DE-AC02-76CH00016 and DE-AS05-80-ER10742.

EXAFS STUDIES OF THE INTERFACIAL PHASE OF HIGHLY MILLED Sn/Ge COMPOSITES

J.K.D.S. Jayanetti, S.M. Heald and Z. Tan, Brookhaven National Laboratory

Mechanically milled Sn/Ge powders have been shown to undergo a depression of melting point in Sn and further studies show that there is a decrease in the enthalpy of fusion with the decreasing Sn content^{1,2}. We have made EXAFS and X-ray diffraction measurements on Sn/Ge powders having compositions ranging from 20 to 50 vol.% Sn. Our measurements show the formation of an amorphous Sn phase which increases with the decreasing Sn content. At 20 vol.% Sn essentially all of Sn is found to be in this phase. Initial results indicate that this phase is due to the possible formation of α -Sn and/or an α -Sn/Ge alloy which is stable even at temperatures above Sn(β -phase) melting point.

1.J.S.C. Jang and C.C. Koch, J. Mater. Res., Vol. 5, No. 2, Feb. 1990.

2.C.C. Koch, J.S.C. Jang and S.S. Gross, J. Mater. Res., vol. 4, No.3, May/Jun. 1989.

The k^2 weighted Fourier transformed EXAFS data obtained at 20[K] for α -Sn, 20 vol.%Sn and β -Sn samples respectively.

This work supported by the U.S. Dept. of Energy, Div. of Mat. Science under contracts Nos. DE-AC02-76CH00016 and DE-AS05-80-ER10742. We also thank Dr. C.K. Saw of Hoechst-Celanese for his help in making diffraction measurements.

 σ^2 -DERIVATIVE METHOD FOR DISORDER IN PbTiO₃

M. JOO, A.M. EDWARDS AND D.E. SAYERS (N.C. State U.)

Temperature-dependent XAFS above the Pb L(III) edge (13055 eV) in PbTiO₃ has been measured between 10K and 300K using a Displex refrigerator. A single shell contribution of the nearest neighbor Pb-O bond in PbTiO₃ to the XAFS spectrum was calculated by inverse Fourier filtering and the relative Debye-Waller factors (DWFs) of the first shell at different temperatures were calculated by exponential fitting. A new analysis technique, the σ^2 -derivative method[1], was then used to get the information about the locations of the phase transition temperatures and their natures. We can confirm two phase transitions below the room temperature in PbTiO₃ using this new analysis technique. A step at about 200K and a peak at about 100K in the first derivative of the total DWFs (Fig.1) indicate that these phase transitions are of second order and of first order respectively. The phase transition temperatures in our results are in good agreement with those of others[2]. However, the natures of these low temperature phase transitions were not reported before.

[1]M. Joo, A.M. Edwards and D.E. Sayers (To be published).

[2]J. Kobayashi, Y. Uesu and Y. Sakemi, Phys. Rev. B28, 1866 (1983).

Acknowledgement:

This work was performed at X-11A Beamline at the National Synchrotron Light Source and is supported by the U.S. Department of Energy under Contract DE-FG05-89ER45384.

Fig.1 First derivative of the total DWFs of the nearest-neighbor Pb-O bond in PbTiO₃: a step at 200K indicates a second order phase transition and a peak at 100K indicates a first order phase transition.

M. JOO, A. EDWARDS, AND D. SAYERS (N.C. State U.)

The Curie temperature of solid solutions of $(\text{Sr,Pb})\text{TiO}_3$ was previously reported to vary as a function of mixture concentration[1], while the Curie temperature of pure SrTiO_3 has been reported to be 45K[2]. In this current work, the solid solution of $\text{Sr}_{0.8}\text{Pb}_{0.2}\text{TiO}_3$ has been studied by temperature-dependent EXAFS in order to investigate the phase transition phenomena of this alloy. The sample was made from SrTiO_3 and PbTiO_3 powders. These were mixed together in the ratio of molar weight 80%:20%, pressed into a pellet and then calcined at 1100°C. X-ray absorption spectra on the Sr-K-edge (16105 eV) were measured in the temperature range between 25K and 250K using a Displex refrigerator. The relative Debye-Waller factors (DWFs) of the first shell of $\text{Sr}_{0.8}\text{Pb}_{0.2}\text{TiO}_3$ were calculated using the exponential fitting method. In Fig.1, the changes in the slope of the DWFs of SrTiO₃ are obvious at around 100K and 40K as previously reported[3]. However, the only evident change in the plot for $\text{Sr}_{0.8}\text{Pb}_{0.2}\text{TiO}_3$ occurs at 200K, which is in good agreement with the prediction of Nomura and Sawada[1]. Further measurements and detailed analysis are currently in progress.

[1] S. Nomura and S. Sawada, J. Phys. Soc. Japan **10** (1955) 108

[2] H. Weaver, J. Phys. Chem. Solids **11** (1959) 274

[3] K. Müller and W. Berlinger, Phys. Rev. Lett. **26** (1971) 13

Acknowledgement:

This work was performed at X-11A Beamline at the National Synchrotron Light Source and is supported by the U.S. Department of Energy under Contract DE-FG05-89ER45384.

Fig.1 Relative experimental DWFs of the first shell of titanate compounds; SrTiO_3 (solid-line) and $\text{Sr}_{0.8}\text{Pb}_{0.2}\text{TiO}_3$ (dotted-line).

M. JOO, A.M. EDWARDS AND D.E. SAYERS (N.C. State U.)

Temperature-dependent XAFS above the Sr K edge (16105 eV) in SrTiO_3 has been measured in the temperature range between 25K and 300K using a Displex refrigerator. The relative Debye-Waller factors (DWFs) of the nearest-neighbor Sr-O bond in SrTiO_3 were calculated by inverse Fourier filtering and then exponential fitting. Using a new analysis technique, the σ^2 -derivative method[1], information about not only the location but also the nature of the phase transitions can be obtained without necessitating separation of the structural and thermal DWFs. We can confirm at least two phase transitions below the room temperature in SrTiO_3 using this new analysis technique. A step at about 105K and a peak at about 65K in the first derivative of the total DWFs (Fig.1) indicate that these phase transitions are of second order[2] and of first order[3] respectively.

[1] M. Joo, A.M. Edwards and D.E. Sayers (To be published).

[2] K.A. Müller and W. Berlinger, Phys. Rev. Lett. **26**, 13 (1971).

[3] F.W. Lytle, J. Appl. Phys. **35**, 546 (1964).

Acknowledgement:

This work was performed at X-11A Beamline at the National Synchrotron Light Source and is supported by the U.S. Department of Energy under Contract DE-FG05-89ER45384.

Fig.1 First derivative of the total DWFs of the nearest-neighbor Sr-O bond in SrTiO_3 : a step at 105K indicates a second order phase transition and a peak at 65K indicates a first order phase transition.

Measurement of Internal Strains Using Embedded Micromarkers

E.H. Jordan, S.C.U. Ochi, D.M. Pease, and J.I. Budnick

Mechanical Engineering, Physics Department, and Institute of Materials Science, University of Connecticut.

We have developed a new experimental method which will allow measurement of strains in the interior of composite materials and other real structural material. The method involves the use of the synchrotron x-ray source to record images of positions of micromarkers (10-40 microns) embedded in a real structural medium before and after deformation of the medium. High contrast radiographs are recorded by using monochromatic x-rays tuned to the absorption edge of the marker material. The images thus recorded consists of white dots representing the images of the micromarkers which absorb most of the x-rays and a dark background representing the transmitted x-rays throughout the relatively transparent matrix.

To achieve the desired strain measurement it is necessary to measure changes in particle spacing to better than 1 part in 10,000. The radiographs are first optically enlarged and the particles' spacings measured. A method of measuring particle spacings has been developed that uses high resolution x-z stages combined with centering detection devices (diodes). An example of a measurement in uniformly strained thin (1.25mm) aluminum plate is shown in Fig. 1. Onto the surface of the experimentally determined strain was compared to that measured by an extensometer and reasonable agreement is observed. The system is now to be used in studying composite materials' behavior.

Strain Versus Load Direction

Fig. 1. Strain vs. load direction of strained aluminum plate. The x marks denote strain determined from position shifts of internal micromarkers.

This work was supported by the U.S. Department of Energy, the UConn School of Engineering, and the UConn Research Foundation.

XANES OF CONVERSION COATINGS ON ALUMINUM ALLOYS

M. Kendig (Rockwell International Science Center), A. Davenport (BNL), H. Isaacs (BNL)

Corrosion protection of light metal alloys such as those of Al includes the application of a chromate conversion coating produced by the reaction of the alloy with an acidic solution containing dichromate and fluoride ions. The Al reduces the hexavalent Cr to form a highly protective hydrated Cr_2O_3 film.

Concern over the environmental hazards presented by Cr(VI) demand the replacement of the chromate conversion coatings by more environmentally benign processes. There are no entirely suitable replacements for chromate conversion coatings. To replace the hexavalent Cr process, it is important to understand the corrosion-inhibiting mechanism of chromate conversion coatings. It has been hypothesized that the Cr_2O_3 coating retains a certain quantity of unreacted Cr(VI) which serves as a source of dichromate to repair breaks and cracks in the film. We have examined this hypothesis by determining to what extent Cr(VI) remains in chromate conversion-coated Al.

XANES analysis of the chemical speciation of the conversion coatings has been chosen since X-ray photoelectron spectroscopy (XPS) presents some drawbacks to a precise evaluation of the ratio of Cr(VI) to the total Cr due to a photodecomposition reaction of the Cr(VI) occurs in the UHV environment as most recently characterized by Schrott and coworkers.¹

Fig. 1 shows the XANES for conversion coatings on Al-2024-T3 that were processed for the indicated times in a conversion coating bath. A pre-edge peak at 4 eV uniquely characterizes the tetrahedral coordination typical of the +6 oxidation state. For all process times, the pre-edge peak appears. Use of standards allowed the generation of a calibration curve to quantify the ratio of tetrahedral Cr(VI) to total Cr in the films. As shown in Fig. 3, the quantity of Cr(VI) in the conversion coatings varies from 10-22% of the total Cr depending on the processing time. For processing times greater than 3 min the % Cr(VI) rises to about 22% and becomes invariant with processing time. Hence, unreacted hexavalent Cr represents a significant component of the conversion coating and most likely provides a source of Cr(VI) for dynamic repair of the coating thereby giving the chromate coating a unique corrosion prevention property.

Fig. 1 XANES for chromate conversion coated Al-2024-T3.

Fig. 2 % Cr(VI) as a function of process time.

1. A. G. Schrott, G. S. Frankel, A. J. Davenport, H. S. Isaacs, C. V. Jahnke and M. Rusak, *Surf. Sci.*, **250**, 139 (1991)

This work supported by Rockwell International IR&D, and the U.S. Dept. of Energy Office of Basic Energy Sciences under contract DE-AC02-76CH00015.

A TIME-RESOLVED DIFFRACTION STUDY OF THE Ta-C COMBUSTION SYSTEM

X-11A

E.M. Larson, J.B. Holt, J. Wong, P.A. Waide, B. Rupp and L. Terminello
Lawrence Livermore National Laboratory, Univ. of California, Livermore, CA 94551

The real time chemical changes in the formation of TaC and Ta₂C by combustion synthesis have been monitored by time-resolved x-ray diffraction using synchrotron radiation as the x-ray source. These reactions have been followed with a time resolution of as little as 50 milliseconds, which is a factor of 10 better than the time resolution of conventional x-ray diffraction. The adiabatic temperatures (i.e. assuming no heat loss) of these reactions are: (i) Ta + C → TaC and (ii) Ta₂C + C → Ta₂C₂ are calculated from known thermodynamic data to be 2452°C and 2378°C respectively. These maximum reaction temperatures are well below the melting points of tantalum (3020°C), carbon (~3800°C), the corresponding products TaC (~3825°C), Ta₂C (~3300°C) and eutectic temperatures (2825°C & 3375°C) known in the binary Ta-C phase diagram. Hence there is no formation of a liquid phase and the syntheses are truly solid combustions. The chemical kinetics of these reactions are deduced by monitoring the disappearance of reactant and the appearance of product Bragg diffraction peaks. Though TaC is most stable, the results for I show the concurrent formation of the subcarbide (Ta₂C). The subcarbide peak persists for a couple of seconds, then decreases in intensity as TaC attains its maximum concentration. In case II, the reaction proceeds directly to the product with no discernable intermediate Ta-C phase within a 50 ms time frame. A simple mechanism is supported for the formation of TaC by carbon diffusion into the Ta bcc lattice resulting in a fcc TaC cell with edge along the face diagonal of the original Ta lattice and an elongation along the 'c' axis.

COMPARISON BETWEEN THORIUM AND CERIUM DOPING AND STUDY OF THE ROLE OF REDUCTION FOR n-TYPE SUPERCONDUCTOR RELATED SYSTEMS

X11A

G. Liang, Yang Guo, D. Badresingh (Sam Houston State U.), M. Croft, J. Chen, J. Peng (Rutgers U.), Renato F. Jardim*, E. A. Early, M. B. Maple (U. C. at San Diego) Beom-hoon O, J. T. Markert (U. of Texas at Austin), M. Nagoshi, and T. Suzuki (NKK Corp., Japan)

Sample variation due to different preparation conditions can have serious effects on the shape of Cu K-edge spectra and thus can possibly obscure the interpretation of the spectral features. In order to clarify these effects, we have carried out an extensive x-ray absorption study for R_{2-x}A_xCuO_{4-s} (with A=Th, Ce, R=Nd, Pr, Sm, Eu, Gd and 0 ≤ x ≤ 0.18) compounds prepared under different conditions, such as: annealed in oxygen or air and reduced in helium, argon or vacuum. We studied the less-studied Th-doped samples in order to compare the electron-doping process between A=Ce (potentially less than tetravalent) and A=Th (tetravalent) compound series.

We have observed the following by comparing the spectra between the helium reduced samples and the corresponding oxygen annealed ones. 1) For Ce-doped series with R=Gd, Eu, 0 ≤ x ≤ 0.18 and R=Pr, Nd, x ≥ 0.1, the Cu K-edge spectra for samples reduced in helium atmosphere are identical to those of corresponding oxygen oxidized samples; for other concentrations and for R=Sm, spectral feature changes induced by helium reduction were observed. 2) These feature changes induced by helium reduction also occur for Th-doped compound series with R=Nd, Pr for all x values and they appear larger than that for the corresponding Ce-doped compounds. 3) Detailed analyses show that the spectral feature changes due to reduction are quite different from those produced solely by electron-doping. We found that the Cu K-edge spectra for selected samples reduced by electron-doping (i. e. by annealing samples in helium, argon, or vacuum) are basically identical. The above observations indicate two things: first, the conclusion (which we made previously^[1,2]) that the doped electrons are injected into Cu-sites is not altered; second, the reduction appears not directly associated with electron-doping into the Cu-3d orbital.

The Cu K-edge spectra show that, if one excludes the spectrum change induced by reduction, the fractions of electronic charge doped from each Ce and Th atom into the Cu-sites are close to each other. This observation indicate that the electron-doping processes are very similar in both the Ce and Th doped compound systems.

[1] J. M. Tranquada et al., Nature (London) 337,720(1989).

[2] G. Liang et al., Phys. Rev. B40, 2646(1989).

Acknowledgments: This work was supported by Research Corporation, the Faculty Research Funds of Sam Houston State University, the Robert A. Welch Foundation Grant No. F-1191. *FAPESP (Brazil) fellowship, and U. S. Department of Energy under Grant No. DE-FG03-86ER45230.

G. Liang, Wei Xu (Sam Houston State U.), M. Croft, J. Chen, J. Peng (Rutgers U.),
Boom-hoan O., and J. T. Markert (U. of Texas at Austin)

We have measured polarization x-ray-absorption spectra at the Cu K-edge from $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$ single crystals with $x=0.15$ and 0.20 . The objective is 1) to examine if the proposed electron-doping and band-shifting mechanism^[1] occurs the same way in the Ce-doped R_2CuO_4 with $\text{R}=\text{Nd}$ as in the reported $\text{R}=\text{Pr}$ system,^[1] and 2) to extend the polarized Cu-K-edge study from low- x (doping-level) region to the high- x region where the superconducting transition occurs.

Fig. 1 shows that the features (labeled by letters A through D) of the Cu K-edge spectrum of the polycrystalline powder sample $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$ can be well interpreted from the combination of the $\theta \approx 15^\circ$ (or nearly $E \parallel c$) and $\theta \approx 80^\circ$ (or nearly $E \perp c$) polarization spectra according to the assigned well and poor screened $1s \rightarrow 4p_x$ (for A and B) and $1s \rightarrow 4p_y$ (for C and D) transitions.^[2] The feature A' located at -1.2 eV for both the powder spectra and the single crystal $E \parallel c$ spectra is lower than the $4p_x$ features (not shown here) of both Cu_2O (at 0.0 eV) and Cu metal (at -0.8 eV). In Fig. 2 we show the polarized Cu K-edge spectra for $\text{Nd}_{2-x}\text{Ce}_x\text{CuO}_4$ with $x = 0.0$ spectra taken from ref. [1] for comparison. Combined with results in ref. [1], we found that the intensity of feature A' roughly increases linearly with x up to $x = 0.2$. The $E \perp c$ spectra in Fig. 2 shows that there appears an about 0.9 eV almost rigid edge-shift between the spectra of the undoped and $x = 0.2$ samples if the weak intensity at -1.2 eV (feature A') could be removed. This intensity could be associated with θ being set closer to 80° (rather than 90°) to avoid the Bragg reflection features in the near-edge region. Our results support that, up to the T^* -phase stability limit $x = 0.2$, the Ce doping donates electrons continuously to the Cu $3d$ states to form Cu^{2+} and also simultaneously shifts the unoccupied $4p$ band downward.

[1] Zhengquan Tan et al., Phys. Rev. B 42, 1037 (1990). [2] J. M. Tranquada et al., Nature (London) 337, 720(1989).

*This work is supported by Research Corporation, the Faculty Research Funds of Sam Houston State University, and the Robert A. Welch Foundation Grant No. F-1191.

X-RAY ABSORPTION STUDY OF ELECTRON-DOPED HIGH- T_c
SUPERCONDUCTOR-RELATED COMPOUNDS $\text{R}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$

G. Liang, Yang Guo, D. Badreisingh, Wei Xu (Sam Houston State U.), M. Croft, J. Chen, J. Peng (Rutgers U.), Boom-hoan O., and J. T. Markert (U. of Texas at Austin)

We have carried out systematic x-ray absorption measurements at the Cu K-edge and Ce $L_{2,3}$ -edge on electron-doped high- T_c superconductors (HTS) $\text{R}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$ with $\text{R}=\text{Nd}$, Pr , Sm , Eu , Gd (not HTS) and $0 \leq x \leq 0.18$. This study was intended to provide electronic structure information in understanding the decrease of T_c with R varying from light to heavier rare-earth elements.

We have observed from Cu K-edge data that the intensity of $1s \rightarrow 4p_x$ [$1s$] transition in a $\text{Cu-}3d^{10}$ ground-state configuration (labeled by letter A in Fig. 1) increases quite linearly with the increase of x in all of the $\text{R}_{2-x}\text{Ce}_x\text{CuO}_{4-y}$ series. This intensity increase is accompanied by two effects: 1) a decrease of the intensities of the $\text{Cu-}3d^9$ ground-state configuration $1s \rightarrow 4p_x$ (labeled by A, B) and $1s \rightarrow 4p_y$ (labeled by C, D) transitions; and 2) a downward edge-shift.^[3] A typical example is shown in Fig. 1 for $\text{R}=\text{Eu}$ series. The measured Ce- L_3 spectra show that Ce is in a valent state similar to that in CeO_2 . The above observations indicate that the electrons doped by Ce atoms are injected into the Cu- $3d$ sites in the same manner in all of the series with R varying from Nd to Gd, irrespective of whether the series exhibits superconductivity or not. Further more, we found that for the same doping level x , there is an about 0.5 eV upward almost rigid edge-shift when the rare-earth elements R vary from $\text{Pr} \rightarrow \text{Nd} \rightarrow \text{Sm} \rightarrow \text{Eu} \rightarrow \text{Gd}$. Fig. 2 shows a typical example of such rigid shift for $x = 0.15$. We interpret this trend as due to a shift of the in-plane $4p$ -states to a higher energy upon the a-axis contraction of the compounds when R varies from Pr to Gd. This band shift might be associated with the annihilation of superconductivity in the electron-doped HTS systems just opposite to what occurs when the Ce substitutes for Nd or Pr (though both electron-doping and the accompanying a-axis expansion), where the in-plane $4p$ band is shifted downward to induce superconductivity in the $\text{R}=\text{Nd}$ and Pr systems.^[3]

[1] J. M. Tranquada et al., Nature (London) 337, 720(1989). [2] G. Liang et al., Phys. Rev. B 40, 2646(1989). [3] Z. Tan et al., Phys. Rev. B 42, 1037(1990).

*This work was supported by Research Corporation, the Faculty Research Funds of Sam Houston State University, and the Robert A. Welch Foundation Grant No. F-1191.

THERMAL VIBRATIONS AND MELTING FROM A LOCAL PERSPECTIVE

P. Livins, E. A. Stern, Z. Zhang
Dept. of Physics, University of Washington, Seattle, Washington 98195

We report a study of the thermal vibrations of lead as a function of temperature up to and through the melting transition. Analysis of X-ray absorption fine structure (XAFS) measurements, obtained on beam line X-11A at NSLS, provides information of short range correlations between the absorbing lead atom and its near neighbors. In applying the log-ratio method to the first shell, we obtain the first four moments of the radial distribution for the melting transition by fitting the cumulant expansion. The temperature dependence below the melting transition is summarized in table I.

By modeling the thermal vibrations with an anharmonic oscillator the above mentioned moments define a radial density function for the first shell. The same moments are also related to the constants defining the anharmonic potential

$$V(z) = \frac{1}{2}kz^2 - gz^3 - fz^4.$$

Here z represents the displacement from the minimum of V at R_0 . These force constants are summarized in table II.

Due to the thermal vibrations at high temperatures, the XAFS signal is only significant at small wave vectors, where the determination of the XAFS background has traditionally been thwarted. However, at high temperatures this low k data is all that is available for analysis. The XAFS for lead is fortunately dominated by the first shell, in which case careful alignment of the data with different temperatures yields the XAFS zero crossings at low k where the phase changes with temperature are insignificant. The energies of the zero crossings then provide constraints for a smooth background fit, which determines the background down to about 3 eV above threshold, approximately 1.7 inverse Angstroms when noting the free electron Fermi wave-vector.

Perhaps the most interesting results of the analysis has been in the interpretation of what information the XAFS technique yields for the liquid state. XAFS continues to be observable on going through the solid to liquid phase transition. However, it should be interpreted to provide information only about the sharp features of the radial distribution. The meaning for the liquid state is that moments obtained, as in the above analysis for the solid, apply only to the vibrational motion between the hops occurring during diffusion in the liquid state. We find that in the liquid state the lead neighbors spend half their time vibrating as opposed to hopping.

Table I.

ΔR	$T(\text{\AA} \times 10^{-5})$	σ^2	$T(\text{\AA} \times 10^{-5})$	$c/T^2(\text{\AA}^2 \times 10^{-8})$	$b/T^3(\text{\AA}^3 \times 10^{-12})$
8.0 ± 3.4	9.9 ± 1.2	1.9 ± 2	6.6 ± 3.3		

Lowest: order temperature coefficients of the first four cumulants for Pb, obtained for temperatures up to the melting point. ΔR , σ^2 , c and b are the first four cumulants respectively.

Table II.

k	g	f
0.87 ± 0.1	0.28 ± 0.09	-0.15 ± 0.17

The harmonic force constant k with the cubic, g , and quartic, f , anharmonic constants obtained for Pb.

References:

1. E. A. Stern, P. Livins and Z. Zhang Phys. Rev. B 43 (in print) (1991).
- Acknowledgements: This research was funded by DOE contract number DE-FG06-84ER45163 We also thank the NSLS, supported under DOE contract number DE-AC0276CH00016.

X-11-A

XANES AND EXAFS STUDY OF MN DOPED CaF_2 DOSIMETRIC PHOSPHOR

A. N. Mansour and J. H. Barkyoub, NSW

Room temperature K -edge XANES and EXAFS studies were carried out on CaF_2 powdered samples that were doped from 0.1 to 5 atomic % with manganese. This material serves as a thermoluminescence radiation dosimeter and the nature of the local environment around the manganese luminescent center is important to the performance of the phosphor.

The Ca K -edge measurements were performed in transmission mode and the Mn K -edge measurements were done in fluorescence (dilute sample limit).

Comparison of the Ca K and Mn K $\chi(k)$ and Fourier transforms shows that the manganese is substituting for Ca on Ca^{2+} sites (Fig. 1). From the position of the Mn K edge it appears that the manganese ion is predominantly in the $2+$ state. These results are what would be expected from optical studies of the crystal-field split d states of the Mn ion.¹ EXAFS parameters derived from both fitting to MnF_2 as a model compound and from a theoretical curved-wave formalism² show a contraction of the first shell around the Mn^{2+} ion of 0.15 Å from the Ca-F distance of 2.365 Å.

Fig. 2 shows the Ca K -edge XANES and Mn K -edge XANES plotted on a common energy scale with $E = 0$ at the first pre-peak in the spectra. From a one-electron band structure of CaF_2 , one can assign the first pre-peak to a $1s$ - $3d$ transition and the higher shoulders to p -like states in the conduction band.³

1. P. J. Alonso and R. Alcala, J. Lumin. 22, 321 (1981).
2. J. J. Rehr, J. Mustre de Leon, S. I. Zabinsky, and R. C. Albers, J. Am. Chem. Soc. 113, 5136 (1991).
3. R. A. Heaton, C. C. Lin, Phys. Rev. B 22, 3629 (1980).

This work is supported by DOE under Contract Number DE-AS05-80ER10742.

CHARACTERIZATION OF A Cu/Al₂O₃ BURIED INTERFACE USING GLANCING-ANGLE X-RAY REFLECTIVITY AND XAFS STUDIES

R. A. Mayanovic, B. A. Bunker, C. Zhong, and A. Anagnostopoulos, Physics Dept., U. of Notre Dame.

Characterization and local structure studies of Cu/Al₂O₃ buried interfaces are important in relation to electronic studies of tunnel barrier junctions. These studies are also significant within the context of our polarization-dependent studies on atomic thermal-motion disorder at solid-solid interfaces. The sample was fabricated by evaporating ~4000Å of Cu on a 5.1cm Si wafer substrate inside an evacuated belljar at ~10⁻⁶ torr. The sample was then quickly transferred to another high vacuum chamber where ~700Å of Al₂O₃ was RF sputtered over the Cu film. Glancing-angle x-ray reflectivity and XAFS measurements were made on the sample at room temperature. The reflectivity measurements were made at 8.7 and 9.2KeV and the Cu K-edge XAFS measurements were made for several angles, ranging from 4-6mrad. The sample was studied under two orientations: Having the x-ray polarization vector ϵ either parallel or perpendicular to the Cu/Al₂O₃ interface. Fig. 1 shows x-ray reflectivity data measured at 8.7KeV with ϵ parallel to the interface. The data was reasonably well fitted using the Fresnel equation modified for interface roughness, as described by Heald et al.¹ Several models for the layered sample were tried during the fitting, including one having an intermediate CuAl₂O₃ alloyed layer sandwiched between the Cu and Al₂O₃ layers: Best results were obtained by assuming a sharp Cu/Al₂O₃ interface. This is verified by preliminary analysis of the polarization-dependent EXAFS data. The value used for the Al₂O₃ surface r.m.s. roughness was 42Å in the fit was 28Å and the value used for the Cu/Al₂O₃ interface r.m.s. roughness was 42Å. The slight discrepancy between the fit and the data, especially near 4mrad, can be explained in terms of long range distortions of the sample, which were unaccounted for during fitting, originating with the Si wafer substrate. Taking this into account, we estimate the true r.m.s. Cu/Al₂O₃ interface roughness to be between 30-40Å.

Fig. 1 X-ray reflectivity data (dots) measured at room temperature, 8.7KeV, and having ϵ parallel to the interface. The fit to the data (line) was obtained using a modified Fresnel equation.

I. S.M. Heald, H. Chen, and J.M. Tranquada, Phys. Rev. B 38, 1016 (1988)

Supported by Office of Naval Research under contract N00014-89-J-1108.

GLANCING-ANGLE XAFS STUDIES OF PREMELTING AT Cu/Al₂O₃ INTERFACIAL REGIONS

R. A. Mayanovic, B. A. Bunker, and N. Jisrawi, Physics Dept., U. of Notre Dame.

Studies on metals have shown that premelting at the surface occurs considerably below the bulk melting temperature, indicating that the wet surface of a solid may act as a nucleation source during melting. Our aim in this work is to study the atomic thermal-motion disorder and vacancy formation associated with the precursors to melting at metal-insulator interfaces using XAFS. Cu K-edge XAFS spectra were taken on Cu/Al₂O₃ multilayers, grown using RF sputtering on optically flat Si wafers having 20 layers each of ~100Å of Cu and ~100Å of Al₂O₃, using the glancing-angle of incidence technique in the fluorescence mode. Measurements were taken at several angles of incidence, from ~3 to 13 mrad, at room temperature and 420 ± 2°C. The multilayers were studied under two orientations: the x-ray polarization vector parallel (P orientation) and perpendicular (S orientation) to the Cu/Al₂O₃ interface. Analysis of the polarization-dependent EXAFS data reveals a directional anisotropy in the EXAFS Debye-Waller factor, occurring both at room temperature and in a much more pronounced fashion at 420 ± 2°C, indicating that the amplitude of relative Cu atom in-plane motion is larger than for out-of-plane motion at the interface (see Fig. 1). As shown in Fig. 2, we have obtained preliminary results for the incident angle dependence of the relative in-plane EXAFS Debye-Waller factor, measured at 420 ± 2°C, indicating a steady decrease of vibrational amplitudes with depth. This, along with an estimate for the average in-plane vibrational amplitude at that temperature of approximately 3-4% of the nearest-neighbor distance for Cu atoms down to a depth of about 30Å from the interface, indicates that we are close to the premelting regime. The implication of this study is that the interface very likely undergoes premelting via a shearing mechanism rather than the recoil mechanism used to explain premelting at a free surface, for which the directional anisotropy in atomic vibrational amplitudes is reversed. The shearing model may ultimately help to explain superheating of encapsulated solids.

Fig. 1 Normalized EXAFS oscillations for the Cu K-edge, measured at a Cu/Al₂O₃ interface, at 3.9 mrad incidence angle and 420 ± 2°C, in S (solid line) and P (dashed line) orientations.

Fig. 2 The Cu atom in-plane EXAFS Debye

-Waller factor (P orient.), as a function of incident angle, measured relative to the out-of-plane component (S orient.) obtained at 3.9 mrad.

I. J. W. M. Frenken and J. F. van der Veen, Phys. Rev. Lett. 54, 134 (1985); P. H. Fuoss, L. J. Norton, and S. Brennan, Phys. Rev. Lett. 60, 2046 (1988).

Supported by Office of Naval Research under contract N00014-89-J-1108.

EXAFS INVESTIGATIONS OF Pt-Au/ γ -Al₂O₃ CATALYSTS

K.I. Pandya (BNL), S.M. Heald (BNL), J. Fraissard (Paris) and L. Petrakis (BNL)

X-ray absorption measurements were carried out at the Pt-L₃ and Au-L₃ edges for Pt_{1-x}Au_x/ γ -Al₂O₃ catalysts with $0 \leq x \leq 0.8$. The samples were reduced at 673 K and the EXAFS measurements were carried out at about 77 K. Figure 1 shows the k^3 -weighted $\chi(k)$ and the corresponding radial structure function for the Pt_{0.8}Au_{0.2}/ γ -Al₂O₃ sample. As the Pt and Au L₃ edges are separated by 350 eV only, the Pt EXAFS are superimposed on the Au EXAFS. Hence the analysis of Au EXAFS is complicated by the presence of Pt EXAFS. A modelling and spectral subtraction approach will be taken to isolate the Au EXAFS. The EXAFS results will be used to derive a model for these catalysts.

Fig. 1. (a) Pt EXAFS and (b) Radial Structure Function for the Pt_{0.8}Au_{0.2}/ γ -Al₂O₃ sample

This work was supported by the U.S. Dept. of Energy, Division of Mat. Science under contract Nos. DE-AC02-76CH00016 and DE-AC05-80-ER10742.

D.M. Pease

Physics Department and Institute of Materials Science, University of Connecticut

There are presently conflicting experimental results for L₃ and L₂ core hole widths and 3d transition metals, (TM) as obtained from x-ray photoemission spectroscopy (XPS) (1,2). In addition, there is disagreement between experimental and theoretical widths (3) which may result from "non lifetime" solid state contributions to the experimental widths (1). The 2p $\rightarrow$ 3s soft x-ray emission spectra are not effected by core hole-valence electron exchange effects and the L₃/L₂ ratio for such spectra therefore is determined by the Auger and Coster-Kronig widths of the L₃ and L₂ holes. An expression is derived for this ratio, in terms of true lifetime widths, and the experimental spectrum (4) is compared to a superposition of Lorentzian functions with L₃/L₂ weights determined according to three different models: (a) theoretical widths only (which do not contain the Coster-Kronig contribution) (b) experimental widths only, and (c) theoretical L₃ widths but experimental difference between L₂ and L₃ widths used to obtain the L₂ width. For elements Cu through Fe, it is found that good agreement with experiment is obtained if the data of reference (1) is used in conjunction with model (c).

Iron 2p $\rightarrow$ 3s emission spectra. Solid line represents smoothed data and dotted lines are theoretical curves, based on models a, b, or c as discussed above. Model C gives good agreement for other TM's as well.

- (1) J.C. Fuggle and S.F. Alvarado, Phys. Rev. A22, 1615 (1980).
- (2) R. Nyholm, N. Martensson, A. Lebugle, and V. Axelsson, J. Phys. F. Metal Physics, 1727 (1981).
- (3) B. Crasemann, Proc. 2nd Workshop on High Energy Ion Atom Collisions, Debrecen, 1984; M.H. Chen, B. Crasemann, and H. Mark, Phys. Rev. A24, 1977 (1981).
- (4) S. Hanzely, Ph.D. dissertation, New Mexico State University, 1968.

This work is supported by the U.S. Department of Energy.

The Structural Characterization of Metal Oxide Clusters in Zeolite-Y and Zeolite-L Hosts, Produced by the Photo-Oxidation of Group(VI) Metal Hexacarbonyls, Followed by Various Degrees of Subsequent Reduction

R.A. Prokopowicz, A. Malck, M. Sieck, A. Stein, G.A. Ozin, (U of Toronto)

Group(VI) metal hexacarbonyls have been used as precursors in the synthesis of highly organized assemblies of molecular-dimension metal oxide clusters, encapsulated within the void structure of various zeolite frameworks. To date, the most extensively studied specimens include various W and Mo oxides in zeolite Y. In the synthesis of these novel materials, volatile $M(CO)_6$ is sublimed into the zeolite hosts where it becomes irreversibly anchored, exclusively in the large α -cages, at a saturation maximum of two molecules per cage (sixteen per unit cell). The carbonyl is next converted in an O_2 atmosphere to the M^{4+} oxide by photo-oxidation, and may be subsequently thermally reduced in vacuum to yield MO_{x-4} clusters (where $0 \leq x \leq 1$). The reduction is reversible by heating in an O_2 atmosphere at $300^\circ C$. Following photo-oxidation, half of the α -cage void volume in the host is freed so that subsequent precursor (saturation level) impregnations / photo-oxidations can be carried out; the stepwise loading proceeds as 16, 24, 28, 30, ..., 32 per unit cell.

The zeolite-encapsulated hexacarbonyl metal(0) precursors were the first samples examined by EXAFS. It was determined that the structural integrity of the free-state precursors was maintained upon anchoring in the framework. Coordination numbers, M-C and M-(C-O) distances agreed with those obtained crystallographically for bulk $M(CO)_6$. Evidence for anchoring of the precursor via the oxygen end of two trans-carbonyl ligands to α -cage extraframework alkali metal cations was also obtained in this study.

Results to date indicate that in Na_xY , WO_3 and $WO_{2.5}$ moieties are formed exclusively as dimers (i.e. W_2O_4 and W_2O_5) which are each anchored to two of the four cations located in an α -cage. This structure is valid at loading levels of 16 molecules per unit cell (one per α -cage), 28 and up to a fully loaded 32 (two per α -cage) for $WO_{2.5}$. For $WO_{2.5}$ dimers are formed at the 16 loading level, but these assume a tetrameric structure (W is coordinated to three other W) at 32 per unit cell. The EXAFS results have indicated that W atoms in the dimers are linked by long bridging O bonds, either one in the $WO_{2.5}$ dimer or two in the WO_3 dimer, and have two short terminal O bonds which interact with the extraframework cations. At loading levels of 16, 28 and 32, the WO_3 species are consistently monomeric; O coordination numbers of 4 (equal bonds) are explained in terms of a monomeric WO_3 moiety interacting with two framework O atoms (which are possibly part of an α -cage oxygen 4-ring).

A group of rubidium Y samples at the 16 per unit cell loading level has also been studied by EXAFS. The Rb_xY analogs of the 16 /unit cell materials were determined to differ structurally, in that both the WO_3 and $WO_{2.5}$ moieties were monomeric. The $16WO_{2.5}Rb_xY$ sample did however, form a dimeric structure; the unexpected result for WO_3 is attributed to the much larger size of the Rb^{+} cations causing the dimer structure to be sterically unfavorable within the smaller space between them. An examination of backscattering due to Rb (from the W-L₂₃ EXAFS) and also the W backscattering contribution in the Rb-K edge EXAFS, indicated that the WO_3 monomer is anchored in close proximity to Rb cations. Anchoring of the WO_3 monomers was, as in the Na^+ case, determined to involve two additional framework O atoms.

The Mo oxide samples which have been examined (these are at a loading level of 16 in NaY) show that only monomeric MoO_3 and $MoO_{2.5}$ moieties are formed. These are structurally similar, in terms of framework (likely 4-ring) anchoring, to the WO_3 monomer analog. This result is consistent with the observation that a stable Mo_2O_7 cluster could not be synthesized in this system.

Funding for this project was provided by the Natural Sciences and Engineering Research Council of Canada

SPONTANEOUS ORDERING EXAFS STUDY OF $Ga_{0.5}In_{0.5}P$ THIN FILMS DEPOSITED ON $GaAs$

Anne Tabor-Morris, Kenneth Kemner, Robert Mayanovic, and Bruce Bunker, University of Notre Dame

Interest in spontaneously ordered alloys deposited on thin films has been sparked due to the observation of ordering and the accompanying anomalous lowering in band gap. Electron and x-ray diffraction have indicated the existence of spontaneous ordering, but controversy still exists as to the nature of this ordering, especially in terms of short-range structures and interatomic distances.¹ EXAFS is suited to the study of the positions and coordination numbers of first, second, and third nearest neighbors around Ga in $Ga_{0.5}In_{0.5}P$, in these samples deposited 2 degrees off of the (100) axis of the $GaAs$ substrate by metal organic chemical vapor deposition. Using the inherent polarization of synchrotron x-ray radiation, we investigate the positions and coordination numbers of the nearest neighbors to indicate the presence and type of structural ordering, specifically probing to see if the type of atom in the second shell is preferentially ordered along axes of the $Ga_{0.5}In_{0.5}P$ single crystal overlayer.

These measurements are made by utilizing a liquid nitrogen cooled total electron yield detector. Data indicates that there is a marked difference in the EXAFS signal amplitude due to the second and third nearest neighbors depending upon the orientation of the sample, as displayed in Figure 1. The difference in the k-dependence of the EXAFS oscillations due to the second and third shells can be attributed to Ga preferentially ordering along the (1,1,0) direction and in along the (1,-1,0) direction, consistent with the evidence of ordering found in other studies. More detailed analysis is currently in progress.

Figure 1. Total Electron Yield Data: Filtered data for second and third nearest neighbors. Solid line: 0 degrees (x-ray beam polarization along (1,1,0) axis). Dashed line: 90 degrees (sample rotated, polarization along (1,-1,0) axis).

1. Kurtz, S., Olsen, J., Kibbler, A., Appl. Phys. Lett., **54** (8) 718 (1989).

* Supported by the Office Of Naval Research under contract N00014-89-J-1108.

† Samples provided by K. Bertness, Solar Energy Research Institute, Golden, CO.

Zhengquan Tan and S. M. Heald (Brookhaven Nat'l Lab); S.-W. Cheong and A. S. Cooper (AT&T Bell Labs); J. I. Budnick (Univ. of Connecticut).

Charge carrier doping strongly affects the magnetic, transport and superconducting properties of the T-, T*- and T*-phase R_2CuO_4 (R=rare earths) compounds. The nature of doping and the doping level depend not only on the heterovalent cation substitution and oxygen stoichiometry but also on the rare earth valence. We have studied the valence of various rare earths in the R_2CuO_4 materials using rare earth L_3 x-ray absorption spectra. Similar to CeO_2 , Ce is trivalent in both T-phase $Nd_{2-x}Ce_xCuO_4$ and T*-phase $Pr_{2-x-y}Ce_ySr_yCuO_4$ (Fig. 1). Pr is trivalent in the T and T* structures (Fig. 2). In T*-phase $Pr_{1.6}Sr_{0.4}CuO_4$ and $Pr_{2-x-y}Ce_ySr_yCuO_4$, Pr has an average valence greater than 3+ (Fig. 2), very likely arising from an inhomogeneous mixture of trivalent and tetravalent ions,¹ with the latter occupying the eightfold-oxygen coordinated position. Tb in T*-phase $La_{0.9}Tb_{0.1}Sr_{0.9}CuO_4$ compounds behaviors similarly to Pr. Our results suggest that Pr and Tb in the T* compounds compensate the attempted hole doping by Sr, leading to the observed nonmetallic conductivity and lack of superconductivity. Other lanthanide rare earths are trivalent in the T, T* and T* phases. We refer to Ref. 1 for details.

1. Z. Tan *et al.*, Phys. Rev. B (Oct 1 issue, 1991); Z. Tan *et al.*, Phys. Rev. B (submitted).

*We acknowledge the support from the U. S. Department of Energy under Contract Nos. DE-AC02-76CH00016 and DE-AC05-80-ER10742.

Fig. 1 (above) Ce L_3 x-ray absorption spectra.

Fig. 2 (right) Pr L_3 (L_2) x-ray absorption spectra.

Zhengquan Tan and S. M. Heald (BNL), S.-W. Cheong, H. Hwang, A. S. Cooper and L. W. Rupp, Jr. (AT&T Bell Labs)

The occurrence of superconductivity in the $La_2CaCu_2O_6$ type structure appears to depend critically on the lanthanum and alkali-earth composition and requires high-pressure oxygen synthesis. In some cases, no superconductivity is observed even though the samples exhibit metallic conductivity and appear to have sufficient carrier concentration. In an effort to understand the local structure and their relationship with physical properties in this type of material, we have undertaken an x-ray-absorption study of the cation local environment in $LaSrCu_2O_6$ and $LaSrCu_2O_{6-y}$ with R=Pr, Nd, Sm and Gd.

In $LaSrCu_2O_{6-y}$, as the rare earth varies from Pr to Gd, both the a- and c-axis lattice constants decrease and the oxygen content decreases from 6.2 for R=Pr to slightly less than 6.0 for R=Sm and Gd. Our La and rare earth XANES data (Fig. 1) and La L_3 - and Sr K -edge (Fig. 2) EXAFS data indicate that from Pr to Gd the rare earth ions increasingly prefer to occupy the site between the two copper-oxygen planes (M_1 site) and increasingly more Sr(La) ions populate the M_2 site in the rock-salt layer. The Sr and La population at the M_1 site clearly correlates with the oxygen content when the type of rare earth ion varies. The Cu K -edge EXAFS data appear to suggest that some Cu atoms are coordinated with six oxygen atoms in addition to the fivefold pyramidal coordination. Our lanthanide L_3 - and Cu K -edge data lead us to believe that the oxygen in excess of 6.0 is present near the M_1 -site Sr(La) ions between the two CuO_2 planes. The detailed oxygen coordination configuration around these Sr(La) ions remains to be determined.

Fig.1 Lanthanide L_3 XANES.

Fig. 2 Fourier transform magnitude of Sr EXAFS.

* We acknowledge the support from the U. S. Department of Energy, Office of Basic Energy Sciences, under Contract Nos. DE-AC02-76CH00016 and DE-AC05-80-ER10742.

EXAFS STUDY OF FE CORE STRUCTURE IN RECONSTITUTED FERRITINS

Qiwu Wang*, Ann Edwards*, Dale Sayers* and Elizabeth Thell#
 Department of Physics* and Biochemistry#, North Carolina State University,
 Raleigh, NC, 27695

Ferritin, a unique protein, serves as a reserve of iron for iron-proteins as well as providing a safe site for the detoxification of excess iron. Its structure involves a protein coat around a core of hydrous, ferric oxide. The process of core formation can be investigated by directly examining the local environment of the Fe in reconstituted ferritin. In a previous study of the very early stage of the process, 10 Fe atoms per protein molecule (10 Fe/protein) were added to the apoferritin and followed by oxidation. The EXAFS results showed that a cluster of Fe was bound to carboxylate-like ligands with a near-neighbor environment of O at 1.99 Å, O/C at 2.79 Å and Fe at around 3.50 Å. However, in natural horse spleen ferritin the attached Fe oxide aggregate is of the order of a few thousand Fe atoms. In order to obtain a better understanding of the core formation at higher Fe concentrations, this current study involves reconstituted ferritins with 50 and 100 Fe/protein.

For Fe50, both the Fe-O/C model for the second shell and Fe-O+Fe-O/C model for the first and second shells together yield not only reasonable parameters, but also the smallest fitting errors. Modelling the second and third shells using the Fe-O/C+Fe-Fe model gave the best fit, indicating once again that an Fe-O/C interaction is the best description for the second shell. The fit also yielded a third shell Fe-Fe coordination of 4-5 Fe atoms at a distance of 3.60 Å. Thus, the environment is still like that for 10 Fe/protein.

The Fourier transform of the EXAFS data for Fe100 has much higher second and third peaks than Fe50 suggesting a larger coordination number and/or stronger back-scattering from near-neighbors. In contrast to the Fe50 case, for the second shell of the Fe100 the Fe-Fe model gives nearly a four times smaller fitting error than the Fe-O model. The suitability of this model was confirmed by the use of the Fe-O+Fe-Fe model for the first two shells and the Fe-Fe models for the second and third shells between 2.08 and 3.90 Å.

When 100 Fe atoms are added to the protein coat of ferritin, the core has properties more like the bulk than like the cluster formed with smaller numbers (10 or 50) of Fe atoms. The bulk properties of the ferritin iron core therefore appear to require at least 100 Fe atoms.

Fig. 1 Fitting $k^2\chi(k)$ of Fe100 with the Fe-O(---), Fe-O+Fe-Fe(- · -) and Fe-Fe(····) models. The experimental data is shown by the unsmoothed solid line.

This work is supported by the NIH under contract DK20251.

EXAFS STUDY OF LOCAL STRUCTURE OF FERROELECTRIC $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ ALLOYS

Z. Wang, R. Maruyama, B. Bunker, U. Debska and J. Furdyna, U. of Notre Dame.

EXAFS has been used to study the local structure of the disordered $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ alloys. Based on the temperature variation of the dielectric constant and from the hysteresis loop at elevated temperatures, R. Weil et al.¹ reported a ferroelectric transition in this alloy system for $x=0.04$ to 0.40. From this work, however, it is unclear what is happening *structurally* through this transition. Considering the large atomic size mismatch between Zn and Cd atoms ($Zn^{2+}=0.83\text{Å}$, $Cd^{2+}=1.03\text{Å}$), it is reasonable that Zn ions may be off-center in the cage of surrounding Te ions, and could move in a multi-well potential. If the ions are indeed off-center, the alloy may undergo an order-disorder ferroelectric phase transition (PPT) as found in $\text{Pb}_{1-x}\text{Ge}_x\text{Te}^2$ and $\text{PbS}_{1-x}\text{Te}_x^3$.

Single crystals of $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ for $x=0.10, 0.20$ and two standards ZnTe and CdTe were grown by Bridgman method, and then were ground to 400 mesh powders. We have performed XAFS measurements on these samples at Zn K-edge in fluorescence mode and Cd K-edge in transmission mode, at temperature ranging from 300K to 500K and 300K for two edges respectively. Figure 1 and 2 show the k -weighted first shell χ data for $x=0.10$ sample at two different edges at 300K. Obviously there is almost no difference in the first shell Cd-Te bonds between the alloy and pure CdTe from figure 1. But as we can see from figure 2, there is a significant difference between the χ data for the alloy and the one for pure ZnTe. However the most important difference is the k -dependence of the amplitudes, which corresponds to about a 0.0007 Å² difference of the EXAFS Debye-Waller factor between the alloy and pure ZnTe. Results for $x=0.20$ sample are essentially the same as for $x=0.10$ sample and are independent of temperature for both. Our results imply: 1) there is no relative shifts between Cd and Te sublattices; 2) First shell Zn-Te bonds have some distortion. If assuming Zn ion is inside an undistorted Te tetrahedron cage, we found Zn ion may be off-center and the displacement is smaller than 0.1 Å. This means the distorted Zn-Te bond could be the source of ferroelectricity of $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ alloys. Polarization XAFS investigation on polarized single crystal samples is our next step.

Fig. 1 Cd K-edge first shell χ data of $\text{Zn}_{0.10}\text{Cd}_{0.90}\text{Te}$ (dash) and CdTe (solid) at 300K

Fig. 2 Zn K-edge first shell χ data of $\text{Zn}_{0.10}\text{Cd}_{0.90}\text{Te}$ (dash) and ZnTe (solid) at 300K

1. R. Weil, R. Nkum, E. Muranevich, and L. Benguigui, Phys. Rev. Lett. **62**, 2774 (1989).
2. Q. T. Islam and B. A. Bunker, Phys. Rev. Lett. **59**, 2701 (1987).
3. B. A. Bunker, Z. Wang and Q. Islam, Proc. of XAFS-VI, York, England, Aug. 1990.

Supported by Office of Naval Research under contract N00014-89-J-1108

EXAFS AND X-RAY EDGE STUDIES OF COBALT SUBSTITUTED

BACTERIORHODOPSIN: IRREVERSIBLE LIGHT INDUCED CATION UPTAKE. M.D. WITT, M.R. Chance, and M. Engelhard. * Dept. of Chemistry, Georgetown U., Washington, DC 20057, Max-Planck Institute for Ernährungsphysiologie, Dortmund, W. Germany.

Bacteriorhodopsin (BR) is the only protein found in the purple membrane of *Halobacterium halobium* and provides the energy for the organism based on its function as a light-driven proton pump. Native BR contains Ca^{2+} or Mg^{2+} ions necessary for normal function. Removal of these metals and reconstitution with other metal ions has allowed a number of spectroscopic measurements that have defined at least two high affinity cation binding sites. Substitution of these metals with cobalt has attracted interest due to its irreversible binding upon illumination. To identify the specific side chains responsible for cobalt coordination, illuminated and unilluminated forms of BR are reconstituted with $[\text{Co}(\text{NH}_3)_4\text{H}_2\text{O}]^{2+}$ in either 3 Co/BR or 1 Co/BR ratios. Illumination of the 3 Co/BR form produces a structural change consistent with carboxylate incorporation. However, no structural change is noted for the 1 Co/BR species, indicating that sites 2 and 3, and not site 1, correspond to the side chains responsible for cobalt coordination. X-ray edge spectra for the $[\text{Co}(\text{NH}_3)_4\text{COO}^-]$ model and the light adapted 3 Co/BR species show a 0.5 eV decrease in the cobalt edge energy (corresponding to a reduction of effective nuclear charge on cobalt) when compared to other BR species (Table 1). The 1 Co/BR species (occupying site 1) shows no edge shift upon illumination, providing evidence consistent with COO^- substitution for water in sites 2 and 3.

TABLE 1. X-ray edge energies for cobalt BR species and model compounds.

COMPOUND	ENERGY (eV)
$\text{Co}(\text{NH}_3)_4\text{H}_2\text{O}$	7720.0
$\text{Co}(\text{NH}_3)_4\text{COO}^-$	7719.5
1 Co/BR	7720.0
1 Co/BR + hv	7720.0
3 Co/BR	7720.0
3 Co/BR + hv	7719.5

EXAFS data for the four BR complexes give average Co-N distances of $1.97 \pm .02$ Å with a Co-O distance of $1.90 \pm .03$ Å. Fourier transforms of the model and all four protein compounds are similar and show no higher shell contributions. This demonstrates that the $[\text{Co}(\text{NH}_3)_4\text{H}_2\text{O}]^{2+}$ species retains its overall integrity when complexed to the protein. Unfortunately, the oxygen of the carboxylate ligand and that of the water could not be distinguished in the analysis. We have provided direct evidence that in sites 2 and 3 the water ligand of $[\text{Co}(\text{NH}_3)_4\text{H}_2\text{O}]^{2+}$ is replaced consistent with carboxylate coordination; where the carboxylate side chains of sequences Glu-Ala-Glu and Asp-Ala-Asp, derived from earlier biochemical characterization, are reasonable coordinating groups. This research is supported by PRF Grant #23257-G3.

X-RAY ABSORPTION NEAR-EDGE STUDY OF SEVERAL BI- AND TI-BASED HIGH- T_c SUPERCONDUCTOR SYSTEMS

Wei Xu, G. Liang, (Sam Houston State U.), M. Croft, J. Chen, J. Peng (Rutgers U.), R. S. Liu (IRC, UK), B. Jayaram (U. Nebraska), J. M. Tarascon (Bell Core)

We have measured x-ray absorption spectra at the Cu K-edge, Ti, Pb, Bi, and Ce L₃-edge for three Ti- and Bi-based high- T_c superconductor systems: $(\text{Ti}_{0.5}\text{Pb}_{0.5})(\text{Ca}_{1-x}\text{Y}_x)\text{Sr}_2\text{Cu}_2\text{O}_{7-s}$, $\text{Bi}_2\text{Sr}_2\text{Ca}_{1-x}\text{Gd}_x\text{Cu}_2\text{O}_{8+y}$, and $\text{Bi}_{2-x}\text{Pb}_x\text{Sr}_2(\text{Sm}_{0.88}\text{Ce}_{0.12})_2\text{Cu}_2\text{O}_{10+y}$. The purpose of this study is to reveal correlations between the electronic/structural changes and the superconducting properties in these materials.

The Cu K-edge spectra of the $(\text{Ti}_{0.5}\text{Pb}_{0.5})(\text{Ca}_{1-x}\text{Y}_x)\text{Sr}_2\text{Cu}_2\text{O}_{7-s}$ series show that the edge-energy does not shift and Cu valence remains unchanged upon the Y^{3+} substitution for Ca^{2+} . The intensity of the $1s \rightarrow 4p_e$ [1] features increase and the feature positions shift downwards in energy with the increase of x . This result can be correlated to the variation of lattice parameters in this series and is consistent with the fact that the normal state resistivity changes gradually from metallic to semiconducting behavior with the increase of x . The Ti and Pb L₃-edge spectra of this compound series show that the intensity of the $2p \rightarrow 6s$ transition feature decreases with the increase of yttrium concentration. This intensity decrease is accompanied with an intensity increase of the $2p \rightarrow 6d_{xy}$ (antibonding) transition peak but the intensity of the $2p \rightarrow 6d_{xy}$ (nonbonding) transition feature remains unchanged [2]. These observations indicate the valences of Ti and Pb decrease somewhat with Y^{3+} substitution for Ca^{2+} but are close to +3 and +4, respectively. Thus, the Y^{3+} substitution for Ca^{2+} could reduce both the $0-2p_{3/2}$ and Ti/Pb $6s$ hole populations.

For the $\text{Bi}_2\text{Sr}_2\text{Ca}_{1-x}\text{Gd}_x\text{Cu}_2\text{O}_{8+y}$ series, the Bi L₃-edge spectra show that Bi valence (close to +3) is basically unaffected by the substitution of Gd^{3+} for Ca^{2+} for both nitrogen and air annealed samples. We found that Gd substitution for Ca changes the Cu K-edge spectra in a similar manner as that induced by Y substitution for Ca in $(\text{Ti}_{0.5}\text{Pb}_{0.5})(\text{Ca}_{1-x}\text{Y}_x)\text{Sr}_2\text{Cu}_2\text{O}_{7-s}$ series. This result is consistent with the results from the transport measurements [3] that the system changes from metallic to semiconducting behavior upon Gd substitution for Ca.

The Ce L₃-edge spectra of the $\text{Bi}_{2-x}\text{Pb}_x\text{Sr}_2(\text{Sm}_{0.88}\text{Ce}_{0.12})_2\text{Cu}_2\text{O}_{10+y}$ with $0.0 \leq x \leq 1.0$ indicate that Ce valent state is close to that in CeO_2 . The lack of the $2p \rightarrow 6s$ feature in both the Pb and Bi L₃-edge spectra indicates that Pb and Bi in this series have constant valences +2 and +3, respectively. This result is consistent with the conclusion from structural analysis [4]. We observed that the edge-energy (and thus the Cu valence) of the Cu K-edge spectra of this series is peaked up at $x = 0.6$ where the superconducting fraction is also a maximum.

[1] J. M. Tranquada et al., Nature 337, 720(1989).

[2] F. Studer et al., Physica C159, 609(1989).

[3] J. Yuan et al., Phys. Rev. B43, 8030(1991).

[4] K. Remsching et al., Physica C170, 284(1990).

[5] B. Jayaram et al., Phys. Rev. B43, 5444(1991).

*This work was supported by Research Corporation, the Faculty Research Funds of Sam Houston State University.

TRENDS IN PROTEIN STRUCTURE CHANGES AROUND THE TRANSITION METAL SITE IN CONCAVALIN A WHEN THE CRYSTALLIZED PROTEIN IS DISSOLVED*

Y. Zhang, E.A. Stern, S.L. Lin (U of Washington), and A.J. Kalb(Gilboa (Weizmann Inst. of Sci.))

A conformational change about the Zn site in Zn-Ca concanavalin A protein was found using XAFS when crystallized protein was dissolved [1]. This phenomenon has now been investigated with different transition metal atoms substituted for the Zn, namely, Ni-Ca, Mn-Ca and Co-Ca concanavalin A, both in crystalline and solution forms. All the experiments were done by measuring the XAFS about the K-edge of the metal atom substituted for Zn.

The log ratio of amplitudes and difference in phase between the crystal and solution of Mn-Ca and Ni-Ca concanavalin A showed that the solution is more ordered by 0.001 Å² and there is no change of coordination number or distances for the 1st neighbors, unlike the case of Zn-Ca concanavalin A protein. The fit of both the crystal and solution forms of the proteins confirmed this with the best models for both being five oxygens and one nitrogen.

For the Co set, the story is different. The log ratio (Fig. 1) between solution and crystal showed the amplitude of solution is about 7% bigger than the crystal which indicates a corresponding smaller coordination number. We get a good fit for the solution of Co-Ca concanavalin A using five oxygen and one nitrogen and for the crystal with 4.5 oxygens and 1 nitrogen. This could be understood as crystal molecules sometimes losing a water oxygen ligand.

The above experiments showed that the local structure surrounding the transition metal site of crystals of concanavalin A depends on the metal atom present: with Zn, a major conformational change occurs on crystallization; with Co a more minor variation occurs with partial loss of water ligands; while with Mn and Ni, the stresses of crystallization only slightly disorder the bond lengths.

Fig. 1.

[1] Lin, S.L., Stern, E.A., Kalb(Gilboa), A.J., and Zhang, Y. (1990) Biochem. 22, 3599-3603; (1991) Biochem. 30, 2323-2332.

* Research funded by NSF grant DMB-8613948.

STRUCTURE AND PHOTO-INDUCED STRUCTURAL CHANGES IN NON-STOICHIOMETRIC a-As_xS_{1-x}

Weiqing Zhou, M. A. Paesler, and D. E. Sayers (NCSU)

X-ray absorption fine structure data show that chemical ordering plays an important role in non-stoichiometric amorphous (a-) As_xS_{1-x} films at 0.4 ≤ x ≤ 0.5. The mixture of different chemical bonding behaviors of As-S and As-As in an a-As₂S₃ random covalent network and an a-As₄S₄ disordered molecular solid lead the structure to a more disordered state, which is shown in the present EXAFS studies by the Debye-Waller factor maximum for 0.4 < x < 0.5. The structure at a composition around x = 0.43 in a-As_xS_{1-x} represents a maximum of flexibility. As with their glassy counterparts, the structure in amorphous annealed films exhibits a distinct compositional trend. Photo-illumination results in (1) a more disordered state as indicated by the increase in the static Debye-Waller factors which are related to the creation of As-As homopolar bonds and (2) modifications in the network structure, such as the increase of the As-As radial distances in the second shells. The relative changes in structural parameters induced by photo-illumination are listed in table 1. In non-stoichiometric a-As_xS_{1-x} (0.4 < x < 0.5), the presence of As₄S₄ molecules in the As₂S₃ network tends to break this network up into small segments. This decoupling of pieces of the As₂S₃ network (comprised of S-bridged As₂S₃ pyramids and As-As wrong bonds) may lead to increased steric freedom which is related to a more disordered state that appears to be associated with photoinduced structural changes in amorphous arsenic sulfide.

Table 1. Relative structural parameters for light-soaked films with respect to annealed films

SAMPLE	SHELL	BOND	Δ(N-Na)/Na	Δ(R-Ra)(Å)	Δσ(10 ⁻⁴ cm ⁻²)
As _{0.42} S _{0.58}	1	As-As	0.5±0.2	0.00±0.01	32±12
	2	As-As	0.0±0.2	0.02±0.01	19±15
As _{0.45} S _{0.55}	1	As-As	0.1±0.2	0.00±0.01	15±12
	2	As-As	0.0±0.2	0.03±0.02	66±15
	2	As-S	0.0±0.2	-0.01±0.02	-12±15

ACKNOWLEDGMENTS

This work is supported by U.S. Department of Energy, Division of Materials Sciences under contract DE-FG05-89ER45384.

X-RAY DIFFRACTION STUDIES OF LIQUID CRYSTALLINE POLYMERS

P. Cebe †, Y. Aihara †, P. Huo †, C. You †, H. Gibson ‡, A. Bluhm ‡, and W. Yeomans ‡

† M.I.T., Dept. of Materials Science and Engineering, Cambridge, MA 02139

‡ U.S. Army Natick Research, Development and Engineering Center, Natick, MA 01760

A new series of liquid crystalline polycarbonates has recently been synthesized [1]. The liquid crystalline polymers (LCPs) contain methyl-substituted stilbene mesogen, and flexible units with carbonate and (CH₂)_n linkages. The number of CH₂ linkages, *n*, varied from *n* = 4 to 12.

Wide angle x-ray scattering (WAXS) allows the determination of the packing arrangement and interchain distances of the molecular segments of the LCPs. We performed real time high temperature WAXS to observe the phase transitions from isotropic melt to nematic, and from nematic to three dimensional crystal. To perform these high temperature studies, a Mettler FP80 hot stage was supported in the x-ray beam path by an aluminum holder. The beam passed through the sample which was encapsulated between two pieces of Kapton tape. The one dimensional gas filled wire detector was positioned to cover the angular range from about 10° to 30° two theta. Wavelength used was that of gold, 0.91 Å.

WAXS proved to be a very sensitive indicator of structure development from the isotropic melt state. The first appearance of the interchain reflections correlated very well with endothermic analysis in which a step in heat capacity was seen at the same temperature.

[1] H. Gibson, A. Bluhm, W. Yeomans, unpublished.

Sponsorship: U.S. Army Summer Faculty Research Program.

SAXS STUDY OF THE FOLDING OF A GLOBULAR PROTEIN IN THE PRESENCE AND ABSENCE OF CHAPERONINS

Authors: John Flanagan*, Mikio Kataoka†, Malcolm Capelt, Donald Engelmann*, *Dept. Molecular Biophys. and Biochem., Yale U., †Physics Dept., Faculty of Science, Tohoku University, and ‡Biology Dept., Brookhaven National Laboratory.

We have initiated SAXS studies to examine the conformation of unfolded *S. nuclease* denatured at low pH, high concentrations of urea, or genetically by deletion of the carboxy terminal 13 amino acid residues of nuclease. Additionally, we have also looked at the solution structures of several of the *E. coli* chaperonins (GroEL, GroES, and DnaK) which are thought to be involved in the *in vivo* folding reaction. To date we have examined the equilibrium denatured conformation of nuclease present at low pH (R_g = 25 Å), in 6M urea (R_g = 35 Å) and by genetic deletion of the carboxy terminal 13 residues (R_g = 20 Å). Additionally, information from the extended scattering curves suggests that the structure of denatured nuclease does not approximate a random coil and, in fact, differs greatly depending upon how the protein is denatured. We have also examined the effects of single amino acid substitutions in the nuclease sequence which are known to effect the protein's thermal stability. We have found several single amino acid substitutions which appear to exert their effect by altering the size and structure of the denatured state of nuclease. These studies provide us with a basis for subsequent time resolved folding/unfolding studies of nuclease.

In a preliminary experiment we have examined the scattering curve of the chaperonin GroEL, a 910 kDa protein composed of 14 monomers, 65 kDa each. Electron microscopy has shown that the subunits are arranged as two rings containing 7 monomers each. Our preliminary scattering curves show numerous subsidiary maxima indicative of its generally spherical shape and possibly due to interference between the individual monomers. The radius of gyration is large, 62 Å and the I(0) is consistent with the scattering species having a molecular weight of 910 kDa. In subsequent experiments we will look at GroES, the ~70 kDa putative polypeptide release factor for GroEL (6 or 7 monomers/molecule), and GroEL liganded to ATP, ADP, GroES, and denatured polypeptide. The experiments should help to place constraints on how GroEL and GroES function in the cell.

THE ROLE OF HISTONE H1 IN CHROMATIN HIGHER-ORDER STRUCTURE

V. Graziano, M. Capel and V. Ramakrishnan (Biology Department, BNL).¹

The nature of the organization of nucleosomes into chromatin higher-order structure is one of the central problems in chromatin structure today (1). The linker histone H1 plays an essential role in this organization. Histone H1 consists of three main domains, a central globular domain flanked by two more extended domains. The globular domain is highly conserved, and interacts with the nucleosome. We are engaged in a study of the role of the various domains of histone H1 by the production of H1 mutants in *E. coli* that lack increasing amounts of the C-terminal tail. In this experiment, as a first step, we look at the effect of removing linker histones H1/H5 from chromatin and reconstitution with recombinant H1. The purpose is to ensure that the recombinant protein is fully capable of functioning as well as the native protein from chicken erythrocytes. The small-angle camera at the NSLS is required, because we require extremely good collimation to be able to measure spacings of the order of 1000 Å or more, in order to get a substantial Guinier region for fibers of the size of 300 Å. Figure 1 shows cross-sectional Guinier plots of chromatin reconstituted with recombinant H1, in increasing ionic strengths. These plots are very similar to those obtained from native H5 (2) and native H1 (3). Nucleosome core particles were used to calibrate the absolute intensity. Using this and the slope and intercepts from Figure 1, the mass per unit length and the cross-sectional radius of gyration were calculated as a function of ionic strength (Figure 2). These values agree very well with those obtained for native chromatin, and for chromatin reconstituted with H1 or H5, thus establishing the use of the recombinant H1 as a basis to study the structure and function of H1.

(1) Felsenfeld, G. and McGhee, J. *Cell* 44, 375 (1986).

(2) Graziano, V. *et al. J. Mol. Biol.* 203 997 (1988).

(3) Graziano, V. and Ramakrishnan, V. *J. Mol. Biol.* 214, 897 (1990).

¹ This work was supported by NIH grant GM 42796 and the Office of Health and Environmental Research, U.S. Department of Energy.

X-RAY DIFFRACTION STUDIES OF BLENDS OF POLY(BUTYLENE TEREPHTHALATE) WITH POLYARYLATE

P. Huo, C. You, Y. Aihara, and P. Cebe

Department of Materials Science and Engineering
Massachusetts Institute of Technology
Cambridge, MA 02139

High intensity synchrotron x-ray sources provide the opportunity to monitor the crystal structure development as a function of temperature in the blend system, poly(butylene terephthalate) with polyarylate, PBT/PAr. Homopolymer PAr is completely non-crystalline, while homopolymer PBT is semicrystalline. In blends of PBT with PAr, crystallization kinetics and degree of crystallinity will depend on the composition ratio of PBT to PAr.

Wide angle x-ray scattering (WAXS) was used to study the nonisothermal crystallization kinetics of PBT/PAr blends. WAXS intensity profile was taken in real time as the temperature increased from room temperature to 230°C, during which we observed the crystallization of originally amorphous PBT/PAr blends of varying composition. The high temperature studies were performed by putting the sample into a Mettler FP80 hot stage. The consistency between x-ray scattering data and kinetics determined by thermal analysis appears to be very good.

For melt and cold crystallized PBT/PAr blends, we monitored the change of WAXS scattering peak angular position and intensity as a function of temperature up to the melting point of the blends. From the peak angle shift, we are able to calculate the thermal expansion coefficient of PBT crystals in the PBT/PAr blends.

Sponsorship: U.S. Army Research Office, Grant DAAL03-91-G-0132.

X-RAY DIFFRACTION FROM MIXTURES OF COLLAGEN TYPES I AND III

V. James, J. McConnell (U of NSW); and M. Capel (BNL)

The conclusions from our use of NSLS in 1989 were reported in James et al., BBA 1078 (1991) 19-22. We have established that an increase in the percentage of collagen type III in a types I and III mixture in mitral heart valves does not of itself produce a decrease in the gap-overlap or d-spacing of the collagen molecule, but we observed a definite decrease in d with increase in age. In 1990, we exposed a greater age range of collagenous materials containing types I and III mixtures, both normal and diseased. The additional meridional reflections recorded have confirmed the original findings. We were also able to gather from the equatorial reflections, information regarding the lateral packing of collagen fibrils, to be used in conjunction with earlier work reported [James et al., Diabetes 40 (1991) 391-394].

Together with results obtained by us elsewhere, we are now heavily engaged in a program to extend our investigations into a whole range of other collagenous materials, such as human breast ducts (normal and cancerous), skin (normal and diabetic), arteries and the corneal material of eyes. All of these materials have approximately the same collagen type III content as heart valves, and should enable us to establish whether our observed changes are tissue-specific.

Our efforts to date have brought requests for collaboration with five laboratories concerned with biological materials, two from USA and one each from UK, USSR and Australia.

One of our ultimate objectives relates to the nature of the spatial relationship between collagen types I and III in the combined forms, and eventually the structure of the collagen type III molecule itself. The latter is considerably longer than type I, due to a longer telopeptide region, so the fact that its presence produces such small changes in d-spacing has led to a considerable amount of speculation, upon which we have begun to shed some light.

Research sponsored by the Australian Research Council and the Commonwealth Department of Industry, Technology and Commerce.

PHASE STUDY OF AOT/WATER/DECANE DENSE MICROEMULSION

E.Y. Sheu*, M.M. De Tar*, and M. Capel†, * Texaco R&D, Beacon, N.Y. 12508; † BNL, Dept. of Biology

A series of AOT/water/decane microemulsions were studied using Beamline X12B small angle spectrometer. The volume fraction of AOT + water ranged from 0.76 to 0.94, and the temperature spanned from 12 to 35 °C. The molecular ratio of water to AOT was fixed at 40.8. Figure 1 shows two typical scattering spectra for 0.8 volume fraction at 25 °C (1), and 35 °C (2) respectively. At 25 °C, the system is in the L_1 phase, consisting of droplets with water as the core, coated with AOT surfactants. As the temperature increased to 35 °C, the system evolved into a coexistence region where the L_1 phase was present along with the L_3 phase, a bilayer structural phase. A comprehensive scattering study of this system is given in Figure 2, where the open circles represent data obtained from a rotating anode small angle X-ray spectrometer at Oak Ridge National Laboratory, and the filled circles represent data from X12B.

The L_3 phase observed in this study (see III in Fig. 2) differs from the L_3 phase observed previously, in which a considerable amount of salt was necessary for this phase to form. In our study the L_3 phase formed in a salt-free environment, implying that this phase can form without added counterions. This finding is significant for theorists to predict micro-emulsion phases based on thermodynamic free energy arguments.

Fig. 1. Scattering intensities for AOT/water/decane for $\phi = 0.8$ at 25 and 35 °C.

Fig. 2. Phase diagram of AOT/water/decane ($[water]/[AOT] = 40.8$).

E.Y. Sheu, M.M. DeTar, D.A. Storm, Texaco Research, 4001 Gulfport, Beacons, NY 12508 and M.S. Capel, Dept Biology, Brookhaven National Laboratory, Upton, NY 11973.

The residual material (vacuum residue, VR) following the refining of crude oil forms colloidal particles in organic solvents. These colloids have been known to hinder further refining and are the major roadblock for VR upgrading. In this experiment, we characterized the properties of a VR derived from a Neutral Zone crude oil, in terms of their structure and polydispersity. We fractionated this VR into two fractions, a pentane soluble (CSS) fraction, and a heptane insoluble (C7I, or asphaltene) fraction.

The extracted CSS was then dissolved in a series of n-alkane solvents from pentane to hexadecane, and the C7I fraction was dissolved in a series of mixed solvents containing toluene and pyridine of various ratios. The SAXS intensity distributions of these systems showed strong small angle scattering, indicating a substantial amount of suspended colloidal particles. A typical $\ln(I(Q))$ vs $\ln(Q)$ plot is shown in Fig. 1. The final slope is much less than four, indicating that the particles are nonspherical, the particle surface is rough, or the system is polydisperse. A detailed analysis showed that the structure is approximately spherical (in agreement with rheological and SANS studies [1]). The polydispersity is about 15% for most of the CSS cases and 25% for C7I cases. The solubility of CSS was found to increase as a function of carbon chain length. This result agrees with Ali and Saleem's study [2] using a fractionation method. We also qualitatively analyzed the particle's surface. Fig. 2 shows a typical Porod plot. No distinctive limit at large Q was obtained. In fact, the slope is significant, meaning that although the particles are approximately spherical, the surfaces are very 'rough'.

[1] E.Y. Sheu, M.M. DeTar, and D.A. Storm, "Asphaltenes in Polar Solvents", to appear in *J. of Noncrystalline Solids*.

[2] M.F. Ali and M. Saleem, *Fuel Science and Technology International*, 6 541 (1988).

Fig. 1

Fig. 2

SAXS STUDIES OF POROUS CARBONS

J. Wegrzyn (BNL-DAS*), A. N. Goland (BNL-DAS), M. Capel (BNL-Biology), and J. Schwartz (Syracuse U.)

Carbon is an element with the unique ability to chemically bond to itself. This results in an immense variety of structures and properties for carbon. For example, it is well known that different carbon pore structures help govern its affinity for adsorbing gases. Additionally, porous activated carbons exhibit one of the strongest low-angle scattering patterns ever recorded, which makes SAXS a powerful tool in classifying the internal structure of activated carbon. Structural studies on porous materials are critical to the further development of a practical low-pressure adsorbent storage system for natural gas. Therefore, five commercially available carbons were studied from which strict Porod scattering ($I(q) \sim A/q^4$) was not observed. Instead, the scattered intensity patterns were complex with each carbon type generating a reproducible but distinct scattering pattern that can be described approximately for $q > 0.1$ (q -scattering vector) as damped quasi-periodic. A circular scattering pattern typical for powdered samples has been assumed for analysis purposes. Figures 1, 2, and 3 show some of the results, where the incremental radial intensity $2\pi R \Delta R I(q)$ was plotted as a function of q . These results indicate the presence of oscillations in scattered intensities that are proportional to either the sine or cosine functions with a period of $(q\delta/2\pi)$. Oscillations in scattered intensity have been observed previously by others and have been discussed by Guinier and colleagues [1]. They can be related to the average square of the material's structure factor through the expansion of the correlation function $\gamma(r)$. A more recent paper [2] has given a geometrical interpretation to the δ term as being the distance between sharp boundary, parallel internal surfaces. This finding has direct application to the characterization of activated carbons since their microporous structures are convoluted two-dimensional graphic layers bound together by strong crosslinking in the third dimension. Further work is planned in the next year with the replacement of the 1D detector by a 2D detector, which is better suited for measuring the circular scattering patterns associated with powdered samples. This replacement will further help to resolve the complex issues of determining the structures of porous carbons as described by SAXS theory.

1. Guinier, A., et al., *Small Angle Scattering of X-rays*, p. 17, J. Wiley & Sons, 1955.
2. Cicciariello, S., *Phys. Rev. A* 44, No. 5, 2975 (1991).

FIG.1 ACTIVATED CARBON
WESTACO 89F-1FIG.2 ACTIVATED CARBON
NORIT 89F-1FIG.3 ACTIVATED CARBON
CALSON 89F-6

*Work supported by the U.S. Department of Energy, Office of Alternative Fuels, Contract No. DE-AC02-76CH00016.

STRUCTURE DETERMINATION AND REFINEMENT OF BOVINE LENS LEUCINE AMINOPEPTIDASE AND ITS COMPLEX WITH BESTATIN

S. K. Burley (Harvard U.), P. R. David (Harvard U.), R. M. Sweet (NSLS), A. Taylor (Tufts U.) and W. N. Lipscomb (Harvard U.)

The three-dimensional structure of bovine lens leucine aminopeptidase (EC 3.4.11.1) complexed with bestatin, a slow-binding inhibitor, has been solved to 3.0 Å resolution by the multiple isomorphous replacement method with phase combination and density modification. In addition, this structure and the structure of the isomorphous native enzyme have been refined at 2.25 and 2.32 Å resolution, respectively, with crystallographic R factors of 0.180 and 0.159, respectively. The current structural model for the enzyme includes the two zinc ions and 481 of the 487 amino acid residues comprising the asymmetric unit. The enzyme is physiologically active as a hexamer, which has 32 symmetry, and is triangular in shape with a triangle edge length of 115 Å and maximal thickness of 90 Å. The secondary structural composition is 35% α -helix and 23% β -strand. The N-terminal domain (160 amino acids) mediates trimer-trimer interactions and does not appear to participate directly in catalysis, while the C-terminal domain (327 amino acids) is responsible for catalysis and binds the two zinc ions, which are less than 3 Å apart. The positions of zinc ions were confirmed using anomalous scattering. The zinc ion that has the lower temperature factor is coordinated by one carboxylate oxygen atom from each of Asp-255, Asp-332 and Glu-334, and the carbonyl oxygen of Asp-332. The other zinc ion, presumed to be readily exchangeable, is coordinated by one carboxylate oxygen atom of each of Asp-273 and Glu-334 and the side chain amino group of Lys-250. The active site also contains two positively-charged residues, Lys-262 and Arg-336. Bestatin binds to one of the active site zinc ions, and its phenylalanine and leucine side chains occupy hydrophobic pockets adjacent to the active site.

The full paper has been accepted by the Journal of Molecular Biology.

This research was supported by the National Institutes of Health, GM06920.

THREE-DIMENSIONAL STRUCTURE OF T7 LYSOZYME

X. Cheng, J.W. Pflugrath (Cold Spring Harbor Laboratory), X. Zhang, and F.W. Studier (Brookhaven National Laboratory)

T7 lysozyme is a bi-functional protein: it is an N-acetyl-muramyl-L-alanine amidase that cuts a bond on the bacteria cell wall, and it can specifically bind and inhibit transcription by T7 RNA polymerase. The three-dimensional structure of T7 lysozyme will provide the structural basis for the unique substrate specificity of this lysozyme, and the mechanism for its interaction with T7 RNA polymerase.

Extensive trials to obtain crystals of wild type T7 lysozyme have been unsuccessful. We constructed a mutant lysozyme which deletes four amino acids from the N-terminal. This mutant lysozyme has the equivalent cell wall cutting activity as wild type but completely lacks the activity of transcription inhibition. The mutant protein crystallizes in space group C222₁ with cell dimensions $a=46.5$ Å, $b=62.5$ Å, and $c=110.0$ Å. We have grown crystals as large as 0.5 by 0.6 by 1.5 mm overnight, which diffracted to at least 2.0 Å resolution at the National Synchrotron Light Source.

We used the facilities at beamline X12-C at the Brookhaven National Laboratory with the assistance of Dr. R.M. Sweet. The optics and intense tunable radiation of the Synchrotron Light Source let us collect data sets from native crystals and different heavy atom soaked crystals using a wavelength of 1.10 Å. Using this native data set, the data set from potassium tetrachloroplatinate (K_2PtCl_6) soaked crystals, and a data set from mercury chloride ($HgCl_2$) cocrystals (which was collected in Cold Spring Harbor Laboratory with a rotating anode generator), we obtained a single platinum site (Fig. 1) and a single mercury site (Fig. 2) in isomorphous difference Patterson maps. These derivatives along with mercury anomalous scattering signals constituted the basis of our approach to solve the phase problem. Further analysis has resulted in an interpretable electron density map (Fig. 3).

Fig.1 Harker section at $w=1/2$. Fig.2 Harker section at $w=1/2$. Fig.3 A section of electron density map at 3.0Å.

Peaks show platinum sites. Peaks show mercury sites.

This work is supported in part by the W.M.Keck Foundation to X.C. and J.W.P.

ABSTRACT

Structural Studies of Tryptophanyl-tRNA Synthetase Using Multiwavelength Anomalous Dispersion from Selenomethionine-Substituted Protein Crystals. S. Doublié, R.M. Sweet* and C.W. Carter, Jr., Department of Biochemistry Campus Box 7260, UNC-Chapel Hill, Chapel Hill, NC 27599-7260 and (*) Department of Biology, BNL Associated Universities, Upton, NY 11973. June 1990 - July 1991

We are working on the X-ray crystal structure of the tryptophanyl-tRNA synthetase from *B. stearothermophilus*. Loss of isomorphism has been a critical obstacle to phasing this structure using heavy atom derivatives. Recent evidence that substantially correct native phases can be estimated by triangulation using Bijvoet difference measurements at several different wavelengths for a number of smaller proteins lead us to pursue substitution of the 10 methionines in TrpRS with selenomethionine.

Crystals of selenated protein diffracted to quite high resolution (better than 1.7 Å in 1.5 degree oscillation photographs taken at beamline X12C). The X12C facility presents a nearly unique capability for us to do this kind of multiwavelength experiment. The unit cell of the crystals presents a number of difficulties for data collection. It is very asymmetric (60 Å x 60 Å x 234 Å). The CAD4 goniostat permits us to prealign these crystals rapidly enough to insure that all crystals are being rotated for data collection with a mirror plane relating all reflections measured; hence Bijvoet related reflections can be measured simultaneously. Moreover, the beamline optics at X12C provide adequate intensities at the three wavelengths necessary to include the absorption edge of Selenium, and the adjustment of the monochromator can be done rapidly enough to facilitate collection of intensities for all three wavelengths from a single crystal. To our knowledge, no other facility can provide this particular instrumental combination of optics, intensity, goniometry, and detector.

Three different visits to the X12C beamline at the National Synchrotron Light Source at Brookhaven National Laboratories provided very encouraging evidence that multiwavelength data can be collected from these crystals. An early attempt to solve for the selenium positions by Direct Methods from amplitudes estimated from an initial (1990) three wavelength experiment produced a single acceptable phase set for which the electron density map revealed ten peaks above 5σ. All but one of these lay very close to the molecular envelope, which had previously been determined (1). Unfortunately, this solution was consistent only with a subset of the best amplitudes, and was inconsistent with the majority of the amplitudes. Several reasons for this inconsistency emerged from a retrospective analysis of statistical properties of the data, together with studies carried out by Peter Lee at BNL. These implicated several instrumental problems that could not easily be rectified post hoc. Two additional trips to Brookhaven in 1991 allowed us to correct many of these problems, and we recollected anomalous data for one wavelength at the absorption edge for selenium using lengthened exposure times, which should contribute to a stronger signal. We are now integrating and reducing these data.

1. Carter, C.W., Jr., Crumley, K.V., Coleman, D.E., Hage, F., and Bricogne, G.: Direct Phase Determination for the Molecular Envelope of *Bacillus stearothermophilus* Tryptophanyl-tRNA Synthetase by X-ray Contrast Variation. *Acta Crystallographica*, A46:57-68, 1990.

STRUCTURAL STUDIES OF THE MALIC ENZYME FROM THE PARASITIC NEMATODE ASCARIS SUUM AND THE INTERLEUKIN-1 RECEPTOR ANTAGONIST PROTEIN.

H. M. Einspahr and B. C. Finzel, The Upjohn Company, Kalamazoo MI 49001.

The mitochondrial malic enzyme (ME) from the parasitic nematode *Ascaris suum* is a tetramer of 65 kD monomers that catalyzes the conversion of malate to pyruvate and is critical for muscle function under anaerobic conditions. We crystallize the protein, a natural isolate, from solutions containing magnesium sulfate and NAD, cofactors of the enzyme, together with sodium tartrate, the anion of which is a substrate analog inhibitor. The crystals are trigonal, space group P3₁21 or its enantiomer, with $a = 131.2(7)$ Å, $c = 152.6(9)$ Å, and two monomers per asymmetric unit. We have collected several data sets from malic enzyme crystals on the x-ray data collection facility on beamline X12-C at NSLS. These data sets establish a practical resolution limit for our crystals of at least 3 Å. Data collected at NSLS also played a role in establishing the 222 symmetry of the ME tetramer and defining the orientations of the symmetry axes by self rotation function analyses (see example below).

The interleukin-1 receptor antagonist protein (IRAP) is a member of the interleukin-1 family of cytokines. Interleukins-1α and 1β, the other members of this family, are 15 kD proteins that, while distantly related in sequence, are very similar in tertiary structure. IRAP is also about 15 kD and sequence comparisons suggest that it too will have the interleukin-1 tertiary fold. But while its relatives bind to the interleukin-1 receptor and produce agonist responses, IRAP competes for binding to the receptor and elicits no response that has been detected. The IRAP protein we use is produced in *E. coli* cells by recombinant methods. We crystallize IRAP by PEG precipitation at 4°C. A data set collected at X12-C was the first we were able to collect of this protein. For this data set, we used the chilled N₂ system to collect at 4°C and only one crystal was required. The X12-C data revealed the crystal to be tetragonal, with space group P4₁2₁2 or its enantiomer and cell dimensions of $a = 71.1$ Å and $c = 116.4$ Å. The resolution limit of this data is 2.2 Å. We have since learned to adapt IRAP crystals to room-temperature data collection in our home laboratory and have begun the search for derivatives.

PROBABLE AMYLASE/PROTEASE INHIBITOR-B FROM RICE SEEDS

Kwang Yeon Hwang, Kyeong Kyu Kim, and Se Won Suh

Department of Chemistry, College of Natural Sciences,
Seoul National University, Seoul 151-742, Korea

A new protein, called Probable Amylase/Protease Inhibitor (PAPI), was purified from rice seeds and its amino acid sequence was determined. It is strongly homologous with an α -amylase inhibitor I-2 from seeds of ragi, Indian finger millet and a 9 kDa barley seed protein, also called a probable amylase/protease inhibitor (PAPI). The rice PAPI did not inhibit any of the proteases and amylases tested. This is in agreement with the recent suggestion that the PAPI may serve as phospholipid transfer proteins on the basis of striking sequence similarities. Proteins of this family are predicted to have a novel tertiary structure with little α -helix but no X-ray structure has been determined.

Large crystals of PAPI-B have been grown from ammonium sulfate solutions. The crystals are tetragonal and belong to the space group P4₁2₁2 (or P4₃2₁2) with unit cell dimensions of $a = b = 38.02$ Å and $c = 98.08$ Å. The asymmetric unit contains one molecule of PAPI-B, with a crystal volume per protein mass (V_M) of 1.97 Å³/Da. The crystals diffract beyond 1.7 Å Bragg spacing when exposed to synchrotron X-rays and they are reasonably stable in the X-ray beam. Native data have been collected to 2.0 Å Bragg spacing. Potential heavy atom derivatives have been prepared.

Publications

Kwang Yeon Hwang, Kyeong Kyu Kim, Sangsoo Kim, Robert M. Sweet, and Se Won Suh (1991), manuscript in preparation.

Supported by Korea Ministry of Education and Pohang Light Source

THERMOSTABLE α -AMYLASE FROM *B. LICHENIFORMIS*

Kwang Yeon Hwang, Kyeong Kyu Kim, and Se Won Suh

Department of Chemistry, College of Natural Sciences,
Seoul National University, Seoul 151-742, Korea

Thermostable α -amylases from *Bacillus* species are industrially important in production of corn syrups and dextrose. α -Amylase from *B. licheniformis*, consisting of 483 amino acids with relative molecular mass of 55,200, shows a remarkable heat stability: its optimum temperature is 90°C and only 10 % of activity is lost after heat treatment at 90°C for 30 min. However, there is no clear understanding of the factors responsible for the thermostability of these enzymes, partly owing to the lack of knowledge about their three-dimensional structures. At present crystal structures of only two fungal and a mammalian α -amylases have been determined: Taka-amylase A from *Aspergillus oryzae*, acid α -amylase from *Aspergillus niger*, and α -amylase from porcine pancreas. And no structures of bacterial α -amylases have been determined.

α -Amylase from *B. licheniformis* has been crystallized by the hanging-drop vapor diffusion method in the presence of calcium ions using ammonium sulfate as precipitant. The crystals are tetragonal, belonging to the space group P4₁2₁2 (or P4₃2₁2), with unit cell dimensions of $a = 119.9$ Å and $c = 85.4$ Å. The asymmetric unit contains one molecule of α -amylase, with a crystal volume per protein mass (V_M) of 2.78 Å³/Da. The crystals diffract to better than 2.0 Å Bragg spacing when exposed to synchrotron X-rays and they are reasonably stable in the X-ray beam. Native data have been collected to 2.4 Å Bragg spacing. Potential heavy atom derivatives have been prepared.

Publications

Suk Yeong Lee, Sangsoo Kim, Robert M. Sweet, and Se Won Suh (1991) Arch. Biochem. Biophys. 291 (in press)

Supported by Korea Ministry of Education and Pohang Light Source.

X-RAY CRYSTALLOGRAPHY OF ARYLESTERASE

Kyeong Kyu Kim, Kwang Yeon Hwang, and Se Won Suh

Department of Chemistry, College of Natural Sciences,
Seoul National University, Seoul 151-742, Korea.

Esterases (EC 3.1), widely distributed in nature, catalyze the hydrolysis of ester bonds. Arylesterases (EC 3.1.1.2) have little action on aliphatic esters and are insensitive to DFP. They are characterized by their preference for aromatic carboxylic acid esters, carbamates, paraoxon, and any other organophosphates. These arylesterases play a role in removing selectively the toxicity of organophosphates in human beings. Despite their potential applications, only a few esterases have been extensively purified and characterized biochemically. And detailed structural information is lacking. Since this enzyme is an interesting target for protein engineering studies, information about its tertiary structure and action of the enzyme reaction would be of great significance. X-ray data of arylesterase from *Pseudomonas fluorescens* have been collected.

Molecular weight: 54,000 dimer of identical subunits
Cell parameters: $a = 147.25 \text{ \AA}$, $c = 131.29 \text{ \AA}$
Space group: Trigonal, $P3_1$ or $P3_112$ or $P3_121$
(or enantiomorph);
2 or 3 molecules per asymmetric unit
($V_M = 3.8$ or 2.5)

Crystal size: Plate, $0.8 \times 0.8 \times 0.3 \text{ mm}$
Maximum resolution: 2.4 $\text{\AA}$
Data collection: Native to 2.65 $\text{\AA}$
Radiation damage: Crystal life is a little less than average.
Previous history: Crystal can be translated in the beam.
Maximum resolution was 3.2 $\text{\AA}$ on the FAST diffractometer system with rotating anode.

Publications

Kyeong Kyu Kim, Kang-Duk Choi, Ock Jun Yoo, Sangsoo Kim, Robert M. Sweet, and Se Won Suh (1991), manuscript in preparation.

Supported by Korea Science and Engineering Foundation and Pohang Light Source

X-RAY CRYSTALLOGRAPHY OF LIPASE

Kyeong Kyu Kim, Kwang Yeon Hwang, and Se Won Suh

Department of Chemistry, College of Natural Sciences,
Seoul National University, Seoul 151-742, Korea

Lipases (EC 3.1.1.3), present in several organisms including animals, plants, fungi, and bacteria, show a wide variety of sizes, substrates and positional specificities, and catalytic rates. They catalyze the hydrolysis of triglycerides into free fatty acids and glycerols. Lipases are active at the interface of water and water-insoluble lipid. They are useful for transesterification, synthesis of esters and peptides, and resolution of racemic mixtures to produce various optically active compounds. They have been classified as serine hydrolases owing to their inhibition by DFP (diisopropyl fluorophosphate), and a catalytic triad of Asp-His-Ser (or Glu-His-Ser) has been observed in the crystal structures of two fungal and human pancreatic lipases. Despite their potential importance in industry, no X-ray structures of bacterial lipases are currently available.

Monoclinic crystals of lipase from *Pseudomonas cepacia* have been grown using MPD as precipitant and native X-ray data to 2.2 $\text{\AA}$ have been collected.

Molecular weight: 33,000 monomer
Cell parameters: $a = 84.91 \text{ \AA}$, $b = 47.33 \text{ \AA}$, $c = 86.00 \text{ \AA}$,
 $\beta = 116.09$

Space group:

$P2_1$;
2 or 3 molecules per asymmetric unit
($V_M = 2.35$ or 1.57)

Crystal size:

Plate, $0.45 \times 1.0 \times 1.0 \text{ mm}$

Maximum resolution:

1.55 $\text{\AA}$

Data collection:

Native to 2.2 $\text{\AA}$

Radiation damage:

Crystal life is longer than average. Crystal can be translated in the beam several times.

Previous history:

Maximum resolution limit was 2.3 $\text{\AA}$ with rotating anode X-rays.

Publications

Kyeong Kyu Kim, Hye Sung Jeon, Sangsoo Kim, Robert M. Sweet, and Se Won Suh (1991), manuscript in preparation.

Supported by Korea Science and Engineering Foundation and Pohang Light Source

J. KURIYAN^{1,2}, T.S.R. KRISHNA^{1,2}, B. GUENTHER¹, X.P. KONG¹ & R.M. SWEET³

¹Laboratory of Molecular Biophysics and ²Howard Hughes Medical Institute, The Rockefeller University, 1230 York Avenue, New York, NY 10021

³Department of Biology, Brookhaven National Laboratory, Upton, NY 10021

We are working on the mechanisms of two related redox enzymes, *E. coli* thioredoxin reductase (ThioR) and *Cryptosporidium parvum* trypanothione reductase (TrypR). The refinement of the structure of ThioR against data to 2.0 Å resolution (measured at Beamline X12-C) has been completed. This structure reveals a different evolutionary solution to the problem of NADPH/disulfide reduction than that represented by enzymes such as glutathione reductase (GR).¹ A detailed description of the high-resolution refinement is in preparation.² TrypR is specific to the protozoan parasites that cause African trypanosomiasis, Chagas' disease and leishmaniasis. It is highly homologous to glutathione reductase, but the two enzymes are mutually exclusive for substrate. We have determined the structure of TrypR in order to understand this enzyme's unique substrate binding properties, with the eventual aim of designing specific inhibitors. A dataset was measured by us at BNL, and structure determination by molecular replacement (using that of human GR as a search model) was successfully accomplished.³ The core structure surrounding the catalytic machinery is almost identical in the human and parasite enzymes. However, significant differences are found at the substrate binding sites. These results provide a detailed view of the residues that can interact with potential inhibitors, and complement previous modelling and mutagenesis studies on the two enzymes.

Figure. Active site region of trypanothione reductase compared with that of the human enzyme glutathione reductase (Karpus and Schulz), showing the relative displacements of the FAD and central domains that result in a more open active site in the parasite enzyme. Thick lines: TrypR, Thin lines, GR. C α atoms are shown for the most of the FAD and central domains (marked FAD/central (1)), a small part of the NADPH domain (NADPH(1)) and part of the interface domain of the other molecule in the dimer (Interface(2)). Chain breaks are indicated by solid hexagons, and the N-terminus of the first molecule and the C-terminus of the 2nd molecule are marked N and C.

References:

1. J. Kuriyan *et al.*, (1991) *Nature (London)* 352, 172-174.
2. J. Kuriyan, T.S.R. Krishna, G. Waksman, R.M. Sweet, in preparation.
3. J. Kuriyan, X.-P. Kong, T.S.R. Krishna, R.M. Sweet, N. J. Murgolo, H.M. Field, A. Cerami & G.B. Henderson (1991) *Proc. Natl. Acad. Sci. (USA)*, 8764-8768.

Support from DOE-OHER, the Pew Scholars Program and the NIH is gratefully acknowledged.

THE STRUCTURES OF DNA AND GM-CSF

Judith LaLonde (U. Penn.), K. Swaminathan (U. Penn.), Robert M. Sweet (NSLS), and Donald Voet (U. Penn.)

We are engaged in a program to determine the X-ray structures of duplex DNAs consisting of non-complementary oligonucleotides. We have synthesized, purified, and crystallized a number of these DNAs. In general, the crystals rapidly decay in the X-ray beam and, moreover, we usually obtain no more than a handful of these crystals. The use of the NSLS allows us to determine the quality of the crystals with a few minutes work and, if the crystals turn out to be of diffraction quality, to collect X-ray intensities so as to minimize the rate of radiation decay with respect to the rate of data collection. Unfortunately, we have not been able to obtain DNA crystals that diffract to high resolution although we are continuing our efforts to obtain such crystals.

A second project is to determine the high resolution X-ray structure of human granulocyte-macrophage colony-stimulating factor (GM-CSF), a protein growth factor for certain types of white blood cells that has considerable medical importance for treating immunosuppressive conditions such as AIDS and the side effects of cancer chemotherapy and radiation therapy. The crystals of this protein that we have obtained diffract X-rays to a radiation limit of about 2.6 angstroms. We have therefore used the NSLS to extract intensities to the maximum possible resolution limit of the crystals. GM-CSF crystals are particularly reluctant to bind heavy atoms as is prerequisite to the use of the method of isomorphous heavy atom replacement to solve the crystal structure. The NSLS has been especially useful for determining whether or not a crystal that has been soaked in a solution containing a particular heavy atom has, in fact, formed an isomorphous heavy atom derivative. This structural determination project is still in progress so that no publications have yet resulted from the work at NSLS.

The DNA project was sponsored by NASA grant NAG8-770, Growth of DNA Crystals in a Microgravity Environment.

The GM-CSF project was sponsored by NIH grant GM-42571, X-Ray Structure of GM-CSF.

X12C
STRUCTURAL ANALYSIS OF THE *E. COLI* TRP REPRESSOR L-TRYPTOPHAN BINDING POCKET: WHY 5-METHYLTRYPTOPHAN IS A BETTER COREPRESSOR

Catherine L. Lawson and Robert M. Sweet, Biology Department, BNL

The *trp* aporepressor, a small dimeric ligand-activated DNA-binding protein, is a key component in the transcriptional regulation of L-tryptophan synthesis in *E. coli*. The orthorhombic crystal form of the active repressor (*trp* aporepressor complexed with the natural ligand L-tryptophan, space group $P2_12_12_1$, $a=53.2$ Å, $b=53.4$ Å, $c=33.2$ Å) will readily exchange L-tryptophan for analogs with similar affinity for the binding pocket¹. Our research project entails crystallographic determination of aporepressor/tryptophan analog complex structures. The goal is to correlate structural properties of the ligand binding pocket with observed tryptophan analog binding affinities and biological activities. Preliminary results with our first analog, 5-methyl-DL-tryptophan (5Metrp), are presented here. The activity of 5Metrp is commonly attributed to the L-enantiomer. 5Metrp is superior to L-tryptophan as a corepressor (i.e., 5Metrp is better at promoting repression of transcription)², and the compound binds with 4-fold higher affinity to the protein (based on L-enantiomer concentration)³.

Diffraction measurements were recorded from two crystals: aporepressor/L-tryptophan and aporepressor/5Metrp. High quality data sets ($R_{\text{merge}}=4.5-5.5\%$) were obtained from each crystal to a resolution of 1.6 Å, with 85-87% complete collection of the unique region of reciprocal space. Optimal phases were obtained by refining a previous model for this crystal form⁴ against the aporepressor/L-tryptophan data. The resulting $(|F_{\text{obs}}(5\text{Metrp})| - |F_{\text{calc}}(\text{L-trp})|)$ difference Fourier map (Fig. 1) suggests that the methyl substitution produces a shift of the ligand indole ring of the along its plane in the binding pocket, which may allow better Van der Waal's and hydrogen bond contacts, as well as lower ligand torsional strain. In contrast, no changes are indicated in the amino-acid portion of the ligand, supporting the assumption that only the L-enantiomer of 5Metrp binds. Independent refinement of the aporepressor/5Metrp structure will be required for full interpretation of the structural differences.

Figure 1. L-tryptophan model superimposed on the $(|F_{\text{obs}}(5\text{Metrp})| - |F_{\text{calc}}(\text{L-trp})|)$ difference Fourier. Thin contours represent positive density ($+5\sigma$), thick contours, negative density (-5σ). The largest positive density is the probable position of the methyl group in 5Metrp.

¹Lawson and Sigler (1988) Nature 333, 869-871; ²Rose et al. (1973) Nature New Biol. 245, 133-137; ³Marmorstein et al. (1987) J. Biol. Chem. 262, 4922-4927; ⁴Lawson et al. (1988) Proteins 3, 18-31.

This work is supported by the Office of Health and Environmental Research of the D.O.E.

X12C

"Structural Basis of Latency in Plasminogen Activator Inhibitor-1". James Mottensen, Anne Strand, Jindrich Symersky, Robert M. Sweet, Dennis E. Danley, Kieran F. Geoghegan, Robert D. Gerard, and Elizabeth J. Goldsmith. The Departments of Biochemistry and Internal Medicine* The University of Texas Southwestern Medical Center 5323 Harry Hines Blvd. Dallas, Texas 75235-9050, *Department of Biology Brookhaven National Laboratory, Upton, New York 11973, †Molecular Genetics Research Department, Central Research Division, Pfizer, Inc., Eastern Point Road Groton, Connecticut 06340

The structure of intact, latent human plasminogen activator inhibitor-1 (PAI-1) has been solved by single crystal x-ray diffraction to 2.6 Å resolution. PAI-1, the fast acting inhibitor of tissue plasminogen activator and urokinase, is a member of the serpin family of protease inhibitors. Serpins normally form very slow-dissociating complexes with their target protease, but are cleaved on dissociation and fold into a highly stable inactive state in which the residues that flank the scissile bond (P1 and P1') are separated by ~70 Å. PAI-1 also spontaneously folds into a stable, ¹⁰ inactive state in the absence of cleavage; this state has been termed "latent" because inhibitory activity can be restored through denaturation and renaturation. The three-dimensional structure of latent PAI-1 reveals that residues on the N-terminal side of the primary recognition site are inserted as a central strand of the largest β-sheet, in positions similar to the corresponding residues in the cleaved form of the serpin α1-proteinase inhibitor (α1-PI). Residues C-terminal to the recognition site occupy positions on the surface of the molecule distinct from those of the corresponding residues in cleaved serpins or in the intact, inactive, serpin homolog ovalbumin and its cleavage product, plakalbumin. We propose that the structure of latent PAI-1 is similar to one formed after cleavage in other serpins, and that the latency transition may represent a regulatory slow-folding step which, under normal physiological conditions, is kinetically inaccessible to other intact serpins.

THREE-DIMENSIONAL STRUCTURE OF cAMP-DEPENDENT PROTEIN KINASE

J.W. Pflugrath, J. Kuret, X. Cheng (CSHL) and R.M. Sweet (NSLS)

Protein kinases are important and conserved components of many regulatory pathways because they can integrate several input signals and coordinate a response through the phosphorylation of a limited number of substrate L-proteins. The three-dimensional crystal structure of the cAMP-dependent protein kinase will let us define the structural basis of enzyme/substrate recognition and assess the implications of this information for the protein kinase family in general.

We have crystallized the cAMP-dependent protein kinase catalytic subunit from *Saccharomyces cerevisiae*. The kinase catalytic subunit crystallizes in space group P6₃22 with cell dimensions $a = b = 61 \text{ \AA}$, $c = 320 \text{ \AA}$. We have grown crystals as large as 0.6 by 0.6 by 1.5 mm which have diffracted beyond $2.5 \text{ \AA}$ resolution at the National Synchrotron Light Source.

We have collected partial X-ray diffraction data in our laboratory with a FAST area detector and rotating anode generator by rotating the crystals for 40° around the a axis with c^* initially oriented toward the beam at a crystal to detector distance of 70 mm . Initial screening in this way yielded a dataset from p -chloromercuriphenyl sulfonate (PCMBS) soaked crystals which showed a single mercury site in both isomorphous difference Patterson maps and in anomalous Patterson maps. This derivative with its anomalous scattering constitutes the basis of our approach to solving the phase problem. In order to obtain complete datasets with reflections separated along l when c^* is oriented near the plane of the detector, we would need to use better optics with a relatively long crystal to detector distance of 320 mm with $\text{CuK}\alpha$ radiation.

To overcome the limitations of our laboratory X-ray source, we used the facilities at beamline X12-C at the Brookhaven National Laboratory National Synchrotron Light Source. The optics and intense tunable radiation of X12-C let us use a wavelength of $0.95 \text{ \AA}$ and a crystal to detector distance of 360 mm to separate the reflections, minimize air absorption and enhance the anomalous signal from the mercury atoms. Since a complete diffraction dataset cannot be obtained from individual crystals due to thermal and radiation damage, we have collected data from over 170 different crystals. Sixteen native crystals produced 9,220 unique reflections from 59,181 observations with an overall Rmerge of 4.7% and a completeness of 97.3% in the 3.0 to $2.8 \text{ \AA}$ shell. More than 37,000 measurements from four kinase crystals soaked in PCMBS resulted in 9613 unique reflection between 20 and $2.5 \text{ \AA}$ with an overall Rmerge of 6.1% . The data is 93.5% complete in the 3.0 - $2.8 \text{ \AA}$ shell. Data collected from numerous other crystals soaked in different heavy atom compounds did not reveal substitution.

Single isomorphous replacement with anomalous scattering refinement of the PCMBS data gave us a trial set of phases which were then used to calculate a $3.2 \text{ \AA}$ resolution electron density map. Regions of the map could be interpreted as α helices. Experiments are underway to improve the interpretability of the map. These include (1) new crystal harvesting conditions which require that all diffraction data be re-collected, (2) additional heavy atom soaks, (3) site-directed mutations in the protein sequence to introduce cysteine residues followed by purification, crystallization and soaking in mercury compounds, and (4) multiwavelength anomalous dispersion experiments with the PCMBS derivative.

This work is supported by NIH grants GM 44806 (J.K. & J.W.P.) and GM 48216 (J.K.).

CRYSTAL STRUCTURE DETERMINATION OF RIBOSOMAL PROTEIN S5

V. Ramakrishnan, R.M. Sweet (Biology, BNL) and S. White (Duke U).

The ribosome is the organelle on which the genetic code is translated into functional protein. It consists of two subunits, a large and a small, each of which consists of RNA and protein components. There is very little knowledge of the high resolution structure of components of the ribosome, and none of any of the components of the small subunit. We have cloned the gene for ribosomal protein S5 from the thermophilic bacterium *B. stearothermophilus*, and have over-expressed it in *E. coli*. The recombinant protein crystallizes isomorphously with the native. Its space group is $P3_121$, and the crystals diffract to $2 \text{ \AA}$ resolution. We used a single gold derivative to phase the X-ray diffraction data. Data were collected at a wavelength just below the L_{III} edge of gold in order to maximize anomalous differences within Bijvoet pairs. Data were collected by rotating the crystal about the a axis, so that anomalous pairs would be measured nearly simultaneously on the FAST area detector. The quality of the anomalous scattering measurements can be seen in a Harker section of an anomalous Patterson function of the data (Fig 1). The data were phased using isomorphous difference and anomalous scattering. The final phasing calculations gave an R_{calc} of 0.47 and an R_{meas} of 0.10 , with a mean figure of merit of 0.76 to $3 \text{ \AA}$ resolution. The data were about 90% complete when only those reflections that had both an anomalous and isomorphous difference contribution were included. The map generated by these calculations was subjected to several rounds of solvent flattening, which increased the figure of merit to 0.89 . A $10 \text{ \AA}$ section of this map is shown in Fig 2. The $3 \text{ \AA}$ map shows clear secondary structure elements, including several α helices and strands of β sheets. Interpretation of the electron density is under way. This work was supported by a grant from the National Institutes of Health (5 R01 GM42796-03) and by the Office of Health and Environmental Research of the U.S. Department of Energy.

Fig.1

Fig.2

STRUCTURE OF THE GLOBULAR DOMAIN OF HISTONE H5

V. Ramakrishnan, V. Graziano and R.M. Sweet (Biology, BNL)

The formation of chromatin higher-order structure requires the presence of a protein, histone H1 (or its analogue H5 in avian erythrocytes). This protein binds to the nucleosome and is essential for their organization into the 30 nm filament, the predominant form of chromatin in the cell. H5 consists of a highly conserved central globular domain flanked by extended arms. Using recombinant DNA methods, we have expressed the gene for the globular domain, and crystallized the resulting protein. The crystals diffract to about 2 Å resolution. Initial attempts to obtain heavy atom derivatives were unsuccessful. We therefore tried to obtain phases using the multi-wavelength anomalous dispersion method (MAD). Crystals were obtained of the protein in which the internal methionine had been replaced by seleno-methionine. X-ray diffraction data were collected at three wavelengths, two of which were near a Se absorption edge at 0.98 Å, and the third was at a remote point, 0.915 Å.

Initial analysis revealed peaks in the anomalous Patterson function, which also helped to identify the space group as $P2_12_12$. A solution to the Patterson was obtained, but the initial map looked uninterpretable. We are in the process of reanalyzing the data to obtain a better map.

This work was supported by a grant from the NIH (5 R01 GM42796-03) and the Office of Health and Environmental Research of the U.S. Department of Energy.

CRYSTALLOGRAPHIC STUDIES OF COMPLEXES OF BOVINE SOMATOTROPIN WITH A SOLUBLE FRAGMENT OF ITS RECEPTOR, RENIN AND EPSP SYNTHASE: Huey-Sheng Shieh, Roderick A. Siegelman, Anna M. Stevens and William C. Stallings (Monsanto Corporate Research, St. Louis), Robert M. Sweet (NSLS).

Growth hormone molecules (somatotropins) mediate various biological activities. The structure of porcine somatotropin has been determined at high resolution, but information from the structure on how the hormone functions has not been definitive. We have grown small crystals of the homologous bovine somatotropin complexed with the binding domain of its receptor. Visualization of the complex structure may allow us to understand the specificity of the interactions and of the activities elicited different somatotropins. The crystals, less than 100 µm in any dimension, were grown from polyethylene glycol, but it was not possible to characterize these crystals using conventional x-ray sources. The crystals were characterized at NSLS beamline X12C using the MADNES suite of programs for autoindexing. Lattice parameters are space group C2 with $a = 117.4$, $b = 143.0$, $c = 67.0$ Å, $\beta = 108.7^\circ$. If a 1:1 complex is assumed, with two complexes in the asymmetric unit ($Z=8$), the value of the Mathews coefficient, V_M , is $2.7 \text{ Å}^3/\text{D}$ and the estimated solvent content is 54% of the lattice volume. The crystals showed order to almost 3-3.5 Å resolution at X12C and data sets from three crystals were collected at the facility. The data were scaled and merged and, after deletion of the weakest reflections, they yielded a set of 7,101 unique intensities with $R_{\text{mer}} = 0.042$; the data are complete at 4.5 Å resolution. A preliminary self-rotation function (MERLOT) did not indicate the relative orientation of two unique complexes.

Renin is an enzyme which cleaves the decapeptide, angiotensin I, from the amino terminus of angiotensinogen. Angiotensin I is subsequently clipped to the octapeptide, angiotensin II which is a potent vasoconstrictor. Therefore, inhibition of renin activity has therapeutic potential in the control of elevated blood pressure. Data on a deglycosylated form of human renin complexed with two inhibitors were collected from four crystals to a nominal resolution of 2.5 Å. Reduction of the intensities to structure factor amplitudes has been a time-consuming process which continues.

5-*enol*-pyruvyl shikimate-3-phosphate synthase (EPSP synthase) is an enzyme on the pathway toward the biosynthesis of aromatic amino acids in plants, bacteria and fungi; it is the target of the non-selective herbicide, glyphosate. Small crystals of a putative ternary complex of the enzyme with substrate, shikimate-3-phosphate and inhibitor, glyphosate were characterized at the facility. The space group is $P2_1$ and the monoclinic lattice parameters are: $a = 65.4$, $b = 85.1$, $c = 79.6$ Å and $\beta = 97.1^\circ$. Assuming $Z = 4$, the Mathews coefficient, V_M , is $2.5 \text{ Å}^3/\text{D}$ and there are two complexes in the asymmetric unit of the lattice. The estimated solvent content is 50%.

MONSANTO CORPORATE RESEARCH

X12C/X25
A COMPARISON OF HIGH-RESOLUTION TRYPSIN CRYSTAL STRUCTURES OBTAINED BY WHITE-BEAM LAUE AND MONOCHROMATIC X-RAY DIFFRACTION USING SYNCHROTRON RADIATION.
P.T.Singer (BNL Biology), R.P.Cary (SUNY Downstate), L.Berman (NSLS), and R.M.Sweet (BNL Biology)

Crystals of the benzamidine-trypsin complex were subjected to X-ray diffraction analysis using both monochromatic and white-beam synchrotron radiation. Diffraction data were obtained over the range of 5.0 to 1.5 Å using the monochromatic beam, and 5.0 to 1.8 Å using the Laue method. The orthorhombic crystals belonged to space group P2₁2₁2₁, with unit cell dimensions a = 63.74, b = 63.54, c = 68.93 Å. Monochromatic reflections were collected on a FAST (Enraf Nonius) area detector. Observed reflections were fitted to an existing model of guanidinobenzoyl-trypsin and refined using the least-squares refinement program, TNT. Two separate sets of crystals, treated at either pH 5.5 or 8.9, were refined to crystallographic R-factors of 19.3% and 20.5%, respectively. White-beam, or "Laue," diffraction data were collected from a single crystal mounted in a flow cell using Kodak DEF film; typical exposure times were 50 ms. Data were collected with the crystal alternately equilibrated against buffers of pH 5.5 and 8.9. The structural models were refined to crystallographic R-factors of 17.2% and 18.5%, for benzamidine-trypsin complex at the low and high pH, respectively.

Comparison of the molecular models of these four structures revealed major differences in the definition of the hydration shell around the protein. While the Laue diffraction data showed the presence of 147 and 95 water molecules in the pH 5.5 and 8.9 structures, respectively, the high-resolution monochromatic data indicated the presence of 182 and 129 water molecules in the putatively identical structures, respectively. Many of the additional water molecules indicated exclusively in the monochromatic data sets were part of a second and third hydration shell. Protein atoms for all four data sets were virtually identical except for the Ser195 side chain in the pH 8.9 Laue structure, which is swiveled away from His57, resulting in the loss of the H-bond between O and NE2 of the histidine. The benzamidine ligand is better resolved in the monochromatic model, probably due to the more extensive data collection in the monochromatic series compared to the Laue data.

The two methods of data collection yield virtually identical models of the benzamidine-trypsin complex. However, the white-beam data seems to result in better maps for sidechain atoms, while the backbone atoms are more clearly defined in maps derived from the monochromatic data, presumably due to the systematic absence of low resolution data in the "Laue" data sets.

This work was sponsored by the U.S. DOE. PTS was supported by an Alexander Hollaender Postdoctoral Fellowship awarded by OHER.

Beam Line X12C

Multiwavelength Data Collection on BamHI

T. Strzelecka (Columbia U.), L. Dornier (NEB), I. Schildkraut (NEB) and A. Aggarwal (Columbia U.)

BamHI is a type II restriction enzyme that specifically recognizes the DNA sequence 5'-GATC-3', and in the presence of Mg²⁺ cleaves it as indicated. As a model system for a better understanding of protein-DNA interactions, we have prepared crystals of native BamHI that diffract to 1.5Å, and crystals of BamHI with its 5 methionine residues replaced by selenomethionine (BamHI-se) residues. A multiwavelength anomalous diffraction (MAD) experiment performed on BamHI-se crystals has the potential to provide the necessary phase information for a complete three dimensional structure determination of this enzyme.

In collaboration with Dr. R. Sweet (Brookhaven), we collected data from BamHI-se crystals at wavelengths corresponding to the peak, the inflection point, and a remote point in the absorption spectra of Se. The data was collected from a total of 23 crystals, spread over 3 visits (each visit ~ 4 - 6 days). Successful application of the MAD method requires precision in data measurement and in wavelength selection. After careful pruning of the data, we retained data from 16 crystals for MAD analysis. From these, we have managed to extract phases for ~ 45% of the possible reflections at 3.0Å. Statistics for these phases appear reasonable. We are in the process of combining these phases with phases derived from other sources for completion of our structural analysis.

STRUCTURE DETERMINATION OF STAPHYLOCOCCAL ENTEROTOXIN B

S. Swaminathan, W. Furey, J. Pletcher and M. Sax,
 Biocrystallography Laboratory, VA Medical Center, University
 Dr. C., Pittsburgh, PA 15240 and R.M. Sweet, NSLS.

Staphylococcal enterotoxins are responsible for the most common food-borne illnesses in the United States. These toxins are known to induce emesis, diarrhea and mitogenesis and it is believed that the toxic, mitogenic and antigenic activities are all effected from different sites on the toxin molecule. In addition, Staphylococcal enterotoxin B (SEB) is known to be a powerful T cell stimulant in mouse and man. It has been described as a "Superantigen" from which insight into the mechanisms of self-tolerance may be obtained. Currently, three-dimensional structure information is not available for any Staphylococcal enterotoxin or superantigen. We have crystallized SEB and are currently determining its 3-D structure.

X-ray data were collected to at least 2.5Å resolution on a Xentronics area detector with a rotating anode source for both native and derivative crystals, but the anomalous scattering data for the platinum derivative was usable only up to 3.5Å resolution. Since the wavelength of the synchrotron radiation could be tuned to be near the absorption of the heavy metal, thereby enhancing the anomalous signal, data were recollected at the NSLS facility at Brookhaven National Laboratory. The synchrotron data extended up to 2.2Å resolution and the anomalous Patterson map was good up to the limit of data collected. Currently, we are trying to use this data in the crystal structure determination.

This work is supported by the VA Medical Center, program #185-12-8499 and by the US Army Medical R&D Command, #87PP7852. The synchrotron data collection was supported by the NSLS facility at Brookhaven National Laboratory.

DIFFUSE X-RAY SCATTERING FROM THE ALLOY Fe-25%Ni-1%C

B.D. Butler (Northwestern U.), P.R. Zschack (ORNL), H. Yalcinkaya (Northwestern U.), J.B. Cohen (Northwestern U.)

Diffuse x-ray scattering on the alloy Fe-24.8%Ni-0.88%C was measured in absolute units in a full volume of reciprocal space. The measurements were made using a bent graphite analyzing monochromator and position sensitive detector. This alloy is austenitic (FCC) at room temperature and transforms, displacively, to martensite (BCT) below $\sim -40^{\circ}\text{C}$.

Motivation for this study is to determine the state of ordering/clustering of the C atoms on the octahedral interstitial sublattice in the austenitic phase. This will be inherited by the product martensite phase and so represents the state of C ordering/clustering at the beginning of the aging process of martensite. The degree of short range order on the Fe-Ni lattice will also be determined from the data.

Measurements were made at two energies. The first at 8800 eV which is above the K-absorption edges of both Fe and Ni. At this energy the dispersion correction for Ni is slightly higher than that of Fe and so the scattering contrast between these elements is minimized. The data will be interpreted as if from a binary "Fe"-C alloy using the method of Georgopoulos-Cohen instead of as a ternary Fe-Ni-C alloy, greatly simplifying the analysis. Measurements were also made just below the Fe K-edge where the contrast between Fe and Ni is maximized. This experiment will be used to determine the degree of short range order on the Fe-Ni sublattice. Observations of the raw diffuse data show no apparent diffuse scattering around the superlattice positions for ordering as would be found in the alloy Ni₃Fe. A quantitative analysis of the degree of ordering on the Fe-Ni sublattice is currently underway.

The very weak scattering associated with C along with the need to tune the energy required that these measurements be made at a synchrotron x-ray source. The excellent x-ray flux and unique detection system make this beam line particularly well suited to this study.

This work is sponsored by the National Science Foundation under Grant No. DMR-8920567
 Beam Time Provided on X14 by the Div. of Materials and Chemical Sciences. D.O.E.

STUDIES OF THE STRUCTURE OF AMORPHOUS MELANINS

J. Cheng, M. Eisner and S. C. Moss (U. of Houston); P. Zschack (ORNL)

Melanin is a ubiquitous biological pigment which may be synthesized from various precursors (L-dopa, tyrosine) or extracted from natural sources (sepia melanin). Because it has never been crystallized, its structure must be studied in the amorphous state after complete removal of the adsorbed water. It is currently believed to be a randomly connected amorphous biopolymer with pronounced graphite-like layer features and extended intralayer linking of 6- and 5-membered rings cross-linked at oxygen and hydrogen bonded sites. Careful data collection requires an accurate estimate of absorption and the removal of Compton scattering which was accomplished here using the X14 parafocussing graphite analyzer/PSD detector. The corrected and normalized data below indicate a similarity, clearly between the synthetic forms and roughly with the natural sepia melanin. Transforms have been taken and structural modelling, for both Q- and r-space fits, is underway.

Research supported by the DOE, Division of Basic Energy Sciences on DE-FG05-87ER45325.

ISOTHERMAL THICKENING AND THINNING PROCESSES IN LOW MW PEO FRACTIONS OBSERVED FROM SYNCHROTRON SAXS

S. Z. D. Cheng, A.-Q. Zhang, J. S. Barley, J.-H. Chen (Akron U.) A. Hubenschuss, and P. R. Zschack (DRNL)

Based on synchrotron time-resolved small angle X-ray scattering (SAXS) and differential scanning calorimetry (DSC) experiments, a nonintegral folding chain (NIF) crystal can be found in low molecular weight poly(ethylene oxide) (PEO) fractions. This crystal grows first as a transient state, and integral folding chain (IF) crystals form later through an isothermal thickening or thinning process. In a PEO fraction with MW=4250, it is observed that either isothermal thickening to the IF (n=0) crystal or thinning to the IF (n=1) crystal occurs depending upon the thermodynamic stabilities of the NIF crystal. At low crystallization temperatures, the main process is isothermal thinning; at intermediate temperatures, both thickening and thinning processes are observed; only isothermal thickening can be seen at high temperatures. This observation indicates that the NIF crystal exists as an initial step of the crystal growth, even though its thermodynamic stability is greater than that of the IF (n=1) crystal. Furthermore, molecular weight dependence of the isothermal thickening and thinning processes have also been investigated. This study focuses on: (1) Can we still identify isothermal thickening or thinning during the transition of NIF to IF crystals with increased molecular weight? (2) Where is the critical molecular weight for PEO fractions above which the thinning is forbidden? (3) How does molecular weight influence the transition kinetics?

Acknowledgement

This research was supported by Exxon Education Foundation. Research carried out (in part) at the NSLS at BNL, which is supported by the US DOE, DMS and MCS.

POWDER DIFFRACTION STUDY OF MAGNETOELASTIC STRAIN BROADENING IN NANOCRYSTALLINE DYSPROSIUM

M.R. Fitzsimmons (LANL), J.A. Eastman (ANL), and L. Thompson (ANL)

Nanocrystalline materials have been shown to possess many unique physical properties, which are attributed to the presence of the large volume fraction of interfacial regions in the material. Some of the unique properties include, increased values of thermal expansion [1], heat capacity [2] and fracture stress [3] compared to the values found for bulk materials. In order to put the properties of nanocrystalline materials to practical use, a better understanding of their microstructure and its influence, if any, on the physical properties of these materials is required. The motivation for this experiment is to determine whether magnetoelastic effects can alter the strain field within a ferromagnetic nanocrystalline material by observing changes of the Bragg reflection profiles as the temperature of the material is increased above its Curie temperature, T_c . Once the ferromagnet becomes nonmagnetic above T_c , internal stresses caused by magnetostriction of the crystallites disappear; therefore, a change of the strain fields in a nanocrystalline material is expected.

Powder diffraction data from both compacted and uncompacted nanocrystalline samples, as well as coarse-grained control specimens, were obtained in the presence of a 2kG magnetic field at approximately two dozen different temperatures ranging from 40K to 298K. The intensity data was fitted to a profile consisting of 50 pseudo-Voigt peaks and a polynomial background. The diffraction patterns observed from the specimens was consistent with the hcp crystal structure at all temperatures. The lattice parameter parallel to the basal plane, a_0 , and normal to the basal plane, c_0 , of a compacted nanocrystalline dysprosium specimen for several temperatures is shown in the figures below. From the linear increase of a_0 with temperature, the thermal expansion coefficient of compacted nanocrystalline Dy parallel to the basal plane is $36 \times 10^{-6} \text{K}^{-1}$ in the measured temperature range. The abrupt variation of the c_0 parameter at approximately 98K and 197K suggests the occurrence of second order phase transformations—the ferromagnetic and antiferromagnetic transformations. The literature values of the Curie and Néel temperatures for single crystals of Dy are 86K and 179K, respectively [4].

- [1] H.J. Klam, H. Hahn and H. Gleiter, *Acta Metall.*, **35**, 2101 (1987).
- [2] R. Birringer, U. Herr and H. Gleiter, *Trans. of Jap. Inst. of Metals*, (1986).
- [3] H.G.F. Wilsdorf, *J. Appl. Phys.*, **48**, 478 (1977).
- [4] F.J. Darnell and E.P. Moore, *J. of Appl. Phys.*, **34**, 1337 (1963).

X-ray Scattering Study of Ag/Si(111) Buried Interface Structures

Hawoong Hong, R. D. Aburano, D.-S. Lin, Haydn Chen, T.-C. Chiang (U. of Illinois-Urbana Champaign) P. Zschack and E. D. Specht (ORNL)

Various interface structures formed between Si(111) and a 260Å of Ag overlayer are investigated by grazing-incidence x-ray diffraction. The samples were prepared in a MBE chamber and brought to NSLS in a desiccator. During the x-ray experiment, the samples were kept in an evacuated Be dome. When the 7x7 reconstructed Si(111) surface is covered with a room-temperature deposited Ag film, diffraction spots from the 7x7 structure are observed. (Fig. 1) When this interface is annealed, the 7x7 fractional order peaks disappear at 250°C and the interface becomes 1x1. (Fig. 2) The $\sqrt{3} \times \sqrt{3}$ Ag/Si(111) is formed by annealing 2 monolayers of Ag on Si(111). We observe the $\sqrt{3} \times \sqrt{3}$ reconstruction to disappear when we deposited a thick Ag film. From rod scans the 7x7 interface appears to remain sharp during annealing. This is contrasted by the $\sqrt{3} \times \sqrt{3}$ interface which is very rough.

Fig. 1 Rodding curves for in-plane scattering from Ag thick film-Si(111) 7x7

Fig. 2 Evolution of the (6/7, 0) peak after annealing to the temperature indicated

This work was supported by the DOE Grant #DEFG02-91ER45439 and #DE-AC05-84OR21400. Acknowledgment is also made to the State of Illinois for a DCCA Technology Challenge Grant.

INELASTIC X-RAY SCATTERING SPECTROSCOPY IN LITHIUM HYDRIDE

E.D. Isaacs, (AT&T); G.E. Ice (ORNL); P. Zschack (ORAU).

Inelastic x-ray scattering measurements in LiH using a unique, high-resolution spectrometer have been successfully performed at the National Synchrotron Light Source bending magnet beamline X14. Energy resolution from 1.5 to 6 eV with momentum transfers in the range $0.4 - 6 \text{ \AA}^{-1}$ have been achieved. Either a spherically bent (not pressed) Graphite (006) mosaic crystal or a spherically bent (plastically deformed) LiF (004) mosaic crystal operating near backscattering with $E_{\text{inc}} = 6 \text{ keV}$ was used for energy dispersion of the scattered beam. For small momentum transfers ($q = 0.8 \text{ \AA}^{-1}$) and about 3 eV resolution in LiH we have observed three relatively sharp ($< 3 \text{ eV}$) band features at around 15 eV associated with the Li^+ (2s) and H^- (1s) electrons, which were previously unobserved (see figure). In addition, we observed the $\text{Li} (1s^2)$ core state at an energy shift of about 55 eV with an as yet unexplained exciton-like doublet feature at a slightly lower energy. At large momentum transfers ($q > 4 \text{ \AA}^{-1}$) the expected single-particle peak was observed.

Our success in LiH has opened up the opportunity to study many other interesting systems. For example, we have investigated the inelastic spectrum in single crystals of aluminum, beryllium and powder samples of undoped Bucky balls.

Figure. Inelastic x-ray scattering spectrum of LiH. $E_{\text{inc}} = 6.3 \text{ keV}$ and $q = 0.8 \text{ \AA}^{-1}$. Note the three features; the elastic peak, the band states at around 15 eV and the $\text{Li} (1s^2)$ core states near 55 eV.

CRYSTALLOGRAPHIC STUDY OF A SINGLE QUASICRYSTAL OF Al-Cu-Fe

X. B. Kan and S. C. Moss (U. of Houston); J. L. Robertson (NIST); T. Ishimasa and M. Mori (U. of Nagoya); D. Gratias (Vfry); V. Eiser (Cornell); P. Zschack (ORNL)

Large single quasicrystalline grains, both rapidly cooled from the melt (Gratias group) and annealed on cooling at -830°C (Ishimasa and Mori), were ground approximately into spheres (they are quite brittle) for crystallography with a resultant diameter of $\sim 0.2 \mu\text{m}$. A sample was then attached to the end of a glass rod inside an evacuated Be dome to remove air scattering at low angles where significant (small $q_{\perp}$) peaks occur. Using $\lambda = 1 \text{ \AA}$ more than 1300 (inequivalent) reflections were collected and are currently being analyzed. An important aspect of the data collection was the simultaneous measurement of the (Cu and Fe) fluorescence channel which allows us to correct for both the non-sphericity of the sample and the hot spot variations in the incident beam. The data collection was done in hierarchies of $q_{\perp}$ (the perpendicular momentum transfer) so as to cover increasingly weaker intensities in the densely populated reciprocal space. A very interesting aspect of our Bragg scans is shown below in which the radial (θ - 2θ) scan in (a) reveals a nearly perfect Gaussian profile (solid curve) while the omega (ω) scan (b) is essentially a pure Lorentzian (solid curve). The q width (FWHM) of the radial scan is $0.009 \text{ \AA}^{-1}$ which is close to instrumental and is not particularly dependent on $q_{\perp}$. The FWHM in omega is 0.06° . The unusual omega lineshape must be associated with a true mosaic spread as it is essentially constant in angle at all Bragg peaks, independent of $q_{\perp}$. Further work has shown that the Lorentzian profile has been substantially enhanced by the mechanical grinding.

Research sponsored by the NSF on DMR-8903339 (Houston) and the David and Lucille Packard Foundation FDN89-1606 (Cornell).

X-RAY DETERMINATION OF SITE OCCUPATION PARAMETERS IN ORDERED TERNARIES Cu(Au₂M₁₋₂)M-Ni-Pd: ANOMALOUS SCATTERING AND RIETVELD REFINEMENT*

R. Kumar, C. J. Sparks, E. D. Specht, P. Zschack, G. E. Ice (ORNL), T. Shiraishi (Kyushu U., Japan), and K. Hisatsune (Nagasaki U., Japan)

X-ray scattering data obtained for multiple wavelengths with synchrotron radiation were analyzed by the Rietveld [1,2] method to determine Ni and Pd distributions on the Cu(000) and Au(111) sites in the CuAuI tetragonal P4/mmm structure. Figure 1 depicts the structure of these alloys. Powdered alloys of CuAu₂M₁₋₂ containing 6 at. % Ni or 10 and 25 at. % Pd were processed to obtain maximum ordering. Analysis showed that Ni is predominantly found on the Cu site and most all the Pd is found on the Au site. The results for Ni are given in Table 1. This use of variable wavelength x rays with simultaneous refinement of the corresponding data is capable of distinguishing site occupations even between two elements of almost equal scattering factor as, for example, Cu and Ni atoms in this investigation. Chemical phase stability is related to the site occupation parameters.

Fig. 1. The atom positions of the CuAuI tetragonal superlattice with the primitive unit cell, having a Cu atom at (0,0,0) and a Au atom at (1/2, 1/2, 1/2).

Table 1. Site occupation distribution in Cu₉₉Au₁Ni₁₂

Crystallographic site	Free refinement	χ^2	Constrained refinement	χ^2
Cu(000)	0.983 (12)	0.0077	0.986 (8)	0.0082
Au(000)	0.008 (1)	0.0077	-0.004 (1)	0.0082
Ni(000)	0.103 (13)	0.0077	0.118 (8)	0.0082
Cu(111)	0.135 (12)	0.0077	0.112 (8)	0.0082
Au(111)	0.874 (1)	0.0077	0.986 (1)	0.0082
Ni(111)	0.018 (13)	0.0077	0.002 (8)	0.0082
$\chi^2 = (R_p/R_{wp})^2$	2.25		2.24	
R_{wp}	13.05%		13.06%	

$8\sigma^2 u_{iso}$ = isotropic thermal correction factor.

$R_p = \sum |F_o| - \sum |F_c|$ = Weighted profile R-factor, and $M = \sum (F_o - F_c)^2$.

*Research managed by the Division of Materials Sciences, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

¹H. M. Rietveld, J. Appl. Cryst., 2, 65 (1969).

²A. C. Larson and R. B. Von Dreele, Los Alamos National Laboratory Report No. LAUR 86-748 1989.

DIRECT DETERMINATION OF PRECIPITATE STRAIN AND SIZES USING HIGH-ORDER X-RAY DIFFUSE SCATTERING*

B. C. Larson, M. D. Galloway, G. E. Ice and P. Zschack (ORNL)

High-order diffuse-scattering measurements on coherent precipitates in (aged) Cu 1% Co crystals have been combined with numerically calculated diffuse scattering cross-section calculations to determine coherency strains and the sizes of the precipitates. Scattering measurements were made near the (8 0 0), (10 0 0), and (12 0 0) Bragg reflections using synchrotron X rays at the ORNL-UIUC beam line at the National Synchrotron Light Source, and calculations were made using anisotropic elasticity. The figure shows diffuse scattering measured near the (8 0 0) reciprocal lattice point in which direct Bragg scattering from the precipitates is present in the form of a completely resolved subsidiary peak. The separation of the "precipitate" scattering peak from the "host lattice" Bragg peak provides a direct measure of the lattice parameter of the precipitates relative to the host lattice. In addition, the diffraction width of the precipitate scattering peak provides a direct measure of the characteristic size of the precipitates when analyzed using the numerical calculations.

The strain within the cobalt precipitates (relative to the host lattice) was found to be -1.4%, in good agreement with the -1.42% strain inferred indirectly from previous low-order diffuse scattering measurements. A value of -1.41% was calculated using elasticity and the -1.8% unconstrained lattice parameter difference between fcc cobalt and copper. The characteristic radius of the precipitates was determined to be 68 Å; however, the use of the precipitate scattering alone does not provide detailed information on the distribution of sizes.

Fig. 1. Diffuse scattering measured in the radial direction near the (8 0 0) Bragg reflection on aged (containing precipitates) and quenched (no precipitates) Cu 1% Co.

*Research sponsored by the Division of Materials Sciences, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

MAGNETIC X-RAY SCATTERING IN DY/LU SUPERLATTICES

M.B. Salamon, R.S. Beach, B.A. Everitt, K. Pettit, A. Matheny, C.P. Flynn;
U. of Illinois at Urbana-Champaign

The aim of our work at the Light Source is to study magnetic scattering in Dy/Lu superlattices. To our run of July 11, 1991 we brought analyzer crystals chosen to reflect X-rays at $2\theta=90^\circ$ on the Dy and Lu L_{III} absorption edges, and arranged to borrow a polarization analyzer from the BNL X-ray scattering group. We lost time to hardware problems, but with the analyzer set to receive $\sigma-\pi$ scattering we believe we observed magnetically scattered photons from the superlattice Dy layers. The real difficulty in the experiment is that, in a superlattice, magnetic intensity necessarily sits on a background of charge scattering. Because the charge scattering cross section is $10^4 \times$ the magnetic cross section, even on resonance, the polarization must be nearly in the same ratio to effect separation of the two. In the future we will incorporate a pre-polarizer which makes use of the Bormann effect; we conducted brief but promising tests of a prototype during this cycle.

We devoted an additional segment of time to investigating the effect of a magnetic field on the (previously detected) splitting of peaks in the basal plane. This splitting seems to vanish in a 600 G field. The system is epitaxial; exactly how the superlattice distorts to yield this surprising result has not yet been resolved.

This work was supported in part by NSF Grant DMR 88-20888 and in part by a State of Illinois DCCA grant.

X-RAY STUDIES OF DYSPROSIUM/LUTETIUM SUPERLATTICES

M.B. Salamon, R.S. Beach, C.P. Flynn, A. Matheny (U. of Illinois-Urbana) and J. Borchers (NIST)

This brief (4-day) preliminary experiment on superlattices had three goals: 1) to use resonant scattering to increase the contrast between Dy and Lu for analysis of the superlattice structure; 2) to search for magnetic scattering (already well characterized by neutron diffraction); and 3) to search for an orthorhombic distortion accompanying the low-temperature development of ferromagnetism.

Goal 1) was readily met (see figure). The sample gave easily measured superlattice sidebands to seventh order, and beyond. The quality of the superlattice, however, made it very difficult to achieve goal 2) inasmuch as strong (third order) superlattice charge scattering overwhelms the magnetic scattering, even when resonance absorption at the Dy L_{III} edge was used with polarization analysis. The analyzer used was not optimized for the Dy edge. Further, the large acceptance angle of X-14 admits a significant amount of elliptically polarized light, giving strong $\sigma-\pi$ charge scattering. Future experiments will use superlattices of smaller period to separate magnetic and charge scattering. A dedicated polarization analyzer/detector will be constructed.

Preliminary data were obtained on the development of orthorhombic distortion at low temperature. Epitaxial constraints prevent the free distortion of the crystal shape. It appears that the sample breaks up into small, orthorhombic domains (as evidenced by the increased width of the split (1010) reflections. This work will be continued on a future experimental run.

This experiment marked one of the first uses of the X-14 Displex, requiring considerable setup time. Modifications of the sample holder and vacuum shroud will make such experiments simpler on future runs.

Fig. 1 Satellites of the (0004) Bragg peak of a Dy/Lu superlattice, taken at the L_{III} edge of Dy.

Supported by NSF Grant DMR-8820888 and State of Illinois DCCA.

ANOMALOUS BEHAVIOR OF DIFFRACTION PEAK WIDTHS IN LIPID MONOLAYERS

M.C. SHIH, T.M. BOHANON, J.M. MIKURUT and P. DUTTA, Northwestern University And P. ZSCHACK, Oak Ridge National Laboratory

In 1942, Copeland, Harkins and Boyd [1] claimed, on the basis of viscosity data, that there existed a 'superliquid' phase in Langmuir monolayers. In this phase (or in this region of the phase diagram) the viscosity was anomalously lower than at lower pressures or at higher temperatures.

We have now explored this region in heneicosanoic acid by performing in-plane and 'rod' diffraction scans as a function of temperature and pressure [2]. We find that the in-plane radial peak widths are broadest in the region of the Rotator-I phase; in other words, the peak widths anomalously become narrower if the pressure is decreased or the temperature is increased. This correlates with, and offers a microscopic basis for, the anomalous behavior of the viscosity. The analogous Rotator-I phase in lamellar bulk paraffins is also known to be poorly ordered.

FIG. 1. Horizontal-plane widths of Peak I (●) and Peak II (□), the two first-order diffraction peaks, as a function of temperature at 31 dynes/cm. (top), and as a function of pressure at 20°C (bottom). (The hexagonal Rotator-II phase has only one first-order peak.) The resolution function (FWHM $0.008\text{\AA}^{-1}$) has not been deconvoluted; the narrowest peaks in the above data are resolution-limited.

1 L.E. Copeland, W.D. Harkins and G.E. Boyd, *J. Chem. Phys.* 10, 357 (1942)

2 M.C. Shih, T.M. Bohannon, J.M. Mikrut, P. Zschack and P. Dutta, submitted to *Phys. Rev. A*
[Work supported by DOE grant no. DE-FG02-84ER45125]

EFFECT OF pH ON A FATTY ACID MONOLAYER WITH CALCIUM IONS IN THE SUBPHASE

M.C. SHIH, T.M. BOHANON, J.M. MIKURUT and P. DUTTA, Northwestern University And P. ZSCHACK, Oak Ridge National Laboratory

It is generally known that the presence of metal ions in the subphase 'condenses' a Langmuir monolayer, i.e. increases its density. This was studied in some detail by Spink and Sanders [1], who measured isotherms as a function of pH and found that the area/molecule at zero pressure decreased with increasing pH.

We have performed X-ray diffraction studies of heneicosanoic acid monolayers with calcium ions in the subphase [2]. Fig. 1 shows the location of typical diffraction peaks in terms of the in-plane and normal-to-plane momentum transfer vectors. It can be seen that the off-plane peak moves to higher K_{xy} and lower K_z as the pH is increased, which means that the area/molecule decreases (consistent with the isotherm data of Ref. 1) and also that the molecular tilt angle decreases. Moreover, the in-plane peak position remains essentially fixed; in other words, the pH-induced distortion of the lattice is primarily uniaxial.

- 1 J.A. Spink and J.V. Sanders, *Trans. Faraday Soc.* 51, 1154 (1955)
- 2 M.C. Shih, T.M. Bohannon, J.M. Mikrut, P. Zschack and P. Dutta, *J. Chem. Phys.* to be published

FIG. 1: Diffraction intensities, plotted as contours in the K_x - K_y plane, at zero pressure in the D phase at various pH values.

[Work supported by DOE grant no. DE-FG02-84ER45125]

HIGH-PRESSURE PHASES OF A LIPID MONOLAYER AND THEIR RELATIONSHIP TO ISOTHERM ANOMALIES

M.C. SHIH, T.M. BOHANON, J.M. MIKRUT and P. DUTTA, Northwestern Univ.

Langmuir monolayers of many fatty acids and alcohols show the same curious feature: within a narrow temperature range, the highest-pressure discontinuity changes from a kink to a flat section and then back to a kink (Fig. 1). Because of anomalous flow properties, this is often referred to as the 'superliquid' region.

We have studied monolayers of heneicosanoic acid in this region using grazing-incidence X-ray diffraction. Along a high-pressure isobar, between the known distorted-hexagonal and hexagonal phases, we see a slightly-distorted phase in a narrow temperature range (Fig. 2). These phases correspond to the 'crystal' (distorted), Rotator II (hexagonal) and Rotator I (intermediate) phases seen in paraffins; the transitions between these phases are also like those in paraffins, with the 'crystal' to Rotator I transition being first order and the Rotator I to Rotator II being either continuous or very weakly first order.

There is an unexpected correspondence between the high-pressure phases and the isotherm features: where 'crystal' or 'Rotator II' phases appear, the isotherm shows a kink, but where the 'Rotator I' phase appears, there is a flat section.

FIG. 1: Isotherms of heneicosanoic acid monolayers in the region where the highest-pressure transition goes from a kink to a flat section and back to a kink. (The lower-pressure discontinuity is a 'swirling' transition between tilted phases.)

FIG. 2: Bit and area/mol. as a function of temperature for a monolayer of heneicosanoic acid along a 31 dyn/cm isobar. The lines are guides to the eye.

TIME-DOMAIN SPECTRUM OF RESONANTLY FILTERED X-RAYS FROM A NUCLEAR BRAGG MONOCHROMATOR

J. Z. Tischler, B.C. Larson, P. Zschack (ORNL)

The use of a Mössbauer resonant beam in a scattering experiment requires a means of analyzing the scattered beam to determine changes in its energy spectrum. The simplest type of experiment, for instance, requires separating the elastic scattering (due to static structural variations) from x-rays scattered inelastically (by dynamical phenomena), where inelastic means an energy shift $\geq 10^{-7}$ eV. Here we show how a powdered filter can provide a convenient means for making such measurements.

The figure shows the measured time spectra of x-rays reflected from the 777 reflection (a magnetic superlattice reflection) of a mosaic $^{57}\text{Fe}_2\text{O}_3$ crystal with and without a powdered $^{57}\text{Fe}_2\text{O}_3$ absorber. The filtered data in the figure were normalized to remove the effects of non-resonant absorption in the filter. Note that the shapes of the two curves are quite different; the relative difference between filtered and unfiltered peaks varies from peak to peak, in particular note the peaks at 18 ns and 28 ns. Since the final time spectrum in a resonant fraction measurement must be a linear combination of the two curves in the figure, this simple filter should provide a convenient analyzer for measuring the resonant fraction in a beam scattered from some sample.

Research sponsored by the Division of Materials Sciences, U.S. Department of Energy, under contract DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc.

[Work supported by USDOE grant no. DE-FG02-84ER45125]

THERMAL EFFECTS ON SOLID NEON

C. T. Venkataraman, R. O. Simmons, M. A. Fradkin, I. Fujita
(University of Illinois, Urbana-Champaign)

In this experiment we are attempting to gain some understanding of thermal effects in solid neon. Experiments on solid neon have been previously performed by our group, using a lab x-ray source, to study the vacancy concentration by the simultaneous measurement of macroscopic and x-ray lattice expansions. The group has also been involved in the measurement of the single particle kinetic energy of the Ne atoms through neutron scattering. Our ultimate goal in doing further scattering experiments using synchrotron radiation is to relate this single particle kinetic energy to the x-ray Debye-Waller factor for neon. The realization of this goal is complicated by the fact that Ne is a highly anharmonic solid.

We prepared the sample in situ by cooling the condensed gas through its freezing point under its own vapor pressure. The cooling was achieved by mounting the beryllium sample cell to a Displex which in turn was mounted on the X-14a Huber diffractometer. We grew four samples, one with gradual cooling and three with rapid cooling, to examine the effect of the cooling rate on the crystal growth. However, in all four cases we obtained a polycrystalline sample composed of large crystallites.

Once the sample was grown, we performed several scans of various Bragg reflections concentrating primarily on the (111) and (220) which are the most intense. We took omega and two-theta scans of the peaks at several temperatures ranging from 12K to 18K, in the hope of characterizing the effect of temperature on the lattice parameter. Synchrotron radiation with its high energy resolution is well suited to observing temperature induced lattice parameter variations of the order of 10^{-4} nm which is the previously observed shift.

We are currently in the process of extracting the thermal diffuse scattering contribution from the Bragg peaks. Once the TDS contribution has been subtracted we will be able to examine the effect of temperature on the integrated Bragg intensity.

This work is supported by the State of Illinois under the grant SCCA 90-82136 and the US Department of Energy BES Materials Science under Contract No. DEFG02-91ER45439.

STUDY OF ALKOXIDE-DERIVED AMORPHOUS TiO_2 POWDER

Q. J. Wang and S. C. Moss (U. of Houston); M. Shalzi and A. M. Glaeser (UC Berkeley);
H. W. Zandbergen (Delft); P. Zschack (ORNL)

Spherical TiO_2 particles ($\sim 0.35 \mu\text{m}$), prepared by controlled hydrolysis of titanium tetraethoxide in ethanol, are porous agglomerates of an approximately 6nm primary structure. We consider here the internal atomic structure of these agglomerates, hitherto considered to be amorphous. X-ray scans on a dried and pressed powder disk indicate a broad oscillatory diffuse scattering pattern with an approximate correlation range of 1nm. The final $S(Q)$ was obtained using both transmission and reflection geometries on a thin pressed disk in an evacuated Be dome. Compton scattering was experimentally removed using the Ice-Sparks analyzer PSD detector (resolution $\sim 30\text{eV}$). Preliminary modelling of this diffuse pattern indicates a similarity to the Debye scattering function for the brookite crystalline phase with approximately a 1nm particle size. The Fourier transform, $G(r)$, also shows distances and coordinations attributable to the (local) octahedrally coordinated crystalline order in brookite (see below). Initial high resolution electron microscope (HREM) studies, on the other hand, reveal only occasionally ordered regions in an otherwise rather amorphous structure and it therefore appears that this material may serve as an interesting prototype for a discussion of nanocrystalline vs. amorphous organization in which the brookite phase - the most distorted of the crystalline polytypes - serves as a guide to the actual structure.

Research supported by the DOE/BES on DE-FG05-87ER45325.

STRUCTURAL STUDY OF SI-ALKALI METAL INTERFACE BY XSW

S. Lagomarsino, F. Scarinci, P. Castrucci (I.E.S.S.), C. Giannini (C.N.R.S.M.), J.R. Patel and E. Fontes (AT&T)

A structural study of Si-alkali metal interface has been carried out by means of X-ray Standing Wave (XSW) technique. In XSW experiment the crystal reflectivity and fluorescence emission from an adsorbate are simultaneously measured for a certain Bragg reflection as a function of incidence angle. The position F_{hkl} and the coherent fraction F_{hkl} obtained by fitting the experimental fluorescence yield with the theoretical one are, in one position model, the distance of the adsorbed atoms to the diffracting plane (hkl) in units of lattice spacing and the relative number of atoms at P respectively. The aim of this work was to determine the alkali metal position on $Si(111)7 \times 7$ reconstructed surface. The AT&T Bell Labs ultra high vacuum chamber at beam line X15A allows in situ film deposition, surface characterization by LEED and Auger spectroscopy and XSW measurements. Cs atoms have been evaporated from a carefully outgassed SAES dispenser on a $Si(111)7 \times 7$ reconstructed surface held at room temperature. Samples with different coverages (0.05, 0.1, 0.5 ML) have been investigated. Their LEED pattern still showed 7×7 symmetry. Fig.1 shows fluorescence yield and reflectivity spectra for (111) and (-111) diffracting planes relative to 0.1 ML coverage. For each diffracting plane

Fig.1

Fig.2

Top site H_{3F} H_{3U} T_{4F} T_{4U} H_{4F} H_{4U}

0.2-0.4 0 0.4-0.5 0 0.2-0.3

F=faulted U=unfaulted

the XSW measurements gave similar results as a function of coverage. F_{111} ranges between 0.36 and 0.53, F_{-111} between 0.70 and 0.76 and F_{111} between 0.20 and 0.31. The quite low coherent fraction, as is present in all the samples, is due to adsorption on different specific sites and not to a random positioning. It is also well known that the alkali metal undergoes a saturation at 1 ML or less, excluding therefore the possibility of building up islands. As it is impossible to fit the experimental values considering a configuration where the Cs atoms are bound to the Si 19 dangling bonds, the threefold hollow site ($H3$), the threefold filled site ($T4$) and the top site are considered as possible absorption sites. Fig.2 shows their

positions with respect to $Si(111)7 \times 7$ surface following the Qian and Chadi atomic position calculation for the DAS model reconstruction. Tab.1 shows the distribution of alkali atoms on the Si reconstructed cell, where distinction has been made between the faulted and the unfaulted half of the cell. The bond length is found to range between 3.27 and 3.67 Å. Our experimental results clearly indicate that alkali metal adsorption does not remove the stacking fault in the $Si(111)7 \times 7$ reconstructed cell.

X-RAY STANDING WAVES: $YBa_2Cu_3O_7$ ON $SrTiO_3$ (100) AND Si (111)

J. Zegenhagen and H.-U. Habermeyer (Max-Planck-Inst., FRG), T. Siegrist, J.R. Patel and E. Fontes (AT&T), and L.E. Berman (NSLS)

We investigated ultrathin films of $YBa_2Cu_3O_7$ (YBCO) on $SrTiO_3$ (100) ($d_{200} = 1.95$ Å) and $Si(111)$ ($d_{111} = 3.14$ Å) with the X-ray standing wave (XSW) technique. Structural information is obtained by analyzing the shape of the fluorescence yield curves within the range of Bragg reflection. Because of the energy dispersive detection of the fluorescence, the lattice location of the individual constituents of a film can, in principle, be analyzed structurally independently.

Figure 1 shows schematically c -axis oriented (c -axis = 11.68 Å) YBCO in $SrTiO_3$ and Si . The standing wave pattern (thin lines) from a (200) reflection is almost in phase with the YBCO a - b planes in case of YBCO/ $SrTiO_3$ since the lattice constants of film and substrate match well, i.e., $11.68/1.95 \approx 6$. In case of YBCO/ $Si(111)$ the wavefield ((111) reflection) goes rapidly out of phase with the a - b planes of the film since $11.68/3.14 = 3.72$.

Thin films of YBCO were prepared by MBE and laser deposited on flux grown $SrTiO_3$ (100) and Czochralski grown $Si(111)$, respectively, and subsequently exposed to ambient conditions. XSW measurement were performed in air using $Si(111)$ and $SrTiO_3$ (200) reflections. Fluorescence from the sample was detected with $Si(Li)$ solid state detector.

The results of two XSW measurements with ≈ 100 Å YBCO on $Si(111)$ and 110 Å YBCO on $SrTiO_3$ (100) are shown in Fig. 2 and 3, respectively.

For YBCO on $Si(111)$ analysis of the angular dependent fluorescence yield of the metal components of the films revealed their uncorrelated distribution with respect to the wavefield planes — a result which is expected because of the unfavorable lattice mismatch. Films thinner than 30 Å will have to be examined, most likely *in-situ*, to provide structural information.

For YBCO on $SrTiO_3$ analysis of the Sr and Ti fluorescence proved the excellent crystalline quality of the substrate. The Ba L-fluorescence from the film could not be separated from the Ti fluorescence. Analysis of the fluorescence of Cu and Y demonstrated their correlation with the wavefield diffraction planes. This does not only indicate good epitaxial quality but also a preferred interfacial bonding geometry.

(1)

(2)

(3)

We have observed sustained intensity oscillations during deposition of Pd on Si(111) at room temperature, as shown in Fig. 1. We presume this is due to a periodic roughening of the overlayer, which produces laterally broad and diffuse diffraction features for partial and fully completed layers, respectively, quite analogous to the mechanism for "RHEED oscillations". An intriguing feature is that the LEED pattern is completely diffuse, so the periodic growth is presumed to take place at the buried interface, where freshly arriving metal reacts with substrate silicon. We obtained data for several different temperatures and also for the CoSi(111) system. Oscillations in the latter are more strongly damped with coverage.

Lateral scans through the diffraction "rods" are shown in Fig. 2 for several different coverages. We find that strained (0001)Pd₂Si/(111)Si forms for coverages below about 7 Å (Fig. 2a). At higher coverages, the Pd₂Si is relaxed with a surface mesh spacing of $k=1.025$ (Fig. 2b). The original phase is consumed by the further growth. A fascinating aspect of the problem is the fact that only a small fraction of the low coverage silicide (below 7Å) is epitaxial. We suspect that the majority of this precursor phase is amorphous, and that we are seeing a thickness induced amorphous to crystalline transition.

This result provides entirely new and unexpected information concerning low temperature silicide formation, with implications for heteroepitaxial growth of metastable layers, in general.

Work supported in part by DOE Fac. & Stud. award, and NSF #DMR8921124.

K. Evans-Lutterodt, E. Fontes, B.J. Hinch, L.H. Dubois and J. Patel, AT&T Bell Laboratories, Murray Hill, NJ 07974.

We have studied the melting of an incommensurate monolayer of Pb on Si(111) with synchrotron x-ray diffraction, with substrate temperatures from T=172K to T=548K. Between T=172K and 273K the Pb lattice constant and half width at half maximum (HWHM) are independent of temperature; the Pb lattice constant is 2.7% larger than the substrate.

Between T=273K and 423K the Pb overlayer lattice constant is a continuous function of temperature, varying between 2.7% and 3.6% incommensurability. The longitudinal and transverse HWHM are temperature independent. Curiously we find that the second harmonic of the incommensurate structure is 3 times as broad as the first harmonic. Clearly, while this phase has long, solid-like correlation lengths (550Å), it does not have conventional long range order. Above T=423K the halfwidths increase with temperature, implying a melting of the 2 dimensional overlayer. Interestingly, we find that the even though the longitudinal correlation length varies between 550Å and 57Å, the ratio of longitudinal to transverse correlation lengths is relatively constant. In other words, the data in the melted phase shows no evidence of the orientational fluctuations expected for a hexatic phase.

R.Felici and C. Ottaviani (ISM-CNR,Italy), and I.K.Robinson and P.Eng (AT&T, Bell Labs.)

The study of the adsorption of metals on the surface of semiconductors is fundamental both from theoretical and from technological points of view because the transport properties of the near surface region are strongly influenced by the geometry and binding site of the deposited metal atoms.

We have performed a surface diffraction experiment to determine the bonding geometry of Antimony onto the Si(001) 2 x 1 reconstructed surface. The measurements have been carried out using the UHV diffractometer installed at the X16A beamline. The samples were prepared *in situ* under high vacuum conditions with a base pressure of 8×10^{-11} torr. Silicon wafers were degassed and cleaned by a resistive heating at 1000 K for about 2 hours followed by a 30 s flash at 1370 K and a slow cooling to room temperature. After this procedure, that was previously tested in a different UHV chamber with LEED and AUGER techniques, a clear Si (001) surface reconstructed 2x1 was obtained.

X-ray diffraction measurements were performed on the clean Si surface. We could observe and measure diffraction peaks corresponding to the 2x1 reconstruction. Two domains were present and the corresponding peaks had a similar intensity. In fig. 1 we present the measured intensity for the (0,1/2,1/2) peak. This means that the miscut of the surface did not induce any preferential orientation in the surface reconstruction. We also looked for the presence of peaks corresponding to different reconstructions (e.g. 2x2). No evidence for such peaks was found.

Fig.1 - Reconstruction peaks (0,1/2,1/2) for clean Si (001) 2x1 surface (a) and for Sb on Si (001) 2x1 (b). The intensities are normalized to the monitor counts. The lines are Lorentzian fit to the experimental data.

Antimony deposition was performed keeping the Si substrate at 800 K and followed by a further annealing at the same temperature for 10 min. This procedure is known to desorb all the Antimony but a monolayer. The amount of Antimony deposited on the surface was also checked by Auger analysis. The surface 2x1 reconstruction peaks are still present after this procedure but with different relative intensities. Data analysis to determine the geometry of the bonds between Antimony and Silicon are still in progress.

The structure of K monolayers on Cu(100)

H.L. Meyerheim, V. Jahns, W. Moritz, and H. Schulz (U. Munich)
I.K. Robinson and P.J. Eng (AT&T Bell Lab., Murray Hill)

The study of alkali adsorption on metal surfaces has found considerable interest from both, theoretical and experimental point of view (1,2). In the LEED study of the adsorbate system K/Cu(100) two commensurate phases labeled C1 and C2 are suggested (3), both of them forming quasi hexagonal structures.

In our first experiment K was evaporated at T=350 K on clean Cu(100). A quasi hexagonal LEED pattern was observed after this procedure. Subsequently grazing incidence x-ray diffraction (NRD) measurements were performed in the radial direction along the a^* axis of the rec. lattice as shown in the upper panel of Fig. 1. In the lower panel the corresponding XRD spectra (h 0 0.3) are shown. The reflex positions of the C1 and the C2 phase are indicated by arrows. Immediately after evaporation only background could be observed. After about 15 min. a superlattice reflection appears at $h = -0.644$ corresponding to an incommensurate phase between C1 and C2.

A continuous shift along a^* was observed with time as is illustrated in Fig. 1. Simultaneously, the reflection intensity rapidly decreases. This might be induced by desorption, clustering of K or even by contamination. In the transverse direction the structure was found to be commensurate within $10^{-4} b^*$ as shown in the center panel. The peak shift along a^* indicates a continuous transition to the less dense C1 structure without the presence of lock-in phases. The analysis in the vicinity of the $(-0.5 \ 0.33)$ reflection led to the same result.

Fig. 1

An attempt was made to fit the measured reflection profiles using a Lorentzian powerlaw function convoluted with a Gaussian resolution function. Depending on the peak position no significant changes of the linewidth and lineshape could be observed, however better quality data are needed in order to make a definitive statement.

In further experiments K was evaporated using a very small evaporation rate while simultaneously performing radial scans as described above. The position of the first peak to appear was found to be $h = -0.614$ shifting upwards with increasing time. In no case the formation of the commensurate phases was observed. The formation of satellite reflections.

The analysis of truncation rod data is still in progress. We (H.L.M. and V.J.) would like to thank I.K. Robinson and AT&T for their hospitality. The support of this work by the BMFT under grant # 05461AB8 is gratefully acknowledged. (1) see e.g. E. Bauer in: Structure and Dynamics of Surfaces (Springer, 1987). (2) T. Aruga et al., PRB 34, 8237 (1986). (3) T. Aruga et al., Surf. Sci. 158, 490 (1985)

I. K. Robinson (Bell Labs), R. Feidenhans'l (Riso), F. Besenbacher and F. Jensen (University of Århus)

We have made 2D crystallographic measurements of various ordered phases of O and S on Cu(111). S induces a well-known $\sqrt{7}\times\sqrt{7}$ phase with a unit cell denoted I in the Figure. The same structure was obtained either by segregation of S impurities from the bulk or by H_2S dosing, although the latter was more reliable.

The O structures (II and III) are more complicated. These were first identified by Scanning Tunneling Microscopy (STM) in Århus, and confirmed with LEED although the pattern is very weak. We used the same dosing and annealing conditions as were used in the STM study and obtained sharp diffraction peaks in the predicted positions.

Both structures II and III have monoclinic symmetry and therefore have six twin domains. We found that the fraction of the surface in each twin varied substantially, but always reproduced. We concluded that surface steps play an important role in selecting the preferred orientation, as they do on Cu(100).

Structure II indexes on a reciprocal lattice 19×19 that of the bulk, while structure III is 44×44 . We have extensive datasets on each structure and crystallographic analysis is under way.

X-RAY DIFFRACTION UNDER GRAZING INCIDENCE ON Si⁺ IMPLANTED SILICON AFTER THERMAL ANNEALING

X16A

S. Rugel, H. Metzger and J. Robinson⁺
U. München, Germany, ⁺ AT&T Bell Labs, Murray Hill

We investigated silicon single crystals with (100) surface orientation after Si⁺ implantation (100 keV, $1e15/cm^2$) and thermal annealing for 10sec at 670°C and 1000°C. Monochromatic X-rays (1.4 Å) were diffracted under grazing incidence by the (022) lattice planes which are perpendicular to the surface. The diffracted intensity as a function of the exit angle α_1 was detected by a position sensitive detector at different incident angles α_0 . In addition, diffuse scattering was measured in the vicinity of the Bragg peak, where the in-plane scattering vector Q was changed by the deviation q from the reciprocal lattice point parallel to Q .

The Bragg diffracted intensity versus α_1 (fig. 1) shows broadening and shift of the peak for α_0 near the critical angle of total reflection α_c , which is typical for dynamic scattering. This indicates amorphous induced perfect recrystallization of the amorphous subsurface layer created by ion implantation. Diffuse scattering intensity is shown in figure 2 on a double logarithmic scale versus q . The intensity is integrated over q_z determined by a narrow region in α_1 . In the regime of q near the reciprocal lattice point the in scattering contribution arises from the long wave region of the defect distortion field. After annealing at 1000°C diffuse intensity shows a q^{-4} behaviour which is due to extended defects, while the sample annealed at 670°C varies as q^{-2} , indicating point defects.

At larger q (here $q = 0.064 \text{ Å}^{-1}$) the distortion field in the vicinity of the defect center contributes to the diffuse intensity. Figure 3 shows α_1 -spectra from both samples taken at $q = 0.064 \text{ Å}^{-1}$. After annealing at 1000°C a distinct maximum of the scattering intensity in α_1 is observed. Computer simulations of scattering due to (111) dislocation loops situated at the original amorph/crystalline interface show good agreement with the experimental data when a loop radius of $175\pm 30 \text{ Å}$ is assumed. This is consistent with TEM investigations taken at the same sample. After annealing at 670°C (o in fig. 3) the scattering intensity at $q = 0.064 \text{ Å}^{-1}$ hardly exceeds the background as expected for small defects.

Supported by BMFT under contract number 03PE112MU

Fig.1: α_0 -resolved (022) Bragg diffraction after annealing at 670°C for different α_0 . Solid lines: dynamical calculations.

Fig.2: Diffuse scattering intensity close to (022) as a function of q on a double log scale

Fig.3: α_1 -resolved diffuse scattering intensity at $q=0.064 \text{ Å}^{-1}$ compared with model calculations for dislocation loops with different loop radii r : $r=125 \text{ Å}$, $---$; $r=175 \text{ Å}$, $---$; $r=200 \text{ Å}$

D.-M. Smilgies (MPIF Göttingen), I.K. Robinson (AT&T Bell Labs), N. Onda and H. von Känel (ETH Zürich)

We have investigated iron silicide films grown by MBE on Si(111) substrates. The films have been grown near room temperature at a stoichiometry of 1:2 (FeSi_2). They crystallize in a metastable cubic phase with metallic properties. We have investigated three samples of thicknesses 72Å, 73Å, and 105Å. The aim of our study was to characterize the structural quality of the films and of the interface between film and substrate. The silicide films are terminated by the native oxide and are stable under normal conditions, so that the measurements could be performed in air.

The figure shows a scan along the (10L) truncation rod. Centered about the (102) position, the thickness fringes of the film can be seen. At (101) and (104) the bulk Bragg reflections of the substrate appear. Note that there are no fringes at the (105) position. In the vicinity of the (101) spot the bulk truncation rod and the fringes are seen to interfere; this will provide structural information about the interface. The thickness oscillations are more strongly attenuated than on other silicide films. This effect might be due to the growth mode in which diffusion is limited so that not all defects produced during the deposition can anneal. The observed pattern of absent reflections, such as the (105), reveals that the film has the CsCl structure and B stacking. This result is surprising because the stoichiometry of the films is known to be FeSi_2 . Further investigations of samples using modified growth conditions are in preparation and should clarify this apparent contradiction.

HYDROCARBON FILMS ON SILICON I: THE STRUCTURE OF WETTING FILMS.

Ian M. Tidwell, Thomas A. Rabedeau, Seth D. Kosowsky, and Peter S. Pershan (Harvard U.)

The wetting of polished silicon wafers by cyclohexane and perfluoro(methylcyclohexane) was studied using x-ray specular and off-specular scattering. The substrates were held in a sealed cell, temperature controlled near room temperature to better than 2mK, and wet through the vapor, which was maintained in saturated equilibrium with the liquid. Differentially heating the substrate relative to the liquid reservoir over the temperature range ΔT from 0K to 1.3K resulted in wetting film thicknesses of between 200Å and 15Å. The film thicknesses, densities and interface widths were determined from x-ray specular reflectivity scans; a typical set are shown in the accompanying figure for cyclohexane films 160Å (○) and 55Å (▽) thick. The data for film thickness l versus temperature differential ΔT are consistent with the $l_0(\Delta T)^{-1/3}$ power law predicted for Van der Waals interactions.

Off-specular scans along a direction with $q_z = q_i/500$, shown in the figure for the 160Å (○) and 55Å (▽) thick cyclohexane films respectively, measure diffuse scattering associated with the roughness of both the substrate/liquid and liquid/vapor interfaces. Oscillations, resulting from correlations between the roughnesses of these two interfaces, decrease with increasing film thickness. For films thicker than approximately 100Å scattering from the two interfaces are uncorrelated.

This work supported by the National Science Foundation through grant No. NSF DMR-89-20490.

Fig. 1. Specular and off-specular data for a 160Å thick cyclohexane film (○ and ● respectively) and a 55Å thick film (▽ and ▼) on silicon.

Ian M. Tidwell, Thomas A. Rabedeau, Seth D. Kosowsky, Peter S. Pershan, John P. Folkers and George M. Whitesides (Harvard U.)

The structure of several C₁₈ alkylsiloxane monolayers were studied using x-ray grazing in-plane diffraction (GID) and reflection (both specular (SR) and off-specular (OSR)). Alkylsiloxane monolayers were formed by self-assembly from solution onto Si(001) wafers coated with the native silicon-oxide. The GID measurements of complete alkylsiloxane monolayer films on polished silicon wafers detected only one peak associated with the hydrocarbon film, at a position corresponding to close packing of the chains with a molecular area of about 20.2 ± 0.2 Å². No higher order peaks were detected. The peak has a Lorentzian line shape, the width of the peak indicating correlation lengths of about 35 ± 5 Å. OSR scans taken along a direction q_z = q_z/500 show structure that is qualitatively similar to the SR, indicating that the roughness of the two interfaces of the hydrocarbon film are correlated.

SR measurements of incomplete alkylsiloxane monolayers confirmed that the hydrocarbon layer has a density similar to the fully formed monolayers, but the molecules tilt, reducing the thickness of the films. GID measurements of these films show that although there were larger intersample variations, the molecular areas were similar to those for fully formed films (20.5 ± 0.5 Å²) and the integrated intensities of the GID peaks and the correlation lengths were both significantly reduced.

Thermal effects were probed by cooling and heating a complete alkylsiloxane monolayer. With increasing temperature, the peak center moved to lower q (the peak shift corresponding to a coefficient of linear expansion of 5 × 10⁻⁴ K⁻¹) and the integrated area of the peak decreased (by a factor of approximately 2 from 0°C to 100°C). The correlation lengths decreased on both heating and cooling from room temperature. A possible explanation for this is that since the monolayers were formed by 'self assembly' at room temperature, temperature changes lead to differential expansion of the hydrocarbon chains and the silicon substrate. While changes on cooling the sample were reversible, the intensity and width of the peak, corresponding to the local order of the alkane chains, were not fully reversible on heating, suggesting reduced alkane order.

Finally, since the samples suffered x-ray damage, the data was taken by translating the sample perpendicular to the beam, to expose a fresh region, approximately every 30 minutes.

This work supported by the National Science Foundation through grant No. NSF DMR-89-20490

Fig. 1. GID peak for a complete alkylsiloxane at 26°C (○) and 100°C (▽).

F.A. Dilmanian¹, E. Nachaliel¹, R.F. Garrett^{1a}, W.C. Thomlinson², L.D. Chapman², N.F. Gmit², N. Lazar², H.R. Moulin², T. Overlitz², H. Rarback², D.N. Slattin¹, P. Spanne³, P.M. Stefan², A.C. Thompson⁴, N.D. Volkow¹, and H.D. Zeman⁵

¹Medical Department¹, NLS², and DAS³, BNL, Center for X-Ray Optics, LBL⁴, and Dept. of Biomed. Eng., U. of Tenn, Memphis⁵

Preliminary phantom studies were carried out with a Multiple Energy Computed Tomography (MECT) system being developed at X17B. The system consisted of the unstable section of the MECT/Materials Sciences monochromator in the Bragg-Bragg mode, and a 120-element Si linear array detector of 0.25 mm element width and 3 mm crystal depth obtained on loan from the Coronary Angiography Project. The data acquisition system was also that of the Angiography Project.

The phantom was a 27-mm Plexiglas cylinder with 5-mm diameter axial holes. The holes were filled with KOH solutions of 200 mmolar (holes 1), 39.2 mM (holes 2), 7.7 mM (holes 3), 1.5 mM (holes 4), and 0.0 mM (holes 5). Images were taken at 20 and 38 keV, using crystal detuning to provide a yield of 50% to 80% of the L_α peak. Data were taken in the continuous mode at time intervals of 1/60 s and at angular steps of 0.25°, yielding a complete 360° measurement in 24 seconds. Corrections were applied to the data before and after tomographic reconstruction, mainly to remove artifacts caused by experimental imperfections. Because of the remaining artifacts in the center, the quantitative analysis included only the outer holes. The SNARK tomographic reconstruction routine was used. Figs. 2 and 3 show the 20 keV and the 38 keV images.

The dual-photon absorptiometry (DPA) formalism was applied to obtain separate images of the low-Z element group (i.e., water) and the intermediate-Z element group (i.e., K). Because of the beam harmonic contamination, the effective beam energy values were obtained first by using the knowledge of the KOH concentrations in the holes. The values obtained were 22 and 39 keV. Figs. 4 and 5 show the low-Z and intermediate-Z DPA images obtained, respectively. While in Fig. 5 the contrast between different K concentrations is enhanced compared to Figs. 2 and 3, the contrast in Fig. 4 shows no dependence on K concentration, as it is expected. Pixel averaging in Fig. 5 shows that the 39.2 mM KOH solution, which is equivalent to 50% of brain normal K concentration, is distinct from water.

Figure 1

Fig. 1: Pattern of the holes in the phantom. The KOH concentration is 200, 39.2, 7.7, 1.5, 0 for holes 1, 2, 3, 4, 5 respectively.

Figure 2

Figure 3

Figure 4

Fig. 2: Reconstructed image at 20 keV.

Fig. 3: Reconstructed image at 38 keV.

Fig. 4: Image of low-Z elements.

Fig. 5: Image of intermediate-Z elements.

^aNow at Australian Nuclear Science and Technology Organization. The authors thank J.B. Hastings and D.P. Siddons for their help in design of the monochromator, and C. Britz, R. Greene, A. Lenhard, and M. Shleifer for their technical support. Research supported under DOE contract DE-AC02-76CH0006.

BIOLOGICAL EFFICACY OF THE AUGER EFFECT

B.H. Laster, W.C. Thomlinson, C. Gordon, N. Gmur, N. Lazarz, V. Benary, L. Warrentien and R. G. Fairchild

Iodoxyuridine (IdUrd), an analog nucleoside to the natural base thymidine (Thd), is incorporated in cellular DNA during cell replication. In addition to its radiosensitizing properties, the presence of iodine in DNA could provide even more ionization within the DNA itself if activated by photons above the K absorption edge of iodine (33.2 keV). Enhancement of dose would accrue from the induction of the photoelectric effect in iodine and the concomitant release of Auger electrons. The X17B1 beam line of the National Synchrotron Light Source (NSLS) at Brookhaven National Laboratory (BNL) delivered monochromatic photons above and below the K edge of iodine to V-79 Chinese hamster cells with and without IdUrd in DNA to evaluate this phenomenon. Results indicated a 40% enhancement of dose between iodinated cells irradiated above and below the K absorption edge, thus showing significant biological efficacy of the Auger effect. The sensitization enhancement ratio (SER) for iodinated and noniodinated cells was 3.0. These findings indicate that Photon Activation Therapy (PAT), using DNA-incorporated IdUrd with Fairchild sources (samarium-145 emitting 40 keV photons) via brachytherapy techniques is a promising new modality for the treatment of malignant brain tumors. Additionally, the use of X17B1 for the external irradiation of superficial tumors with DNA-incorporated IdUrd should be considered for tumors that cannot be adequately resected.

IN-SITU X-RAY DIFFRACTION STUDY ON MODIFIED SPINEL (β) AND SPINEL PHASES (γ) OF Mg_2SiO_4 AT HIGH PRESSURE AND

HIGH TEMPERATURE

Y. Meng, D. J. Weidner, G. D. Gwanmesia, R. C. Lieberman, M. T. Vaughan, R. E.

Pacalo, A. Yeganeh-Haeri, K. Leinenweber, Y. Wang, Y. Zhao

CHPR, and Dept. of Earth and Space Sciences, State University of New York, Stony Brook, NY 11794

The P-V-T equation of state of the β and γ -phases of Mg_2SiO_4 were studied up to 67 kbar and 1000 °C in a DIA-type cubic-anvil press (SAM-85) installed at X17B. In order to have a direct comparison, both specimens were loaded in the same cell, in which NaCl was used as a pressure medium and also as an internal standard. Unit cell parameters were measured to 67 kbar at room temperature and up to about 670 °C while keeping the ram load constant. The pressure in the sample chamber changes by less than 2 kbar when the ram load is constant. Phase transformation to the olivine structure (α -phase) occurs in the temperature range from 670 to 790 °C in about 15 minutes. The results on β and γ -phase were complicated by non-hydrostatic stress; however, the phase transition seems to have the effect of reducing this stress. Reliable data were obtained on the transformed α -phase of Mg_2SiO_4 at 67 kbar and high temperature. The average volume expansion is $30 \times 10^{-6}/K$, compared to $38 \times 10^{-6}/K$ at 1 bar (Hazen, 1976). The average value of the Anderson-Gruneisen parameter δ_T of the α -phase is calculated to be 5.1 over the range from room temperature to 1000 °C.

Performance of the MECT/Materials Science Monochromator in the Laue-Laue Diffraction Mode

E. Nachev¹, F.A. Dilmanian¹, R.F. Garrett¹, T. Overstulzen²,
L.E. Berman², C. Brite², L.D. Chapman², J.B. Hastings², N. Lazar²,
A. Lenhard², H. Raback², M. Shleifer², D.P. Siddons²,
P.M. Stefan², and W.C. Thomlinson²

Medical Department¹ and the NSLS², BNL

A high-energy wide-beam tunable monochromator was earlier developed for use in the Multiple Energy Computed Tomography (MECT) system and in the Material Science research. The design was a Golovchenko-type. The monochromator was originally prepared for the Bragg-Bragg mode. In this mode the monochromator performed satisfactorily in the 31-92 keV range and was used for preliminary MECT phantom studies, as well as for XRF experiments. The general design of the MECT/Materials Science monochromator was subsequently changed to consist of two Laue-Laue monochromators, one tunable and one at the fixed energy of 100 keV (see Fig. 1), in which the Golovchenko monochromator serves as the tunable section. In this design, the first crystal of the first section acts as a semi-transparent reflector that splits the beam into the two branches. For the MECT's dual-photon absorption (DPA) method, these two branches will be tuned to 40 keV and 100 keV. The two beams will be simultaneously detuned and ready. Switching from one to another will be merely a matter of shutting of one and opening the other, which should not take more than a few seconds.

The Tunable Section, equipped with two Si<220> Laue crystals, 70 mm long, 7 mm high, and 1 mm thick, was tested in the X17B hutch in October 1991. The first crystal was water-cooled using a molybdenum cooling block thermally coupled to the widened base of the Si crystal with *in*-Ga liquid metal. The electrorestric devices used for the Laue tuning and for the tilt had 30 micron and 105 micron stroke, respectively.

The measurements showed that the beam non-uniformity along its 65-mm horizontal width at detuning level with a yield of 10% of the Laue peak is not worse than 50%, which is acceptable for preliminary MECT imaging. The Laue curves measured (Fig. 2) agree with the expected Darwin widths.

Fig. 1: Conceptual design of the monochromator. L1 and L2 are crystals of the tunable, low-energy section, and H1 and H2 are those of the fixed, high-energy section (probably 100 keV).

Fig. 2: Rocking curves measured with the tunable section of the monochromator.

¹Golovchenko et al., Rev. Sci. Instr. 52, 509-516, 1981.
²Now at Australian Nuclear Science and Technology Organization.
The authors thank R. Greene, M. Nelson and M. Rivers for their assistance.
Research supported under DOE contract DE-AC02-76CH0006.

PRELIMINARY STUDIES OF THE POTENTIAL FOR STRUCTURE REFINEMENT AT HIGH PRESSURES USING THE DIA APPARATUS: SiO₂, STISHOVITE, AT HIGH PRESSURE.

John B. Parise, K. Leinenweber, D. J. Weidner, J. Rakovan, M. Vaughan,
G. Gwanmesia and R. C. Liebermann (SUNY, Stony Brook).

Very fine grained stishovite (SiO₂) powder (Gwanmesia *et al.*, 1990) has been studied as a function of pressure in the DIA apparatus. An example of raw data obtained is shown in Figure 1 along with that normalized to the absorption-corrected incident X-ray intensity and that calculated using the parameters of Ross *et al.*, 1990. The qualitative agreement suggests that, with appropriate precautions to minimize the effects of preferred orientation and deviatoric stress, energy dispersive data may be useful for the refinement of crystal structure in this apparatus.

Figure 1. Raw and normalized data for stishovite collected at 4.5 GPa at room temperature. The normalization involves correction for the absorption of optical elements such as Be windows, sample cell assembly and air, along with the Compton shifted incident beam spectrum. During data collection the apparatus was translated to minimize the effects of preferred orientation. The top window contains a pattern calculated using the cell parameters and atomic parameters at 1.5 GPa reported by Ross *et al.* (1990).

Chapman, D. (1990) *Nuc. Inst. Methods in Phys. Res.* A291, 205-208
Gwanmesia, G. D. *et al.*, *Geophys. Res. Letters* 17, 1331-1334
Ross, N. L. *et al.* (1990) *Am. Mineral.* 75, 739-747

Computed Microtomography of Meteorites using a Linear Array Detector

S.R. Sutton^{1,2}, P. Spanne², M.L. Rivers^{1,2}, and K.W. Jones²

¹Dept. of Geophysical Sciences, The Univ. of Chicago, Chicago, IL 60637;

²Dept. of Applied Science, Brookhaven National Laboratory, Upton, NY 11973

Computed microtomography (CMT) is a useful technique for imaging the internal structures of objects with high spatial resolution. The principal value in planetary sciences is the ability to "see" the internal structures of extraterrestrial objects, such as rare Antarctic meteorites, prior to developing sampling strategies and allocation plans. In particular, the locations of clasts are determined, allowing planes with maximum number of clast intersections (potential cut planes) to be visualized.

We have been developing CMT techniques using bending magnet and wiggler synchrotron radiation at the NSLS. Our previous work has used primarily a pencil beam-transmission approach with a scintillation detector and a 2-dimensional CMT image was made on a fragment of the Allende carbonaceous chondrite with this apparatus on the X17 wiggler source. We report here our first attempts to use a linear array detector to improve the acquisition speed. In this approach, a slit source (horizontally larger than the object but vertically narrow) irradiated an entire plane in the object simultaneously. A linear position-sensitive array detector was then used to simultaneously measure the integrated absorption along many parallel beam paths through the sample. The particular array used in this work was a 1024-element photodiode array with a total width of 25 mm (each element = 25 μ m) and covered by a thin, Gd_2O_3 phosphor. Optical blooming by the phosphor limited the effective spatial resolution of the device to about 100 μ m.

Forty (40) horizontal slices within the 1 cm Allende fragment were imaged in only 30 minutes, a decrease of about a factor of 200 in analytical time compared to the pencil beam approach. Each slice was separated vertically by 100 μ m. The resulting reconstructed tomogram clearly shows chondrules, inclusions and fractures with a spatial resolution of about 100 μ m.

Potential detrimental consequences of the CMT irradiation must be studied before the method can be considered for routine use on rare extraterrestrial objects as part of the preliminary examination activities of the meteorite curator. That is, it is important to ensure that applications of radiation-sensitive analytical techniques (e.g., thermoluminescence) are not compromised by the CMT imaging. Radiation doses delivered by the wiggler synchrotron beam under the CMT conditions (1.5 mm Ta filter on incident beam) are estimated to be 10-100 rads/sec or a total dose per CMT slice (10 sec exposure) of about 100-1000 rads. This dose is relatively low when compared with the typical cosmic radiation exposure ($> 10^5$ rads) and the equivalent dose of meteorite thermoluminescence (10^4 - 10^5 rads). The use of monochromatic radiation rather than filtered white radiation is expected to reduce the required radiation dose by about an order of magnitude.

(Research supported by NASA NAG9-106, DoE DE-AC02-76CH00016, NSF EAR89-15699.)

More on the High-Pressure, High-Temperature Equation of State of $MgSiO_3$ Perovskite

D. J. Weidner, Y. Wang, R. C. Liebermann, M. T. Vaughan, K. Leinenweber, Y. Meng, R. E. G. Pacalo, A. Remsburg, A. Yeganeh-Haeri, Y. Zhao
(CHiPR and Department of Earth and Space Sciences, State University of New York at Stony Brook, Stony Brook, NY 11794)

Following our previous high-temperature, high-pressure study on $MgSiO_3$ perovskite (Wang *et al.*, *Science*, 251, 410, 1991), we have carried out additional *in situ* synchrotron x-ray diffraction studies in the DIA-6 cubic-anvil apparatus (SAM-85) at X17 beamline of Brookhaven National Light Source. The polycrystalline perovskite specimen was synthesized in the 2000-ton Uniaxial Split-Sphere apparatus (USSA-2000) at 26 GPa and 2073 K. In order to minimize deviatoric stress, the sample was embedded in NaCl, which also served as an internal pressure standard. Energy-dispersive x-ray spectra were collected at 7.2 GPa on both heating and cooling cycles. With increasing temperature, a sudden increase in the unit-cell volume was once again observed at about 620 K, but with a smaller magnitude than we previously reported. The pressure remained constant within 0.2 GPa during heating to 900 K. Upon cooling, unit-cell parameters were reversible to 623 K, below which pressure decreases about 0.5 - 1.0 GPa were observed from the NaCl standard. At 7.2 GPa, the high-temperature ($T > 623$ K) thermal expansion is reversible; the room-temperature unit-cell volume also appears reversible within the experimental error after pressure correction. Above 620 K, the data on unit-cell volume can be fitted rather well by a Suzuki-type formula with effects of pressure taken into account. The volumetric thermal expansion (α_v) thus obtained fits our previous high-temperature data almost equally well, yielding $\alpha_v \approx 1.6$ (10^{-5} K⁻¹) at 650 K and 1.8 near 1000 K. Our new data thus confirmed the low volumetric thermal expansion we previously reported.

The Pressure Dependence of Transition Temperature for Structural Phase Transition in NaMgF_3 Perovskite

Yusheng Zhao, Donald J. Weidner, Robert C. Liebermann, Xing Liu, Jaidong Ko, Yue Meng, Rosemary E. G. Pacalo, John B. Parise, Michael T. Vaughan, Yanbin Wang, Amir Yeganeh-Haeri
 CHiPR, Mineral Physics Institute, Dept. of Earth and Space Sciences, State University of New York, Stony Brook, NY 11794

An *in-situ* high pressure/temperature X-ray powder diffraction study is conducted on NaMgF_3 perovskite using synchrotron radiation (X17B superconducting wiggler port, NSLS, Brookhaven National Laboratory). High pressure/temperature is achieved by a multi-anvil press (SAM-85, High Pressure Lab. Stony Brook). The observation of superlattice diffractions related to in-phase octahedral tilting ϕ^+ and anti-phase octahedral tilting ϕ^- , such as $\{121, 103, 211\}_{Pbnm}$ and $\{120, 210\}_{Pbnm}$ & $\{122, 212\}_{Pbnm}$, indicates that the crystal structure of NaMgF_3 perovskite transforms directly from orthorhombic $Pbnm$ to cubic $Pm\bar{3}m$ without any tetragonal intermediate phase. The result suggest that the order parameters, namely the octahedral tiltings $\phi=\phi_z^+$ and $\theta=\phi_{xy}^-$, are strongly coupled together during the structural phase transition process of the NaMgF_3 perovskite. The same kind of phenomena are also observed at high pressure. The transition temperature is observed increase with increasing pressure and having a positive slope of about 6°C/Kbar (see attached diagram). This result support the Samara's theory that the structural phase transition driven by zone-boundary phonon softening has the positive slope in the P - T space.

MONOCHROMATOR HARMONIC CONTENT MEASUREMENTS AND CALCULATIONS AT ENERGIES ABOVE 26 KEV

D. Chapman, H. Moutin (NSLS); R. Garrett (BNL Med.)

We have measured and calculated the harmonic content from single and double crystal monochromators at x-ray energies above 20 keV. The measurement was done by using the Compton scattering from a thin aluminum plate in the monochromatized synchrotron beam from the X17 Superconducting Wiggler beamline. An intrinsic germanium detector was used to measure the scattered energy spectrum from this plate at about 90° scattering angle. The advantage of this technique is that the fundamental and harmonics appear at their Compton shifted energies and thus problems with pulse pile-up can be avoided. This allows an unambiguous assignment of the intensity of the fundamental and harmonics. These measurements were done with a variety of absorbers with good agreement with calculated intensities.

Measurements and calculations were done for both a single crystal monochromator (Bragg Case $\text{Si}(111)$ with 2.5° asymmetry) at 33 keV and for a double crystal monochromator (Bragg Case $\text{Si}(220)$'s) operating in the energy range of 20 to 80 keV. Based on the 33 keV single monochromator measurements, we have modelled with great success the response of a $\text{Si}(\text{Li})$ array detector in the monochromatized beam with a variety of absorbers present. Some of these results are shown below in Figures (a) and (b). The spectrum in Figure (a) is taken with no absorbers present. Figure (b) has 145 mm of water attenuating the incident beam. The solid squares represent the modelled intensities. This information will be used to correct the response of the detector for beam hardening effects in the Digital Subtraction Coronary Angiography Project images of the human heart. In the case of the double crystal monochromator, effects on the fundamental and harmonic content as a function of the second crystal detuning were also measured and calculated. This monochromator will be used for the Multiple Energy Computed Tomography project. Detuning the monochromator will be used to control the amount of harmonic contamination, a factor responsible for image degradation.

Work funded by DOE under Contract No. DE-AC02-76CH00016.

FIRST HUMAN TRANSVENOUS CORONARY ANGIOGRAPHY IMAGES AT THE NSLS

W. Thomlinson, N. Gmür, D. Chapman, N. Lazarz (NSLS); R. Garrett (BNL Med.); J. Morrison, P. Reiser, V. Padmanabhan, L. Ong, S. Green (NSU); A. Thompson (LBL); H. Zeman (U. of TN); G. Brown (SSRL); R. Hofstadter, E. Rubenstein (Stanford U.); J. Giacomini, H. Gordon (PAVH).

The Synchrotron Medical Research Facility (SMERF) at the NSLS became fully operational with the successful imaging of the coronary arteries in two patients on Oct. 5 and 9, 1990. The approach eliminates those hazards of the conventional procedure which arise from the injection of the iodine contrast agent directly into the coronary arteries. The tuneable energy and high intensity of the synchrotron radiation allow the visualization of the coronary arteries following the far safer injection of the contrast agent into the venous system. The patient is imaged by two simultaneous x-ray fan beams (0.5 mm x 140 mm) above and below the iodine k-edge allowing image digital subtraction and elimination of artifacts in the final image.

The project started at the Stanford Synchrotron Radiation Laboratory (SSRL) under the direction of Dr. Edward Rubenstein and Dr. Robert Hofstadter. The first synchrotron transvenous coronary angiograms were obtained at the SSRL in 1986, followed by further imaging runs in 1987 and 1989. During the NSLS Phase II expansion, Dr. William Thomlinson headed the construction of a superconducting wiggler beamline and the development of the world's first synchrotron medical research facility. As a result of the advantages of having a year-round operating schedule, a permanent medical facility, and eventually higher flux, the angiography project hardware was moved to SMERF from SSRL. Over the past year, intensive commissioning of the beamline and safety reviews have occurred.

The image below is a left anterior oblique (LAO 50°) view of a human heart showing the right coronary artery (arrow). The clinical importance of this image is that the artery is still widely patent four years after balloon angioplasty was performed. This image was obtained using a right atrial power injection of 36 ml of contrast agent (Angiostart 370; Bertex) at 15 ml/s. The x-ray ring energy was 2.5 GeV and current was 203 mA, with a wiggler field strength of 4.0 T. The radiation exposure was 0.4 Rad/image frame.

The immediate goal of the project is to study the optimal imaging parameters of the system, determine those regions of the coronary arterial system which can be imaged in this manner, and to upgrade the beamline optics to allow research quality imaging. The long range goal of the project is to carry out clinical research on the causes and possible cures of human coronary arterial disease.

1. W. Thomlinson et al. The Lancet, 337, 360 (1991).

Work funded by DOE Contract No. DE-AC02-76CH00016(BNL); DOE Contract No. DE-FG03-87ER60527 and NIH Account No. NIH 5R01 HL 39253-04(Rubenstein).

IMAGING AN OCCLUDED ARTERY USING TRANSVENOUS CORONARY ANGIOGRAPHY

W. Thomlinson, N. Gmür, D. Chapman, N. Lazarz (NSLS); R. Garrett (BNL Med.); J. Morrison, P. Reiser, V. Padmanabhan, L. Ong, S. Green (NSU); A. Thompson (LBL); H. Zeman (U. of TN); G. Brown (SSRL); R. Hofstadter, E. Rubenstein (Stanford U.); J. Giacomini, H. Gordon (PAVH).

A third patient was imaged at the NSLS Synchrotron Medical Research Facility (SMERF) on February 12, 1991. The patient selected was previously diagnosed as having a total occlusion in the mid-portion of the right coronary artery. This condition was well characterized and documented by conventional x-ray angiography techniques using 35 mm CINE film images and allowed a direct comparison with the transvenous digital subtraction coronary angiography x-ray images produced on X17B2.

Since the first set of October 1990 images were taken, a number of upgrades, aimed at improving image quality, have taken place.

1. The superconducting wiggler field was increased from 4.0 to 4.4 Tesla.
2. The X17B2 monochromator silicon (111) crystals, originally cut with a 0.9° asymmetry, were replaced with Si(111) 2.5° asymmetry crystals.
3. The x-ray ring global orbit feedback system¹ was made operational in order to stabilize the vertical movement of the synchrotron beam.

The image below is a left anterior oblique (LAO 30°) view of the coronary artery of the patient described above. Arrow #1 shows the right coronary artery (RCA) descending toward the right ventricle. Arrow #2 shows where contrast agent is blocked due to total occlusion of the RCA. A small (<1 mm diam.) marginal branch of the RCA, with its origin just above the total occlusion, can be seen spiraling down the right ventricle (Arrow #3). The tubular object descending into the middle of the image is the catheter behind the heart. The dark areas to the right are the aorta (upper right) and left ventricle (middle right). The image was obtained using a right atrial power injection of 36 ml of contrast agent (Angiostart 370; Bertex) at 15 ml/sec and an x-ray ring current of 178 mA. The radiation exposure was 0.7 Rad/image frame. Note the improved arterial details in this image compared to the October 1990 image (see accompanying abstract).

1. J. Galayda, Synchrotron Radiation News, 3, 22 (1990).

Work funded by DOE Contract No. DE-AC02-76CH00016(BNL); DOE Contract No. DE-FG03-87ER60527 and NIH Account No. NIH 5R01 HL 39253-04(Rubenstein).

H.D. Zeman and F.A. Di Bianca
Department of Biomedical Engineering, University of Tennessee, Memphis, TN

W.C. Thomlinson
National Synchrotron Light Source, Brookhaven National Laboratory, Upton, NY

A kinesthetic charge detector is proposed for improving the efficiency of X-ray flux utilization for intravenous coronary angiography at the National Synchrotron Light Source at Brookhaven National Laboratory. At the desired spatial resolution of 0.25 mm, the use of a kinesthetic charge detector to replace the one-dimensional Si(Li) detector used in the earlier studies, will lead to an increase in the utilized X-ray flux of at least a factor of eight, while maintaining the same patient scanning speed of 12 cm/sec.

The kinesthetic charge detector can achieve 0.25 mm resolution in the scanning direction with ~ 2 mm high X-ray beam by producing a positive ion charge image of the patient in a gas medium and moving the ion image with an electric field at the same speed and in the same direction that the patient is scanned through the X-ray beam. X-rays enter the detector perpendicular to an electric field generated between a high voltage electrode and a grid. In this drift region of the detector, the ions drift at constant velocity towards the grid which shields the detecting finger electrodes from the charge of the drifting ions. The field is set 3-5 times higher in the region between the grid and the finger electrodes in order that none of the ions stop on the grid, and in order to minimize the loss in spatial resolution produced by the transit time in this region. In this way an 8 times longer image integration time can produce an 8 times higher detected X-ray flux in the detector.

The use of krypton gas in the kinesthetic detector will allow high spatial resolution and high quantum efficiency to be achieved at reasonably attainable gas pressures. Krypton is chosen over xenon for use with monochromatic X-rays at the iodine (33.169 keV), gadolinium (50.239 keV), or ytterbium (61.332 keV) K-edge, because the shorter range of the K-emission X-rays reduces greatly the veiling glare. A detector with a 2 cm gas thickness and 75 atm would have at least

75% quantum efficiency even at the Yb edge based on calculations using the mass attenuation coefficients of Kr.

The spatial resolution achievable in a gas detector is limited by the range of the photoelectrons produced by the incident X-ray beam in the gas. For 75 atm of Kr, 0.25 mm resolution is achievable, even at the Yb K-edge based on calculations using range-energy data for electrons.

The prototype kinesthetic detector was brought from UT, Memphis, to NSLS to demonstrate the spatial resolution capabilities of a Kr gas detector when used with monochromatic synchrotron radiation X-rays. The prototype detector has 256 detecting electrodes with an 0.15 mm center-to-center spacing. The detector was filled with 25 atm of Kr, and the X-ray beam used for imaging had an energy just above the iodine X-edge at 33.169 keV. Images were made with various test objects moving at 12 cm/sec on the patient chair, with a data acquisition rate sufficient to achieve 0.25 mm resolution in the scanning direction. Images taken of bar pattern test objects, phantoms containing dilute iodine containing contrast agent, and of a human vertebral column demonstrated spatial resolutions up to 0.14 mm and high contrast resolution.

The assistance of T. Hallahan at UT and R. Green at NSLS in the preparation of the prototype detector tests is greatly appreciated. These tests were performed under the auspices of the US Department of Energy under contract No. DE-AC02-76CH-00016 to Brookhaven National Laboratory, and under the auspices of the University of Tennessee, Department of Biomedical Engineering.

X17C DIFFRACTION FROM SINGLE CRYSTALS OF SUB-ATTOLITER VOLUME AND A NEW PHASE OF BISMUTH

E. F. Skelton (NRL), L. W. Finger (CIW), S. B. Gadri (NRL), J. Z. Hu (CIW), J. D. Ayers, (NRL), and H. K. Mao (CIW)

Previously, we reported x-ray diffraction from a sub-femtoliter volume single crystal of Bi.^{1,2} We have continued these studies and successfully examined similarly prepared samples with smaller diameters: 0.091, 0.067, and 0.042 μ m. We have obtained useful diffraction data from a crystal volume of 28 attoliters; the scattering power, as defined by Eq. (1) in Ref. 2, associated with our weakest detectable reflection from the smallest sample, is 1.8×10^{10} .

One question that interested us appertained to the continuity of these single crystals. In the 0.091 μ m diameter filament, we followed the (1,0,4)-peak along the full length of the sample, approximately 4 mm. Although this does not ensure that the sample is a single crystal of this length, since we find that the Bi-1 (1,0,4) planes always grow approximately parallel to the fiber axes. In Fig. 1, three rocking curves of the (1,0,4) peak from the same sample are plotted, each recorded at a different position along the length of the fiber. They each have a half-width of about 0.3° in ω ; they also each exhibit a different grade of crystal quality, suggesting a variation of defects or the presence of small angle grain boundaries along the length of the fiber.

Fig. 1: Rocking curves of the (1,0,4)-peak as measured at three different positions along the length of the sample

Single crystal data collected in 19 diffraction spectra from the 0.067 μ m diameter sample are incompatible with all known structures of Bi. We believe this to be a new phase of E, which we have tentatively identified as Bi-X, since nine phases of Bi have already been reported.³ Bi-X has a face-centered cubic lattice and the crystal structure has been determined. The details of this structure determination will be reported elsewhere.

¹E. F. Skelton, H. K. Mao, L. W. Finger, S. B. Gadri, J. Z. Hu, N. E. Moulton, and J. D. Ayers, NSLS Annual Rept. 1990, p. 290

²E. F. Skelton, J. D. Ayers, S. B. Gadri, N. E. Moulton, K. P. Cooper, L. W. Finger, H. K. Mao, and J. Hu, Science **253**, 1123 (1991)

³C. G. Homan, J. Phys. Chem. Solids **36**, 1249 (1975).

X17C
A METHOD FOR PROBING VARIATIONS IN δ IN $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ SINGLE CRYSTALS USING ENERGY DISPERSIVE DIFFRACTION

E. F. Skelton (NRL), S. B. Qadri (NRL), J. Z. Hu (CIW), M. S. Osofsky (NRL), J. L. Cohn[†] (NRL), and T. A. Vanderah (NWC)

A method is proposed for assessing the variance in δ in single crystals of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$. The basic idea is to employ the empirical relationship based on the work of Parks et al.¹ augmented by our own studies,² whereby δ has been correlated with the c-axis lattice parameter through the following empirical equation: $c(\text{\AA}) = 12.8714 - 0.1728(7-\delta)$. A single crystal of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ was mounted on a four-circle diffractometer and illuminated with the polychromatic radiation emanating from the superconducting wiggler on X17. The incident beam was apertured to a $30 \mu\text{m} \times 10 \mu\text{m}$ cross-section; the scattered radiation was analyzed with an intrinsic Ge. energy-sensitive detector equipped with a receiving collimator with a square window $50 \mu\text{m}$ on a side and set at a fixed 2θ angle of 15° . The (0,0,l) diffraction spectra ($l \leq 19$) were recorded in the usual fashion;³ an example is shown in Fig. 1. These spectra have been measured as a function of spatial position across the face of the crystal; they will be examined in search for changes in the c-axis between the central region and the edges. Each of the (0,0,l) peaks in the spectrum can be used to calculate c. We find systematic variations in these values which we believe to be due to two effects, both related to the increased photon energies associated with the higher order reflections: (1) small displacements of the effective scattering center and (2) possible variations in the oxygen content between the surface region and the interior. We believe this may be a novel method for assessing variations in δ with a spatial resolution on the order of $10 \mu\text{m}$ in these $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ single crystals.

Fig. 1: Spectrum of (0,0,l) reflections from a $30 \mu\text{m} \times 10 \mu\text{m}$ area of a single crystal of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$

[†]Postdoctoral Associate from Office of Naval Technology

¹M. E. Parks, A. Navrotsky, K. Mocala, E. Takayama-Muramachi, A. Jacobson, and P. K. Davis, J. Solid State Chem. **79**, 53 (1989).

²M. S. Osofsky, J. L. Cohn, E. F. Skelton, M. M. Miller, R. J. Soulen, Jr., S. A. Wolf, and T. A. Vanderah, to be pub.

³E. F. Skelton, Acta Cryst. **A 32**, 467 (1976)

X18A

EARLY STAGES OF PRECIPITATION OF δ' IN Al-3.2 wt% Li

V. Mahadev, K. Mahalingam and G. L. Liedl, Purdue University.

A study of the early stages of δ' phase formation in a binary Al-3.2 wt%Li alloy was performed by X-ray diffuse scattering experiments. The study was conducted on single crystal samples of the alloy in the as quenched state and after short isothermal aging treatments at 433K. The short range order parameters and the average atomic displacements were determined. Measured values of the displacements are very small, indicating the strains associated with the precipitates are very small. The measured values of the short range order parameters indicate that the δ' phase has a very high degree of order in the as quenched state itself. A simple model was proposed wherein perfectly ordered δ' particles were distributed in a random α matrix. The model included as parameters the volume fraction of δ' and the mean particle size. A Gaussian distribution of sizes from a fourth of the mean to three times the mean was assumed. The diffuse intensity due to ordering was calculated using kinematical theory of x-ray diffraction. The short range order parameters determined by Fourier inversion of calculated diffuse intensities from the simple model was within statistical uncertainty of the experimentally determined values. These results indicate that the δ' particles possess a high degree of order even in the as quenched state. The presence of the δ' phase in the as quenched state may suggest that the δ' is formed by spinodal decomposition. But the absence of satellite peaks around the diffraction peaks disallows spinodal decomposition as the mechanism of precipitation of δ' . Further, the observed increase in the volume fraction of the δ' indicates a growth behavior. One possible explanation for the existence of the δ' in the as quenched state may be the existence of precursors at high temperatures which aid in the rapid precipitation of the δ' phase. Such a precursor could be a strong tendency for clustering similar to the ordering configuration of aluminum and copper at high temperatures. However, the rapid formation of the δ' phase is possible even in the absence of such precursors because of the large diffusivity of the lithium atoms in the alloy. In either case our study indicates a nucleation and growth type of formation for δ' in the early stages. This research was sponsored by DOE (contract No. DE-FG02-85ER 45813).

A STUDY OF SUBSTRATE-INDUCED ALIGNMENT OF SMECTIC LC PHASES: PRESENCE OF SURFACE OR SUBSTRATE STABILIZED PHASES NEAR THE LC-AIR INTERFACE

L. J. Martínez-Miranda and E. Smela (U. of Penn.)

The molecular arrangement in a liquid crystal film differs in the bulk compared to surfaces and interfaces. The ordering in LC films is in general stronger than in bulk samples, and in freely suspended films a quasi-smectic phase appears. An LC film deposited onto a grooved substrate experiences a competition between the aligning forces at the LC-air interface, which force the molecules to align normal to the surface, and the aligning forces at the LC-substrate interface, which forces the molecules to align parallel to the plane of the substrate.

The aim of our project is to expand the studies of LC films deposited on weak interaction substrates. We are interested in studying two aspects of this problem: a) how the LC alignment and structural conformation vary as a function of increasing distance from the air-LC interface to the LC-substrate interface; and b) how the systematic change in the parameters of the aligning grooved substrate (e.g., the period between grooves, the groove depth and thickness) affect the LC molecular alignment. The substrates were prepared by chemically etching gratings into glass slides. These gratings were treated with a silane before deposition of the LC film. The liquid crystal used is cyanobiphenyl (8CB), which exhibits two mesophases in the bulk: a smectic A₂ with a layer spacing of 31.6 Å, and a nematic. Our studies at beamline X-18A used 12keV x-rays, focused to a spot size of 4mm² (first run) and 2mm² (second run) at the sample position. We performed a series of experiments on samples with different grating periods, with "egg-carton" surfaces, and with flat surfaces. In our results from the inplane scattering, that is, from molecules parallel to the surface, θ -2 θ scans consistently showed diffraction peaks from molecules which were lying parallel to the grating with a layer spacing of approximately 33.5 Å. These peaks appeared at incident angles between $\theta = 0.2$ to 0.4° . We never saw diffraction perpendicular to the grating, nor did we observe these peaks from LC films on flat or "egg-carton" surfaces. There is very strong evidence, therefore, that these gratings induced alignment in some regions of the films. Furthermore, our most recent results indicate that the onset and range of parallel alignment depend on the film thickness, with thinner films exhibiting purely perpendicular alignment. All cut of plane scans in all films show the appearance of a peak centered at 2.1° , which corresponds to a layer spacing of 28.2 Å. There is evidence that this peak evolves from a layer spacing of 30.3 Å to its final value for angles of incidence that vary from 0° to 0.5° . This peak reappears close to the glass interface.

Our future studies will concentrate on the detailed study of the effect of film thickness and grating size on the presence and range of in-plane peaks, as well as on the evolution of the 28.2 Å out of plane peak.

This research was supported by the IBM Program of Support for the Materials and Processing Sciences Grant, NSF-MRL Grant DMR-88-19885, and the NSLS Faculty-Student Research Support Program.

INVESTIGATION OF THE LOCAL ATOMIC STRUCTURE IN HgCdTe

D.Q. Na, J.B. Cohen and P. Georgopoulos, Dept. of Materials Science and Engineering, Northwestern U., Evanston, IL.

Semiconductors based on the pseudobinary compound Hg_{1-x}Cd_xTe have been studied since the 1960's for use in infrared detectors and solar cells. EXAFS measurements have shown^{1,2} that the Hg-Te and Cd-Te bond lengths in these systems are relatively constant throughout the entire composition range despite the change in lattice parameter and that Hg-Hg, Hg-Cd and Cd-Cd distances are equal to the average lattice parameter of the alloy. Durbin's³ standing wave results on Cd_{0.72}Mn_{0.28}Te, however, suggest that the distances are unequal. Quintana⁴ in a recent study has found diffuse peaks in the diffraction pattern of both bulk Hg_{1-x}Cd_xTe and Cd_{1-x}Mn_xTe, indicative of significant local displacements, as well as local order.

A diffuse scattering study was undertaken to help resolve some of these discrepancies and to explore the local structure of this class of materials more fully. A diffuse peak was observed in all specimens examined, which included two thin film HgCdTe samples grown on differently oriented substrates (Figs 1 and 2), as well as one thin film of InGaP (Fig 3). Data analysis is still in progress.

1. Mayanovic, R.A., W.-F. Pong, and B.A. Bunker (1990). Phys. Rev. B, 42, 11174-11182
2. Balzarotti, A., N. Motta, A. Kisiel, M. Zinnal-Sternawska, M.T. Czyrk, M. Podgurny. (1987). Phys. Rev. B 31, 7626-7639.
3. Durbin, S.M. (1988). J. Appl. Phys. 64(5), 2312-2315.
4. Quintana, J.P. (1991) PhD Thesis, Northwestern U.

Figure 1

Figure 2

Figure 3

*This research was sponsored by NSF

X-RAY CHARACTERIZATION OF 1.0 nm SN FILM ON SI (111)

K.F. Peters, J.B. Cohen and P. Georgopoulos, Dept. of Materials Science and Engineering, Northwestern U., Evanston, IL.

Tin was evaporated by resistive heating from a tantalum boat onto a polished Si single crystal (111). Both surfaces were rinsed in de-ionized water and isopropyl alcohol. The deposition was monitored and interrupted after 1.01 nm nominal thickness. Atomic force microscopy (AFM) revealed that Sn formed islands with typical diameters of 20-60 nm and height of 15 nm.

These samples were evaluated at the Brookhaven National Laboratory, National Synchrotron Light Source, MATRIS beamline X18A on a Huber 4-circle diffractometer using normal diffraction geometry. An x-ray energy of 7100 eV ($\lambda = 1.74621 \text{ \AA}$) was selected. In all, about 25 2θ , χ and ω scans were performed to characterize six different Sn peaks.

It was quickly determined that the x-ray scattering from 1.0 nm of Sn allows clear and easy determination of integrated intensities. The accompanying figure shows three different 2θ scans in the vicinity of the Si 111 peak. One scan goes through the Si 111 maximum; the two others represent tilts of $\pm 1^\circ$ in χ . Sn 200 is clearly visible on the tail of the Si peak.

The Sn 200 intensity was found to be sensitive to changes in χ , and its maximum lies at a different χ value than the Si 111 peak. The χ sensitivity indicates a strong orientation relationship with the substrate, which is the reason for the strong intensity of the Sn 200 peak. Further, since Sn 200 and Si 111 have slightly different χ values the corresponding crystal axes are not exactly parallel.

Applying the Scherrer equation, the particle size was found to be 1.41 to 1.49 nm for the 400 and 200 Sn peaks, respectively. The difference from the gravimetric thickness measurement of 1.01 nm during deposition can probably be attributed to the island morphology of the Sn film.

This research was sponsored by DOE

X-RAY STUDY OF SUBSTRATE-INDUCED ALIGNMENT OF A SMECTIC-A LIQUID CRYSTAL X18A

E. Smela and L. J. Martinez-Miranda (U. of Penn.)

We performed a structural study of the liquid crystal (LC) octyloxybiphenyl (8CB), deposited on gratings, grids and flat surfaces, using high resolution x-ray scattering at low incidence angles. 8CB is a room temperature smectic A γ with a layer spacing of 31.6 Å. Glass and silicon were used as substrates. These were treated with either one of the organic surfactants MAP or DMOAP. Surface tension forces cause the liquid crystal molecules to align perpendicularly with respect to the plane of the substrate at the air interface. Competing with the LC-air interface, which is a strong aligner, a grating at the LC-substrate interface produces distortions in the smectic layering with an excess elastic energy, which favors alignment parallel to the substrate and the grooves. The studies used 12keV x-rays focused to a spot size of 2mm^2 at the sample position.

Out-of-plane scans were done on most of the samples, showing very bright, resolution-limited peaks at a 2θ angle of 1.88° , corresponding to $q = 0.199 \text{ \AA}^{-1}$. This scattering geometry detects layers parallel, or molecules perpendicular, to the plane of the substrate. The second harmonic of this peak was quite strong also. The inplane scans, which detect the smectic layers perpendicular to the plane of the substrate, were done at angles $\phi = 0^\circ$ and 90° with respect to the gratings to ascertain the molecular orientation. None of the samples with a film thickness of less than approximately 30 μm showed any in-plane scattering. This was true regardless of the surface pattern. Furthermore, the substrate material and surface treatment had no effect on the results.

Films on the flat surfaces had very small inplane peaks: the height was only twice the background noise. The peaks appeared at both $\phi = 0^\circ$ and 90° , showing random azimuthal alignment. Like the flat surfaces, the grid of holes showed small inplane peaks at both $\phi = 0^\circ$ and 90° , although the peaks were slightly larger.

In stark contrast, every sample consisting of a film greater than 30 μm on a grating showed inplane scattering at $\phi = 0^\circ$, and either absolutely no scattering or scattering at substantially lower intensity at $\phi = 90^\circ$. This is illustrated in Figure 1. The magnitude of the inplane scattering was at least 50 times smaller than it was in the out of plane scans, and the peak positions were shifted consistently to slightly lower q than in the out of plane scans. We observed the inplane Bragg peak even at incidence angles close to the critical angle (0.16°) for 8CB, which implies that regions of parallel alignment may exist close to the air interface. It is not clear, though, if the inplane scattering is due mostly to numerous small regions distributed inside the film, or to one very thin region of greater extent located deeper within the film.

This research was supported partially by the IBM Program of Support for the Materials and Processing Sciences Grant, NSF-MRL Grant DMR-88-19885, and the NSLS Faculty-Student Research Support Program.

Figure 1. Inplane scattering from 8CB on a 9 μm grating etched on Si: a) 17 μm thick film; b) 67 μm thick film.

MULTILAYER STRUCTURE AND GROWTH OF XENON ON Ag(111)

S.-K. Wang^(a), P. Dai^(a), S. N. Ehrlich^(b), T. Augot^(a), J. Z. Larese^(c), and H. Taub^(a)
 (a) U. of Mo., Columbia; (b) Purdue U.; (c) BNL

Currently, there is considerable interest in the multilayer structure and growth of physisorbed films on solid substrates.¹ Basic issues of concern are the layer-by-layer structure of the solid film and its ultimate thickness. We have undertaken a series of experiments to address these questions in the case of xenon films adsorbed on the Ag(111) surface. Comprehensive low-energy electron diffraction (LEED) experiments on rare gas films by Webb and co-workers² have shown the Ag(111) surface to be of prototypical quality for physisorption studies while Xe provides a strongly scattering film for x-ray diffraction.

Our experiments utilize an ultra-high vacuum chamber which has been especially designed for x-ray diffraction from physisorbed films.³ The chamber mounts on the γ -circle of the standard Huber diffractometer at the X18a beam line. It has a low-temperature capability ($T \geq 28$ K) and features a low-current LEED system which allows diffraction patterns to be obtained without moving the sample from the x-ray scattering position. The LEED system can be conveniently used to perform an *in situ* coverage determination. The Ag(111) substrate is dosed with Xe at a constant rate and a step-wise attenuation of the Ag(10) LEED spot is observed upon condensation of the first and second solid Xe layers.²

Our preliminary x-ray results are summarized in the figure at the left comparing the Xe-Xe atom-spacing in the film (circles) and the bulk⁴ (triangles) vs temperature. The film data is taken cooling while dosing at a pressure of $\sim 3 \times 10^{-7}$ Torr. We see that upon monolayer completion the Xe-Xe spacing has reached the bulk value to within the experimental uncertainty. As verified in other runs, there is virtually no thermal contraction of the film after adsorption of the bilayer. Upon cooling below 50 K, a Xe (111) Bragg peak appears indicating nucleation of isotropically-oriented bulk particles. We have not yet conducted out-of-plane scans to determine the film thickness. However, an intensity increase in scans parallel to the surface through the Xe (10) rod at temperatures just above where the bulk peak appears suggests that the film thickness exceeds three layers.

¹See, for example, G. B. Hess, in *Phase Transitions in Surface Films 2*, edited by H. Taub, G. Torzo, H. J. Lauter, S. C. Fain, Jr., NATO ASI Series, Vol. B 267 (Plenum, New York, 1991), p. 357.
²J. Unguris, L. W. Brich, E. R. Moog, and M. B. Webb, *Surf. Sci.* **87**, 415 (1979).
³J. R. Dennison, S.-K. Wang, P. Dai, T. Augot, S. N. Ehrlich, and H. Taub, in preparation.
⁴M. L. Klein and J. A. Venables, eds., *Rare Gas Solids*, Vol. II (Academic Press, New York, 1977).

Supported by NSF Grant Nos. DMR-8704938 and DMR-9011069, DOE Grant No. DE-FG02-85ER45183 of the MATRIX PRT at the NSLS, and DOE Division of Materials Science Contract No. DE-AC02-76CH00016.

IN SITU X-RAY DIFFRACTION STUDY OF YBCO THIN FILMS

J.O. Zheng, S. Williams, X.K. Wang, T. Rippert, S. Maglic,
 D. Segel, R.P.H. Chang, and J.B. Ketterson
 Northwestern University
 T. Roberts, Y. Lin, R.T. Knapik, and K. Gray
 Argonne National Laboratory

The initial experiments we conducted at the NSLS used *in situ* x-ray diffraction to study the effects of substrate temperature, deposition rate, and oxygen partial pressure on the growth of YBCO films on SrTiO₃. The most recent work examined YBCO growth on LaAlO₃ and MgO.

A substrate temperature study was conducted with YBCO films on LaAlO₃ with T_s ranging from 660°C to 765°C. The other parameters were held at the same values used for films grown on SrTiO₃ (deposition rate 1 Å/s, 0 partial pressure 30 mTorr, Ar partial pressure 60 mTorr). The films grew with a c-axis orientation much more readily than those grown on SrTiO₃. When the substrate temperature was only 660°C, c-axis growth was comparable to a-axis growth until the film reached a thickness of about 2500 Å. At temperatures of 730°C and higher, no a-axis growth was observed. The c-axis films were tilted relative to the substrate by 15° to 20° . *In plane* x-ray diffraction studies reveal only the (110) peak was present (Figure 1). This indicated that the films were of a single crystal-like nature.

The thermal expansion of YBCO films was also examined. It was found that the a-axis of the films expanded at roughly the same rate as LaAlO₃, while the c-axis expanded at twice the rate.

Two different studies were made of the growth of 18a-00y on (100) MgO substrates. The first set of experiments varied the deposition rate, while holding all other parameters constant. The lowest sputtering rate used was .3 Å/s and the highest was 6 Å/s. C-axis growth was dominant at all rates. The rocking curve of the film steadily improved as the deposition rate was increased. The film peaks seem to show a slight increase in angle as the rate is increased, indicating the lattice parameter is shrinking. However, the data is not conclusive.

The second study varied the substrate temperature while the other parameters constant. C-axis growth was observed at temperatures above 600°C, a mixture of a and c-axis growth occurred at 550°C, and only a-axis growth occurred between 500 and 525°C. Figure 2 shows the growth of the (006) peak at 670°C. In contrast to data obtained on SrTiO₃ and LaAlO₃, the peak position does not shift as the film thickens, indicating that any stress that occurs at nucleation is not being relieved as the film grows.

Figure 3 shows the growth of the (006) peak for various substrate temperatures. The lattice parameter is becoming smaller as the temperature increases. This is consistent with a study conducted by Michikami and Asahi in which the substrate position was varied relative to the sputtering guns. When the substrate was near the guns, the lattice parameter was closer to that of the bulk value of YBCO. When the substrate was farther away, the lattice parameter was closer to MgO. Higher kinetic energy, whether due to a high substrate temperature or proximity to the guns, appears to enable the lattice parameter of the film to approach the bulk value.

1. Osamu Michikami and Masayoshi Asahi, YBCO Film Formation by Magnetron Sputtering with Facing Targets, *Japanese Journal of Applied Physics*, vol. 30, No. 5, pp. 939-44, May 1991.

Research sponsored by the NSF Science and Technology Center for Superconductivity.

X18-A

Figure 1

Figure 2

Figure 3

X-Ray Reflectivity Study of Si/C Multilayers*

J.M. Bai, Y.L. Chan, H. Homma (Brooklyn College of CUNY), and P.A. Montano (Argonne National Laboratory).

We have measured using x-ray reflectivity the structure of Si/C multilayers. The multilayers were prepared using the rf-sputtering technique. The modulation wavelengths of the samples were from 14 to 62 Å, and the total thicknesses between 900 to 2000 Å. We used sapphire and float glass as substrates for the samples. The x-ray energy was 24.5 Kev, see figure 1.

We employed in our analysis of the data the matrix formalism developed by B. Vidal and P. Vincent [1] for the x-ray specular reflectivity of multilayers. The analysis of the data gave us information on the electron density profile and roughness at the interfaces. We found that the samples grown on a sapphire substrate possess a sharper interface than those grown on float glass substrates. We observed the formation of a carbide phase at the silicon-carbon interface for samples heated in UHV to 873 K.

REFERENCE

- [1] B. Vidal and P. Vincent, Applied Optics, Vol. 23, No. 11, June 1984.

*This work is supported by the U. S. Department of Energy, BES-Materials Sciences, under contract No. #W-31-109-ENG-38.

X-Ray Reflectivity Study of Ti/C and W/C Multilayers*

Y. L. Chan, J. M. Bai (Brooklyn College, City University of New York) P. A. Montano (Argonne National Laboratory), E. Ziegler (ESRF), I. K. Schuller (University of California, San Diego).

We have investigated the structure of W/C and Ti/C multilayers over a wide range of periodicities using x-ray reflectivity measurements. The samples were prepared by plasma rf sputtering techniques on a Si(111) substrate and the modulation and lattice structure were determined by x-ray diffraction and electron microscopy. Electron energy loss spectroscopy was used to obtain a non-destructive composition depth profile of the material by changing the impinging electron energy. The measurements were carried out under UHV conditions. All the major electron energy loss peaks of the multilayers were identified. The measurements were performed as a function of temperature, between 293 and 973 K.

We employed in our analysis of the data the matrix formalism developed by B. Vidal and P. Vincent [1] for the x-ray specular reflectivity of multilayers. The analysis of the data gave us information on the electron density profile and roughness at the interfaces. We observed the formation of a carbide at the W/C interface. The structure of the interface in Ti/C does not show the presence of a carbide phase, there is strong evidence of the formation of thin titanium oxide interface in all the Ti/C multilayers. We also investigated the stability of the multilayers in an oxidizing environment and at high temperatures.

X-ray Reflectivity of a Ti/C Multilayer

X-ray Reflectivity of a W/C Multilayer

*This work is supported by the U. S. Department of Energy, BES-Materials Sciences, under contract No. #W-31-109-ENG-38.

R.A. Prokopowicz, A. Stein, G.A. Ozin, (U of Toronto)

A series of *novel* sodalite-based materials has been synthesized by Ag^+ melt ion exchange of hydrothermally prepared sodium halo-sodalites (this framework is constructed of entirely space filling bcc cubo-octahedral β -cages). In terms of the unit cell structure of the host lattice (two β -cages per unit cell, each of which can contain a halide anion and four cations arranged in a tetrahedron with the anion at the centre), the materials can be described by the general formula $\text{Na}_4\text{Ag}_{(4-x)}\text{X}_x\text{SOD}$ (where $x = 0.8$; $n = 0.2$; and $\text{X and X}^+ = \text{Cl, Br, I}$).

As a class of materials, these represent monodispersed clusters of bulk semiconductor species within an insulating aluminosilicate framework. In mixed halide systems, one can produce monodispersed narrow band-gap semiconductor clusters embedded within a supralattice of a wide band-gap semiconductor clusters. There also exists an ability to control the concentration and spatial distribution of various β -cage species, for example the extent of Ag exchange of the original Na cations, and/or the concentration of Ag, Br or Ag, I tetrahedra in a mostly Ag, Cl supralattice. As a result of being able to control the extent and nature of electronic coupling between these encapsulated clusters, such materials might function as resonant tunnelling quantum dot transistors and high gain heterojunction multiple quantum dot laser arrays. Another possible area in which this class of materials is expected to be significant, is that of reversible optical data storage. In principle, through the control of the compositional parameters listed above, these materials could be used to achieve an improvement both in storage density and switching speed over existing technology.

This group of materials has been examined by several techniques such as: bulk elemental analysis, electron microscopy, mid and far-FTIR, UV-Visible and Laser Induced Luminescence spectroscopies, DOR-MAS-NMR, and High Resolution Powder XRD with Rietveld Refinement of the data. In order to gain further structural information, and especially to resolve ambiguities in the Rietveld results, EXAFS has been used to examine a group of five samples. These are: (1) a fully Ag exchanged mixed chloro-iodo-sodalite (45 at% I, 55 at% Cl); (2) a fully Ag exchanged sample containing 75% Ag, 25% Ag, Br clusters; (3) a low silver bromo sodalite (0.3 Ag, 7.7 Na per unit cell); (4) a fully Ag exchanged sample containing only Ag, Br clusters; and (5) a silver bromo-sodalite having incomplete Ag exchange (i.e. 2.5 Ag and 5.5 Na per unit cell). Data were collected for the Ag K-edge only. In the mixed halide material for example, two Rietveld refinements gave equally adequate structural determinations: one having two different Ag-X distances, and another in which both silver halide clusters were found to have the same distance (the average of the two distances in the other model). EXAFS results gave evidence that the former model was most appropriate, and confirmed many of the other structural details of the Rietveld refinements such as silver-halide and silver-framework oxygen distances. It was further determined that the clusters found in mixed halide systems (eg. Ag, I and Ag, Br) had structures which were virtually identical to those existing in samples containing only one halide.

Funding for this project was provided by the Natural Sciences and Engineering Research Council of Canada.

EXPERIMENTAL CHARACTERIZATION OF CIRCULAR FRESNEL ZONE PLATES

W. Yun, J. Chrzas, B. Lai, D. Legnini, J. Viccaro (ANL), R. Bionta, K. Skulina (LLNL)

Development of microfocusing x-ray optics of submicron spatial resolution and high focusing efficiency will have a significant impact on many microfocusing-based x-ray techniques, such as microimaging, microanalysis, microdiffraction, microspectroscopy, and holography. We have established a collaborative effort developing transmission Fresnel zone plates (TFZP) for hard x-ray focusing applications. Here we present the experimental characterization of a TFZP of 40-cm focal length. The TFZP, consisting of alternate Cu and Al zones, was fabricated by a sputtering/slicing technique. The parameters of interest are: (1) the spatial resolution of the TFZP, (2) the focusing efficiency as a function of x-ray energy, and (3) important zone plate geometries, such as zone plate thickness, aspect ratio of zones in a period, and accuracy of zone registry. A large amount of data necessary for characterization of the TFZP was obtained; data analysis is in progress. In Figs. 1 & 2 we present some highlights of the experimental results. Fig. 1 is an image of a Au grid obtained at the 3rd order focal plane using 8.2 keV x-rays. The image indicates that a spatial resolution of about 2 microns was obtained using the TFZP. The spatial resolution is currently limited by the geometrically demagnified image of the effective x-ray source size, that is by the electron beam size in the storage ring. This limitation can be removed by installing a small aperture upstream of the zone plate. A spatial resolution of about 0.5 micron may be possible when a proper aperture is installed. Fig. 2 shows the focusing efficiency of the zone plate as a function of x-ray energy measured in the energy range between 5-11 keV. A maximum focusing efficiency of about 21% was obtained at 8.8 keV x-ray energy. Work is supported in part by the Department of Energy, BES-Materials Science, under contract W-31-109-ENG-38.

Fig. 1

Fig. 2

In Situ Investigation of Passive Films on Fe-Cr Alloys¹

J.A. Bardwell (NRC), A.J. Davenport (BNL), H.S. Isaacs (BNL), G.I. Sproule (NRC), B. MacDougall (NRC) and M.J. Graham (NRC)

The valence state of chromium in passive oxide films on Fe-Cr alloys has been successfully studied with XANES *in situ* in an electrochemical cell with the sample under potential control. The cell design is similar to that used to study Al-Cr alloys.² At low potentials, Cr(III) is present in the oxide film. As the potential is increased, the appearance of the distinctive Cr(VI) pre-edge peak was correlated with the "transpassive" increase in current found in the cyclic voltammogram. Its disappearance coincided with the corresponding reduction wave. Figure 1, below shows XANES spectra obtained at potentials below and above the transpassive peak. The formation and reduction of Cr(VI) was reversible, and only small amounts were detected in the film.

The Cr(VI) in the passive film is not indefinitely stable, and is completely absent after a number of hours of air exposure. This result is shown in Figure 2, where the pre-edge peak is completely absent after 16.5 hr. of air exposure. This demonstrates the importance of carrying out *in situ* experiments to study the chemistry of passive films present on metals in aqueous environments.

Fig. 1. *In Situ* XANES spectra of a Fe-26Cr thin film electrode under potential control. The broken line shows the position of the pre-edge peak characteristic of Cr(VI).

Fig. 2. *In Situ* and *ex situ* XANES spectra of a Fe-26Cr thin film electrode after various times of air exposure.

1 "In Situ XANES Detection of Cr(VI) in the Passive Film on Fe-26Cr", J.A. Bardwell, A.J. Davenport, H.S. Isaacs, G.I. Sproule, B. MacDougall, and M.J. Graham, submitted to J. Electrochem. Soc.

2 A.J. Davenport, H.S. Isaacs, G.S. Frankel, A.G. Schrott, C.V. Jahnes and M.A. Russak, J. Electrochem. Soc., 138, 337 (1991).

Acknowledgement: This work was carried out in part under the auspices of the US Department of Energy, Division of Materials Sciences, Office of Basic Energy Science under Contract No. DE-AC02-76CH00016.

Fe K-EDGE EXAFS STUDIES OF NITROGENASE.

J. Chen (LBL), S.J. George, J. Christiansen, and S. P. Cramer (UC Davis), R. Titusworth and B. Hales (LSU).

Nitrogenases are enzymes responsible for the biological fixation of dinitrogen to ammonia. Their active center comprises an iron-sulfur cluster of unknown structure which contains one other metal, either Mo, V or Fe. We have probed the structure of the Mo form of this enzyme using Fe K-edge x-ray absorption spectroscopy.

Figure 1 shows the Fourier transformed data of 1.5 dithionite reduced and thionine oxidized MoFe proteins (transform range: $k = 2-15.3 \text{ \AA}^{-1}$). For both oxidized and reduced samples, the spectra are similar and are dominated by Fe-S and Fe-Fe interactions at 2.29 $\text{\AA}$ and 2.61 $\text{\AA}$, as well as the long Fe-Fe distance 3.7 $\text{\AA}$. Preliminary analysis also suggests that there are some longer interactions at 4.2 $\text{\AA}$ and 4.7 $\text{\AA}$.

Figure 2 shows curve-fitting results for the reduced sample. An r -space window from 1 to 3 $\text{\AA}$ is used in the back transformation to filter out the longer shell EXAFS contributions. The fit is carried out by using previously described programs and functions (1). The best fit was obtained with four components, Fe-O, Fe-S, Fe-Fe, and Fe-Mo at 1.74, 2.29, 2.61, and 2.75 $\text{\AA}$, respectively. Compared with the oxidized sample, the Fe-S distance is a little bit longer in the reduced protein. We think that this is probably due to the weaker Fe-S bond strength in the reduced sample.

This work was supported by DOA and NIH under grants no. DOA-91-37305-6514 and NIH-GM-44380-2.

Figure 1. Fourier transformed data of nitrogenase samples.

Figure 2. Experimental and fitted EXAFS for reduced MoFe protein. (a) Spectra, (b) Fourier transforms.

(1) Cramer, S. P. et. al.; J. Am. Chem. Soc. 1984, 106, 1467-1471.

BEAM LINE X-19A

X-RAY ABSORPTION STUDIES OF 4d TRANSITION METAL ELEMENTS

J.Chen, Y.Jeon, F.Lu and M.Croft
 Department of Physics, Rutgers University, Piscataway, NJ 08855
 P.H.Ansari
 Department of Physics, Seton Hall University, South Orange, NJ 07079

Work by our group[1] and others has shown that the L_2 and L_3 "white line" (WL) strength in the late 5d-row transition metal elements scales with the 5d-orbital hole count. Moreover these elemental studies proved to be of great use in determining bonding induced 5d-orbital occupation changes in compounds[2]. We are extending these studies to the 4d-row transition metal elements, and show their L_3 (figure 1) and L_2 (figure 2) near edge spectra below.[All spectra are normalized to the continuum.] The increase in WL strength with increasing 4d-hole count between Ag and Mo is both systematic and dramatic. Closer inspection reveals that the L_3 -WL is stronger than the L_2 -WL feature in Rh and Pd, consistent with spin-orbit induced anisotropy in $d_{5/2}$ to $d_{3/2}$ hole counts near top of the 4d band. These results are in analogy with our 5d-row studies[1], however the generally higher resolution and in some cases the resolved WL features stand in contrast to the 5d-row studies. These elemental studies will be used as standards for compound studies.

Reference
 [1] Boyun Qi, I.Perez, P.H.Ansari, F.Lu and Mark Croft, Phys.Rev.B 36,2972(1987)
 [2] Y.Jeon, B.Qi, F.Lu and M.Croft, Phys.Rev.B 40,1538(1989)

X19A

IN SITU XANES STUDY OF Cr VALENCY CHANGES IN PASSIVE OXIDES ON AlCr THIN FILMS UNDER ELECTROCHEMICAL CONTROL

A. J. Davenport (BNL), H. S. Isaacs (BNL), G. S. Frankel (IBM), A. G. Schrott (IBM), C. V. Jahnes (IBM), and M. A. Russak (IBM)

The valence state of chromium in passive oxide films on sputtered AlCr in aqueous environments has been studied *in situ* using XANES¹. The alloy was present as a thin film ~40Å in order to minimize signal from the underlying metal. Measurements were made in fluorescence geometry in the cell shown in Fig.1. A 13-channel solid state detector (Canberra) was used to detect Cr Kα radiation to discriminate against the scattered incident beam. Initially, chromium was present in the metallic state (Fig.2(a)). When a high potential was applied (2V vs. mercurous sulfate electrode), it was oxidized to the 6-valent state (Fig.2(b)), characteristic pre-edge peak. The soluble 6-valent state was trapped in the film and did not dissolve. The chromium was electroactive and could be reversibly reduced to the 3-valent state by reduction at a low potential (-1.5V).

Fig.1 Schematic plan of experimental apparatus.
 Fig.2 Solid lines: Al-12Cr (a) at open circuit, (b) 2V(MSE), (c) -1.5V(MSE); broken lines (a) Cr metal, (b) K_2CrO_4 (c) Cr_2O_3 .

1 A. J. Davenport, H. S. Isaacs, G. S. Frankel, A. G. Schrott, C. V. Jahnes, and M. A. Russak, *J. Electrochem. Soc.*, **138** 337 (1991).

This work was performed in part under the auspices of the U.S. Department of Energy, Division of Materials Sciences, Office of Basic Energy Science under Contract No. DE-AC02-76CH00016.

MICROBIAL TRANSFORMATIONS OF URANIUM: SPECIATION STUDIES

A. J. Francis, C. J. Dodge and J. B. Gillow (ENL) and F. Lu (U. KY)

Microorganisms play a major role in the oxidation/reduction of various elements in nature and affect their valence state and solubility. Although much is known of the microbial transformations of toxic metals, there is a paucity of information on the transformations of the actinides. Fundamental knowledge of the mechanisms of transformations is needed for development of novel concepts to treat transuranic wastes as well as use in the predictive biogeochemical transport models. Reduction of uranium by an anaerobic bacterium *Clostridium sp.*, isolated from uranium waste, was investigated. The bacterium was allowed to grow in a prereduced nutrient medium containing 0.2M uranyl acetate. The culture medium was centrifuged and the resulting pellet was analyzed for uranium by XANES. Samples were placed on Mylar film and the oxidation state of uranium was determined using an electron yield detector. Standard uranium compounds were analyzed at the following edges: $M_{II} 3p_{3/2} \rightarrow 6d(4299\text{eV})$, $M_{IV} 3d_{3/2} \rightarrow 5f(3720\text{eV})$ and $M_{V} 3d_{5/2} \rightarrow 5f(3545\text{eV})$. The XANES at the M_V edge provided data with least interference (Fig. A). The sample before bacterial treatment (control) showed a maximum absorption at 3551.05eV confirming the uranium 6^+ oxidation state (Fig. B). The sample subjected to bacterial activity showed a shift in the absorption energy to 3550.14eV which is higher than the uranium metal, U (3549.59eV) but lower than uranium dioxide, U^{4+} (3550.34eV). These results show that bacteria catalyze the reduction of soluble U^{6+} to an insoluble U with a lower oxidation state than previously suggested.

T. Francis, A.J., C.J. Dodge, J.B. Gillow, and J.E. Cline. Radiochimica Acta 52/53, 311-316 (1991).

Research was supported in part by Subsurface Science Program and R&D Laboratory Technology Transfer Program, Office of Energy Research, Department of Energy.

X-RAY ABSORPTION MEASUREMENTS ON RT_2Si_2 COMPOUNDS

Y. Jeon, J. Chen, F. Lu, M. Croft, Rutgers University, Piscataway, NJ 08855
S. A. Shaheen, Florida State University, Tallahassee, FL 32306
P. H. Ansari, Seton Hall University, South Orange, NJ 07079

Previously we have studied RT_2X_2 compounds with R = rare earth, T = 5d row element and X = Si or Ge. These studies indicated that $T \leftrightarrow X$ charge exchange occurs in these 1:2:2 materials in a way analogous to that observed in $T-X$ binary compounds [1]. Namely strong $T-X$ bonding leads to shifting of T 5d states above E_F and an increase in the effective T 5d-hole count results. Indeed, the electronic structure calculations of Hoffman and coworkers has emphasized that strongly hybridized/bonded $T-X$ planes are the basic building block of the 1:2:2 structure [2].

We are currently involved in extending such studies to 4d-row compounds. In Fig. 1 we show that L_3 edge of selected RT_2Si_2 compounds, where T = 4d row element. Not only do these results underscore the presence of similar $T-Si$ bonding effects in these 1:2:2 compounds, but the better energy resolution at the 4d L_3 edge allow the identification of partially resolved 4d states that have been split off to higher energies (see feature A in Fig. 1). We tentatively associate these states with the antibonding $T(4d)-Si$ states. In Fig. 2 we have observed similar split-off $T-4d$ states in $PrRhB_n$ compounds ($n = 1, 2$) and attribute them also to antibonding $Rh-B$ states that have been shifted away from the Fermi energy.

Fig. 1. $T-L_3$ spectra for a series of 4d row 1:2:2 compounds. Note the resolved WL feature (A) split away from lowest energy peak.

Fig. 2. $Rh-L_3$ spectra for $PrRh_3B_n$ ($n = 0, 1, 2$) compounds. Note the resolved WL feature (A) split away from lowest energy peak.

References:

- [1] Y. Jeon, B. Qi, F. Lu, and M. Croft, Phys. Rev. B 40, 1538 (1989).
- [2] C. Zheng and R. Hoffman, J. Phys. Chem. 89, 4175 (1985).

Speciation of Uranium Compounds by XANES

F. Lu (U. Kentucky), C. J. Dodge and A. J. Francis (BNL)

Uranium exists in the 3+, 4+, 5+, and 6+ oxidation states, with 4+ and 6+ being the predominant forms. As a highly correlated metal, it also forms non-stoichiometric mixed valence compounds. Speciation of uranium in natural systems and in mixed waste materials provide a basis to determine their stability or mobility in the environment¹.

XANES has been used to study uranium valence states and charge transfer. The core level absorption edge energy shifts reflect charge screening effect, the charge localized at the uranium site, as well as the average valence². M_{IV} and M_V edges correspond to the transition from 3d^{3/2} to the unoccupied 5f levels and are of particular interest. However, the uranium absorption length at these edges is only about 0.5 μm. This results in significant thickness effects in both the fluorescence and transmission modes³.

XANES spectra in the fluorescence and electron yield mode for a selected uranium compound is presented in Fig. A. The thickness effect in the fluorescence mode is evidenced by peak broadening (5.6 eV width). This effect is negligible in the electron yield mode since the secondary electrons were removed from the top 0.1 μm of the sample (3.0 eV width). The peak energy positions in the electron yield mode for uranium metal, UO₂, and UO₃ are 3549.59 eV, 3550.33 eV, and 3551.05 eV, respectively (Fig. B).

1.) Francis, A.J. *Experientia* 46, 840-851 (1990).2.) Kalkowski, G., G. Kaindl, M. Brewer, and K. Krone. *Phys. Rev.* B35, 2667 (1987).

3.) F. Lu, et al. To be published.

Research was supported in part by Subsurface Science Program and R&D Technology Transfer Program, Office of Energy Research, Department of Energy.

MULTI-SHELL FITTING OF EXAFS DATA IN COMPLEX VANADIUM COMPOUNDS

F. Lytle (Boeing), M. Thompson (Pacific Northwest Laboratories) and R. Friedman and J. Ebner (Monsanto)

A major goal of this research is to accurately measure the V-O bond distances in the first coordination shell of various V⁴⁺/V⁵⁺ vanadium phosphate catalysts. The presence of the short vanadyl bond creates distorted coordination shells and a severe test of the EXAFS technique. This report discusses the accuracy that we have demonstrated to date. We were successful in deriving from V₂O₅ an accurate phase shift for the V-O atom pair and O backscattering envelope so that distance determination by least squares fitting could be attempted. Reasonable EXAFS data were available for the reference compounds of known structure listed in Table 1. VPO is (VO)₂P₂O₇. These compounds contain V in valences from V³⁺ to V⁵⁺ including mixed valence and are a significant test of the EXAFS technique. The known bond distances were obtained from the paper by Wong, et al. (1) and for VPO from Thompson and Ebner (2). Each data set was Fourier filtered to select the region of the first coordination sphere and then fitted in K space using different trial subshells, with appropriate coordination numbers and bond distances, sufficient to account for the known crystallography. During the fits the coordination numbers were fixed while the bond distances R and relative disorder Δσ² were allowed to vary freely. The fits were of good quality as shown by the small standard deviation (SD) listed in Table 1. The value of Δσ² was small and generally varied randomly about zero. When it was large as in the short bond in V₄O₇ it may indicate additional disorder in the short bond, perhaps due to the known mixed valence. The difference between the measured and known distance, ΔR, is listed in the far right column and was used to estimate the average (weighted) error in R of 0.02 ± 0.05 (SD) Å. Therefore, this reference set indicates that the bond distances in these complicated compounds can be determined to ±0.05 Å. Note the structural correspondence between the crystal/ne VPO and active catalysts prepared from aqueous and organic precursors (3).

1. J. Wong, F. W. Lytle, R. P. Messmer and D. H. Maylotte, *Phys. Rev.* B30, 5596 (1984).

2. M. R. Thompson and J. R. Ebner, manuscript in preparation.

3. J. R. Thompson and M. R. Thompson, in *Structure-Activity Relationships in Heterogeneous Catalysts*, Eds. R. K. Grasselli and A. W. Sleight, Elsevier (1991), p. 31.

Table 1. Determination of bond distances in complex V compounds by fitting EXAFS data.

Reference	SD of Fit	Δσ ²	Coord. No.	R, meas. (Å)	R, known (Å)	ΔR (Å)
V ₂ O ₃	0.625	0.0009	3	1.983	1.960	0.023
V ₂ O ₄	0.542	0.0053	3	2.053	2.060	- 0.007
		- 0.0007	1	1.639	1.760	- 0.121
V ₄ O ₇	0.485	0.0050	2	1.846	1.865	- 0.019
		0.0054	3	1.952	2.030	- 0.078
		0.0106	2	1.782	1.890	- 0.108
		- 0.0013	3	1.951	2.004	- 0.053
V ₂ O ₅	0.951	- 0.0002	1	2.110	2.101	0.009
		- 0.0054	1	1.590	1.585	0.005
		- 0.0020	1	1.791	1.780	0.011
		0.0010	2	1.849	1.878	- 0.029
VPO	0.657	- 0.0002	1	2.016	2.021	- 0.005
green		- 0.0027	1	1.613	1.604	0.009
crystals		0.0019	4	1.970	1.951	0.019
		0.0020	1	2.167	2.070	0.097
				ΔR (wt. avg.) =		- .02 ± .05
VPO	0.861	- 0.0021	1	1.627		
organic		0.0046	4	1.997		
		- 0.0005	1	2.302		
VPO	0.726	- 0.0031	1	1.598		
aqueous		- 0.0042	4	1.981		
		- 0.0004	1	2.263		

The work of M. Thompson was funded in part by the USDOE Office of Conservation and Renewable Energy, Advanced Industrial Concepts Division.

EXAFS OF Fe^{3+} IN NITRILE HYDRATASE

M. Nelson and H. Jin (Du Pont), and R. Scharrow, G. Grove, and C. Buck (Haverford C.)

We have performed x-ray fluorescence spectroscopic studies of the iron site in nitrile hydratase from *Brevibacterium* sp. R312. This enzyme catalyzes the hydration of nitriles to amides and is unusual for two reasons: first it has a green color imparted by a 710 nm absorption band that we separately have shown represents a transition with cysteine thiolate to iron charge transfer character; and second, the ferric ion is in the low-spin electronic configuration, rare for Fe^{3+} coordinated by protein-derived ligands.

Analysis of the pre-edge region of the iron k-edge x-ray fluorescence spectra showed a shoulder at 7112 eV corresponding to the $1s \rightarrow 3d$ transition. The small integrated intensity of this absorption suggests that the metal ion exists in a nearly centrosymmetric environment, likely a distorted octahedral ligand field. This is consistent with analysis of the epr spectrum of the protein.

The data in the EXAFS region of the spectrum were fit using a model that included contributions from a total of six coordinated ligands. No reasonable fits to the data could be obtained without including contributions from sulfur, corroborating the coordination of cysteine to the iron. The model that best fit the data included three low Z ligands (N or O) at a distance of 1.98 Å and three S ligands at a distance of 2.21 Å. The accuracy in the number of ligands is estimated to be ± 1 , and in the distances ± 0.03 Å. These distances are reasonable in terms of known low-spin ferric thiolate complexes.

In combination with other data, this suggests a structure for the metal in this enzyme in which the ferric ion is coordinated by three cysteine thiolates, two histidine imidazoles, and a water. This water might be an exchangeable ligand, representing a site at which enzyme substrates or products might coordinate the iron during catalysis. Consequently, we have obtained x-ray fluorescence spectra of samples of enzyme prepared in the presence of isobutyronitrile (a slow substrate) or acetamide (a product). Although each of these samples showed subtle changes in their epr spectra, our preliminary analysis reveals no significant changes in their x-ray fluorescence spectra as compared to the native enzyme. Unfortunately few good models for the low-spin ferric case exist to enable us to predict what to expect to see upon substitution of a nitrile or amide for a water at the metal.

Research sponsored by a grant from the Howard Hughes Medical Institute through the Undergraduate Biological Sciences Education Initiative (Haverford C.).

X-ray Absorption Studies of Organo-disulfide Redox Polymer Electrodes.

X. Q. Yang, K. H. Xue, H. S. Lee, and J. McBreen(DAS/BNL), T. A. Skotheim, (Moltech Co.), F. Lu, (CFRLS, University of Kentucky)

Three systems of organo-disulfide compounds were studied by NEXAFS: (1) 2,5-Dimercapto-1,3,4-thiadiazole, its lithium and potassium salts and polymer; (2) Trithiocyanuric acid, its lithium and potassium salts and polymer; (3) poly(vinylmercaptan), and its lithium and potassium salts.

The NEXAFS spectra of poly(trithiocyanuric acid) and related compounds are plotted in figure 1. The spectrum of poly(trithiocyanuric acid), lithium trithiocyanurate, and potassium trithiocyanurate are marked as curve a, b, and c respectively. The spectrum of poly(trithiocyanuric acid) in discharged and recharged states are marked as curve d and e. In comparison with the spectrum of poly(trithiocyanuric acid) in curve a, the spectra for the alkali-metal salts (curve b and c) showed of new features labeled as 1, 4, and 6. Feature 3, the characteristic peak of poly(trithiocyanuric acid) spectrum is disappeared. All of these new features are more pronounced in curve c when lithium has been replaced by potassium. We refer these new features as characteristic peaks of sulfur-alkali metal bonding. Peak 1 is the π transition related to S=C bond as a result of the $\text{S}^{+}\text{Li}^{+}$ bonding. All of the new features related to $\text{S}^{+}\text{Li}^{+}$ bonding are apparent in curve d after the poly(trithiocyanuric acid) cathode has been discharged. At the same time, the intensity of peak 2 is decreased and the peak width is narrowed. This gives us strong evidence that the electrochemical reduction is indeed taking place through the depolymerization in which the S-S bonds were broken and replaced by $\text{S}^{+}\text{Li}^{+}$ bonds. The electrochemical reversibility of this reaction could be seen in curve e, which was taken after recharging the battery. In curve e, all the features related to $\text{S}^{+}\text{Li}^{+}$ bonding are weakened while the feature 2 related to S-S bonding grows stronger and is broadened compared to curve d. This is a clear indication of the forming of S-S bonds by electrochemical polymerization. The other two systems showed similar effects and will be discussed in a separate paper. Our NEXAFS study provides a clear spectroscopic evidence that the charge-discharge process in lithium-disulfide polymer batteries is realized by the electrochemical polymerization and depolymerization through the S-S bond formation and breaking.

NEXAFS spectra of

- (a) poly(trithiocyanuric acid)
- (b) lithium trithiocyanurate
- (c) potassium trithiocyanurate
- (d) poly(trithiocyanuric acid) discharged
- (e) poly(trithiocyanuric acid) recharged

Work supported by U. S. DOE under Contract No. DE-AC02-76CH00016.

X-19C
X-RAY TOPOGRAPHIC STUDIES OF THE PHASE TRANSFORMATION OCCURRING AT -143°C IN *p*-TERPHENYL SINGLE CRYSTALS.[†]

R. DiSalvo, M. Dudley, J. Wu (SUNY Stony Brook), and D. Gordon-Smith (University of Warwick).

A specially designed cold-finger cryostat was constructed to perform *in situ* synchrotron white beam topography studies on the low temperature phase transition occurring in *p*-terphenyl crystals.¹ Figure 1 (a) shows the room temperature Laue pattern of the monoclinic crystal. As the crystal was progressively cooled through the transition temperature, obvious changes in the transmission Laue pattern became evident. Figure 1(b) shows a Laue pattern recorded at -167°C, showing what appear to be "double images". The average positions of the pairs of reflections can be related to the pseudo-monoclinic structure proposed by Baudour et al.² The image doubling corresponds to the coexistence of two possible low temperature structures, related by a crystallographic glide plane.² The presence of these two structures is indicated in figure 1(b) by the subscripts, α and β . Many of the diffraction images appear distorted as a result of strain induced during the phase transformation. Examination of the crystal upon reheating showed that the original Laue pattern is restored with no apparent differences in the microstructure.

Figure 1.

References: 1. M. Dudley, R. DiSalvo, J. Wu, D. Gordon-Smith, and W. Jones, to appear in *Mol. Cryst. Liq. Cryst.*, (1991); 2. J.L. Baudour, H. Calleau, and W.B. Yelon, *Acta Cryst.*, B33, 1773-1780, (1977).

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

X-19C
CHARACTERIZATION OF CHEMOMECHANICALLY POLISHED CdTe and CdZnTe SINGLE CRYSTAL WAFERS BY SYNCHROTRON X-RAY WHITE BEAM TOPOGRAPHY.[†]

Thomas Fanning, Michael Dudley (SUNY Stony Brook), Don Di Marzio and Louis Casagrande (Grunman Corporate Research Center).

CdTe and CdZnTe single crystal wafers are important substrate materials for HgCdTe based infrared detectors. A series of chemomechanically polished CdTe and CdZnTe single crystal wafers from a number of manufacturers were compared to Grunman grown CdTe. All the wafers were oriented in the [111] direction and the (333) was taken as the main reflection with the x-ray film oriented as to eliminate any aspect distortion in the topographic image ($\theta_B = 20^\circ$, $\lambda = 0.85\text{\AA}$). Although optically similar in the visible and the IR (except for precipitate densities), the x-ray topographic images varied widely between the various manufacturers. All show precipitate strain fields of varying sizes and densities. Figure 1(a) is a manufacturer's polished CdZnTe 1x1 cm wafer, showing a pronounced precipitate structure. It also exhibits a low angle grain boundary not visible optically (IR). X-ray rocking curve maps confirm the strained nature of this region. This is compared to an image of a vertical Bridgman grown Grunman CdTe wafer in figure 1(b). Note the twin lamella, indicated by T. The wafer exhibits a well-defined cellular structure of dislocations. Typical x-ray rocking curve widths in these regions range from 10 to 15 arcseconds FWHM.

Figure 1.

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded by Grunman Corporate Research Center.

MODIFICATION OF PRECIPITATE STRAIN FIELDS ACCOMPANYING RTP OF HIGH CARBON CONTENT SILICON.[†]

T. Fanning, M. Dudley, F.F.Y. Wang (SUNY Stony Brook), D. Gordon-Smith (University of Warwick) and D.T. Hodul (Varian Research Center).

Synchrotron white beam topography has been used to study the effect of Rapid Thermal Processing (RTP) on the defect microstructure of high carbon content silicon. Among the effects observed is the apparent partial relaxation of the elastic strain fields associated with precipitates which existed prior to RTP.^{1,2} This is demonstrated by comparison of figures 1(a) and (b), recorded respectively before and after RTP ($g=333$, $\lambda=0.68\text{\AA}$). This relaxation is likely to be due to the formation of dislocations in the vicinity of the precipitates although in this particular case, dislocation source operation appears to occur on a scale below the resolution limit of the technique. However, in others, loop formation around precipitates is observable.³

Figure 1.

References: 1. M. Dudley, F.F.Y. Wang, T. Fanning and D. Gordon-Smith, in "Materials Reliability Issues in Microelectronics", M.L. Green, J.C. Gelpey, J. Wortman, and R. Singh (Eds.), Mat. Res. Soc. Symp. Proc., 224, 61-66, (1991); 2. T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith and D.T. Hodul, in "Rapid Thermal and Integrated Processing", J.R. Lloyd, T.S. Ho, C.T. Sah, and E. Yost (Eds.), Mat. Res. Soc. Symp. Proc., 225, 301-306, (1991); 3. T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith and D.T. Hodul "Observation of RTP Induced Dislocation Nucleation at Precipitates in High Carbon Content Silicon", in this volume.

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

OBSERVATION OF COMBINED CLIMB AND GLIDE INDUCED BY RTP OF HIGH CARBON CONTENT SILICON.[†]

T. Fanning, M. Dudley, F.F.Y. Wang (SUNY Stony Brook), D. Gordon-Smith (University of Warwick) and D.T. Hodul (Varian Research Center).

Synchrotron white beam topography has been used to characterize the nature of dislocation sources created as a result of Rapid Thermal Processing (RTP) of silicon samples with high carbon concentrations.^{1,2} The specimen studied here was of (111) orientation and was 100mm in diameter. When examined prior to RTP it was found to be essentially defect free. Examination of the same sample following RTP revealed the operation of multiple dislocation sources. Examples of a commonly observed dislocation configuration, marked C, are shown in figure 1 ($g=333$, $\lambda=0.68\text{\AA}$). Figure 2(b) is an enlargement of the configuration on the extreme left of figure 1. Figure 2(b) shows a different topograph recorded of this same configuration ($g=022$, $\lambda=1.42\text{\AA}$). In this latter image, the diffraction geometry is such that the operative slip planes (in this case $\{111\}$) are viewed "edge on". This image reveals that slip has occurred on two parallel slip planes separated by a distance of some $12\mu\text{m}$ over which climb has occurred.

Figure 1.

Figure 2.

References: 1. M. Dudley, F.F.Y. Wang, T. Fanning and D. Gordon-Smith, in "Materials Reliability Issues in Microelectronics", M.L. Green, J.C. Gelpey, J. Wortman, and R. Singh (Eds.), Mat. Res. Soc. Symp. Proc., 224, 61-66, (1991); 2. T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith and D.T. Hodul, in "Rapid Thermal and Integrated Processing", J.R. Lloyd, T.S. Ho, C.T. Sah, and E. Yost (Eds.), Mat. Res. Soc. Symp. Proc., 225, 301-306, (1991).

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

X-19C OBSERVATION OF RTP INDUCED DISLOCATION NUCLEATION AT PRECIPITATES IN HIGH CARBON CONTENT SILICON.[†]

T. Fanning, M. Dudley, F.F.Y. Wang (SUNY Stony Brook), D. Gordon-Smith (University of Warwick) and D.T. Hodul (Varian Research Center).

Synchrotron white beam topography of high carbon content silicon wafers following Rapid Thermal Processing (RTP) has revealed the relaxation of the elastic strain fields associated with precipitates which existed prior to RTP.^{1,2,3} This relaxation is likely to be due to the formation of dislocations in the vicinity of the precipitates. In some cases dislocation source operation appears to occur on a scale below the resolution limit of the technique. However, in others, loop formation around precipitates is observable. An example of this is shown in figures 1(a) and (b) recorded respectively before and after RTP. Note the small dislocation loops emanating from the precipitate.

Figure 1.

References: 1. T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith and D.T. Hodul "Modification of Precipitate Strain Fields Accompanying RTP of High Carbon Content Silicon", in this volume; 2. M. Dudley, F.F.Y. Wang, T. Fanning and D. Gordon-Smith, in "Materials Reliability Issues in Microelectronics", M.L. Green, J.C. Gelpey, J. Wortman, and R. Singh (Eds.), *Mat. Res. Soc. Symp. Proc.*, 224, 61-66, (1991); 3. T. Fanning, M. Dudley, F.F.Y. Wang, D. Gordon-Smith and D.T. Hodul, in "Rapid Thermal and Integrated Processing", J.R. Lloyd, T.S. Ho, C.T. Sah, and E. Yost (Eds.), *Mat. Res. Soc. Symp. Proc.*, 225, 301-306, (1991).

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

X-19C STUDIES OF RTP INDUCED DAMAGE IN HIGH CARBON DOPED SILICON.[†]

T. Fanning, M. Dudley, F.F.Y. Wang (SUNY Stony Brook), D. Gordon-Smith (University of Warwick) and D.T. Hodul (Varian Research Center).

Synchrotron white beam topography has been used to investigate the damage induced in silicon single crystals by Rapid Thermal Processing (RTP). Samples with a range of carbon concentrations were examined, although results presented here pertain to a specimen with a carbon concentration of 3.15ppma. This specimen, which was of (111) orientation and 100mm in diameter, was examined prior to RTP and found to be essentially defect free. Examination of the same sample following RTP revealed propagation of multiple dislocation sources, nucleated at the crystal edge, as well as formation of precipitates. A topograph illustrating this behavior is shown in figure 1 ($g=333$, $\lambda=0.68\text{\AA}$). This behavior can be contrasted to that of nominally carbon-free silicon which showed only RTP induced glide of existing dislocations, with no nucleation of new sources.¹

Figure 1.

References: 1. M. Dudley, F.F.Y. Wang, T. Fanning, G. Tolis and J. Wu, *Mater. Letts.*, 10, 87, (1990).

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

X-19C CHARACTERIZATION OF DEFECT STRUCTURES CLOSE TO GRAIN BOUNDARIES IN ICE CRYSTALS.†

F. Liu and I. Baker (Dartmouth College), and G.-D. Yao and M. Dudley (SUNY Stony Brook).

X-ray topography has been used extensively to study single crystal ice¹. Naturally, ice exists not as single crystals but as polycrystals. Therefore, understanding of defect structures and their behavior in ice polycrystals is of considerable interest to the ice mechanics community. In this report we present results of a preliminary study of dislocations and grain boundaries in high-purity polycrystalline ice containing columnar grains using synchrotron white beam topography.

Screw, 30° and 60° basal dislocations with Burgers vector $<11\bar{2}0>$ were found to far outnumber other dislocations. Figure 1 ($g=2\bar{1}\bar{1}0$, $\lambda=0.69\text{\AA}$) shows an image recorded from a single grain from a polycrystalline specimen. Thickness fringes can be seen associated with the boundary. Note the dislocations which propagate up to the boundary. The detailed interactions between dislocations and grain boundaries are currently under investigation.

Figure 1.

References: 1. A. Higashi, "Lattice Defects in Ice Crystals, X-ray Topographic Observations", Hokkaido University Press, Sapporo, Japan, (1988).

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by NASA under contract #NAS8-38147.

X-19C OBSERVATION OF FAULTS PRODUCED BY VACANCY CONDENSATION IN ICE SINGLE CRYSTALS.†

F. Liu and I. Baker (Dartmouth College), and G.-D. Yao and M. Dudley (SUNY Stony Brook).

Synchrotron white beam topography has been used to investigate the formation of stacking faults via vacancy condensation in large-grain ice polycrystals. Figure 1 shows an example of a topograph recorded following quenching. An interesting feature is the absence of stacking faults in the vicinity of the boundary B, despite the high density of faults within the grain. The formation of this depleted zone, which is of the order of 0.5mm wide, can be understood by considering the grain boundaries as sinks of water molecule interstitials and vacancies. The size of the depleted zone is found to be a function of the angle between the basal plane and the grain boundary. For example, a 1mm thick grain with the grain boundary parallel to the large surface, and with the basal plane nearly perpendicular to the boundary did not show any evidence for stacking faults, even following rapid quenching. This implies that diffusion in ice occurs mainly on the basal plane, and that an ice sample as thin as a TEM sample can not represent the microstructure of bulk ice.

Figure 1.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Work supported in part by the Army Research Office through grant No. DAA-G29-85-K-0251, and by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

X-19C

SYNCHROTRON TOPOGRAPHIC INVESTIGATION OF DAMAGE DISTRIBUTIONS
RESULTING FROM ELECTROMAGNETIC BREAKDOWN OF EPTAXIAL SILICON
p-n JUNCTIONS.†

G. Tolis, M. Dudley (SUNY Stony Brook), C. Fazi (Harry Diamond Laboratory) and D. Gordon-Smith (University of Warwick).

Grazing Bragg-Laue white beam topographic imaging has been used to determine the distribution of damage as a function of depth, resulting from the electromagnetic breakdown of p-n junctions on silicon. Damage resulting from breakdown in both forward and reverse bias was studied, and clear differences were revealed. Damage in the reverse bias case was much more extensive in the regions close to the specimen surface than in the forward bias case. This is in qualitative agreement with predictions of damage distributions based on electrothermal models. Figures 1-3 show details from three grazing Bragg-Laue topographs, recorded from a p-n junction which failed in reverse bias. Each image has a different kinematical penetration depth (t_k^2): Figure 1, $g=313$, $\lambda=0.85\text{\AA}$, $t_k^2=3.9\mu\text{m}$; Figure 2, $g=313$, $\lambda=0.9\text{\AA}$, $t_k^2=4.89\mu\text{m}$; Figure 3, $g=313$, $\lambda=0.99\text{\AA}$, $t_k^2=10.48\mu\text{m}$. Note the contrast changes which become apparent as the penetration depth increases.

Figure 1.

Figure 2

Figure 3.

X-19C

ASSESSMENT OF DAMAGE INDUCED BY ELECTROMAGNETIC BREAKDOWN IN
SILICON BI-POLAR DIODES BY SYNCHROTRON WHITE BEAM X-RAY TOPOGRA-
PHY†

S. Wang, M. Dudley (SUNY Stony Brook), D. Gordon-Smith (University of Warwick) and C. Fazi (Harry Diamond Laboratory).

Synchrotron white beam x-ray topographic (SWBXT) imaging has proven to be a powerful technique for the assessment of structural damage resulting from electromagnetic breakdown in Si bi-polar diodes. Unlike optical microscopy, previously applied to such damage assessment, SWBXT imaging is a highly-strain sensitive technique which can distinguish differences between structural damage in the single crystal epilayer region due to the electromagnetic breakdown, and other surface irregularities occurring on the metallization and silicon oxide layers. Assessment of damage using optical microscopy can be misleading, as demonstrated below. Figure 1(a) is an optical micrograph of a typical Si diode after electromagnetic breakdown in reverse bias. There are two spot-like features visible on the metallization, one located on the circumference and the other in the interior. However, in the corresponding x-ray diffraction topograph (figure 1(b), $g=242$, $\lambda=0.78\text{\AA}$), only the feature on the circumference gives significant contrast, and is in fact due to a region of lattice distortion resulting from the breakdown. On the other hand, the center spot is virtually invisible, indicating that it is just a surface feature on the metallization.

(a)

(b)

0.1mm

Figure 1.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded by U.S. Army Harry Diamond Laboratories.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded by U.S. Army Harry Diamond Laboratory.

X-19C

GRAZING BRAGG-LAUE SYNCHROTRON X-RAY TOPOGRAPHIC IMAGING OF
NEAR-SURFACE STRUCTURAL DAMAGE IN SILICON BI-POLAR DIODES AFTER
FAILURE WITH MICROWAVE PULSES†

S. Wang, M. Dudley (SUNY Stony Brook), D. Gordon-Smith (University of Warwick) and C. Fazi (Harry Diamond Laboratory).

Structural damage in Si bi-polar diodes resulting from electromagnetic breakdown is confined to a very shallow region close to the surface of the junction.^{1,2} The grazing Bragg-Laue white beam topographic imaging technique has been employed to determine the character of the damage for the case of damage caused by microwave pulses. Since the x-ray penetration depths can be controlled and varied by simple rotations of the Si wafer in the grazing Bragg-Laue technique, structural information from shallow regions beneath the electrode can be obtained from the x-ray diffraction topographs. Figures 1 (a)-(c) show x-ray diffraction topographs ($g=242$) recorded with three different penetration depths: Figure 1(a) ($\lambda=0.76\text{\AA}$), $4.4\mu\text{m}$; Figure 1(b) ($\lambda=0.78\text{\AA}$), $8.9\mu\text{m}$; Figure 1(c) ($\lambda=0.82\text{\AA}$), $18\mu\text{m}$, showing the distribution of damage in a triangular shaped diode after electromagnetic breakdown with microwave pulses of $10\mu\text{s}$ duration. Apparently, as the penetration depth increases from 4.4 to $18\mu\text{m}$, the "pencil-point" contrast, due to the structural damage appears increasingly diffuse. This behavior is attributable to the existence of a "filamentation" region resulting from a reverse bias failure, rather than a forward bias type of failure.^{1,2}

Figure 1.

References: 1. G. Tolis, Masters Thesis, SUNY Stony Brook, (1990); 2. M. Dudley, G. Tolis, D. Gordon-Smith, and C. Fazi, "Government Microcircuit Applications Conference Digest of Papers", Vol. XVI, 651, (1990).

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded by U.S. Army Harry Diamond Laboratory

X-19C

CHARACTERIZATION OF DISLOCATION STRUCTURES AROUND INDENTS ON
{111} INDIUM ANTIMONIDE SURFACES.†

J. Wu and M. Dudley (SUNY Stony Brook)

Synchrotron white beam x-ray topography in the transmission Laue geometry has been used to characterize defect structures around indents on single crystals of InSb. Dislocation mobility in InSb is large enough that individual dislocations can easily be resolved using x-ray topography. Microindentation was produced on the {111} surfaces by a Vickers profile indenter with various loads from 10g, 25g, 50g, 100g, 200g, 300g, 500g, to 1Kg at a temperature of 390°C , this being above the brittle-to-plastic transition temperature. Figures 1(a) and (b) show white radiation synchrotron topographs ($g=422$ and 400 , respectively) recorded following indentation. The contrast from the severely deformed area caused by the highest load indent, and the dislocations emitted from the lower load indentation sites are clearly shown. The dislocation configurations in this III-V semiconductor material are very strongly influenced by the existence of large differences between the mobilities of screw and 60° segments of dislocation loops. Further research is being conducted to investigate these differences.

Figure 1.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund which is administered by the American Chemical Society.

X-19C
CHARACTERIZATION OF GEL-GROWN CRYSTALS BY SYNCHROTRON WHITE
BEAM TOPOGRAPHY. †

J. Wu, S.-P. Wang, and M. Dudley (SUNY Stony Brook).

The gel growth technique has several advantages over other single crystal growth techniques. It can effectively cut down the nucleation density and the crystal growth rate. It can also minimize stress on the crystal during the growth process. In order to evaluate this technique, calcium tartrate single crystals ($\text{CaC}_2\text{O}_6 \cdot 4\text{H}_2\text{O}$) were grown in silica gels made of Na_2SiO_3 solutions. Synchrotron white radiation topography was used to characterize the quality of the crystals grown. Figure 1 shows a transmission topograph of a gel-grown calcium tartrate crystal. Growth veils, growth sector boundaries and grown-in dislocations are observed. It was also observed that the crystal is sensitive to the x-ray beam, especially at high currents. The crystal damaged after irradiation at ≈ 200 mA for around 1 minute. Some micro-cracks were subsequently developed.

Figure 1.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund, which is administered by the American Chemical Society.

X-19C
ANALYSIS OF DEFECT STRUCTURES IN NdGaO_3 SINGLE CRYSTALS.†

G.-D. Yao, S.-Y. Hou and M. Dudley (SUNY Stony Brook), and J.M. Phillips (A.T. & T. Bell Labs.).

Synchrotron White Beam Topography in conventional Bragg geometry has been used to characterize the defect content of NdGaO_3 single crystals. These crystals are potential candidates for use as substrates for High T_c superconductor epilayers*. As can be seen below in Figure 1, the crystal studied had a low dislocation density, and showed no evidence for twinning. The dislocation marked D can be seen to be made up of well defined straight segments, each lying along a low energy direction on the operative slip plane. Correlation between the defect structure in these substrate crystals and the superconducting behavior of high T_c films grown on them is currently being investigated.

Figure 1.

References: 1. H.M. O'Bryan, P.K. Gallagher, G.W. Berkstresser, and C.D. Brandle, J. Mater. Res., 5, 183, (1990).

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund, which is administered by the American Chemical Society. Crystals supplied by C.D. Brandle at A.T. & T. Bell Labs.

ANALYSIS OF TWIN STRUCTURES IN LaAlO_3 SINGLE CRYSTALS.[†]

G.-D. Yao, S.-Y. Hou and M. Dudley (SUNY Stony Brook), and J.M. Phillips (A.T. & T. Bell Labs.).

Synchrotron White Beam Topography in conventional Bragg geometry has been used to characterize the defect content of LaAlO_3 single crystals. These crystals are potential candidates for use as substrates for High T_c superconductor epilayers.[†] As can be seen below in Figure 1 ($g=220$, $\lambda=0.99\text{\AA}$), the crystal has a detailed structure of interpenetrating twins. Two general kinds of twinning are observed, marked T_1 and T_2 . Complete analysis of the twinning operations has been performed. The influence of the twin structures on the superconducting behavior of the high T_c films is currently under investigation.

Figure 1.

References: 1. H.M. O'Bryan, P.K. Gallagher, G.W. Berkstresser, and C.D. Brandle, *J. Mater. Res.*, 5, 183, (1990).

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund, which is administered by the American Chemical Society. Crystals supplied by C.D. Brandle at A.T. & T. Bell Labs.

X-19C
COMPARISON BETWEEN OBSERVATION AND CALCULATION OF SURFACE RELAXATION EFFECTS ON DISLOCATION STRAIN FIELDS IN THIN FILMS.[†]

G.-D. Yao, J. Wu, and M. Dudley (SUNY Stony Brook), and V. Shastri and P. Anderson (Ohio State University).

The influence of surface relaxation on the imaging of dislocations in thin single crystal films, using white beam synchrotron radiation topography in grazing Bragg-Laue geometry, has been assessed.[†] Agreement between predicted and observed visibility of interfacial dislocation images could only be obtained by incorporating surface relaxation effects. Calculated equi-effective-misorientation contours for various diffraction geometries are presented in figures 1 (without surface relaxation) and 2 (with surface relaxation). t indicates kinematical penetration depth. The influence of surface relaxation effects is evident. Figure 3 shows grazing Bragg-Laue topographs recorded with diffraction geometries identical to those used in calculation of figures 1 and 2. The specimen consists of a $3\mu\text{m}$ thick epilayer of GaAs grown by MBE on a $600\mu\text{m}$ thick Si substrate.

Figure 1.

Figure 2.

Figure 3.

References: 1. G.-D. Yao, J. Wu, M. Dudley, V. Shastri and P. Anderson, to appear in *Mat. Res. Symp. Proc.*, 209, (1991).

[†] Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by Texas Instruments.

IMAGING OF DEFECT STRUCTURES IN CdZnTe SINGLE CRYSTALS.†

G.-D. Yao, M. Dudley, T. Fanning (SUNY Stony Brook) and D. Larson (Grueman Corporate Research Center).

Synchrotron white beam topography in conventional Bragg geometry has been used to characterize defect structures in CdZnTe single crystals. 180° rotation twins are prevalent in this material. In the white beam technique these become visible via orientation contrast. Most twinned regions are macroscopic in size, although some microscopic twin lamellae are observed, as can be seen in figure 1. Also visible in this image are a grain boundary and dislocation sub-structures. Correlations between observed defect configurations and growth parameters are currently under investigation.

Figure 1.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by NASA under contract #NAS8-38147.

X-19C SYNCHROTRON TOPOGRAPHY OBSERVATIONS OF THE SECOND ORDER PHASE TRANSITION IN LANTHANUM ALUMINATE SINGLE CRYSTALS.†

G.-D. Yao, S.-Y. Hou, M. Dudley, (SUNY Stony Brook), and J.M. Phillips (A.T. & T. Bell Labs.).

Synchrotron white beam topography has been used to monitor the second order, rhombohedral to cubic phase transition which occurs on heating LaAlO_3 through $\approx 435^\circ\text{C}$. At temperatures approaching that of the transition, the angular shift associated with the presence of twins in this material at room temperature is seen to gradually diminish (see figure 1(a) and (b), recorded respectively at temperatures of 20°C and 200°C ($g=220$)). Eventually this shift goes to zero as the operative twin plane in the rhombohedral phase becomes a mirror plane in the high temperature cubic phase. This is demonstrated by the absence of twins in figure 1(c), recorded at 435°C . On cooling, twins are seen to nucleate in the center of the crystal and grow towards the edge. The final twin configuration, shown in figure 1(d), recorded at room temperature, is seen to be less dense than the starting one. This reduction in twin density may have beneficial effects on the use of this material as a substrate for high T_c superconductor thin films.

Figure 1.

† Acknowledgements: Work performed on the Synchrotron Topography Project Beamline X-19C, which is supported in part by DOE under Grant No. DE-FG02-84ER45098. Research funded in part by the donors of the Petroleum Research Fund which is administered by the American Chemical Society. The authors are grateful to C.D. Brandle of A.T. & T. Bell Labs. for providing the crystals.

B.J. Factor (Applied Physics Dept., Stanford U.), T.P. Russell and M.F. Toney (IBM Research).

Figure 1

Figure 2

Recently, grazing incidence x-ray diffraction has been used to discern enhanced ordering at the surface of an aromatic polyimide¹. A 2600 Å film which had been cured at 420°C was examined by measuring the diffraction at several different values of q_0 , the component of the scattering vector perpendicular to the surface, which corresponded to x-ray penetration depths ranging from 50 Å to the film thickness. In this work, a 400 Å film cured at 300°C was examined using the same technique. As can be seen in Figure 1, the diffraction pattern changes dramatically as the x-rays become surface sensitive. For bulk penetration ($q_0 > 0.024 \text{ Å}^{-1}$), this diffraction is characteristic of the liquid crystalline ordering found in bulk films². When the x-rays are surface sensitive ($q_0 < 0.024 \text{ Å}^{-1}$), the diffraction is characteristic of the crystalline ordering observed at the surface of the 2600 Å film cured at 420°C.

The first peaks for the liquid-crystalline and crystalline states are centered at 0.434 Å^{-1} and 0.397 Å^{-1} , respectively, with a full width at half maximum which decreases from 0.009 Å^{-1} to 0.06 Å^{-1} . The lineshape profiles were taken to be a superposition of these two contributions, and the integrated intensity of each contribution was calculated separately³. Assuming the scattered intensity was proportional to the overlap of the concentration of diffracting material with the x-ray field (as determined by linear optics⁴), the integrated intensities of the separate contributions from the liquid-crystalline and crystalline states were determined.

In Figure 2, the integrated intensity of reflections corresponding to the liquid-crystalline and crystalline states are modeled with two different concentration profiles. The solid line was fitted to a model with crystalline ordering in a 40 Å layer near the air surface, and a liquid-crystalline bulk. The dotted line corresponds to a model where a 40 Å crystalline layer exists at both the air and substrate surfaces. The two curves for the 0.434 Å^{-1} data describe the data nearly equally well. Therefore, it is clear that the air surface affects the polymer in a way which promotes enhanced ordering nearby.

However, for the 0.397 Å^{-1} data, the model for a 40 Å layer at the substrate surface clearly overestimates the data, while that for no ordering at the substrate is somewhat low. The best fit was for a 10 Å layer near the substrate surface (not shown). Nonetheless, comparison of the models shows that the influence of the substrate and air interfaces is clearly different. This could be due to polymer adsorption at the silicon substrate or, more simply, that the polymers cannot order as easily at a hard wall surface as at a free surface.

This research is supported by the Department of Energy through the Office of Basic Energy Sciences (DOE-BES), and IBM. B.J.F. acknowledges support from the DOE-BES, Division of Materials Science through SSRL.

1. B.J. Factor, T.P. Russell, M.F. Toney, *Phys. Rev. Lett.*, **66**, 1181 (1991).
2. Tabahashi, J.; Yoon, D.Y.; Parrish, W. *Macromolecules*, **17**, 2583 (1984).
3. B.J. Factor, *Ph.D. Thesis*, Stanford University, 1991.
4. Max Born and Emil Wolf, *Principles of Optics*, Pergamon, Oxford, 1958.

J.P. Hill, P. Westbrook, M. Greven, R.J. Birgeneau (MIT) and T.R. Thurston (BNL).

As part of an ongoing research program into the 3d Random Field Ising Model (RFIM), we report experiments carried out on two crystals of $\text{Mn}_{0.75}\text{Zn}_{0.25}\text{F}_2$.

In a previous magnetic x-ray scattering experiment¹ on the first of these two crystals, sample #1, we observed a long range ordered (LRO) ground state, in apparent contradiction with all previous neutron scattering work. This led to the hypothesis that there was a surface relaxation into the true ground state, while the bulk remained frozen in a domain state.

In an effort to gain further insight into the mechanism for surface relaxation, we studied two crystals cut from the same boule. Both crystals were of excellent sample homogeneity and crystallographic quality (mosaic $\sim 0.003^\circ$ HWHM). For the two samples $T_N(0)$ was 46.1K and 46.0K respectively. The transition was sharp, any rounding being significantly less than 50mK , ruling out any surface concentration gradient effects.

Representative field cooled transverse scans of the (100) magnetic peak are shown in the figure. The dashed line in each case is the same resolution function. The two crystals exhibited dramatically different magnetic ordering on cooling in a field. Sample #2 did not exhibit LRO, but formed the domain state that neutrons had previously observed in the bulk. The solid line shown is a fit to a Lorentzian squared. (It is worth noting that this is the first time that broadened peaks have been observed with magnetic x-ray scattering).

Sample #1 again showed a resolution limited Bragg peak indicative of LRO, but also present was a much smaller short range ordered component. The solid line in the figure is a delta function plus a Lorentzian squared with the correlation length fixed at the value obtained from #2. Preliminary analysis suggests that the LRO appears at a second order transition about 1K below the metastability boundary.

Mechanically the two samples differ only in one important regard, their surface preparation. A visual inspection of the two crystals

reveals more scratches on sample #1 than on #2, which appears perfect. This difference manifests itself in x-ray reflectivity data which showed that sample #1 had a reflectivity half that of #2, suggesting the surface was indeed significantly rougher. Real space images taken with an environmental SEM (ESEM) confirmed this. This technique has a resolution of better than $1\mu\text{m}$. The surface of #1 was markedly more scratched than #2 which appeared almost perfect in these images.

This work suggests that it is possible that surface imperfections nucleate the LRO. Further work is proceeding.

1. J.P. Hill *et al.*, *Phys. Rev. Lett.* **66**, 3281 (1991).

Figure 3

X-RAY SCATTERING STUDIES OF STRAIN AT THE $\text{SiO}_2/\text{Si}(001)$ INTERFACE

S.D. Kosowaly, T.A. Rabedeau, P.S. Pershan (Harvard U.), J. Bevk, B.S. Freer (AT&T Bell Labs)

We are conducting a study of the role of strain in the spatial dependence of structural changes between crystalline $\text{Si}(001)$ and an amorphous silicon oxide overlayer. Current theories assert that strain dependent energetics factor into the structure and stoichiometry of the oxide/silicon interface. To investigate these effects, we induce a variable external strain at the $\text{SiO}_2/\text{Si}(001)$ interface by dilating the Si lattice to match a host lattice, in particular, a $\text{Si}_x\text{Ge}_{1-x}$ (001) substrate ($0 \leq x \leq 1$). In this initial run, two monolayers of Si were epitaxially deposited on a Ge(001) substrate (i.e., $y=0$). The sample was then oxidized in dry O_2 at room temperature.

Preliminary analysis of the grazing incidence diffraction (GID) $\text{SiO}_2/\text{Ge}(001)$ scattering data indicates the presence of interfacial order comprised of coexisting 2×1 and 1×2 epitaxial domains. Analysis of the line shapes for the GID data indicates exponential decay of in-plane order with ~ 200 Å correlation length. The unit cell responsible for the 2×1 symmetry structure is sufficiently well ordered that large in-plane momentum transfer peaks (such as the $(3/2, -9/2)$) are clearly evident (see figure). Specular reflectivity data, however, indicate that the interface is atomically rough. Future analysis will reveal the nature of any interfacial order, in particular, whether a structured SiO_2 layer exists at the oxide/semiconductor interface.

Work supported by the DOD Joint Services Electronics Program under grant N00014-89-J-1023 and by AT&T Bell Laboratories.

Transverse scans through several in-plane surface peaks at $q_z \approx 0.03 \text{ Å}^{-1}$. (a) $(1/2, -1/2)$ (b) $(1/2, -7/2)$ (c) $(3/2, 3/2)$ (d) $(3/2, -9/2)$

Nitrogen Induced (2×3) Reconstruction of $\text{Cu}(110)$

A. R. Sandy, C. You, and S. G. J. Mochrie (MIT); A. Baddorf, D. M. Zehner (ORNL); L. D. Gibbs (BNL).

There is a great deal of controversy in the recent literature concerning the nature of the reconstruction of the nitrogen covered $\text{Cu}(110)$ surface. Recent STM¹ and LEIS² measurements claim a missing-row structure while other LEIS³ measurements suggest an added-row structure. In an attempt to resolve this controversy we have performed x-ray reflectivity measurements of the $\text{N}/\text{Cu}(110)$ system at 300 K. Measurements of both reconstruction and crystal truncation rods have been made on an absolute scale and are displayed in Fig. 1. The data are shown as solid circles; the best fit to the profiles as solid lines. The agreement is quite good and suggests that a buckled, added-row structure is correct.

Fig. 1

¹H. Niehus *et al.*, Phys. Rev. B 43, 12619 (1991).
²R. Spitz *et al.*, Surf. Sci. Lett. 250, L355 (1991).
³M. J. Ashwin *et al.*, Surf. Sci. 237, 108 (1990).

ELECTROCHEMICALLY ADSORBED BISMUTH MONOLAYERS ON SILVER (111): SURFACE CRYSTALLOGRAPHY OF THE UNIAXIAL COMMENSURATE STRUCTURE

M.F. Toney, J.G. Gordon, M.G. Samant, G.L. Borges, D.G. Wiesler, (IBM Research Division, Almaden Research Center), D. Yee and L.B. Sorensen (U. Washington)

We describe in-situ surface X-ray scattering measurements of electrochemically deposited Bi monolayers on Ag(111). Bi forms an unusual structure: a two-dimensional *rectangular* lattice that is uniaxially commensurate with the *hexagonal* Ag(111) surface along the Ag [211] direction.¹ The two-dimensional structure is shown in the figure. There are two Bi adatoms per rectangular unit cell and one adatom is displaced from the centered-rectangular position by 0.35Å along the commensurate direction. This displacement shortens two of the Bi-Bi bond distances but lengthens two others and it reflects the tendency toward covalent bonding in Bi. The monolayer structure is close to that of the close-packed or (102) planes of bulk Bi, and we believe that the formation of a commensurate layer is due to a fortuitously close match between the atomic spacings of the Ag substrate and the (102) Bi planes.

To determine the registry of the commensurate Bi rows, we conducted measurements of the silver truncation rods. These indicate that the commensurate Bi rows lie along the rows of three-fold hollow sites in the substrate (see Figure). In addition, our analysis of the intensities from the Bi peaks shows that the disorder in the monolayer is anisotropic: the in-plane root-mean-square displacement amplitude in the commensurate direction is 0.2Å, while in the incommensurate direction, it is 0.3Å. This anisotropy reflects the tendency of the substrate to lock the adatoms in position along the commensurate direction, while along the incommensurate direction, this tendency is absent.

The Bi-Ag(111) structure that we observe in-situ (under the electrolyte) is the same as that observed in ex-situ emersion experiments.² In these experiments, the electrode surface is examined with surface science methods following its emersion (removal) from solution,³ and a key question to be addressed is whether the structure determined after emersion is the same as the structure in-situ. Our results for Bi-Ag(111) provide an existence proof that the in-situ structure can be preserved in an ex-situ emersion experiment.¹

1. M.F. Toney et al., *Langmuir* 7, 796 (1991).
2. J. Laguerre-Davidson et al., *Langmuir* 4, 224 (1988).
3. A.F. Hubbard, *Chem. Rev.* 88, 633 (1988).

Structure of Bi-Ag(111). The open circles represent the surface atoms of the Ag substrate and the shaded circles represent the Bi adatoms. The relative sizes of these circles correspond to the near neighbor spacing of bulk Ag (2.89Å) and bulk Bi (3.07Å). The Bi monolayer is commensurate in the *a* direction, (*a* = 5.005Å = $\sqrt{3}$ times Ag nearest-neighbor distance), but is incommensurate in the *b* direction.

This work was partially supported by the Office of Naval Research.

STRUCTURAL AND MAGNETIC X-RAY SCATTERING MEASUREMENTS OF EPITAXIAL DY THIN FILMS

M.F. Toney, D.G. Wiesler, B.D. Ilersmiller, and R.F. Farrow (IBM Research Division, Almaden Research Center)

We describe structural and magnetic X-ray scattering measurements of a 2000Å thick Dy film, sandwiched by LaF₃ films on a GaAs(111) substrate. The structure was grown by molecular beam epitaxy with the c-axis of the Dy parallel to the LaF₃ c-axis and GaAs[111] axis.¹ The goal of the experiments was to show the feasibility of magnetic X-ray diffraction measurements on these thin films and to compare their behavior with bulk crystals.

Between 85 and 179K, bulk Dy is antiferromagnetic with an hcp lattice. The magnetic moments lie in the basal plane and form a helix about the c-axis. The magnetic wavevector is $\tau = 2\pi/d$, where *d* is the helix period and is incommensurate with the lattice. Bulk Dy transforms into a ferromagnet at 85K.

For the magnetic X-ray scattering, a polarization analyzer² was used to greatly reduce the background. This was a pyrolytic graphite crystal set to diffract the (006) planes ($2\theta = 90.8^\circ$) and was aligned to diffract perpendicular to the scattering plane. We used an X-ray energy of 7790 eV, since this provided the maximum resonant enhancement³ of the magnetic scattering from the Dy film.

The figure shows typical scans of the (00 2 + τ) magnetic diffraction peak in the antiferromagnetic phase for this 2000Å thick film. With increasing temperature, τ increases and the intensity decreases, although the peak width remains constant. We have quantified the temperature dependence of the peak position and intensity^{4, 5} and find the temperature dependence of τ is similar to bulk, although a weaker dependence is observed below ~110K. The c-axis lattice constant of the Dy film also has a similar temperature dependence to bulk Dy from 300K to about 110K, but below this, the film is strained. This suggests that the different magnetic behavior is caused by magnetoelastic effects.

The transition to ferromagnetic ordering at 85K exhibits temperature hysteresis as is evident in the structural and magnetic X-ray data and in SQUID magnetometry data. We believe this hysteresis arises from a combination of the strain-energy barrier accompanying the transition and magnetic inhomogeneities in the film.^{4, 5}

1. R.F. Farrow, S.S. Parkin, and V.S. Speriosu, *J. Appl. Phys.* 64, 5315 (1988).
2. D.B. Gibbs, *Physica* 3, 159, 145 (1989).
3. E.D. Isaacs et al., *Phys. Rev. B* 40, 9336 (1989).
4. R.C. Farrow et al., *J. Appl. Phys.*, in press (1991).
5. M.F. Toney et al., *Proc. Mater. Res. Soc.* 231, xxx (1991).

UNDERPOTENTIALLY DEPOSITED THALLIUM ON SILVER (111): TWO-DIMENSIONAL COMPRESSIBILITY

M.F. Toney, J.G. Gordon, M.G. Samant, G.L. Borges, O.R. Melroy, (IBM Research Division, Almaden Research Center), D. Yee and L.B. Sorensen (U. Washington)

Using in-situ surface x-ray scattering, we have investigated the atomic structure of electrochemically deposited Tl on Ag(111). The monolayers were formed by underpotential deposition (UPD) at electrode potentials positive of the reversible potential for bulk Tl deposition. At potentials between -475 and -680 mV (vs Ag/AgCl), the Tl deposit forms an incommensurate, hexagonal two dimensional (2D) monolayer that is compressed relative to bulk Tl and rotated from the Ag [011] direction by $\Omega = 4.5^\circ$.^{1,2} This structure is shown in Fig. 1. Between -680 mV and -710 mV (the bulk deposition potential), Tl/Ag(111) forms a bilayer and this also has an incommensurate, hexagonal structure.

As Fig. 2 shows, the monolayer compresses with decreasing electrode potential (increasing chemical potential). These data permit a calculation of the 2D monolayer compressibility κ_{2D} ,^{2,3} which is essentially proportional to the slope of the data in Fig. 2. We find that κ_{2D} varies linearly with potential from 2.2 Å²/eV at -480 mV to 0.9 Å²/eV at -650 mV and that the average compressibility is $\kappa_{2D} = 1.5$ Å²/eV. The decrease in κ_{2D} with more negative potentials or smaller near neighbor spacing is expected: As the atomic spacing decreases, the atom-atom repulsive force becomes increasingly stronger, and this makes it increasingly difficult to pack the adatoms closer together. The κ_{2D} estimated for a 2D free-electron gas model^{2,3} (0.44 Å²/eV) is in qualitative agreement with the measured κ_{2D} but is not quantitatively correct. A more quantitative explanation awaits a better calculation of κ_{2D} for these heavy metals.

The angle Ω depends on electrode potential and atomic spacing, but irreversibly decreases with potential cycling (which is possibly due to adsorption of trace impurities).² Despite this irreversibility, the dependence of Ω on atomic spacing qualitatively agrees with theory.⁴

1. M.F. Toney et al., Phys. Rev. B 42, 5594 (1990).
2. M.F. Toney et al., submitted to Phys. Rev. B.
3. O.R. Melroy et al., Phys. Rev. B 38, 10962 (1988).
4. J.P. McTague and A.D. Novaco, Phys. Rev. B 19, 5209 (1979).

This work was partially supported by the Office of Naval Research.

Fig. 1. Schematic real-space representation of Tl on Ag(111).^{1,2} The open circles represent atoms in the Ag(111) surface and the shaded circles represent the Tl atoms.

Fig. 2. Dependence of near-neighbor distance a_m on electrode potential (vs Ag/AgCl). The line is a least-squares fit of a quadratic to the data. For bulk Tl the average a_m is 3.432 Å.²

STRUCTURE AND EPITAXY OF ANODIC TiO₂/Ti(1120)

D.G. Wiesler, M.F. Toney, M.G. Samant, and O.R. Melroy (IBM Research Division, Almaden Research Center), W.H. Smyrl, and C.S. McMillan (U. Minnesota).

Thin anodic metal oxide films have a variety of technological applications, including corrosion inhibition, solar energy conversion, electrocatalysis, and use in capacitors and MOS devices. While such films have been studied by a number of techniques, surprisingly little is known about their structure and epitaxy onto the metal underlayer. This situation is particularly true for anodic films of Ti and the goal of these experiments is to determine the anodic oxide structure and epitaxy and relate this to the oxide properties. Here we report X-ray scattering measurements of the anodic oxide formed on Ti(1120) by ramping the potential at 0.1 mV/sec to 9-10 V in 0.05 M H₂SO₄.^{1,2}

We find only fully oxidized TiO₂ present in the film, 45±15% of which is anatase and the remainder rutile. The rutile is orientationally ordered to within 8° of the Ti substrate, with its tetragonal *a*- and *b*-axes in the Ti [1100] and [0001] directions, respectively, and the *c*-axis along the surface normal. This orientational relationship suggests a possible growth model for the rutile.¹ The diffraction peak widths correspond to an average in-plane domain size of 40 Å, which is quite small. The rutile is on average strain-free, but there is some suggestion of inhomogeneous strain of ≈1.2%.

Anatase has domains twice this size, but it is orientationally ordered to a lesser extent than rutile; this is illustrated in the Figure. We have analyzed these and similar data. We find the anatase forms a textured powder with a dominant orientation that has the anatase (013) planes in the surface, rotated azimuthally so that the near-neighbor Ti-Ti bonds are matched as closely as possible between the Ti and the oxide. Because both rutile and anatase have well-defined orientations with respect to the Ti substrate, but not to each other, we propose that both oxides are formed at the metal/oxide interface.

1. M.F. Toney, in *The Application of Surface Analysis Methods to Environmental Materials Interactions*, edited by D.R. Baer et al., (The Electrochemical Society, 1991), p. 200.
2. D.G. Wiesler et al., submitted to Surf. Sci.

This work was partially supported by the Department of Energy.

Rocking curves at different χ for the Anatase (101) peaks. Scans are offset by a variable amount for clarity.

IN-SITU X-RAY SCATTERING DURING ELECTRODEPOSITION UNDER CONTROLLED HYDRODYNAMICS

M.J. Armstrong, G.M. Whitney (IBM Storage System: Products Division), and M.F. Toney, (IBM Research Division, Almaden Research Center).

Previous in-situ X-ray scattering experiments of metal/electrolyte interfaces have been performed in stagnant electrolyte.^{1,2} This limits the processes under investigation to those that do not involve the depletion of ions from the electrolyte. Agitation, however, is desirable for the study of many important processes, such as electrodeposition and corrosion. Here we describe a cell design for X-ray scattering measurements of a metal/electrolyte interface during electrodeposition under controlled hydrodynamics and show results of X-ray reflectivity measurements during NiFe electrodeposition.

The cell design (see Fig. 1) uses an impinging jet flow and the flow pattern is established by three channels cut into a Teflon spacer, which is sandwiched between two Teflon plates.³ The electrolyte enters through the center channel and exits by the two side channels. The substrate is positioned opposite to the end of the center channel. The X rays pass through Kapton windows held in place by the Teflon plates. By changing the thickness of the Teflon spacer, the X-ray path-length through the electrolyte can be varied from 1-10 mm. This allows optimization of the path length for the maximum signal.

Reflectivity measurements were performed during the electrodeposition of NiFe. The substrates consisted of a thin NiFe 'seed layer' (≈100 nm) on smooth Si bars. Figure 2 shows the reflectivity after one hour with a partial current for NiFe deposition of 0.17 mA/cm². The expected deposit thickness was 200 nm. However, the reflectivity indicates the total NiFe film thickness (seed layer plus deposit) is 110 nm, and this thickness varied little during 2.5 hours of deposition. Furthermore, the specular reflectivity decreased and the off-specular scattering increased during NiFe deposition. This suggests that deposition occurred by the formation and growth of small clusters of NiFe.

1. M.F. Toney et al., *Langmuir* 7, 796 (1991), and references therein.
2. B.M. Ocko et al., *Phys. Rev. Lett.* 65, 1466 (1990).
3. M.J. Armstrong, G.M. Whitney, and M.F. Toney, in *X-ray Methods in Corrosion and Interfacial Electrochemistry*, edited by A.J. Davenport and J.G. Gordon, in press.

Fig. 2. In-situ X-ray reflectivity of a NiFe deposit. The energy was 17.4 keV.

Fig. 1. Teflon spacer to establish flow pattern.

THE COMMENSURATE AND INCOMMENSURATE PHASES OF BILAYER XENON ON SINGLE CRYSTAL GRAPHITE.

W.J. Nuttall, K.P. Fahey, M.J. Young, R.J. Birgeneau (MIT)

We continue our investigations of multilayer rare gas films adsorbed on single crystal graphite substrates. It is of fundamental interest to probe the structural properties of such a multilayer film by layer. One might anticipate that the layer nearest the substrate would behave in a way reminiscent of the much studied monolayer and sub-monolayer films. Those layers somewhat distant from the graphite, however, might have a periodicity and structure characteristic of bulk crystals of the adsorbing species.

In order to probe the nature of this crossover we investigated the phase behavior of a bilayer (2.05ML) film of xenon on a single Kish crystal of graphite (mosaic < -0.008° HWHM). We have placed much experimental effort, particularly using scanning angle microscopy, to determine which crystals coming out of the manufacturing process are free of silica contamination. We made a final in situ determination of our substrate's surface cleanliness and crystallographic quality by adsorbing a commensurate $\sqrt{3}\times\sqrt{3}$ monolayer of krypton, ($\xi=2000\text{\AA}$).

For bilayer xenon we found a high temperature (-100K) aligned phase of one distinct incommensurability ($\xi=750\text{\AA}$). On cooling the incommensurability decreases and the peak broadens. At around 56.5K a second xenon peak corresponding to a commensurate structure appears. On cooling the next few degrees the

scattering at the commensurate position increases with a concomitant decrease in the incommensurate signal. Below around 50K, however, the situation appears to have stabilized with a permanent low Q shoulder on a slightly larger commensurate peak. We observe that at low temperatures, say -45K we have ξ_c -250Å and ξ_{cc} -130Å.

One outstanding advantage of a single crystal substrate is the ability to probe orientational and out of plane structure. At several temperatures we scanned up the rod which results from the near 2-D nature of our xenon sample. Such scans allow one to determine such things as the xenon planar stacking sequence or the layer by layer phase structure.

Although data analysis is ongoing it is clear that at 70.2K our incommensurate rod shows a dip at the (Q_c , 0, 0) position, showing that there is a bilayer component to this incommensurate phase. Below the transition we find that the out of plane data at either the commensurate or incommensurate position are monolayer in form.

One of us (WIN) acknowledges the generous support of S.E.R.C. (U.K.).

Polarization Measurements on a 3d Ising system

J.P. Hill, P. Westbrook, R.J. Birgeneau
Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
T.R. Thurston, and Doon Gibbs
Brookhaven National Laboratory, Upton, New York 11973

the data a Ge(440) crystal and an energy of $E=8.766\text{KeV}$ were used. The tighter q -space resolution provided by the germanium crystal discriminates against the multiple scattering peaks which were the major source of the background.

The data clearly show that for $\psi=0^\circ$ the polarization of the (100) peak is rotated by 90° relative to the unchanged (200) peak, thus conclusively demonstrating that at $H=0\text{T}$ all the scattering at the (100) position, which comes in at $T_N=46.4\text{K}$, is of magnetic origin. The solid line in figure 1 is the prediction of Blume and Gibbs shown for comparison. It should be noted that this is the first time that the explicit polarization dependence of the scattered beam on the spin angular momentum has been measured.

As was first realized by Platzman and Tsouar, magnetic x-ray scattering can change the polarization of the scattered radiation. In an elegant paper by Blume and Gibbs, the general case was calculated for the first time, explicitly including the effect of the orbital angular momentum on the final polarization. In MnF_2 , there is negligible orbital angular momentum and the cross section simplifies such that at the (100) Bragg peak the scattering from the Ising spins can be separated into two components. The component of the spin perpendicular to the scattering plane, produces the dominant contribution to the signal but does not alter the polarization of the x-ray beam, while the component in the plane, appears in the off-diagonal matrix elements and connects a horizontally, σ , polarized initial state with a vertically, π , polarized final state and vice versa. Unfortunately the intensity contribution from these elements is reduced by a factor of $\sin^2\theta$ relative to the diagonal elements which makes the experiment technically much harder.

The essence of this experiment was to rotate the spins into the scattering plane and measure the degree of linear polarization of the scattered beam as a function of this rotation angle. Simultaneous measurements of the charge (200) peak provide an excellent check on this procedure.

This test was carried out on the MIT-IBM beamline X20C, at NSLS which has a vertically and horizontally focusing mirror upstream of a double bounce Si(111) monochromator. A crystal of $\text{Mn}_{0.95}\text{Zn}_{0.05}\text{F}_2$ with a mosaic of 0.002° FWHM was used. The degree of linear polarization, $P' = \frac{I_{\sigma} - I_{\pi}}{I_{\sigma} + I_{\pi}}$ (where I is the integrated intensity) is measured by selecting each of the two components independently with the use of an analyzer with a 90° reflection, which is set to diffract in the scattering plane (horizontal polarization, σ detected) or at 90° to it (vertical polarization, π detected). Initially a $\text{Cr}(008)$ crystal and an x-ray energy of $E=10.45\text{KeV}$ were used for the $\psi=90^\circ$ and $\psi=60^\circ$ data. However for the rest of

RAPID CRYSTALLIZATION OF AMORPHOUS CO2B STUDIED BY TIME RESOLVED X-RAY SCATTERING

S. Brauer, H.E. Fischer, J.O. Ström-Olsen, M. Sutton (McGill University); G.B. Stephenson (IBM);

To investigate the crystallization of metallic glasses on short time scales, ribbon samples are abruptly heated to a prescribed temperature while x-ray scattering patterns are repeatedly acquired to monitor the transformation. Increasing the temperature accelerates the reaction. For sufficiently short crystallization times, the path of transformation changes such that new non-equilibrium phases are produced. Although much has been learned from the evolution of these metastable structures, we have long sought a material with a simple single-step transformation from the amorphous to the equilibrium crystal state. Such a model system would provide a more direct test of nucleation and growth models of the crystallization process. The alloy Co_{73}B crystallizes directly into the equilibrium tetragonal phase, even at the fastest transformation rates which we can observe.

The scattering patterns shown in Figure 1 show how the initial broad amorphous scattering evolves into Bragg peaks of the equilibrium crystal phase. The data was collected in reflection with a linear photodiode array position sensitive detector and an incident flux of $\approx 10^{13}$ photons per second at the 1mm^2 beam-spot. Diffraction patterns with 256 pixels can be acquired as quickly as every 11 ms. The solid line fit is a linear combination of the amorphous and crystalline scattering patterns which is used to extract the crystalline volume fraction as a function of time as shown in Figure 2 for seven different crystallization temperatures. The brittle crystalline samples generally break before the transformation is complete. The same data is shown in Figure 3 with the time of each run renormalized. Scaling behavior is clear. Work is underway to understand the resulting master curve in light of nucleation and growth models for the solid state.

Figure 1

Figure 2

Figure 3

Supported in part by the Natural Sciences and Engineering Research Council of Canada.

EVIDENCE FOR TWO-STEP DISORDERING OF THE Au(110) 1x2 RECONSTRUCTION

D. T. Keane*, P. A. Bancel, J. L. Jordan-Sweet, G. A. Held (IBM Watson Research Center, Yorktown Heights, NY), and A. Mak and R. J. Birgeneau (MIT)

We have carried out an x-ray scattering study of the thermal disordering of an Au(110) 1x2 reconstructed surface at the IBM/MIT beamline X-20C. Observing the temperature dependence of the in-plane superlattice and integral order surface peaks, as well as the $(\ell \neq 0)$ specular reflectivity, we find that at $T_c = 735K$ the 1x2 surface undergoes a reversible deconstruction transition. In Fig. 1 below are shown representative scans along the $(0.06 \ 0.06 \ \ell)$ direction. As may be inferred from Figs. 1(a) and (b), below 734K the equilibrium structure of our Au(110) surface is a 1x2 reconstruction. However, above 735K, the $(0.06 \ 0.06 \ \ell/2)$ superlattice peaks broaden significantly and exhibit a sharp drop in peak intensity (Fig. 1(c)). Yet at temperatures as high as 748K, the positions of these peaks remain commensurate with the bulk crystalline lattice. Also, throughout this transition the width and intensity of the $(0.06 \ 0.06 \ 1)$ peak is observed to remain constant. These observations indicate that the transition is characterized by a proliferation of antiphase defects with no measurable change in the density of surface steps. Further, we find that the transition is described by critical exponents consistent with a 2-d Ising transition.

Above $T = 784K$, we find that the widths of both the (001) surface peak and the (110) specular reflectivity have broadened significantly. Also, the $(0.06 \ 0.06 \ 1.5)$ superlattice peak has shifted by $0.015\text{\AA}$ from the bulk commensurate position (Fig. 1(d)). These observations indicate that by this temperature there has been a proliferation

ation of surface steps as well as antiphase defects: this is consistent with a surface roughening transition, the onset of which lies between 752 and 784K.

The MIT component of this work was supported by the Joint Services Electronics Program under Contract No. DAAL-86-K0002.

*Also, Dept. of Physics, Princeton University, Princeton, NJ 08544

1. I. K. Robinson, E. Vlieg, and K. Kern, Phys. Rev. Lett. **63**, 2578 (1989).
2. D. T. Keane, et al, Surf. Sci. **250**, 8 (1991).

X-RAY RESONANCE MAGNETIC SCATTERING IN $\text{Eu}(\text{As}_{0.02}\text{P}_{0.98})_3$

T. Chattopadhyay (ILL), G. Grübel (PRL) and L. Rebersky (PRL)

We have performed X-ray resonance magnetic scattering investigations on the sine-wave and helimagnetic phases of $\text{Eu}(\text{As}_{0.02}\text{P}_{0.98})_3$ at the L_{III} absorption edge of europium. A (001) single crystal plate of $\text{Eu}(\text{As}_{0.02}\text{P}_{0.98})_3$ of dimensions of about $8 \times 8 \times 1 \text{ mm}$ with a mosaic spread of about 0.02° was mounted inside a helium cryostat of the four-circle diffractometer with its monoclinic b axis parallel to the ϕ axis of the diffractometer. Search for magnetic intensity at $Q = (1.27, 0, 2.221)$ at 6 K was not successful when the X-ray energy was far away from the L absorption edges. Also search at this Q did not reveal any magnetic intensity with X-ray energy tuned to the L_{III} absorption edge. But when the X-ray energy was tuned to the L_{III} absorption edge sharp magnetic reflection was observed. Fig. 1 shows the background and the integrated intensity of the $1.27, 0, 2.221$ reflection of the helimagnetic phase of $\text{Eu}(\text{As}_{0.02}\text{P}_{0.98})_3$ as functions of the X-ray energy about the L_{III} absorption edge of europium. The resonance line shape is asymmetrical with shoulder at the low energy side. We have investigated the sine-wave-to-helimagnetic-transition in $\text{Eu}(\text{As}_{0.02}\text{P}_{0.98})_3$ by X-ray resonance magnetic scattering. Fig. 2 shows the temperature variation of the intensity of the $1.27, 0, 2.22$ magnetic reflection and the z component k_z of the wavevector parallel to c close to the sine-wave-to-helimagnetic phase transition. The intensity and the wavevector show abrupt change at $T = 7.5 \text{ K}$ showing the first-order sine-wave-to-helimagnetic phase transition.

Figure 1

Figure 2

X-RAY RESONANCE MAGNETIC SCATTERING IN EuTe

T. Chattopadhyay (ILL) and G. Grübel (BNL)

We have investigated the magnetic ordering of the type-II antiferromagnet EuTe by X-ray resonance magnetic scattering at the L_{II} and L_{III} absorption edges of europium with synchrotron radiation. A (100)-single crystal plate of EuTe of dimensions of about 8 mm x 8 mm x 2 mm with a mosaic spread of 0.03° was mounted inside the helium cryostat of the four-circle diffractometer with one of its cubic axis parallel to the ϕ -axis of the diffractometer. Search for magnetic intensity at $Q = (1/2, 1/2, 9/2)$ was not successful when the X-ray energy was far away from the L absorption edges. However, sharp magnetic reflection was observed at this Q when the X-ray energy was tuned close to the L_{II} and L_{III} absorption edges. Fig. 1 shows the temperature variation of the intensity of $1/2, 1/2, 9/2$ magnetic reflection at $E = 7610$ eV. The intensity of this reflection decreases continuously and becomes zero at $T_N = 10$ K confirming its magnetic origin. Fig. 2 shows the energy variation of the background and the integrated intensity of the $1/2, 1/2, 9/2$ magnetic reflection at $T = 7.0$ K at the L_{III} absorption edge. The resonance line shape is asymmetric and has a shoulder at the lower energy side as was also observed in EuAs, previously by us [1]. However, the resonance line width in the metallic EuAs₃ is much larger than that of the insulating EuTe. The two experiments were performed under almost identical conditions and therefore we suspect that the resonance line widths of these compounds might contain quantitative information about their magnetic electron structures. Detailed calculations to extract such information are in progress.

1. T. Chattopadhyay, G. Grübel, J.D. Axe and D. Gibbs, J. Mag. Mater. (in press)

Figure 1

Figure 2

X-RAY REFLECTIVITY STUDY OF INTERFACE GROWTH DYNAMICS: IN SITU SPUTTER DEPOSITION OF Au ON SILICON*

R.P. Chiarello, H.K. Kim, and H. You (ANL)

Dynamic scaling behavior of an interface during growth and effect of growth conditions on the nature of interface have been the subject of many theoretical and computational studies [1]. Motivated by the computational and theoretical studies we experimentally study the interface growth behavior using the sputter deposition technique.

A brief description of the chamber used in this study is given in the report for X6B beamline of this volume. During the measurements the plasma was on but the deposition rate was reduced so that no detectable amount of deposition can occur during the x-ray scans. Measurements were made at several substrate temperatures and at several partial pressures of Ar. For each condition a series of reflectivity scans were made at various film thicknesses.

Three pairs of reflectivity scans for different thickness of films are shown in the figure. Two scans in each pair are for the same thickness but for low (upper scan) and high (lower scan) Ar partial pressures. The second pair shows most clearly the difference that we want to point out. In the upper curve the amplitude of oscillation is small but the oscillation persists to $0.5 \text{ \AA}^{-1}$ while in the lower curve the amplitude is large for the small Q range, but rapidly damps out and shows no oscillation for higher Q . Without any speculation, it seems clear from the data that x-ray reflectivity scans manifest the nature of the interface profile. Data analysis is in progress for a real space description of the interfaces at various phases of interface growth.

[1] for an example see C. Tang, S. Alexander, and R. Bruinsma, Phys. Rev. Lett. **64**, 772 (1990).

*Supported by DOE under Contract # W-31-109-Eng-38

RESONANT MAGNETIC SCATTERING IN EUROPIUM COMPOUNDS

G. Grübel, J. D. Axe, L. Rebersky, Doon Gibbs (BNL), and T. Chattopadhyay (CEN and ILL, Grenoble)

Resonant magnetic scattering experiments were performed at beamline X22B in semimetallic antiferromagnetic EuAs_3 and semiconducting, antiferromagnetic $\text{Eu}(\text{P}_{0.88}\text{As}_{0.02})_3$ at the L_{II} or L_{III} absorption edges of Eu . When the energy of the incident x-rays was tuned close to the L_{II} or L_{III} absorption edges a strong resonant enhancement of the magnetic scattering intensity was observed in both systems. Figure 1 shows the results for EuAs_3 . The absorption coefficient (top) was calculated from the measured fluorescence signal. The lower panel (filled symbols) shows the integrated intensity (corrected for absorption) of the $(1\ 0\ 5/2)$ magnetic reflection at 6.2K as a function of the incident x-ray energy. The lineshapes of both resonances are asymmetric. A polarization analysis at the L_{II} edge shows that the polarization of the scattered beam is unrotated (π - π). The open symbols in Figure 1 show the π - π component of the magnetic scattering (scaled by a factor 7.5). This result is in agreement with the predictions of the X-Ray Resonance Exchange Scattering Model (XRES) for the polarization response in EuAs_3 and incident π polarization. The asymmetric energy dependence of the scattering intensity can be described as a result of the interference between the π components of the non-resonant and resonant cross-sections. The solid line in Figure 1 (bottom) is a fit to the data assuming a model that employs only dipolar transitions ($2p_{3/2} - 5d$) for the resonant cross-section. The experiments show no evidence for a quadrupolar contribution to the observed resonances.

POTENTIAL INDUCED (1×3) RECONSTRUCTION OF THE $\text{Au}(110)$ SURFACE

B. M. Ocko, G. Helgesen, B. Schardt, J. Wang (BNL), and A. Hamelin (CNRS)

In vacuum, the clean $\text{Au}(110)$ surface has been shown to form a (1×2) reconstruction. The ability to induce a reconstructed $\text{Au}(110)$ surface, in an electrochemical cell, has been suggested on the basis of the hysteresis in the "double layer" capacitance and from the frequency dispersion of the alternating signal in a limited range of potential. In order to investigate the structure of the $\text{Au}(110)$ electrode surface, we have carried out surface x-ray diffraction measurements in a variety of salt solutions as a function of the applied potential. At sufficiently negative potentials, in 0.1M NaF , NaCl , NaBr , LiCl , and CsCl , we observe a (1×3) reconstruction with (111) microfacets. Above a critical threshold potential, the (1×3) reconstruction vanishes and the surface forms a (1×1) structure. The threshold potential depends on the electrolyte, and in all solutions the transition between these two structures is completely reversible. X-ray diffraction peaks from the reconstructed surface are observed at $L = 1/3, 5/3, 7/3$, and $11/3$ in the grazing incidence geometry (see Fig. 1). Therefore, the $\text{Au}(110)$ surface in 0.1M NaCl exhibits a three-fold increase in the size of the unit cell. Analysis of the diffraction profiles provides a direct measure of the size of the correlated reconstructed domains. The best fit in 0.1M NaCl gives a correlation length of $50 \pm 5\text{ \AA}$ along the ridge direction and $250 \pm 25\text{ \AA}$ along the row direction. The phase behavior has been measured by varying the potential at fixed reciprocal space positions (see Fig 2).

Fig. 1 The intensity difference along $(0\ 1\ 0\ .1\ .1)$ between profiles at -0.3 and 0.0 V in 0.1 M NaCl . Peaks are at $L=1/3, 5/3$, and $7/3$.

Fig. 2 (A) Differential capacitance from the $\text{Au}(110)$ surface in 0.1 M NaCl , in the positive going sweep, at 8 mV/sec at $8(\text{top})$, $12, 20, 80, 160\text{ Hz}$. (B-D) Potential dependence of the scattered intensity from the $\text{Au}(110)$ electrode in 0.1 M NaCl at 1 mV/sec at three wavevectors.

X-RAY CRYSTAL TRUNCATION ROD ANALYSIS OF UNTWINNED $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ SINGLE CRYSTALS: TERMINATION FINGER PRINT

H. You, U. Welp, G. W. Crabtree, and Y. Fang (ANL), S.K. Sinha and J.D. Axe (BNL), and X. Jiang and S.C. Moss (U. Houston)

X-ray scattering from a half infinite three dimensional single crystal, known as Crystal truncation rod (CTR), [1] was measured for untwinned single crystals of $\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$. CTR results from the abrupt termination of electron density at the surface of the single crystal and can be expressed as a convolution of x-ray reflectivity and Bragg reflections. [2] In this expression the scattering amplitudes from every Bragg reflections with the same $Q_{\parallel}$ components are summed with appropriate phase factors which depend on the position of the terminating plane, thereby, signifying the position of terminating plane under a favorable condition.

The measurements were carried out at 10 K with a wavelength of 1.55 Å in a vertical scattering geometry. Data were taken at various L values in a geometry where the angles of incident and exit x-rays with respect to the sample surface were same.

The data (open circles) along (00L) and (10L) rods and fits (solid lines) to them are shown in Fig. 1. The fits to both (00L) and (10L) data independently find a Y termination out of six possible metal containing layers in unit cell. To be more specific the terminating plane is found to be slightly higher than that of the bare Y layer indicating that Y atoms could have been oxidized. This is an example where a careful CTR analysis can be used in determining the top growth plane without elaborate surface preparations.

[1] I.K. Robinson, Phys. Rev. B33, 3830 (1986).

[2] H. You, in *Proceedings of the 2nd International Conference on Surface X-ray and Neutron Scattering*, Bad Honnef, Germany, 1991, edited by H. Zabel and I.K. Robinson (Pergamon Press).

*Supported by DOE W-31-109-Eng-38, DOE DE-AC02-76CH00016, NSF-8809854, and NSF DMR-8903339

X-ray Diffuse Scattering From Poly(bromostyrene) PBs/Toluene Solutions
W. Zhao, X. Zhao, M. Rafailovich, J. Sokolov (Queens College), M. Sanyal (BNL), S.K. Sinha (BNL, Exxon)

The diffuse scattering from PBs (molecular weight = 1M) /toluene solutions as a function of PBs concentration up to 10% weight/weight) was measured for q_z (normal to sample surface) = 0.1 to 0.3 Å⁻¹. Diffuse scattering from small-molecule liquids such as water and ethanol is well understood in terms of thermally excited capillary waves[1]. However, polymer solution exhibit strong viscoelastic effects and have a rich variety of hydrodynamic modes not present on simple liquids [2].

Initial measurements were made for the pure solvent, toluene, which is expected to have similar scattering to ethanol. Using the known surface tension of 21.7 dyne/cm, calculations based on a capillary wave model were used to fit the data. Good agreement between theory and experiment was found and the results are shown in Fig. 1. The scattering from various PBs/toluene solutions was measured which showed that the basic line shape of diffuse scans for each q_z was similar to that of toluene but the variation of overall intensity with q_z increased with increasing polymer concentration.

Calculations of the surface height-height correlation function $\langle Z(R)Z(0) \rangle$ were made as a function of the elastic shear modulus G . For moderate G (100-1000 dyne/cm²), it can be shown that

$$\langle Z(R)Z(0) \rangle = \frac{K_B T}{2\pi\gamma} \left[\ln\left(\frac{R}{2}\right) - \gamma_E - 2\ln\left(\frac{4G}{\kappa\gamma}\right) \right]$$

where K_B = Boltzman constant, T = temperature, γ = surface tension, κ = gravitational cutoff and γ_E = Euler's constant, 0.5772. For $G = 0$, only the last two terms are present which is the capillary wave result. With this approximation $\langle Z(R)Z(0) \rangle$ has the same functional form as for capillary waves but has an added G -dependent constant.

The values of γ were measured independently (in collaboration with B. Sauer of DuPont) and comparison of calculated and experimental diffuse spectra yields values for G of 120 dyne/cm² and 180 dyne/cm² for 3.3% and 5.75% PBs/toluene solutions, respectively. Fig. 2 shows the results, using best-fit values of G . The values for G are in reasonable agreement with those extrapolated from light scattering experiments on a lower molecular weight (90K) 18% PBs/toluene solution [3].

[1] Schwartz et al. Phys. Rev. A31, 5687 (1990), Sanyal et al. Phys. Rev. Lett. 66, 628 (1991).

[2] Harden et al. J. Chem. Phys. 94, 5208 (1991).

[3] Cao et al. to be published.

This work is supported by the NSF(DMR-8921556), DOE (DE-FG02-90ER45437) and the Petroleum Research Fund.

D. L. Abernathy, S. G. J. Mochrie (MIT), D. M. Zehner (ORNL), G. Grubel, D. Gibbs (BNL)

Previous work has shown that the Pt(001) surface reconstructs below 1820K to form a close-packed, hexagonal layer on top of the bulk (001) planes. Above 1820K, the in-plane diffraction pattern exhibits only the square symmetry of the truncated bulk. New transverse lineshape measurements across the specular rod show that there is a dramatic change in surface morphology on going from the reconstructed, lower temperature phase to the high temperature phase. [below left] At 1790K the transverse lineshape at $Q_z(001)$ is resolution limited, indicating that the surface in the reconstructed phase is atomically smooth on length scales exceeding 5000Å. At 1850K, however, the lineshape is broad and diffuse, showing that the surface is rough. The scattering from the surface for out-of-plane momentum transfer of $0.5 < Q_z < 4.5$ in the high temperature phase can be modeled using a logarithmically divergent height-height correlation function, which is a slightly modified version of that used for liquid surfaces. The roughness exponent extracted from fitting the lineshapes is shown at right below. As expected, it is a periodic function of Q_z with minima at the Bragg reflections and maxima at the anti-Bragg positions. One surprising result is that the capillary modes cutoff wavevector corresponds to a length of ~170Å.

This work is supported by US DOE DE-AC0276CH00016, US DOE DE-AC05-84OR21400, NSF 8806591 and JSEP DAAL03-89-C-0001.

ORDER, DISORDER AND STRUCTURE OF CRYSTALS OF C₆₀/C₇₀

J. Bohr (RISO), Doon Gibbs (BNL), S. K. Sinha (EXXON), W. Krätschmer (Max-Planck), G. Van Tendeloo (RUCa), E. Larsen (RISO), H. Egsgaard (RISO), and L. E. Berman (BNL)

Single crystals with two different morphologies, black and brown, of C₆₀ with about 12% C₇₀ have been studied by synchrotron x-ray diffraction. Intensities from 8 reflections show that the charge distribution of the C₆₀ cluster is shell-like giving rise to an oscillatory behaviour in the diffracted intensities. From the intensities, the distance between diametrically opposite carbon atoms has been determined to be 7.24 ± 0.22 Å. The thickness of the charge distribution of the shell can be estimated as 3.06 Å and the cavity within a C₆₀ cluster to have a diameter of about 4.18 Å. The finite longitudinal width of the diffraction peaks from black crystals show lack of long-range crystalline order. The rocking curves are about 7° broad and depict an unusually smooth behaviour. This may be indicative of a glassy or hexatic phase. High resolution electron microscopy allows identification of small crystallites with a relatively well defined orientation relationship. On the contrary, brown crystals have long-range order.

MAGNETIC X-RAY SCATTERING STUDY OF Tb

P. M. Gehring, L. Rebersky, Doon Gibbs, and G. Shirane
Brookhaven National Laboratory, Upton, New York 11973

The temperature and energy dependence of the magnetic scattering in the ferromagnetic spiral phase of Tb has been studied between 220 K and 235 K by high resolution x-ray scattering techniques using synchrotron radiation. In order to see the magnetic scattering it was necessary to exploit the resonant enhancement of the magnetic x-ray cross section by tuning the incident photon energy to the Tb L_{III} edge. A lower bound for this enhancement is of order 20.

The magnetic modulation wave vector τ_m decreases continuously with decreasing temperature over most of the range between the Néel and Curie points in agreement with prior neutron diffraction measurements. No evidence was found for any lock-in behavior above T_c , as has been observed in other rare-earths. However spin-slip scattering was seen at $4 - \tau_c$ ($\tau_c = 12\tau_m - 2$) as shown in Fig. 1.

A very strong hysteresis effect is present in Tb below T_N , as the value of the modulation wave vector depends on the rate of cooling from the paramagnetic regime. In addition, the radial linewidth of the magnetic satellite at $(002)^+$ is not resolution-limited and increases as $T \rightarrow T_c^+$ as shown in Fig. 2. By comparison, that of the charge peak at (002) remains constant. No conclusive evidence was found for scattering at higher harmonics.

Fig. 1

Fig. 2

Research at Brookhaven is supported by the Division of Materials Sciences, U. S. Department of Energy under Contract No. DE-AC02-76CH00016.

Reconstruction of the Pt(111) Surface.
G. Grübel, K.G. Huang, Doon Gibbs(BNL*), D.M. Zehner(ORNL**),
A.R. Sandy, and S.G.-J. Mochrie(MIT)

The structure of the clean Pt(111) surface has been studied between 300K and 0.90T_m (T_m=2045K) via x-ray scattering. The surface is unreconstructed for temperatures less than 0.65T_m, but reconstructs at higher temperatures to form a layer isotropically compressed and incommensurate with the underlying bulk (111)-planes. A disordered arrangement of discommensurations separates regions with ideal fcc stacking from regions with faulted stacking. With increasing temperature the compression of the surface layer increases. Figure 1 (closed circles) shows the incommensurability δ as a function of the normalized temperature (T/T_m). Between 0.65T_m and 0.75T_m the surface layer forms a translationally disordered discommensuration fluid. As shown in the inset, the incommensurability follows a one-third order power law versus reduced temperature (T-0.65T_m)/(0.65T_m). Above 0.75T_m orientational order appears and grows continuously with further increase in temperature. This is illustrated in Figure 2 showing the scattering intensity taken on a circular path (radius δ) around the crystal truncation rods. The modulation in these scans indicates the development of orientational order and is described by $I(\theta) = (1 + A_0 \cos(6\theta))$. The temperature dependence of the orientational order parameter A_0 is also displayed in Fig. 1.

This work is supported by U.S. DOE contract No. DE-AC02-76CH00016, and DMS, Office of Basic Energy Sciences, U.S. DOE Contract No. DE-AC05-84OR21400 with Martin Marietta ES, Inc., NSF Grant No. DMR-8806591

In-plane structural modulation of the CaF_2 thin film on $\text{Si}(111)$

K. G. Huang (BNL)*; J. Zegenhagen, J. R. Patel and J. M. Phillips (AT&T Bell Labs)

The in-plane structure of a thin CaF_2 film (50 Å thick) epitaxially grown on $\text{Si}(111)$ has been studied using glancing incidence x-ray diffraction. The sample was grown in an MBE chamber and studied with x-rays under normal ambient conditions.

From x-ray diffraction in glancing incidence geometry, the average in-plane lattice mismatch between the CaF_2 overlayer and the Si substrate was determined to be 1.6% which is larger than the bulk value (0.6% at RT). This is illustrated in Fig. 1 where the peak separation corresponds to an incommensurability of 0.016 between the overlayer and the substrate. The in-plane tensile strain (1%) of the thin film results from the presence of interfacial misfit dislocations. In addition, the in-plane diffraction pattern of the CaF_2 overlayer exhibits an hexagonal symmetry with the overlayer diffraction peaks decorating each of the substrate rod. This suggests that the strain of the overlayer is nearly isotropic within the (111)-plane. Fig. 2 shows a circular scan containing six overlayer peaks, which was taken around the (01)_A substrate crystal truncation rod with a radius of the incommensurability (0.016). The existence of the satellite peaks (see Fig. 1) can be understood by the fact that the overlayer is not simply incommensurate to but modulated in two dimension by the Si substrate. The detail data analysis is underway.

* Work supported by US DOE contract No. DE-AC02-76CH00016.

* Now at MSD, Argonne National Laboratory.

Fig. 1

Specular and Diffuse Scattering From ErAs Buried Epitaxial Layers

P. F. Miceli (Bellcore), M. Sanyal (BNL), S. K. Sinha (Exxon) and C. J. Palmström (Bellcore)

Conventional low angle reflectivity has found many uses in studying the structure of laminar systems. Recently, it was demonstrated^{1,2} that the extended range of x-ray reflectivity, which is measured from low angles through the regions of Bragg scattering at higher angles, provides a new powerful method to investigate buried crystalline layers and interfaces. Reflectivity, however, measures interface fluctuations along the film normal and information regarding lateral correlations of these fluctuations is contained in the scattering transverse to the reflectivity rod.

This work on the ErAs/GaAs epitaxial system initiates a new program to study specular and diffuse scattering, measured across the extended reflectivity, of buried epitaxial layers and interfaces. ErAs is ideal for these studies since Er is a strong scatterer of x-rays and has an extremely high melting temperature, prohibiting interdiffusion between layers. Furthermore, various stages of lattice relaxation can be achieved depending on the film thickness.

In the figure below we show a transverse scan for 3 monolayers of ErAs buried by 100 Å of amorphous Si. The data were collected at $2.02 \text{ \AA}^{-1}$ which corresponds to Bragg scattering coming from the thin ErAs layer. From the fit shown by the solid curve, we find there are both specular and diffuse components to the line shape. Data such as these should allow a detailed study of interfaces and our work is in progress.

1. X-ray Extended Reflectivity of Heteroepitaxial Layers, P. F. Miceli, C. J. Palmström and K. W. Moyers, submitted.
2. X-ray Bragg Reflectivity of ErAs Epitaxial Films, P. F. Miceli, C. J. Palmström and K. W. Moyers, Surface X-ray and Neutron Scattering, Springer-Verlag, in press.
3. X-ray Scattering Study of Lattice Relaxation in ErAs Epitaxial Layers on GaAs, P. F. Miceli, C. J. Palmström and K. W. Moyers, Appl. Phys. Lett. **58**, 1602 (1991).

Au(113) SURFACE STUDIES

A. R. Sandy, D. L. Abernathy, and S. G. J. Mochrie (MIT); D. M. Zehner (ORNL); L. D. Gibbs (BNL).

High-resolution x-ray-scattering measurements of the phase behavior of the Au(113) surface have been performed between 300 K and the bulk melting temperature of Au (1337.4 K). The (113) face of a face-centered-cubic (fcc) crystal lies almost exactly between the low-index (111) and (001) faces so we expect that studies of the thermal behavior of this surface will provide information on the equilibrium crystal shape of Au. The ideal (113) surface consists of parallel, closest-packed rows separated from one another by $\sqrt{11}/2$ nearest-neighbor separations. We find that below 700 K the surface is reconstructed and that the reconstruction is incommensurate with the underlying bulk. Fig. 1 shows a scan obtained by varying the momentum transfer perpendicular to the closest-packed rows while keeping the momentum transfer along the surface normal direction constant. Peaks at even integers are the bulk truncation rods, while those marked by the vertical dashed lines are due to the surface reconstruction. Further analysis will reveal the nature of the reconstruction.

Fig. 1

ANOMALOUS REFLECTIVITY MEASUREMENT OF ELECTRON DENSITY PROFILES IN THIN FILMS

M.K. Sanyal, S.K. Sinha, A. Gibaud, K.G. Huang (BNL); M. Rafailovich, J. Sokolov, X. Zhao, W. Zhao (Queens College, NY); S.A. Schwarz (Bellcore, NJ)

X-ray reflectivity techniques are being increasingly used to determine electron density profiles of surfaces and interfaces. However, since only intensity information is available, the current methods necessitate fitting of an assumed model to the specular reflectivity profiles. A new technique, termed anomalous reflectivity, has been developed and used to determine in a model independent way the electron density profile across a thin film of 15% polybromostyrene (PBS) [M.W.=20K] dissolved in polystyrene (PS) [M.W.=4K], in order to determine the preferential absorption of the PBS at the film surfaces. The film was deposited on a highly polished (100) surfaces of a single crystal germanium wafer.

The measurement of the specular reflectivity close to and away from an x-ray absorption edge of the substrate is shown, using the Distorted Wave Born Approximation for the scattering, to yield a unique model-independent solution for the electron density profile of the film. The results obtained are in good qualitative agreement with the results of SIMS measurements, but have better spatial resolution.

The method we use is based on the fact that the reflectivity of a thin film on a substrate involves an interference between the reflectivity of the film and that of the substrate and hence contains phase information. In order to obtain both the real and imaginary parts of $\rho(q_z)$, which is the Fourier transform in the z-direction of the scattering density, one needs to measure reflectivity at and away from an x-ray absorption edge of the substrate. Finally the electron density $n(z)$ in real space is determined by Fourier transforming the obtained $\rho(q_z)$. The experiments were carried out in X22C beamline and further work using this technique is underway.

DETERMINATION OF ELECTRON DENSITY PROFILE OF A THIN FILM USING ANOMALOUS REFLECTIVITY

M. K. Sanyal (BNL), S. K. Sinha (Exxon), A. Gibaud (U. du Maine), K. G. Huang (ANL), B. L. Carvalho (MIT), M. Rafailovich, J. Sokolov, X. Zhao, and W. Zhao (Queens).

The methods used to determine the scattering density profiles from a reflectivity curve rely on fitting of a model form for the density profile. It is not obvious that the solution obtained by this fitting procedure is unique, since it is an intensity which is being fitted and phase information is absent. We present a method of obtaining a model independent electron density profile for a thin film deposited on a substrate using x-ray reflectivity measurements carried out at energies close to, and away from a convenient absorption edge of the substrate. One is then able to obtain the real and imaginary parts of $\rho(q_z)$, the Fourier transform of the (x, y) averaged electron density of the film. In order to test the method conceptually, we considered a Ge substrate and modelled absorbed films of various electron density profiles such as (a) a uniform bilayer of two different densities, (b) an exponentially decreasing density profile, (c) a parabolic density profile, and (d) a linear density profile. We then simulated "measured" reflectivities by calculating these using the "slicing" method at the two different energies used in the actual experiments. In Fig. 1, these model profiles along with the calculated profiles obtained from this method are shown.

We now illustrate the method with a synchrotron x-ray experiment carried out on a film consisted of $3\frac{1}{2}$ bilayers of Cd -arachidate Langmuir-Blodgett film deposited on a polished germanium substrate. Fig. 2 shows the electron density profile obtained by our method, together with the estimated error bars as calculated from the statistical errors in the observed counts. The resulting profile shows three peaks due to the Cd counterions associated with the carboxylic head groups, while the peak at the Ge -surface is not evident as it is presumably extremely sharp and merges with the sudden rise in electron density at the Ge /film interface. This histogram was used to generate the reflectivity curves at and away from the absorption edge by means of the slicing method which were then folded with the instrumental resolution. These curves are shown together with the experimental data in Fig. 3.

Figure 1

Figure 2

Figure 3

RESONANCE X-RAY MAGNETIC SCATTERING STUDY OF ERBIUM

M. K. Sanyal, D. Gibbs (BNL), J. Bohr (RISØ), and M. Wulff (RISØ and ESRF)

In continuation of our x-ray magnetic scattering studies of rare earth metals, we have measured the polarization and resonance properties of the magnetic cross-section of erbium as a function of temperature. The energy dependence of the line shape of this scattering at the magnetic wave vector τ_m was analyzed near the L_{III} absorption edge. This analysis was done using a previously described formalism $1/1$ with the assumption that the observed magnetic scattering arises from a structure combining both c -axis and basal-plane components.

Figure 1

The evolution of this line shape with respect to temperature is determined by the tilt angle of the moment. In Fig. 1, we summarize our results as a function of the temperature and fitted tilt angle. These are in good agreement with results obtained by neutron scattering $/2/$. We do not observe similar line shapes at temperatures lower than 20K, in the so-called conical phase. In contrast, our study shows that scattering intensity at τ_m exhibits peaks and valleys as a function of energy extending even 2keV below the L_{III} absorption edge. We also observe unusual behaviour in the energy dependence of the scattering at $2\tau_m$ at temperature 40K (shown in Fig. 2). Detailed analysis of this data are in progress.

$/1/$ D. Gibbs, G. Grübel, D. R. Harshman, E. D. Isaacs, D. B. McWhan, D. Mills, and C. Veier. Phys. Rev. B **43**, 5663 (1991).

$/2/$ M. Habenschuss, C. Stassis, S. K. Sinha, H. N. Deckman, and F. H. Spedding. Phys. Rev. B **10**, 1020 (1974).

Figure 2

Magnetic Critical Scattering in Holmium

T.R. Thurston (BNL), Doon Gibbs (BNL), J.P. Hill (MIT) and G. Shirane (BNL)

A number of studies have shown that the order parameter of a phase transition can be directly measured using magnetic x-ray scattering techniques.¹ However, until now no measurement of the short range critical correlations above a magnetic transition has been performed. This report describes preliminary measurements of the short range magnetic correlations in the spiral antiferromagnet holmium.

X-ray scattering measurements of the critical correlations offer several advantages over neutron scattering measurements, the standard technique. Principle among these is the complete integration over energy of the fluctuations possible with x-rays. In some problems the better spatial resolution and surface sensitivity afforded with x-rays can also be advantageous.

Holmium was studied because of its large moment (~10 Bohr magnetons) and because it has already been carefully examined in previous² resonant scattering work. The system undergoes a number of magnetic phase transitions, but the one of current interest occurs at ~138K. The spin system is disordered above this temperature, and below it orders in an incommensurate spiral structure.

The figure shows the half width at half maximum (HWHM) of the magnetic peak in both the longitudinal and transverse directions for temperatures above the transition. The photon energy was set at 8067eV, the *L*_{III} absorption edge for holmium. It is clear that the peak broadens significantly above the

J.P. Hill *et al.* Phys. Rev. Lett. 66, 3281 (1991).
 D. Gibbs *et al.* Phys. Rev. B 43, 5663 (1991).

transition, indicating that only short range magnetic correlations exist.

Although further studies are necessary in order to make quantitative comparisons between x-ray and neutron results for the correlation length, these results suggest that magnetic x-ray scattering techniques will be viable for this type of measurement. Among the issues to be addressed in future studies is whether the resonant magnetic x-ray cross-section measures a 2-spin or 4-spin correlation function.

Magnetic structure in U_{0.85}Th_{0.15}Sb using resonant X-ray scattering

C.C. Tang, W. Stirling (U. of Keele, UK), G.H. Lander (EITU, Karlsruhe),
 D. Gibbs (Physics Dept., BNL), O. Vogt (ETH, Zürich)

Resonant X-ray scattering in uranium compounds^{1,2} is sufficiently strong at the *M*_{IV} edge (*E* = 3.73 keV) that detailed studies of magnetic structures can be performed. The scans shown in the figure below were taken at this energy from a single crystal of U_{0.85}Th_{0.15}Sb in the duplex arrangement at X22C. Special evacuated beam tubes (constructed at Keele) were installed to minimize absorption and allowed reproducible integrated intensities to be obtained. Although work on this system with neutrons was performed some years ago³, we have discovered some new effects.

The material initially orders at *T*_N = 192K with an incommensurate wavevector *q* = (0, 0, 0.433). As *T* is decreased a number of other *q* vectors appear including (at *T* ~ 170K) a component at the commensurate value of *q* = 1/2. At *T* ~ 145K there is the sudden appearance of the *q* = 1/4 and 3/4 commensurate peaks. By 10K, see (b), this phase dominates the pattern, although a strong *q* = 1/2 peak is still seen.

High-resolution scans with *E* = 8keV + analyzer have provided evidence for a volume discontinuity and a substantial increase in the mosaicwidth (transverse scans) at *T* ~ 155K. No change is seen in the radial width, indicating the material is not two phase in nature.

Initial indications are that the ratios *I* (*q* = 1/4) / *I* (*q* = 1/2) are substantially different as measured by X-rays and neutrons. This could be due to surface effects as the X-rays have a penetration depth of less than 5000 Å. Further neutron experiments are planned.

Figure Scans with *E* = 3.73 keV along the (00*l*) direction at (a) 155K and (b) 10K from a single crystal of U_{0.85}Th_{0.15}Sb. The hatched peaks are nonmagnetic arising (mostly) from higher order light.

References:

1. D. McWhan *et al.*, Phys. Rev. B 42, 6007 (1990)
2. C.C. Tang *et al.*, submitted to Phys. Rev. B (April 1991)
3. J. Rossat-Mignod *et al.*, "Handbook of the Actinides", Vol.1, p.469

SPATIALLY SELECTIVE LOCAL STRUCTURAL INFORMATION FOR InGaAs MULTILAYERS USING DIFFRACTION ANOMALOUS FINE STRUCTURE

C.E. Bouldin, J. Woicik (NIST)
H. Stragier, D. Yee, C. Bitz, L.B. Sorensen, K. Kuhn (U. of Washington)

We have used Diffraction Anomalous Fine Structure (DAFS) to obtain XAFS-like local structural information (i.e., near-neighbor bond lengths, coordination numbers and bond-length disorders) for spatially selected regions of an InGaAs multilayer sample.

The sample contained six $\text{In}_x\text{Ga}_{1-x}\text{As}$ layers with increasing In concentrations grown on a (001) GaAs substrate. The top two layers were each 5000 Å thick and had In concentrations of $x = 0.39$ and $x = 0.25$, respectively. As shown below, each layer produced a distinct diffraction peak, allowing the Ga and As K edge DAFS signals for each layer to be measured without interference from neighboring layers even though those layers also contain Ga and As. The Fourier transforms of the As edge DAFS signals measured using the (004) Bragg reflections of the top two layers are compared below with the fluorescence XAFS signal from the sample (which averages over all the layers). The first-shell peak, at $R = 2\text{Å}$, reflects the first-neighbors of the As atoms, which are In and Ga. A large In shoulder near 2.5Å is visible for the top layer with $x = 0.39$, and a much smaller In shoulder is visible for the $x = 0.25$ second layer. There are also visible changes in the higher shells.

These preliminary results show that it is possible to obtain spatially selected local structural information using DAFS.

X-ray Diffuse Scattering Study of Liquid Polymer Films
W. Zhao, X. Zhao, M. Rafailovich, J. Sokolov (Queens College), M. Sanyal (BNL), S.K. Sinha (BNL, Exxon)

We have studied the temperature dependence and the influence of film thickness on the x-ray diffuse scattering of several amorphous liquid polymer systems: polyethylene propylene (PEP), polystyrene (PS), polybutadiene (PBd), and PB/PBd interfaces, all mounted on highly polished Si wafers and measured in-situ in both the liquid and quenched states. For q_z of $0.1 - 0.3\text{Å}^{-1}$, diffuse scattering rocking curves were taken as well as specular scans up to $q_z = 0.5\text{Å}^{-1}$. Sample temperature was varied from room temperature to approximately 300°C for PEP this range is considerably above the bulk glass transition temperature T_g of $\sim 55^\circ\text{C}$ while for PBd and PS T_g is $\sim 110^\circ\text{C}$. Films were spun-cast onto Si substrates with thicknesses from about 150Å (which is the same order as the radius of gyration, R_g , of a single polymer chain in the melt) up to 3000Å .

Films of PEP show a strong temperature dependence, with a substantial decrease of the intensity of diffuse scattering in the near-specular region with increasing temperature, the effect being larger for larger values of q_z . Fig. 1 shows rocking scan spectra for $q_z = 0.25\text{Å}^{-1}$ for $T = \text{RT}$ and 235°C , the decrease in diffuse scattering at the higher temperature reaches nearly an order of magnitude. This behavior is cyclical; samples could be heated and cooled and reheated with the spectra exhibiting good reproducibility. There was also reproducibility from sample to sample. The diffuse scattering is sensitive to the correlations of surface fluctuations [1] and correlations calculated in a capillary-wave model were tried but fail to predict the observed temperature dependence. It is probable that the viscoelastic properties of the polymers have to be taken into account. One explanation may be that the polymer entanglement length, L_e (which is a measure of the average distance between interchain contact points) provides a short wavelength cutoff ($\sim 30-50\text{Å}$) which suppresses more modes as L_e increases. For PEP, estimates of L_e vs. temperature are available from neutron spin-echo scattering experiments [2] which give $L_e = 35\text{Å}$ at room temperature and $L_e = 50\text{Å}$ at 300°C , a trend consistent with the x-ray scattering results. Similar x-ray experiments performed on PS over the same temperature range show very little change in the diffuse scattering. At this point, however, no direct experimental results are available for L_e of PS.

Thin PEP films ($\sim 150\text{Å}$) were also observed to have strong temperature effects, but in addition showed very long relaxation times on the order of hours. Fig. 2 shows diffuse scans for a thin PEP film of molecular weight 290K . The original room temperature spectrum shows a large amount of diffuse scattering relative to thicker films while the high temperature point is similar to the thick film data. However, upon cooling to room temperature, the original spectrum is not recovered, even after one day. These long relaxation times reflect the strong polymer-substrate interactions which make the dynamics sluggish.

Model calculations are now being developed to include the elastic effects on correlations. This is simplest for polymer solutions where the surface wave spectrum is still mostly capillary like. Such calculations give good agreement with experiment for PBdS/toluene solutions and are now being extended to the more difficult case of melts.

[1] Sinha et al. Phys. Rev. B38, 2297 (1988).

[2] Butera et al. Phys. Rev. Lett. 66, 2088 (1991).

This work is supported in part by the NSF (DMR-8921556), DOE (DE-FG02-90ER45437) and the Petroleum Research Fund.

LOCAL STRUCTURAL DETERMINATIONS USING DIFFRACTION ANOMALOUS FINE STRUCTURE (DAFS) MEASUREMENTS

J.O. Cross, H. Stragier and L.B. Sorensen (U. of Washington)
C.E. Bouldin and J. Woicik (NIST)

We have demonstrated a new x-ray spectroscopy, Diffraction Anomalous Fine Structure (DAFS), for obtaining the short-range order information usually obtained from X-Ray Absorption Fine Structure (XAFS). DAFS is compared with XAFS below; (A) and (B) are the DAFS intensities versus the incident photon energy for the Cu (111) and Cu (222) Bragg reflections from a 2000 Å thick Cu film grown epitaxially on mica and (C) shows the corresponding Cu XAFS spectrum.

The DAFS shown contains both the real (f_1) and imaginary (f_2) components of the anomalous scattering amplitude, as well as a small contribution from the absorption in the Cu film. The real and imaginary components of the scattering amplitude are related by causality, resulting in a $\cos(2kr + \delta(k))$ dependence of the f_1 fine structure and a $\sin(2kr + \delta(k))$ dependence of the f_2 fine structure, as observed in XAFS. For a thin, mosaic, monoatomic crystalline Cu film, the diffraction intensity is $I = |f|^2(1 - \exp(-\mu t))$, where $f = f_0 + f_1 + if_2$ is the full atomic scattering amplitude, t is the film thickness, and μ is the x-ray absorption coefficient. The full anomalous scattering amplitude, $f_1 + if_2$, is the sum of the usual free atom scattering amplitude, $f_0 + if_2^0$, and the fine structure correction due to the scattering atoms, $f_2(\chi_1 + i\chi_2)$; χ_1 has the $\cos(2kr + \delta(k))$ dependence and χ_2 the $\sin(2kr + \delta(k))$ dependence mentioned above.

Because the DAFS and XAFS differ only in phase, the Fourier transform magnitudes of the DAFS and XAFS should be identical. The Fourier magnitudes obtained from (A), (B) and (C) (after background removal) show that the spectral components are identical.

These measurements show that DAFS measurements can be used to provide local structural information identical to that obtained from XAFS.

ATOMIC EXCHANGE AT SEMICONDUCTOR SUPERLATTICE INTERFACES

K. M. Kemner, B. A. Bunker, N. Samarth, H. Luo, J. K. Furdyna (Notre Dame)

XAFS has been used to investigate atomic ordering effects in the $\text{Ga}_{1-x}\text{In}_x\text{As}_y\text{Sb}_{1-y}$ ($y=0.05$ and 0.10) bulk crystal system. Results from data taken at the As edge and Ga edges showed a Ga-As and In-Sb pairing preference implying that this is a chemically-driven system as opposed to one driven by strain. The results imply the system is exhibiting either incipient ordering or micro-clustering.¹

To extend this work, we have used polarization-dependent grazing-incidence XAFS with electron-yield detection to study semiconductor alloy superlattices. The particular samples used for this study are two different $\text{ZnTe} / \text{CdSe}$ (001) superlattices, with 250 periods of $2/2$ layers (i.e. $\text{Zn-Te-Zn-Te-Cd-Se-Cd-Se}$) and $5/5$ layers. In this preliminary work, we find Zn-Se and Cd-Te coordination numbers increased above that expected for sharp interfaces in the superlattice. In this system, strain would appear to suppress interdiffusion; these results are therefore reminiscent of the chemically-driven quaternary III-V system mentioned above.

We are currently working on a detailed comparison between our XAFS results (polarization-dependent coordination numbers, distances, and Debye-Waller factors) and valence-force-field strain-minimization calculations using a supercell model.

Se edge, measured at room temperature, in a $2/2$ ZnSe/CdTe superlattice (filled circles), ZnSe (+), and CdSe (x).

¹ S. I. Islam and B. A. Bunker, Physics Letters **156**, 247 (1991).

We gratefully acknowledge discussions with M. Weidmann and K.E. Newman. This work is supported in part by the US Office of Naval Research under contracts N00014-89-J-1108 and N00014-90-J-1782. The X23-A2 beamline at Brookhaven National Laboratory is supported by the National Institute of Standards and Technology.

Total Electron Yield EXAFS at Low Temperatures

Kenneth M. Kemner, Z. Wang, R. Mayanovic, B. A. Bunker, University of Notre Dame.

Besides obtaining EXAFS data in the fluorescence mode, Total Electron Yield (TEY), using a He-gas-filled detector, has been shown to have advantages for EXAFS data acquisition for some single crystal and thin film samples. This is primarily due to the decrease in number and amplitude of the Bragg diffraction peaks from the single crystal sample or substrate when using TEY as compared to fluorescence measurements. Coupled with this is a desire to perform EXAFS experiments at low temperatures (around 77 K) so as to reduce the thermal vibrations contributing to the EXAFS Debye-Waller factor.

We have solved these two problems by designing a free-flowing He-gas-filled TEY detector that operates close to the liquid nitrogen temperature (89 ± 2 K). The detector consists of an approximately 1/2 inch thick 2 1/2 inch radius aluminum disk with 1/16 inch thick, 1/2 inch high plexiglass sides and a 1/16 inch plexiglass lid all used to create a disc of space approximately 1/2 inch high and 2 1/2 inches radius where the sample sits. He gas travels through a Cu tube submerged in liquid nitrogen and then into this chamber to aid in the cooling of the sample as well as an amplifier of the electrons emanating from the sample (connected to the negative terminal of a 66.7 volt battery) and traveling up to an aluminum collector plate (connected to a Keithley current amplifier). To further lower the temperature of the sample, Cu cold fingers were attached underneath the aluminum base. The detector can then be lowered into a liquid nitrogen bath up to the middle of the aluminum base.

We performed Transmission EXAFS at 77K on an Au foil (6.8 mg/cm^2) as well as a polycrystalline GaAs sample (400 mesh, 4 layers). We then performed low angle (5 to 9 degrees) TEY EXAFS measurements on the same Au foil and a single crystal GaAs wafer. A minimum of two scans were performed at a minimum of three different angles perpendicular to the azimuthal axis so as to differentiate between the EXAFS oscillations and the small amplitude Bragg diffraction peaks which were deglitched. One scan from each orientation was converted into chi space and then merged, and all subsequent analysis of the Transmission and TEY data was performed in an identical fashion (i.e. same window widths, weighting factors etc.). After running a ratio between Transmission and TEY envelope data, assuming the Transmission data to be correct, coordination numbers for 1st, 2nd and 3rd shells were found to be off by no more than 2 percent. Debye-Waller factors were always within $0.0004 \text{ \AA}^2$, and radial distances agreed to better than $0.004 \text{ \AA}$ for 1st shell data and $0.013 \text{ \AA}$ for 2nd shell data.

Fig. 1 TEY (solid line) and Transmission (dotted line) chi data for GaAs at 89K. TEY spectrum is sum of four scans.

Fig. 2 TEY (solid line) and Transmission (dotted line) chi data for Au at 89K. TEY spectrum is sum of three scans.

STUDY OF PHASE TRANSITIONS IN POTASSIUM NIOBATE

K. H. Kim (NRL-NRC), W. T. Elam (NRL) and M. I. Bell (NRL)

Niobium K-edge EXAFS spectra of potassium niobate (KNbO_3) have been measured across several phase transitions using the transmission method. A small furnace was used to raise the sample temperature and a closed cycle helium refrigerator was used for cooling.

KNbO_3 has the perovskite structure and shows four phases: cubic, tetragonal, orthorhombic and rhombohedral. Previous EXAFS study of this material showed that the phase transitions show characteristics of order-disorder behavior. [1] To study the nature of phase transitions more carefully, we took data around each transition temperature in small steps.

Standard EXAFS analysis methods were used to get chi data and r-space Fourier transforms (FT) for data around the highest temperature, cubic - tetragonal transition. The figure below shows the FT at temperatures above and below the transition temperature. In the FT, the region corresponding to Nb-Nb distance was carefully observed, and it showed no sudden change across the transition temperature. This is somewhat surprising in view of the order-disorder type transition for the material. One possible explanation comes from the possibility of correlation between neighboring Nb atoms. [2]

We have made theoretical calculations for this region in FT according to the order-disorder type structure, with and without correlations between the neighboring Nb atoms. As this region in FT includes multiple scattering paths involving oxygen atoms between the Nb atoms, rigorous theoretical calculation is difficult. We made some qualitative comparisons, however, including both single and multiple scatterings, and the results show that if there were no correlation, the change in the FT would be greater than the experimental data show.

Data around other transition temperatures will be analyzed in the same way.

[1] K. H. Kim, W. T. Elam and E. F. Skelton, *Mat. Res. Soc. Symp.* **172**, 291 (1989)

[2] M. I. Bell, K. H. Kim, and W. T. Elam, *Ferroelectrics* **120**, 103 (1991)

Figure: Magnitude of Fourier transform of KNbO_3 below and above the cubic-tetragonal transition temperature.

CHARACTERIZATION OF A $\text{Cu}/\text{Al}_2\text{O}_3$ BURIED INTERFACE USING GLANCING-ANGLE X-RAY REFLECTIVITY AND XAFS STUDIES

R. A. Mayanovic, B. A. Bunker, C. Zhong, and A. Anagnostopoulos, Physics Dept., U. of Notre Dame.

Characterization and local structure studies of $\text{Cu}/\text{Al}_2\text{O}_3$ buried interfaces are important in relation to electronic studies of tunnel barrier junctions. These studies are also significant within the context of our polarization-dependent studies on atomic thermal-motion disorder at solid-solid interfaces. The sample was fabricated by evaporating $\sim 4000\text{\AA}$ of Cu on a 5.1 cm Si wafer substrate inside an evacuated belljar at $\sim 10^{-6}$ torr. The sample was then quickly transferred to another high vacuum chamber where $\sim 700\text{\AA}$ of Al_2O_3 was RF sputtered over the Cu film. Glancing-angle x-ray reflectivity and XAFS measurements were made on the sample at room temperature. The reflectivity measurements were made at 8.7 and 9.2 KeV and the Cu K-edge XAFS measurements were made for several angles, ranging from 4-6 mrad. The sample was studied under two orientations: Having the x-ray polarization vector ϵ either parallel or perpendicular to the $\text{Cu}/\text{Al}_2\text{O}_3$ interface. Fig. 1 shows x-ray reflectivity data measured at 8.7 KeV with ϵ parallel to the interface. The data was reasonably well fitted using the Fresnel equation modified for interface roughness, as described by Heald et al.¹ Several models for the layered sample were tried during the fitting, including one having an intermediate CuAl-CuO alloyed layer sandwiched between the Cu and Al_2O_3 layers: Best results were obtained by assuming a sharp $\text{Cu}/\text{Al}_2\text{O}_3$ interface. This is verified by preliminary analysis of the polarization-dependent EXAFS data. The value used for the Al_2O_3 surface r.m.s. roughness was 42Å. The slight discrepancy between the fit and the data, especially near 4 mrad, can be explained in terms of long range distortions of the sample, which were unaccounted for during fitting, originating with the Si wafer substrate. Taking this into account, we estimate the true r.m.s. $\text{Cu}/\text{Al}_2\text{O}_3$ interface roughness to be between 30-40Å.

Fig. 1 X-ray reflectivity data (dots) measured at room temperature, 8.7 KeV, and having ϵ parallel to the interface. The fit to the data (line) was obtained using a modified Fresnel equation.

I. S.M. Heald, H. Chen, and J.M. Tranquada, Phys. Rev. B 38, 1016 (1988)

Supported by Office of Naval Research under contract N00014-89-J-1108.

GLANCING-ANGLE XAFS STUDIES OF PREMELTING AT $\text{Cu}/\text{Al}_2\text{O}_3$ INTERFACIAL REGIONS

R. A. Mayanovic, B. A. Bunker, and N. Jisrawi, Physics Dept., U. of Notre Dame.

Studies on metals have shown that premelting at the surface occurs considerably below the bulk melting temperature, indicating that the wet surface of a solid may act as a nucleation source during melting.¹ Our aim in this work is to study the atomic thermal-motion disorder and vacancy formation associated with the precursors to melting at metal-insulator interfaces using XAFS. Cu K-edge XAFS spectra were taken on $\text{Cu}/\text{Al}_2\text{O}_3$ multilayers, grown using RF sputtering on optically flat Si wafers having 20 layers each of $\sim 100\text{\AA}$ of Cu and $\sim 100\text{\AA}$ of Al_2O_3 , using the glancing-angle of incidence technique in the fluorescence mode. Measurements were taken at several angles of incidence, ranging from ~ 3 to 13 mrad, at room temperature and $420 \pm 2^\circ\text{C}$. The multilayers were studied under two orientations: the x-ray polarization vector parallel (P orientation) and perpendicular (S orientation) to the $\text{Cu}/\text{Al}_2\text{O}_3$ interface. Analysis of the polarization-dependent EXAFS data reveals a directional anisotropy in the EXAFS Debye-Waller factor, occurring both at room temperature and in a much more pronounced fashion at $420 \pm 2^\circ\text{C}$, indicating that the amplitude of relative Cu atom in-plane motion is larger than for out-of-plane motion at the interface (see Fig. 1). As shown in Fig. 2, we have obtained preliminary results for the incident angle dependence of the relative in-plane EXAFS Debye-Waller factor, measured at $420 \pm 2^\circ\text{C}$, indicating a steady decrease of vibrational amplitudes of approximately 3-4% of the nearest-neighbor distance for Cu atoms down to a depth of about 30Å from the interface, indicates that we are close to the premelting regime. The implication of this study is that the interface very likely undergoes premelting via a shearing mechanism rather than the recoil mechanism used to explain premelting at a free surface, for which the directional anisotropy in atomic vibrational amplitudes is reversed. The shearing model may ultimately help to explain superheating of encapsulated solids.

Fig. 1 Normalized EXAFS oscillations for the Cu K-edge, measured at a $\text{Cu}/\text{Al}_2\text{O}_3$ interface, at 3.9 mrad incidence angle and $420 \pm 2^\circ\text{C}$, in S (solid line) and P (dashed line) orientations.

Fig. 2 The Cu atom in-plane EXAFS Debye-Waller factor (P orient.), as a function of incident angle, measured relative to the out-of-plane component (S orient.) obtained at 3.9 mrad.

I. J. W. M. Frenken and J. F. van der Veen, Phys. Rev. Lett. 54, 134 (1985); P. H. Fuoss, L. J. Norton, and S. Brennan, Phys. Rev. Lett. 60, 2046 (1988).

Supported by Office of Naval Research under contract N00014-89-J-1108.

EXAFS STUDY OF STRUCTURAL CHANGES OF THE Ni₃Al ORDERED INTERMETALLIC BY MECHANICAL MILLING

T. Nasu, A.M. Edwards*, D.E. Sayers* and C.C. Koch
 Department of Materials Science and Engineering, and
 *Department of Physics, NCSU, Raleigh, NC, 27695

X-ray absorption fine structure data from the Ni₃Al ordered intermetallic were taken at various stages of its structural evolution induced by mechanical milling.

Least-squares fitting results of the EXAFS data show that both the Ni-Ni and the Ni-Al atomic distances increase slightly with milling time. During the first 5 hours of milling, the Ni-Ni coordination decreased while the Ni-Al coordination increased. This observation is indicative of an order-disorder transition, which was confirmed by XRD and TEM. Subsequent milling produced a remarkable increase in the Ni-Ni coordination number indicative of some kind of Ni clustering. Simultaneously, the Ni-Al coordination number decreased markedly; the total Ni coordination always remaining roughly constant (see table below).

These EXAFS results suggest that segregation of Al from Ni₃Al occurred during the mechanical milling process and, eventually, a partially amorphous structure evolved. After the longest milling time, a Ni-rich nano-crystalline phase was surrounded by the Al-rich amorphous phase.

Milling time (hours)	N Ni-Ni	R Ni-Ni (Å)	N Ni-Al	R Ni-Al (Å)
0	8.0	2.47	4.0	2.47
2	7.4	2.48	4.6	2.49
5	7.0	2.48	4.8	2.49
23	10.4	2.49	1.5	2.49
50	11.0	2.49	0.8	2.49

T. Nasu and C.C. Koch were partly supported by the National Science Foundation under grant number DMR-8620394-02. A. M. Edwards and D. E. Sayers were supported by the Department of Energy under contract number DE-FG05-89ER45384.

Intermediate-Range-Order Around Copper in Crystalline and Amorphous Silicates using EXAFS

David A. McKown, NIST, Gaithersburg, MD 20899

Recent neutron scattering studies of Ca-silicate glasses have presented results that imply intermediate-range-ordering of the network modifying Ca environments, which link together to form sheets of edge sharing Ca-octahedra [1]. From EXAFS, ordering of network modifying Cu environments is found for an amorphous Cu-silicate. Chrysocolla, relative to a crystalline Cu-silicate of similar composition, diopase (Cu₂Si₂O₇·6H₂O). The crystal structure of diopase [2], the standard for this study, is comprised of isolated 6-membered rings that are "glued" together by Cu²⁺ and H₂O. The Cu is coordinated by four oxygens at bond distances that average 1.96 Å. Each CuO₄ unit links together with one CuO₄ unit by edge sharing (Cu-Cu at 2.99 Å) and with two other CuO₄ units by corner sharing (Cu-Cu at 3.17 Å). Since all oxygens in the CuO₄ units are shared with SiO₄ tetrahedra there are four Cu-Si distances that average 3.17 Å.

The second shell in the partial RDFs (near 2.80 Å in Fig. 1) shows significant increase in area for chrysocolla compared to diopase; it is unusual for an amorphous material to show this behavior compared with its crystalline counterpart. Fitting results for the backtransforms of the various shells indicate that there is little difference in the Cu-O (first shell) and Cu-Si (second shell) environments between diopase and chrysocolla. The Cu-Cu (second shell) pair correlations comparing diopase with chrysocolla indicate that the bond distance increases from 3.10 to 3.13 Å, the coordination number increases from 3.0 to 4.1, and the Debye-Waller factor decreases from 0.0100 to 0.0059 Å². A possible structural adjustment from diopase to chrysocolla is the replacement of one edge sharing CuO₄ unit, or the short Cu-Cu distance, with two corner sharing CuO₄ units, or two longer Cu-Cu distances, so that each CuO₄ unit shares corners with four other CuO₄ units and all Cu-Cu distances are similar. These results indicate that crystallographic constraints which impose two sets of Cu-Cu distances in diopase can be relaxed in chrysocolla to produce one set of longer Cu-Cu distances; this gives the impression that the second shell in chrysocolla is more ordered than in diopase.

References

- [1] P.H. Gaskell, M.C. Eckersley, A.C. Barnes, P. Chieux, *Nature*, **350**, 675-677 (1991).
- [2] K.-H. Breuer, W. Eysel, R. Muller, *Zeits. Krist.*, **187**, 15-23 (1989).

Fig. 1. Fourier transform of the diopase and chrysocolla $k^2\chi(k)$ data.

ORIENTATION-DEPENDENT XAFS STUDIES OF HIGH-T_c SUPERCONDUCTING Pb-Bi-Sr-Ca-Cu-O SYSTEM.

W.F. Peng,¹ P.K. Tseng,¹¹ J.B. Shi,¹² H.C. Ku,¹³ R.A. Mayanovic,¹⁴ B.A. Bunker,¹⁵ Phys. Dept. Tamkang U., Natl. Taiwan U.,¹¹ Natl. Chiao Tung U.,¹² Natl. Tsing Hua U.,¹³ U. of Notre Dame,¹⁴

We have used polarization-dependent XAFS to study the high-T_c superconductors (Bi_{1-x}Pb_x)₂Sr_{1.5}Ca_{0.5}O_{10-δ}. Bulk samples were ground into powders which were subsequently uniaxially oriented in a epoxy-embedded solid solution using a 9.4T Tesla magnetic field at room temperature. X-ray diffraction measurements indicate that the uniaxial orientation direction for the particles is coincident with the crystallographic c axis; the a and b axes for each particle are randomly oriented with respect to one another.

We have measured the Bi, Pb L_{2,3}-edge and Sr K-edge polarization-dependent XAFS spectra at room temperature and Cu K-edge spectra at temperatures ranging from near 10K to room temperature using a Displex cryostat. Preliminary results show that Pb, Bi L_{2,3}-edge and Sr K-edge XAFS spectra do not vary substantially for different sample orientation with respect to the x-ray polarization vector E. In addition, we infer that Pb ions substitute for Bi ions in the Bi-Sr-Ca-Cu-O system due to the fact that there is no significant difference between the Pb and Bi L_{2,3}-edge XAFS spectra, respectively. The Cu K-edge XAFS spectra, measured at several orientations of the (Bi_{1-x}Pb_x)₂Sr_{1.5}Ca_{0.5}O_{10-δ} sample's c axis relative to E at room temperature, are shown in Fig. 1: An orientational dependence of the XAFS structure is observed in this case revealing a pronounced directional anisotropy of the electronic structure at the Cu site. It is evident from Fig. 1 that the first shoulder along the XANES edge for the (Bi_{1-x}Pb_x)₂Sr_{1.5}Ca_{0.5}O_{10-δ} sample becomes most prominent when c is parallel to E and least prominent when c is perpendicular to E or when the sample is unoriented. We believe that this feature can be attributed to dipole-allowed direct transitions of the photoelectron to Cu 4p-like final states since these are split into out-of-plane (4p_z) (along c axis) and in-plane (4p_x) states by crystal-field effects. Further polarization-dependent XAFS studies of these high-T_c superconductors versus temperature are in progress.

Fig. 1. Normalized Cu K near-edge absorption for (Bi_{1-x}Pb_x)₂Sr_{1.5}Ca_{0.5}O_{10-δ} samples. θ is the angle between polarization vector E and c axis. (a) Cu₂O (b) CuO (c) θ=0° (d) θ=45° (e) θ=90° (f) unoriented sample.

* Supported by the National Science Council (Taiwan,ROC) under grand No. NSC81-0208-N-032-13.

X-ray Absorption Investigation of Amorphous Platinum Pyrimidine Derivatives

R. Serimaa, V. Eteläniemi, T. Laitalainen*, T. Paakkari and I. Mutilainen*
University of Helsinki, Department of Physics, Helsinki, Finland
* University of Helsinki, Department of Chemistry, Helsinki, Finland

A series of amorphous platinum complexes were formed in a reaction of cis-diaquodiamineplatinum sulphate with pyrimidine nucleosides. These complexes have shown promising antitumor activity. We have investigated uridine complexes Pt-green, Pt-violet, Pt-blue and Pt-black synthesized in different conditions. Wide angle x-ray scattering (WAXS) measurements have been carried out using conventional x-ray sources to characterize average radial atomic distribution function. Further investigations by transmission EXAFS were done to determine the local structure of Pt atoms surrounded by O, N and Pt atoms. Fourier transform of the EXAFS oscillation (see figure, uncorrected for phase shift) shows differences in the coordination number of the first coordination cell (Pt-O, Pt-N). Platinum atoms are expected to form single chains and the Pt-Pt distance is about 3 Å in a view of the WAXS results. Both EXAFS and WAXS studies show clearly higher Pt-peak in the case of Pt-violet than in the other cases indicating longer chains. Further analysis is in progress.

Research supported by Neste Foundation, by the Technology Development Centre and by the Academy of Finland, Finland.

SITE SELECTIVE LOCAL STRUCTURE INFORMATION FOR YBaCuO USING DIFFRACTION ANOMALOUS FINE STRUCTURE

H. Stragier, C. Bitz, D. Yee and L.B. Sorensen (U. of Washington)
C.E. Bouldin, J. Woick (NIST)
J.M. Tranquada (BNL)
K. Ludwig (Boston U.)

We have used Diffraction Anomalous Fine Structure (DAFS) to determine the local structure of the two copper sites in $\text{YBa}_2\text{Cu}_3\text{O}_7$. By measuring DAFS for different Bragg reflections, we obtained DAFS spectra with different weights for each of the two copper sites. This allows the local structure to be determined for each site.

We measured the Cu K edge DAFS for a $\text{YBa}_2\text{Cu}_3\text{O}_7$ thin film grown on a LaAlO_3 substrate with the c axis parallel to the surface normal. A two-circle spectrometer with a vertical scattering plane was used to minimize polarization corrections. We measured the DAFS for the (003), (005), (006), and (007) Bragg reflections; the figures below show the Fourier transforms. The differences between the DAFS spectra for these Bragg reflections are dramatic, especially for the higher shells which contain contributions of the Y, Ba, and Cu neighbors. The copper to Y, Ba and Cu coordinations are quite different for the two copper sites as reflected in the Fourier transforms shown below. The relative contributions from each copper site to the (00L) reflection intensities have been calculated from known atomic positions; they are given by $\text{Ncos}(Q_z Z)$, yielding: (003) $\text{Cu}(1)=1.0$, $\text{Cu}(2)=1.84$; (005) $\text{Cu}(1)=1.0$, $\text{Cu}(2)=0.31$; (006) $\text{Cu}(1)=1.0$, $\text{Cu}(2)=1.37$; (007) $\text{Cu}(1)=1.0$, $\text{Cu}(2)=-2.0$.

These preliminary results show that it is feasible to use DAFS to obtain site selective local structural information in the 123 superconductors.

EXAFS STUDY OF SPONTANEOUS ORDERING IN $\text{Ga}_{0.5}\text{In}_{0.5}\text{P}$ THIN FILMS DEPOSITED ON $\text{GaAs}^{\dagger}$

Anne Tabor-Morris, Kenneth Kemner, Robert Mayanovic, and Bruce Bunker, University of Notre Dame

Alloys deposited in thin films using epitaxial growth techniques have shown a spontaneous ordering and a lowering in band gap under certain growth conditions.¹ One such thin film is $\text{Ga}_{0.5}\text{In}_{0.5}\text{P}$ deposited 2 degrees off of the (100) axis of the GaAs substrate by metal organic chemical vapor deposition.¹ Electron and x-ray diffraction studies have indicated the existence of long range ordering.² Short range structures and interatomic distances are studied here using polarized X-ray beam EXAFS at the Ga K-edge. A liquid nitrogen cooled total electron yield detector produces data in which the Ga nearest neighbors are specifically probed to see if the type of atom in the second shell is preferentially ordered along axes of the $\text{Ga}_{0.5}\text{In}_{0.5}\text{P}$ single crystal overlayer.

As illustrated in Figure 1, there is a marked difference in the amplitude and phase of the Fourier transform of the EXAFS chi data in higher shell region, depending upon the orientation of the sample. This difference in the second and third shells is thought to be attributable to Ga preferentially ordering along the (1,1,0) direction and In along the (1,-1,0) direction, consistent with the evidence of ordering found in the other studies. Further analysis is necessary and currently in progress.

Figure 1. Total Electron Yield Data: Fourier transforms, k^3 weighted. There are two scans shown for each polarization. Solid line: 0 degrees (x-ray beam polarization along (1,1,0) axis). Dashed line: 90 degrees (sample rotated, polarization along (1,-1,0) axis).

1. Wei, S.-H., Zunger, A., Appl. Phys. Lett. **56** (7) 662 (1990).
2. Kurtz, S., Olsen, J., Kibbler, A., Appl. Phys. Lett. **57** (18) 1922 (1990).
3. Kurtz, S., Olsen, J., Kibbler, A., Appl. Phys. Lett. **54** (8) 718 (1989).

* Supported by the Office Of Naval Research under contract N00014-89-J-1108.

† Samples provided by K. Bertness, National Renewable Energy Laboratory, Golden, CO.

EXAFS STUDY OF GOLD-INDUCED GERMANIUM CRYSTALLIZATION

Zhengquan Tan and S. M. Heald (Brookhaven Nat'l Lab);

C. E. Bouldin and J. C. Woicik (Nat'l Inst. of Standards and Technology)

When in contact with metals, amorphous silicon (*a*-Si) and germanium (*a*-Ge) crystallize at temperatures significantly lower than in their bulk form. In order to understand the role of the contact metal, we have studied the crystallization process of *a*-Ge/Au bilayers and co-evaporated *a*-Ge_{1-x}Au_x (with nominal *x*=0.02 and 0.12) films by measuring the Ge and Au EXAFS. The amorphous Ge shows only the nearest neighbor contribution in the EXAFS, as it crystallizes, distant neighbors appear (Fig. 1a). Upon isothermal annealing for 10 minutes, the Ge/Au bilayer crystallizes at 170 °C, and in the meantime, a large amount of Au appears in the top layer, very likely in the grain boundaries. For co-evaporated films, the *x*=0.12 sample crystallizes at 130 °C and the *x*=0.02 sample at 330 °C. A pure *a*-Ge film does not crystallize at 330 °C under similar conditions. In the *x*=0.02 and 0.12 *a*-Ge_{1-x}Au_x films, each Au atom on the average is bound with one to two Ge atoms at 2.40 Å, but with no Au atom. As the films crystallize, Au transforms into fcc gold in the *x*=0.12 sample but remains bound with Ge in the *x*=0.02 sample (Fig. 1b). Although the Au solubility in Ge is extremely low (0.01%) in equilibrium, our results suggest that Au-Ge bonding can be significant in the *a*-Ge and may play an important role in lowering the Ge crystallization temperature.

*Work supported by the U. S. Department of Energy under Contract Nos. DE-AC02-76CH00016.

Fig. 1 Fourier-transform magnitudes of (a) Ge K-edge EXAFS, and (b) Au L₃-edge EXAFS for *a*-Ge_{1-x}Au_x films. The spectrum of bulk gold is included for comparison.

EXAFS STUDY OF FERROELECTRIC PbS₂Te_x ALLOYS

Zhihai Wang and Bruce A. Bunker, University of Notre Dame.

Ferroelectric phase transitions (FPT) induced by off-center ions have attracted considerable attention in the past decade because of the general interest in disordered system and cooperative phenomena. Highly polarizable crystals containing off-center impurities, such as IV-VI semiconductors, are particularly interesting. Continuing our EXAFS investigation on local structure in ferroelectric IV-VI alloys, we have extended to PbS₂Te_x system. Previous studies were concentrated on cation-substituted solid solutions, for example Pb_{1-x}Ge_xTe, in which at least one of the end-member binary compounds is ferroelectric. Here we focus on an anion-substituted system, in which both two end-member binary components are paraelectric. The anomalous properties of this alloy system, which include the temperature dependence of resistivity, strong nonlinearity of concentration dependence of T_c and exhibition of the "quasi-Kondo" effect, suggest that there is a second order phase transition and the Sulfur atoms play an important role in the transition. EXAFS studies on the Pb site in these alloys have been performed above and below the transition temperatures on three samples, *x*=0.10, *x*=0.18 and *x*=0.30. We found a single Pb-Te distance but two distinct S distances separated by approximately 0.2 Å. The bond lengths do not depend on temperature if normal thermal expansions taken into consideration. Figure 1 summarizes the first shell Pb-Te and Pb-S bond length information for three samples. The EXAFS results imply that S atoms are displaced about 0.6 Å from the cell center at all temperatures and concentrations in both the low- and high-temperature phases. These findings suggest that the S atoms may undergo an order-disorder transition as observed in Pb_{1-x}Ge_xTe. By comparing EXAFS Debye-Waller (DW) factors of Pb-Te bonds of the alloys with that of pure PbTe measured at the same temperature, we can get additional structural information. The results for three samples are shown in Figure 2. We found that the relative DW factors increase with decreasing of temperature in low temperature phase. We believe this is due to the relative shift between the Pb and Te sublattices causing a splitting of Pb-Te distance in the alloys. This indicates another character of this phase transition — dispersive character.

Fig. 1 The first shell Pb-Te and Pb-S bond lengths as function of S concentration. Note the two distinct S distances.

Fig. 2 The difference in first-shell EXAFS Debye-Waller factors $\Delta\sigma^2$ associated with the Pb edge, relative to PbTe measured at the same temperature, as a function of temperature T.

I. Q. T. Islam and B. A. Bunker, Phys. Rev. Lett. 59, 2701 (1987).
Supported by Office of Naval Research under contract N00014-89-J-1108

EXAFS STUDY OF THE FERROELECTRIC $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ ALLOYS

Z. Wang, R. Mayanovic, B. Bunker, U. Debska and J. Furdyna, U. of Notre Dame.

Weil et al.¹ reported a ferroelectric transition in the $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ alloy system, for $x=0.04$ to 0.40 , based on the temperature variation of the dielectric constant and from the hysteresis loop at elevated temperatures. From their work, however, it is unclear what is happening *structurally* through this transition and what mechanism may be driving the transition. Since there is a large ionic size mismatch between Zn and Cd cations (Zn^{2+} : $0.83\text{\AA}$, Cd^{2+} : $1.03\text{\AA}$), it is possible that Zn ions may be off-center along the symmetric axis of and inside the Te tetrahedral cage. If the ions are indeed off-center, the alloy may undergo an orientational order-disorder ferroelectric phase transition (FPT) as found in $\text{Pb}_x\text{Ge}_{1-x}\text{Te}^2$ and $\text{PbS}_x\text{Te}_{1-x}^3$.

We reported previously⁴ our Zn K-edge EXAFS results for the powder $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ samples. In that study, we found a $0.0007\text{\AA}$ difference of the first shell EXAFS Debye-Waller factor between the alloys and pure ZnTe, which means the Zn-Te bonds are slightly distorted. Assuming Zn ions are inside an undistorted Te tetrahedral cage, this implies that Zn ions may be off-center by less than $0.1\text{\AA}$. In order to test that this is caused by the off-center displacement of Zn ions, we have performed a polarization-dependent XAFS investigation on E-field polarized single crystal samples at the Zn K-edge. Bridgman-method grown single crystals of $\text{Zn}_x\text{Cd}_{1-x}\text{Te}$ ($x=0.10$ and 0.20), were cut and polished to a $1\text{mm} \times 10\text{mm} \times 10\text{mm}$ plate, having the $\langle 110 \rangle$ direction normal to the sample surface and the $\langle 111 \rangle$ direction parallel to one of the sample edges. The EXAFS measurements were performed in fluorescence mode with the polarization vector of x-rays either parallel or perpendicular to the $\langle 111 \rangle$ sample direction, along which a 5 to 15 Volt potential was applied, and the sample tilted about 5 degrees relative to the beam. In this way, if the Zn ions are off-center, they can be uniaxially oriented along the $\langle 111 \rangle$ direction by the external field, in order for us to see a difference in the EXAFS spectra between parallel and perpendicular measurements. Fig. 1 shows the Zn K-edge χ data for $x=0.10$ sample at room temperature. Because of unwanted diffraction peaks, it was necessary to rotate the sample slightly from one position to another, resulting in the two sets of data being slightly different. This difference is comparable to the possible effect caused by the off-center Zn ions. Thus no conclusive result about the Zn ions being off-center along the $\langle 111 \rangle$ direction can be deduced from the present experiment. Further studies involving modulation experiments are in progress.

Fig. 1 Zn K-edge χ data of $\text{Zn}_{0.10}\text{Cd}_{0.90}\text{Te}$ at $\langle 111 \rangle$ parallel (solid line) and perpendicular (dashed line) to the polarization of x-rays at 300K.

1. R. Weil, R. Nkum, E. Muranevich, and L. Benguigui, Phys. Rev. Lett. 62, 2774 (1989).
2. Q. T. Islam and B. A. Bunker, Phys. Rev. Lett. 59, 2701 (1987).
3. B. A. Bunker, Z. Wang and Q. Islam, Proc. of XAFS-VI, York, England, Aug. 1990.
4. Z. Wang, R. Mayanovic, and B. Bunker, Bull. Amer. Phys. Soc. 36, No.3 762 (1991).

Supported by Office of Naval Research under contract N00014-89-J-1108

IN-SITU OBSERVATION OF DEFECT FORMATION IN CdTe CRYSTALS AT HIGH TEMPERATURES DUE TO IMPOSED MECHANICAL STRESSES

R. Balasubramanian (Clarkson Univ.) and W. R. Wilcox (Clarkson Univ.)

In the last two decades, virtually every known method of crystal growth has been used to grow high quality CdTe crystals. No technique has yet produced such crystals. For use in device fabrication or as substrates for epitaxial growth, CdTe single crystals still have to be "mined" from larger polycrystalline ingots. Further, the crystals obtained currently have a high incidence of defects like twins, low-angle grain boundaries, dislocations and precipitates.

Thermal and mechanical stresses experienced by an ingot during the growth process are thought to be important factors in defect formation and multiplication. Therefore, in order to improve the techniques of crystal growth of CdTe , it is necessary to expand our understanding of the mechanism of defect formation and apply this knowledge in the control of the crystal growth process. The real-time, diffraction imaging of the process of defect formation, propagation and multiplication is being carried out at beamline X23A3. We have constructed an apparatus for in-situ imaging of crystals at high temperatures (1250 K) and under loads of up to 7 kilograms. Experiments have been carried out successfully with single crystal samples of silicon. The study of defect formation and multiplication on cadmium telluride is planned for the near future.

This work is being supported by the Center for Commercial Crystal Growth in Space, a NASA CCDS.

David R. Black and Harold E. Burdette

In collaboration with Maurice Landstrass at Crystallume Inc. we have used high resolution diffraction imaging to examine a homoepitaxial diamond film and substrate. The sample was prepared by CVD growth of approximately 120 μm of diamond film on a natural 1x1x0.1 mm type IIa substrate. Diffraction images were recorded for the following 5 orientations of the sample: (111) and (220) asymmetric transmission, (220) and (400) symmetric transmission and (400) symmetric surface. In three of these orientations a weak secondary image was observed, an example of a pair of images is shown in figure 1. The misorientation between the two images is about 0.2 degrees, which is the same misorientation seen between the seed and new growth in high temperature/high pressure diamond growth. Since the substrate crystal was not available for examination prior to growth, the assignment of these two images to substrate and film requires an examination of the image quality when the diffracted beam exits the film side or the substrate side, as shown in figure 2. Our preliminary conclusion is that the substrate is the primary diffraction and the film is the weak secondary diffraction. The narrow diffracted area from the film indicates that it is probably highly strained.

Figure 1. (111) asymmetric transmission topographs taken at 8 keV showing the primary diffraction on the left and a weak secondary diffraction on the right. The diffracted beam exits from the substrate side.

Figure 2. (040) symmetric transmission topographs taken at 8 keV. In this case the beam exits from the film side. Compare the image crispness to figure 1.

David R. Black and Harold E. Burdette

In collaboration with scientists from General Electric, the defect microstructure of single crystal man-made diamond is being investigated by high resolution x-ray diffraction imaging. Until recently, the highest quality diamonds were considered to be natural type IIa (nitrogen concentration < 50 ppm) crystals. Since the growth conditions for these diamonds are unknown and uncontrolled, each crystal is an individual with a unique set of defects and properties. The high temperature/high pressure process patented by GE has been developed to the point that man made type IIa diamonds can be grown which are superior to natural crystals. The correlation between growth conditions and crystalline perfection is being studied to take advantage of this opportunity to grow reproducible high quality diamond. The change in crystal perfection as a function of growth is shown in figure 1. An isotopically pure ^{12}C crystal was sectioned perpendicular to the growth direction and each section examined. It was found that the new growth was initially highly defected, figure 1a. As the crystal grew the center section was almost perfect, figure 1b-c and then deteriorated near the termination of growth, figure 1d. It was also discovered that the new growth is misoriented from the seed crystal by about 0.2°. Whether this misorientation is required for growth to occur is currently being investigated.

Figure 1. (111) asymmetric transmission topographs taken at 8 keV from four slices of a man made diamond crystal.

DIFFRACTION IMAGING OF ISOTOPICALLY PURE MAN-MADE DIAMOND

D. E. Black (NIST) and H. E. Burdette (NIST),

X-ray diffraction imaging has been used to compare the defect structure and overall crystalline perfection of a sample of isotopically pure ^{12}C man-made diamond and a similar sample containing the naturally occurring isotopic concentration of ^{13}C . These samples were supplied by Bill Hanholzer of G.E.'s Corporate Research Center and were grown by the temperature gradient method invented by G.E. The samples are thin sections, approximately 0.6 - 0.7 mm thick, cut perpendicular to the growth direction from larger stones that are typically in excess of 1 carat. The x-ray topographs were taken using 8 keV x-rays in transmission through the samples.

Figure 1a shows the sample with natural isotope abundance and figure 1b the isotopically pure sample. The high concentration of defects radiating from a central point seen in figure 1a is typical of man-made type IIa diamonds. The lack of defect images in the central region of figure 1b shows that the isotopically pure sample is a more perfect crystal. The randomly oriented (non-crystallographic) dark lines are residual damage from polishing. Whether the observed improved crystal perfection is a result of isotopic purity alone cannot be determined from an examination of one sample. The relationship between crystal perfection, isotope concentration and other growth conditions is presently being investigated.

Figure 1. X-ray topographs of a man-made diamond with natural isotope abundance in (a) and isotopically pure in (b). Figure 1a is a (111) diffraction and figure 1b is a (111) diffraction (bold indicates negative index).

ABSOLUTE CALIBRATION OF SMALL-ANGLE X-RAY SCATTERING

P. R. Jemian (ANL), G. G. Long (NIST), S. Krueger (NIST) and D. R. Black (NIST)

Absolute calibration of small-angle scattering data is a prerequisite to the derivation of most of the physically relevant microstructural parameters available from such experiments. Examples of the parameters which can be derived include the total volume fraction of particles or pores, their number distribution, and their total surface scattering area. Sensitive modeling of the scatterers also requires absolute calibrated data.

One of a widely-used set of calibration standards used at the Oak Ridge National Laboratory small-angle scattering facility is a collection of aluminum single crystals with small percentages (0.4 - 0.8%) of voids induced by neutron irradiation. At low scattering vector ($h < 0.014\text{\AA}^{-1}$), the scattering is isotropic and obeys the Guinier formula. The radius of gyration R_g has been determined to be 203 - 210 Å. Since the NIST SAXS facility on X23A3 is self-calibrated by means of a primary calibration technique, a measurement of the Oak Ridge sample designated "Al-8" was made for verification of consistency.

SAXS measurements on Al-8 were performed using 8 keV and 10 keV incident photons. The scattering curves are shown below in Figure 1 and the volume distribution of diameters, determined using a maximum entropy analysis, is shown in Figure 2. The agreement with the Oak Ridge analysis of the scattering by this sample is excellent.

The authors acknowledge with gratitude support for this work from the U.S. Army Research Office under contract no. MPR-ARO 102-90 and from the U.S. Department of Energy under contract no. DE-FG02-86 ER-5229.

MORPHOLOGY OF L-ARGININE-PHOSPHATE CRYSTALS FOR NONLINEAR OPTICAL DEVICES

G. G. Long (NIST), D. R. Black and H. E. Burdette

L-arginine-phosphate (LAP) is a new nonlinear optical crystal which shows promise for future applications in optical devices. The crystals for this study were grown from aqueous solution by a temperature lowering technique in a reciprocating crystallizer by S. M. Rao, A. K. Batra, C. Cao and R. B. Lal at the Alabama A&M University. Prior to the x-ray topography study reported here, defects in the LAP crystals were investigated by the growers using an etch pit technique to reveal dislocations and using birefringency interferometry. The effect of a high dislocation density region is to distort the fringes when viewed in the birefringency interferometer.

Five (001) crystal slices from two boules were investigated by monochromatic x-ray topography. Defects observed included slip lines, regions of single dislocations, liquid phase inclusions, and microcracking. Since the time of this study, the growth geometry was modified so that liquid phase inclusions have been eliminated and microcracking is much reduced.

Fig. 1 is an (002) reflection topograph showing regions of good crystal perfection and single dislocations. These results have been correlated directly with the etch pit results. Fig. 2 is an (020) transmission topograph showing slip line defects and liquid phase inclusions.

Fig. 1

Fig. 2

X-RAY DIFFRACTION IMAGING OF L-ARGININE PHOSPHATE (LAP)

G. G. Long (NIST), D. R. Black (NIST), H. E. Burdette (NIST), S. M. Rao (Alabama A&M) and R. B. Lal (Alabama A&M)

L-arginine phosphate is a new nonlinear optical crystal belonging to the monoclinic system with space group $P2_1$ and with two formula units per cell. Crystals grown from aqueous solutions exhibit pinacoidal habit with a prominent basal pinacoid. In the past, etch figures have been made on different naturally growing pinacoidal faces of LAP crystals. Such studies have revealed the symmetry of different faces, and different dislocation densities depending on the location of the crystal slice relative to the seed.

In the present study, monochromatic x-ray diffraction imaging was carried out on LAP crystals in an attempt to measure the dislocation density and to observe the growth habit of LAP. Figure 1 shows a (002) reflection topograph of a slice of a solution-grown LAP crystal containing the seed. Individual images of dislocations near the surface can be seen, as well as indications of the growth habit.

This work is supported by the Office for the Commercialization of Space, NASA, CCGS/Clarkson University.

This work was supported by the Center for Commercial Crystal Growth in Space, Clarkson University, NY

STRUCTURE OF MICROPOROUS FOAMS

C. G. Long (NIST), S. Krueger (NIST), B. J. Olivier (SNL), R. Lagasse (SNL) and D. W. Schaefer (SNL)

Sandia National Laboratories has developed several classes of microporous foams. Many of these foams have pore sizes less than 10 μm and very high surface areas (10-500 m^2/g). These foams are produced by several processes mostly involving phase separation. The structure of these foams depends on a variety of factors including gelation, thermal history, polymer tacticity and solvent thermodynamics. The purpose of the small-angle x-ray scattering measurements is to relate the structure of these foams to processing parameters and establish the essential parameters that control foam structure.

Previous small-angle scattering has generally shown near power-law scattering profiles with slopes near -4 in the Porod regime, indicative of smooth pore boundaries. One exception to this rule was found for deuterated isotactic polystyrene for which an intermediate power-law was found with a slope of approximately -2. These experiments indicated that the pore geometry could only be established by means of exceedingly high-resolution Bonse-Hart-type measurements.

The SAXS instrument on the X23A3 beamline was used to perform feasibility studies on a pyrolyzed ($\sim 1000^\circ\text{C}$) polyacrylonitrile foam. Pore anisotropy is known to exist in this system. The approximate foam composition was measured to be: 95% carbon, 4% nitrogen and 1% oxygen. The thickness was 2.1 mm and the density was 43.2 mg cm^{-3} . The sample exhibited gradations in color with an approximate 4 mm periodicity. SAXS measurements on dark and light regions of the sample indicated different amounts of broadening in the central portion of the SAXS scan near h ($= 4\pi \sin \theta / \lambda$) = 0. Further it was found that the broadening was wavelength dependent. This made it possible to explore a multiple-small-angle x-ray scattering analysis of the data where the size region probed by the MSAXS formalism is 0.1-10 μm . Application of the multiple scattering formalism indicated that there are different average radii of the pores in this foam in the light and the dark regions.

HIGH-RESOLUTION SAXS STUDY OF THE STRUCTURE OF MICROPOROUS FOAMS

B. J. Olivier (Sandia Nat. Labs.), D. W. Schaefer (Sandia Nat. Labs.), S. Krueger (NIST), and G. Long (NIST)

Sandia National Laboratories has developed several classes of microporous foams. Many of these foams have exceedingly small pore sizes ($< 10 \mu\text{m}$) and very high surface areas (10-500 m^2/g). These foams are produced by several processes mostly involving phase separation. The structure of these foams depends on a variety of factors including gelation, thermal history, polymer tacticity and solvent thermodynamics. It is our goal to relate the structure of these foams to processing parameters and establish the essential parameters that control foam structure.

For many of these foams, previous small-angle scattering studies ($Q \geq 0.005 \text{ \AA}^{-1}$) have shown near power-law scattering profiles with slopes near -4, indicative of smooth pore boundaries. On occasion, however, there has been an exception to this rule where an intermediate power-law regime was found with a slope of about -2. This observation was attributed to sheet-like lamellae.

The foams currently under investigation were prepared by gel and solvent crystallization methods. Several of the small-angle scattering spectra show novel features for $Q < 0.001 \text{ \AA}^{-1}$, indicative of an additional length scale of the order of $\sim 1 \mu\text{m}$. Fig. 1 shows one of the scattering spectra, which has been placed on an absolute scale and desmeared. This spectrum is remarkable in that it displays a slope of about -3 over three decades in scattering vector. At very small Q ($\sim 10^{-4}$), many of the foams also display noticeable broadening as a function of wavelength. The data are currently being analyzed further to determine if the Berk formalism (1) for multiple small-angle scattering by neutrons (2) carries over into the x-ray regime.

Figure 1. SAX spectrum, plotted on a log-log scale, of one of the foams.

1. N. F. Berk and K. A. Hardman-Rhyme, *J. Appl. Cryst.* **18**, (1985) 467-472 and *J. Appl. Cryst.* **21**, (1988) 645-651.
2. S. Krueger, G. G. Long and R. A. Page, *Acta Cryst.* **A47**, (1991) 282-290.

This work was supported by the U.S. DoE under contract number DE-AC04-76DP00789.

Work performed at Sandia National Laboratories was supported by the U.S. Department of Energy under contract number DE-AC04-76DP00789.

Images, in the form of photographs, video tape records, and digital electronic outputs from a CCD camera computer, for selected Bragg reflections, were obtained in transmission using 11.4 keV monochromatic radiation. This study serves to deduce the history of growth, impurity intake and fracture of a Zambian α -quartz single crystal in a general context of metallogenic and mineral deposition processes. At first, translation— as well as section—topographies, in O—beam and several complementary H—beam settings— covering the entire, 1mm thick, crystal plate, were recorded on film. The images show the partitioning of the crystal section, cut in {101} orientation, into domains of differing and variable diffraction contrast. Most conspicuous are parts of the crystal associated with repeated optical (Brazil-law) and electrical (Dauphine-law) twinning. The individually varying contributions of each part of the crystal to the integrated intensity determine the spread $\Delta\theta_0$ of reflections about the precise Bragg angle θ_0 . These variations were determined in detail by a series of photographic O— and H—beam images, taken in discrete steps of 0.003 – 0.006° through $\Delta\theta_0$ for 2020, 3031 and 1120 reflection settings. The successive distributions of changing and vanishing contrasts, contributed to the integrated intensity by different parts of the crystal, map out its 3-dimensional lattice strains ($\Delta d/d_{hkl}$) and/or rotations $\delta\theta_{hkl}$. Enhanced reflectivity, evidently related to growth sectors, is found to originate from optical twin lamellae, dispersed impurity distributions, and large particulate impurity inclusions, aligned, some with preferred orientation, along apparent, probably healed, fracture surfaces. The high resolution x-ray microscope (designed and devised by Kunizuma, Dobbyn and Spal, 1990, at N.I.S.T.) was used next to obtain images of 0.16 – 0.25 mm² areas, with 20x magnification, at 27 closely neighbouring sites along a path within 3mm radius from a conspicuous crack bifurcation. During the 1st run (in May), with the crystal set for a 3031 reflection, photographic and digital records were taken only of the O—beam images. The digital records lend themselves supremely to quantitative topography and analysis by derived image representations. These include contour and vector maps, lattice strains and stresses, at all or specified levels, as well as their derivatives and resolved components in specific crystallographic directions. The strain fields of an arrested crack tip and of a large, crystallographically bounded, fluid inclusion (negative crystal) with a bubble inside it, have been of particular interest. Therefore most of a 2nd run (in July) was used to investigate more closely with the diffraction microscope at 20x magnification, the lattice strains associated with the negative crystal and those due to apparently relaxed crack tip stresses. The O—beam and 3031 H—beam images of the 0.5mm wide field containing the negative crystal were sampled in 0.002° steps through a double diffraction peak of -0.0175° half-width. Peak doubling and vanishing contrasts for the inclusion at one position within $\Delta\theta_0$ confirm its negative crystal habit. A similar series, through a sharp single peak, of the O— and H—beam images was obtained for the crack tip strains.

During a few remaining hours of beamtime, an experiment of real time, video topography of a TGS-crystal undergoing the ferro—paraelectric transition with heating, was run with white beam radiation. A transition was observed but the crystal became severely radiation—damaged.

Support of this work by the Nat'l Science & Engin'g Research Council Canada, the National Inst. of Science & Technology, and B.N.L. is gratefully acknowledged.

In previous radiographic studies using the asymmetric Bragg diffraction microscope, anomalously high contrast was sometimes observed in regions a few microns wide along object edges. The effect was surmised to be Fresnel diffraction, but the experimental conditions were unfavorable for quantitative analysis. Here, the effect is observed at the edge of a thin film, and quantitatively demonstrated to be phase contrast Fresnel diffraction. The phase contrast qualifier is crucial, because absorption contrast produces a much smaller effect. This study establishes that the spatial resolution of the microscope is limited by Fresnel diffraction.

The film was a $0.15\mu\text{m}$ thick deposit of Pd covering a $100 \times 100\mu\text{m}^2$ area on a 15 mil thick Si wafer. The deposited side of the wafer was optically polished, but not the other side, unfortunately (see below). A film edge was examined under the microscope, using 80X magnification, 1 Å radiation, and 500 sec exposure. Fig. 1 shows the image intensity, normalized to its value on the wafer's bare surface, as a function of distance from the film edge, with $- (+)$ distance representing the interior (exterior) of the film. Each point represents integrated intensity on an $88 \times 0.25\mu\text{m}^2$ strip parallel to the film edge. The absorption contrast is the difference between asymptotic values on the left and right. The anomalous contrast is the amount by which the difference between peak values exceeds the absorption contrast.

The curve in Fig. 1 is the calculated Fresnel diffraction pattern of the film edge, with no adjustable parameters. The anomalous contrast depends on the phase shift introduced by the film, which was 7.4° . Larger shifts could give much larger contrast, and zero shift gives negligible anomalous contrast. Because the edge was oriented vertically, the diffraction pattern depends on the horizontal source size. Its previously measured 2σ value of 0.84 mm [R. Spal, 4/89, unpublished] was used. Compared to results for a point source, the anomalous contrast is reduced by a factor of 4.3, and secondary Fresnel fringes are absent. Finally, the scatter of points about the curve is due to wafer surface roughness, not counting statistics.

Fig. 1. Fresnel diffraction pattern of thin film edge.

X23A3
THE EFFECT OF MICROGRAVITY ON THE MESOSCOPIC STRUCTURE OF LEAD TIN TELLURIDE:
HIGH RESOLUTION MONOCHROMATIC DIFFRACTION IMAGING OF CRYSTALS GROWN IN
MICROGRAVITY ON SPACE SHUTTLE STS 61A AND ON THE GROUND

Bruce Steiner, Ronald C. Dobbryn, David Black, Harold Burdette, Masao Kuriyama,
Richard Spal (NIST), Richard Simchick,* and Archibald Fripp (NASA Langley
Research Center)

High resolution x-ray diffraction images of two directly comparable crystals of lead tin telluride, one Bridgman-grown on Space Shuttle STS 61A and the other terrestrially Bridgman-grown under similar conditions from similar material, present similar grain structure, but different subgrain structure. In the terrestrial sample, the appearance of an elaborate array of subgrains is closely associated with the apparent intrusion of one or more additional phases that are out of diffraction in all of the various images (Figure 1). The formation of this elaborate subgrain structure with affiliated intruding phases is inhibited by growth in microgravity (Figure 2). The intrusion of a second phase is predicted by the full phase diagram for this material,¹ in spite of the complete miscibility along the lead-tin axis.² Suppression in microgravity of the formation of the second phase may be associated with the absence of thermosolutal instability present in full gravity.

Figure 1. High resolution (220) diffraction image of terrestrial lead tin telluride crystal.

Figure 2. High resolution (220) diffraction image of lead tin telluride crystal grown on STS 61A.

- * Lockheed Engineering and Sciences Co., Hampton
1 J. F. Butler and T. C. Harman, "Metallic Inclusions and Cellular Substructure in $Pb_{1-x}Sn_xTe$ Single Crystals," J. Electrochem. Soc. 116, 260-263 (1969).
2 T. C. Harman, "Control of Imperfections in Crystals of $Pb_{1-x}Sn_xTe$, $Pb_{1-x}Sn_xSe$, and $PbS_{1-x}Se_x$," J. Nonmetals 1, 183-194 (1973).

This work was supported by the Microgravity Sciences & Applications Division of NASA

X23A3
THE MESOSCOPIC STRUCTURAL TRANSITION FROM CZOCHRALSKI TO BRIDGMAN GROWTH OF
GALLIUM ARSENIDE: HIGH RESOLUTION MONOCHROMATIC DIFFRACTION IMAGING

Bruce Steiner, Ronald C. Dobbryn, David Black, Harold Burdette, Masao Kuriyama,
Richard Spal (NIST), David Matthiesen, and Brian Ditchek (GTE Laboratories)

The transition from the Czochralski-grown seed to Bridgman growth has been observed by high resolution monochromatic diffraction imaging (Figures 1 & 2). The demarcation of the Czochralski seed from the new Bridgman growth is delineated in two ways. First, toward the periphery of the boule the lattice on the two sides of the boundary diffracts at slightly different angles. Either the lattice constant, or orientation, or both differs in the new growth. Second, the mesoscopic structure of the growth is observed to be transformed at the interface. The cellular structure of the seed is characteristic of diffraction images of Czochralski-grown undoped gallium arsenide. In the new Bridgman growth, the formation of cells is completely suppressed. Nevertheless, the two lattices mate without visible interruption. However, lattice mismatch appears to set up a gradual warping of the crystal lattice.

Figure 1. High resolution (220) diffraction image of seed - new growth interface region in lightly selenium-doped gallium arsenide crystal.

Figure 2. Enlargement of region in a high resolution (220) diffraction image of the seed - new growth interface region in lightly selenium-doped gallium arsenide crystal.

This work was supported by the Microgravity Sciences & Applications Division of NASA.

X23A3
THE MESOSCOPIC STRUCTURE OF MERCURIC IODIDE CRYSTALS GROWN IN SPACELAB III AND
ON THE GROUND: HIGH RESOLUTION MONOCHROMATIC DIFFRACTION IMAGING

Bruce Steiner, Ronald C. Dobbyn, David Black, Harold Burdette, Masao Kuriyama,
Richard Spal (NIST) and Lodewijk van den Berg (EC&G Energy Measurements)

High resolution monochromatic diffraction images of three mercuric iodide crystals show mesoscopic structural differences that correlate with growth conditions, chemical analysis, and device performance. Features that appear to be an impurity phase or phases (non diffracting inclusions) are found throughout crystals terrestrially grown from material similar to that used in Spacelab III (Figure 1). However, the formation of these features is suppressed both by growth in microgravity (Figure 2) and by terrestrial growth from higher purity material. The performance of radiation detectors made from these various materials, moreover, is correlated with the appearance of the multiple phase features. Given generally high quality crystals, the attainment of the maximum charge carrier mobility and lifetime appears to be substantially more dependent on the absence of mesoscopic (i.e. 1-100 micrometer) impurity multiphase regions than on macroscopic (i.e. >1 millimeter) lattice regularity within a few arc seconds over a major fraction of the crystal.

Figure 1. High resolution (0 1 1) diffraction image of terrestrial mercuric iodide crystal.

Figure 2. High resolution (1 0 10) diffraction image of Spacelab III mercuric iodide crystal.

This work was supported by the Microgravity Sciences & Applications Division of NASA

X23A3
STRUCTURAL CONTINUITY IN TRIGLYCINE SULFATE CRYSTALS GROWN IN SPACELAB III: HIGH
RESOLUTION MONOCHROMATIC DIFFRACTION IMAGING

Bruce Steiner, Ronald C. Dobbyn, David Black, Harold Burdette, Masao Kuriyama,
Richard Spal (NIST), Ravindra B. Lal, and Ashok Batra (Alabama A&M University)

A normal slice from a disc-shaped terrestrial triglycine sulfate seed crystal with additional growth on Spacelab III diffracts into images that appear in closely spaced groups as the crystal is rotated about its [100] and [001] axes. The appearance of images in groups at similar diffraction angles indicates that this crystal consists of layered grains whose lattices are similar but rotated with respect to one another. Since each image is nearly "complete," the grain boundaries are roughly parallel to the (010) crystal surface. Space growth occurred in the [001] direction along that one edge of the seed. The absence of a clear demarcation between the seed and new growth in most of the images contrasts with the terrestrial growth of comparable material. However, this interface is clearly visible in Figure 1, as defects in the seed grain appear not to propagate into the new growth in the central portion of the disc. Toward the edge of the disc, the irregularity observed in 2.1 of the grains is consistent with rapid growth anticipated from the defects observed near the seed edge.

Figure 1. High-resolution (200) diffraction image in Laue geometry from grain B1 of TGS crystal from Spacelab III.

Figure 2. High-resolution (001) diffraction image in Laue geometry from grain A3 of TGS crystal from Spacelab III.

This work was supported by the Microgravity Sciences and Applications Division of NASA

Ag $L_{2,3}$ EDGE XANES STUDY OF CHARGE REDISTRIBUTION AT THE Ag SITE IN Au-Ag ALLOYS

A. Bzowski, Y.M. Yiu and T.K. Sham (U. Western Ontario)

We have carried out X-ray absorption measurements at the Ag $L_{2,3}$ edge of a series of Au_x-Ag_{1-x} alloys (x varies from .75 to .25) using a total electron yield method in which the sample is in a He atmosphere. This work complements earlier Au $L_{2,3}$ edge results.

Fig. 1 L_3 XANES of Ag (top) and Au (bottom) in alloy compared with pure metal.

Although, it is well known that Au loses d and gains s-conduction charge when alloyed with Ag. The situation at the Ag site, is not clear. We want to determine whether or not Ag gains d charge locally upon alloying using Ag $L_{2,3}$ XANES which is sensitive to the d hole population above the Fermi level (dipole selection rule). Fig. 1 shows the comparison of the Ag and Au L_3 XANES of the 50-50 alloy with that of the pure metal. It is apparent from fig. 1 that while the Ag L_3 edge of Ag metal has stronger absorption than that of the alloy just above the threshold, Au metal absorbs less than the alloy. This observation indicates that Ag gains d character (loses d holes) upon compound formation. A trend is found among all the alloys and in the L_2 edge spectra.

Work supported by OCMR and Au and Ag (top) compared with pure metal. NSERC of Canada

STRUCTURE IN MULTILAYER SYSTEMS

C.L. Foiles, J.A. Coven, D. Koppel and H.-K. Sung (Michigan State U.), T.I. Morrison (IIT)

Multilayers with a repeating bilayer unit composed of Fe and a non-magnetic spacer (Fe/X) have quite different magnetic properties as the thicknesses of the bilayer decreases. The importance of structure for these behaviors is being investigated in three systems -- Fe/Si, Fe/V and Fe/Mo.

For Fe/Si with the Si layer fixed at 50 Å, ferromagnetism becomes weaker and disappears when the Fe layers are < 10 Å. Reflection X-ray diffraction data are broadened by the small thickness of the Fe layer and can only confirm that layering has occurred. Trends in transmission electron diffraction (TED) data suggest the Fe becomes amorphous but a persistent sharpness of the <110> Bragg peak raises some questions. Earlier EXAFS studies answered these questions; Fe layers of < 14 Å are amorphous and layers of 20 Å are largely amorphous [1].

In Fe/V the decrease in saturation magnetization for Fe and V layers of equal but decreasing thicknesses is gradual and the loss of magnetism for one monolayer (ML) of Fe is a matter of debate. TED data indicate an in-plane structural transition to a single BCC lattice with no apparent effect upon magnetization [2]. EXAFS data for both Fe and V indicate crystalline environments down to 2 ML's. The differences in backscattering are small and we are attempting to develop a self-consistent model using the EXAFS data for both elements.

For Fe/Mo the in-plane transition to a single BCC lattice and the loss of ferromagnetism occur near the same thickness. The raw EXAFS data for Fe in samples down to 4 ML/4 ML thicknesses have crystalline features and give evidence of an increasing importance of Mo backscattering in the Fe signal. A sequence of samples where the Fe layer is fixed at 3 ML and the Mo layer increases in thickness produce similar data. However, Fourier transforms of these data have a dominant first peak and an irregular pattern of secondary peaks. This result suggests the presence of a large, static disorder. Measurements below room temperature and subsequent use of a quantitative analysis which includes such large disorder are planned.

- [1] C.L. Foiles, Metall. Trans. (in press); C.L. Foiles, T.I. Morrison, H. Chen and H.-K. Sung in preparation (1991).
- [2] C.L. Foiles, Ultramicro. 29, 80 (1989).

Supported in part by MSU-Center for Fundamental Materials Research and by the Research Corporation.

EXAFS STUDIES OF STRUCTURAL ANISOTROPY IN Fe-Tb AMORPHOUS FILMS

V.G. Harris*, K.D. Aylesworth*, N.C. Koon, and W.T. Elam, Naval Research Laboratory, Washington, D.C. 20375

EXAFS spectra were collected for four Fe-Tb amorphous films in both the as-deposited state and after various heat treatments. Spectra, encompassing both the Fe and Tb absorption edges, were acquired with the sample positioned (i) normal to the incidence photons, and (ii) at a 150 glancing angle to the incident photons. Because of the polarization nature of synchrotron radiation these sample orientations allow for the sampling of element-specific atomic structure parallel to the film plane and perpendicular to the film plane, respectively. Magnetic characterization using a vibrating sample magnetometer (VSM) show the three alloys richest in Tb to have strong magnetic anisotropy fields aligned perpendicular to the film plane increasing in magnitude with Tb content. The fourth alloy, most dilute in Tb, was found to display nearly isotropic magnetization curves with respect to field orientation indicative of a formation of a second phase. Fourier transformed EXAFS data of both the Fe and Tb absorption edges illustrate an anisotropy in the data between sample orientations with respect to incident photons. This anisotropy is defined as a change in peak-to-peak amplitude in k-space, and a significant change in amplitude of the NN Fourier peak in r-space. The magnitude of this EXAFS anisotropy correlates closely with the measured magnetic anisotropy strongly suggesting that this structural anisotropy is the dominant contributing mechanism to magnetic anisotropy in these alloys. If this indeed is the case, low temperature heat treatments should be effective in reducing both the measured magnetic anisotropy and the detected anisotropy in the EXAFS data. The Fourier transform of the Fe EXAFS data is presented for the Fe₇₄Tb₂₆ alloy in Figures

(a), (b) and (c) for the as-deposited state and after low temperature heat treatments. Room temperature VSM measurements indicate the magnetic anisotropy has decreased significantly in the annealed samples. Figure (a) illustrates the Fourier transformed data of this alloy in its as-deposited state for both sample orientations. Clearly an anisotropy exists in the data. Figures (b) and (c) illustrate similar data acquired of samples after a 200°C and 300°C (t=1 hour) heat treatment, respectively. These transformed data illustrate significant reduction in the structural anisotropy observed in the as-deposited sample. In addition, the anisotropy seen in (b), over the range 2.5-4.0 Å and the shift in the nearest neighbor peak distance, is nearly eliminated after the heat treatment at 300°C (see (c)). These results illustrate unequivocally the existence of a structural anisotropy between the in-plane and out-of-plane structure in these alloy films. In addition, heat treatment trends with magnetic anisotropy strongly suggest that this structural anisotropy is the dominant mechanism contributing to the magnetic anisotropy in non-S-state rare earth-transition metal alloys.

* Supported by the National Research Council Fellowship Program

EXAFS STUDIES OF AMORPHOUS-TO-CRYSTALLINE TRANSITION IN Co₇₀Fe₃₀P₁₀Si₁ THIN FILM SPECIMENS

V.G. Harris*, S.A. Oliver, P. Dorsey and C. Victoria, Northeastern University, Boston, MA 02115
K.H. Kim and W.T. Elam, U.S. Naval Research Laboratory, Washington, D.C. 20375-5000

The well documented magnetic, mechanical, electrical and electrochemical properties of transition-metal-based metallic glasses often provide them advantages over polycrystalline alloys for many applications. Our interest in these materials stems from their potential application in thin film microwave devices.

Our previous studies of these alloys have shown they possess a metastable amorphous structure in the as-deposited state. Heat treatment procedures have proven to be effective in stabilizing this phase for application purposes. However, we found cobalt-rich alloys of this system to experience a unique sensitivity to these procedures, often indicating evidence of magnetic nonuniformities in their FM spectra. To investigate the origins of this behavior several specimens cut from a single cobalt-rich alloy film were subjected to various heat treatment procedures and subsequent EXAFS analysis.

A 150-nm-thick film was sputter-deposited from a Co₇₀Fe₃₀P₁₀Si₁ pressed powder target onto rotating quartz substrates. Approximately 10 samples were cored from the resultant film and subjected to magnetic field annealing procedures at 500 Oe. A field is applied parallel to the film plane at temperatures from 125 °C to 400 °C. The x-ray absorption spectra detailing the k-edge extended fine structure for transition metal constituents were collected on X23B using a conversion electron XAFS analysis chamber. Following standard XAFS analysis procedures the energy above the edge was isolated and a pre-edge linear background removed. The data was then further smoothed by cubic spline fitting and converted to wave vector space. Fourier transforms of this data illustrate the local atomic environments about the Co ion are presented in Figure 1 (transforms have been normalized and vertically stepped to allow direct comparison). A transform of Fe ion XAFS data for iron is presented for qualitative comparison. Figure 1 illustrates the onset of long range order in films annealed as low as 300°C. This is unexpected in this alloy whose crystallization temperature has been widely reported near 400-425°C. In addition, Fourier peaks appearing >3 Å, arising from this ordering, closely resembles the bcc signature profile of the iron.

EXAFS modeling, using FEFF version 3.0 FORTRAN code, allowed for quantitative analysis of this evolution. Results of these procedures indicate a low temperature clustering of boron about the transition metal ions. It is believed that a selected removal of cobalt from the amorphous matrix facilitates an acceptable ratio of Fe₂ to Fe₃ allow for the formation of bcc solid solution. Similar studies of an iron-rich alloy EXAFS data reveal similar behavior. However, the formation of the solid solution (bcc Fe-Ni in this case) did not occur until near the expected crystallization temperature (~425°C). The premature formation of the Fe-C solid solution is believed to be the origin of the detected magnetic inhomogeneities in cobalt-rich alloys.

Figure 1. Fourier transformed data of cobalt ion in specimens of alloy Co₇₀Fe₃₀P₁₀Si₁ (bcc iron is presented for comparison purposes only).

EXAFS STUDIES OF 18S Fe-Tb ALLOY FILMS

V.G. Harris, K.H. Kim and W.T. Elam
U.S. Naval Research Laboratory, Washington, D.C. 02375-5000

A series of Fe-Tb alloy films were ion beam sputter-deposited onto thin glass substrates to investigate relations between magnetic and structural properties as a function of heat treatment and composition. Presented here are preliminary findings of electron-yield EXAFS measurements and analyses performed on as-deposited Fe-Tb alloy films.

Room temperature EXAFS spectra were recorded on the NRL materials analysis beam line X23B using an electron-yield technique. Sample electron-yield currents, normalized to the incident photon flux, were recorded for x-ray energies extending 100 eV below the Fe K-edge to 600 eV above the Tb L_{III}-edge. In this manner, one scan was sufficient in recording XAFS data for both ions in the binary films.

Following standard EXAFS analysis procedures the spectrum above each edge was isolated and a linear background subtracted. The data were then converted to k-space and subjected to cubic-spline smoothing. Fourier transformation of the k-space data provided information of the central ion's local atomic structure. As the Fe data is limited to k_{max} 9.75 Å⁻¹ due to the presence of the Tb L_{III}-edge all transformed data has a k-space range of 2 to 9.75 Å⁻¹ (k²-weighted).

Presented in Figures 1(a) and (b) are transformed data of the Fe and Tb ions in FeTb alloys, respectively. The FeTb alloy compositions are approximated (a 5 at/o) by qualitative, sputter-yield to surface area calculations and ratioed absorption step heights to be 60Fe-40Tb (alloy 2), 70Fe-30Tb (alloy 3), 80Fe-20Tb (alloy 4) and 90Fe-10Tb (alloy 5). The Fourier transforms of Fe in these alloys (Fig. 1(a)) display significant nearest neighbor (NN) peak amplitude increases with increased Fe-content indicating that the NN shell is largely dominated by Fe-Fe correlations. For alloys 4 and 5, two smaller peaks resolved near 3 and 5 Å are observed, possibly indicating the existence of crystallites in these films. The transforms of XAFS data for Tb in alloys 2, 3, and 4 (Fig. 1(b)) display features near the NN Fourier peak. The relative amplitude changes with respect to alloy composition indicate the shoulders near 1.8 Å and 3.5 Å are dominated by Fe-Fe correlations. The central NN peak is also dominated by Fe-Fe correlation but may have contributions from Tb as well. The transform of alloy 5 has a large peak near 1 Å arising from low signal-to-noise in k-space data. The appearance of resolved peaks > 4 Å are suspect because of the low signal-to-noise but could arise from coherent photoelectron scattering in crystallites also observed in the Fe transforms.

Fig. 1 Fourier transforms of Fe K-edge XAFS data (a) and Tb L_{III}-edge XAFS data (b) in FeTb alloy films. All transforms were deduced from k-space data having k²-weighting and a range of 2 to 9.75 Å⁻¹.

Fe THICKNESS EFFECT ON STRUCTURE AND MAGNETISM IN LOW-DIMENSIONAL Fe/Tb MULTILAYERED STRUCTURES

V.G. Harris*, K.H. Kim*, N.C. Koon, and W.T. Elam, Naval Research Laboratory, Washington, D.C. 20375-5000

R. Coehoorn and W. Hoving, Philips Research Laboratories, Eindhoven, The Netherlands

X-ray absorption spectra encompassing both the Fe K and the Tb L_{III} absorption edges were collected using a conversion-electron technique on the Naval Research Laboratory's materials analysis beamline X23B. Following established EXAFS procedural steps the extended fine structure (EFS) was isolated and normalized to the edge step height, a non-oscillatory background was removed by fitting to a cubic spline, and the data was converted to wave vector space. These data in turn were Fourier transformed to provide direct information of the atomic environment about the central atom. Although electron phase shifts have not been accounted for in this stage of the analysis transform comparisons can be made to discern trends and relative changes in structure. Figure 1 are the Fourier transformed data of the Fe EXAFS acquired from normal incidence absorption spectra having the radiation's electric field vector in the film plane. The Fourier transform of the Fe EXAFS of a bcc Fe film standard is presented for comparison purposes. All transforms have been calculated using a k space range of 2.8 to 9.75 Å⁻¹ (k²-weighting).

The Fourier transformed data of MLs having Fe thicknesses less than 15 Å display amorphous-like transform profiles. Specifically, they illustrate a broad, central nearest neighbor peak centered near 2 Å and very little structure at radial distances greater than 3 Å.

The transform corresponding to the ML having t_{Fe}=15 Å illustrates additional structure at higher radial distances in the form of a broad peak at 4-4.5 Å. In addition, the split peak seen in the amorphous spectra appears to merge to a single very broad peak over the range 3-4 Å. In comparison to the bcc Fe standard these characteristics provide a qualitative match to the signature peaks seen in the bcc profile. This evidence suggest that an abrupt structural transition takes place in samples having t_{Fe} between 12 Å and 15 Å. The bcc-like structure is more pronounced in samples having larger t_{Fe} with peaks shifting to radial distances closer to those of the bcc film standard and additional Fourier peaks appearing at higher radial distances.

Magnetic properties reflect this phase transition as the magnetic moment of Fe is measured to increase significantly from the amorphous-like state to the ordered state. In addition the measured Kerr rotation angle was found to experience a minima during this transition. The coercive field is measured to decrease with increased t_{Fe} due to increases in magnetic anisotropy fields. Diffraction peaks stemming from the silicon substrate and sample periodicity appear in the Tb EXAFS spectra making it difficult to provide corresponding data.

* Supported by the National Research Council Post-doctoral Fellowship program

Figure 1. Fourier transformed Fe EXAFS data collected for a series of Fe/Tb multilayered structures. Fourier transform of Fe EXAFS from a bcc Fe film standard is presented for comparison.

X-24A Experimental Techniques: Flux Measurements from Grating-Multilayer Monochromator Combinations for Low Energy Studies

T. Jach (NIST), S. Scuthworth (NIST), P. L. Cowan (ANL), R. C. C. Perera (LBL), and B. W. K. in (BNL)

In the interest of obtaining useful x-ray flux at lower x-ray energies, we have replaced the two crystals in our right-angle type monochromator¹ with a synthetic multilayer followed by an organic crystal. The multilayer is able to withstand exposure to the white beam and the organic crystal provides the energy resolution. The multilayer is composed of 100 bilayers of W and C, with a total d-spacing of 26.25Å. The organic crystal is RAP (Rubidium Acid Phthalate) and its (100) d-spacing of 13.06Å is matched to the second order reflection of the multilayer.

The combination has been evaluated in the energy range of 750–1500eV. The measurements were made in the standard X-24A geometry of a premirror, monochromator, and toroidal focussing mirror. Measurements were made using different detectors: a silicon PIN photodiode and a channel plate for analog current measurements, and a spiraltron for photon pulse counting.

Measurements made at 750eV by both the channel plate and the PIN photodiode obtained a flux at the sample of $1.6 \cdot 10^7$ photons/s. Fig. 1 shows the relative reflectivity of the Ni-coated mirrors scanned over an energy range which includes the Ni L-edges.

P. L. Cowan, J. B. Hastings, T. Jach, and J. P. Kirkland, Nucl. Inst. & Methods 208, 349 (1983).

Fig. 1. Relative reflectivity of toroidal mirror with Ni coating in the energy range 800–900 eV as measured with a PIN diode detector. The data are unnormalized.

"X-RAY ADSORPTION ISOTHERMS"

J.L. Jordan-Sweet (IBM), B.D. Swanson and L.B. Sorensen (Univ. of Washington)

We have demonstrated the feasibility of using the fluorescence yield to measure layer-by-layer adsorption of an inert gas on single crystal surfaces.

The traditional methods used to measure layer-by-layer deposition of gases onto substrates are volumetric and gravimetric adsorption isotherms. The number of gas atoms (or molecules) adsorbed is measured as a function of the equilibrium 3D vapor pressure in a closed cell. In the first case, this is done by pressure differences and in the second by changes in the adsorbate-substrate combined weight. Both techniques require high specific surface area substrates to obtain measurable signals. From a series of isotherms recorded at different temperatures, the 2D phase diagram of the adsorbate-substrate system can be obtained.

In our experiment we simultaneously recorded the volumetric isotherm of argon on graphite foam and measured the argon fluorescence yield (FY) for argon adsorbed on a single crystal of graphite in the same cell at 66 K. Figure 1 shows the first layer step as a vertical riser in the volumetric isotherm at ~0.01 Torr. Figure 2 shows the x-ray FY from the argon L edge at 2960 eV, for the argon adsorbed onto the single crystal. A linear contribution from the 3D gas has been subtracted. Note that the vertical riser can be seen at the same 3D pressure.

This demonstrates that x-ray fluorescence can be used to measure isotherms, and coverages, of adsorbed gases on single crystal surfaces where it would be impossible to measure conventional isotherms. This will be valuable for x-ray scattering studies of mono- and few-layer adsorbed films on single surfaces.

Figure 1:
Argon/Graphite Foam
Volumetric Isotherm

Figure 2:
Argon/Graphite(001)
Fluorescence Isotherm

X-RAY, SOFT X-RAY, AND VUV BEAM POSITION MONITOR*

B. A. Karlin
National Synchrotron Light Source, Brookhaven National Laboratory, Upton NY 11973

P. L. Cowan** and J. C. Woicik
National Institute of Standards and Technology, Gaithersburg MD 20899

ABSTRACT

An x-ray beam position monitor has been developed and is currently operational at the NIST beamline X-24A of the National Synchrotron Light Source, Upton, NY. It was developed to accurately measure the vertical position of the synchrotron beam and to allow precise positioning of beamline optics. It is an area type monitor positioned 5.6 meters from the Tangent Point and precedes all beamline instrumentation. Vertical beam position is given by the differential photoemission current from two grids mounted on a Macor frame. The monitor also senses the total photon beam flux.

* Work performed under the auspices of the U.S. Department of Energy, under contract DE-AC02-76CH00016.

** Current affiliation: Argonne National Laboratory, Argonne IL 60439

X-RAY STANDING-WAVE STUDY OF Sb ON InP(110), GaP(110) and GaAs(110)

T. Kendelewicz*, J. C. Woicik**, K. E. Miyano*, P. L. Cowan*, B. A. Karlin***, C. E. Bouldin**, P. Pianetta*, and W. E. Spicer* --*Stanford U; ** NIST; ***NLSLS

The soft X-ray standing-wave technique has been used to study ordered monolayers of Sb on InP(110), GaP(110) and GaAs(110). Using the back-reflection diffraction geometry from (220) planes, we determine the vertical distances of Sb atoms to the III-V semiconductor surface and compare this to theoretical calculations and elastic low energy electron diffraction determinations. We have also addressed the issue of the Sb overlayer buckling and show how this quantity scales with the covalent radii of substrate atoms. An example of the data for the Sb/InP(110) interface is shown below and the fit parameters are enclosed in the table.

ELDED++	THEORY++	XSW
D	Δ	D
1.11	.25	1.09
0.99	.75	1.12
		.58

+ C. Mailhot et al., PRB 31, 2213 (1985); ++ C. B. Duke et al. Surf. Sci. 163, 391 (1985)

J. C. Levin (U. Tennessee/ORNL), D. W. Lindle (NIST), N. Keller (UT/ORNL), R. D. Miller (UT/ORNL), Y. Azuma (ANL), N. Berrah Mansour (ANL), H. G. Berry (ANL), and I. A. Sellin (UT/ORNL)

We have measured the ratio of double-to-single photoionization of helium far above the double-ionization threshold using time-of-flight techniques during NSLS single-bunch operations. We find $\text{He}^{++}/\text{He}^+ = 1.6 \pm 0.3\%$ at $h\nu = 2.8 \text{ keV}$. This ratio, which has been the subject of many detailed studies near threshold (e.g. ref. 2), has not previously been measured accurately at photon energies above 200 eV. Helium is the simplest atom which exhibits electron-electron correlation. As a result, photoionization of He has been used as a testing ground for understanding correlation phenomena, such as autoionization, correlation-satellite production (shakeup) and double photoionization. Theoretical prediction of the energy dependence of the double photoionization of helium is a fundamental problem in atomic physics which requires solution of the Coulomb three-body problem. In the independent-particle framework, in which the photon can interact directly with only one electron, double photoionization can proceed only by electron-electron correlation. Therefore, the optical treatment of electron-electron correlation is of fundamental importance in calculations of double photoionization. Further, determination of the ratio of double-to-single photoionization of helium at high energy is of critical importance in clarifying the relationship of electron-electron correlation in double photoionization to that in double ionization by charged particles.

FIG. 1. Time-of-flight spectrum of helium following photoionization by 2.8 keV photons.

1. J. C. Levin, D. W. Lindle, N. Keller, R. D. Miller, Y. Azuma, N. Berrah Mansour, H. G. Berry, and I. A. Sellin, *Phys. Rev. Lett.* **67**, 968(1991).
2. H. Kossmann, V. Schmidt, and T. Andersen, *Phys. Rev. Lett.* **60**, 1266 (1988).

This work was supported in part by the NSF and by U. S. DOE, Division of Chemical Sciences, under Contract No. DE-AC05-84OR21400, with Martin Marietta Energy Systems, Inc., and by U. S. DOE under Contract No. W-31-109-ENG-38.

S. H. Southworth (NIST) and P. L. Cowan (ANL)

We have used a quadrant PIN silicon photodiode as a position monitor of the monochromatized x-ray beam in beamline X-24A [1]. The active area of the commercially available diode is divided into four electrically isolated quadrants. The diode was chosen with the additional feature of a 1.3 mm diameter hole in the center, which offers the opportunity to pass a focused beam through the detector and onto a sample, while intercepting the edges of the beam for position detection.

The detector was mounted in vacuum at the focal plane of the beamline. The center of a focused beam of 3640 eV x-rays was passed through the hole in the diode, while the edges of the beam were intercepted and produced photocurrent signals of approximately 10 μA per quadrant. These signals were processed externally using standard transresistance amplifiers, and the sum and difference signals were compared to determine the relative beam positions. Signal levels proportional to the horizontal and vertical beam positions and to the total intensity intercepted are derived simultaneously.

The detector was determined to be quite sensitive to the angular and translational motion of the x-ray beam arising from movement of beamline optical components such as the focusing mirror and double-crystal monochromator. This provides diagnostic information on the performance of the beamline optics and suggests that the quadrant detector may be useful as a component in feedback loops used to maintain beam position on target. A full report on this beam position monitor is given in Ref. [2].

[1] P. L. Cowan, S. Brennan, R. D. Deslattes, A. Henins, T. Jach, and E. G. Kessler, *Nucl. Instr. and Meth.* **A246** (1986) 154; P. L. Cowan, S. Brennan, T. Jach, D. W. Lindle, and B. A. Karlin, *Rev. Sci. Instr.* **60** (1989) 1603.

[2] S. H. Southworth and P. L. Cowan, Proceedings of the 7th National Conference on Synchrotron Radiation Instrumentation, Baton Rouge, Louisiana, 1991, to be published in *Nucl. Instr. and Meth.*

X-RAY STANDING-WAVE STUDY OF RARE GAS ATOMS PHYSORBED ON SINGLE CRYSTAL GRAPHITE

Brian D. Swanson, Cecilia Bitz, Dennis Yee, Larry B. Sorensen (U. of Washington), Jean Jordan-Sweet (IBM), Paul Cowan (NIST), and Barry Karlin (NSLS)

X-ray back-reflected standing wave (XBRSW) measurements were used to determine the adsorbate-substrate separation of rare gases physisorbed on graphite single crystals.

The experimental apparatus consisted of a sample cell with beryllium and optical windows connected to a Displex closed-cycle refrigerator, gas dosing tube, and diode temperature sensors. Inside the sample cell, the graphite crystal was mounted in a tweezer assembly which allowed the graphite crystal to be cleaned in situ by quick resistive heating to above 1000° C. The gas coverage was determined from simultaneous volumetric isotherms of the graphite foam ballast in the cell and by direct x-ray fluorescence measurements of the physisorbed gas on the graphite single crystal surface.

The fluorescence yields of krypton at 1585 eV and argon at 2960 eV were measured while scanning the incident x-ray energy through the back-reflected graphite (004) Bragg peak at 3697 eV. The figure below shows the graphite reflectivity and the monolayer krypton fluorescence yield at 26 K. A fit to the data indicates the krypton separation to be 0.67 (modulo 1) of the graphite (004) spacing (1.67 Å) with a coherent fraction of 0.72. Our results demonstrate that the XBRSW method can be used to accurately determine how the krypton-graphite separation changes with temperature and coverage e.g., whether the krypton layer becomes lower and flatter as the atoms go into the graphite wells during the incommensurate-commensurate transition.

Determination of the Sb/Si(111) interfacial structure by back reflection x-ray standing waves and surface extended x-ray absorption fine structure

J.C. Woick⁽¹⁾, T. Kendelewicz⁽²⁾, K.E. Miyano⁽²⁾, P.L. Cowan⁽¹⁾,
C.E. Bouldin⁽¹⁾, B.A. Karlin⁽³⁾, P. Pianetta⁽²⁾, and W.E. Spicer⁽²⁾

⁽¹⁾National Institute of Standards and Technology, Gaithersburg MD 20899.

⁽²⁾Stanford Electronics Laboratory, Stanford University, Stanford CA 94305.

⁽³⁾National Synchrotron Light Source, Brookhaven National Laboratory, Upton NY 11973.

Based on the surface extended x-ray absorption fine structure (SEXAFS) absolute surface coordination numbers, Woick et al. [Phys. Rev. B 43, 4331 (1991)] argued that Sb trimers adsorb on the Si(111) surface in a modified 'bridge' configuration rotated by 60° relative to the atop 'milk stool' geometry proposed by Abukawa et al. [Surf. Sci. Lett. 201, 513 (1988)]. Using the accurate near neighbor bond lengths as determined from SEXAFS, these models predict perpendicular distances of 1.76 and 2.60 Å between the top Si atoms and the Sb layer, respectively. New back reflection x-ray standing wave (BRXSW) data find the Sb atoms to reside 2.53 ± 0.1 Å above the Si surface, in clear support of the milk stool geometry. Polarization dependent SEXAFS relative coordination numbers support the milk stool model as well.

Top

Side

X-ray standing wave determination of the clean InP(110) surface reconstruction

J.C. Woick⁽¹⁾, T. Kendelewicz⁽²⁾, K.E. Miyano⁽²⁾, P.L. Cowan⁽¹⁾,
C.E. Bouldin⁽¹⁾, B.A. Karlin⁽³⁾, P. Pianetta⁽²⁾, and W.E. Spicer⁽²⁾

- 1) National Institute of Standards and Technology, Gaithersburg MD 20899
- 2) Stanford Electronics Laboratory, Stanford CA 94305
- 3) National Synchrotron Light Source, Brookhaven National Laboratory, Upton NY 11975

The x-ray standing wave (XSW) technique through detection of surface sensitive Auger electrons has been used to determine the surface reconstruction of clean InP(110). Using the back reflection diffraction geometry from (220) planes, we find the perpendicular displacements of the surface P and In atoms to be $+0.18 \pm 0.1$ Å and -0.48 ± 0.08 Å, respectively, from their unrelaxed bulk positions. The use of low energy x-rays combined with a novel method of data analysis opens the general problem of surface reconstruction to study by this technique.

EFFICIENCY OF SOFT X-RAY DETECTOR PHOSPHORS

E. L. Benitez, D. E. Husk, and S. E. Schnatterly (U. of Virginia)

Phosphors are used in detectors to convert soft x-rays to visible light. We have performed photoluminescent efficiency measurements in the 3-50 eV range on a variety of phosphors: $Y_2O_3:S:Eu$, $Y_2O_3:Eu$, $La_2O_3:Tb$, $Gd_2O_3:Tb$, $ZnO:Zn$, and $Sr_5Cl(PO_4)_3:Eu$.

We have modeled the phosphor efficiencies using a formula based on surface recombination of electron-hole pairs and inelastic electron scattering data for the absorption and reflection coefficients of the materials. In previously published papers^{1,2} we have fit the model to the measured efficiencies up to 160 eV. We have obtained excellent fits which provide values for the diffusion length, surface recombination velocity, and bulk quantum efficiencies of the materials. Fig. 1 shows the absorption coefficient of $La_2O_3:Tb$ as measured by inelastic electron scattering. Fig. 2 shows our previously published measured efficiency of the phosphor (squares) and the efficiency obtained from the model and the absorption coefficient (line). The efficiency of the phosphors below 40 eV is of particular interest since the absorption coefficients in that energy region are on the order of the diffusion length and contain multiple peaks. In addition, removing the effects of surface recombination from the data may reveal structure due to other processes affecting the photoluminescence near the bandgap energy.

At X24C it is now possible for us to reduce the uncertainties in our earlier data and to extend our measurements down to 3 eV. Initial analysis indicates that the lower energies surface recombination remains the dominant source of structure in the energy dependence of the efficiencies of these phosphors.

1. E. L. Benitez, D. E. Husk, C. Tarrío, and S. E. Schnatterly, *Appl. Phys. Lett.* 59, 396 (1991).
2. E. L. Benitez, D. E. Husk, S. E. Schnatterly, and C. Tarrío, *J. Appl. Phys.* 70, 3256 (1991).

Research supported by National Science Foundation Grant DMR-88-19052.

COMBINED PULSED LASER AND SYNCHROTRON RADIATION MEASUREMENTS OF PHOTOCHEMICAL DECOMPOSITION OF GaAs (110)

J. P. Long, S. S. Goldenberg, and M. N. Kabler (NRL)

We have recently shown that low-level laser irradiation of cleaved GaAs causes decomposition and the formation of Ga islands on the surface. PES measurements show three distinct manifestations of this process which appear and grow progressively with radiation: 1) a satellite at lower binding energy on the Ga 3d core spectrum, 2) a metallic edge in the band gap, and 3) increased band bending near the surface. A detailed fitting procedure has been developed and EDC's analyzed into surface, bulk, and metallic components in order to characterize the process. Fluence dependence is consistent with a 2-photon process. This is, in turn, consistent with arguments suggesting that single 2.43-eV photons are not sufficiently energetic to decompose GaAs. Post mortem SEM photographs show island morphologies entirely consistent with the PES data. The figure shows islands on cleaved GaAs subjected to 7.7×10^6 pulses at 3.9 mJ/cm^2 incident fluence. The viewing angle is 20 degrees grazing. The distribution of islands is evidently affected by nucleation at an unseen cleavage step.

— 0.1 μm

PHOTODECOMPOSITION STUDY OF HYDROGEN ADSORPTION ON VANADIUM DIOXIDE NEAR THE SEMICONDUCTOR - METAL PHASE TRANSITION

V.M. Benardes (NRL), R.T. Williams (Wales Forest U.), J.P. Long (NRL), R.K. Reed (Wales Forest U.), and P.H. Klein (retired)

Vanadium Dioxide (VO_2) is one of several oxides of first-row transition metals which are known to exhibit a semiconductor-to-metal phase transition (SMT) at some critical temperature T_c . Below $T_c \approx 340 \text{ K}$, VO_2 is a semiconductor (bandgap $\approx 0.7 \text{ eV}$); above T_c it is a metal.

It has been suggested that adsorbed hydrogen is responsible for the stabilized metallic phase in VO_2 films which have undergone electrochemical cathodic reduction. Because the SMT is accompanied by a characteristic shape-change of the V3d-derived feature at the top of the valence band (VB), UPS provides an attractive means for studying the relationship between adsorption and the SMT at the surface under carefully controlled conditions. We find that adsorbed hydrogen, rather than stabilizing the metallic phase, impedes it.

For the $\sim 1 \times 10^5 \text{ m}^2$, irregularly shaped, samples available in this study, ordinary cleaving failed to produce clean surfaces, most likely as a consequence of fracture along contaminated grain boundaries. However, abrading the surfaces yielded regions characteristic of a clean surface as shown in Fig. 1: a strong V3d VB feature, relatively little OH adsorption, and a strong SMT signature, in particular the appearance of a distinct Fermi edge in the high temperature metallic phase. Spectra were retaken as a function of temperature after various dosings with hot-filament-activated hydrogen, H_2 .

Results for surfaces exposed to 10^4 Langmuirs of H_2 are given in Fig. 2. Figure 2b documents the absence of the SMT as a consequence of dosing. The loss of signal intensity after the dose may be caused by a temporary, dose-induced decrease in electron multiplier gain; signal intensity recovered over time.

Primary conclusions from this study are: (1) H_2 partially reduces the surface with formation of OH in the adsorbed layer, as can be inferred from normalized spectra (not shown), where it is apparent that dosing increased the relative strength of the OH-derived VB feature; (2) Hydrogen adsorption quenches the SMT within the photoemission sampling depth (~ 5 lattice constants), possibly as a consequence of increased V-O π -bonding.

¹ A. Chenevas-Paule, J. Phys. (Paris) 37, C4-75 (1976).

PHOTOLESSON SPECTROSCOPY OF SEMICONDUCTORS EXHIBITING SURFACE-INDUCED INHOMOGENEITIES

J.P. Long, S.S. Coldenberg¹, and M.N. Kohler
Naval Research Laboratory

For those semiconductors where the intrinsic surface state density in the gap is insufficient to pin the Fermi level irrespective of extrinsic perturbations such as cleavage defects or deposited metal clusters, band bending varies transversely across the surface, leading to undesirable inhomogeneous broadening of photoelectron spectra. Until we developed a new line-fitting scheme, such inhomogeneities hindered extraction of the weak metallic gold component originating from Ga islands which formed on GaAs(110) after laser-induced photochemical decomposition.

The inhomogeneities are characterized by recording time-resolved spectra during the 5-ns laser pulse from a copper vapor laser ($h\nu=2.43$ eV). As seen in Fig. 1, the Ga 2d spectrum from an n-type sample is not only shifted by the laser-induced surface photovoltage (SPV), but also narrowed because the SPV reduces electrostatic inhomogeneities. Crucial to these experiments is the reduction in laser-synchrotron duty factor mismatch which ensues when the storage ring operates with a single electron bunch, the light pulses from which overlaps the laser pulse.

In the fits of Fig. 1a and Fig. 2, which shows ordinary spectra in the dark, inhomogeneities are incorporated by adding to each Voigt singlet used to model the bulk and surface spin-orbit doublets, a related, asymmetric, Voigt-derived singlet. The related singlets are derived from each parent singlet with the same parameters, which then constitute new fitting variables. The scheme was tested successfully against lines broadened theoretically with the aid of a Poisson-solving code which gave potential distributions both near circular islands and near linear features intended to mimic cleavage steps and chains of islands. The improved fits enabled the growth rate of Ga islands to be determined which led to the conclusion that a 2nd order process underlies the laser-induced decomposition.

¹ Sachs/Freeman Assoc., Landover, MD

C-1s NEXAFS AND EXAFS OF GRAPHITE, NATURAL DIAMOND, AND SYNTHETIC DIAMOND THIN FILMS

J. Nithianandam and J.C. Rife (NRL)

We have made reflectance, transmission, and photoemission measurements of the carbon 1s transitions in bulk graphite and bulk and thin film diamond. Data extend over a spectral range from 200 eV to 600 eV and measurements of the absorption edge region near 290 eV are at resolving powers better than 1200. The beamline optics were glow discharge cleaned to minimize carbon contamination flux losses. Background suppression and correction were required to obtain the highest quality data. Exciton features have been observed in synthetic thin films as well as bulk. Quantitative sp^2 to sp^3 bonding ratios, useful for understanding deposition processes, have been determined for a number of diamond thin films.

The figure shows the absorption coefficient of a unique diamond thin film with 30 nm diameter crystallites in the energy region just below the much stronger sp^3 edge at 290 eV. The 2.5 μ m thick thin film was grown by microwave CVD with 0.5% methane in hydrogen at 10 Torr pressure and 800C substrate temperature. Also shown are the absorption of the (2x1) reconstructed clean diamond (111) surface from photoemission measurements [Morar et al, Phys. Rev., B33, 1346 (1986)] and the absorption of graphite from electron energy loss measurements [Fink, Sol. St. Comm., 47, 687 (1983)]. We believe that the diamond thin film spectral features arise from

absorption at internal crystallite interfaces resembling the clean diamond (111) or (100) surface with negligible disorder induced effects.

Process-Dependent Spectral Reflectance Studies of thermal SiO₂ on Si
H. F. Wei, A. K. Henning, T. Hochwitz (Dartmouth College), J. Slinkman (IBM)
J. Rife (NRL), W. R. Hunter (Sachs/Freeman Associates)

A matrix of 64 device-quality p-type (100) 5" silicon wafers were RCA cleaned then dry oxidized at 800 °C, 900 °C, and 1000 °C (some wafers also received annealing in Ar) at IBM in Essex Junction, VT. Variable-angle-of-incidence spectral reflectance experiments were performed on 6 of these oxidized wafers at photon energies from 5 eV to 12 eV for a total of 5 angles of incidence at each photon energy. The purpose of this study is to extract absorption spectra from the reflectance spectra, then deconvolve the absorption spectra into individual subgap defect densities of SiO₂ films as a function of process variables. Process-dependent defect densities in thermal SiO₂ films is one of the essential elements in a ULSI device reliability simulator we are building.

Figure 1 shows the process-dependent reflectance spectra for 4 samples oxidized at 800 °C but each received different amount of Ar annealing (0 min, 10 min, 100, and 1000 min respectively) at 900 °C.

Figure 2 shows the extracted absorption spectrum for one of the samples, using the least square fitting of the Fresnel reflectance equations. Various defect states contributed to the absorption spectrum in excess to the Urbach tail which is resulted from the disorder states.

Figure 3 shows the refractive index spectrum for the same sample along with the refractive index of glass [1]. Based on the Lorentz-Lorenz analysis to figure 3, it is verified that the thermal SiO₂ film is more densified than glass.

[1] H. R. Philipp, in Handbook of Optical Constants of Solids, ed. E. D. Palik, (Academic Press, Inc., 1985), p. 749
This work was supported by an IBM-Shared University Research Grant through the General Technology Division, Essex Junction, VT 05452-4299.

PERFORMANCE OF AN ADAPTIVE JET COOLED SILICON MONOCHROMATOR IN A HIGH POWER DENSITY X-RAY BEAM

J. E. Berman (NSLS), M. Hart (U. Manchester), and S. Sharma (NSLS)

The X25 wiggler white beam, when doubly-focussed using a toroidal platinum-coated silicon mirror, has a power of 50 W at 130 mA in the hutch. The FWHM dimensions of the focal spot are 0.8 mm horizontal by 0.4 mm vertical, giving a power density in excess of 150 W/mm². This is the highest x-ray power density available to a synchrotron radiation user, and is comparable to the densities which optics at next-generation synchrotron source undulator beam lines will have to withstand. Hence X25 offers a unique test facility for such optics. For crystal monochromators to diffract x-rays efficiently, surface strains within the beam footprint, whether thermal or mechanical, must be kept much smaller than the Bragg reflection Darwin width. For all materials at room temperature, this level of performance is impossible to achieve for a conventional geometry in the presence of the X25 doubly-focussed power density. Adaptive techniques, however, may be used to compensate the thermal strain field. The schematic below right shows the crystal design which we have tested. The silicon crystal is a hollow cylinder capped by a 150 µm thick membrane. The lower (open) end of the cylinder is glued to a stainless steel single-water-jet manifold; the jet tube is aimed at normal incidence at the underside of the membrane, which is the diffracting surface. The crystal is surrounded by a can fitted with the beryllium windows (not shown); by pressurizing the can with helium, the membrane can be forced to bow into the cylinder, compensating the outwardly convex thermal distortion. Shown below are measured rocking curves using (a) the Si(111) fundamental reflection at 6.17 keV and (b) the Si(333) harmonic at 18.51 keV, as a function of can overpressure, at 160 mA. Whereas the theoretical rocking curve widths are 13 and 1 arcsec for the fundamental and harmonic respectively, the measured widths for the uncompensated crystal (zero overpressure) are 208 and 165 arcsec. These improve to 23 and 13 arcsec respectively for the compensated crystal (80 kPa overpressure), with concomitant gains in peak diffracted intensity.

This work was supported in part by the DOE, Contract No. DE-AC02-76CH00016.

PERFORMANCE OF BE CRYSTALS IN A HIGH POWER DENSITY X-RAY BEAM

A.K. Freund, S. Jokschi¹, K. Kawata², G. Marot, E. Ziegler (ESRF), L.E. Berman and J.B. Hastings (NSLS).

Beryllium mosaic crystal are currently proposed as monochromators and beam splitters for synchrotron X-ray beam lines. The mosaic spread should be about 1 arcmin for this application. For test measurements at high heat load we used the focused beam provided on X25 by a platinum coated toroidal mirror that had recently been installed at a glancing angle of 3 mrad (cut-off around 20 keV) and that condenses part of the wiggler beam into an area of about 1 mm (H) x 0.5 mm (V). The power measured at the focus is 75 W at 230 mA ring current which corresponds to a power density of 150 W/mm². A 12 mm diameter and 2 mm thick Be single crystal was kept in contact with a cooled Be support using springs. The whole assembly was enclosed in a He filled box to avoid deterioration of Be and put into the white beam. The third harmonic of the 110 reflection diffracted a beam of 18.3 keV that was analyzed by a Si perfect crystal (333 reflection) in almost dispersion-free geometry. An "ideal" mosaic rocking curve of 58 arcsec width was obtained for the "cold" beam increasing to 73 arcsec with the "hot" beam (150 W/mm²). No deterioration of the crystal was observed. The rocking curve broadening was also due to a nonuniform thermal contact between the Be crystal and the cooled Be substrate. With an improved cooling scheme and smaller thickness Be crystals will be capable of meeting the requirements specifications.

1 also HASYLAB 2 on leave from Photon Factory

PERFORMANCE OF VERY THIN SI CRYSTALS UNDER HIGH HEAT LOAD

A.K. Freund, G. Marot, K. Kawata¹, S. Jokschi², E. Ziegler (ESRF), L.E. Berman and J.B. Hastings (NSLS).

For test measurements at high heat load we used the focused beam provided on X25 by a platinum coated toroidal mirror that had recently been installed at a glancing angle of 3 mrad (cut-off around 20 keV) and that condenses part of the wiggler beam into an area of about 1 mm (H) x 0.5 mm (V). The power measured at the focus is 75 W at 230 mA ring current which corresponds to a power density of 150 W/mm². An 11 µm thin and 100 mm diameter Si membrane was glued at its periphery onto an aluminum ring at 30°C and then mounted on a box that could be pressurized with He gas. This box was supported by a copper manifold supplying cooling water to an optically polished Be block which was brought in contact with the Si foil. The same cooling loop controlled the temperature of the box and thus the Al ring. By increasing the water temperature above 30°C it was possible to apply a stretching force to the Si foil through thermal expansion of the Al ring. Applying pressure to the box permitted to push the Si foil against the Be substrate and to establish direct thermal contact. It turned out that surface imperfections or dust particles between the two surfaces produced such a big rocking curve broadening that the Si foil had to be slightly detached, leaving a He gap of a few µm between Si and Be. Then a 3.3 arcsec wide (333) diffraction profile was obtained with the "cold" beam increasing to 3.9 arcsec with a "warm" beam (10 W/mm²). The "hot" beam (100 W/mm²) destroyed the crystal, which showed that the thin He layer was not sufficient to conduct the heat away across the He gap to the Be substrate.

1 on leave from Photon Factory
2 also HASYLAB

INELASTIC X-RAY SCATTERING SPECTROSCOPY IN ALUMINUM

E.D. Isaacs, H. Williams, P.M. Platzman (AT&T); P. Zschack (ORAU); G.E. Ice, (ORNL).

The plasmon dispersion and lineshape in a single crystal of aluminum has been measured at the NSLS X25 wiggler beamline source. For small momentum transfers, $q \leq 1.6 \text{ \AA}^{-1}$ ($q_f = 1.75 \text{ \AA}^{-1}$), the inelastic spectrum consists of a single plasmon peak which disperses quadratically (see figure) as expected from RPA calculations. The integrated count rate of the plasmon peak at $q = 1 \text{ \AA}^{-1}$ is 50 cts/sec for a sample volume of $1 \times 1 \text{ mm}^2 \times 0.025 \text{ mm}$ thick. The lineshape in the low momentum transfer region is gaussian with a FWHM of 3 eV. This width is mostly the 2.2 eV resolution width of the spherically bent, LiF(004) scattered-beam analyzer, but also includes the incident beam band-width (1 eV) as well as the natural linewidth of the plasmon (1 eV). As q approaches q_f the linewidth broadens and evolves continuously into a peak with a double-bump structure, in the single particle continuum ($2 \text{ \AA}^{-1} < q < 5 \text{ \AA}^{-1}$). The center-of-mass of the scattering intensity in this intermediate momentum region shows a negative dispersion (see figure), which is not predicted by RPA and has been observed previously in other light metals such as beryllium, graphite, and sodium. Further studies in Aluminum will be performed with emphasis on the plasmon lineshape in the intermediate momentum transfer regime and with improved analyzer resolution.

Figure. Energy dispersion and linewidth of the plasmon in a single crystal of aluminum. Note the sub-quadratic dispersion above q_f . Cross - center-of-mass position of plasmon peak; box - gaussian width of plasmon peak; Solid line - quadratic fit to data.

HIGH RESOLUTION X-RAY ABSORPTION SPECTROSCOPY

K. Härmäläinen, D. P. Siddons, J. B. Hastings, L. E. Berman, C.-C. Kao, V. Stojanoff and S. P. Cramer (U. of California, Davis), NSLS, Bldg 725D, Brookhaven National Laboratory

XANES and EXAFS can give with a relative simple experimental set-up valuable information about the density of unoccupied states, which is sensy to bonding, oxidation and coordination of the system, and also local structural information even in disordered system. The resolution of absorption spectroscopy has not been limited by the experimental methods but mainly due to the natural lifetime broadening of the core hole. However, if the energy of the fluorescence photon following the absorption process can be measured with a better resolution than the lifetime width the lifetime broadening can be overcome [1].

We have developed a high resolution fluorescence spectrometer using a Johann geometry in a backscattering arrangement [2]. It is based on a spherically bent Si(440) crystal and the resolution is 0.3 eV at 6.5 keV. The figure below on left shows the measured dysprosium nitrate L α_1 fluorescence spectrum. The fluorescence photons within energy range corresponding to the analyzer resolution (dashed line on left) were monitored while scanning the incident energy through the Dy LIII-absorption edge. The resulting spectrum is shown by the solid line on the right and is compared with a conventional transmission measurement from the same sample and with same incident beam resolution (dashed line). The enhancement of the resolution is dramatical revealing fine structure of the edge which is totally invisible in the conventional XANES.

[1] K. Härmäläinen, D. P. Siddons, J. B. Hastings and L. E. Berman, Phys. Rev. Lett., 68, 2850 (1991)

[2] V. Stojanoff, K. Härmäläinen, D. P. Siddons, J. B. Hastings, L. E. Berman, S. Cramer and G. Smith, to be published in Rev. Sci. Instrum. (SRI 1991 Proceedings).

This work was supported by the DOE under Contract No. DE-AC02-76CH00016 and the Finnish Academy.

PERFORMANCE OF A GALLIUM COOLED 85° INCLINED SILICON MONOCHROMATOR FOR A HIGH POWER DENSITY X-RAY BEAM

W. K. Lee, A. T. Macrander, D. M. Mills, C. S. Rogers and R. K. Smith (APS, ANL) and L. E. Berman (NSLS)

The focussed wiggler X-ray beam at X-25 delivers about 24 W of power at 120 mA ring current into a spot size of FWHM $0.4 \times 0.8 \text{ mm}^2$, resulting in an average power density of about 75 W/mm^2 . An uncooled piece of silicon placed in the beam will be highly strained due to the thermal distortions. We have made double crystal rocking curve measurements of gallium cooled Si(111) at 5 keV in both the normal (which we will refer to as flat) geometry and the new inclined geometry. The inclined crystal geometry spreads the beam footprint out while maintaining a $b = -1$ symmetric Bragg reflection. At an 85° inclination angle, the beam footprint is 11.5 times larger than for the flat geometry. Figure 1 below shows the rocking curves for Si(111) at 5 keV for both the flat and the inclined crystal case. The sharp peak seen in the inclined crystal case is due to the flat 15 keV third harmonic registering in the ion chamber. Figure 2 shows the rocking curves for the 15 keV third harmonic Si(333). The data was taken at 145 mA ring current and at a constant gallium pumping rate of 2 gallons per minute. The advantage of the inclined geometry is clearly seen. The residual strain observed for the 15 keV Si(333) reflection (measured FWHM ~ 2.7 arc sec compared with the theoretical value of 1.1 arc sec) is totally due to the mounting of the crystals and not the heating from the X-ray beam. The same rocking curve width was obtained by placing a piece of aluminium absorber upstream of the monochromator. The thermal responses of the crystal were recorded with an IR camera. In the flat geometry case, the peak temperature was about 50°C above the nominal silicon temperature; while in the inclined geometry case, it was only 2°C above the nominal silicon temperature.

This work was supported by the U.S. Dept. of Energy, BES-Materials Sciences, under contract number W-31-109-ENG-38.

HIGH RESOLUTION INELASTIC X-RAY SCATTERING STUDY OF THE BORON K-ABSORPTION EDGE

S. Manninen, Department of Physics, University of Helsinki, Finland
K. Hämäläinen, NSLS, Bldg 725D, Brookhaven National Laboratory

XANES is a well-established technique for obtaining information about the band structure in solids. However, in the case of light elements conventional absorption measurements are difficult because the edge energies correspond to the UV or soft x-ray region and therefore very thin samples and clean surfaces are needed. Inelastic x-ray scattering can be used as an alternative method to get bulk information in these cases [1].

We have used inelastic x-ray scattering to study the near edge structure of beta-rhombohedral boron K-absorption edge. High resolution (0.7 eV incident, 0.3 eV analyzer [2]) combined with bright, intense synchrotron radiation make it possible to probe the fine structure of the density of unoccupied electron states. The figure below shows the measured spectrum as a function of incident energy while monitoring a fixed scattered photon energy with a high resolution analyzer. Our result does not show the predicted and previously observed peak due to the $1s$ to d transition. Work to improve the experimental setup and analyzer for better signal-to-noise ratio is in progress.

(Left) Measured inelastically scattered spectrum from boron sample, (right) K-edge region at 6676 eV.

- [1] W. Schulte, U. Bonse, H. Nagasawa, A. Kaprielian and A. Bertoldi, Phys. Rev. B, 38, 2112 (1988)
- [2] K. Hämäläinen, D. P. Siddons, J. B. Hastings and L. E. Berman, Phys. Rev. Lett., 68, 2850 (1991)
- [3] A. D. Zetis and D. K. Papadimitrou, Phys. Rev. Lett., 60, 61 (1988)

This work was supported by the DOE under Contract No. DE-AC02-76CH00016 and the Finnish Academy.

CRYOGENIC COOLING OF SI MONOCHROMATORS

G. Marot, A.K. Freund, M. Rossat, S. Jokschi¹, K. Kawata², E. Ziegler (ESRF), L.E. Berman, D. Chapman, J.B. Hastings, M. Iarocci G. van der Laske and N. Lazare (NSLS).

For test measurements at high heat load we used the focused beam provided on X25 by a platinum coated toroidal mirror that had recently been installed at a glancing angle of 3 mrad (cut-off around 20 keV) and that condenses part of the wiggler beam into an area of about 1 mm (H) x 0.5 mm (V). The power measured at the focus is 75 W at 230 mA ring current which corresponds to a power density of 150 W/mm². Two cooling schemes were studied inside a vacuum cross at a fixed Bragg angle of about 18 degrees: a directly cooled Si crystal of 100 x 80 x 16 mm³ with two INVAR manifolds glued onto its side faces with cryogenic glue, and a side-cooled crystal of 100 x 80 x 50 mm³ with two cooled copper blocks attached to the 100 x 80 mm² side faces via indium foils and using a spring loaded mechanism. When strongly attenuating the incident beam ("cold" beam) the double crystal (333) rocking curve width was about 0.4 arcsec broader than the perfect crystal one with a repeatability of 0.2 arcsec that was also the mechanical resolution of the rotation stage. During cooling down to -190 C the rocking curve width very slightly decreased but essentially remained constant, i.e. around 1.2 arcsec. With the unattenuated "hot" beam the width progressively decreased from about 58 arcsec at room temperature to 1.7 arcsec at -180 C (see figures). From the latter result we deduce a thermal slope error of about 1 arcsec at a power density of about 150 W/mm² normal to the beam or 50 W/mm² on the crystal surface. Both samples performed equally well.

1 also HASYLAB

2 on leave from Photon Factory

KINETICS OF THE ELECTROCHEMICAL INDUCED RECONSTRUCTION OF THE Au(111) SURFACE

Ben Ocko, Jia Wang, Hugh Isaacs, Alison Davenport, and Lonny Berman (BNL)

We have carried out surface x-ray diffraction studies of the Au(111) surface in an electrochemical cell. Charging the surface negative induces the (23x1) striped surface reconstruction which exhibits the diffraction pattern shown in Fig. 1. The reconstructed surface exhibits four extra spots surrounding the (0,1) reflection which are arranged in a hexagonal pattern. We label the direction connecting one of the primary reconstruction spots and the (0,1) reflection as the q_r axis. At sufficiently positive potentials these additional spots vanish and this is associated with the transition to (1x1) symmetry.

Figure 1

Information on the nucleation and kinetics of the electrochemically induced striped phase is contained in the time evolution of the scattering profiles after a step change in potential. Measurements were carried out in 0.01M NaCl with 0.4M NaF solution. The scattered intensity is recorded as the potential is continuously cycled between 0.5V (1x1) at $t=0$ sec. and -0.3V (reconstructed) at $t=20$ sec. with the cycle shown in Fig. 2 (top). The position along q_r is advanced after every potential cycle in order to acquire the scattering along q_r versus time. In Fig. 2, equal intensity contours lines ($\times 0.65$) are shown. Within several seconds after raising the potential to +0.5V, the intensity contours associated with the striped phase, $q_r=0.038[\text{\AA}^{-1}]$ vanish and there is a subsequent increase in the intensity at $q_r=0$. About five seconds after the potential is lowered to -0.3V there is a rather uniform increase in the scattered intensity all the way from $q_r=0.01[\text{\AA}^{-1}]$ until $0.05[\text{\AA}^{-1}]$ with no apparent peak. We associate this behavior with a disordered striped phase in which an ordered strip length has not been established. At about 10 seconds after lowering the potential to -0.3V an ordered striped phase is established yet the correlation length is rather broad. In principle, the behavior of the stripe correlation length would provide invaluable information on the growth mode of the striped phase. Detailed analysis is ongoing.

Figure 2

M. Sutton, E. Dufrane (McGill University), S.G.J. Mochrie (MIT), L.E. Berman (NSLS), G. Held, G.B. Stephenson (BNM)

One effect observed when using coherent light to illuminate a sample containing disorder is a graininess in the scattered light called speckle. Speckle results whenever randomly distributed regions of a material introduce different phase shifts into the scattering of coherent incident light. If the arrangement of these regions evolves in time, the speckle pattern will also change. Observation of the intensity fluctuations at a single point in the speckle pattern provides a direct measure of the time correlations of these fluctuations. This leads to a technique^[1], alternately called light beating spectroscopy, dynamic light scattering or intensity fluctuation spectroscopy. This technique is commonly used with laser light and by extending it to x-rays, one can probe smaller length scales and one can probe opaque materials. The observation of speckle in the (100) Bragg peak of Cu_3Au was previously demonstrated^[2]. Here we report on our progress in performing x-ray intensity fluctuation spectroscopy.

We used X25 without a mirror and with a double crystal Si (111) monochromator set to 1.77 Å. The incident beam was collimated by a 6 μm diameter pinhole placed 28 m from the source. Figure 1 shows the Fraunhofer diffraction pattern in the 2θ direction for this pinhole. With the installation of the monochromator upstream of the hutch, the background radiation has been greatly reduced, and we can easily observe the wiggles in the diffraction pattern over the 10^{-4} drop in intensity shown.

To observe intensity fluctuations we use the (100) peak in Cu_3Au . The experiment is carried out by first annealing the crystal at a temperature in the disordered phase, then quenching it to a temperature in the ordered phase and observing the scattering as a function of time. Figure 2 shows the result of one such quench. Conventional x-ray measurements^[3] show a monotonic increase in scattering. The observed non-monotonic behavior with our coherent x-ray beam is due to the intensity fluctuations predicted. These results have been confirmed during the July 1991 undulator run at CHESS. Many thousands of similar time series were collected by using a 2D detector array to collect data in parallel.

[1] P.N. Pusey, in *Photon Correlation Spectroscopy and Velocimetry*, E.Z. Cummins and E.R. Pike eds., Plenum, New York (1977), 45-141.

[2] M. Sutton et al, *Nature*, **352**, 608 (1991)

[3] S.E. Nagler et al, *Phys. Rev. Lett.*, **61**, 718 (1988).

Figure 1: Fraunhofer diffraction of a 6 micron pinhole. An 8.5 micron diameter pinhole at 1 meter was used to collimate the diffracted beam. The solid line represents a guide to the eye.

Figure 2: The intensity of scattered x-rays as a function of time after a quench from 395°C to 375°C at time $t=0.0s$. The diffracted beam was collimated by a 33 μm diameter pinhole 1m from the sample. This pinhole was centered on the (100) Bragg peak of Cu_3Au .

X-RAY SEARCH FOR CDW SATELLITES IN $\text{Ba}_{1-x}\text{K}_x\text{BiO}_3$.

P. Wochner, Q. J. Wang and S. C. Moss (U. of Houston)*; S. K. Sinha (Exxon); G. Grübel, H. Chou, L. E. Berman and J. D. Axe, (Brookhaven National Lab); C.-K. Loong (Argonne National Lab); J. Z. Liu, (U. of California, Davis)

Two flux-grown single crystals of $\text{Ba}_{1-x}\text{K}_x\text{BiO}_3$ were studied on the wiggler insertion line: one in the middle of the semiconducting regime ($x=0.23$) studied at 300K and one near the boundary ($x=0.35$) between semiconductor and metal studied at 300K and 11K. The proposed charge density wave (CDW) satellites, seen earlier with electron diffraction,^{1,2} had been attributed possibly to electron-beam effects² even though their position seemed to vary smoothly (linearly) with K content.¹ A careful search, both with and without a Si(111) analyzer crystal, in all equivalent $\langle 110 \rangle$ directions of the pseudocubic reciprocal lattice, revealed no satellite structure in either of the samples. We therefore conclude that the electron diffraction results may well be induced through electron-beam effects.²

1. S. Pei et al., *Phys. Rev. B* **39**, 811 (1989); *ibid* **41**, 4126 (1990).

2. E. A. Hewat et al., *Physica C* **157**, 228 (1989).

*Research supported by the NSF on DMR-8903339 and the State of Texas at the Texas Center for Superconductivity.

HIGH HEAT LOAD STUDY OF MULTILAYER MONOCHROMATORS AT ROOM AND LOW TEMPERATURES

E. Ziegler, A.K. Freund, G. Marot, S. Jaksch¹, K. Kawata², M. Rossat, (ESRF),
L.E. Berman, J.B. Hastings, M. Iarocci, G. van der Laske and N. Lazarz (NSLS).

For test measurements at high heat load we used the focused beam provided on X25 by a platinum coated toroidal mirror that had recently been installed at a glancing angle of 3 mrad (cut-off around 20 keV) and that condenses part of the wiggler beam into an area of about 1 mm (H) x 0.5 mm (V). The power measured at the focus is 75 W at 230 mA ring current which corresponds to a power density of 150 W/mm². Three samples deposited on Si substrates were studied in vacuum (W/Si, W/B₄C, produced by Ovonic, and Fe/C, provided by Philips) in order to test their resistance to high radiation doses. They were contacted to a cooled Si support with vacuum grease and positioned at about 2° glancing angle where they reflect an X-ray energy of 6.1 keV in the first order. A perfect Si(111) crystal was used to analyze the reflected beam in the +, + dispersive setting. The W/B₄C and Fe/C multilayers were exposed for four hours to the "hot" beam and did not show any alteration to the rocking curves recorded with the perfect Si analyser in a θ -2 θ scan mode. The W/Si sample was exposed for ten hours and, during that time, also cooled down to liquid nitrogen temperature in order to see whether it withstood cryocooling. Again, no changes of the rocking curves were observed during and after exposure. This shows that the cryogenic cooling technique can also be applied to multilayers on Si substrates and that multilayers are capable of resisting to high power density X-ray beams, at least for several hours. Longer exposures are now required.

1 also HASYLAB
2 on leave from Photon Factory

Lunar Volcanic Glasses: Trace Element Abundances in Individual Spheres

J. W. Delano¹, S. R. Sutton^{2,3}, and J. V. Smith²

¹Dept. of Geological Sciences, SUNY, Albany, NY 12222

²Department of the Geophysical Sciences and Consortium for Advanced Radiation Sources, The University of Chicago, Chicago, IL 60637

³Department of Applied Science, Brookhaven National Laboratory, Upton, NY 11973.

Mare volcanic glasses appear to be primary magmas from the lunar mantle [e.g., 1]. Many questions remain to be answered about the mare volcanic glasses, but one of the most interesting ones deals with the source(s) of volatile elements (e.g., Cu, Zn, Ga, Pb) associated with these magmatic eruptions [e.g., 2]. Although details remain complex [e.g., 3], these volatiles may have implications for not only the Moon's earliest thermal history but also its origin. Analysis of mare volcanic glasses by synchrotron x-ray fluorescence provides data on the trace abundances of lithophile and volatile elements that bear on these scientific issues.

A total of 13 spheres was analyzed - 10 Apollo 15 green, 1 Apollo 17 orange and 2 Apollo 15 yellow. The detected trace elements included Ni, Cu, Zn, Ga, Rb, Sr, Y, Zr and Nb. The initial results are in agreement with analyses of other investigators of individual glass spherules by instrumental neutron activation [Ni, Rb, Sr; e.g., 4] and ion microprobe [Zn, Sr, Zr; e.g., 5] with the exception of Cu. Our result for the Apollo 17 orange glass gives < 1 ppm Cu, whereas ion microprobe analyses yielded abundances of about 450 ppm [6]. To the best of our knowledge, our results for Ga, Y and Nb are the first reported for the interiors of individual spheres.

Figure 1 is a ternary diagram involving non-volatile, geochemically incompatible elements (Ti, Zr) and a volatile, incompatible element (Zn). Due to the large differences in abundances among these elements, coefficients have been applied (e.g., 500 * Zn) to give comparable weighting to each element. Our data show that the abundance of Zn in the high-Mg magmas represented by the mare volcanic glasses (i.e., not surface-correlated Zn) is strongly enriched compared to the magmas represented by the crystalline mare basalts. This figure supports the view that the volcanic glasses were derived from generically different mantle source-regions than the mare basalts.

Acknowledgments: NASA NAG-78 (JD) and NAG-106 (SRS), NSF EAR-86-18346, DOE DE-AC02-76CH00016, NSLS staff, U. of C. CARS Startup Grant, State of Ill. Tech. Challenge Grant. References: [1] Delano (1986) PLPSC 16, D201-D213; [2] Chon et al. (1975) PLSC 6, 1701-1727; [3] Tatsumoto et al. (1987) PLPSC 17, E361-E371; [4] Hughes et al. (1988) GCA 52, 2379-2391; [5] Galbreath et al. (1990) GCA 54, 2565-2575; [6] Shearer et al. (1990) EPSL, in press

Figure 1: Ti-Zr-Zn ternary diagram of the mean volcanic glass compositions and the field of mare basalt compositions. The volcanic glasses are enriched in the volatile element Zn relative to the crystalline mare basalts.

DETECTION AND QUANTIFICATION BY SRXRF OF SOME TRACE METALS IN CORN BRAN AND CORN BRAN PASSED THROUGH THE PIG GASTROINTESTINAL TRACT

F. Dintzis (NCAUR, ARS, USDA) AND K. Jones (NSLS)

The objective of this experiment was to use SRXRF to detect and quantitate some trace metals in corn pericarp tissues retrieved from the GI tract of pigs. Corn pericarp tissues for use as controls were loaded in-vitro with Fe, Zn and Cu. Concentrations of 100 PPM Fe and Zn and 70 PPM Cu in the wet ashed bulk controls were measured by atomic absorption.

Tissue samples were embedded in IR White Acrylic Resin, soft grade, and cut to a thickness of 2 microns with a microtome on a dry glass knife. The sectioned tissues were fastened at each end with Elmer's glue to a backing of thin Kapton film. A beam aperture of 20 x 20 microns was used with counting times of 2 minutes to collect x-ray data. X-ray counts from filter paper loaded with known concentrations of elements provided a standard to estimate the concentrations of minerals detected in the corn pericarp sections.

Concentrations estimated from x-ray measurements were quite heterogeneous, even when obtained from the same specimen. Concentration ranges of Zn, 0-57 PPM, and Cu, 0-12 PPM, were estimated from SRXRF data for the in-vitro loaded specimens. These values were significantly lower than the bulk values measured by atomic absorption. Values of Fe in these specimens varied from 0-6 PPM, but are considered to be unreliable because of suspected contamination associated with the target matrix.

As estimated by SRXRF data, concentration ranges of Zn, 0-100 PPM, Fe, 0-50 PPM, Cu, 0-5 PPM, and Mn, 0-30 PPM, were found in the pericarp tissues retrieved from various locations in the GI tract of pigs. Concentrations were highly heterogeneous, as in the in-vitro loaded tissues.

This experiment demonstrates that SRXRF can be useful in detecting trace minerals of nutritional importance in food components exposed to mammalian digestive processes. Methods to accurately quantitate such measurements require further attention.

Cosmic Class Stratospheric Particles: Trace Elements and Zn Depletions

G. J. Flynn

Department of Physics, SUNY-Plattsburgh, Plattsburgh, NY 12901

S. R. Sutton

Department of Applied Science, Brookhaven National Laboratory, Upton, NY 11973.

Minor and trace element abundances were measured by synchrotron x-ray fluorescence on eight stratospheric particles: five "cosmic" (C) and three "possibly cosmic" (C?) in the Johnson Space Center (JSC) preliminary classification. The particles had typical dimensions of 10 micrometer, making the present set of particles smaller than most we have previously analyzed [1]. Our main goals in the present study were to better define the constituents of the C and C? classes. One of the more interesting observations is that Zn concentrations in three of the chondritic particles are extremely low. One possibility is that these depletions result from partial heating of the particles.

Zn was depleted by more than a factor of 10 in 3 of the particles which exhibited chondritic minor/trace element patterns in our analyses. Since Zn is quite volatile, with a loss temperature of 600°C in heating experiments on meteorites [2], the low Zn abundances in particles exhibiting otherwise chondritic trace element patterns may result from particle heating, as might occur on atmospheric entry, for example.

The Fe/Ni ratio, when coupled with major element abundances, appears to be a useful discriminator of cosmic particles. The C-type particle without detectable Ni in its JSC EDX spectrum exhibited a clearly non-chondritic minor/trace element abundance pattern, as have the previous low-Ni particles we have analyzed (1). Two of the C? particles having Fe/Ni peak height ratios similar to Allende in their JSC EDX spectra also exhibited chondritic minor/trace element abundance patterns, suggesting they are extraterrestrial.

Acknowledgments: NASA NAG-459 (GF), NAG-106 (SRS), DOE DE-AC02-76CH00016, NSF EAR86-18346, NSLS staff, U. of Chicago CARS Startup Grant, State of Ill. Tech. Challenge Grant. References: [1] Flynn, G.J. and Sutton, S.R., Proc. 20th Lunar Planet. Sci. Conf., 335-342 (1990). [2] Krammuddin, M. et al., Geochim. Cosmochim. Acta, 41, 1247 (1977).

Figure 1: Trace element abundances for chondritic particle W7029B5 showing a large Zn depletion.

IDENTIFICATION AND QUANTIFICATION OF TOXIC TRACE ELEMENTS IN MID-CONTINENTAL ATMOSPHERIC AEROSOLS

Richard H. Grant and Darrell G. Schulze (Purdue University, West Lafayette, IN)
Stephen R. Sutton and Mark L. Rivers (University of Chicago, Chicago, IL)

Industrial sites, unvegetated mine spoils, and landfills are potential fugitive sources for heavy metals in atmospheric aerosols. To measure the dispersion of aerosols from such sites, one must know the chemical composition of the aerosols in the ambient air. The "ambient" aerosol composition can vary, however, depending on the source region(s) of the air masses arriving at the site. The purpose of this study is to relate the aerosol chemistry of air masses sampled in the Midwest to the air mass trajectories and source regions. Only a few 10's of micrograms of aerosol may be available for analysis from size-fractionating air samplers and microanalytic techniques are necessary to analyze such small samples. The elemental composition of individual particles is also of interest.

We studied atmospheric aerosols collected near Vincennes, Indiana which were representative of air arriving from Canada (1 sample), the Atlantic Ocean (1 sample), or the Gulf of Mexico (3 samples) within 3 days, or which had been stagnant over the Midwest for >7 days (3 samples). Trace element composition of the 3.3 - 7 μ m aerosol fraction was analyzed on the synchrotron x-ray fluorescence microprobe on beamline X26A. Elemental concentrations were normalized with respect to Fe.

The concentrations of Ca, K, Ti, Mn, Sr, and Rb were generally similar to those of average continental crust, indicating little or no enrichment from anthropogenic sources. Calcium contents of one Gulf and one Midwest sample were exceptions, suggesting a local source of Ca. Copper, Zn, Br, and Pb, however, were enriched in the aerosols relative to the crustal average. Of the total flux of Cu and Zn into the atmosphere, about 75% of the Cu and 90% of the Zn is from anthropogenic sources (Nriagu, 1979; Nriagu and Davidson, 1980). Thus, the elevated Cu and Zn levels are indicators of enrichment from anthropogenic sources. The elevated levels of Pb are presumably the result of anthropogenic sources as well. Elevated Br levels may be the result of either anthropogenic or natural marine sources.

The Cu content of the Canadian air mass was near the crustal average and was at least an order of magnitude lower than the other air masses. The low Cu content, coupled with low Pb and Zn contents, suggests that the Canadian air was relatively unmodified by anthropogenic pollutants. If Canadian air masses are generally lower in anthropogenic pollutants such as Cu and Zn, aerosol emissions from point sources can best be evaluated when Canadian air is arriving in the study area. We plan to include additional Canadian samples in future microprobe runs to test this hypothesis.

Nriagu, J. O., and C. I. Davidson. 1980. Zinc in the atmosphere. *In* J. O. Nriagu (ed.) Zinc in the environment I. Ecological cycling. Wiley, NY.

Nriagu, J. O. 1979. Copper in the atmosphere and precipitation. *In* J. O. Nriagu (ed.) Copper in the environment I. Ecological cycling. Wiley, NY.

We gratefully acknowledge travel support from the Department of Energy's Division of University and Industry Programs, Office of Energy Research, NSLS Faculty/Student Research Support Program (Grant) and from the Purdue Agricultural Experiment Station (Schulze).

Chromium Oxidation State in Individual Lunar Olivine Grains using MicroXANES

S. R. Sutton^{1,2}, K. W. Jones¹, B. Gordon¹, M. L. Rivers^{1,2}, and J. V. Smith²

¹Department of Applied Science, Brookhaven National Laboratory, Upton, NY 11973.

²Department of the Geophysical Sciences and Consortium for Advanced Radiation Sources, The University of Chicago, Chicago, IL 60637

The oxidation states of minor and trace elements in lunar minerals provide important clues on the physico-chemical history of the Moon. Chromium in olivine has been particularly intriguing since lunar olivines are enriched in Cr compared with terrestrial olivines [1,2]. Steele and Smith [1] suggested that minor element contents were most readily explained if both Cr^{II} and Cr^{III} are present, the latter residing in a couple with Al. However, direct evidence that Cr exists in a reduced chemical form in lunar olivine has been difficult to obtain. We report here our first attempts to apply x-ray absorption near edge structure (XANES) in a microprobe mode in order to obtain spectra on individual co-existing phases in conventional rock thin sections.

The synchrotron x-ray fluorescence microprobe on beamline X26 at the National Synchrotron Light Source (Brookhaven National Laboratory, NY) was used in its normal configuration (see e.g., [3]) with the addition of a Si(220) channel-cut monochromator on the incident radiation. A beam size of 200 micrometers was used in the present experiments. Each XANES measurement consisted of scanning the incident beam energy from just below to just above the Cr K absorption edge and recording the Cr K alpha count rate at each incident energy. Typically, a scan of 100 eV was made in 500 steps (0.2eV/step) and each energy was counted for 5 seconds for a total scan time of about 1 hour. The resulting spectra were then compared with those of reference materials to infer valence state.

MicroXANES spectra are shown in the figure for lunar rock 15555 olivine, Cr^{II} in CrCl₂, and Cr^{III} in a synthetic forsterite containing 1.25% Cr₂O₃. The mean olivine edge is intermediate to those of Cr^{II} and Cr^{III} supporting the hypothesis that Cr in lunar olivine exists as both Cr^{II} and Cr^{III}. Additional high resolution work is needed to quantify this ratio.

Acknowledgments: NASA NAG9-106, DOE DE-AC02-76CH00016, NSF EAR86-18346, NSLS staff, U. of C. CARS Startup Grant, State of Ill. Tech. Challenge Grant. References: [1] Steele and Smith (1975) Proc. Lunar Sci. Conf. 6th, 451-467; [2] Smith and Steele (1976) American Mineralogist 61, 1059-1116; [3] Flynn and Sutton (1990) Proc. 20th Lunar Planet. Sci. Conf., 335-342.

Figure 1: Cr K microXANES spectra for lunar rock 15555 olivine, Cr^{II} in CrCl₂, and Cr^{III} in synthetic forsterite. The intermediate absorption edge of the lunar olivine suggests the presence of Cr^{II} and Cr^{III}.

Experimental Determination of the X26 Bending Magnet X-ray Source Size

S. R. Sutton and M. L. Rivers
Department of the Geophysical Sciences and Consortium for Advanced Radiation Sources,
The University of Chicago, Chicago, IL 60637;
Department of Applied Science, Brookhaven National Laboratory, Upton, NY 11973.

The X26 bending magnet x-ray source size was determined experimentally using a pinhole camera approach. Source size measurements are of interest in predicting the minimum synchrotron beam size attainable with apertures and in verifying theoretical machine parameters.

The experimental apparatus consisted of a beam-defining aperture (combination V and slit), an ion chamber (Io monitor), a pinhole collimator, a pure silica scatterer and solid state x-ray detector. The V-slit aperture resided in the beamline 6 meters from the synchrotron source and 3 meters from the pinhole in the hutch (ion chamber between the aperture and pinhole). The scatterer was positioned 3 cm downstream from the pinhole with its surface vertical and at 45 degrees to the beam. The detector was at about 60 degrees to the incident beam (i.e., looking somewhat upstream) and within the storage ring plane. In this experiment, the V-slit opening was set at about 1000 square micrometers, so that the source could be imaged in the hutch. The hutch lift table supporting the pinhole, scatterer and detector was then scanned vertically in 10 micrometer steps. The scattered radiation intensity at each vertical table position was used to map the vertical beam profile.

The scattered intensity exhibited a Gaussian dependence with pinhole (table) height (see figure). For scattered radiation between 14 and 16 keV, the standard deviation of the profile was 58 micrometers. Since the source-to-aperture distance (6 meters) was twice the aperture-to-pinhole distance (3 meters), the system had an inherent demagnification of 2:1 (about 10% reduction in apparent source size). Consequently, the inferred vertical source size σ_y was 116 micrometers (reasonably consistent with the NSLS estimate of the typical vertical source size of about 150 micrometers). Assuming an emittance ϵ of 0.102E-06 (NSLS X-ray Ring) and a vertical betatron function β_y of 15 m (estimate for X26 port), an emittance coupling χ of about 1% was calculated.

Diffraction effects were tested for by making measurements at 3 different aperture sizes and by measuring the scatter at different energies. Decreasing aperture size by a factor of 4 resulted in a 10% reduction in apparent source size. The energy dependence of the source size was small, about 5% from 5 to 30 keV.

Acknowledgments: NASA NAG9-106 (SRS), NSF EAR86-18346, DOE DE-AC02-76CH00016, NSLS staff, Univ. of Chicago CARS Startup Grant, State of Illinois Technology Challenge Grant.

Analysis of Oceanic Fluid Inclusions by Synchrotron X-ray Fluorescence

David A. Vanko, Dept. of Geology, Georgia State U., Atlanta, GA 30303-3083
S.R. Sutton, and M.L. Rivers Dept. of Applied Science, Brookhaven National Laboratory, Upton, NY 11973

Approximately thirty hours of beamtime was used in May, 1990, to perform a number of reconnaissance experiments on the X26A Synchrotron X-Ray Fluorescence (SXRF) microprobe, with the aim of obtaining x-ray spectra from minute natural and synthetic fluid inclusions in quartz. Encouraged by the success of Frantz et al. (Analysis of fluid inclusions by x-ray fluorescence using synchrotron radiation, Chemical Geology, 69, 235-244, 1988) in their analysis of synthetic fluid inclusions bearing calcium and zinc chloride, we attempted to determine to what extent SXRF was suitable for studying natural inclusions. We were particularly interested in testing the ability of the microprobe to "see" variations in major cation composition of bulk fluid inclusions, and to detect minor elements.

Natural fluid inclusions from several extinct seafloor hydrothermal systems were examined. Brine-filled inclusions, typically 10 micrometers across and containing halite-saturated solutions, were known on the basis of prior studies to contain aqueous fluids of mineralized sodium and calcium chloride. Spectra obtained by SXRF revealed the presence of Ca and, in some cases, K. Na was not detected because the Na Kalpha radiation is absorbed by the sample itself and by the air path to the detector. At higher energies, spectra from these inclusions showed significant peaks for Mn, Fe, Cu, and Zn. In a few cases, Br and Sr were also detected.

These reconnaissance results are exciting because of the potential for quantifying the concentrations of both the major and minor cations in hydrothermal fluids. The calcic nature of the brines and their metal-carrying capacity are both significant geochemical characteristics that speak to the chemical evolution of seawater as it permeates oceanic crust, and the ability of the fluid to transport and deposit massive sulfide deposits on the seafloor. Further studies are planned to develop a protocol for quantitative analysis of Ca, K, Mn, Fe, Cu, Zn, Cl and Br in natural and synthetic fluid inclusions in quartz.

ANALYSIS OF SYNTHETIC FLUID INCLUSIONS IN QUARTZ BY SXRF MICROPROBE

D.A. Vanko (Georgia St. U.), S.R. Sutton and M.L. Rivers (BNL)

We are systematically analyzing a set of synthetic NaCl-CaCl₂-H₂O fluid inclusions (each about 5-10 μm in diameter) hosted by natural Brazilian quartz. The synchrotron beam is stopped down to approx. 10-15 μm across, and inclusions within about 15 μm of the quartz surface are irradiated. Characteristic Ca and Cl x-rays from the fluid are detected by a solid-state Si(Li) system. After peak fitting, one may determine the intensity ratio Ca/Cl.

The intensity ratio is corrected for differential absorption by the fluid in the inclusion, the quartz, the airpath between sample and detector, and the detector's Be window. Relative fluorescence yields are calculated with the program NRLXRF. Careful measurement of the quartz thickness is critical in this correction: we are investigating the application of confocal imaging microscopy as a technique to make this measurement accurately.

The elemental Ca/Cl ratio of the inclusion fluid can presently be determined to +/- 20%. Combined with independent knowledge of the total freezing point depression of the inclusion fluid (obtained by freezing-stage microscopy), the bulk composition can be solved. Results are useful in modelling fluid-rock equilibria in hydrothermal systems, with applications in ore deposits, petrology, and geochemistry.

Research sponsored in part by NSF EAR-9103530 and DOE DE-AC02-76CH00016.

ANALYSIS OF TITANIUM IN BORON NITRIDE COATING ON A BERYLLIUM WINDOW

I. Gmür (NLS) and M. Rivers (U. of Chicago)

A commercially available boron nitride (BN) aerosol (Combat® Boron Nitride Aerosol Spray, The Carborundum Company, Electronic Ceramics Division, P.O. Box 664, Niagara Falls, NY 14302) has been found to protect white x-ray beamline beryllium windows, whose surfaces are exposed to atmosphere, from contamination and degradation. Experience on NLS beamline X25 showed that when a detector was mounted in-line with a 6 μm pinhole and scanned in 10 μm steps across an 8 keV beam transmitted through a BN coated Be window, beam intensity varied up to 25%, point-to-point (Brian Stephenson, IBM, and Mark Sutton, McGill U. - priv. comm.). Previous EDS and EELS analyses revealed the presence of titanium in the BN aerosol, possibly used as a binding agent. Since such variations in beam intensity could not be due to absorption by BN alone, the Ti was suspected.

Fluorescence analysis over 300 seconds of a single area on a BN coated Be window revealed a strong Ti K_α peak (Fig. 1 - other peaks due to Be substrate and Ar in air). The maximum measured intensity corresponds to ~160 μg/cm² Ti (based on Ti reference sample). This thickness is calculated to absorb 8 keV by only 3.5%. The X25 window would have to have been sprayed 7 times thicker than the window we analyzed to explain the 25% intensity modulations. A Ti fluorescence scan of 10 μm steps over 0.5 mm at 5 sec step showed >140% variations in Ti concentration over 10 μm distances and factors of 6 over 100 μm distances. Fig. 1 shows that the Fe concentration is greater than Ti and thus would absorb 8 keV by a greater percentage; however, the Fe is uniformly distributed in the Be window and would thus not be responsible for variations in beam transmission.

While not correlating directly with each other, due to variations in aerosol spray thickness and uniformity of spray on the Be windows, the analyses above show that the variation in transmitted beam intensity could indeed be caused by non-uniform distribution of Ti in the BN coating.

These findings indicate that the decision whether or not to use BN on Be windows should be based, in part, on a) the need for uniform beam transmission at lower x-ray energies across the window, and b) the use of small pinholes or detector pixel sizes.

I. Gmür, N.F. October 1991. Boron Nitride Protective Coating of Beryllium Window Surfaces. NLSL, BNL, Upton, NY. BNL Informal Report 46597. 21 pp.

Fig. 1
Work funded by DOE Contract No. DE-AC02-76CH00016 and NSF EAR86-18346 (MLR).

MEASUREMENTS OF COMPTON-EJECTED ELECTRONS FOR Ne GAS USING FILTERED "WHITE LIGHT" OF 10-15 keV

N. A. Guardala, J. L. Price, M. F. Stumborg, D. G. Simons (Naval Surface Warfare Center); G. A. Glass, Jr. (U. of Southwestern Louisiana); B. M. Johnson, K. W. Jones (BNL)

Initial studies of the double differential cross sections of Compton-ejected electrons were undertaken this past August at the NSLS. Beam line X26C was used for the measurements. The "white light" x rays originating from the storage ring were filtered twice in order to achieve the bandwidth quoted above--first with a thick Be window, which produces a low-energy cutoff at 3 keV and then by the use of a thick 3-mm Kapton filter. This arrangement ensured that the majority of electrons were ejected from the K shell of Ne.

The ejected electrons were energy analyzed using a spherical sector electrostatic spectrometer having an energy resolution of ca. 3%, an angular resolution of 5°, and a maximum attainable electron kinetic energy of 4.5 keV. The electrons were detected with a channel electron multiplier placed at the exit aperture of the spectrometer. A gas jet was used to provide a steady supply of Ne target atoms into the x-ray interaction region. The spectrometer focal point was positioned at approximately the center of the gas jet/x-ray interaction region. In this set of experiments the spectrometer was placed at 90° with respect to the incoming x-ray beam. The voltage in the sectors of the electron analyzer was swept in a periodic manner from 0-2000 V. Typically, this was done with 30-second scans involving steps of 4 V. The applied voltage was incremented using a CAMAC-controlled data acquisition system. K-shell Auger electrons were detected as a means of normalization so that values of the absolute cross sections could be obtained.

The next set of experiments is scheduled for early December. Modifications to the gas jet, the scattering chamber, and the electron analyzer have been made in order to improve the energy and spatial resolution. These experiments will also incorporate measurements at 0° and 180°. Plans are under way to design and build a magnetic spectrometer which will be capable of analyzing electrons with kinetic energies of up to 100 keV. Presumably, this will allow for experiments at higher x-ray energies with the attendant higher inelastically-scattered electron energies.

Research supported in part by US DOE Office of Basic Energy Sciences, Chemical Sciences Division, Contract No. DE-AC02-76CH00016.

DEVELOPMENT OF WIRE SCANNER TO DETERMINE VERTICAL PROFILES OF INTERSECTING PHOTON AND ION BEAMS.

B.M. Johnson, D. H. Lee, M. Franklin^(a), R. Hedeman^(b) (DAS, BNL)

An experimental apparatus is under development to perform crossed-beams measurements on the inner-shell photoionization of ions. The initial experiments will focus on the measurement of charge-state distributions following K-shell photoionization of Ar⁺ and Ar²⁺ ion beams. Maximum counting rates will be achieved when the ion and photon beams completely overlap at their point of intersection in the horizontal plane. To ensure this condition, the vertical profiles of each beam must be determined.

Shown below are three perspectives of a simple wire scanner which was developed to determine vertical profiles of the ion and photon beams. Note that each wire is parallel to the other two, but at 45° to both beam directions, and only one wire at a time is in the plane of the two intersecting beams.

Also shown below are typical results for several vertical scans of the focussed photon beam. A 1:1 cylindrical focussing mirror is positioned at 9.8 meters from the synchrotron radiation source point and is designed to produce a focussed image at 19.6 meters when the mirror-pitch angle is 4.4 mr. Each scan, numbered 1 through 8 is recorded at 19.2 meters down the beam line for mirror-pitch angles from 4.45 to 4.80 mr, respectively, in steps of 0.05 mr. The number of counts is linearly proportional to the current generated when photons impact the wire to produce photoelectrons, as well as secondary electrons and ions. Note that although scan number 4 (pitch angle of 4.60 mr) yields the highest peak current, the width (FWHM) of all peaks is less than 0.04 inches (1 mm).

Similar scans of the ion beam show that a vertical beam diameter of less than 1 mm can be readily achieved. By scanning first each beam with the other blocked and then both beams together, optimum overlap of the intersecting ion and photon beams is achieved.

^(a)Present address Yale University, New Haven, CT.

^(b)Present address The Hotchkiss School, Lakeville, CT.

(Research supported by the Division of Chemical Sciences, Office of Basic Energy Sciences, US Department of Energy under Contract No. DE-AC02-76CH00016.)

Jeanne Paquette (Dept. ESS, S.U.N.-Y at Stony Brook)
Synthetic calcite crystals (300-700 micrometers across) grown in the presence of Mn and Sr display non-equivalent 1014 (r type) and 0112 (f type) faces and associated growth sectors. Under cathodoluminescence (CL) the crystals show concentric, sectoral and intrasectoral zoning of Mn. The samples were mounted in Epotek on Optisil silica disks and polished down to thicknesses of 30-50 micrometers. Mn and Sr contents (500-1300ppm range) were determined from each zone and each type of growth sector by electron probe microanalysis (EPMA). SXRF spectra were collected to compare the resolution, accuracy and precision of this technique to those of EPMA, to determine if other elements were present and to correlate their distribution with that of Mn and Sr. The synchrotron beam size was approximately 7x12 micrometers. Kapton (18mm) was used as filter. CL photographs and reflected-light photographs were used to position the beam next to EPMA contamination spots and in areas of homogeneous luminescence. Carbonate standards were used to evaluate in-run beam stability and analytical precision between runs.

Core-to-edge profiles were obtained from each sector type. "Standardless" element concentrations determined along traverses within r and f sector types reveal related enrichment trends for Mn and Sr that are inversely correlated, an observation consistent with EPMA data. Traces of Zn and Pb (< 50 ppm) were also detected in the crystal cores. These two elements were not added intentionally during growth runs; rather, they appear to have been leached from the reaction vessel or were present as impurities in the reagent-grade reactants. The relative enrichment trends of Zn and Pb between sector types correlate with those of Mn and Sr, respectively. Fe was not detected and, therefore, cannot account for the inverse relationship of cathodoluminescence and Mn intensity observed between Mn-bearing sectors of the f and r types.

There were discrepancies between absolute element concentrations and elemental ratios determined by EPMA and SXRFMA. Contributing factors include variable sample thickness, interference from diffraction peaks, limited beam resolution (interaction volume probably includes different zones), and edge effects at the periphery of crystals (where the interaction volume of the beam and the sample may not be totally contained in the crystal). Standards with lower trace element concentrations (about 1000 ppm) are probably needed. These factors require further investigation before the SXRFMA can be applied as a quantitative geochemical probe of carbonate minerals. However, these preliminary results highlight its potential as a non-destructive, semi-quantitative mapping technique for metal impurities.

DYNAMIC STUDIES OF HIGH-RESOLUTION TRYPsin CRYSTAL STRUCTURES DURING CATALYSIS OBTAINED BY WHITE-BEAM LAUE X-RAY DIFFRACTION USING SYNCHROTRON RADIATION.

P.T.Singer (BNL Biology), R.P.Cary (SUNY Downstate), K.Jones (BNL DAS), W.F. Mangel (BNL Biology) and R.M.Sweet (BNL Biology)

Crystals of the metastable catalytic intermediate *p*-guanidinobenzoyl-trypsin (GBT) were subjected to X-ray diffraction analysis using white-beam ("Laue") synchrotron radiation. Diffraction data were obtained over the range 5.0 to 1.8 Å using the Laue method. Orthorhombic crystals belonging to space group P2₁2₁2₁ had with unit cell dimensions a = 63.74, b = 63.54, c = 68.93 Å. White-beam, or "Laue," diffraction data were collected from a single crystal mounted in a flow cell using Kodak DEF film; typical exposure times were 1 sec. Data were collected with the crystal equilibrated against pH 5.5 buffer. At this pH, the GBT is quite stable, with a lifetime of several days. The pH of the buffer flowing through the flow cell was raised to 8.9, and additional data sets were collected after 3 and 80 minutes. In preliminary kinetic experiments, the lifetime of GBT was measured to be ~ 1 hr under equivalent conditions.

In preliminary crystallographic studies, it was shown that such a pH jump had no effect on the structure of benzamide-inhibited trypsin, with the exception of various polar sidechains located on the surface of the protein molecule. Most notable was the Ser195 side chain in the pH 8.9 structure, which is swiveled away from His57, resulting in the loss of the H-bond between O and NE2 of the histidine.

While the models have not been refined to fit the present data, difference Fourier maps between the time zero (pH 5.5) and the first pH 8.9 (i.e. 3 min. timepoint) data sets clearly show movement of atoms near the active site of the enzyme. The most striking is the ~2 Å shift of a water molecule previously identified by this group as important in the stability of GBT. Furthermore, densities in the difference map suggest that backbone atoms may also be moving in and around the active site, in direct contrast to the results obtained in the previous study of the effect of such a pH jump on benzamide-inhibited trypsin.

This work was sponsored by the U.S. DOE. PTS was supported by an Alexander Hollaender Post-doctoral Fellowship awarded by OHER.

Laue Diffraction Studies at X26C

E. Westbrook, M. Westbrook (ANL); I. Schlichting, D. Ringe (Brandeis); K. W. Jones, P. Singer, P. Spanne, R. Sweet (BNL)

In partnership, Argonne National Laboratory's Structural Biology Center (SBC) and BNL/DAS staff are developing a Laue diffraction capability at beam line X26C where a wide energy bandwidth of x rays, directed upon a protein crystal, stimulates a significant fraction of its entire diffraction pattern in a very brief period of time. Photographic plates and other detectors are being installed to enhance data collection, speed, accuracy, and convenience of use. During 1991 crystal diffraction studies were done by members of the Brookhaven Biology Department who determined three crystal structures of trypsin-inhibitor complexes by Laue diffraction methods on X26C. In addition, members of the Brandeis University faculty have established a strong research program at the beam line and have collected Laue diffraction data for five separate projects on: cytochrome P450, carbonmonoxy-myoglobin, catalase, chymotrypsin, and glutathione synthetase.

Figure 1 is a Laue diffraction photograph of glutathione synthetase taken by the Brandeis group.

Research supported in part by US Department of Energy, Contract No. W-31-109-ENG-38 (ANL) and Contract No. DE-AC02-76CH00016 (BNL).

END

**DATE
FILMED**

7 / 21 / 92