12
5-26-92 JSD

PNL-8098
UC-940

Development and Validation of a Cleanup Method for Hydrocarbon Containing Samples for the Analysis of Semivolatile Organic Compounds

**E. W. Hoppe
R. W. Stromatt
J. A. Campbell**

**M. J. Steele
T. E. Jones**

April 1992

**Prepared for the U.S. Department Energy
under Contract DE-AC06-76RLO 1830**

**Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute**

PNL-8098

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor Battelle Memorial Institute, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST LABORATORY
operated by
BATTELLE MEMORIAL INSTITUTE
for the
UNITED STATES DEPARTMENT OF ENERGY
under Contract DE-AC06-76RLO 1830

Printed in the United States of America

Available to DOE and DOE contractors from the
Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831;
prices available from (615) 576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service,
U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

PNL--8098

DE92 013759

DEVELOPMENT AND VALIDATION OF A CLEANUP METHOD FOR
HYDROCARBON CONTAINING SAMPLES FOR THE ANALYSIS
OF SEMIVOLATILE ORGANIC COMPOUNDS

E. W. Hoppe
R. W. Stromatt
J. A. Campbell
M. J. Steele
T. E. Jones

April 1992

Prepared for
the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830

Pacific Northwest Laboratory
Richland, Washington 99352

MASTER

EXECUTIVE SUMMARY

Samples obtained from the Hanford single shell tanks (SSTs) are contaminated with normal paraffin hydrocarbon (NPH) as hydrostatic fluid from the sampling process or can be native to the tank waste. The contamination is usually high enough that a dilution of up to several orders of magnitude may be required before the sample can be analyzed by the conventional gas chromatography/mass spectrometry methodology. This can prevent detection and measurement of organic constituents that are present at lower concentration levels. To eliminate or minimize the problem, a sample cleanup method has been developed and validated and is presented in this document.

The sample is first extracted following Pacific Northwest Laboratory procedure PNL-ALO-120 or PNL-ALO-344. An aliquot of the extract is added to water, which is then made basic, and the base-neutral compounds are extracted into methylene chloride. The aqueous phase is made acidic, and the acidic compounds are extracted into methylene chloride. The base-neutral fraction is solvent exchanged into cyclohexane, and is eluted through a commercial pre-packed silica gel column. The NPH is eluted from the column with pentane, which is followed by elution of the base-neutrals with successively more polar solvents. This fraction and the acid fraction are concentrated, and are ready for analysis.

The effectiveness of this cleanup method was demonstrated for all of the U.S. Environmental Protection Agency's (EPA) semivolatile target list compounds. Both blanks and single shell tank samples spiked with all target compounds and NPH were used in this development and validation study. The NPH removal averaged 94% for samples from SST B110 and 97% for samples from tank U110. The recoveries were good for most of the target compounds and satisfactory for all, although they were lower for the more troublesome compounds. Advisory quality control limits were established for the surrogate and matrix spike compounds. As more data are obtained using this cleanup method, required QC limits will be put in place, and a better definition of cleanup effectiveness can be established.

CONTENTS

EXECUTIVE SUMMARY	iii
INTRODUCTION	1
INITIAL PNL PROCEDURE GC/MS ANALYSES	3
EPA SW-846 METHOD 3630 PROCEDURE GC/MS ANALYSES	4
FINAL PNL PROCEDURE GC/MS ANALYSES	4
EXPERIMENTAL	5
PROCEDURES	5
Preliminary Investigation	5
Initial Development and Validation Study	6
Final Development and Validation Study	8
ANALYSIS	8
RESULTS.	11
PRELIMINARY INVESTIGATION	11
INITIAL DEVELOPMENT AND VALIDATION STUDY	11
FINAL DEVELOPMENT AND VALIDATION STUDY	13
DISCUSSION	15
PRELIMINARY INVESTIGATION	15
INITIAL DEVELOPMENT AND VALIDATION STUDY	15
FINAL DEVELOPMENT AND VALIDATION STUDY	19
METHOD ADVISORY CONTROL LIMITS	20
CONCLUSIONS	23
REFERENCES	25

APPENDIX A - PNL-ALO-120, PROCEDURE FOR EXTRACTION OF SINGLE SHELL TANK SAMPLES FOR THE ANALYSIS OF SEMIVOLATILE ORGANIC COMPOUNDS	A.1
APPENDIX B - PNL-ALO-122, CLEANUP OF HYDROCARBON CONTAINING SAMPLES FOR THE ANALYSIS OF SEMIVOLATILE ORGANIC COMPOUNDS, Revision 1.0	B.1

FIGURES

1	Flow Diagram of the Initial Development and Validation Study	26
2	Simplified Schematic of the Column Cleanup Procedures	27
3	GC/FID Chromatogram of Normal Paraffin Hydrocarbon	28
4	GC/FID Chromatogram of NPH Through Silica Column Eluted with Hexane	29
5	GC/FID Chromatogram of Methylene Chloride: Hexane Elution of NPH Following Hexane Elution	30
6	GC/FID Chromatogram of the Standard Mix Containing NPH, Base Neutral Mix 2, and AHs Through Silica Column Eluted with Hexane . .	31
7	GC/FID Chromatogram of the Standard Mix Containing NPH, Base Neutral Mix 2, and PAHs Through Silica Column Eluted with Methylene Chloride: Hexane and then Methylene Chloride After Original Elution with Hexane	32

TABLES

1	Calculated Percent Recoveries for Selected Compounds by GC/FID, Preliminary Investigation	33
2	Percent Recovery Results for Reagent Blank, Initial PNL Cleanup Method	34
3	Percent Recovery Results for Method Blank, Initial PNL Cleanup Method	36
4	Percent Recovery Results for Blank Spike, Initial PNL Cleanup Method	38
5	Percent Recovery Results for Blank Spike Duplicate, Initial PNL Cleanup Method	40
6	Percent Recovery Results for Method Blank, EPA 3630 Cleanup Method	42
7	Percent Recovery Results for Blank Spike, EPA 3630 Cleanup Method	44
8	Percent Recovery Results for Blank Spike Duplicate, EPA 3630 Cleanup Method	46
9	Percent Recovery Results for Blank Spike, Initial PNL Cleanup Method, Followed by Increased Methylene Chloride Elution	48
10	Percent Recovery Results for Blank Spike with NPH, Initial PNL Cleanup Method, Followed by Increased Methylene Chloride Elution	50
11	Percent Recovery Results for Blank Spike with NPH Duplicate, Initial PNL Cleanup Method, Followed by Increased Methylene Chloride Elution	52
12	Percent Recovery Results for Blank Spike, EPA 3630 Cleanup Method, Followed by Methylene Chloride Elution	54
13	Percent Recovery Results for Blank Spike with NPH, EPA 3630 Cleanup Method, Followed by Methylene Chloride Elution	56
14	Percent Recovery Results for Blank Spike Duplicate with NPH, EPA 3630 Cleanup Method, Followed by Methylene Chloride Elution	58
15	Results for Single Shell Tank B110 Using Initial PNL Cleanup Method	60

16	Percent Recovery Results for Single Shell Tank B110 Matrix Spike, Initial PNL Cleanup Method	62
17	Percent Recovery Results for Single Shell Tank B110 Matrix Spike Duplicate, Initial PNL Cleanup Method	64
18	Results for Single Shell Tank U110 Using Initial PNL Cleanup Method	66
19	Percent Recovery Results for Single Shell Tank U110 Matrix Spike, Initial PNL Cleanup Method	68
20	Percent Recovery Results for Single Shell Tank U110 Matrix Spike Duplicate, Initial PNL Cleanup Method	70
21	Results for Single Shell Tank B110 Using Initial PNL Cleanup Method	72
22	Percent Recovery Results for Single Shell Tank B110 Matrix Spike, EPA 3630 Cleanup Method	74
23	Percent Recovery Results for Single Shell Tank B110 Matrix Spike Duplicate, EPA 3630 Cleanup Method	76
24	Comparison of the Effect of NPH in Blank Spikes on Cleanup Method Recoveries Including Methylene Chloride Elution	78
25	Percent Recovery Results for Blank Spikes, Following Soil Extraction Procedure Prior to Final PNL Cleanup Method	80
26	Percent, Accuracy, and Control Limit Data from Blank Spikes Following Soil Extraction Procedure and Prior to Final PNL Cleanup Method	82
27	Percent Recovery Results for Blank Spikes Numbers 1 and 2 Using the Final PNL Cleanup Method	84
28	Percent Recovery Results for Blank Spikes Numbers 3, 4, and 5 Using the Final PNL Cleanup Method	86
29	Percent Recovery Results for Blank Spikes Numbers 6, 7, 8 and 9 Using the Final PNL Cleanup Method	88
30	Precision Accuracy, and Control Limit Data from Blank Spikes Using the Final PNL Cleanup Method	90
31	Recovery Loss from the Use of Final PNL Cleanup Procedure	92

32	Calculation for Estimating Mean Recoveries for Compounds Spiked in Single Shell Tank Samples B110 and U110 Using the Final PNL cleanup Procedure (Mean Recovery = $P2BN - (NMS - BN -$ Blank BN) + Acid)	94
33	Estimated Precision, Accuracy and Control Limits for SST Semivolatile Organic Analysis	96

INTRODUCTION

Analysis of samples for organic compounds as part of the single shell tank (SST) chemical characterization has been compromised by the presence of large amounts of normal paraffin hydrocarbons (NPH). NPH is made up of straight chain fully saturated hydrocarbons primarily in the C12 to C14 range and used as a hydrostatic fluid to keep the bore hole open as core sample segments are removed from the waste tanks. The use of NPH as a hydrostatic fluid leads to contamination of the core segments. In addition, many tank wastes contain substantial quantities of NPH which are native to the sample itself. The presence of the high NPH concentration in the samples, whether introduced by sampling or native to the sample, can lead to significant problems in the analyses performed by gas chromatography/mass spectrometry (GC/MS). When performing GC/MS analysis, dilutions often up to several orders of magnitude were necessary to prevent overloading the analytical system. Thus, detection of many less prominent sample constituents was not possible.

Attempts to resolve the difficulties associated with high levels of NPH must be twofold. One approach is to attempt to find another suitable hydrostatic fluid that does not impact sample analyses or to identify a new sampling process that does not require a hydrostatic fluid. This would eliminate NPH as a contaminant and would allow determination of NPH native to the sample. The second approach is to develop a laboratory separations cleanup process that allows removal of native and contaminant NPH from SST samples without compromising the semivolatile organic analyses. This document deals with minimizing the NPH problems associated with the semivolatile analysis utilizing a sample cleanup process.

The approach for separation of NPH from the analytes of interest in the semivolatile fraction is based on selective retention of certain components within one phase while the other component is transferred to a second phase. Ideally, one would obtain a complete separation of NPH from the analytes of interest. However, since the separations are based on thermodynamic and kinetic processes that are dependent upon the chemical characteristics of the various compounds and surfaces of the substrates, one must balance the

complete removal of NPH against the losses of the analytes of interest. Thus, as the chemical characteristics of the analytes of interest approach those of NPH, primarily polarity in this case, the separation becomes more difficult. The approach then must be to remove adequate amounts of NPH and evaluate the recovery of the analytes of interest.

The cleanup method described in this document is employed following extraction of the waste or soil using other procedures such as those provided by the U.S. Environmental Protection Agency (EPA). Extraction procedures used at the Analytical Chemistry Laboratory at Pacific Northwest Laboratory (PNL)^(a) are derived directly from EPA Contract Laboratory Program (CLP) Statement of Work (SOW) 2/88 (e.g., PNL method PNL-ALO-344) or from EPA Solid Waste Division methods SW-846 (e.g., method PNL-ALO-120). In the PNL cleanup method, an aliquot of the sample extract obtained from a preceding extraction procedure is added to a volume of reagent water. The pH of the water is adjusted to >11, and the base-neutral fraction is extracted into methylene chloride. The pH of the aqueous phase is then adjusted to <2, and the acid fraction is extracted into methylene chloride. Following the extract's concentration and exchange into cyclohexane, the base-neutral fraction is eluted through a silica gel column (an amorphous silica with weakly acidic properties) with pentane. The pentane containing the NPH fraction is discarded, and the analytes are eluted from the column with methylene chloride, methylene chloride/acetone, and acetone. This fraction and the acid fraction from the first cleanup step are concentrated and are then ready for analysis.

The study described in this document consists of a preliminary investigation performed by the Chemical Sciences Department followed by two sets of development and validation experiments performed by the Analytical Chemistry Laboratory at PNL. The preliminary investigation qualitatively assessed the removal of NPH and the recovery of a limited set of semivolatile compounds. The first validation portion of the work presented in this document consisted of testing the proposed procedure based on the preliminary study by performing an extraction, cleanup, and analyses on blank-spikes, blank-spikes with NPH,

(a) Operated for the U.S. Department of Energy by Battelle Memorial Institute Under Contract DE-AC06-76RLO 1830.

and two samples with spike and NPH. For these studies, the spike contained all the EPA CLP SOW 2/88 semivolatile organic target compounds. The final methylene chloride/acetone and acetone column wash steps described in the preceding paragraph were not used in the first portion of the validation studies. However, a final wash step using additional methylene chloride in an attempt to improve analyte recoveries was added for selected samples. In addition, the EPA SW-846 method 3630 was performed along with the PNL method to provide a comparison. The results were favorable, but conditions were not optimum. Consequently, in the second portion of the study, the PNL procedure was slightly modified and tested on nine blank-spikes with NPH. Changes tested were the use of commercial pre-packed silica columns and use of a more polar column wash in the final step of the column elution.

The GC/MS analyses were performed on aliquots of sample residues after appropriate steps of the procedures such that analyte recoveries could be identified at each key step of the procedure. A flow diagram of the study and the sampling points for analysis during the procedure is shown in Figure 1. These analyses were performed as follows:

INITIAL PNL PROCEDURE GC/MS ANALYSES

- after the initial extraction and concentration
- after liquid/liquid base-neutral partition and concentration
- after liquid/liquid acid partition and concentration
- after solvent exchange and silica gel column cleanup steps and concentration
- after the additional methylene chloride elution step of the cleanup column and concentration when designated.

EPA SW-846 METHOD 3630 PROCEDURE GC/MS ANALYSES

- after the initial extraction and concentration
- after liquid/liquid base-neutral partition followed by liquid/liquid neutral back-extraction and concentration
- after liquid/liquid acid partition and concentration

- after solvent exchange and silica gel column cleanup steps and concentration
- after the additional methylene chloride elution step of the cleanup column and concentration when designated.

FINAL PNL PROCEDURE GC/MS ANALYSES

In the second set of experiments using the modified PNL procedure, not all of the analyses were performed as for the first set. The analyses after base-neutral partitioning was omitted because assessment of this step had already been adequately defined. Analyses performed for this work were as follows:

- after the initial extraction and concentration
- base-neutral fraction after all solvent exchange and cartridge cleanup steps and concentration
- acid fraction after the liquid/liquid partition and concentration.

EXPERIMENTAL

The following discussion describes the experiments that were conducted during the preliminary investigation, and the initial and final studies for the development and validation of a method for removing interfering hydrocarbons from samples prior to analysis for semivolatile organic compounds. Included are descriptions of the sample processing procedures and analysis operations.

PROCEDURES

The sample preparation and cleanup procedures utilized in these studies are described in this section. Included in this description are the procedures used for the preliminary investigation, the PNL and the EPA procedures in the initial development and validation studies, and the modified PNL procedure used in the final studies.

Preliminary Investigation

In the following discussion, a description of the sample preparation and cleanup used in the preliminary investigation to establish the feasibility of the method is given.

Sample Preparation: Standards were obtained from Supelco, Inc. (Bellefonte, Pennsylvania) in methylene chloride solvent. They were concentrated by nitrogen blow down and redissolved in hexane. NPH was obtained from Westinghouse Hanford Company. A 5 mL aliquot was diluted with 20 mL of hexane.

Cleanup: 25-30 grams of 100-200 mesh silica gel (Mallinckrodt) was heated at 120°C for 12 hours. After cooling, 20 mL of methylene chloride was added to form a slurry. The slurry was then added to a 2.5 cm I.D. X 25 cm glass column and eluted with an additional 50 mL of methylene chloride. The column was then washed with 100 mL of hexane. Approximately 10 grams of sodium sulfate, dried at 400°C overnight, was placed at the top of the column. After the sample, in hexane, was placed on the column, 75 mL of hexane was passed through the column and collected for NPH analysis. Then 70 mL of

methylene chloride/hexane (2:3) was passed through the column and then finally 30 mL of methylene chloride was added. The methylene chloride elutions were combined and concentrated for analyte analysis.

Initial Development and Validation Study

The sample preparation and cleanup procedures for both the initial PNL and the EPA method are described in the following discussion. The initial PNL cleanup described in this section is essentially that proposed by the preliminary investigation.

Initial extraction: The initial extraction followed the PNL method PNL-ALO-120 which is based on EPA methodologies. The PNL-ALO-120 method, as applied to SST samples, assumes a medium level extraction because of sample radioactivity and limited sample availability. The sample pH is adjusted as necessary to within 5-11. Sodium sulfate and the appropriate spiking compounds are added, and then the methylene chloride. After ultrasonic extraction, the extract is separated from the sample by filtration or centrifugation. The sample extract is aliquoted for the subsequent cleanup. A copy of method PNL-ALO-120 is included in Appendix A.

Initial PNL cleanup: An aliquot of the sample extract obtained from the initial extraction is added to a volume of reagent water. The pH of the water is adjusted to >11, and the base-neutral fraction is extracted three times into methylene chloride. The pH of the aqueous phase is then adjusted to <2, and the acid fraction is extracted three times into methylene chloride. Following the extract's concentration and exchange into hexane, the base-neutral fraction is added to a pentane pre-washed silica gel column packed as described previously in the initial study. Pentane is then added to remove the NPH, and after elution is complete, the hydrocarbon fraction is discarded. The analytes are eluted from the column with successive additions of pentane/methylene chloride and methylene chloride. For selected samples, an additional column elution of methylene chloride was employed because of observed remaining color. This fraction and the acid fraction from the first cleanup step are concentrated, and are then ready for analysis.

A simplified schematic of the cleanup procedure is shown in Figure 2. The PNL cleanup procedure is given in detail in Appendix B of this document. The procedure as it is written represents this initial development and validation study. The procedure will be modified later to represent conditions developed by the second development and validation test set, and to reflect experience gained with additional sample analyses.

EPA Method 3630 cleanup: This method does not contain specific procedures for liquid/liquid partitioning. However, the following describes a partitioning similar to that used in various EPA procedures where the intent is to isolate analytes based on their relative polarities.

An aliquot of the sample extract obtained from the initial extraction is added to a volume of reagent water. The pH of the water is adjusted to >11 , and the base-neutral fraction is extracted three times into methylene chloride. The pH of the aqueous phase is then adjusted to <2 , and the acid fraction is extracted three times into methylene chloride and is ready for concentration and subsequent analysis. The methylene chloride containing the base-neutral fraction is then added back to the aqueous phase which is still at pH <2 . The neutral fraction is then extracted three times and is concentrated, exchanged in cyclohexane, and is ready for the silica gel column cleanup procedure. The aqueous phase, following the extraction of the neutral components and still containing the bases, is made basic to >11 and extracted again three times with methylene chloride. This organic phase containing the basic compounds is concentrated and is ready for analysis.

The neutral fraction contained in 2 mL of cyclohexane is then passed through a 1.0 X 25 cm column containing about 10 grams 100-200 mesh pre-activated silica gel that was pre-washed with pentane. The column is eluted with 40 mL pentane, which is then discarded. The neutral fraction is then eluted from the column with methylene chloride/pentane. This solution is concentrated and is then ready for analysis.

Final Development and Validation Study

The cleanup procedure described in the preceding section was modified to eliminate problems observed when performing the initial development and validation studies. The modified cleanup procedure is described in the following discussion.

Initial extraction: The initial extraction was performed in the same manner as for the initial validation study.

Final PNL Cleanup: An aliquot of the initial extraction was separated into acid and base-neutral fractions employing liquid/liquid partitioning as described in the initial validation study. For the column cleanup portion of the procedure, pre-packed disposable polypropylene columns (Analytichem International-Varian) were employed. These columns are essentially the body of a 60 cc disposable syringe, approximately 2.5 cm I.D. X 15 cm long, packed with 10 grams of 100-200 mesh silica gel. The columns were activated at 120°C for 12 hours. The column void volume, about 20 mL, was determined by weight using methylene chloride to determine the proper eluting volumes to be used during the cleanup procedure. About 1 cm, approximately 4 grams, of sodium sulfate that had been conditioned at 400°C for 8 hours was added to the column. The column was then pre-eluted using 25 mL of methylene chloride followed by 40 mL of pentane, which were discarded. The base-neutral extraction residue that had been exchanged into about 2 mL hexane was added to the column. The column was then eluted with 25 mL of pentane and this eluent containing the NPH was discarded. The column was then eluted with 25 mL each of pentane/methylene chloride (1:1), methylene chloride, methylene chloride/acetone (1:1), and finally acetone. All of these fractions were combined and concentrated for analysis.

ANALYSIS

The instrumentation used in the preliminary investigation and the initial and final PNL sample cleanup studies are described in the following section. Also included are descriptions of the instrumentation calibrations, the reagents used, and the analysis procedures.

Preliminary Investigation Instrumentation: A Hewlett-Packard 5880 gas chromatograph equipped with a flame ionization detector was utilized to analyze these samples. The GC injector and detector were operated at 300°C. The oven temperature program began at 50°C for 1 minute, then ramped to 300°C at a rate of 8°C/minute, and held at 300°C for 10 minutes.

Initial and Final Development and Validation Instrumentation: The GC/MS used for this work was a Hewlett-Packard system consisting of an HP-5890 GC (using a DB-5, 30-meter by 0.25 mm ID, 0.25 micron film thickness, capillary column), an HP-5790 MS, an HP-7673 autosampler, the HP-1000 series A system controller/data acquisition/data processor system, and associated peripherals.

Calibration: The mass spectrometer mass calibration and tune parameters were confirmed with a DFTPP analysis according to EPA CLP SOW 2/88 specifications. A single level continuing calibration was performed for the CLP target list compounds each day of operation.

The calibration for the NPH was performed at three concentrations: 50, 100, and 150 µg/mL total NPH. The extracted ion used for the quantitation was mass 43. Mass 57 was used to resolve ambiguities. The sum of the peak areas for the NPH constituents was used to calculate a single response factor for the NPH, since individual constituent concentrations for the NPH used as spike material for this work were not known.

The area percentages for the NPH constituents were:

dodecane - 18%

tridecane - 48%

tetradecane - 33%

pentadecane - 1+

The internal standard 1,4-Dichlorobenzene-d₄ was used for NPH quantitation because its non-deuterated analog compound, 1,4-Dichlorobenzene, behaved similarly in the cleanup procedure to the NPH.

Reagents: All the solvents used in this work were of pesticide quality or equivalent. The target compounds for the spikes and instrument calibration were prepared from Supelco Analytical Reference Standards for EPA Contract

Laboratory Program Methods mixes. The stock spike solution was prepared by combining the Supelco mixes such that each analyte was at 200 $\mu\text{g/mL}$. The NPH was obtained from Westinghouse Hanford Company. The NPH was prepared for use by diluting to a final concentration of 6025 $\mu\text{g/mL}$ in acetone. Additional information on the reagents is provided in the procedure in Appendix B.

Apparatus and materials: These are described in the procedure in Appendix A.

Analysis procedure: Prior to analysis, 300 μL of a sample residue (maintained at $4^{\circ}\text{C} \pm 2^{\circ}\text{C}$ until ready for analysis) was transferred to an autosampler vial, and made up to 40 $\mu\text{g/mL}$ with internal standards using the Supelco internal standards mix. The GC/MS analysis followed the EPA CLP SOW 2/88 conditions. The linearity of the instrument was verified from an initial 5-point calibration. The mass calibration and tune check was performed and the instrument was calibrated using a single level continuing calibration. Each series of residue analyses was followed by analysis of a twofold dilution of the stock spike solution.

RESULTS

In the following discussion, a description of the tabulated analysis data obtained in these studies is presented. Included are data from the preliminary investigation and the initial and final development and validation studies.

PRELIMINARY INVESTIGATION

Figure 3 shows a chromatogram obtained with the GC/flame ionization detector of NPH. Figure 4 is a chromatogram of NPH eluted through a silica column using hexane, and Figure 5 is of NPH eluted through the same column with methylene chloride/hexane (2:3).

Figure 6 is a chromatogram of the hexane fraction eluted from a silica gel column containing NPH, base-neutral mix 2, and polycyclic aromatic hydrocarbon (PAH) mix. Figure 7 shows a chromatogram of the fraction collected from the same column eluted with the methylene chloride/hexane mix. The calculated percent recoveries for selected compounds are shown in Table 1.

INITIAL DEVELOPMENT AND VALIDATION STUDY

Tables 2 through 5 show the results obtained from various stages of the extraction, liquid/liquid partitions, and cleanup activities for the reagent blank, a method blank, a blank spike, and a spike duplicate utilizing the PNL cleanup. Blanks in tables indicate that no detectable quantity was found. The spikes for these studies include the complete semivolatile target compound list specified in EPA CLP-SOW 2/88 and the method surrogate compounds. The steps at which aliquots for analysis were taken are presented in the flow diagram in Figure 1. The result from each step is expressed in terms of percent recovery. Step "Re-Step 1" is the re-analysis of the residue obtained from step one. The total recovery column shown is the sum of the recovery after the cleanup stage, in this case step P3BN, and the acid partition stage, step P2A.

Tables 6 through 8 show the results obtained for the method blank, a blank spike, and a spike duplicate through the various procedural steps

utilizing the EPA SW-846 method 3630 as the column cleanup. The total recovery is obtained by the sum of steps E2A, E2B, and E3N.

Tables 9 through 11 show the results obtained for the blank spike, a blank spike with NPH added, and a NPH-added duplicate employing the PNL cleanup. Additionally, step P3BN2 was used in this portion of the study. Step P3BN2 is the elution of the column with additional methylene chloride following step P3BN. The total recovery column is the sum of results obtained for steps P2A, P3BN, and P3BN2.

Tables 12 through 14 show the results obtained for the blank spike, a blank spike with NPH added, and a NPH-added duplicate employing the EPA cleanup. Additionally, step E3N2 was used in this portion of the study. Step E3N2 is the elution of the column with additional methylene chloride following step E3N. The total recovery column is the sum of results obtained for steps E2A, E2B, E3N, and E3N2.

Tables 15 through 23 show results for SST samples B110 and U110. The NPH in these samples are native to the samples. No additional NPH was spiked into the samples for this set of experiments.

Tables 15 through 17 show the results obtained for a sample from B110, a B110 spike, and a spike duplicate utilizing the PNL cleanup procedure. Results for the un-spiked sample are expressed in terms of percent expected recovery had the sample been spiked. Ostensibly, the results for target compounds are comparable to units that are approximately 6 times $\mu\text{g/g}$. For example, a result of 100 percent as shown on the table is approximately a concentration of 600 $\mu\text{g/g}$ in the sample. For NPH, the level is 60 times $\mu\text{g/g}$. Therefore, 100 percent is approximately equal to a concentration of 6000 $\mu\text{g/g}$ of NPH.

Tables 18 through 20 show the results obtained for a sample from U110, a U110 spike, and a spike duplicate employing the PNL cleanup procedure.

Tables 21 through 23 represent results obtained for a sample from B110, a B110 spike, and a spike duplicate utilizing the EPA cleanup procedure.

Table 24 summarizes the effect of the presence of NPH on target compound recoveries for the PNL and EPA cleanup methods.

FINAL DEVELOPMENT AND VALIDATION STUDY

Table 25 shows the recoveries obtained from a blank and nine blank spike samples that have undergone the medium level soils extraction process.

Table 26 summarizes these results by tabulating the mean recoveries from each of these analyses.

Tables 27 through 29 exhibits the recoveries obtained from the blank and the nine blank spikes to which the remainder of the cleanup process was applied as described previously in the procedures portion of this report. The total recovery is based on the results obtained for the acid fraction and the cleaned base-neutral fraction. Table 30 summarizes these results by tabulating the mean recoveries from each of these analyses.

DISCUSSION

The results of the preliminary investigation, and the initial and final development and validation PNL cleanup studies, are interpreted in the following discussion. In addition, the method advisory control limits for the final PNL cleanup procedure is described.

PRELIMINARY INVESTIGATION

The results indicate that the NPH is eluted in a hexane elution of the silica gel column. Figure 4 shows a GC/FID chromatogram of NPH eluted with hexane. Figure 5 shows the same column eluted with methylene chloride/hexane (2:3). Approximately 90% of the NPH was recovered from the hexane fraction. The remaining 10% is shown to be recovered in the methylene chloride/hexane fraction.

The limited set of semivolatile analytes were seen to favor the methylene chloride/hexane fraction. Figure 6 is a chromatogram of the hexane fraction eluted from a silica gel column containing NPH, Supelco base-neutral mix 2, and Supelco PNA mix. While NPH is quite prevalent as expected, none of the analytes can be detected in the chromatogram. Figure 7 illustrates the recovery of the analytes and Table 1 provides the calculated recoveries. The poor recoveries of the more polar, oxygen-containing compounds indicated a final elution of the column using a more polar solvent was in order. A more thorough method of developing the column cleanup at this point would have been to perform the cleanup using a solvent gradient and collect fractions or to employ high-performance liquid chromatography (HPLC) methods as an analog for development purposes. However, time and funding did not allow further development as the primary task. Any further development was limited to modifications made during the validation studies.

INITIAL DEVELOPMENT AND VALIDATION STUDY

The results of the work described in this document are summarized in Tables 2 through 23. The recoveries obtained from various stages of the extraction, liquid/liquid partitions, and cleanup activities for all the

experimental work are shown for both the PNL and EPA cleanup methods on blanks and samples, spiked blanks and samples, and blank spikes with and without NPH added. The steps at which aliquots for analysis were taken are presented in the flow diagram in Figure 1. The result from each step is shown to represent the successive losses expressed in terms of percent recovery from the initial standard. The results under column "Step 1" are the results from the soil extraction procedure. "Re-Step 1" is the re-analysis of the residue obtained from step one and is presented here to indicate the reproducibility of the analysis. Columns "P2BN" and "P2A" are results from the liquid/liquid partitioning steps of the procedure and include losses from "Step 1" as well. Column "P3BN" shows the recovery obtained following the PNL column cleanup, and includes the losses from the partitioning step "P2BN" and the soils extraction stage "Step 1." The total recovery column shown is the sum of the recovery after the cleanup stage, in this case step P3BN, and the acid partition stage, step P2A. The results for the acids are obviously not affected by the column cleanup step.

Results are summarized in Tables 2 through 5 for the PNL cleanup method for a reagent blank, a method blank, a blank spike, and a blank spike duplicate. These tables show that generally the substituted aniline compounds and 4-Chlorophenyl-phenylether exhibited the greatest losses during the soils extraction step. The dienes, substituted anilines, and phenols suffered the greatest losses during the partitioning steps. The results for the PNL cleanup step seem inconsistent. One would expect good results for the PAH compounds considering experience with the rather well established EPA cleanup procedure.

Based upon these observations, it is believed that the inconsistency arises from the column not adequately being flushed during the PNL cleanup. Column void volume for the PNL cleanup is almost equal to the volume of the elution solvent. Consequently, there is little margin for error during the elution process. Much of the unexpected additional retention of analytes on the column, which results in low recoveries, is likely due to holdup on the column caused by inadequate elution volume. This problem is seen in results described later in this discussion.

The EPA cleanup method results for the method blank, a blank spike, and a blank spike duplicate are summarized in Tables 6 through 8. The total recovery values for this method are obtained from the sum of steps E2A, E2B, and E3N. In general, recoveries are fair for the phenols and PAH but not for the chlorinated compounds, neutrals, and phthalates. The partitioning steps and cleanup steps are relatively self consistent as are the total recoveries for the two blank spikes.

The results for the PNL cleanup for the blank spike, a blank spike with NPH added, and an NPH-added duplicate, along with results for step P3BN2, are summarized in Tables 9 through 11. Step P3BN2 is the elution of the column with the additional methylene chloride following step P3BN. This step was added because of the observed color in the column after step P3BN. The total recovery column is the sum of results obtained for steps P2A, P3BN, and P3BN2. Significant improvement in recovery was achieved for many of the compounds due to the added methylene chloride elution step as shown in the P3BN2 column in those tables. This improvement is directly related to the poor wash volume to column void volume ratio after step P3BN. The presence of the NPH also enhanced the recoveries as seen in the improved total recoveries of compounds for which no P3BN2 improvements were observed. Low recoveries for some of the bases, phthalates, and a few neutrals still exists. The general improvement in recoveries is also seen in the partitioning step. Improved column performance is likely a consequence of increased migration of analytes in the column due to the NPH before becoming stationary on the silica phase. Less methylene chloride mobile phase is then required to begin analyte elution from the column. The effect of the NPH on recoveries is summarized in Table 24. An average of 73% of the NPH was removed in these experiments.

The results for the EPA method, including step E3N2, for the blank spike, a blank spike with NPH added, and an NPH-added duplicate are summarized in Tables 12 through 14. Step E3N2 is the elution of the column with methylene chloride following step E3N. Step E3N2 is not part of the EPA procedure, but was added because of the color remaining in the column after step E3N. The total recovery column is the sum of results obtained for steps E2A, E2B, E3N, and E3N2. The EPA method did not benefit as much as the PNL method from

the additional methylene chloride elution step, because of the lower void volumes of the columns used in the EPA method. Recoveries were slightly improved for only Nitrobenzene, Nitrobenzene-d5, 2,6-Dinitrotoluene and N-Nitrosodiphenylamine. The effect of the NPH upon recovery was generally negative as seen in Table 24. Only the Nitrobenzenes and N-Nitrosodiphenylamine benefitted. In addition, the reproducibility of the duplicate results was worse. The average NPH removal for these duplicates was about 94%.

Results are summarized in Tables 15 through 17 for SST sample B110, B110 spike, and the spike duplicate utilizing the PNL cleanup procedure. Results for the un-spiked sample are expressed in terms of percent expected recovery had the sample been spiked. These samples did not go through the P3BN2 step, and the results are parallel to the results shown in Table 5 for the blank spike duplicate. When comparing the results in Table 4, inconsistencies are noted, especially for the Dichlorobenzenes. This is probably due to the small elution volume. As noted for the effect of NPH on the blank spikes, the NPH also generally enhances recoveries for the sample spikes. Although the reproducibility is poor for sample spike duplicates, the results are parallel. NPH removal was 82% for the unspiked sample and averaged 97% for the duplicates.

The results summarized in Tables 18 through 20 for SST sample U110, U110 spike, and the spike duplicate employing the PNL cleanup procedure also show inconsistencies. The chlorinated compounds are particularly inconsistent. Otherwise the results are similar to the blank spike results in Tables 4 and 5. Again, there is a general enhancement of recoveries that is attributable to the NPH. There was 99.8% removal of the NPH for the sample, and an average of 99.4% removal for the spiked sample duplicates.

The EPA cleanup results for SST sample B110, B110 spike, and spike duplicate are summarized in Tables 21 through 23. The duplicate sample spikes agree well, and they parallel the EPA method results for the blank spikes with a few exceptions. In general, the compound recoveries were higher for the sample spikes than for the blank spikes, with the exception of the substituted anilines, Nitrobenzenes, and N-Nitrosodiphenylamine. The recovery differences, both positive and negative, are likely due to sample matrix effects and

may have nothing to do with the presence of NPH. The NPH removal was 94% for the sample and averaged 88% for the sample spike duplicates.

Table 24 summarizes the effect of the NPH on recoveries in using the PNL and EPA cleanup methods. There is a clear improvement in recoveries of all but a very few compounds using the PNL method in the presence of NPH, while the opposite is true for the EPA method in the presence of NPH. As discussed previously, the improved PNL cleanup recoveries with NPH present is likely due to further migration of the analytes into the column, and hence the analytes are flushed more quickly from the column. The EPA cleanup, by virtue of a substantially smaller column and proportionally larger solvent elution volumes, is not expected to exhibit the same effect. The reason for diminished recoveries for the EPA cleanup due to the presence of the NPH is not clear; the effect may be an inherent problem with the method as used. Explaining these discrepancies is outside the scope of this study.

FINAL DEVELOPMENT AND VALIDATION STUDY

The inconsistent results observed in the initial PNL cleanup method are thought to be primarily a consequence of the excessively large void volume of the cleanup column. Therefore, the use of the smaller, commercial pre-packed columns were tested in the final study. In addition, the low and inconsistent recoveries observed for some of the more polar and oxygen containing compounds in the initial work suggested the need for a final column elution with a more polar solvent. To this end, elution steps with methylene chloride/acetone (1:1) and acetone were added to the method.

In Table 25, the recoveries of the spike compounds and NPH are summarized for a blank and nine blank spikes with NPH, that have undergone a medium level soils extraction procedure without sample cleanup. Table 26 provides the mean recoveries and standard deviations $STD(N-1)$ for all the compounds and NPH. In addition, the Mean Recovery $\pm 3 \times STD(N-1)$ values for the surrogate and matrix spike compounds in the CLP SOW 2/88 method for the semivolatiles are included.

Recoveries for these compounds as well as for the NPH in a blank and the nine blank spikes with NPH after completion of the cleanup procedure are

summarized in Tables 27 through 29. The mean recoveries and STD(N-1) values for the compounds and NPH are shown in Table 30 along with the mean recovery $\pm 3 \times \text{STD}(N-1)$ values for the surrogate and matrix spike compounds. Table 31 shows the effect of the cleanup procedure on the recoveries. The largest effects are noted for the diene and chlorinated compounds. The improvement for 2-Nitroaniline is probably an anomaly; there were considerable chromatography problems with the anilines in some of the samples. About 73% of the NPH was removed by the cleanup.

In comparing the results summarized in Table 31 (Mean After Cleanup column) and Tables 10 and 11 (Total Recovery columns), the improvement in performance using the "final" PNL method is striking: there are no inconsistencies, the reproducibility is better, and the recoveries are improved. There are no recoveries in the "teens" or less, as there were in the initial studies. Use of the more polar solvent for the final wash and the use of columns with the smaller void volume most certainly played an important role in this improvement.

METHOD ADVISORY CONTROL LIMITS

Application of this procedure to further SST analysis requires that meaningful advisory control limits are established. The final PNL cleanup method employing disposable cartridge columns is certainly the process of choice based upon cost, performance, and level of effort. To date, however, only the partition segment of the cleanup has been performed on actual waste samples while the cartridge cleanup has only been applied to blank spike samples. It can easily be inferred that the final PNL cartridge cleanup will perform well with actual samples. The reasons for this are several fold:

- Both the initial and final PNL cleanup methods used identical preparation processes; therefore, no differences exist prior to the actual column cleanup step.
- Data for blank spike samples, including the NPH interferant, from both cleanup methods can be compared to provide a good baseline understanding of process performance.
- Data from smaller columns have been obtained through assessment of the EPA procedure. Column capacity is not an issue.

A quantitative derivation of the advisory control limits for SST wastes can be developed. As stated previously, the results for the samples up to the actual column cleanup step may be used as there was no difference in preparation to that point. If the recovery from the blank spike up to the column cleanup step employing the initial PNL cleanup method can be subtracted from the recovery of the final PNL cleanup method, the result will be the final cleanup step's effect on recovery apart from the influence of the previous steps. The resultant difference effect can then be subtracted from the recovery results for SST samples in the initial study prior to column cleanup. This may be expressed as follows:

$$\text{Mean Recovery (SST sample)} = \text{P2BN} - (\text{NMS-BN} - \text{Blank BN}) + \text{Acid}$$

$$\text{Standard Deviation (N-1)} = [(P2BN)^2 + [ABS(NMS-BN)^2 - (\text{Blank BN})^2] + (\text{Acid})^2]^{1/2}$$

where: Mean Recovery is the percent recovery calculated for a given compound in SST samples.

Standard Deviation is calculated as N-1

ABS is the absolute value

P2BN is the average recovery obtained from SST samples for a given compound through step P2BN.

NMS-BN is the average recovery obtained from blank spike + NPH samples for a given compound prior to the initial PNL column cleanup.

Blank BN is the average recovery obtained from blank spikes + NPH after final PNL column cleanup.

Acid is the average recovery obtained from SST samples for a given compound through the acid partitioning step.

The calculation neglects any matrix effect that may occur on the cleanup column, although this is believed to be small. Table 32 shows the data used for this calculation. Table 33 provides the precision, accuracy, and calculated advisory control limits for surrogate and spiking semivolatile compounds.

CONCLUSIONS

As discussed previously, a PNL method has been developed and validated for use on samples that contain difficult levels of NPH. The sample is first extracted following PNL procedure PNL-ALO-120 or PNL-ALO-344. An aliquot of the extract is added to water, which is then made basic, and the base-neutral compounds are extracted into methylene chloride. The aqueous phase is made acidic, and the acidic compounds are extracted into methylene chloride. The base-neutral fraction is solvent exchanged into cyclohexane and eluted through a commercial pre-packed silica gel column. The NPH is eluted from the column with pentane, which is followed by elution of the base-neutrals with successively more polar solvents. This fraction and the acid fraction are concentrated and are ready for analysis.

The effectiveness of this cleanup method was demonstrated for all of the semivolatile target list compounds in the EPA CLP SOW 2/88. Both blanks and SST samples spiked with all target compounds and NPH were used in this development and validation study. The NPH removal averaged 94% for samples from SST B110 and 97% for samples from tank U110. The recoveries were good for most of the target compounds and satisfactory for all, although they were lower for the more troublesome compounds. Advisory quality control limits were established for the surrogate and matrix spike compounds. As more data are obtained using this cleanup method, required QC limits will be put in place, and a better definition of cleanup effectiveness can be established.

The NPH contamination was considered to be the overwhelming factor affecting GC/MS analysis. If other less conspicuous analytical difficulties exist, they may only now be detectable. For example, siloxanes have been detected in past analyses when large amounts of NPH were present. Following the cleanup procedure, siloxanes have still been detected during analysis of semivolatile residues. The detrimental contribution siloxanes may have, if any, has not been determined at this time.

Capillary gas chromatography as an analytical tool is well documented and offers high chromatographic resolution and sensitivity, and is specified in the current regulatory literature for semivolatile analysis. However, it

lacks the ruggedness often required when characterizing complex wastes. Complementary preparation and analyses employing more robust separation techniques such as liquid chromatography should be developed and will certainly have broad application in waste tank characterization and should be explored in the near future.

REFERENCES

EPA Contract Laboratory Program, Statement of Work for Organic Analysis, Multi-Media, Multi-Concentration, SOW No. 2-88.

EPA Contract Laboratory Program, Statement of Work for Organic Analysis, Multi-Media, Multi-Concentration, Document Number OLM01.0, March 1990.

EPA Office of Solid Waste, Test Methods for Evaluating Solid Waste, SW-846, Third Edition, 11-86.

PNL-ALO-344, Procedure for Sample Preparation of Samples for GC/MS Analysis for Semivolatile Organic Compounds, Rev. 0, PNL Technical Procedure, 4-26-91.

PNL-ALO-345, GC/MS Analysis of Extractable Semivolatile Organic Compounds, Rev. 0, PNL Technical Procedure, 8-3-89.

FIGURE 2. Simplified Schematic of the Column Cleanup Procedures

FIGURE 3. GC/FID Chromatogram of Normal Paraffin Hydrocarbon (NPH)

FIGURE 5. GC/FID Chromatogram of Methylene Chloride: Hexane Elution of NPH Following Hexane Elution

FIGURE 7. GC/FID Chromatogram of the Standard Mix Containing NPH, Base Neutral Mix 2, and PAHs Through Silica Column Eluted with Methylene Chloride: Hexane and then Methylene Chloride After Original Elution with Hexane

TABLE 1. Calculated Percent Recoveries for Selected Compounds
by GC/FID, Preliminary Investigation

Sample: Prepared Standard No. Compound Name	Percent Recovery
7) 1,3-Dichlorobenzene	70
25) 1,2,4-Trichlorobenzene	87
32) Hexachlorocyclopentadiene	88
46) 2,4-Dinitrotoluene	90
48) 4-Chlorophenyl-phenylether	0
56) Hexachlorobenzene	95
59) Anthracene	91
63) Pyrene	100
65) Butylbenzylphthalate	0
68) Chrysene	100
75) Indeno(1,2,3-cd)pyrene	84

TABLE 2. Percent Recovery Results for Reagent Blank, Initial PNL Cleanup Method

Sample: Battelle Reagent Blank						
No.	Compound Name	Step 1	Re-Step 1	P2BN	P2A	Total Recovery
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol					
3)	#Phenol-d5					
4)	Phenol					
5)	bis(2-Chloroethyl)ether					
6)	2-Chlorophenol					
7)	1,3-Dichlorobenzene					
8)	1,4-Dichlorobenzene					
9)	Benzyl alcohol					
10)	1,2-Dichlorobenzene					
11)	2-Methylphenol					
12)	bis(2-Chloroisopropyl)ether					
13)	4-Methylphenol					
14)	N-Nitroso-di-n-propylamine					
15)	Hexachloroethane					
16)	*Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5					
18)	Nitrobenzene					
19)	Isophorone					
20)	2-Nitrophenol					
21)	2,4-Dimethylphenol					
22)	Benzoic acid					
23)	bis(2-Chloroethoxy)methane					
24)	2,4-Dichlorophenol					
25)	1,2,4-Trichlorobenzene					
26)	Naphthalene					
27)	4-Chloroaniline					
28)	Hexachlorobutadiene					
29)	4-Chloro-3-methylphenol					
30)	2-Methylnaphthalene					
31)	*Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene					
33)	2,4,6-Trichlorophenol					
34)	2,4,5-Trichlorophenol					
35)	2-Chloronaphthalene					
36)	#2-Fluorobiphenyl					
37)	2-Nitroaniline					
38)	Dimethylphthalate					
39)	Acenaphthylene					
40)	2,6-Dinitrotoluene					
41)	3-Nitroaniline					

TABLE 2. (contd)

Sample: Battelle Reagent Blank							Total
No.	Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Recovery
42)	Acenaphthene						
43)	2,4-Dinitrophenol						
44)	4-Nitrophenol						
45)	Dibenzofuran						
46)	2,4-Dinitrotoluene						
47)	Diethylphthalate						
48)	4-Chlorophenyl-phenylether						
49)	Fluorene						
50)	4-Nitroaniline						
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol						
53)	N-Nitrosodiphenylamine						
54)	#2,4,6-Tribromophenol						
55)	4-Bromophenyl-phenylether						
56)	Hexachlorobenzene						
57)	Pentachlorophenol						
58)	Phenanthrene						
59)	Anthracene						
60)	DI-n-butylphthalate			0.6			
61)	Fluoranthene						
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene						
64)	#Terphenyl-d14				0.1	0.1	0.2
65)	Butylbenzylphthalate						
66)	3,3'-Dichlorobenzidine						
67)	Benzo(a)anthracene						
68)	Chrysene						
69)	Bis(2-Ethylhexyl)phthalate						
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71)	DI-n-octylphthalate					0.9	0.9
72)	Benzo(b)fluoranthene						
73)	Benzo(k)fluoranthene						
74)	Benzo(a)pyrene						
75)	Indeno(1,2,3-cd)pyrene						
76)	Dibenz(a,h)anthracene						
77)	Benzo(g,h,i)perylene						
NPH (Paraffin Hydrocarbon, C12-C14)							

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 3. Percent Recovery Results for Method Blank, Initial
PNL Cleanup Method

Sample: Battelle Method Blank						
No.	Compound Name	Step 1	Re-Step 1	P2BN	P2A	Total P3BN Recovery
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol	87.4	87.8	0.3	27.4	27.4
3)	#Phenol-d5	94.4	95.7	4.2	44.1	44.1
4)	Phenol					
5)	bis(2-Chloroethyl)ether					
6)	2-Chlorophenol					
7)	1,3-Dichlorobenzene					
8)	1,4-Dichlorobenzene					
9)	Benzyl alcohol					
10)	1,2-Dichlorobenzene					
11)	2-Methylphenol					
12)	bis(2-Chloroisopropyl)ether					
13)	4-Methylphenol					
14)	N-Nitroso-di-n-propylamine					
15)	Hexachloroethane					
16)	*Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5	78.4	79.5	65.4	0.2	0.2
18)	Nitrobenzene					
19)	Isophorone					
20)	2-Nitrophenol					
21)	2,4-Dimethylphenol					
22)	Benzoic acid					
23)	bis(2-Chloroethoxy)methane					
24)	2,4-Dichlorophenol					
25)	1,2,4-Trichlorobenzene					
26)	Naphthalene					
27)	4-Chloroaniline					
28)	Hexachlorobutadiene					
29)	4-Chloro-3-methylphenol					
30)	2-Methylnaphthalene					
31)	*Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene					
33)	2,4,6-Trichlorophenol					
34)	2,4,5-Trichlorophenol					
35)	2-Chloronaphthalene					
36)	#2-Fluorobiphenyl	77.4	84.5	68.8	0.7	46.3
37)	2-Nitroaniline					
38)	Dimethylphthalate					
39)	Acenaphthylene					
40)	2,6-Dinitrotoluene					
41)	3-Nitroaniline					

TABLE 3. (contd)

Sample: Battelle Method Blank							Total
No.	Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Recovery
42)	Acenaphthene						
43)	2,4-Dinitrophenol						
44)	4-Nitrophenol						
45)	Dibenzofuran						
46)	2,4-Dinitrotoluene						
47)	Diethylphthalate						
48)	4-Chlorophenyl-phenylether						
49)	Fluorene						
50)	4-Nitroaniline						
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol						
53)	N-Nitrosodiphenylamine						
54)	#2,4,6-Tribromophenol	107.1	114.0		15.3		15.3
55)	4-Bromophenyl-phenylether						
56)	Hexachlorobenzene						
57)	Pentachlorophenol						
58)	Phenanthrene						
59)	Anthracene						
60)	Di-n-butylphthalate						
61)	Fluoranthene						
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene						
64)	#Terphenyl-d14	87.3	97.6	73.0	0.8	66.6	67.4
65)	Butylbenzylphthalate						
66)	3,3'-Dichlorobenzidine						
67)	Benzo(a)anthracene						
68)	Chrysene						
69)	Bis(2-Ethylhexyl)phthalate						
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate						
72)	Benzo(b)fluoranthene						
73)	Benzo(k)fluoranthene						
74)	Benzo(a)pyrene						
75)	Indeno(1,2,3-cd)pyrene						
76)	Dibenz(a,h)anthracene						
77)	Benzo(g,h,i)perylene						
NPH (Paraffin Hydrocarbon, C12-C14)							

* Compound is an Internal Standard

Compound is a Surrogate Spike

**TABLE 4. Percent Recovery Results for Blank Spike, Initial
PNL Cleanup Method**

Sample: PNL Blank Spike No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	98.6	93.3	0.1	37.3		37.3
3) #Phenol-d5	95.2	91.0	1.9	40.8		40.8
4) Phenol	94.8	90.6	1.9	35.3		35.3
5) bis(2-Chloroethyl)ether	79.7	76.6	62.6	0.2		0.2
6) 2-Chlorophenol	92.3	88.3	0.4	36.7		36.7
7) 1,3-Dichlorobenzene	88.3	85.9	63.4	0.1	2.3	2.4
8) 1,4-Dichlorobenzene	88.9	87.6	61.4	0.1	2.3	2.4
9) Benzyl alcohol	92.3	85.1	62.4	0.5		0.5
10) 1,2-Dichlorobenzene	90.6	82.6	58.2	0.1	2.2	2.3
11) 2-Methylphenol	87.2	82.8	10.8	32.3		32.3
12) bis(2-Chloroisopropyl)ether	101.6	93.8	72.1	0.2		0.2
13) 4-Methylphenol	93.8	88.3	7.5	34.0		34.0
14) N-Nitroso-di-n-propylamine	96.9	90.8	70.3			
15) Hexachloroethane	89.2	85.0	59.2	0.1	1.4	1.4
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	82.7	83.9	66.3	0.1		0.1
18) Nitrobenzene	88.3	85.3	63.5	0.1		0.1
19) Isophorone	89.9	84.1	61.2	0.2		0.2
20) 2-Nitrophenol	89.7	88.4		39.6		39.6
21) 2,4-Dimethylphenol	85.8	84.7	33.8	16.4		16.4
22) Benzoic acid	89.5	85.1		25.1		25.1
23) bis(2-Chloroethoxy)methane	91.0	90.6	66.5	0.2		0.2
24) 2,4-Dichlorophenol	89.2	89.0	0.3	43.2		43.2
25) 1,2,4-Trichlorobenzene	83.0	83.0	62.6	0.2	2.9	3.1
26) Naphthalene	84.2	86.7	62.1	0.6	3.3	3.9
27) 4-Chloroaniline	44.5	31.7	18.7	0.0		0.0
28) Hexachlorobutadiene	82.4	83.7	58.9	0.2	2.3	2.4
29) 4-Chloro-3-methylphenol	94.3	91.1	6.9	39.4		39.4
30) 2-Methylnaphthalene	86.0	87.7	64.3	0.3	3.1	3.4
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	86.8	85.1	49.4		0.2	0.2
33) 2,4,6-Trichlorophenol	97.2	97.8		45.3		45.3
34) 2,4,5-Trichlorophenol	87.4	90.5		43.4		43.4
35) 2-Chloronaphthalene	84.8	87.0	67.3	0.2	2.9	3.2
36) #2-Fluorobiphenyl	83.4	88.9	69.7	0.3	3.5	3.7
37) 2-Nitroaniline	98.3	92.9	76.2	0.1		0.1
38) Dimethylphthalate	94.7	92.9	71.9	0.3		0.3
39) Acenaphthylene	88.1	90.7	67.8	0.3	3.1	3.4
40) 2,6-Dinitrotoluene	94.9	89.1	73.2			
41) 3-Nitroaniline	96.2	64.0	37.8			

TABLE 4. (contd)

Sample: PNL Blank Spike							Total
No.	Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Recovery
42)	Acenaphthene	92.4	91.9	72.8	0.3	3.7	4.0
43)	2,4-Dinitrophenol	106.8	93.7		25.7		25.7
44)	4-Nitrophenol	106.2	97.5		40.9		40.9
45)	Dibenzofuran	90.3	92.0	71.8	0.3	3.7	4.0
46)	2,4-Dinitrotoluene	93.8	94.0	72.8	0.3		0.3
47)	Diethylphthalate	99.0	95.5	79.3	0.4	0.1	0.5
48)	4-Chlorophenyl-phenylether	58.6	57.4	44.7	0.2	2.3	2.5
49)	Fluorene	91.1	91.1	72.7	0.3	3.6	3.9
50)	4-Nitroaniline	63.0	86.8	62.4	0.1		0.1
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol	100.3	96.2		37.9		37.9
53)	N-Nitrosodiphenylamine	101.4	107.6	71.8	0.2		0.2
54)	#2,4,6-Tribromophenol	98.5	105.2		45.7		45.7
55)	4-Bromophenyl-phenylether	88.2	91.4	66.4	0.3	3.6	3.9
56)	Hexachlorobenzene	92.8	93.8	69.3	0.4	3.9	4.2
57)	Pentachlorophenol	96.0	102.4		45.8		45.8
58)	Phenanthrene	92.4	93.1	72.9	0.4	4.6	5.0
59)	Anthracene	91.1	92.1	68.0	0.3	4.0	4.3
60)	DI-n-butylphthalate	92.3	93.0	72.6	0.5	0.3	0.8
61)	Fluoranthene	91.8	90.0	68.5	0.4	5.1	5.5
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene	94.9	88.7	72.0	0.4	4.8	5.1
64)	#Terphenyl-d14	81.2	81.9	67.9	0.4	5.6	6.0
65)	Butylbenzylphthalate	95.9	88.6	75.8	0.4	0.2	0.6
66)	3,3'-Dichlorobenzidine	100.0	71.4	73.5	0.1		0.1
67)	Benzo(a)anthracene	96.2	90.5	76.3	0.5	5.6	6.1
68)	Chrysene	88.7	84.4	75.2	0.5	6.2	6.7
69)	Bis(2-Ethylhexyl)phthalate	88.4	87.4	72.0	0.4		0.4
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71)	DI-n-octylphthalate	84.0	93.7	69.5	0.3	0.1	0.4
72)	Benzo(b)fluoranthene	96.4	98.0	75.3	0.4	4.5	4.9
73)	Benzo(k)fluoranthene	70.7	89.9	69.3	0.5	4.7	5.2
74)	Benzo(a)pyrene	91.3	96.4	70.0	0.4	4.1	4.5
75)	Indeno(1,2,3-cd)pyrene	107.4	104.2	76.3	0.3	3.3	3.6
76)	Dibenz(a,h)anthracene	85.9	92.4	68.2	0.3	2.8	3.1
77)	Benzo(g,h,i)perylene	99.7	98.1	78.6	0.4	2.9	3.3
NPH (Paraffin Hydrocarbon, C12-C14)							

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 5. Percent Recovery Results for Blank Spike Duplicate, Initial
PNL Cleanup Method

Sample: PNL Blank Spike Duplicate						
No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	107.3	104.9	0.2	39.4		39.4
3) #Phenol-d5	101.4	102.0	1.9	42.5		42.5
4) Phenol	101.6	103.6	2.0	39.0		39.0
5) bis(2-Chloroethyl)ether	85.1	87.4	64.8	0.1		0.1
6) 2-Chlorophenol	98.7	98.4	0.6	40.7		40.7
7) 1,3-Dichlorobenzene	100.7	97.2	65.4	0.1	34.3	34.4
8) 1,4-Dichlorobenzene	91.5	96.4	63.6	0.1	37.1	37.2
9) Benzyl alcohol	100.2	96.5	66.5			
10) 1,2-Dichlorobenzene	95.6	91.4	62.4	0.1	36.5	36.6
11) 2-Methylphenol	93.1	94.8	12.1	35.8		35.8
12) bis(2-Chloroisopropyl)ether	108.5	103.9	78.2	0.1		0.1
13) 4-Methylphenol	103.2	100.1	9.0	37.1		37.1
14) N-Nitroso-di-n-propylamine	102.8	99.7	75.2	0.1		0.1
15) Hexachloroethane	95.0	96.1	62.7	0.0	27.3	27.4
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	89.0	95.5	69.8	0.1		0.1
18) Nitrobenzene	91.9	93.4	68.1	0.1		0.1
19) Isophorone	96.1	95.7	65.1	0.1		0.1
20) 2-Nitrophenol	96.9	100.4	0.2	43.7		43.7
21) 2,4-Dimethylphenol	92.4	93.8	34.2	17.7	0.3	18.0
22) Benzoic acid	100.1	99.9		19.5		19.5
23) bis(2-Chloroethoxy)methane	95.9	99.4	69.2	0.2		0.2
24) 2,4-Dichlorophenol	93.6	102.5	0.5	46.9		46.9
25) 1,2,4-Trichlorobenzene	86.9	92.3	64.5	0.1	50.4	50.6
26) Naphthalene	91.0	96.4	65.4	0.7	51.9	52.6
27) 4-Chloroaniline	49.7	37.3	19.6	0.1		0.1
28) Hexachlorobutadiene	87.6	94.4	62.0	0.1	42.4	42.5
29) 4-Chloro-3-methylphenol	99.4	101.6	7.8	42.8		42.8
30) 2-Methylnaphthalene	92.0	97.5	68.0	0.2	60.5	60.7
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	93.3	96.3	50.4		14.0	14.0
33) 2,4,6-Trichlorophenol	102.0	108.0		48.6		48.6
34) 2,4,5-Trichlorophenol	93.8	99.9		45.9		45.9
35) 2-Chloronaphthalene	90.6	96.3	68.5	0.2	62.9	63.0
36) #2-Fluorobiphenyl	88.3	99.0	72.9	0.2	63.3	63.5
37) 2-Nitroaniline	106.7	104.1	80.2	0.1		0.1
38) Dimethylphthalate	98.9	105.5	73.6	0.2		0.2
39) Acenaphthylene	93.1	101.3	69.1	0.1	64.3	64.4
40) 2,6-Dinitrotoluene	98.9	101.2	77.2			
41) 3-Nitroaniline	100.0	71.2	40.4			

TABLE 5. (contd)

Sample: PNL Blank Spike Duplicate							Total
No.	Compound Name	Step 1	Re--Step 1	P2BN	P2A	P3BN	Recovery
42)	Acenaphthene	97.0	102.3	75.7	0.2	71.3	71.5
43)	2,4-Dinitrophenol	119.9	112.5		22.8		22.8
44)	4-Nitrophenol	97.9	87.2		44.7		44.7
45)	Dibenzofuran	97.0	101.1	74.8	0.2	71.0	71.1
46)	2,4-Dinitrotoluene	99.6	105.0	79.0	0.1		0.1
47)	Diethylphthalate	104.9	106.7	81.6	0.2	0.3	0.5
48)	4-Chlorophenyl-phenylether	61.4	63.7	45.2		47.1	47.1
49)	Fluorene	95.3	103.2	77.0	0.2	73.6	73.7
50)	4-Nitroaniline	69.6	100.0	67.4			
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol	112.4	110.3		40.8		40.8
53)	N-Nitrosodiphenylamine	106.3	116.1	75.2	0.1	0.1	0.2
54)	#2,4,6-Tribromophenol	104.6	118.1		49.0		49.0
55)	4-Bromophenyl-phenylether	93.4	99.6	67.9	0.1	76.5	76.6
56)	Hexachlorobenzene	95.0	103.6	73.0	0.1	68.9	69.0
57)	Pentachlorophenol	105.6	116.0		49.4		49.4
58)	Phenanthrene	98.9	101.6	75.6	0.2	80.7	80.9
59)	Anthracene	96.1	100.3	72.5	0.2	76.0	76.2
60)	Di-n-butylphthalate	99.7	101.8	76.3	0.2	0.0	0.3
61)	Fluoranthene	96.8	98.9	73.2	0.2	85.9	86.1
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene	91.1	100.1	81.5	0.2	91.1	91.3
64)	#Terphenyl-d14	79.4	90.8	76.9	0.3	83.6	83.9
65)	Butylbenzylphthalate	92.6	99.7	86.4	0.2		0.2
66)	3,3'-Dichlorobenzidine	99.9	78.1	83.8			
67)	Benzo(a)anthracene	93.9	100.6	87.1	0.3	91.6	91.9
68)	Chrysene	84.9	91.8	83.2	0.3	96.2	96.5
69)	Bis(2-Ethylhexyl)phthalate	86.9	96.2	81.0	0.3	1.9	2.1
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate	90.3	105.7	74.5	0.2	0.1	0.2
72)	Benzo(b)fluoranthene	112.7	115.1	77.6	0.2	76.5	76.7
73)	Benzo(k)fluoranthene	81.4	117.9	75.9	0.2	70.2	70.4
74)	Benzo(a)pyrene	99.1	109.2	74.6	0.2	76.3	76.5
75)	Indeno(1,2,3-cd)pyrene	118.1	125.0	81.4	0.1	81.0	81.1
76)	Dibenz(a,h)anthracene	93.4	104.9	72.1	0.1	75.6	75.7
77)	Benzo(g,h,i)perylene	107.4	114.1	83.8	0.2	81.7	81.8

NPH (Paraffin Hydrocarbon, C12-C14)						
-------------------------------------	--	--	--	--	--	--

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 6. Percent Recovery Results for Method Blank, EPA 3630 Cleanup Method

Sample: EPA Method Blank							
No.	Compound Name	Step 1	Re-Step 1	E2N	E2A	E2B	Total Recovery
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol	99.9	101.4	0.1	44.0		44.0
3)	#Phenol-d5	118.4	115.5	0.4	47.8		47.8
4)	Phenol						
5)	bis(2-Chloroethyl)ether						
6)	2-Chlorophenol						
7)	1,3-Dichlorobenzene						
8)	1,4-Dichlorobenzene						
9)	Benzyl alcohol						
10)	1,2-Dichlorobenzene						
11)	2-Methylphenol						
12)	bis(2-Chloroisopropyl)ether						
13)	4-Methylphenol						
14)	N-Nitroso-di-n-propylamine						
15)	Hexachloroethane						
16)	*Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5	92.1	95.9	28.9	0.7	0.2	1.8
18)	Nitrobenzene						
19)	Isophorone						
20)	2-Nitrophenol						
21)	2,4-Dimethylphenol						
22)	Benzoic acid						
23)	bis(2-Chloroethoxy)methane						
24)	2,4-Dichlorophenol						
25)	1,2,4-Trichlorobenzene						
26)	Naphthalene						
27)	4-Chloroaniline						
28)	Hexachlorobutadiene						
29)	4-Chloro-3-methylphenol						
30)	2-Methylnaphthalene						
31)	*Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene						
33)	2,4,6-Trichlorophenol						
34)	2,4,5-Trichlorophenol						
35)	2-Chloronaphthalene						
36)	#2-Fluorobiphenyl	92.3	102.7	37.7	2.2	2.1	39.0
37)	2-Nitroaniline						
38)	Dimethylphthalate						
39)	Acenaphthylene						
40)	2,6-Dinitrotoluene						
41)	3-Nitroaniline						

TABLE 6. (contd)

Sample: EPA Method Blank							Total
No.	Compound Name	Step 1	Re-Step 1	E2N	E2A	E2B	E3N Recovery
42)	Acenaphthene						
43)	2,4-Dinitrophenol						
44)	4-Nitrophenol						
45)	Dibenzofuran						
46)	2,4-Dinitrotoluene						
47)	Diethylphthalate	1.7	1.8				
48)	4-Chlorophenyl-phenylether						
49)	Fluorene						
50)	4-Nitroaniline						
51)	*Phenanthrene--d10	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol						
53)	N-Nitrosodiphenylamine						
54)	#2,4,6-Tribromophenol	110.8	117.5		78.7		78.7
55)	4-Bromophenyl-phenylether						
56)	Hexachlorobenzene						
57)	Pentachlorophenol						
58)	Phenanthrene						
59)	Anthracene						
60)	Di-n-butylphthalate						
61)	Fluoranthene						
62)	*Chrysene--d12	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene						
64)	#Terphenyl--d14	88.7	94.9	53.1	8.8	2.2	62.4 73.3
65)	Butylbenzylphthalate						
66)	3,3'-Dichlorobenzidine						
67)	Benzo(a)anthracene						
68)	Chrysene						
69)	Bis(2-Ethylhexyl)phthalate						
70)	*Perylene--d12	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate						
72)	Benzo(b)fluoranthene						
73)	Benzo(k)fluoranthene						
74)	Benzo(a)pyrene						
75)	Indeno(1,2,3-cd)pyrene						
76)	Dibenz(a,h)anthracene						
77)	Benzo(g,h,i)perylene						
NPH (Paraffin Hydrocarbon, C12-C14)							

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 7. Percent Recovery Results for Blank Spike, EPA 3630 Cleanup Method

Sample: EPA Blank Spike No. Compound Name	Step 1	Re-Step 1	E2N	E2A	E2B	E3N	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	98.6	93.3	0.2	54.5			54.5
3) #Phenol-d5	98.2	91.0	1.0	60.3	0.3		60.5
4) Phenol	94.8	90.6	0.9	55.4			55.4
5) bis(2-Chloroethyl) ether	79.7	76.6	29.0	0.4	3.8		4.2
6) 2-Chlorophenol	92.3	88.3	0.3	55.1			55.1
7) 1,3-Dichlorobenzene	88.3	85.9	32.8	0.4	4.7	1.2	6.3
8) 1,4-Dichlorobenzene	88.9	87.6	35.2	0.4	4.8	1.8	7.0
9) Benzyl alcohol	92.3	85.1	36.0	1.6	8.6		10.3
10) 1,2-Dichlorobenzene	90.6	82.6	35.2	0.6	5.0	4.5	9.9
11) 2-Methylphenol	87.2	82.8	7.5	49.4	1.2		50.6
12) bis(2-Chloroisopropyl) ether	101.6	93.8	44.1	0.7	6.0		6.6
13) 4-Methylphenol	93.8	88.3	6.2	54.4	0.8		55.2
14) N-Nitroso-dl-n-propylamine	96.9	90.8	43.1	0.9	5.5		6.4
15) Hexachloroethane	89.2	85.0	33.5	0.4	4.1	0.1	4.6
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	82.7	83.9	42.6	0.5	5.8	5.2	11.5
18) Nitrobenzene	88.3	85.3	43.5	0.6	6.4	6.7	13.7
19) Isophorone	89.9	84.1	42.9	1.3	6.2		7.5
20) 2-Nitrophenol	89.7	88.4	0.2	65.8			65.8
21) 2,4-Dimethylphenol	85.8	84.7	23.7	26.6	4.0	0.3	31.0
22) Benzoic acid	89.5	85.1		42.9			42.9
23) bis(2-Chloroethoxy)methane	91.0	90.6	47.3	1.0	7.6		8.6
24) 2,4-Dichlorophenol	89.2	89.0	0.3	64.1	0.0		64.1
25) 1,2,4-Trichlorobenzene	83.0	83.0	39.9	0.7	7.0	1.5	9.2
26) Naphthalene	84.2	86.7	42.2	0.7	7.5	38.7	46.9
27) 4-Chloroaniline	44.5	31.7	2.0	0.2	100.0		100.2
28) Hexachlorobutadiene	82.4	83.7	39.4	0.6	6.4		7.0
29) 4-Chloro-3-methylphenol	94.3	91.1	5.9	64.3	0.8		65.1
30) 2-Methylnaphthalene	86.0	87.7	46.4	0.8	8.9	44.2	53.8
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	86.8	85.1	25.7	0.2	2.3		2.5
33) 2,4,6-Trichlorophenol	97.2	97.8		67.8			67.8
34) 2,4,5-Trichlorophenol	87.4	90.5		64.3			64.3
35) 2-Chloronaphthalene	84.8	87.0	47.2	0.8	9.2	41.2	51.2
36) #2-Fluorobiphenyl	83.4	88.9	49.9	1.0	9.9	50.0	60.9
37) 2-Nitroaniline	98.3	92.9	56.4	1.1	8.0		9.1
38) Dimethylphthalate	94.7	92.9	55.2	1.8	8.1		9.9
39) Acenaphthylene	88.1	90.7	49.1	0.9	9.0	48.8	58.7
40) 2,6-Dinitrotoluene	94.9	89.1	53.5	0.9	6.7		7.6
41) 3-Nitroaniline	96.2	64.0	37.5	0.6	61.7		62.3

TABLE 7. (contd)

Sample: EPA Blank Spike							
No. Compound Name	Step 1	Re-Step 1	E2N	E2A	E2B	E3N	Total Recovery
42) Acenaphthene	92.4	91.9	54.3	1.1	10.4	54.2	65.7
43) 2,4-Dinitrophenol	106.8	93.7		58.7			58.7
44) 4-Nitrophenol	106.2	97.5		69.4			69.4
45) Dibenzofuran	90.3	92.0	51.2	1.2	10.7	55.1	66.9
46) 2,4-Dinitrotoluene	93.8	94.0	54.0	1.6	6.8		8.4
47) Diethylphthalate	99.0	95.5	59.6	2.2	8.5	0.2	10.9
48) 4-Chlorophenyl-phenylether	58.6	57.4	33.2	0.8	7.1	36.8	44.8
49) Fluorene	91.1	91.1	56.3	1.4	11.2	57.4	70.0
50) 4-Nitroaniline	63.0	66.8	45.1	0.5	11.8		12.3
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	100.3	96.2		68.7			68.7
53) N-Nitrosodiphenylamine	101.4	107.6	67.7	0.8	12.6	6.0	19.4
54) #2,4,6-Tribromophenol	98.6	105.2		77.7			77.7
55) 4-Bromophenyl-phenylether	88.2	91.4	54.7	1.5	11.2	59.5	72.2
56) Hexachlorobenzene	92.8	93.8	55.0	1.7	9.3	1.4	12.4
57) Pentachlorophenol	96.0	102.4		74.3			74.3
58) Phenanthrene	92.4	93.1	59.2	1.8	11.3	63.6	78.7
59) Anthracene	91.1	92.1	56.2	1.7	10.7	56.8	69.2
60) Di-n-butylphthalate	92.3	93.0	61.8	2.6	9.6		12.3
61) Fluoranthene	91.8	90.0	58.6	2.0	10.0	67.9	79.9
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	94.9	88.7	56.8	2.1	9.2	66.3	77.6
64) #Terphenyl-d14	81.2	81.9	53.8	2.4	9.6	61.6	73.6
65) Butylbenzylphthalate	95.9	88.6	59.8	2.7	8.9		11.5
66) 3,3'-Dichlorobenzidine	100.0	71.4	29.1		24.8		24.8
67) Benzo(a)anthracene	96.2	90.5	57.7	2.2	9.0	66.6	77.8
68) Chrysene	88.7	84.4	57.0	2.5	9.7	68.5	80.6
69) Bis(2-Ethylhexyl)phthalate	88.4	87.4	158.5	2.9	39.2	0.2	42.2
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	84.0	93.7	66.5	2.6	11.6		14.2
72) Benzo(b)fluoranthene	96.4	98.0	59.1	1.8	8.7	55.4	65.9
73) Benzo(k)fluoranthene	70.7	89.9	56.5	2.5	9.7	55.3	67.4
74) Benzo(a)pyrene	91.3	96.4	58.9	2.0	8.8	53.6	64.4
75) Indeno(1,2,3-cd)pyrene	107.4	104.2	62.3	1.9	8.6	56.6	67.1
76) Dibenzo(a,h)anthracene	85.0	92.4	54.6	1.7	7.5	53.8	62.9
77) Benzo(g,h,i)perylene	99.7	98.1	60.9	1.9	8.3	51.7	61.8
NPH (Paraffin Hydrocarbon, C12-C14)							

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 8. Percent Recovery Results for Blank Spike Duplicate, EPA 3630 Cleanup Method

Sample: EPA Blank Spike Duplicate								Total
No.	Compound Name	Step 1	Re-Step 1	E2N	E2A	E2B	E3N	Recovery
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol	107.3	104.9	0.2	50.9			50.9
3)	#Phenol-d5	101.4	102.0	1.3	54.6	0.1		54.7
4)	Phenol	101.6	103.6	1.3	52.1			52.1
5)	bis(2-Chloroethyl)ether	85.1	87.4	35.3	0.2			0.2
6)	2-Chlorophenol	98.7	98.4	0.3	51.6			51.6
7)	1,3-Dichlorobenzene	100.7	97.2	42.6	0.2	1.7	1.5	3.4
8)	1,4-Dichlorobenzene	91.5	96.4	45.1	0.2	1.6	2.3	4.1
9)	Benzyl alcohol	100.2	96.5	47.8	1.2	4.2		5.3
10)	1,2-Dichlorobenzene	95.6	91.4	45.6	0.2	1.7	5.3	7.1
11)	2-Methylphenol	93.1	94.8	9.3	45.4	0.4		45.8
12)	bis(2-Chloroisopropyl)ether	108.5	103.9	58.1	0.3	2.1		2.4
13)	4-Methylphenol	103.2	100.1	8.4	49.0	0.3		49.2
14)	N-Nitroso-di-n-propylamine	102.8	99.7	50.7	0.5	1.8		2.2
15)	Hexachloroethane	95.0	96.1	45.0	0.1	1.5		1.6
16)	*Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5	89.0	95.5	51.1	0.2	1.8	6.7	8.7
18)	Nitrobenzene	91.9	93.4	52.3	0.2	1.9	8.7	10.0
19)	Isophorone	96.1	95.7	49.6	0.8	2.0		2.8
20)	2-Nitrophenol	96.9	100.4	0.1	55.1			55.1
21)	2,4-Dimethylphenol	92.4	93.8	27.1	22.5	1.2		23.7
22)	Benzole acid	100.1	99.9		42.5			42.5
23)	bis(2-Chloroethoxy)methane	95.9	99.4	55.1	0.5	2.3		2.8
24)	2,4-Dichlorophenol	93.6	102.5	0.3	55.7			55.7
25)	1,2,4-Trichlorobenzene	86.9	92.3	47.0	0.3	2.0	2.0	4.3
26)	Naphthalene	91.0	96.4	49.9	0.8	2.2	39.5	42.6
27)	4-Chloroaniline	49.7	37.3	1.8	0.4	95.0		95.4
28)	Hexachlorobutadiene	87.6	94.4	47.0	0.3	2.0		2.3
29)	4-Chloro-3-methylphenol	99.4	101.6	6.3	55.7	0.2		55.9
30)	2-Methylnaphthalene	92.0	97.5	51.6		2.1	42.9	45.0
31)	*Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene	93.3	98.3	32.7		0.4		0.4
33)	2,4,6-Trichlorophenol	102.0	108.0		58.7			58.7
34)	2,4,5-Trichlorophenol	93.8	99.9		55.7			55.7
35)	2-Chloronaphthalene	90.6	96.3	50.3	0.5	2.1	32.7	35.3
36)	#2-Fluorobiphenyl	88.3	99.0	50.4	0.5	2.1	44.1	46.7
37)	2-Nitroaniline	106.7	104.1	59.3	0.7	1.5		2.2
38)	Dimethylphthalate	98.9	105.5	54.9	1.3	2.2		3.4
39)	Acenaphthylene	93.1	101.3	51.5	0.6	2.0	44.7	47.3
40)	2,6-Dinitrotoluene	98.9	101.2	54.5	0.6	1.2		1.8
41)	3-Nitroaniline	100.0	71.2	46.6	0.4	47.6		48.0

TABLE 8. (contd)

Sample: EPA Blank Spike Duplicate							
No. Compound Name	Step 1	Re-Step 1	E2N	E2A	E2B	E3N	Total Recovery
42) Acenaphthene	97.0	102.3	58.0	0.7	2.4	49.6	52.8
43) 2,4-Dinitrophenol	119.9	112.5		55.0			55.0
44) 4-Nitrophenol	97.9	87.2		65.6			65.6
45) Dibenzofuran	97.0	101.1	53.2	0.8	2.3	50.5	53.6
46) 2,4-Dinitrotoluene	99.6	105.0	54.8	1.2	1.2		2.3
47) Diethylphthalate	104.9	106.7	59.5	1.5	2.5	0.2	4.2
48) 4-Chlorophenyl-phenylether	61.4	63.7	34.5	0.6	1.5	33.9	36.0
49) Fluorene	95.3	103.2	57.9	1.0	2.2	51.4	54.6
50) 4-Nitroaniline	69.6	100.0	61.5	0.4	3.6		3.9
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	112.4	110.3		66.5			66.5
53) N-Nitrosodiphenylamine	106.3	116.1	66.7	0.6	2.9	7.3	10.7
54) #2,4,6-Tribromophenol	104.6	118.1		68.9			68.9
55) 4-Bromophenyl-phenylether	93.4	99.6	55.5	1.2	2.3	56.1	59.5
56) Hexachlorobenzene	95.0	103.6	57.0	1.2	2.5	0.5	4.1
57) Pentachlorophenol	105.6	116.0		67.2			67.2
58) Phenanthrene	98.9	101.6	58.5	1.4	2.5	60.6	64.5
59) Anthracene	96.1	100.3	54.8	1.3	2.3	57.0	60.6
60) Di-n-butylphthalate	99.7	101.8	62.8	1.8	3.0	0.0	4.9
61) Fluoranthene	96.8	98.9	57.4	1.5	2.5	64.9	68.8
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	91.1	100.1	54.5	1.5	2.3	67.3	71.1
64) #Terphenyl-d14	79.4	90.8	51.3	1.7	2.4	63.4	67.5
65) Butylbenzylphthalate	92.6	99.7	60.1	2.0	2.7		4.8
66) 3,3'-Dichlorobenzidine	99.9	78.1	23.0		5.2		5.2
67) Benzo(a)anthracene	93.9	100.6	55.0	1.7	2.4	66.4	70.4
68) Chrysene	84.9	91.8	55.4	1.9	2.6	72.9	77.4
69) Bis(2-Ethylhexyl)phthalate	86.9	96.2	54.6	2.0	2.7	0.2	4.9
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	90.3	105.7	59.3	2.0	2.5		4.5
72) Benzo(b)fluoranthene	112.7	115.1	55.6	1.4	2.2	54.4	58.0
73) Benzo(k)fluoranthene	81.4	117.9	62.4	1.9	2.7	55.3	59.9
74) Benzo(a)pyrene	99.1	109.2	56.2	1.5	2.3	51.7	55.5
75) Indeno(1,2,3-cd)pyrene	118.1	125.0	60.4	1.3	2.1	57.4	60.8
76) Dibenzo(a,h)anthracene	93.4	104.9	54.2	1.2	1.8	52.4	55.5
77) Benzo(g,h,i)perylene	107.4	114.1	59.2	1.3	2.2	55.1	58.5
NPH (Paraffin Hydrocarbon, C12-C14)							

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 9. Percent Recovery Results for Blank Spike, Initial PNL Cleanup Method, Followed by Increased Methylene Chloride Elution

Sample: PNL Blank Spike NPH (control)					Total
No. Compound Name	Step P2BN	P2A	P3BN	P3BN2	Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	0.2	55.9	0.0	0.0	55.9
3) #Phenol-d5	4.2	58.7			58.7
4) Phenol		54.3		0.1	54.4
5) bis(2-Chloroethyl)ether	71.9	0.5		37.7	38.2
6) 2-Chlorophenol	0.9	57.2			57.2
7) 1,3-Dichlorobenzene	61.8	0.3	34.0		34.4
8) 1,4-Dichlorobenzene	62.3	0.4	37.8		38.1
9) Benzyl alcohol	72.2	1.7			1.7
10) 1,2-Dichlorobenzene	60.9	0.4	37.6		38.0
11) 2-Methylphenol	25.0	43.3		12.0	55.3
12) bis(2-Chloroisopropyl)ether	84.3	0.6		57.9	58.5
13) 4-Methylphenol	19.9	48.4		0.3	48.6
14) N-Nitroso-di-n-propylamine	79.6	0.9			0.9
15) Hexachloroethane	59.7	0.3	15.4		15.7
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	71.5	0.5		57.4	57.8
18) Nitrobenzene	73.6	0.5		59.9	60.4
19) Isophorone	69.3	1.5			1.5
20) 2-Nitrophenol	0.4	67.9			67.9
21) 2,4-Dimethylphenol	53.4	15.3	0.3	23.1	38.7
22) Benzoic acid		52.7			52.7
23) bis(2-Chloroethoxy)methane	75.1	1.1		65.5	66.5
24) 2,4-Dichlorophenol	0.9	70.9			70.9
25) 1,2,4-Trichlorobenzene	68.6	0.7	48.5	0.0	49.2
26) Naphthalene	69.5	1.0	53.3	1.0	55.3
27) 4-Chloroaniline	102.5	0.6		1.2	1.8
28) Hexachlorobutadiene	68.8	0.7	19.9		20.6
29) 4-Chloro-3-methylphenol	18.1	59.4		2.5	61.9
30) 2-Methylnaphthalene	71.0	1.0	59.1	1.4	61.5
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	61.5	0.3	14.5		14.8
33) 2,4,6-Trichlorophenol		71.7			71.7
34) 2,4,5-Trichlorophenol		68.7			68.7
35) 2-Chloronaphthalene	72.0	1.3	61.7	0.1	63.0
36) #2-Fluorobiphenyl	74.4	1.4	59.9	3.4	64.6
37) 2-Nitroaniline	84.2	1.6		65.9	67.5
38) Dimethylphthalate	78.9	2.8			2.8
39) Acenaphthylene	74.3	1.5	60.4	2.6	64.4
40) 2,6-Dinitrotoluene	77.2			70.6	70.6
41) 3-Nitroaniline	84.5	0.4	0.1	3.2	3.8

TABLE 9. (contd)

Sample: PNL Blank Spike NPH (control)					
No. Compound Name	Step P2BN	P2A	P3BN	P3BN2	Total Recovery
42) Acenaphthene	81.5	1.9	66.5	3.1	71.6
43) 2,4-Dinitrophenol		78.9			78.9
44) 4-Nitrophenol		75.8			75.8
45) Dibenzofuran	75.8	2.1	60.7	6.9	69.7
46) 2,4-Dinitrotoluene	77.6	2.4		67.1	69.5
47) Diethylphthalate	81.6	3.2	0.2	0.0	3.5
48) 4-Chlorophenyl-phenylether	48.6	1.5	39.9	5.5	47.0
49) Fluorene	78.4	2.3	63.4	7.1	72.9
50) 4-Nitroaniline	100.0	0.9		25.2	26.1
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol		85.1			85.1
53) N-Nitrosodiphenylamine	91.2	0.8		52.7	53.5
54) #2,4,6-Tribromophenol		85.0			85.0
55) 4-Bromophenyl-phenylether	77.7	2.7	60.1	8.8	71.6
56) Hexachlorobenzene	79.6	2.7	37.2		39.9
57) Pentachlorophenol		95.2			95.2
58) Phenanthrene	78.2	2.8	66.2	6.3	75.3
59) Anthracene	76.2	2.6	63.6	4.7	70.9
60) Di-n-butylphthalate	76.7	3.5	0.0	0.0	3.5
61) Fluoranthene	76.5	2.9	63.3	7.5	73.7
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0
63) Pyrene	77.4	2.9	69.6	7.6	80.0
64) #Terphenyl-d14	72.6	3.0	65.0	9.1	77.1
65) Butylbenzylphthalate	77.5	3.3	0.0	0.0	3.3
66) 3,3'-Dichlorobenzidine	75.3				0.0
67) Benzo(a)anthracene	81.1	3.1	69.4	8.8	81.3
68) Chrysene	78.9	3.4	73.5	10.0	86.9
69) Bis(2-Ethylhexyl)phthalate	74.2	4.0	0.2	62.0	66.2
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	67.9	3.5	0.0	26.6	30.1
72) Benzo(b)fluoranthene	80.8	3.0	63.1	10.0	76.2
73) Benzo(k)fluoranthene	70.5	2.8	68.0	10.4	81.2
74) Benzo(a)pyrene	75.4	2.7	65.0	8.3	76.1
75) Indeno(1,2,3-cd)pyrene	80.4	2.8	64.4	9.7	76.9
76) Dibenz(a,h)anthracene	71.3	2.5	60.1	9.0	71.6
77) Benzo(g,h,i)perylene	80.0	2.8	63.4	9.0	75.2
NPH (Paraffin Hydrocarbon, C12-C14)		0.1			0.1

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 10. Percent Recovery Results for Blank Spike with NPH, Initial PNL Cleanup Method, Followed by Increased Methylene Chloride Elution

Sample: PNL Blank Spike + NPH					
No. Compound Name	Step P2BN	P2A	P3BN	P3BN2	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	0.4	64.7	0.0	0.0	64.7
3) #Phenol-d5	3.6	65.8			65.8
4) Phenol	2.6	59.8			59.8
5) bis(2-Chloroethyl)ether	79.3	2.2		46.2	48.4
6) 2-Chlorophenol	1.0	66.6			66.6
7) 1,3-Dichlorobenzene	69.1	1.4	46.7		48.1
8) 1,4-Dichlorobenzene	73.9	1.6	48.0		49.6
9) Benzyl alcohol	80.4	5.2			5.2
10) 1,2-Dichlorobenzene	69.9	1.6	48.3		49.9
11) 2-Methylphenol	20.7	54.0		7.9	62.0
12) bis(2-Chloroisopropyl)ether	96.6	2.6		70.1	72.7
13) 4-Methylphenol	16.0	57.3			57.3
14) N-Nitroso-di-n-propylamine	91.1	3.6			3.6
15) Hexachloroethane	69.1	1.1	46.2		47.3
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	82.4	2.4		66.9	69.3
18) Nitrobenzene	80.4	2.8		72.6	75.4
19) Isophorone	77.6	5.3			5.3
20) 2-Nitrophenol	0.4	78.9			78.9
21) 2,4-Dimethylphenol	54.0	28.0	0.3	18.8	47.0
22) Benzoic acid		61.7			61.7
23) bis(2-Chloroethoxy)methane	81.9	4.6		76.1	80.8
24) 2,4-Dichlorophenol	0.9	74.7			74.7
25) 1,2,4-Trichlorobenzene	75.0	2.9	64.6	0.1	67.6
26) Naphthalene	75.7	3.4	62.2	5.0	70.6
27) 4-Chloroaniline	24.3	0.6		1.9	2.5
28) Hexachlorobutadiene	73.7	2.5	57.7		60.3
29) 4-Chloro-3-methylphenol	14.9	69.6		1.4	71.1
30) 2-Methylnaphthalene	55.5	4.7	63.5	5.9	74.1
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	62.2	1.3	44.8		46.2
33) 2,4,6-Trichlorophenol		83.9			83.9
34) 2,4,5-Trichlorophenol		81.6			81.6
35) 2-Chloronaphthalene	79.9	5.5	72.8	0.1	78.5
36) #2-Fluorobiphenyl	82.7	6.1	69.1	9.5	84.7
37) 2-Nitroaniline	90.8	6.4		76.3	82.7
38) Dimethylphthalate	81.2	8.2			8.2
39) Acenaphthylene	79.0	6.5	67.5	6.2	80.3
40) 2,6-Dinitrotoluene	82.1	6.1		81.3	87.3
41) 3-Nitroaniline	113.3	2.9		12.9	15.8

TABLE 10. (contd)

Sample: PNL Blank Spike + NPH					Total
No. Compound Name	Step P2BN	P2A	P3BN	P3BN2	Recovery
42) Acenaphthene	88.0	7.0	75.2	8.1	90.4
43) 2,4-Dinitrophenol		83.2			83.2
44) 4-Nitrophenol		80.9			80.9
45) Dibenzofuran	85.1	7.2	70.5	12.2	89.9
46) 2,4-Dinitrotoluene	83.4	7.5		75.3	82.8
47) Diethylphthalate	94.8	8.8	0.0	0.0	8.9
48) 4-Chlorophenyl-phenylether	0.9		0.8	0.1	0.8
49) Fluorene	87.0	7.6	75.5	10.1	93.1
50) 4-Nitroaniline	154.7	3.5		59.3	62.8
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol		85.2			85.2
53) N-Nitrosodiphenylamine	92.8	5.4		63.8	69.2
54) #2,4,6-Tribromophenol		86.8			86.8
55) 4-Bromophenyl-phenylether	76.5	7.0	75.4	11.3	93.6
56) Hexachlorobenzene	78.8	7.4	89.2		96.6
57) Pentachlorophenol		91.9			91.9
58) Phenanthrene	84.7	8.0	77.2	10.3	95.6
59) Anthracene	79.9	7.4	76.2	8.6	92.1
60) Di-n-butylphthalate	82.7	8.6	0.0	0.0	8.7
61) Fluoranthene	78.0	7.9	74.5	10.5	92.9
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0
63) Pyrene	82.8	7.8	71.6	10.8	90.2
64) #Terphenyl-d14	76.5	7.3	69.1	10.2	86.6
65) Butylbenzylphthalate	86.4	8.5		0.0	8.5
66) 3,3'-Dichlorobenzidine	100.0	5.4			5.4
67) Benzo(a)anthracene	86.8	8.2	73.8	10.2	92.2
68) Chrysene	85.6	8.4	75.6	12.4	96.4
69) Bis(2-Ethylhexyl)phthalate	82.5	9.0	0.1	58.9	67.9
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	79.2	8.4	0.1	26.2	34.8
72) Benzo(b)fluoranthene	89.3	7.3	70.7	11.5	89.5
73) Benzo(k)fluoranthene	76.3	8.8	76.7	11.6	97.1
74) Benzo(a)pyrene	79.5	7.3	71.8	9.7	88.7
75) Indeno(1,2,3-cd)pyrene	88.9	7.8	74.5	10.3	92.6
76) Dibenz(a,h)anthracene	79.2	7.0	67.1	10.2	84.3
77) Benzo(g,h,i)perylene	90.8	7.9	72.5	9.8	90.3
NPH (Paraffin Hydrocarbon, C12-C14)	64.8	3.4	21.8		25.2

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 11. Percent Recovery Results for Blank Spike with NPH Duplicate, Initial PNL Cleanup Method, Followed by Increased Methylene Chloride Elution

Sample: PNL Blank Spike + NPH Duplicate					Total
No. Compound Name	Step P2BN	P2A	P3BN	P3BN2	Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	0.6	61.5	0.0	0.0	61.6
3) #Phenol-d5	4.1	66.5			66.5
4) Phenol	0.0	59.0			59.0
5) bis(2-Chloroethyl)ether	80.4	0.8		51.7	52.5
6) 2-Chlorophenol	1.2	64.3			64.3
7) 1,3-Dichlorobenzene	67.3	0.6	42.5	0.3	43.3
8) 1,4-Dichlorobenzene	76.1	0.6	44.4	0.3	45.2
9) Benzyl alcohol	79.0	2.3			2.3
10) 1,2-Dichlorobenzene	69.0	0.6	43.3	0.3	44.2
11) 2-Methylphenol	21.7	54.0			54.0
12) bis(2-Chloroisopropyl)ether	90.8	0.9		73.4	74.3
13) 4-Methylphenol	17.2	57.1			57.1
14) N-Nitroso-di-n-propylamine	86.1	1.1			1.1
15) Hexachloroethane	68.6	0.4	37.7	0.3	38.4
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	83.3	0.7		74.7	75.3
18) Nitrobenzene	79.8	0.7		73.7	74.4
19) Isophorone	76.5	1.6			1.6
20) 2-Nitrophenol	0.8	77.6			77.6
21) 2,4-Dimethylphenol	56.1	24.5	0.4	1.0	25.9
22) Benzoic acid	0.0	65.5			65.5
23) bis(2-Chloroethoxy)methane	84.0	1.3		77.6	78.9
24) 2,4-Dichlorophenol	1.3	75.2			75.2
25) 1,2,4-Trichlorobenzene	76.5	1.0	60.9	0.4	62.2
26) Naphthalene	77.2	1.0	60.4	0.5	61.9
27) 4-Chloroaniline	100.0	0.1		2.4	2.5
28) Hexachlorobutadiene	76.9	0.8	51.0	0.4	52.2
29) 4-Chloro-3-methylphenol	15.9	69.0			69.0
30) 2-Methylnaphthalene	63.4	1.2	64.4	0.4	66.1
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	76.1	0.3	37.2	0.0	37.5
33) 2,4,6-Trichlorophenol	0.0	83.9			83.9
34) 2,4,5-Trichlorophenol	0.0	81.6			81.6
35) 2-Chloronaphthalene	81.0	1.3	71.4	0.4	73.1
36) #2-Fluorobiphenyl	83.7	1.3	73.5	0.5	75.4
37) 2-Nitroaniline	93.6	1.4		86.1	87.5
38) Dimethylphthalate	86.3	2.5			2.5
39) Acenaphthylene	80.0	1.4	73.9	0.6	75.9
40) 2,6-Dinitrotoluene	87.3	1.3		90.0	91.3
41) 3-Nitroaniline	100.0	0.4			0.4

TABLE 11. (contd)

Sample: PNL Blank Spike + NPH Duplicate					Total	
No.	Compound Name	Step P2BN	P2A	P3BN	P3BN2	Recovery
42)	Acenaphthene	90.5	1.6	80.9	0.6	83.1
43)	2,4-Dinitrophenol	0.0	86.5			86.5
44)	4-Nitrophenol	0.0	83.0			83.0
45)	Dibenzofuran	85.1	1.8	79.4	1.0	82.3
46)	2,4-Dinitrotoluene	86.8	2.2		89.3	91.5
47)	Diethylphthalate	97.8	2.9		0.1	3.0
48)	4-Chlorophenyl-phenylether	0.0				0.0
49)	Fluorene	84.3	2.0	83.8	0.9	86.7
50)	4-Nitroaniline	113.8	0.7		12.3	13.0
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol	0.0	90.8			90.8
53)	N-Nitrosodiphenylamine	105.2	1.3		88.9	90.1
54)	#2,4,6-Tribromophenol	0.0	93.1			93.1
55)	4-Bromophenyl-phenylether	89.8	2.1	81.1	1.8	85.0
56)	Hexachlorobenzene	90.7	2.4	81.2	0.4	84.0
57)	Pentachlorophenol	0.0	100.9			100.9
58)	Phenanthrene	90.2	2.5	82.9	0.8	86.3
59)	Anthracene	88.0	2.4	80.5	0.8	83.7
60)	Di-n-butylphthalate	89.0	3.1	0.0	0.0	3.1
61)	Fluoranthene	86.8	2.5	79.5	1.0	83.1
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0
63)	Pyrene	86.8	2.7	80.4	1.0	84.0
64)	#Terphenyl-d14	77.6	2.8	75.4	2.7	80.8
65)	Butylbenzylphthalate	86.5	3.1		0.0	3.1
66)	3,3'-Dichlorobenzidine	86.1				0.0
67)	Benzo(a)anthracene	87.6	3.0	82.3	1.7	86.9
68)	Chrysene	85.1	3.2	87.4	1.8	92.4
69)	Bis(2-Ethylhexyl)phthalate	82.6	3.3	0.1	2.8	6.2
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate	83.0	2.7	0.0	1.0	3.7
72)	Benzo(b)fluoranthene	102.3	2.3	81.6	2.1	86.0
73)	Benzo(k)fluoranthene	108.2	2.8	81.8	2.4	87.0
74)	Benzo(a)pyrene	87.3	2.3	81.1	1.7	85.1
75)	Indeno(1,2,3-cd)pyrene	97.0	2.4	84.3	1.5	88.1
76)	Dibenz(a,h)anthracene	84.3	2.1	71.3	3.2	76.6
77)	Benzo(g,h,i)perylene	96.0	2.5	80.4	1.6	84.4
NPH (Paraffin Hydrocarbon, C12-C14)		71.1	1.0	28.3	0.3	29.6

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 12. Percent Recovery Results for Blank Spike, EPA 3630 Cleanup Method, Followed by Methylene Chloride Elution

Sample: EPA Blank Spike NPH (control)						Total
No. Compound Name	E2N	E2A	E2B	E3N	E3N2	Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	0.2	75.7				75.7
3) #Phenol-d5	3.0	73.5	0.3			73.8
4) Phenol	2.8	64.8	4.3			69.1
5) bis(2-Chloroethyl)ether	55.2	0.4			9.2	9.6
6) 2-Chlorophenol	0.7	70.6				70.6
7) 1,3-Dichlorobenzene	55.0	0.4	4.2	0.3		4.9
8) 1,4-Dichlorobenzene	54.2		4.5	0.4		4.8
9) Benzyl alcohol	54.3	1.8	7.8			9.6
10) 1,2-Dichlorobenzene	53.2	0.4	4.4	1.3		6.1
11) 2-Methylphenol	18.4	53.1	1.5			54.6
12) bis(2-Chloroisopropyl)ether	65.8	0.5	5.1		14.0	19.6
13) 4-Methylphenol	14.1	57.6	1.1			58.7
14) N-Nitroso-di-n-propylamine	62.0	0.7	4.5			5.2
15) Hexachloroethane	54.5	0.3	3.8	0.1		4.2
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	61.2	0.4	5.0	5.4	8.7	19.5
18) Nitrobenzene	61.8	0.5	5.3	7.0	9.4	22.1
19) Isophorone	59.5	1.3	5.0			6.3
20) 2-Nitrophenol	0.3	84.1				84.1
21) 2,4-Dimethylphenol	42.4	20.4	4.0	0.4		24.7
22) Benzoic acid		78.9				78.9
23) bis(2-Chloroethoxy)methane	64.9	0.9	6.2			7.1
24) 2,4-Dichlorophenol	0.6	82.3				82.3
25) 1,2,4-Trichlorobenzene	58.3	0.6	5.8	0.4		6.8
26) Naphthalene	62.3	1.2	6.2	36.4		43.8
27) 4-Chloroaniline	1.2		87.6			87.6
28) Hexachlorobutadiene	57.5	0.5	5.4	0.3		6.2
29) 4-Chloro-3-methylphenol	14.7	67.6	1.1			68.7
30) 2-Methylnaphthalene	63.0	0.8	6.6	40.2		47.7
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	43.9		1.8			1.8
33) 2,4,6-Trichlorophenol		81.3				81.3
34) 2,4,5-Trichlorophenol		74.8				74.8
35) 2-Chloronaphthalene	62.9	1.1	6.8	29.6		37.4
36) #2-Fluorobiphenyl	63.3	1.2	6.8	43.3		51.3
37) 2-Nitroaniline	69.1	1.4	5.2			6.6
38) Dimethylphthalate	66.6	2.2	5.8			7.9
39) Acenaphthylene	63.0	1.2	6.3	44.6	0.0	52.1
40) 2,6-Dinitrotoluene	65.5	1.3	4.3		15.2	20.8
41) 3-Nitroaniline	38.7		91.6	0.3		91.9

TABLE 12. (contd)

Sample: EPA Blank Spike NPH (control)						
No. Compound Name	E2N	E2A	E2B	E3N	E3N2	Total Recovery
42) Acenaphthene	68.6	1.5	7.0	48.9		57.4
43) 2,4-Dinitrophenol		96.6				96.6
44) 4-Nitrophenol		85.5				85.5
45) Dibenzofuran	66.0	1.7	7.8	49.1		58.6
46) 2,4-Dinitrotoluene	66.3	2.0	4.3		9.8	16.2
47) Diethylphthalate	70.8	2.3	5.7	0.2		8.2
48) 4-Chlorophenyl-phenylether	42.2	1.2	4.6	35.9		41.8
49) Fluorene	66.1	1.7	7.3	53.0	0.2	62.2
50) 4-Nitroaniline	52.1		9.9			9.9
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol		97.3				97.3
53) N-Nitrosodiphenylamine	62.4	0.9	7.0	6.9	15.4	30.2
54) #2,4,6-Tribromophenol		92.1				92.1
55) 4-Bromophenyl-phenylether	64.5	2.0	6.9	58.3	0.4	67.5
56) Hexachlorobenzene	64.7	2.2	6.0	0.8		8.9
57) Pentachlorophenol		88.5				88.5
58) Phenanthrene	70.7	2.2	6.9	61.7	0.1	70.9
59) Anthracene	66.2	2.1	6.6	55.8	0.1	64.5
60) Di-n-butylphthalate	72.0	2.7	6.1	0.1	0.0	8.8
61) Fluoranthene	70.5	2.3	6.2	64.0	0.3	72.7
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	73.6	2.4	5.9	61.7	0.3	70.2
64) #Terphenyl-d14	66.9	2.4	5.6	59.8	1.0	68.8
65) Butylbenzylphthalate	75.7	3.0	5.9			8.9
66) 3,3'-Dichlorobenzidine	29.4		9.8			9.8
67) Benzo(a)anthracene	74.2	2.5	5.8	61.9	0.7	70.9
68) Chrysene	76.2	2.8	6.2	66.2	0.9	76.1
69) Bis(2-Ethylhexyl)phthalate	72.8	3.0	5.9	1.2	0.1	10.2
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	68.1	2.7	5.7	1.1	0.0	9.5
72) Benzo(b)fluoranthene	73.3	2.2	5.6	57.6	0.9	66.3
73) Benzo(k)fluoranthene	68.1	2.8	6.1	56.9	0.9	66.7
74) Benzo(a)pyrene	67.8	2.3	5.4	55.7	0.6	64.0
75) Indeno(1,2,3-cd)pyrene	73.3	2.2	5.4	61.2	0.7	69.5
76) Dibenz(a,h)anthracene	65.8	2.0	4.7	58.1	0.9	65.7
77) Benzo(g,h,i)perylene	74.1	2.0	5.1	51.2	0.6	58.9
NPH (Paraffin Hydrocarbon, C12-C14)						

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 13. Percent Recovery Results for Blank Spike with NPH, EPA 3630 Cleanup Method, Followed by Methylene Chloride Elution

Sample: EPA Blank Spike + NPH						Total
No. Compound Name	E2N	E2A	E2B	E3N	E3N2	Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	1.6	68.1				68.1
3) #Phenol-d5	1.6	68.8				68.8
4) Phenol	1.6	61.2				61.2
5) bis(2-Chloroethyl)ether	44.7	0.5	34.9			35.4
6) 2-Chlorophenol	0.5	65.2				65.2
7) 1,3-Dichlorobenzene	59.3	0.5	0.8			1.3
8) 1,4-Dichlorobenzene	62.1	0.5	0.9			1.5
9) Benzyl alcohol	57.3	2.9	3.0			5.9
10) 1,2-Dichlorobenzene	58.3	0.5	1.0			1.5
11) 2-Methylphenol	10.9	55.9	0.3			56.1
12) bis(2-Chloroisopropyl)ether	74.1	0.8	1.4		4.1	6.4
13) 4-Methylphenol	7.7	58.7	0.2			58.8
14) N-Nitroso-dl-n-propylamine	64.7	1.3	1.3			2.6
15) Hexachloroethane	58.0	0.4	0.8			1.2
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	61.9	0.7	1.4	5.8	23.1	31.0
18) Nitrobenzene	61.2	0.8	1.5	7.4	24.0	33.7
19) Isophorone	56.3	2.1	1.6			3.7
20) 2-Nitrophenol		72.6				72.6
21) 2,4-Dimethylphenol	30.8	33.6	1.1	0.1		34.9
22) Benzoic acid		65.6				65.6
23) bis(2-Chloroethoxy)methane	62.8	1.7	2.6			4.3
24) 2,4-Dichlorophenol	0.4	73.0				73.0
25) 1,2,4-Trichlorobenzene	59.7	1.0	2.6			3.6
26) Naphthalene	59.7	1.2	2.6	21.3		25.1
27) 4-Chloroaniline		1.3	82.5			83.8
28) Hexachlorobutadiene	57.7	0.9	2.1			3.0
29) 4-Chloro-3-methylphenol	7.4	70.4	0.3			70.7
30) 2-Methylnaphthalene	59.2	1.9	4.4	23.9		30.2
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	50.0	0.5	0.7			1.2
33) 2,4,6-Trichlorophenol		74.7				74.7
34) 2,4,5-Trichlorophenol		68.3				68.3
35) 2-Chloronaphthalene	60.7	2.0	5.4	16.4		23.8
36) #2-Fluorobiphenyl	62.5	2.2	5.6	27.5		35.2
37) 2-Nitroaniline	68.8	2.3	3.6			5.9
38) Dimethylphthalate	65.3	3.1	3.7			6.7
39) Acenaphthylene	62.6	2.2	5.2	30.4		37.8
40) 2,6-Dinitrotoluene	64.0	1.8	2.7			4.5
41) 3-Nitroaniline	100.0		84.1			84.1

TABLE 13. (contd)

Sample: EPA Blank Spike + NPH						Total
No.	Compound Name	E2N	E2A	E2B	E3N	E3N2 Recovery
42)	Acenaphthene	66.1	2.6	5.9	35.7	44.1
43)	2,4-Dinitrophenol		86.3			86.3
44)	4-Nitrophenol		77.0			77.0
45)	Dibenzofuran	65.9	2.7	7.7	39.5	49.9
46)	2,4-Dinitrotoluene	66.3	2.7	2.7		5.4
47)	Diethylphthalate	69.8	3.2	3.4		6.6
48)	4-Chlorophenyl-phenylether				0.6	0.6
49)	Fluorene	67.3	2.8	7.3	44.9	55.0
50)	4-Nitroaniline	74.1	1.1	6.2		7.3
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol		86.9			86.9
53)	N-Nitrosodiphenylamine	66.7	1.4	6.5	12.2	40.9
54)	#2,4,6-Tribromophenol		92.6			92.6
55)	4-Bromophenyl-phenylether	65.6	3.0	8.1	47.4	58.5
56)	Hexachlorobenzene	65.1	3.0	4.4		7.4
57)	Pentachlorophenol		84.7			84.7
58)	Phenanthrene	66.9	3.3	6.9	51.5	61.7
59)	Anthracene	63.6	3.0	6.9	49.8	59.7
60)	Di-n-butylphthalate	66.6	4.8	3.4	0.0	8.2
61)	Fluoranthene	67.1	3.4	5.0	52.9	61.2
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0
63)	Pyrene	68.6	3.1	4.5	51.2	58.8
64)	#Terphenyl-d14	60.8	3.1	4.0	45.4	0.1
65)	Butylbenzylphthalate	68.1	3.5	3.1		6.6
66)	3,3'-Dichlorobenzidine	34.9	1.1	5.9		7.0
67)	Benzo(a)anthracene	69.3	3.2	4.1	52.8	60.1
68)	Chrysene	69.8	3.6	4.1	55.9	63.6
69)	Bis(2-Ethylhexyl)phthalate	62.7	3.6	2.9	0.1	6.6
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate	60.0	3.5	2.7		6.2
72)	Benzo(b)fluoranthene	65.5	3.4	3.3	50.7	57.3
73)	Benzo(k)fluoranthene	66.4	3.6	3.7	51.8	59.0
74)	Benzo(a)pyrene	63.1	3.1	3.4	50.6	57.1
75)	Indeno(1,2,3-cd)pyrene	69.1	3.2	3.1	52.3	58.5
76)	Dibenz(a,h)anthracene	59.7	2.7	2.7	46.2	51.6
77)	Benzo(g,h,i)perylene	66.8	3.1	2.9	45.3	51.2
NPH (Paraffin Hydrocarbon, C12-C14)		61.6	1.6	5.1		6.7

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 14. Percent Recovery Results for Blank Spike Duplicate with NPH,
EPA 3630 Cleanup Method, Followed by Methylene Chloride
Elution

Sample: EPA Blank Spike + NPH Duplicate						Total
No. Compound Name	E2N	E2A	E2B	E3N	E3N2	Recovery
1) *1,4--Dichlorobenzene--d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2--Fluorophenol	0.2	51.4				51.4
3) #Phenol--d5	1.5	54.9				54.9
4) Phenol	1.5	50.8				50.8
5) bis(2--Chloroethyl)ether	39.6	1.0	28.4	0.3		29.7
6) 2--Chlorophenol		51.9				51.9
7) 1,3--Dichlorobenzene	51.5	0.8	0.5	0.8		2.2
8) 1,4--Dichlorobenzene	52.2	0.9	0.7	1.0		2.6
9) Benzyl alcohol	52.8	3.0	1.6			4.6
10) 1,2--Dichlorobenzene	51.5	0.9	0.6	3.1		4.6
11) 2--Methylphenol	9.0	44.5				44.5
12) bis(2--Chloroisopropyl)ether	66.3	1.4		9.0	15.7	26.1
13) 4--Methylphenol	6.6	47.6				47.6
14) N-Nitroso--di-n--propylamine	57.2		0.6			0.6
15) Hexachloroethane	48.3	0.5	0.6			1.1
16) *Naphthalene--d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene--d5	59.0	1.2	0.4	21.7	62.6	86.0
18) Nitrobenzene	59.3	1.4	0.5	24.6	66.8	93.3
19) Isophorone	59.7	3.2	0.9			4.1
20) 2--Nitrophenol	0.2	56.6				56.6
21) 2,4--Dimethylphenol	30.7	26.0	0.4	0.1		26.6
22) Benzoic acid		22.7				22.7
23) bis(2--Chloroethoxy)methane	65.5	2.3	0.9			3.3
24) 2,4--Dichlorophenol	0.4	63.1				63.1
25) 1,2,4--Trichlorobenzene	60.6	1.7	1.1	0.9		3.7
26) Naphthalene	63.0	1.9	1.1	43.7		46.7
27) 4--Chloroaniline		3.0	71.1			74.0
28) Hexachlorobutadiene	60.3	1.6	1.3			2.9
29) 4--Chloro--3--methylphenol	6.6	58.2	0.0			58.2
30) 2--Methylnaphthalene	65.2	2.4	1.5	48.9		52.8
31) *Acenaphthene--d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	36.2	0.6	0.2			0.8
33) 2,4,6--Trichlorophenol		60.0				60.0
34) 2,4,5--Trichlorophenol		56.5				56.5
35) 2--Chloronaphthalene	63.9	2.7	1.6	29.7		33.9
36) #2--Fluorobiphenyl	66.2	2.7	1.7	51.1		55.5
37) 2--Nitroaniline	69.8	3.4	0.6			4.0
38) Dimethylphthalate	71.6	5.4	1.6			7.0
39) Acenaphthylene	67.2	3.0	1.7	52.4		57.1
40) 2,6--Dinitrotoluene	65.9	2.7	0.5	1.4		4.6
41) 3--Nitroaniline	97.4		60.6	0.1		60.7

TABLE 14. (contd)

Sample: EPA Blank Spike + NPH Duplicate						
No. Compound Name	E2N	E2A	E2B	E3N	E3N2	Total Recovery
42) Acenaphthene	70.4	3.7	2.3	57.5		63.5
43) 2,4-Dinitrophenol		33.9				33.9
44) 4-Nitrophenol		53.1				53.1
45) Dibenzofuran	69.0	4.0	2.0	55.2		61.2
46) 2,4-Dinitrotoluene	64.9	4.2	0.6			4.8
47) Diethylphthalate	72.5	5.7	1.5	0.0		7.2
48) 4-Chlorophenyl-phenylether						0.0
49) Fluorene	71.4	4.3	2.2	56.5		62.9
50) 4-Nitroaniline	73.2	2.0	1.5			3.6
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol		53.4				53.4
53) N-Nitrosodiphenylamine	79.3	4.4	2.2	40.8	111.3	158.7
54) #2,4,6-Tribromophenol		73.9				73.9
55) 4-Bromophenyl-phenylether	73.1	4.6	2.1	60.8		67.5
56) Hexachlorobenzene	74.5	5.3	2.1	0.3		7.6
57) Pentachlorophenol		63.0				63.0
58) Phenanthrene	73.8	5.6	2.4	63.6		71.6
59) Anthracene	70.9	5.2	2.1	60.0		67.3
60) Di-n-butylphthalate	76.4	6.5	1.5	0.0		8.1
61) Fluoranthene	76.5	5.9	2.0	64.7		72.6
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	70.5	5.3	1.8	64.0		71.2
64) #Terphenyl-d14	64.4	5.5	1.9	59.5	0.5	67.4
65) Butylbenzylphthalate	70.7	5.9	1.2			7.2
66) 3,3'-Dichlorobenzidine	45.0		3.2			3.2
67) Benzo(a)anthracene	71.8	5.7	1.9	64.2		71.7
68) Chrysene	73.9	6.4	1.8	67.3		75.5
69) Bis(2-Ethylhexyl)phthalate	69.6	6.2	1.3	0.1		7.6
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	70.8	6.1	1.0	0.0		7.1
72) Benzo(b)fluoranthene	72.0	5.7	1.4	58.2		65.4
73) Benzo(k)fluoranthene	76.3	6.6	1.7	58.6		66.9
74) Benzo(a)pyrene	67.2	5.7	1.4	57.0		64.0
75) Indeno(1,2,3-cd)pyrene	74.5	5.6	1.2	57.1		63.9
76) Dibenz(a,h)anthracene	67.2	4.9	1.0	51.8		57.7
77) Benzo(g,h,i)perylene	71.2	5.5	1.2	55.1		61.8
NPH (Paraffin Hydrocarbon, C12-C14)	57.6	2.5	2.3			4.8

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 15. Results for Single Shell Tank B110 Using Initial
PNL Cleanup Method

Sample: Tank B110					
No. Compound Name	Step 1	P2BN	P2A	P3BN	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	91.1	0.1	26.2		26.2
3) #Phenol-d5	103.0	1.0	32.6		32.6
4) Phenol					
5) bis(2-Chloroethyl)ether					
6) 2-Chlorophenol					
7) 1,3-Dichlorobenzene					
8) 1,4-Dichlorobenzene					
9) Benzyl alcohol					
10) 1,2-Dichlorobenzene					
11) 2-Methylphenol					
12) bis(2-Chloroisopropyl)ether					
13) 4-Methylphenol					
14) N-Nitroso-di-n-propylamine					
15) Hexachloroethane					
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	88.4	58.9	0.3		0.3
18) Nitrobenzene					
19) Isophorone					
20) 2-Nitrophenol					
21) 2,4-Dimethylphenol					
22) Benzoic acid					
23) bis(2-Chloroethoxy)methane					
24) 2,4-Dichlorophenol					
25) 1,2,4-Trichlorobenzene					
26) Naphthalene					
27) 4-Chloroaniline					
28) Hexachlorobutadiene					
29) 4-Chloro-3-methylphenol					
30) 2-Methylnaphthalene					
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene					
33) 2,4,6-Trichlorophenol					
34) 2,4,5-Trichlorophenol					
35) 2-Chloronaphthalene					
36) #2-Fluorobiphenyl	96.2	59.6	0.8	55.4	56.2
37) 2-Nitroaniline					
38) Dimethylphthalate					
39) Acenaphthylene					
40) 2,6-Dinitrotoluene					
41) 3-Nitroaniline					

TABLE 15. (contd)

Sample: Tank B110					
No.	Compound Name	Step 1	P2BN	P2A	Total Recovery
42)	Acenaphthene				
43)	2,4-Dinitrophenol				
44)	4-Nitrophenol				
45)	Dibenzofuran				
46)	2,4-Dinitrotoluene				
47)	Diethylphthalate				
48)	4-Chlorophenyl-phenylether				
49)	Fluorene				
50)	4-Nitroaniline				
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol				
53)	N-Nitrosodiphenylamine				
54)	#2,4,6-Tribromophenol	103.9		54.1	54.1
55)	4-Bromophenyl-phenylether				
56)	Hexachlorobenzene				
57)	Pentachlorophenol				
58)	Phenanthrene				
59)	Anthracene				
60)	Di-n-butylphthalate				
61)	Fluoranthene				
62)	*Chrysene-d12	100.0	100.0	100.0	100.0
63)	Pyrene				
64)	#Terphenyl-d14	98.7	70.3	2.3	70.3
65)	Butylbenzylphthalate				
66)	3,3'-Dichlorobenzidine				
67)	Benzo(a)anthracene				
68)	Chrysene				
69)	Bis(2-Ethylhexyl)phthalate		1.9		1.0
70)	*Perylene-d12	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate				
72)	Benzo(b)fluoranthene				
73)	Benzo(k)fluoranthene				
74)	Benzo(a)pyrene				
75)	Indeno(1,2,3-cd)pyrene				
76)	Dibenz(a,h)anthracene				
77)	Benzo(g,h,i)perylene				
NPH (Paraffin Hydrocarbon, C12-C14)		133.0	71.4	1.2	16.8
					18.0

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 16. Percent Recovery Results for Single Shell Tank B110 Matrix Spike, Initial PNL Cleanup Method

Sample: Tank B110 Matrix Spike					
No. Compound Name	Step 1	P2BN	P2A	P3BN	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	104.3	0.4	42.6	0.0	42.6
3) #Phenol-d5	118.5	3.6	45.5		45.5
4) Phenol	108.6	3.1	39.9		39.9
5) bis(2-Chloroethyl)ether	110.3	59.6	0.3		0.3
6) 2-Chlorophenol	100.4	0.8	43.1		43.1
7) 1,3-Dichlorobenzene	97.9	52.3	0.2	31.9	32.1
8) 1,4-Dichlorobenzene	98.2	51.8	0.2	35.6	35.8
9) Benzyl alcohol	101.5	60.5	0.8		0.8
10) 1,2-Dichlorobenzene	98.6	52.2	0.2	36.7	36.9
11) 2-Methylphenol	99.9	20.3	27.8		27.8
12) bis(2-Chloroisopropyl)ether	106.0	68.4	0.3		0.3
13) 4-Methylphenol	105.2	15.9	31.8		31.8
14) N-Nitroso-di-n-propylamine	99.2	65.4	0.3		0.3
15) Hexachloroethane	97.8	52.4	0.2	13.5	13.6
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	106.4	60.1	0.2	1.0	1.2
18) Nitrobenzene	106.2	56.7	0.5	1.4	1.9
19) Isophorone	104.0	56.5	0.4		0.4
20) 2-Nitrophenol	106.2	0.3	50.6		50.6
21) 2,4-Dimethylphenol	99.0	40.2	0.9	0.2	1.1
22) Benzoic acid	75.8		39.7		39.7
23) bis(2-Chloroethoxy)methane	114.5	61.0	0.4		0.4
24) 2,4-Dichlorophenol	111.7	0.7	52.9		52.9
25) 1,2,4-Trichlorobenzene	109.5	55.8	0.3	46.9	47.2
26) Naphthalene	109.9	54.9	0.4	52.4	52.8
27) 4-Chloroaniline	136.9	84.6			0.0
28) Hexachlorobutadiene	102.9	56.9	0.3	18.8	19.1
29) 4-Chloro-3-methylphenol	113.5	15.3	42.6		42.6
30) 2-Methylnaphthalene	93.8	41.0	0.5	58.2	58.6
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	92.0	50.2	0.1	16.4	16.5
33) 2,4,6-Trichlorophenol	106.3		53.3		53.3
34) 2,4,5-Trichlorophenol	99.9		51.8		51.8
35) 2-Chloronaphthalene	107.4	55.3	0.4	61.1	61.5
36) #2-Fluorobiphenyl	107.4	58.7	0.4	61.8	62.2
37) 2-Nitroaniline	103.0	66.6	0.4		0.4
38) Dimethylphthalate	109.3	61.7	0.6		0.6
39) Acenaphthylene	100.4	53.1	0.4	60.9	61.3
40) 2,6-Dinitrotoluene	99.2	63.5	0.3		0.3
41) 3-Nitroaniline	67.5	73.2			0.0

TABLE 16. (contd)

Sample: Tank B110 Matrix Spike

No. Compound Name	Step 1	P2BN	P2A	P3BN	Total Recovery
42) Acenaphthene	115.1	61.5	0.5	69.1	69.5
43) 2,4-Dinitrophenol	52.7		58.5		58.5
44) 4-Nitrophenol	96.9		55.5		55.5
45) Dibenzofuran	107.6	62.2	0.5	66.4	67.0
46) 2,4-Dinitrotoluene	101.4	64.0	0.5		0.5
47) Diethylphthalate	111.1	66.0	0.7	0.3	0.9
48) 4-Chlorophenyl-phenylether	69.0	36.7	0.3	45.0	45.3
49) Fluorene	108.4	61.7	0.6	71.0	71.6
50) 4-Nitroaniline	151.1	114.4	0.2		0.2
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	78.7		59.3		59.3
53) N-Nitrosodiphenylamine	129.2	63.7	0.4	1.9	2.3
54) #2,4,6-Tribromophenol	102.6		55.1		55.1
55) 4-Bromophenyl-phenylether	105.3	57.0	0.6	71.3	71.9
56) Hexachlorobenzene	103.7	60.7	0.6	39.8	40.4
57) Pentachlorophenol	89.2		69.8		69.8
58) Phenanthrene	109.9	62.2	0.7	74.1	74.7
59) Anthracene	103.1	58.1	0.6	70.7	71.3
60) Di-n-butylphthalate	107.7	63.4	0.7	0.0	0.7
61) Fluoranthene	102.6	59.8	0.6	73.9	74.5
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0
63) Pyrene	107.3	59.8	0.6	75.0	75.6
64) #Terphenyl-d14	98.2	56.9	0.8	73.9	74.7
65) Butylbenzylphthalate	112.9	61.1	0.6		0.6
66) 3,3'-Dichlorobenzidine	78.2	25.6			0.0
67) Benzo(a)anthracene	108.0	60.7	0.7	74.2	74.9
68) Chrysene	114.4	60.5	0.7	78.7	79.4
69) Bis(2-Ethylhexyl)phthalate	116.6	56.4	0.7	0.2	0.9
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	110.3	54.7	0.6	0.1	0.7
72) Benzo(b)fluoranthene	108.2	63.2	0.6	74.7	75.3
73) Benzo(k)fluoranthene	103.2	48.5	0.7	77.5	78.2
74) Benzo(a)pyrene	102.5	55.1	0.6	73.1	73.7
75) Indeno(1,2,3-cd)pyrene	107.5	62.0	0.6	75.7	76.3
76) Dibenz(a,h)anthracene	97.5	56.1	0.5	67.9	68.4
77) Benzo(g,h,i)perylene	103.6	60.4	0.6	72.3	72.8
NPH (Paraffin Hydrocarbon, C12-C14)	168.8	74.7	0.6	4.0	4.6

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 17. Percent Recovery Results for Single Shell Tank B110 Matrix Spike Duplicate, Initial PNL Cleanup Method

Sample: Tank B110 Matrix Spike Duplicate					Total
No. Compound Name	Step 1	P2BN	P2A	P3BN	Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	106.3	0.5	57.3	0.0	57.3
3) #Phenol-d5	120.1	4.4	61.7		61.7
4) Phenol	111.8	4.3	57.6		57.6
5) bis(2-Chloroethyl)ether	111.5	77.1	0.6		0.6
6) 2-Chlorophenol	101.8	1.0	58.7		58.7
7) 1,3-Dichlorobenzene	101.3	68.3	0.4	45.0	45.5
8) 1,4-Dichlorobenzene	101.0	69.5	0.4	47.8	48.2
9) Benzyl alcohol	101.4	79.7	1.9		1.9
10) 1,2-Dichlorobenzene	102.0	69.9	0.4	47.7	48.1
11) 2-Methylphenol	98.7	24.5	44.3		44.3
12) bis(2-Chloroisopropyl)ether	115.7	98.5	0.7		0.7
13) 4-Methylphenol	102.3	19.4	50.8		50.8
14) N-Nitroso-di-n-propylamine	103.2	91.7	0.8		0.8
15) Hexachloroethane	101.8	69.5	0.3	17.8	18.2
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	104.2	81.4	0.5		0.5
18) Nitrobenzene	107.9	81.0	0.6		0.6
19) Isophorone	107.1	77.3	1.2		1.2
20) 2-Nitrophenol	102.9	0.4	71.0		71.0
21) 2,4-Dimethylphenol	92.7	49.7	9.6	0.2	9.8
22) Benzoic acid	83.1		56.6		56.6
23) bis(2-Chloroethoxy)methane	118.3	84.8	1.1		1.1
24) 2,4-Dichlorophenol	112.7	0.9	73.2		73.2
25) 1,2,4-Trichlorobenzene	110.7	74.5	0.7	60.1	60.8
26) Naphthalene	112.1	74.4	1.1	63.2	64.3
27) 4-Chloroaniline	127.9	100.0	0.1		0.1
28) Hexachlorobutadiene	108.8	77.3	0.6	25.5	26.2
29) 4-Chloro-3-methylphenol	114.6	18.8	63.5		63.5
30) 2-Methylnaphthalene	97.2	54.6	1.1	61.5	62.6
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	83.7	62.9	0.3	13.3	13.5
33) 2,4,6-Trichlorophenol	104.7		78.3		78.3
34) 2,4,5-Trichlorophenol	96.6		75.7		75.7
35) 2-Chloronaphthalene	104.6	77.0	1.1	67.1	68.2
36) #2-Fluorobiphenyl	106.1	78.0	1.1	66.3	67.5
37) 2-Nitroaniline	98.8	95.9	1.4		1.4
38) Dimethylphthalate	104.6	83.2	2.1		2.1
39) Acenaphthylene	94.9	69.9	1.1	62.8	63.9
40) 2,6-Dinitrotoluene	95.0	85.8	1.1		1.1
41) 3-Nitroaniline	61.9	100.0	0.6	0.1	0.7

TABLE 17. (cont'd)

Sample: Tank B110 Matrix Spike Duplicate					Total	
No.	Compound Name	Step 1	P2BN	P2A	P3BN	Recovery
42)	Acenaphthene	110.6	86.1	1.4	72.6	74.1
43)	2,4-Dinitrophenol	58.2		77.2		77.2
44)	4-Nitrophenol	88.6		82.1		82.1
45)	Dibenzofuran	106.6	84.4	1.6	68.4	70.0
46)	2,4-Dinitrotoluene	93.5	87.8	1.8		1.8
47)	Diethylphthalate	106.8	90.6	2.4	0.4	2.7
48)	4-Chlorophenyl-phenylether	65.9	49.3	1.1	45.4	46.5
49)	Fluorene	105.9	85.3	1.8	72.0	73.8
50)	4-Nitroaniline	134.1	156.0	1.7		1.7
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol	77.6		85.5		85.5
53)	N-Nitrosodiphenylamine	126.9	87.8	1.4		1.4
54)	#2,4,6-Tribromophenol	98.3		81.6		81.6
55)	4-Bromophenyl-phenylether	105.5	78.6	1.9	72.2	74.0
56)	Hexachlorobenzene	100.5	85.8	2.0	43.5	45.5
57)	Pentachlorophenol	87.3		108.7		108.7
58)	Phenanthrene	110.1	85.2	2.0	78.2	80.3
59)	Anthracene	100.5	77.8	1.6	69.9	71.5
60)	Di-n-butylphthalate	107.5	85.5	2.6	1.0	3.6
61)	Fluoranthene	103.6	82.3	2.1	79.4	81.5
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0
63)	Pyrene	104.3	80.1	2.0	83.0	84.9
64)	#Terphenyl-d14	96.4	73.1	2.2	76.5	78.6
65)	Butylbenzylphthalate	105.1	80.8	2.4	0.5	2.9
66)	3,3'-Dichlorobenzidine	66.2	100.0			0.0
67)	Benzo(a)anthracene	104.4	81.5	2.1	83.9	85.9
68)	Chrysene	112.7	82.6	2.3	85.0	87.3
69)	Bis(2-Ethylhexyl)phthalate	110.1	74.5	2.5		2.5
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate	109.6	77.8	2.5	13.0	15.5
72)	Benzo(b)fluoranthene	105.1	87.3	2.1	73.6	75.7
73)	Benzo(k)fluoranthene	102.3	87.5	2.3	58.3	60.7
74)	Benzo(a)pyrene	98.3	76.5	1.7	63.3	64.9
75)	Indeno(1,2,3-cd)pyrene	105.4	87.3	2.1	77.4	79.5
76)	Dibenz(a,h)anthracene	99.6	78.8	1.9	66.7	68.6
77)	Benzo(g,h,i)perylene	106.9	85.7	2.0	74.8	76.8
NPH (Paraffin Hydrocarbon, C12-C14)		192.7	104.3	1.5		1.5

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 18. Results for Single Shell Tank U110 Using Initial PNL Cleanup Method

Sample: Tank U110							Total
No.	Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Recovery
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol	94.7	98.3		56.0		56.0
3)	#Phenol-d5	109.9	106.2	1.0	62.7		62.7
4)	Phenol						
5)	bis(2-Chloroethyl)ether						
6)	2-Chlorophenol						
7)	1,3-Dichlorobenzene						
8)	1,4-Dichlorobenzene						
9)	Benzyl alcohol						
10)	1,2-Dichlorobenzene						
11)	2-Methylphenol						
12)	bis(2-Chloroisopropyl)ether						
13)	4-Methylphenol						
14)	N-Nitroso-di-n-propylamine						
15)	Hexachloroethane						
16)	*Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5	89.7	89.5	58.5	0.2	3.7	3.9
18)	Nitrobenzene						
19)	Isophorone						
20)	2-Nitrophenol						
21)	2,4-Dimethylphenol						
22)	Benzic acid						
23)	bis(2-Chloroethoxy)methane						
24)	2,4-Dichlorophenol						
25)	1,2,4-Trichlorobenzene						
26)	Naphthalene						
27)	4-Chloroaniline						
28)	Hexachlorobutadiene						
29)	4-Chloro-3-methylphenol						
30)	2-Methylnaphthalene						
31)	*Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene						
33)	2,4,6-Trichlorophenol						
34)	2,4,5-Trichlorophenol						
35)	2-Chloronaphthalene						
36)	#2-Fluorobiphenyl	95.4	89.2	60.7	0.5	57.2	57.7
37)	2-Nitroaniline						
38)	Dimethylphthalate						
39)	Acenaphthylene						
40)	2,6-Dinitrotoluene						
41)	3-Nitroaniline						

TABLE 18. (contd)

Sample: Tank U110						
No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Total Recovery
42) Acenaphthene						
43) 2,4-Dinitrophenol						
44) 4-Nitrophenol						
45) Dibenzofuran						
46) 2,4-Dinitrotoluene						
47) Diethylphthalate						
48) 4-Chlorophenyl-phenylether						
49) Fluorene						
50) 4-Nitroaniline						
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol						
53) N-Nitrosodiphenylamine						
54) #2,4,6-Tribromophenol	109.9	101.9		69.9		69.9
55) 4-Bromophenyl-phenylether						
56) Hexachlorobenzene						
57) Pentachlorophenol						
58) Phenanthrene						
59) Anthracene						
60) Di-n-butylphthalate						
61) Fluoranthene						
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene						
64) #Terphenyl-d14	104.4	93.2	74.3	1.0	83.1	84.1
65) Butylbenzylphthalate						
66) 3,3'-Dichlorobenzidine						
67) Benzo(a)anthracene						
68) Chrysene						
69) Bis(2-Ethylhexyl)phthalate			0.7			
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate						
72) Benzo(b)fluoranthene						
73) Benzo(k)fluoranthene						
74) Benzo(a)pyrene						
75) Indeno(1,2,3-cd)pyrene						
76) Dibenz(a,h)anthracene						
77) Benzo(g,h,i)perylene						
NPH (Paraffin Hydrocarbon, C12-C14)	10.5	12.1	6.5		0.2	0.2

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 19. Percent Recovery Results for Single Shell Tank U110 Matrix Spike, Initial PNL Cleanup Method

Sample: Tank U110 Matrix Spike							Total
No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Recovery	
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0		100.0
2) #2-Fluorophenol	107.7	110.9	0.2	50.7	0.0		50.7
3) #Phenol-d5	117.9	121.6	2.2	57.7			57.7
4) Phenol	114.6	104.7	3.7	53.7			53.7
5) bis(2-Chloroethyl)ether	89.7	121.4	60.8	0.7			0.7
6) 2-Chlorophenol	96.6	101.7	0.4	48.4			48.4
7) 1,3-Dichlorobenzene	102.7	101.9	57.7	0.5	1.1		1.6
8) 1,4-Dichlorobenzene	107.0	104.4	57.0	0.5	2.4		2.9
9) Benzyl alcohol	105.6	120.2	64.1	2.4			2.4
10) 1,2-Dichlorobenzene	98.7	100.9	57.6	0.5	16.3		16.8
11) 2-Methylphenol	91.9	97.7	11.7	40.5			40.5
12) bis(2-Chloroisopropyl)ether	116.8	131.8	83.2	0.9			0.9
13) 4-Methylphenol	102.9	108.0	9.3	47.3			47.3
14) N-Nitroso-di-n-propylamine	118.4	138.8	72.0	1.7			1.7
15) Hexachloroethane	100.2	105.2	57.2	0.4			0.4
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0		100.0
17) #Nitrobenzene-d5	103.3	108.9	66.3	0.7			0.7
18) Nitrobenzene	106.9	109.2	68.5	1.1			1.1
19) Isophorone	104.5	108.8	61.5	2.4			2.4
20) 2-Nitrophenol	112.3	108.9	0.2	62.0			62.0
21) 2,4-Dimethylphenol	57.7	56.3	15.8	7.7	0.1		7.8
22) Benzoic acid	47.8	59.9		36.6			36.6
23) bis(2-Chloroethoxy)methane	107.5	110.3	67.8	2.1			2.1
24) 2,4-Dichlorophenol	108.8	103.9	0.4	63.7			63.7
25) 1,2,4-Trichlorobenzene	102.6	99.8	59.9	1.0	0.8		1.8
26) Naphthalene	104.8	99.5	60.9	2.3	42.7		44.9
27) 4-Chloroaniline	65.1	69.2	36.0	0.1			0.1
28) Hexachlorobutadiene	101.7	99.1	61.5	0.8	0.1		0.9
29) 4-Chloro-3-methylphenol	108.0	115.4	8.7	59.5			59.5
30) 2-Methylnaphthalene	103.3	85.4	55.5	2.0	47.6		49.6
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0		100.0
32) Hexachlorocyclopentadiene	112.5	117.3	48.1	0.6			0.6
33) 2,4,6-Trichlorophenol	112.9	103.9		67.9			67.9
34) 2,4,5-Trichlorophenol	102.5	99.1		65.8			65.8
35) 2-Chloronaphthalene	104.0	98.8	60.4	2.3	51.4		53.7
36) #2-Fluorobiphenyl	107.2	102.6	63.5	2.5	52.8		55.3
37) 2-Nitroaniline	113.3	117.7	75.1	2.5			2.5
38) Dimethylphthalate	113.2	109.9	67.7	3.4			3.4
39) Acenaphthylene	86.6	79.6	50.6	1.9	43.2		45.1
40) 2,6-Dinitrotoluene	109.5	110.4	66.4	2.2			2.2
41) 3-Nitroaniline	75.3	72.7	67.9	1.0	0.1		1.1

TABLE 19. (contd)

Sample: Tank U110 Matrix Spike

No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Total Recovery
42) Acenaphthene	110.7	108.9	68.4	2.8	57.9	60.7
43) 2,4-Dinitrophenol	24.5	35.0		62.1		62.1
44) 4-Nitrophenol	108.4	113.3		75.1		75.1
45) Dibenzofuran	109.1	106.0	66.6	2.9	56.9	59.9
46) 2,4-Dinitrotoluene	112.8	108.6	68.4	2.8		2.8
47) Diethylphthalate	114.0	114.2	71.2	3.6	0.2	3.9
48) 4-Chlorophenyl-phenylether	69.2	67.6	40.3	1.9	38.4	40.2
49) Fluorene	110.7	108.7	68.5	2.9	60.4	63.3
50) 4-Nitroaniline	91.3	80.8	56.5	1.2		1.2
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	43.0	51.9		69.3		69.3
53) N-Nitrosodiphenylamine	120.6	113.0	67.9	2.2		2.2
54) #2,4,6-Tribromophenol	110.4	100.3		67.9		67.9
55) 4-Bromophenyl-phenylether	109.4	100.4	66.5	3.0	60.4	63.4
56) Hexachlorobenzene	112.8	99.9	69.6	3.1	0.2	3.3
57) Pentachlorophenol	109.2	98.1		82.9		82.9
58) Phenanthrene	111.1	104.9	71.8	3.3	63.8	67.1
59) Anthracene	99.0	91.2	57.9	2.3	52.4	54.7
60) Di-n-butylphthalate	110.3	101.6	71.5	3.8	0.0	3.8
61) Fluoranthene	107.1	96.8	69.4	3.2	63.3	66.5
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	101.1	108.5	69.4	2.9	67.6	70.5
64) #Terphenyl-d14	93.1	103.3	66.4	2.9	65.4	68.3
65) Butylbenzylphthalate	101.6	113.0	70.1	3.3		3.3
66) 3,3'-Dichlorobenzidine	55.0	51.0	48.3			0.0
67) Benzo(a)anthracene	98.1	107.5	66.7	2.8	67.4	70.2
68) Chrysene	95.7	108.4	70.6	3.1	73.3	76.4
69) Bis(2-Ethylhexyl)phthalate	94.4	101.1	63.3	3.2	0.3	3.5
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	110.2	95.7	63.5	3.4	0.2	3.7
72) Benzo(b)fluoranthene	118.3	111.0	67.9	3.1	61.0	64.2
73) Benzo(k)fluoranthene	127.2	86.0	65.6	3.2	67.3	70.5
74) Benzo(a)pyrene	99.3	89.0	53.5	2.1	52.1	54.2
75) Indeno(1,2,3-cd)pyrene	145.3	114.5	68.7	3.0	63.8	66.9
76) Dibenz(a,h)anthracene	107.5	98.3	62.1	2.6	58.1	60.8
77) Benzo(g,h,i)perylene	120.3	114.3	66.6	2.8	60.9	63.7
NPH (Paraffin Hydrocarbon, C12-C14)	25.3	27.8	12.9	0.3		0.3

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 20. Percent Recovery Results for Single Shell Tank U110 Matrix Spike Duplicate, Initial PNL Cleanup Method

Sample: Tank U110 Matrix Spike Duplicate							Total
No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Recovery	
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0		100.0
2) #2-Fluorophenol	98.8	101.9	0.3	46.9	0.2		47.1
3) #Phenol-d5	118.7	114.9	1.9	56.2			56.2
4) Phenol	117.4	108.5	2.2	57.6			57.6
5) bis(2-Chloroethyl)ether	80.2	109.2	57.0	0.2			0.2
6) 2-Chlorophenol	94.3	97.5	0.6	48.1			48.1
7) 1,3-Dichlorobenzene	92.3	90.3	50.7	0.2	31.9		32.1
8) 1,4-Dichlorobenzene	97.4	94.2	50.3	0.2	33.7		33.9
9) Benzyl alcohol	103.2	114.7	62.9	0.8			0.8
10) 1,2-Dichlorobenzene	90.9	91.6	51.7	0.2	34.3		34.5
11) 2-Methylphenol	80.8	85.1	9.0	39.7			39.7
12) bis(2-Chloroisopropyl)ether	110.8	123.7	80.1	0.2			0.2
13) 4-Methylphenol	86.8	91.6	7.4	44.0			44.0
14) N-Nitroso-dl-n-propylamine	115.2	128.4	73.3	0.3			0.3
15) Hexachloroethane	91.9	94.1	50.8	0.1	13.1		13.2
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0		100.0
17) #Nitrobenzene-d5	97.1	104.2	66.0	0.2	0.0		0.2
18) Nitrobenzene	133.6	145.0	91.6	0.4			0.4
19) Isophorone	68.7	108.2	64.1	0.3			0.3
20) 2-Nitrophenol	108.3	109.9	0.3	65.8			65.8
21) 2,4-Dimethylphenol	8.2	9.1	4.3	3.0			3.0
22) Benzoic acid	37.5	40.5		15.6			15.6
23) bis(2-Chloroethoxy)methane	104.9	107.7	70.7	0.3			0.3
24) 2,4-Dichlorophenol	103.8	103.4	0.6	65.5			65.5
25) 1,2,4-Trichlorobenzene	97.6	97.5	59.7		49.3		49.3
26) Naphthalene	97.4	99.2	60.6	3.2	52.0		55.2
27) 4-Chloroaniline	41.9		1.0	0.1			0.1
28) Hexachlorobutadiene	95.7	95.4	62.0	0.2	18.6		18.8
29) 4-Chloro-3-methylphenol	103.7	115.0	7.6	62.2			62.2
30) 2-Methylnaphthalene	97.3	91.5	61.2	0.3	58.6		58.9
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0		100.0
32) Hexachlorocyclopentadiene	117.7	118.8	55.9	0.1	16.7		16.7
33) 2,4,6-Trichlorophenol	111.8	100.6		66.7			66.7
34) 2,4,5-Trichlorophenol	104.3	94.9		66.4			66.4
35) 2-Chloronaphthalene	101.0	96.4	62.5	0.3	61.1		61.4
36) #2-Fluorobiphenyl	106.7	98.2	62.6	0.3	61.9		62.2
37) 2-Nitroaniline	96.9	98.6	69.4	0.3			0.3
38) Dimethylphthalate	115.7	109.4	72.6	0.5			0.5
39) Acenaphthylene	70.9	65.0	42.5	0.2	41.8		42.0
40) 2,6-Dinitrotoluene	110.2	109.0	71.8	0.2			0.2
41) 3-Nitroaniline	3.6		5.9				0.0

TABLE 20. (contd)

Sample: Tank U110 Matrix Spike Duplicate

No. Compound Name	Step 1	Re-Step 1	P2BN	P2A	P3BN	Total Recovery
42) Acenaphthene	110.2	105.3	69.2	0.3	66.5	66.9
43) 2,4-Dinitrophenol	42.8	45.4		59.2		59.2
44) 4-Nitrophenol	131.4	133.8		91.9		91.9
45) Dibenzofuran	109.4	103.7	70.2	0.4	67.3	67.7
46) 2,4-Dinitrotoluene	113.2	109.4	74.8	0.4		0.4
47) Diethylphthalate	111.9	107.9	74.3	0.6	0.3	0.8
48) 4-Chlorophenyl-phenylether	69.5	66.6	41.5	0.2	45.9	46.1
49) Fluorene	110.6	104.9	71.4	0.4	70.5	70.9
50) 4-Nitroaniline	29.9	30.3	36.1			0.0
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	62.4	65.4		73.6		73.6
53) N-Nitrosodiphenylamine	61.6	56.3	45.9	0.2		0.2
54) #2,4,6-Tribromophenol	117.4	105.1		72.9		72.9
55) 4-Bromophenyl-phenylether	109.3	102.9	70.6	0.5	71.4	71.8
56) Hexachlorobenzene	112.0	99.5	74.8	0.5	42.2	42.7
57) Pentachlorophenol	119.7	111.6		93.4		93.4
58) Phenanthrene	108.2	102.6	75.0	0.5	74.4	74.9
59) Anthracene	51.4	49.6	33.1	0.5	32.9	33.4
60) Di-n-butylphthalate	106.7	103.3	75.3	0.7	0.0	0.7
61) Fluoranthene	104.8	96.3	71.9	0.5	72.7	73.2
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	95.8	101.6	66.0	0.5	72.4	72.9
64) #Terphenyl-d14	90.0	93.8	62.0	0.6	69.4	70.0
65) Butylbenzylphthalate	99.5	105.5	68.4	0.6		0.6
66) 3,3'-Dichlorobenzidine						0.0
67) Benzo(a)anthracene	84.8	93.0	58.1	0.5	63.8	64.3
68) Chrysene	99.2	102.8	68.1	0.6	76.8	77.4
69) Bis(2-Ethylhexyl)phthalate	94.5	97.1	62.0	0.7	0.1	0.9
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	113.8	100.2	71.8	0.6	0.1	0.7
72) Benzo(b)fluoranthene	119.0	106.4	79.0	0.5	71.2	71.7
73) Benzo(k)fluoranthene	113.9	109.0	66.0	0.5	74.8	75.2
74) Benzo(a)pyrene	35.7	34.6	22.6	0.1	22.4	22.5
75) Indeno(1,2,3-cd)pyrene	87.7	116.1	74.2	0.5	71.6	72.1
76) Dibenzo(a,h)anthracene	105.1	102.9	65.9	0.4	65.4	65.8
77) Benzo(g,h,i)perylene	116.6	112.9	73.7	0.5	70.2	70.6
NPH (Paraffin Hydrocarbon, C12-C14)	20.7	23.6	11.0		0.8	0.8

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 21. Results for Single Shell Tank B110 Using Initial PNL Cleanup Method

Sample: Tank B110						
No. Compound Name	Step 1	E2N	E2A	E2B	E3N	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	98.1	0.2	49.7			49.7
3) #Phenol-d5	104.2	1.8	55.1			55.1
4) Phenol						
5) bis(2-Chloroethyl)ether						
6) 2-Chlorophenol						
7) 1,3-Dichlorobenzene						
8) 1,4-Dichlorobenzene						
9) Benzyl alcohol						
10) 1,2-Dichlorobenzene						
11) 2-Methylphenol						
12) bis(2-Chloroisopropyl)ether						
13) 4-Methylphenol						
14) N-Nitroso-di-n-propylamine						
15) Hexachloroethane						
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	92.6	50.1	0.4	0.3		0.8
18) Nitrobenzene						
19) Isophorone						
20) 2-Nitrophenol						
21) 2,4-Dimethylphenol						
22) Benzoic acid						
23) bis(2-Chloroethoxy)methane						
24) 2,4-Dichlorophenol						
25) 1,2,4-Trichlorobenzene						
26) Naphthalene						
27) 4-Chloroaniline						
28) Hexachlorobutadiene						
29) 4-Chloro-3-methylphenol						
30) 2-Methylnaphthalene						
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene						
33) 2,4,6-Trichlorophenol						
34) 2,4,5-Trichlorophenol						
35) 2-Chloronaphthalene						
36) #2-Fluorobiphenyl	95.4	55.6	0.7	1.4	19.8	21.9
37) 2-Nitroaniline						
38) Dimethylphthalate						
39) Acenaphthylene						
40) 2,6-Dinitrotoluene						
41) 3-Nitroaniline						

TABLE 21. (contd)

Sample: Tank B110							Total
No.	Compound Name	Step 1	E2N	E2A	E2B	E3N	Recovery
42)	Acenaphthene						
43)	2,4-Dinitrophenol						
44)	4-Nitrophenol						
45)	Dibenzofuran						
46)	2,4-Dinitrotoluene						
47)	Diethylphthalate						
48)	4-Chlorophenyl-phenylether						
49)	Fluorene						
50)	4-Nitroaniline						
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol						
53)	N-Nitrosodiphenylamine						
54)	#2,4,6-Tribromophenol	110.8		69.8			69.8
55)	4-Bromophenyl-phenylether						
56)	Hexachlorobenzene						
57)	Pentachlorophenol						
58)	Phenanthrene						
59)	Anthracene						
60)	Di-n-butylphthalate						
61)	Fluoranthene						
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene						
64)	#Terphenyl-d14	96.9	66.6	1.7	4.2	36.2	42.1
65)	Butylbenzylphthalate						
66)	3,3'-Dichlorobenzidine						
67)	Benzo(a)anthracene						
68)	Chrysene						
69)	Bis(2-Ethylhexyl)phthalate		2.1				
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate						
72)	Benzo(b)fluoranthene						
73)	Benzo(k)fluoranthene						
74)	Benzo(a)pyrene						
75)	Indeno(1,2,3-cd)pyrene						
76)	Dibenz(a,h)anthracene						
77)	Benzo(g,h,i)perylene						
NPH (Paraffin Hydrocarbon, C12-C14)		170.8	85.9	1.5	4.4		5.9

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 22. Percent Recovery Results for Single Shell Tank B110 Matrix Spike,
EPA 3630 Cleanup Method

Sample: Tank B110 Matrix Spike						Total
No. Compound Name	Step 1	E2N	E2A	E2B	E3N	Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	99.0	3.5	70.4		0.0	70.4
3) #Phenol-d5	109.9	2.1	75.0	0.1		75.1
4) Phenol	105.0	2.3	64.8			64.8
5) bis(2-Chloroethyl)ether	103.6	54.9	0.4	50.5		50.8
6) 2-Chlorophenol	94.8	0.4	68.9			68.9
7) 1,3-Dichlorobenzene	93.8	51.8	0.3	2.5	0.5	3.3
8) 1,4-Dichlorobenzene	98.4	54.1	0.3	2.5	0.9	3.7
9) Benzyl alcohol	97.5	58.1	1.7	5.8		7.2
10) 1,2-Dichlorobenzene	93.0	52.5	0.3	2.7	4.4	7.4
11) 2-Methylphenol	96.8	11.8	54.6	0.7		55.3
12) bis(2-Chloroisopropyl)ether	106.8	68.2		3.3	0.1	3.4
13) 4-Methylphenol	97.2	9.2	60.8	0.5		61.3
14) N-Nitroso-di-n-propylamine	96.8	63.0	0.6	2.9		3.5
15) Hexachloroethane	93.3	55.1	0.2	2.5		2.7
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	99.9	67.1	0.4	3.2	12.2	15.7
18) Nitrobenzene	102.3	62.3	0.4	3.2	13.6	17.2
19) Isophorone	98.7	60.4	0.9	3.5		4.4
20) 2-Nitrophenol	98.1	0.2	74.9			74.9
21) 2,4-Dimethylphenol	91.5	32.2	18.8	2.0	0.3	21.1
22) Benzoic acid	71.9		64.7			64.7
23) bis(2-Chloroethoxy)methane	109.3	66.6	0.7	4.0		4.7
24) 2,4-Dichlorophenol	110.0	0.5	77.3			77.3
25) 1,2,4-Trichlorobenzene	99.5	59.7	0.5	3.6	0.6	4.7
26) Naphthalene	105.5	63.4	1.1	4.0	51.7	56.8
27) 4-Chloroaniline	114.7			51.2		51.2
28) Hexachlorobutadiene	106.2	61.3	0.5	3.8		4.3
29) 4-Chloro-3-methylphenol	107.2	9.6	70.8	0.4		71.2
30) 2-Methylnaphthalene	90.1	62.7	0.7	4.1	54.2	59.0
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	80.5	48.4		1.2		1.2
33) 2,4,6-Trichlorophenol	101.8		83.0			83.0
34) 2,4,5-Trichlorophenol	91.5		75.7			75.7
35) 2-Chloronaphthalene	99.2	62.9	0.8	3.9	48.7	53.4
36) #2-Fluorobiphenyl	103.2	64.9	0.9	4.2	54.7	59.9
37) 2-Nitroaniline	95.5	54.0	1.3	3.2		4.5
38) Dimethylphthalate	102.2	68.3	1.5	4.2		5.7
39) Acenaphthylene	96.2	62.5		3.8	52.3	56.2
40) 2,6-Dinitrotoluene	87.4	67.6	0.9	2.7		3.6
41) 3-Nitroaniline	61.8	20.4		16.3		16.3

TABLE 22. (contd)

Sample: Tank B110 Matrix Spike						
No. Compound Name	Step 1	E2N	E2A	E2B	E3N	Total Recovery
42) Acenaphthene	108.2	70.5	1.2	4.7	58.2	64.0
43) 2,4-Dinitrophenol	53.9		81.9			81.9
44) 4-Nitrophenol	87.2		85.1			85.1
45) Dibenzofuran	101.8	67.2	1.3	4.3	57.9	63.5
46) 2,4-Dinitrotoluene	90.8	68.7	1.4	3.0		4.5
47) Diethylphthalate	106.5	74.9	1.8	4.7	0.2	6.7
48) 4-Chlorophenyl-phenylether	64.7	44.3	0.8	2.9	40.0	43.7
49) Fluorene	104.5	70.1	1.4	4.4	59.9	65.7
50) 4-Nitroaniline	136.0	107.1	1.2	8.5		9.7
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	76.5		86.3			86.3
53) N-Nitrosodiphenylamine	125.9	71.6	1.3	5.2	14.3	20.7
54) #2,4,6-Tribromophenol	95.4		92.2			92.2
55) 4-Bromophenyl-phenylether	108.5	69.1	1.6	4.5	63.0	69.1
56) Hexachlorobenzene	94.0	70.2	1.6	5.0	0.9	7.5
57) Pentachlorophenol	83.7		95.7			95.7
58) Phenanthrene	106.1	74.5	1.8	5.4	66.0	73.2
59) Anthracene	101.7	69.0	1.6	4.9	60.3	66.8
60) Di-n-butylphthalate	105.4	75.5	2.1	5.1	0.0	7.3
61) Fluoranthene	101.0	74.9	1.9	5.4	68.2	75.4
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	102.6	70.8	1.8	5.3	73.0	80.1
64) #Terphenyl-d14	97.1	67.0	2.0	5.1	68.2	75.2
65) Butylbenzylphthalate	105.4	73.6	2.1	5.0		7.1
66) 3,3'-Dichlorobenzidine	69.6	18.2		3.4		3.4
67) Benzo(a)anthracene	105.0	72.6	2.0	5.1	73.0	80.1
68) Chrysene	106.1	72.6	2.2	5.5	76.4	84.1
69) Bis(2-Ethylhexyl)phthalate	110.1	72.7	2.4	5.3	0.2	7.9
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	108.4	73.4	1.7	5.0		6.7
72) Benzo(b)fluoranthene	101.1	75.2	1.9	4.7	61.7	68.3
73) Benzo(k)fluoranthene	107.9	72.8	2.0	5.1	59.6	66.7
74) Benzo(a)pyrene	98.2	69.8	1.8	4.7	59.0	65.4
75) Indeno(1,2,3-cd)pyrene	105.9	76.2	1.8	4.3	64.6	70.7
76) Dibenzo(a,h)anthracene	95.9	68.5	1.6	3.8	59.3	64.7
77) Benzo(g,h,i)perylene	103.7	76.9	1.1	4.2	58.1	63.4
NPH (Paraffin Hydrocarbon, C12-C14)	259.5	141.7	2.5	12.0	0.5	14.9

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 23. Percent Recovery Results for Single Shell Tank B110 Matrix Spike Duplicate, EPA 3630 Cleanup Method

Sample: Tank B110 Matrix Spike Duplicate						
No. Compound Name	Step 1	E2N	E2A	E2B	E3N	Total Recovery
1) *1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0
2) #2-Fluorophenol	102.9	0.2	63.9		0.0	64.0
3) #Phenol-d5	118.7	2.4	69.8			69.8
4) Phenol	110.0	2.6	62.0			62.0
5) bis(2-Chloroethyl)ether	110.8	51.7	1.0	44.6		45.6
6) 2-Chlorophenol	103.3	0.6	66.2			66.2
7) 1,3-Dichlorobenzene	99.7	48.4	0.8	0.3	3.7	4.8
8) 1,4-Dichlorobenzene	102.2	49.3	0.8	0.3	5.0	6.2
9) Benzyl alcohol	103.4	53.5	2.9	1.4		4.2
10) 1,2-Dichlorobenzene	102.6	49.3	0.8	0.3	8.5	9.6
11) 2-Methylphenol	100.2	13.7	50.9			50.9
12) bis(2-Chloroisopropyl)ether	115.5	63.7	1.1	0.5	0.2	1.8
13) 4-Methylphenol	106.3	10.8	56.4			56.4
14) N-Nitroso-di-n-propylamine	100.2	58.9	1.5	0.4		1.9
15) Hexachloroethane	97.5	50.2	0.7	0.3	0.5	1.5
16) *Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0
17) #Nitrobenzene-d5	102.6	62.3	0.9	0.3	12.2	13.4
18) Nitrobenzene	104.6	61.7	1.1	0.3	13.9	15.3
19) Isophorone	103.4	57.3	2.1	0.6		2.7
20) 2-Nitrophenol	101.8	0.4	71.8			71.8
21) 2,4-Dimethylphenol	92.2	34.6	17.8	0.3	0.3	18.4
22) Benzoic acid	74.2		53.5			53.5
23) bis(2-Chloroethoxy)methane	113.4	64.0	1.8	0.7		2.5
24) 2,4-Dichlorophenol	108.9	0.6	72.9			72.9
25) 1,2,4-Trichlorobenzene	106.9	59.0	1.2	0.7	5.1	6.9
26) Naphthalene	110.3	61.8	1.4	0.7	41.2	43.3
27) 4-Chloroaniline	123.1			47.6		47.6
28) Hexachlorobutadiene	106.6	56.8	1.1	0.8	0.7	2.6
29) 4-Chloro-3-methylphenol	111.2	11.4	66.0			66.0
30) 2-Methylnaphthalene	94.6	61.3	1.8	1.0	45.3	48.1
31) *Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0
32) Hexachlorocyclopentadiene	84.7	45.1	0.6	0.2		0.8
33) 2,4,6-Trichlorophenol	106.4		78.1			78.1
34) 2,4,5-Trichlorophenol	94.4		71.6			71.6
35) 2-Chloronaphthalene	103.8	60.6	1.9	1.2	43.2	46.3
36) #2-Fluorobiphenyl	105.4	61.4	2.1	1.3	46.9	50.2
37) 2-Nitroaniline	99.7	57.4	2.8	0.8		3.6
38) Dimethylphthalate	104.8	63.9	3.3	1.6		4.9
39) Acenaphthylene	98.5	59.2	2.0	1.4	44.9	48.3
40) 2,6-Dinitrotoluene	90.5	63.6	2.3	0.8		3.1
41) 3-Nitroaniline	64.7	22.2	0.5	13.4		14.0

TABLE 23. (contd)

Sample: Tank B110 Matrix Spike Duplicate						
No. Compound Name	Step 1	E2N	E2A	E2B	E3N	Total Recovery
42) Acenaphthene	111.6	67.5	2.6	2.0	51.6	56.2
43) 2,4-Dinitrophenol	54.1		86.7			86.7
44) 4-Nitrophenol	89.7		81.9			81.9
45) Dibenzofuran	106.5	63.7	2.7	1.9	50.3	54.8
46) 2,4-Dinitrotoluene	91.0	65.2	2.9	1.1		4.0
47) Diethylphthalate	107.7	68.6	3.7	2.0	0.2	5.9
48) 4-Chlorophenyl-phenylether	68.3	41.5	1.8	1.5	33.4	36.7
49) Fluorene	107.9	65.9	2.9	2.2	51.0	56.2
50) 4-Nitroaniline	140.8	100.7	2.2	3.5		5.7
51) *Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0
52) 4,6-Dinitro-2-methylphenol	76.4		90.1			90.1
53) N-Nitrosodiphenylamine	127.9	69.0	2.7	2.6	18.1	23.4
54) #2,4,6-Tribromophenol	102.2		84.8			84.8
55) 4-Bromophenyl-phenylether	104.8	65.0	3.2	2.5	52.3	58.0
56) Hexachlorobenzene	99.2	65.5	3.3	2.6	2.1	7.9
57) Pentachlorophenol	88.7		85.3			85.3
58) Phenanthrene	109.7	69.0	3.6	3.0	56.1	62.6
59) Anthracene	103.0	65.6	3.4	2.6	52.0	58.0
60) Di-n-butylphthalate	107.8	70.5	4.2	2.0	0.0	6.3
61) Fluoranthene	102.6	68.1	3.8	2.8	57.1	63.6
62) *Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0
63) Pyrene	103.6	72.5	3.8	2.6	58.1	64.5
64) #Terphenyl-d14	97.8	68.0	3.9	2.5	53.9	60.2
65) Butylbenzylphthalate	105.6	73.2	4.6	1.9		6.5
66) 3,3'-Dichlorobenzidine	70.5	22.4		1.4		1.4
67) Benzo(a)anthracene	106.8	74.5	4.0	2.6	58.3	64.9
68) Chrysene	113.3	74.4	4.4	2.7	60.2	67.3
69) Bis(2-Ethylhexyl)phthalate	110.7	72.8	5.1	2.0	0.1	7.2
70) *Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0
71) Di-n-octylphthalate	107.9	65.9	4.7	1.9	0.0	6.5
72) Benzo(b)fluoranthene	103.3	70.8	3.7	2.1	46.8	52.6
73) Benzo(k)fluoranthene	105.4	64.2	4.3	2.4	49.8	56.5
74) Benzo(a)pyrene	101.0	67.8	3.7	2.2	45.2	51.1
75) Indeno(1,2,3-cd)pyrene	106.6	70.7	3.8	2.0	50.7	56.5
76) Dihenz(a,h)anthracene	100.0	64.0	3.4	1.6	46.5	51.5
77) Benzo(g,h,i)perylene	109.1	70.7	3.8	1.9	45.5	51.2
NPH (Paraffin Hydrocarbon, C12-C14)	232.1	115.7	4.4	4.4	0.7	9.4

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 24. Comparison of the Effect of NPH in Blank Spikes on Cleanup Method Recoveries Including Methylene Chloride Elution

Sample: Blank Spike No. Compound Name	PNL with P3BN2 step			EPA with E3N2 step		
	MS+NPH	MS only	NPH Effect	MS+NPH	MS only	NPH Effect
1) *1,4-Dichlorobenzene-d4	100.0	100.0	0.0	100.0	100.0	0.0
2) #2-Fluorophenol	63.1	55.9	7.2	59.7	75.7	-16.0
3) #Phenol-d5	66.1	58.7	7.4	61.9	73.8	-11.9
4) Phenol	59.4	54.4	5.0	56.0	69.1	-13.1
5) bis(2-Chloroethyl)ether	50.4	38.2	12.2	32.6	9.6	23.0
6) 2-Chlorophenol	65.5	57.2	8.2	58.6	70.6	-12.0
7) 1,3-Dichlorobenzene	45.7	34.4	11.3	1.7	4.9	-3.1
8) 1,4-Dichlorobenzene	47.4	38.1	9.3	2.0	4.8	-2.8
9) Benzyl alcohol	3.7	1.7	2.0	5.3	9.6	-4.3
10) 1,2-Dichlorobenzene	47.0	38.0	9.0	3.0	6.1	-3.0
11) 2-Methylphenol	58.0	55.3	2.7	50.3	54.6	-4.3
12) bis(2-Chloroisopropyl)ether	73.5	58.5	15.0	16.2	19.6	-3.4
13) 4-Methylphenol	57.2	48.6	8.5	53.2	58.7	-5.5
14) N-Nitroso-di-n-propylamine	2.3	0.9	1.4	1.6	5.2	-3.6
15) Hexachloroethane	42.9	15.7	27.2	1.2	4.2	-3.1
16) *Naphthalene-d8	100.0	100.0	0.0	100.0	100.0	0.0
17) #Nitrobenzene-d5	72.3	57.8	14.5	58.5	19.5	39.0
18) Nitrobenzene	74.9	60.4	14.5	63.5	22.1	41.4
19) Isophorone	3.5	1.5	2.0	3.9	6.3	-2.4
20) 2-Nitrophenol	78.2	67.9	10.4	64.6	84.1	-19.5
21) 2,4-Dimethylphenol	36.4	38.7	-2.3	30.7	24.7	6.0
22) Benzoic acid	63.6	52.7	10.9	44.1	78.9	-34.8
23) bis(2-Chloroethoxy)methane	79.9	66.5	13.3	3.8	7.1	-3.3
24) 2,4-Dichlorophenol	74.9	70.9	4.1	68.0	82.3	-14.2
25) 1,2,4-Trichlorobenzene	64.9	49.2	15.7	3.6	6.8	-3.2
26) Naphthalene	66.3	55.3	11.0	35.9	43.8	-7.9
27) 4-Chloroaniline	2.5	1.8	0.7	78.9	87.6	-8.6
28) Hexachlorobutadiene	56.2	20.6	35.6	2.9	6.2	-3.3
29) 4-Chloro-3-methylphenol	70.1	61.9	8.2	64.4	68.7	-4.2
30) 2-Methylnaphthalene	70.1	61.5	8.6	41.5	47.7	-6.2
31) *Acenaphthene-d10	100.0	100.0	0.0	100.0	100.0	0.0
32) Hexachlorocyclopentadiene	41.8	14.8	27.1	1.0	1.8	-0.8
33) 2,4,6-Trichlorophenol	83.9	71.7	12.2	67.4	81.3	-13.9
34) 2,4,5-Trichlorophenol	81.6	68.7	12.9	62.4	74.8	-12.4
35) 2-Chloronaphthalene	75.8	63.0	12.7	28.8	37.4	-8.6
36) #2-Fluorobiphenyl	80.0	64.6	15.4	45.4	51.3	-5.9
37) 2-Nitroaniline	85.1	67.5	17.6	4.9	6.6	-1.6
38) Dimethylphthalate	5.4	2.8	2.5	6.9	7.9	-1.1
39) Acenaphthylene	78.1	64.4	13.6	47.5	52.1	-4.6
40) 2,6-Dinitrotoluene	89.3	70.6	18.7	4.6	20.8	-16.3
41) 3-Nitroaniline	8.1	3.8	4.3	72.4	91.9	-19.5

TABLE 24. (contd)

Sample: Blank Spike No. Compound Name	PNL with P3BN2 step			EPA with E3N2 step		
	MS+NPH	MS only	NPH Effect	MS+NPH	MS only	NPH Effect
42) Acenaphthene	86.7	71.6	15.1	53.8	57.4	-3.6
43) 2,4-Dinitrophenol	84.8	78.9	6.0	60.1	96.6	-36.5
44) 4-Nitrophenol	82.0	75.8	6.2	65.1	85.5	-20.5
45) Dibenzofuran	86.1	69.7	16.4	55.5	58.6	-3.1
46) 2,4-Dinitrotoluene	87.1	69.5	17.6	5.1	16.2	-11.1
47) Diethylphthalate	5.9	3.5	2.4	6.9	8.2	-1.2
48) 4-Chlorophenyl-phenylether	0.4	47.0	-46.6	0.3	41.8	-41.5
49) Fluorene	89.9	72.9	17.0	58.9	62.2	-3.3
50) 4-Nitroaniline	37.9	26.1	11.8	5.4	9.9	-4.5
51) *Phenanthrene-d10	100.0	100.0	0.0	100.0	100.0	0.0
52) 4,6-Dinitro-2-methylphenol	88.0	85.1	2.9	70.1	97.3	-27.1
53) N-Nitrosodiphenylamine	79.7	53.5	26.2	109.8	30.2	79.6
54) #2,4,6-Tribromophenol	89.9	85.0	5.0	83.3	92.1	-8.8
55) 4-Bromophenyl-phenylether	89.3	71.6	17.7	63.0	67.5	-4.5
56) Hexachlorobenzene	90.3	39.9	50.4	7.5	8.9	-1.4
57) Pentachlorophenol	96.4	95.2	1.2	73.9	88.5	-14.7
58) Phenanthrene	90.9	75.3	15.6	66.7	70.9	-4.2
59) Anthracene	87.9	70.9	17.0	63.5	64.5	-1.1
60) Di-n-butylphthalate	5.9	3.5	2.3	8.1	8.8	-0.7
61) Fluoranthene	88.0	73.7	14.3	66.9	72.7	-5.8
62) *Chrysene-d12	100.0	100.0	0.0	100.0	100.0	0.0
63) Pyrene	87.1	80.0	7.1	65.0	70.2	-5.2
64) #Terphenyl-d14	83.7	77.1	6.6	60.0	68.8	-8.8
65) Butylbenzylphthalate	5.8	3.3	2.5	6.9	8.9	-2.0
66) 3,3'-Dichlorobenzidine	2.7	0.0	2.7	5.1	9.8	-4.7
67) Benzo(a)anthracene	89.6	81.3	8.2	65.9	70.9	-5.0
68) Chrysene	94.4	86.9	7.5	69.5	76.1	-6.5
69) Bis(2-Ethylhexyl)phthalate	37.1	66.2	-29.1	7.1	10.2	-3.1
70) *Perylene-d12	100.0	100.0	0.0	100.0	100.0	0.0
71) Di-n-octylphthalate	19.2	30.1	-10.9	6.6	9.5	-2.8
72) Benzo(b)fluoranthene	87.7	76.2	11.6	61.3	66.3	-5.0
73) Benzo(k)fluoranthene	92.1	81.2	10.9	63.0	66.7	-3.7
74) Benzo(a)pyrene	86.9	76.1	10.8	60.5	64.0	-3.5
75) Indeno(1,2,3-cd)pyrene	90.4	76.9	13.4	61.2	69.5	-8.3
76) Dibenz(a,h)anthracene	80.4	71.6	8.9	54.6	65.7	-11.0
77) Benzo(g,h,i)perylene	87.3	75.2	12.2	56.5	58.9	-2.5
NPH (Paraffin Hydrocarbon, C12-C14)	27.4	0.1	27.3	5.7		-2.5

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 25. Percent Recovery Results for Blank Spikes, Following Soil Extraction Procedure Prior to Final PNL Cleanup Method

No.	Compound Name	Fraction	Blank	Ext 1	Ext 2	Ext 3	Ext 4	Ext 5	Ext 6	Ext 7	Ext 8	Ext 9
1)	*1,4-Dichlorobenzene-d4		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol		100.2	111.1	106.7	106.0	113.6	97.5	104.7	110.4	102.9	104.5
3)	#Phenol-d5		101.1	103.9	100.9	98.6	107.4	92.4	96.7	102.5	96.9	96.6
4)	Phenol			93.3	90.7	88.2	99.6	86.0	90.8	95.7	90.6	92.4
5)	bis(2-Chloroethyl)ether			115.3	116.3	117.1	122.4	107.4	115.3	117.6	110.0	119.7
6)	2-Chlorophenol			104.1	97.6	98.2	106.5	88.9	97.5	101.9	96.6	98.4
7)	1,3-Dichlorobenzene			100.9	97.4	98.1	102.7	88.9	98.5	101.3	93.5	100.0
8)	1,4-Dichlorobenzene			94.3	94.3	96.3	105.5	88.5	91.6	96.0	92.6	94.1
9)	Benzyl alcohol			101.4	99.6	99.4	106.9	90.0	98.5	103.8	97.0	99.1
10)	1,2-Dichlorobenzene			100.7	97.7	97.5	104.0	89.8	96.7	100.2	95.9	97.3
11)	2-Methylphenol			96.2	93.0	91.3	99.6	85.3	99.1	95.5	90.5	90.5
12)	bis(2-Chloroisopropyl)ether			105.6	103.0	103.6	114.0	96.1	104.1	109.0	101.8	104.4
13)	4-Methylphenol			97.9	94.0	93.8	102.6	85.7	93.3	98.5	91.1	93.7
14)	N-Nitroso-di-n-propylamine			108.8	106.6	107.7	111.3	97.7	107.5	103.2	101.4	107.1
15)	Hexachloroethane			97.1	98.0	96.6	101.7	88.9	95.1	99.2	95.4	96.9
16)	*Naphthalene-d8		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5		94.8	98.2	92.8	94.6	99.5	86.9	93.4	96.2	92.1	95.7
18)	Nitrobenzene			97.3	93.4	95.3	99.0	87.6	93.6	97.6	92.7	95.5
19)	Isophorone			100.7	96.7	97.6	102.1	89.2	97.1	101.7	95.9	99.3
20)	2-Nitrophenol			98.1	92.0	92.1	98.6	85.5	91.7	96.0	89.9	92.8
21)	2,4-Dimethylphenol			93.7	89.3	90.7	95.8	82.8	87.0	93.7	89.1	91.9
22)	Benzoic acid			91.2	88.8	89.7	93.1	81.3	87.6	92.0	88.7	91.0
23)	bis(2-Chloroethoxy)methane			96.4	92.5	91.1	98.6	84.9	92.4	96.5	91.1	94.8
24)	2,4-Dichlorophenol			100.2	94.5	94.4	100.8	87.4	93.1	98.9	93.6	96.9
25)	1,2,4-Trichlorobenzene			95.3	90.3	90.0	95.4	84.0	89.0	92.3	89.3	90.9
26)	Naphthalene			90.0	87.0	84.1	90.3	79.2	83.9	87.9	83.4	87.1
27)	4-Chloroaniline			100.0	79.7	65.1	90.7	57.2	53.7	88.0	76.1	56.7
28)	Hexachlorocyclopentadiene			94.9	90.1	89.7	95.5	83.1	89.8	89.4	86.4	91.0
29)	4-Chloro-3-methylphenol			98.0	94.6	96.1	99.3	87.4	95.6	99.3	94.2	97.9
30)	2-Methylnaphthalene			77.4	75.0	75.7	78.8	69.7	74.3	77.4	75.2	76.4
31)	*Acenaphthene-d10		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene			98.5	88.8	85.0	92.8	77.4	82.0	81.2	80.0	79.4
33)	2,4,6-Trichlorophenol			102.3	96.0	98.1	101.4	88.7	97.2	98.2	95.0	97.6
34)	2,4,5-Trichlorophenol			100.5	93.0	94.8	99.1	84.8	92.0	94.2	92.9	95.1
35)	2-Chloronaphthalene			98.4	92.9	94.2	97.9	84.4	92.3	93.3	90.4	94.1
36)	#2-Fluorobiphenyl		94.3	99.7	92.7	95.3	99.4	87.2	93.3	93.8	91.9	95.4
37)	2-Nitroaniline			69.9	65.4	67.8	76.3	63.5	64.0	63.5	63.7	64.1
38)	Dimethylphthalate			105.5	96.9	98.4	105.1	89.2	98.0	99.4	94.8	99.4
39)	Acenaphthylene			97.7	91.9	93.2	95.7	83.4	89.7	90.9	89.0	92.4
40)	2,6-Dinitrotoluene			107.0	101.7	103.0	109.3	91.6	100.4	102.8	96.5	102.8
41)	3-Nitroaniline			99.5	92.2	88.9	100.0	80.4	84.3	93.3	89.2	85.1

TABLE 25. (contd)

No.	Compound Name	Fraction	Blank	Ext 1	Ext 2	Ext 3	Ext 4	Ext 5	Ext 6	Ext 7	Ext 8	Ext 9
42)	Acenaphthene			101.7	97.2	97.5	102.4	89.0	96.5	95.9	93.4	96.5
43)	2,4-Dinitrophenol			119.2	108.2	110.2	115.2	97.7	108.6	110.0	103.4	108.5
44)	4-Nitrophenol			118.6	109.7	113.2	118.6	94.4	108.7	112.6	105.1	108.9
45)	Dibenzofuran			98.4	93.5	92.5	98.7	84.8	92.1	93.6	87.7	93.9
46)	2,4-Dinitrotoluene			106.9	99.1	103.3	106.0	90.4	99.8	101.2	96.6	101.4
47)	Diethylphthalate			108.8	101.4	102.1	108.7	93.1	102.7	102.9	98.8	104.5
48)	4-Chlorophenyl-phenylether			99.8	95.5	96.3	99.6	86.6	93.6	97.1	90.8	95.1
49)	Fluorene			103.4	98.0	98.5	104.3	89.8	98.8	98.5	95.4	98.5
50)	4-Nitroaniline			97.6	92.5	95.2	100.0	84.1	95.4	98.8	87.9	95.4
51)	*Phenanthrene-d10		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol			105.2	100.9	101.2	106.5	89.4	101.9	103.7	95.1	100.5
53)	N-Nitrosodiphenylamine			126.1	121.6	122.5	125.5	112.4	121.0	124.1	118.4	118.4
54)	#2,4,6-Tribromophenol		105.2	103.5	95.4	97.2	100.3	84.8	96.8	99.4	92.1	96.2
55)	4-Bromophenyl-phenylether			95.0	91.4	90.7	96.8	85.2	93.6	93.8	87.5	89.1
56)	Hexachlorobenzene			98.9	92.9	96.2	97.2	87.9	94.3	97.3	88.0	94.2
57)	Pentachlorophenol			110.8	99.7	102.0	110.4	93.9	102.2	105.2	100.1	103.4
58)	Phenanthrene			95.9	92.1	92.0	95.8	83.5	91.1	92.5	86.8	89.1
59)	Anthracene			95.9	93.6	92.5	97.1	85.0	93.0	94.7	89.9	91.7
60)	Di-n-butylphthalate			95.7	89.6	90.5	90.3	83.9	90.5	92.0	86.9	89.6
61)	Fluoranthene			90.4	85.3	86.7	89.2	78.5	87.1	88.1	83.5	85.4
62)	*Chrysene-d12		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene			94.2	90.7	91.6	93.1	81.2	93.9	90.4	89.6	90.6
64)	#Terphenyl-d14		77.7	87.3	86.0	86.7	88.1	74.6	87.1	84.9	85.2	85.7
65)	Butylbenzylphthalate			97.2	92.6	92.2	95.1	81.0	95.7	92.9	91.0	92.0
66)	3,3'-Dichlorobenzidine			138.3	125.1	115.2	126.8	98.1	108.5	127.1	120.5	108.0
67)	Benzo(a)anthracene			99.6	94.4	96.8	97.5	83.7	98.5	95.0	91.5	94.6
68)	Chrysene			88.1	87.9	87.3	87.5	76.6	90.1	84.6	86.6	87.2
69)	Bis(2-Ethylhexyl)phthalate			93.0	90.3	90.3	91.1	79.9	90.3	90.6	87.3	90.0
70)	*Perylene-d12		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate			92.9	86.5	88.5	90.2	81.6	87.1	90.9	86.0	85.5
72)	Benzo(b)fluoranthene			100.2	102.5	103.6	101.9	89.6	99.4	103.2	90.3	95.2
73)	Benzo(k)fluoranthene			92.9	89.0	85.4	93.0	84.3	87.5	93.3	86.6	90.1
74)	Benzo(a)pyrene			98.6	94.6	94.5	99.2	86.8	94.9	95.5	92.3	95.2
75)	Indeno(1,2,3-cd)pyrene			98.6	93.5	93.0	95.8	84.3	93.5	94.7	89.5	93.3
76)	Dibenz(a,h)anthracene			93.8	89.3	89.5	88.6	81.5	89.4	91.2	84.6	87.9
77)	Benzo(g,h,i)perylene			97.5	92.8	93.9	94.9	84.6	93.6	97.2	90.0	94.5
NPH (Paraffin Hydrocarbon, C12-C14)												
				64.0	64.7	57.8	63.3	59.1	63.5	65.6	62.3	63.5

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 26. Percent, Accuracy, and Control Limit Data from Blank Spikes Following Soil Extraction Procedure and Prior to Final PNL Cleanup Method

No.	Compound Name	Mean		-3XSTD	+3XSTD
		Recovery	STD (N-1)		
1)	*1,4-Dichlorobenzene-d4	100.0	0.0	100.0	100.0
2)	#2-Fluorophenol	105.8	5.0	90.8	120.7
3)	#Phenol-d5	99.7	4.3	86.7	112.7
4)	Phenol	91.9	4.0	79.9	103.9
5)	bis(2-Chloroethyl)ether	115.7	4.6		
6)	2-Chlorophenol	98.9	5.0	83.8	114.0
7)	1,3-Dichlorobenzene	97.9	4.3		
8)	1,4-Dichlorobenzene	94.8	4.7	80.8	108.8
9)	Benzyl alcohol	99.6	4.7		
10)	1,2-Dichlorobenzene	97.7	3.9		
11)	2-Methylphenol	92.3	4.3		
12)	bis(2-Chloroisopropyl)ether	104.6	4.9		
13)	4-Methylphenol	94.5	4.8		
14)	N-Nitroso-di-n-propylamine	105.7	4.2	93.1	118.3
15)	Hexachloroethane	96.5	3.5		
16)	*Naphthalene-d8	100.0	0.0	100.0	100.0
17)	#Nitrobenzene-d5	94.4	3.5	83.9	104.9
18)	Nitrobenzene	94.7	3.4		
19)	Isophorone	97.8	3.9		
20)	2-Nitrophenol	93.0	4.1		
21)	2,4-Dimethylphenol	90.4	3.9		
22)	Benzoic acid	89.3	3.5		
23)	bis(2-Chloroethoxy)methane	93.1	4.1		
24)	2,4-Dichlorophenol	95.5	4.2		
25)	1,2,4-Trichlorobenzene	90.7	3.5	80.4	101.1
26)	Naphthalene	85.9	3.6		
27)	4-Chloroaniline	74.1	16.8		
28)	Hexachlorobutadiene	90.0	3.8		
29)	4-Chloro-3-methylphenol	95.8	3.7	84.8	106.9
30)	2-Methylnaphthalene	75.5	2.6		
31)	*Acenaphthene-d10	100.0	0.0	100.0	100.0
32)	Hexachlorocyclopentadiene	85.0	7.0		
33)	2,4,6-Trichlorophenol	97.2	3.9		
34)	2,4,5-Trichlorophenol	94.0	4.5		
35)	2-Chloronaphthalene	93.1	4.1		
36)	#2-Fluorobiphenyl	94.3	3.6	83.4	105.2
37)	2-Nitroaniline	66.5	4.3		
38)	Dimethylphthalate	98.5	5.0		
39)	Acenaphthylene	91.5	4.1		
40)	2,6-Dinitrotoluene	101.7	5.2		
41)	3-Nitroaniline	90.3	6.7		

TABLE 26. (contd)

No.	Compound Name	Mean		
		Recovery	STD (N-1)	
			-3XSTD	+3XSTD
42)	Acenaphthene	96.7	4.0	84.6
43)	2,4-Dinitrophenol	109.0	6.2	
44)	4-Nitrophenol	110.0	7.4	87.9
45)	Dibenzofuran	92.8	4.5	
46)	2,4-Dinitrotoluene	100.5	5.0	85.5
47)	Diethylphthalate	102.5	4.8	
48)	4-Chlorophenyl-phenylether	94.9	4.2	
49)	Fluorene	98.3	4.2	
50)	4-Nitroaniline	94.1	5.2	
51)	*Phenanthrene-d10	100.0	0.0	100.0
52)	4,6-Dinitro-2-methylphenol	100.5	5.3	
53)	N-Nitrosodiphenylamine	121.1	4.3	
54)	#2,4,6-Tribromophenol	97.1	5.8	79.7
55)	4-Bromophenyl-phenylether	91.4	3.7	
56)	Hexachlorobenzene	94.1	3.9	
57)	Pentachlorophenol	103.1	5.3	87.2
58)	Phenanthrene	91.0	4.0	
59)	Anthracene	92.6	3.6	
60)	Di-n-butylphthalate	89.9	3.2	
61)	Fluoranthene	86.0	3.5	
62)	*Chrysene-d12	100.0	0.0	100.0
63)	Pyrene	90.6	3.9	78.9
64)	#Terphenyl-d14	84.3	4.5	70.9
65)	Butylbenzylphthalate	92.2	4.6	
66)	3,3'-Dichlorobenzidine	118.6	12.4	
67)	Benzo(a)anthracene	94.6	4.8	
68)	Chrysene	86.2	3.9	
69)	Bis(2-Ethylhexyl)phthalate	89.2	3.8	
70)	*Perylene-d12	100.0	0.0	100.0
71)	Di-n-octylphthalate	87.7	3.4	
72)	Benzo(b)fluoranthene	98.4	5.4	
73)	Benzo(k)fluoranthene	89.1	3.4	
74)	Benzo(a)pyrene	94.6	3.6	
75)	Indeno(1,2,3-cd)pyrene	92.9	4.0	
76)	Dibenz(a,h)anthracene	88.4	3.6	
77)	Benzo(g,h,i)perylene	93.2	4.0	
	NPH (Paraffin Hydrocarbon, C12-C14)	62.6	2.6	

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 27. Percent Recovery Results for Blank Spikes Numbers 1 and 2 Using the Final PNL Cleanup Method

No.	Compound Name	Fraction	Total Rec			Total Rec			Total Rec		
			Blank BN	Blank A	Blank	Ext 1 BN	Ext 1 A	Ext 1	Ext 2 BN	Ext 2 A	Ext 2
1)	*1,4-Dichlorobenzene-d4		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol			88.1	88.1		77.1	77.1		79.5	79.5
3)	#Phenol-d5			74.1	74.1	2.7	79.2	81.9	2.7	78.3	81.1
4)	Phenol						66.5	66.5		65.4	65.4
5)	bis(2-Chloroethyl)ether					68.1			63.1		63.1
6)	2-Chlorophenol						72.4	72.4	0.8	73.6	74.4
7)	1,3-Dichlorobenzene					55.5	0.0	55.5	44.7	0.0	44.8
8)	1,4-Dichlorobenzene					56.0		56.0	45.8		45.8
9)	Benzyl alcohol					76.8		76.8	72.3		72.3
10)	1,2-Dichlorobenzene					60.8	0.0	60.8	55.9	0.0	56.0
11)	2-Methylphenol					16.1	60.3	76.5	16.4	59.0	75.4
12)	bis(2-Chloroisopropyl)ether					91.2		91.2	89.8		89.8
13)	4-Methylphenol					10.8	64.0	74.8	11.5	63.2	74.7
14)	N-Nitroso-di-n-propylamine					83.1		83.1	78.4		78.4
15)	Hexachloroethane					33.3		33.3	25.8		25.8
16)	*Naphthalene-d8		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5		59.4		59.4	75.5	0.0	75.5	70.4		70.4
18)	Nitrobenzene					68.6		68.6	64.8		64.8
19)	Isophorone		0.8		0.8	71.8	0.0	71.8	66.5	0.1	66.6
20)	2-Nitrophenol					0.6	88.2	88.8	0.5	85.2	85.6
21)	2,4-Dimethylphenol					48.2	32.0	80.3	44.0	27.9	71.8
22)	Benzoic acid						80.1	80.1		78.7	78.7
23)	bis(2-Chloroethoxy)methane					85.2		85.2	77.2		77.2
24)	2,4-Dichlorophenol					0.9	85.2	86.1	0.8	82.0	82.7
25)	1,2,4-Trichlorobenzene					61.9		61.9	46.6	0.0	46.7
26)	Naphthalene					78.7	0.0	78.8	72.6	0.1	72.7
27)	4-Chloroaniline					67.9		67.9	47.4		47.4
28)	Hexachlorobutadiene					33.6		33.6	27.0		27.0
29)	4-Chloro-3-methylphenol					10.5	81.8	92.3	11.4	77.9	89.4
30)	2-Methylnaphthalene					69.5		69.5	63.6		63.6
31)	*Acenaphthene-d10		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene					24.0		24.0	16.5		16.5
33)	2,4,6-Trichlorophenol						88.9	88.9		87.7	87.7
34)	2,4,5-Trichlorophenol						84.1	84.1		81.6	81.6
35)	2-Chloronaphthalene					70.2		70.2	62.4	0.1	62.5
36)	#2-Fluorobiphenyl		58.7		58.7	77.1		77.1	71.4	0.1	71.4
37)	2-Nitroaniline					90.6		90.6	86.6		86.6
38)	Dimethylphthalate					88.5		88.5	81.4	0.1	81.5
39)	Acenaphthylene					83.5		83.5	77.8		77.8
40)	2,6-Dinitrotoluene					76.1		76.1	70.5		70.5
41)	3-Nitroaniline					88.9		88.9	79.7		79.7

TABLE 27. (contd)

No.	Compound Name	Fraction	Total Rec			Total Rec						Total Rec	
			Blank A	Blank B	Blank	Ext 1 BN	Ext 1 A	Ext 1	Ext 2 BN	Ext 2 A	Ext 2		
42)	Acenaphthene					88.2	0.1	88.3	82.2	0.1	82.3	88.6	88.6
43)	2,4-Dinitrophenol						91.2	91.2				81.9	81.9
44)	4-Nitrophenol						84.3	84.3				75.7	75.7
45)	Dibenzofuran					76.1		76.1	75.7	0.1			
46)	2,4-Dinitrotoluene					77.4		77.4	73.4				
47)	Diethylphthalate				1.4	85.1	0.2	85.3	80.6	0.2			
48)	4-Chlorophenyl-phenylether	1.4				82.0		82.0	76.8	0.1			
49)	Fluorene					89.8		89.8	84.9	0.1			
50)	4-Nitroaniline					56.4		56.4	74.3				
51)	*Phenanthrene-d10	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol						97.3	97.3				95.4	95.4
53)	N-Nitrosodiphenylamine					105.6		105.6	94.7				
54)	#2,4,6-Tribromophenol		101.1	101.1	101.1		100.0	100.0				93.4	93.4
55)	4-Bromophenyl-phenylether					89.0		89.0	80.9				
56)	Hexachlorobenzene					52.3		52.3	37.0				
57)	Pentachlorophenol						97.7	97.7				100.2	100.2
58)	Phenanthrene					92.0		92.0	83.2				
59)	Anthracene					89.4		89.4	79.8	0.1			
60)	Di-n-butylphthalate	4.1		4.1	4.1	88.2	0.5	88.6	80.5	0.3			
61)	Fluoranthene					86.2	0.1	86.3	77.7	0.1			
62)	*Chrysene-d12	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene					88.1		88.1	91.7	0.1			
64)	#Terphenyl-d14	66.4		66.4	66.4	74.9	0.2	75.1	78.8	0.2			
65)	Butylbenzylphthalate					90.2	0.3	90.5	92.4	0.3			
66)	3,3'-Dichlorobenzidine					76.5		76.5	64.9				
67)	Benzo(e)anthracene					91.8		91.8	94.8				
68)	Chrysene					89.2		89.2	93.5				
69)	Bis(2-Ethylhexyl)phthalate	1.2		1.2	1.2	86.1	0.3	86.3	88.5	0.2			
70)	*Perylene-d12	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate	1.0		1.0	1.0	86.3	0.1	86.4	61.9	0.1			
72)	Benzo(b)fluoranthene					105.6	0.1	105.7	99.9	0.1			
73)	Benzo(k)fluoranthene					87.3	0.1	87.4	89.1	0.1			
74)	Benzo(a)pyrene					90.3	0.0	90.3	85.0	0.1			
75)	Indeno(1,2,3-cd)pyrene					88.3		88.3	84.3				
76)	Dibenz(a,h)anthracene					84.1	0.0	84.1	80.5	0.0			
77)	Benzo(g,h,i)perylene					89.8	0.1	89.9	85.7	0.1			
NPH (Paraffin Hydrocarbon, C12-C14)						11.5	0.0	11.5	13.8	0.1			13.9

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 28. Percent Recovery Results for Blank Spikes Numbers 3, 4, and 5 Using the Final PNL Cleanup Method

No.	Compound Name	Total Rec			Total Rec			Total Rec		
		Ext 3 BN	Ext 3 A	Ext 3	Ext 4 BN	Ext 4 A	Ext 4	Ext 5 BN	Ext 5 A	Ext 5
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol		80.8	80.8		87.2	87.2		85.7	85.7
3)	#Phenol-d5	4.0	80.6	84.6	4.3	87.5	91.8		84.9	84.9
4)	Phenol		68.6	68.6		72.9	72.9		72.4	72.4
5)	bis(2-Chloroethyl)ether	75.3		75.3	81.9		81.9	71.1		71.1
6)	2-Chlorophenol	1.3	75.6	76.9	2.0	81.1	83.0		80.6	80.6
7)	1,3-Dichlorobenzene	66.2		66.2	77.6		77.6	60.9		60.9
8)	1,4-Dichlorobenzene	68.5		68.5	73.0		73.0	57.4		57.4
9)	Benzyl alcohol	83.9		83.9	93.8		93.8	81.0		81.0
10)	1,2-Dichlorobenzene	70.2		70.2	76.4		76.4	65.6		65.6
11)	2-Methylphenol	19.9	61.7	81.7	18.9	67.8	86.6	17.9	64.7	82.6
12)	bis(2-Chloroisopropyl)ether	104.8		104.8	117.4		117.4	102.6		102.6
13)	4-Methylphenol	14.4	65.1	79.4	13.8	71.1	84.9	13.2	68.8	82.0
14)	N-Nitroso-di-n-propylamine	91.9		91.9	106.3		106.3	93.3		93.3
15)	Hexachloroethane	33.7		33.7	53.4		53.4	34.8		34.8
16)	*Naphthalene-d8	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5	79.6		79.6	83.6		83.6	76.1		76.1
18)	Nitrobenzene	71.9		71.9	79.8		79.8	69.4		69.4
19)	Isophorone	73.5	0.0	73.5	81.4	0.0	81.4	73.2		73.2
20)	2-Nitrophenol	0.8	84.8	85.6	1.3	93.8	95.0	0.7	91.3	92.0
21)	2,4-Dimethylphenol	50.3	28.1	78.4	55.3	36.5	91.7	50.1	31.1	81.2
22)	Benzoic acid		76.5	76.5		90.3	90.3		85.6	85.6
23)	bis(2-Chloroethoxy)methane	87.3		87.3	96.0		96.0	85.4		85.4
24)	2,4-Dichlorophenol	1.2	81.6	82.8	1.7	91.5	93.2	1.1	90.2	91.3
25)	1,2,4-Trichlorobenzene	67.5		67.5	75.1		75.1	60.2		60.2
26)	Naphthalene	80.4	0.0	80.5	86.7	0.0	86.7	79.9	0.0	79.9
27)	4-Chloroaniline	76.4		76.4	76.9		76.9	66.8		66.8
28)	Hexachlorobutadiene	30.5		30.5	47.9		47.9	33.5		33.5
29)	4-Chloro-3-methylphenol	14.1	78.6	92.6	13.5	89.8	103.3	13.2	86.8	100.0
30)	2-Methylnaphthalene	69.8		69.8	74.2		74.2	69.8		69.8
31)	*Acenaphthene-d10	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene	24.1		24.1	36.5		36.5	24.1		24.1
33)	2,4,6-Trichlorophenol		88.5	88.5		99.5	99.5		96.8	96.8
34)	2,4,5-Trichlorophenol		85.5	85.5		92.0	92.0		90.9	90.9
35)	2-Chloronaphthalene	71.1		71.1	77.0		77.0	71.3		71.3
36)	#2-Fluorobiphenyl	79.1		79.1	86.0		86.0	77.6		77.6
37)	2-Nitroaniline	95.6		95.6	110.4		110.4	97.4		97.4
38)	Dimethylphthalate	90.1		90.1	102.8		102.8	90.5		90.5
39)	Acenaphthylene	84.7		84.7	94.0		94.0	85.2		85.2
40)	2,6-Dinitrotoluene	78.5		78.5	87.7		87.7	77.6		77.6
41)	3-Nitroaniline	97.0		97.0	100.0		100.0	93.6		93.6

TABLE 28. (contd)

No.	Compound Name	Fraction	Total Rec			Total Rec			Total Rec		
			Ext 3	Ext 3 A	Ext 4 BN	Ext 4 A	Ext 4	Ext 5 BN	Ext 5 A	Ext 5	Total Rec
42)	Acenaphthene		90.4		97.4		97.4	90.0		90.0	90.0
43)	2,4-Dinitrophenol			89.1		108.3	108.3		105.6	105.6	105.6
44)	4-Nitrophenol			89.4	0.5	100.1	100.7	0.4	94.3	94.6	94.6
45)	Dibenzofuran		82.3		79.6		79.6	83.1		83.1	83.1
46)	2,4-Dinitrotoluene		80.8		91.5		91.5	82.8		82.8	82.8
47)	Diethylphthalate		88.4	0.1	96.9	0.2	97.1	86.8		86.8	86.8
48)	4-Chlorophenyl-phenylether		85.5		93.9		93.9	85.6		85.6	85.6
49)	Fluorene		93.5		100.7		100.7	94.0		94.0	94.0
50)	4-Nitroaniline		99.1		64.7		64.7	93.9		93.9	93.9
51)	*Phenanthrene-d10		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol			96.7		110.5	110.5		109.4	109.4	109.4
53)	N-Nitrosodiphenylamine		104.5		115.3		115.3	105.4		105.4	105.4
54)	#2,4,6-Tribromophenol		0.3	92.9	0.7	100.1	100.9	0.4	102.3	102.7	102.7
55)	4-Bromophenyl-phenylether		91.2		100.2		100.2	93.6		93.6	93.6
56)	Hexachlorobenzene		54.7		73.4		73.4	53.5		53.5	53.5
57)	Pentachlorophenol			91.7		105.1	105.1		106.3	106.3	106.3
58)	Phenanthrene		91.2		99.3		99.3	94.7		94.7	94.7
59)	Anthracene		89.2		97.3		97.3	92.4		92.4	92.4
60)	Di-n-butylphthalate		88.1	0.2	99.8	0.4	100.3	90.5	0.4	90.9	90.9
61)	Fluoranthene		86.0	0.0	96.3	0.0	96.3	88.7		88.7	88.7
62)	*Chrysene-d12		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene		91.9	0.0	100.5	0.0	100.5	98.0		98.0	98.0
64)	#Terphenyl-d14		79.1	0.1	79.2	0.1	84.8	85.6	0.1	85.7	85.7
65)	Butylbenzylphthalate		94.2	0.1	94.3	0.4	103.4	97.8	0.2	98.1	98.1
66)	3,3'-Dichlorobenzidine		77.3		85.8		85.8	86.3		86.3	86.3
67)	Benzo(a)anthracene		95.0	0.1	104.9		104.9	101.8		101.8	101.8
68)	Chrysene		90.0		100.5		100.5	100.3		100.3	100.3
69)	Bis(2-Ethylhexyl)phthalate		89.6	0.1	97.8	0.1	97.9	92.0	0.1	92.1	92.1
70)	*Perylene-d12		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate		86.9	0.0	95.7	0.0	95.7	89.7	0.0	89.7	89.7
72)	Benzo(b)fluoranthene		107.6		115.6		115.6	106.1		106.1	106.1
73)	Benzo(k)fluoranthene		88.4		95.9		95.9	96.1		96.1	96.1
74)	Benzo(a)pyrene		92.0	0.0	101.3		101.3	95.9		95.9	95.9
75)	Indeno(1,2,3-cd)pyrene		91.5		102.8		102.8	96.9		96.9	96.9
76)	Dibenz(a,h)anthracene		86.3		93.6		93.6	90.5		90.5	90.5
77)	Benzo(g,h,i)perylene		93.4		104.5		104.5	97.5		97.5	97.5
NPH (Paraffin Hydrocarbon, C12-C14)			11.1	0.0	17.1	0.0	17.1	14.1	0.0	14.2	14.2

* Compound is an Internal Standard
Compound is a Surrogate Spike

TABLE 29. Percent Recovery Results for Blank Spikes Numbers 6, 7, 8 and 9 Using the Final PNL Cleanup Method

No.	Compound Name	Fraction	Total Rec			Total Rec			Total Rec			Total Rec		
			Ext 6 BN	Ext 6 A	Ext 6	Ext 7 BN	Ext 7 A	Ext 7	Ext 8 BN	Ext 8 A	Ext 8	Ext 9 BN	Ext 9 A	Ext 9
1)	*1,4-Dichlorobenzene-d4		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
2)	#2-Fluorophenol			70.3	70.3		84.8	84.8		79.6	79.6		81.6	81.6
3)	#Phenol-d5	3.0		71.8	74.8		82.7	82.7		80.8	80.8	2.5	81.9	84.4
4)	Phenol			60.6	60.6		69.3	69.3		68.7	68.7		68.6	68.6
5)	bis(2-Chloroethyl)ether	62.5		62.5	72.0			72.0	80.1		80.1	74.6		74.6
6)	2-Chlorophenol			66.6	66.6		77.4	77.4		75.1	75.1		76.3	76.3
7)	1,3-Dichlorobenzene	54.0		54.0	72.6		0.1	72.7	73.7		73.7	70.4	0.1	70.4
8)	1,4-Dichlorobenzene	52.9		52.9	71.1			71.1	74.0		74.0	69.2		69.2
9)	Benzyl alcohol	73.5		73.5	84.2			84.2	90.9		90.9	85.5		85.5
10)	1,2-Dichlorobenzene	55.8		55.8	73.0		0.1	73.0	78.0		78.0	71.3	0.1	71.3
11)	2-Methylphenol	16.9		55.4	72.2	18.5	62.1	80.6	18.6	62.4	91.1	14.1	67.3	81.5
12)	bis(2-Chloroisopropyl)ether	89.6		89.6	112.1			112.1	114.8		114.8	106.3		106.3
13)	4-Methylphenol	12.2		57.5	69.7	12.9	67.7	80.5	13.5	56.0	79.5	9.7	68.6	78.3
14)	N-Nitroso-di-n-propylamine	83.1			83.1	104.1		104.1	107.1		107.1	98.4		98.4
15)	Hexachloroethane	37.6			37.6	61.7		61.7	47.4		47.4	49.5		49.5
16)	*Naphthalene-d8	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
17)	#Nitrobenzene-d5	69.7		69.7	83.3			83.3	85.6		85.6	79.6		79.6
18)	Nitrobenzene	63.3		63.3	74.3			74.3	77.0		77.0	71.7		71.7
19)	Isophorone	68.9		68.9	79.2		0.1	79.3	78.9		78.9	75.2	0.1	75.3
20)	2-Nitrophenol	0.7		77.9	78.6	0.9	94.3	95.2	1.0	87.6	88.6	0.6	88.6	92.2
21)	2,4-Dimethylphenol	49.0		27.4	76.4	51.9	32.9	84.8	54.9	31.3	86.1	44.0	42.3	86.3
22)	Benzoic acid			70.6	70.6		93.0	93.0		87.8	87.8		80.2	80.2
23)	bis(2-Chloroethoxy)methane	81.9		81.9	92.9			92.9	92.2		92.2	89.5		89.5
24)	2,4-Dichlorophenol	1.0		78.1	79.2	1.2	90.6	91.8	1.1	87.7	88.9	0.8	86.1	86.9
25)	1,2,4-Trichlorobenzene	60.9		60.9	72.1			72.1	74.6		74.6	71.1	0.1	71.2
26)	Naphthalene	71.3		71.3	84.7		0.1	84.8	87.4	0.0	87.4	83.4	0.1	83.5
27)	4-Chloroaniline	80.3		80.3	85.7			85.7	71.6		71.6	100.0		100.0
28)	Hexachlorobutadiene	40.4		40.4	59.3		0.0	59.3	43.3		43.3	45.1		45.1
29)	4-Chloro-3-methylphenol	13.1		73.0	86.0	13.6	88.5	102.2	13.8	87.2	101.0	9.8	85.0	94.8
30)	2-Methylnaphthalene	66.8		66.8	76.3			76.3	76.6		76.6	74.9		74.9
31)	*Acenaphthene-d10	100.0		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
32)	Hexachlorocyclopentadiene	29.3		29.3	45.3			45.3	31.6		31.6	30.7		30.7
33)	2,4,6-Trichlorophenol			82.8	82.8		97.4	97.4		95.1	95.1		92.5	92.5
34)	2,4,5-Trichlorophenol			76.6	76.6		92.5	92.5		91.1	91.1		88.4	88.4
35)	2-Chloronaphthalene	68.9		68.9	78.4		0.1	78.5	78.6		78.6	73.4		73.4
36)	#2-Fluorobiphenyl	75.0		75.0	85.6		6.1	86.7	85.6		85.6	79.9	0.1	80.0
37)	2-Nitroaniline	100.2		100.2	104.7			104.7	108.1		108.1	99.4		99.4
38)	Dimethylphthalate	93.1		93.1	99.2			99.2	98.4		98.4	94.0		94.0
39)	Acenaphthylene	82.3		82.3	92.8		0.1	92.9	92.5		92.5	88.3	0.1	88.4
40)	2,6-Dinitrotoluene	79.7		79.7	86.0			86.0	83.5		83.5	80.6		80.6
41)	3-Nitroaniline	95.8		95.8	96.1			96.1	97.9		97.9	96.4		96.4

TABLE 29. (contd)

No.	Compound Name	Fraction	Total Rec			Total Rec			Total Rec			Total Rec		
			Ext 6 A	Ext 6 BN	Ext 7 A	Ext 7 BN	Ext 7	Ext 8 A	Ext 8 BN	Ext 9 A	Ext 9 BN	Ext 9 A	Ext 9 BN	Ext 9
42)	Acenaphthene		87.8		115.4	99.8	115.4	0.1	99.9	97.6	0.0	97.6	92.9	93.1
43)	2,4-Dinitrophenol		80.3		80.3		80.3	100.1	100.1	100.1	100.1	94.0	94.0	94.0
44)	4-Nitrophenol		78.1	0.6	78.1	0.6	78.1	96.1	96.1	96.1	96.1	92.6	92.6	93.0
45)	Dibenzofuran		81.1	81.1	81.1	81.1	81.1	90.5	90.5	90.5	90.5	75.2	75.2	75.3
46)	2,4-Dinitrotoluene		84.3	84.3	84.3	84.3	84.3	88.3	88.3	88.3	88.3	84.6	84.6	84.6
47)	Diethylphthalate		90.3	0.1	90.4	94.4	94.6	93.2	93.2	93.4	0.2	93.4	89.8	90.0
48)	4-Chlorophenyl-phenylether		84.0	84.0	84.0	95.0	95.1	93.4	93.4	93.4	93.4	87.9	87.9	87.9
49)	Fluorene		92.2	92.2	92.2	103.3	103.3	103.3	103.3	103.3	103.3	97.1	97.1	97.2
50)	4-Nitroaniline		83.9	83.9	83.9	79.0	79.0	100.0	100.0	100.0	100.0	73.7	73.7	73.7
51)	Phenanthrene-d10		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol		89.3	89.3	89.3	113.2	113.2	108.1	108.1	108.1	108.1	100.8	100.8	100.8
53)	N-Nitrosodiphenylamine		103.5	103.5	103.5	111.3	111.3	110.8	110.8	110.8	110.8	97.8	97.8	97.8
54)	#2,4,6-Tribromophenol		0.3	84.9	85.2	0.4	103.8	102.2	102.2	102.2	102.2	93.7	93.7	93.7
55)	4-Bromophenyl-phenylether		87.4	87.4	87.4	103.4	103.4	98.3	98.3	98.3	98.3	69.5	69.5	69.6
56)	Hexachlorobenzene		62.8	62.8	62.8	82.0	82.0	66.0	66.0	66.0	66.0	96.5	96.5	96.5
57)	Pentachlorophenol		80.4	80.4	80.4	103.8	103.8	101.6	101.6	101.6	101.6	93.3	93.3	93.3
58)	Phenanthrene		88.3	88.3	88.3	100.8	100.8	98.4	98.4	98.4	98.4	91.7	91.7	91.7
59)	Anthracene		85.2	85.2	85.2	100.4	100.4	96.6	96.6	96.6	96.6	90.3	90.3	90.7
60)	Di-n-butylphthalate		86.8	86.8	86.8	94.5	94.5	94.4	94.4	94.4	94.4	89.4	89.4	89.5
61)	Fluoranthene		82.9	82.9	82.9	96.9	96.9	93.9	93.9	93.9	93.9	100.0	100.0	100.0
62)	*Chrysene-d12		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
63)	Pyrene		94.7	94.7	94.7	101.6	101.6	101.7	101.7	101.7	101.7	97.7	97.7	97.8
64)	#Terphenyl-d14		79.7	79.7	79.7	84.1	84.1	84.3	84.3	84.3	84.3	82.2	82.2	82.4
65)	Butylbenzylphthalate		96.5	96.5	96.5	99.7	99.7	99.9	99.9	99.9	99.9	97.7	97.7	97.9
66)	3,3'-Dichlorobenzidine		86.3	86.3	86.3	88.7	88.7	88.8	88.8	88.8	88.8	90.6	90.6	90.6
67)	Benzo(a)anthracene		97.7	97.7	97.7	106.7	106.7	113.6	113.6	113.6	113.6	101.8	101.8	101.9
68)	Chrysene		93.3	93.3	93.3	99.3	99.3	99.5	99.5	99.5	99.5	94.8	94.8	94.8
69)	Bis(2-Ethylhexyl)phthalate		93.0	93.0	93.0	94.1	94.1	94.5	94.5	94.5	94.5	93.2	93.2	93.6
70)	*Perylene-d12		100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
71)	Di-n-octylphthalate		84.5	84.5	84.5	89.1	89.1	89.2	89.2	89.2	89.2	88.6	88.6	88.6
72)	Benzo(b)fluoranthene		103.5	103.5	103.5	117.9	117.9	118.0	118.0	118.0	118.0	107.4	107.4	107.5
73)	Benzo(k)fluoranthene		90.9	90.9	90.9	96.2	96.2	91.4	91.4	91.4	91.4	88.1	88.1	88.2
74)	Benzo(a)pyrene		86.4	86.4	86.4	98.7	98.7	99.4	99.4	99.4	99.4	93.9	93.9	94.0
75)	Indeno(1,2,3-cd)pyrene		87.7	87.7	87.7	98.9	98.9	98.9	98.9	98.9	98.9	93.7	93.7	93.7
76)	Dibenz(a,h)anthracene		81.9	81.9	81.9	92.7	92.7	93.4	93.4	93.4	93.4	87.4	87.4	87.4
77)	Benzo(g,h,i)perylene		85.9	85.9	85.9	99.5	99.5	99.5	99.5	99.5	99.5	93.1	93.1	93.1
NPH (Paraffin Hydrocarbon, C12-C14)			19.1	19.1	19.1	25.6	25.6	17.0	17.0	17.0	17.0	21.5	21.5	21.6

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 30. Precision Accuracy, and Control Limit Data from Blank Spikes Using the Final PNL Cleanup Method

No.	Compound Name	Mean Recovery	STD (N-1)	-3XSTD	+3XSTD
1)	*1,4-Dichlorobenzene-d4	100.0	0.0	100.0	100.0
2)	#2-Fluorophenol	81.5	5.3	65.4	97.5
3)	#Phenol-d5	82.1	5.1	66.8	97.4
4)	Phenol	68.1	3.7	56.9	79.3
5)	bis(2-Chloroethyl)ether	72.1	6.8		
6)	2-Chlorophenol	75.9	4.7	61.7	90.0
7)	1,3-Dichlorobenzene	64.0	10.9		
8)	1,4-Dichlorobenzene	63.1	10.2	32.5	93.7
9)	Benzyl alcohol	82.4	7.3		
10)	1,2-Dichlorobenzene	67.5	8.4		
11)	2-Methylphenol	79.8	4.3		
12)	bis(2-Chloroisopropyl)ether	103.2	10.8		
13)	4-Methylphenol	78.2	4.5		
14)	N-Nitroso-di-n-propylamine	94.0	10.8	61.7	126.3
15)	Hexachloroethane	41.9	11.6		
16)	*Naphthalene-d8	100.0	0.0	100.0	100.0
17)	#Nitrobenzene-d5	76.8	8.6	50.9	102.7
18)	Nitrobenzene	71.2	5.4		
19)	Isophorone	74.3	4.9		
20)	2-Nitrophenol	88.8	5.2		
21)	2,4-Dimethylphenol	81.9	6.0		
22)	Benzoic acid	82.5	7.2		
23)	bis(2-Chloroethoxy)methane	87.5	5.9		
24)	2,4-Dichlorophenol	87.0	4.8		
25)	1,2,4-Trichlorobenzene	65.6	9.2	38.1	93.1
26)	Naphthalene	80.6	5.7		
27)	4-Chloroaniline	74.8	14.4		
28)	Hexachlorobutadiene	40.1	10.1		
29)	4-Chloro-3-methylphenol	95.7	6.1	77.3	114.1
30)	2-Methylnaphthalene	71.3	4.5		
31)	*Acenaphthene-d10	100.0	0.0	100.0	100.0
32)	Hexachlorocyclopentadiene	29.1	8.4		
33)	2,4,6-Trichlorophenol	92.1	5.5		
34)	2,4,5-Trichlorophenol	87.0	5.4		
35)	2-Chloronaphthalene	72.4	5.2		
36)	#2-Fluorobiphenyl	77.7	8.3	52.7	102.7
37)	2-Nitroaniline	99.2	7.8		
38)	Dimethylphthalate	93.1	6.4		
39)	Acenaphthylene	86.8	5.5		
40)	2,6-Dinitrotoluene	80.0	5.3		
41)	3-Nitroaniline	93.9	6.2		

TABLE 30. (contd)

No.	Compound Name	Recovery	STD (N-1)	-3XSTD	+3XSTD
42)	Acenaphthene	91.9	5.7	74.9	108.9
43)	2,4-Dinitrophenol	97.0	11.2		
44)	4-Nitrophenol	90.9	7.8	67.6	114.2
45)	Dibenzofuran	81.7	6.0		
46)	2,4-Dinitrotoluene	83.5	5.7	66.4	100.6
47)	Diethylphthalate	89.6	5.1		
48)	4-Chlorophenyl-phenylether	87.1	6.1		
49)	Fluorene	95.4	6.2		
50)	4-Nitroaniline	80.6	15.2		
51)	*Phenanthrene-d10	100.0	0.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol	102.3	8.3		
53)	N-Nitrosodiphenylamine	106.4	5.8		
54)	#2,4,6-Tribromophenol	98.1	5.9	80.4	115.8
55)	4-Bromophenyl-phenylether	93.1	6.9		
56)	Hexachlorobenzene	61.2	13.4		
57)	Pentachlorophenol	98.1	8.1	73.9	122.4
58)	Phenanthrene	93.5	5.6		
59)	Anthracene	91.3	6.4		
60)	Di-n-butylphthalate	90.7	5.6		
61)	Fluoranthene	88.7	6.3		
62)	*Chrysene-d12	100.0	0.0	100.0	100.0
63)	Pyrene	97.0	6.2	78.5	115.5
64)	#Terphenyl-d14	81.0	7.1	59.6	102.4
65)	Butylbenzylphthalate	98.0	5.5		
66)	3,3'-Dichlorobenzidine	82.8	8.3		
67)	Benzo(a)anthracene	101.0	6.9		
68)	Chrysene	96.8	6.4		
69)	Bis(2-Ethylhexyl)phthalate	93.0	4.5		
70)	*Perylene-d12	100.0	0.0	100.0	100.0
71)	Di-n-octylphthalate	88.3	4.1		
72)	Benzo(b)fluoranthene	108.7	6.0		
73)	Benzo(k)fluoranthene	91.5	3.7		
74)	Benzo(a)pyrene	93.7	5.7		
75)	Indeno(1,2,3-cd)pyrene	93.8	6.4		
76)	Dibenz(a,h)anthracene	87.8	5.0		
77)	Benzo(g,h,i)perylene	94.4	6.5		
	NPH (Paraffin Hydrocarbon, C12-C14)	16.8	4.8		

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 31. Recovery Loss from the Use of Final PNL Cleanup Procedure

No.	Compound Name	Mean Before Cleanup	Mean After Cleanup	Difference	Percent Change
1)	*1,4-Dichlorobenzene-d4	100.0	100.0	0.0	0.0
2)	#2-Fluorophenol	105.8	81.5	-24.3	-23.0
3)	#Phenol-d5	99.7	82.1	-17.6	-17.7
4)	Phenol	91.9	68.1	-23.8	-25.9
5)	bis(2-Chloroethyl)ether	115.7	72.1	-43.6	-37.7
6)	2-Chlorophenol	98.9	75.9	-23.0	-23.3
7)	1,3-Dichlorobenzene	97.9	64.0	-34.0	-34.7
8)	1,4-Dichlorobenzene	94.8	63.1	-31.7	-33.4
9)	Benzyl alcohol	99.6	82.4	-17.1	-17.2
10)	1,2-Dichlorobenzene	97.7	67.5	-30.3	-31.0
11)	2-Methylphenol	92.3	79.8	-12.5	-13.6
12)	bis(2-Chloroisopropyl) ether	104.6	103.2	-1.5	-1.4
13)	4-Methylphenol	94.5	78.2	-16.3	-17.3
14)	N-Nitroso-di-n-propylamine	105.7	94.0	-11.7	-11.1
15)	Hexachloroethane	96.5	41.9	-54.6	-56.6
16)	*Naphthalene-d8	100.0	100.0	0.0	0.0
17)	#Nitrobenzene-d5	94.4	76.8	-17.6	-18.7
18)	Nitrobenzene	94.7	71.2	-23.5	-24.8
19)	Isophorone	97.8	74.3	-23.5	-24.0
20)	2-Nitrophenol	93.0	88.8	-4.1	-4.4
21)	2,4-Dimethylphenol	90.4	81.9	-8.5	-9.4
22)	Benzoic acid	89.3	82.5	-6.7	-7.5
23)	bis(2-Chloroethoxy)methane	93.1	87.5	-5.7	-6.1
24)	2,4-Dichlorophenol	95.5	87.0	-8.5	-8.9
25)	1,2,4-Trichlorobenzene	90.7	65.6	-25.1	-27.7
26)	Naphthalene	85.9	80.6	-5.2	-6.1
27)	4-Chloroaniline	74.1	74.8	0.7	0.9
28)	Hexachlorobutadiene	90.0	40.1	-49.9	-55.5
29)	4-Chloro-3-methylphenol	95.8	95.7	-0.1	-0.1
30)	2-Methylnaphthalene	75.5	71.3	-4.3	-5.6
31)	*Acenaphthene-d10	100.0	100.0	0.0	0.0
32)	Hexachlorocyclopentadiene	85.0	29.1	-55.9	-65.8
33)	2,4,6-Trichlorophenol	97.2	92.1	-5.0	-5.2
34)	2,4,5-Trichlorophenol	94.0	87.0	-7.1	-7.5
35)	2-Chloronaphthalene	93.1	72.4	-20.7	-22.2
36)	#2-Fluorobiphenyl	94.3	77.7	-16.6	-17.6
37)	2-Nitroaniline	66.5	99.2	32.8	49.3
38)	Dimethylphthalate	98.5	93.1	-5.4	-5.5
39)	Acenaphthylene	91.5	86.8	-4.7	-5.2
40)	2,6-Dinitrotoluene	101.7	80.0	-21.7	-21.3
41)	3-Nitroaniline	90.3	93.9	3.6	4.0

TABLE 31. (contd)

No.	Compound Name	Mean Before Cleanup	Mean After Cleanup	Difference	Percent Change
42)	Acenaphthene	96.7	91.9	-4.8	-5.0
43)	2,4-Dinitrophenol	109.0	97.0	-12.0	-11.0
44)	4-Nitrophenol	110.0	90.9	-19.1	-17.4
45)	Dibenzofuran	92.8	81.7	-11.1	-12.0
46)	2,4-Dinitrotoluene	100.5	83.5	-17.0	-16.9
47)	Diethylphthalate	102.5	89.6	-12.9	-12.6
48)	4-Chlorophenyl-phenylether	94.9	87.1	-7.8	-8.2
49)	Fluorene	98.3	95.4	-2.9	-3.0
50)	4-Nitroaniline	94.1	80.6	-13.5	-14.4
51)	*Phenanthrene-d10	100.0	100.0	0.0	0.0
52)	4,6-Dinitro-2-methylphenol	100.5	102.3	1.8	1.8
53)	N-Nitrosodiphenylamine	121.1	106.4	-14.7	-12.1
54)	#2,4,6-Tribromophenol	97.1	98.1	1.0	1.1
55)	4-Bromophenyl-phenylether	91.4	93.1	1.6	1.8
56)	Hexachlorobenzene	94.1	61.2	-32.8	-34.9
57)	Pentachlorophenol	103.1	98.1	-4.9	-4.8
58)	Phenanthrene	91.0	93.5	2.5	2.7
59)	Anthracene	92.6	91.3	-1.2	-1.3
60)	Di-n-butylphthalate	89.9	90.7	0.8	0.9
61)	Fluoranthene	86.0	88.7	2.7	3.1
62)	*Chrysene-d12	100.0	100.0	0.0	0.0
63)	Pyrene	90.6	97.0	6.4	7.1
64)	#Terphenyl-d14	84.3	81.0	-3.3	-3.9
65)	Butylbenzylphthalate	92.2	98.0	5.8	6.3
66)	3,3'-Dichlorobenzidine	118.6	82.8	-35.8	-30.2
67)	Benzo(a)anthracene	94.6	101.0	6.3	6.7
68)	Chrysene	86.2	96.8	10.5	12.2
69)	Bis(2-Ethylhexyl)phthalate	89.2	93.0	3.8	4.3
70)	*Perylene-d12	100.0	100.0	0.0	0.0
71)	Di-n-octylphthalate	87.7	88.3	0.6	0.7
72)	Benzo(b)fluoranthene	98.4	108.7	10.3	10.4
73)	Benzo(k)fluoranthene	89.1	91.5	2.4	2.7
74)	Benzo(a)pyrene	94.6	93.7	-0.9	-1.0
75)	Indeno(1,2,3-cd)pyrene	92.9	93.8	0.9	1.0
76)	Dibenz(a,h)anthracene	88.4	87.8	-0.6	-0.7
77)	Benzo(g,h,i)perylene	93.2	94.4	1.2	1.3
NPH (Paraffin Hydrocarbon, C12-C14)		62.6	16.8	-45.8	-73.2

* Compound is an Internal Standard

Compound is a Surrogate Spike

TABLE 32. Calculation for Estimating Mean Recoveries for Compounds Spiked in Single Shell Tank Samples B110 and U110 Using the Final PNL Cleanup Procedure (Mean Recovery = P2BN-(NMS-BN - Blank BN) + Acid)

Sample: Single Shell Tanks B110, U110	P2BN Mean STD (N-1)	NMS-BN Mean STD (N-1)	Blank BN Mean STD (N-1)	Acid Mean STD (N-1)	Mean Recovery STD (N-1)
No. Compound Name					
42) Acenaphthene	71.3	10.5	89.2	1.8	5.7
43) 2,4-Dinitrophenol	0.0	0.0	0.0	0.0	0.0
44) 4-Nitrophenol	0.0	0.0	0.0	0.0	0.0
45) Dibenzofuran	70.8	9.6	85.1	0.0	81.7
46) 2,4-Dinitrotoluene	73.7	10.4	85.1	2.4	83.5
47) Diethylphthalate	75.5	10.6	96.3	2.1	80.7
48) 4-Chlorophenyl-phenylether	42.0	5.3	0.4	0.6	87.1
49) Fluorene	71.7	9.9	85.7	1.9	95.4
50) 4-Nitroaniline	90.8	54.7	134.2	28.9	80.6
51) *Phenanthrene-d10	100.0	0.0	100.0	0.0	100.0
52) 4,6-Dinitro-2-methylphenol	0.0	0.0	0.0	0.0	0.0
53) N-Nitrosodiphenylamine	66.3	17.2	99.0	8.7	106.4
54) #2,4,6-Tribromophenol	0.0	0.0	0.0	0.0	0.4
55) 4-Bromophenyl-phenylether	68.2	9.0	83.2	9.4	93.1
56) Hexachlorobenzene	72.7	10.5	84.7	8.4	61.2
57) Pentachlorophenol	0.0	0.0	0.0	0.0	0.0
58) Phenanthrene	73.6	9.5	87.4	3.9	93.4
59) Anthracene	56.7	18.3	84.0	5.8	91.3
60) Di-n-butylphthalate	73.9	9.2	85.8	4.4	81.7
61) Fluoranthene	70.8	9.2	82.4	6.2	88.7
62) *Chrysene-d12	100.0	0.0	100.0	0.0	100.0
63) Pyrene	68.8	8.5	84.8	2.8	97.0
64) #Terphenyl-d14	67.2	6.8	77.1	0.8	80.9
65) Butylbenzylphthalate	70.1	8.1	86.5	0.0	97.7
66) 3,3'-Dichlorobenzidine	47.3	45.9	93.0	9.8	82.8
67) Benzo(a)anthracene	66.8	10.5	87.2	0.6	100.9
68) Chrysene	70.4	9.1	85.3	0.4	96.7
69) Bis(2-Ethylhexyl)phthalate	64.1	7.6	82.5	0.0	83.6
70) *Perylene-d12	100.0	0.0	100.0	0.0	100.0
71) Di-n-octylphthalate	66.9	10.1	81.1	2.7	79.6
72) Benzo(b)fluoranthene	74.3	10.9	95.8	9.2	108.6
73) Benzo(k)fluoranthene	66.9	16.0	92.3	22.6	91.5
74) Benzo(a)pyrene	51.9	22.2	83.4	5.6	93.7
75) Indeno(1,2,3-cd)pyrene	73.1	10.7	93.0	5.7	93.8
76) Dibenz(a,h)anthracene	65.7	9.6	81.7	3.6	87.8
77) Benzo(g,h,i)perylene	71.6	10.9	93.4	3.7	94.4
NPH (Paraffin Hydrocarbon, C12-C14)					

* Compound is an Internal Standard
Compound is a Surrogate Spike

TABLE 32. (contd)

Sample: Single Shell Tanks B*10, U110		P2BN	NMS-BN		Blank BN		Acid		Mean	
No. Compound Name		Mean	STD (N-1)	Mean	STD (N-1)	Mean	STD (N-1)	Mean	STD (N-1)	Mean
1) *1,4-Dichlorobenzene-d4		100.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	100.0
2) #2-Fluorophenol		0.3	0.1	0.5	0.2	0.0	0.0	46.6	11.4	46.6
3) #Phenol-d5		2.4	1.4	3.8	0.4	3.2	0.7	52.8	11.6	54.5
4) Phenol		3.3	0.9	1.3	1.9	0.0	0.0	52.2	8.4	54.2
5) bis(2-Chloroethyl)ether		63.6	9.1	79.8	0.8	72.1	6.8	0.4	0.2	56.3
6) 2-Chlorophenol		0.7	0.2	1.1	0.2	1.4	0.6	49.6	6.5	50.5
7) 1,3-Dichlorobenzene		57.2	7.9	68.2	1.3	63.9	10.9	0.3	0.2	53.3
8) 1,4-Dichlorobenzene		57.2	8.7	75.0	1.5	63.1	10.2	0.3	0.2	45.6
9) Benzyl alcohol		66.8	8.7	79.7	1.0	82.4	7.3	1.5	0.8	71.0
10) 1,2-Dichlorobenzene		57.8	8.4	69.4	0.7	67.4	8.4	0.3	0.1	56.2
11) 2-Methylphenol		16.4	7.3	21.2	0.7	17.5	7.2	38.1	7.2	50.7
12) bis(2-Chloroisopropyl)ether		82.5	12.4	93.7	4.1	103.2	10.8	0.5	0.3	92.5
13) 4-Methylphenol		13.0	5.6	16.6	0.9	12.4	1.5	43.5	8.3	52.3
14) N-Nitroso-di-n-propylamine		75.6	11.3	88.6	3.5	94.0	10.8	0.8	0.7	81.8
15) Hexachloroethane		57.5	8.5	68.9	0.4	41.9	11.6	0.2	0.1	30.8
16) *Naphthalene-d8		100.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	100.0
17) #Nitrobenzene-d5		65.2	8.7	82.8	0.7	76.8	8.6	0.3	0.2	59.5
18) Nitrobenzene		74.4	15.1	80.1	0.4	71.2	5.4	0.6	0.3	66.2
19) Isophorone		64.9	8.8	77.1	0.8	66.9	23.7	1.1	0.9	55.8
20) 2-Nitrophenol		0.3	0.1	0.6	0.2	0.8	0.3	62.3	8.7	62.9
21) 2,4-Dimethylphenol		27.5	21.0	55.0	1.5	49.7	4.0	5.3	4.0	27.5
22) Benzoic acid		0.0	0.0	0.0	0.0	0.0	0.0	37.1	16.8	37.1
23) bis(2-Chloroethoxy)methane		71.1	10.0	83.0	1.5	87.5	5.9	1.0	0.8	76.6
24) 2,4-Dichlorophenol		0.7	0.2	1.1	0.3	1.1	1.1	63.6	8.4	64.5
25) 1,2,4-Trichlorobenzene		62.4	8.2	75.8	1.1	65.6	9.2	0.7	0.4	52.9
26) Naphthalene		62.7	8.2	76.5	1.1	80.6	5.7	1.8	1.2	68.6
27) 4-Chloroaniline		55.4	45.4	62.1	53.5	74.8	14.4	0.1	0.0	68.2
28) Hexachlorobutadiene		64.4	8.9	75.3	2.3	40.1	10.1	0.5	0.3	29.7
29) 4-Chloro-3-methylphenol		12.6	5.3	15.4	0.7	12.6	1.6	57.0	9.7	66.7
30) 2-Methylnaphthalene		53.1	8.6	59.4	5.6	71.3	4.5	1.0	0.8	65.9
31) *Acenaphthene-d10		100.0	0.0	100.0	0.0	100.0	0.0	100.0	0.0	100.0
32) Hexachlorocyclopentadiene		54.3	6.6	69.1	9.8	29.1	8.4	0.3	0.2	14.5
33) 2,4,6-Trichlorophenol		0.0	0.0	0.0	0.0	0.0	0.0	66.6	10.2	66.6
34) 2,4,5-Trichlorophenol		0.0	0.0	0.0	0.0	0.0	0.0	64.9	9.9	64.9
35) 2-Chloronaphthalene		63.8	9.3	80.4	0.8	72.4	5.2	1.0	0.9	56.7
36) #2-Fluorobiphenyl		63.8	7.2	83.2	0.7	77.7	8.3	0.9	0.8	59.3
37) 2-Nitroaniline		76.7	13.2	92.2	2.0	99.2	7.8	1.1	1.0	84.9
38) Dimethylphthalate		71.3	9.1	83.7	3.7	93.1	6.4	1.6	1.4	82.3
39) Acenaphthylene		54.0	11.5	79.5	0.7	86.8	5.5	0.9	0.8	62.2
40) 2,6-Dinitrotoluene		71.9	9.9	84.7	3.6	80.0	5.3	0.9	0.9	68.1
41) 3-Nitroaniline		86.6	19.0	106.6	9.4	93.9	6.2	0.0	0.0	73.9

TABLE 33. Estimated Precision, Accuracy and Control Limits for SST Semivolatile Organic Analysis

Sample: Single Shell Tanks B110, U110		Mean			
No.	Compound Name	Recovery	STD (N-1)	-3XSTD	+3XSTD
1)	*1,4-Dichlorobenzene-d4	100.0	0.0	100.0	100.0
2)	#2-Fluorophenol	46.6	11.4	12.3	80.9
3)	#Phenol-d5	54.5	11.7	19.3	89.7
4)	Phenol	54.2	8.7	28.2	80.2
5)	bis(2-Chloroethyl)ether	56.3	11.3		
6)	2-Chlorophenol	50.5	6.6	30.8	70.2
7)	1,3-Dichlorobenzene	53.3	13.4		
8)	1,4-Dichlorobenzene	45.6	13.3	5.6	85.6
9)	Benzyl alcohol	71.0	11.4		
10)	1,2-Dichlorobenzene	56.2	11.9		
11)	2-Methylphenol	50.7	10.3		
12)	bis(2-Chloroisopropyl)ether	92.5	15.9		
13)	4-Methylphenol	52.3	10.1		
14)	N-Nitroso-di-n-propylamine	81.8	15.2	36.1	127.4
15)	Hexachloroethane	30.8	14.4		
16)	*Naphthalene-d8	100.0	0.0	100.0	100.0
17)	#Nitrobenzene-d5	59.5	12.2	22.9	96.1
18)	Nitrobenzene	66.2	16.1		
19)	Isophorone	55.8	25.3		
20)	2-Nitrophenol	62.9	8.7		
21)	2,4-Dimethylphenol	27.5	21.7		
22)	Benzoic acid	37.1	16.8		
23)	bis(2-Chloroethoxy)methane	76.6	11.5		
24)	2,4-Dichlorophenol	64.5	8.4		
25)	1,2,4-Trichlorobenzene	52.8	12.3	15.9	89.6
26)	Naphthalene	68.6	10.0		
27)	4-Chloroaniline	68.1	68.7		
28)	Hexachlorobutadiene	29.7	13.3		
29)	4-Chloro-3-methylphenol	66.7	11.2	33.1	100.3
30)	2-Methylnaphthalene	65.9	9.2		
31)	*Acenaphthene-d10	100.0	0.0	100.0	100.0
32)	Hexachlorocyclopentadiene	14.5	8.4		
33)	2,4,6-Trichlorophenol	66.6	10.2		
34)	2,4,5-Trichlorophenol	64.9	9.9		
35)	2-Chloronaphthalene	56.7	10.7		
36)	*2-Fluorobiphenyl	59.3	11.0	26.2	92.3
37)	2-Nitroaniline	84.9	15.3		
38)	Dimethylphthalate	82.3	10.6		
39)	Acenaphthylene	62.2	12.8		
40)	2,6-Dinitrotoluene	68.1	10.6		
41)	3-Nitroaniline	74.0	20.3		

TABLE 33. (contd)

Sample: Single Shell Tanks B110, U110		Mean			
No.	Compound Name	Recovery	STD (N-1)	-3XSTD	+3XSTD
42)	Acenaphthene	75.2	11.8	39.7	110.6
43)	2,4-Dinitrophenol	64.3	8.7		
44)	4-Nitrophenol	76.6	15.4	30.4	122.8
45)	Dibenzofuran	68.7	11.4		
46)	2,4-Dinitrotoluene	73.5	11.7	38.5	108.5
47)	Diethylphthalate	61.7	30.2		
48)	4-Chlorophenyl-phenylether	129.5	8.1		
49)	Fluorene	82.8	11.6		
50)	4-Nitroaniline	37.9	60.0		
51)	*Phenanthrene-d10	100.0	0.0	100.0	100.0
52)	4,6-Dinitro-2-methylphenol	71.9	10.9		
53)	N-Nitrosodiphenylamine	74.8	18.4		
54)	#2,4,6-Tribromophenol	67.3	10.6	35.4	99.3
55)	4-Bromophenyl-phenylether	79.5	11.1		
56)	Hexachlorobenzene	50.8	14.9		
57)	Pentachlorophenol	88.7	16.4	39.4	138.0
58)	Phenanthrene	81.2	10.4		
59)	Anthracene	65.3	18.5		
60)	Di-n-butylphthalate	71.7	28.9		
61)	Fluoranthene	78.8	9.4		
62)	*Chrysene-d12	100.0	0.0	100.0	100.0
63)	Pyrene	82.4	10.2	51.9	112.9
64)	#Terphenyl-d14	72.5	9.8	43.0	102.0
65)	Butylbenzylphthalate	83.0	9.9		
66)	3,3'-Dichlorobenzidine	37.0	46.2		
67)	Benzo(a)anthracene	82.0	12.6		
68)	Chrysene	83.5	11.2		
69)	Bis(2-Ethylhexyl)phthalate	66.9	30.2		
70)	*Perylene-d12	100.0	0.0	100.0	100.0
71)	Di-n-octylphthalate	67.2	29.6		
72)	Benzo(b)fluoranthene	88.7	13.0		
73)	Benzo(k)fluoranthene	67.8	27.4		
74)	Benzo(a)pyrene	63.3	22.2		
75)	Indeno(1,2,3-cd)pyrene	75.5	11.2		
76)	Dibenz(a,h)anthracene	73.2	10.3		
77)	Benzo(g,h,i)perylene	74.0	12.2		
NPH (Paraffin Hydrocarbon, C12-C14)					

* Compound is an Internal Standard

Compound is a Surrogate Spike

APPENDIX A

PNL-AO-120, PROCEDURE FOR EXTRACTION OF SINGLE SHELL TANK SAMPLES FOR THE ANALYSIS OF SEMIVOLATILE ORGANIC COMPOUNDS

TITLE: PNL-ALO-120, PROCEDURE FOR EXTRACTION OF SINGLE SHELL TANK SAMPLES
FOR THE ANALYSIS OF SEMIVOLATILE ORGANIC COMPOUNDS

APPLICABILITY

This procedure describes the method used for the extraction of single shell tank samples and the subsequent preparation of the extract for the analysis of semivolatile organic compounds. This procedure may be used to support work performed in compliance with EPA Contract Laboratory Program (CLP) or other projects/programs as elected.

DEFINITIONS

None

RESPONSIBLE STAFF

Cognizant Scientist
Analyst

PROCEDURE

- 1.0 This procedure is for extracting nonvolatile and semivolatile organic compounds from solids such as soils, sludges and other waste samples including single shell tank samples. The sonication process ensures intimate contact of the sample matrix with the extraction solvent. The method is divided into two sections based on the expected concentration of organics in the sample. The low concentration method (individual extractable organic components ≤ 20 mg/Kg) uses a larger sample size and a more rigorous extraction procedure. The high concentration method (>20 mg/Kg individual extractable organic components) uses a smaller sample, less extraction solvent and is much simpler to perform. The procedure provides for adjustment of pH for samples which do not meet the pH 5-11 extraction criteria. Extracts are filtered and evaporated to a volume suitable for optional GPC cleanup procedures followed by screening and/or analyses by GC/FID or GC/MS.
- 2.0 Apparatus
 - a) Mortar and pestle.
 - b) 1-mm standard sieve.
 - c) Sonicator - Ultrasonic cell disrupter: Heat Systems - Ultrasonic, Inc. Model W-385 (475 watt) sonicator or equivalent (power wattage must be a minimum of 375 with pulsing capability) using a no. 208 or no. 305 3/4" disrupter horn for low level analyses, and a no. 200 1/2" tapped disrupter horn and no. 419 1/8" standard taper microtip probe.
 - d) Oven, drying.
 - e) Desiccator.

- f) Pasteur pipets, 1-ml.
- g) Beakers 400-ml.
- h) Glass scintillation vials, →20-ml, with screw cap and Teflon liners.
- i) Buchner funnel
- j) Filter paper, Whatman no. 41 or equivalent.
- k) Glass conical funnel
- l) Filter flask, 500-ml.
- m) Concentrator tube, 10-ml graduated (Kontes K-570050-1025 or equivalent).
- n) Evaporator flask, 500-ml (Kontes K-570001-0500 or equivalent).
- o) Snyder columns, three-ball macro (Kontes K-503000-0121 or equivalent) and two-ball micro.
- p) Water bath evaporator, 8-position S-evap, SS boiling water bath Organomation Associates Catalog #12020 or equivalent.
- q) N-evap, 12 position with SS water bath Organomation Associates Catalog #11155 or equivalent.
- r) Boiling Chips, 10/40 mesh (silicon carbide or equivalent) solvent extracted or fired at 450°C for 24 hours, cooled in a desiccator and stored in vial with screw cap.
- s) Balance, capable of accurately weighing 0.01 gram.
- t) Spatula, stainless steel or Teflon.
- u) Drying column, 20-mm ID. Pyrex^R chromatographic column with Pyrex^R glass wool at bottom. Note, the glass column may also be fitted with a Teflon stopcock and a fritted glass disc to retain the drying compound, however they are difficult to decontaminate after highly contaminated extracts have been passed through them. Use a small pad of glass wool to retain the drying compound. Prewash the glass wool with 50 ml of acetone followed with 50 ml of elution solvent prior to packing the column with the drying compound.
- v) Centrifuge.
- w) pH meter.

3.0 Reagents

- a) Sodium sulfate, powdered anhydrous and reagent grade, heated at 450°C for 8 hours, cooled in a desiccator and stored in a glass bottle. Baker anhydrous powder, catalog #73898 or equivalent.
- b) Extraction solvents.
- c) 1:1 sulfuric acid (H₂SO₄) and water
- d) 10 N NaOH

4.0 Standards

- a) Base/neutral acid surrogate solution.

Surrogate standards are added to all samples, blanks, matrix spikes, matrix spike duplicates, and calibration solutions; the compounds specified are:

<u>Base Neutrals (100 µg/1.0 ml)</u>	<u>Acids (200 µg/1.0 ml)</u>
nitrobenzene - d ₅	phenol - d ₅
terphenyl - d ₁₄	2-fluorophenol
2-fluorobiphenyl	2,4,6-tribromophenol
optional - 1 additional	optional - 1 additional

- b) Matrix standard spiking solutions.

<u>Base/Neutrals (100 µg/1.0 ml)</u>	<u>Acids (200 µg/1.0 ml)</u>
1,2,4-trichlorobenzene	pentachlorophenol
acenaphthene	phenol
2,4-dinitrotoluene	2-chlorophenol
pyrene	4-chloro-3-methylphenol
N-nitroso-di-n-propylamine	4-nitrophenol
1,4-dichlorobenzene	

Prepare separate surrogate and matrix spiking solutions at a concentration of 100 µg/1.0 ml for base/neutral and 200 µg/1.0 ml for acids in methanol. Store the solutions at 4°C (±2°C) in amber glass Teflon-sealed containers. The solutions must be replaced after twelve months, or sooner, if comparison with quality control check samples indicate a problem.

5.0 Safety

- a) Observe the general laboratory safety rules.
- b) Follow the requirements of applicable RWP's.
- c) Observe regulations for handling, storage and disposal of hazardous organic chemicals.

6.0 Quality Control

- a) A reagent blank shall be subjected to exactly the same procedure for each set of samples.
- b) A matrix spike and matrix spike duplicate [see section 4.4. (b)] shall be performed on one sample from each set of samples having the same sample matrix and/or to one sample from each group of 20 samples received as a group and/or originating from the same sample matrix. The matrix spike and matrix spike duplicate samples shall be subjected to the same procedure.
- c) Base/neutral and acid surrogate spikes [see section 4.4 (a)] shall be added to all blanks, samples, and matrix spiked samples.

7.0 Analysis

7.1 Initial Sample Preparation

- a) Verify that the sample has been received and logged into the Chemistry and Analysis Section per PNL-SA-34.
- b) Grind and/or mix the sample thoroughly.
- c) Weigh out a subsample, (weight shall be specified by the cognizant scientist) for pH measurement - typically 1 to 2 grams.
- d) Add 1 ml of water per each gram of sample and stir for 1 hour.
- e) Determine pH of sample with glass electrode and pH meter while stirring.

Proceed with steps f and g only if the pH is outside the 5-11 range:

- f) Add 1:1 H_2SO_4 or 10 N NaOH dropwise until the pH is within the 5-11 range. Allow to stir for 5 minutes. Due to sample variability it is recommended the sample be adjusted to pH 7-8.
- g) Record sample weight, volume of water added to sample, pH, and volume of 1:1 H_2SO_4 or 10 N NaOH added on the Shielded Analytical Laboratory (SAL) sample preparation instruction sheet (Refer to PNL-SA-50).
- h) Dispose of the sample used for the pH and acid/base adjustment measurement.

7.2 Extraction of Sample for Medium Level Semivolatile Organics

- a) Weigh out a subsample for extraction. The weight shall be specified by the Cognizant Scientist and shall be recorded on

the SAL sample preparation. (EPA suggested weight for the sample is 1 gram). (Refer to PNL-SA-50)

- b) Add sample to a 20-ml vial. Vial shall be cleaned per PNL-SA-41 and just prior to the extraction the vial should be rinsed with methylene chloride. If required, add a proportional amount of acid or base as calculated in steps 4.7.1 f-h. Mix thoroughly into soil
- c) Add anhydrous sodium sulfate at twice the weight of the combined sample and pH adjustment reagents. For 1 gram of sample, add 2 grams of powdered anhydrous sodium sulfate to the sample and mix well.
- d) Add 1 ml of the surrogate spike solution to the vial. [See standard section 4.4 (a)]
- e) If the sample is chosen for matrix spike or matrix spike duplicate add 1 ml of the matrix standard spiking solution. [See standard section 4.4 (b)]
- f) Add whatever volume of methylene chloride needed to bring the final solution volume to 10.0 ml (considering the added volume of surrogate and matrix spike).
- g) Disrupt the sample with the 1/8 inch tapered microtip ultrasonic probe for 2 minutes at an output control setting 5 in the continuous mode. Note: before extraction mix the sample and make sure any lumps of the sodium sulfate are broken up and are free flowing using a clean glass rod or a clean spatula.
- h) Loosely pack a disposable pipet with 2-3 cm glass wool and 2-3 cm of granular anhydrous sodium sulfate on top. Do not pre-rinse the column.
- i) Filter the extract through the glass wool and collect 5.0 ml in a concentrator tube. Centrifugation may also be used in place of or in conjunction with step (j).
- j) Place a micro-snyder column on the concentrator tube. Prewet the Snyder balls by adding a small amount of methylene chloride through the top of the condenser. Gently heat the tube in about a 70-90°C water bath. At the proper rate of distillation the balls of the column will chatter continuously; but the chamber will not flood with vapor.
- k) When the volume in the concentrator tube is below ~0.5 ml remove the apparatus from the water bath and allow to cool ~ 10 minutes. Optionally, the concentrator tube may be placed in an N-evaporator with the water bath at (35°C) and evaporate the solvent to 0.5 ml using a gentle stream of nitrogen gas. The internal walls of the concentrator tube should be washed down with a small volume of methylene chloride during the operation.

During the evaporation, the tube solvent level must be kept below the water level of the bath. The extract should never be allowed to become dry.

- l) Transfer the 0.5 ml concentrate to a 2 ml screw top vial and bring to 1.0 ml from rinsate of the ampule using quantitative transfer. Label the vial with sample ID.
- m) Submit the sample for GC/FID screening or GS/MS screening or GS/MS analysis.
- n) Fill out the organic extraction bench sheet (PNL-SA-45) and submit it with the sample.

7.3 Extraction of Sample for Low Level Semivolatile Organics

- a) Weigh out a sample specified by the cognizant scientist.
(Note: EPA methodology recommends a 30 gram sample for the low level extraction. Record the sample weight on the SAL sample preparation instruction sheet (Refer to PNL-SA-50).
- b) Add the sample to a clean 400 ml beaker. Beaker should be cleaned per PNL-SA-41 and just prior to the extraction the beaker should be rinsed with methylene chloride.
- c) If the sample pH determined in steps 4.7.1 a-e is outside the allowable range of pH 5-11, add a proportional amount of 1:1 H_2SO_4 or 10 N NaOH as determined in steps 4.7.1 f-h to obtain pH 7-8. Mix thoroughly.
- d) Add 2 times the combined sample and pH adjustment reagents weight of sodium sulfate. For example, add 60 grams of anhydrous powdered sodium sulfate for a 29 gram sample with 1 gram of acid added.
- e) Add 100 ml of 1:1 methylene chloride-acetone or a volume equal to 3.3 times the sample weight to the beaker.
- f) Add 1.0 ml of the surrogate spike to the beaker [see standards section (a)]
- g) If the sample is designated for matrix spiking or matrix spike duplicate, add 1 ml of the matrix standard spiking solution [see standards section (c)].
- h) If the sample extract is to be cleaned up using Gel Permeation Chromatography (GPC) add a volume of spiking solutions specified by the cognizant scientist. Record the volume and source of the spiking solutions on the organic extraction bench sheet (PNL-SA-45).
- i) Place the bottom surface of the 3/4 inch sonicator disruptor horn about 1/2 inch below the surface of the solvent. Sonicate for 1 1/2 minutes with the W-385 using no. 208 3/4 inch

standard disruptor horn with the output control knob set at 10 (or no. 305 3/4 inch tapped high gain Q disruptor horn at 5) and mode switch on "1 sec pulse" and % duty cycle at 50%.

- j) Decant the filter extract through a #41 filter paper approximately half filled with granular sodium sulfate in a glass conical funnel. Centrifugation may also be used instead of or in conjunction with the above filtration. A sodium sulfate drying column must be used if the funnel drying method is not employed.
- k) Repeat the above extraction with 2 more 100 ml or a volume equal to 3.3 times the sample weight portions of the 1:1 methylene chloride-acetone mixture.
- l) Transfer the extracts to a Kuderna-Danish (K-D) concentrator consisting of a 10 ml concentrator tube, a 3 ball Snyder column and 500 ml evaporative flask. The concentrator should be cleaned per PNL-SA-41 and just prior to assembly should be rinsed with methylene chloride. One or two clean boiling chips should be added to the concentrator tube during the assembly to prevent bumping.
- m) Place the K-D apparatus on a hot water bath (80 to 90°C) so that the concentrator tube is partially immersed in the hot water and the entire lower rounded surface of the flask is bathed with hot steam. Prewet the Snyder balls by adding a small amount of methylene chloride through the top of the condenser. At the proper rate of distillation the balls of the column will chatter continuously; but the chamber will not flood with vapor.
- n) When the volume in the concentrator tube is below ~5 ml remove the K-D apparatus from the water bath and allow to cool ~ 10 minutes.
- o) If screening is required, make the volume of the residue to 10 ml, 1 ml of the solution in the concentrator tube can be transferred to an autosampler and/or 2 ml vial and submitted for GC/FID screening or GC/MS screening. Normally low level extracts are concentrated as follows:
- p) Place a micro-snyder column on the concentrator tube. Prewet the Snyder balls by adding a small amount of methylene chloride through the top of the condenser. Gently heat the tube in about a 70-90°C water bath. At the proper rate of distillation the balls of the column will chatter continuously; but the chamber will not flood with vapor.
- q) When the volume in the concentrator tube is below ~0.5 ml remove the apparatus from the water bath and allow to cool ~ 10 minutes. Optionally, the concentrator tube may be placed in an N-evaporator with the water bath at (35°C) and evaporate the solvent to 0.5 ml using a gentle stream of nitrogen gas. The

internal walls of the concentrator tube should be washed down with a small volume of methylene chloride during the operation. During the evaporation, the tube solvent level must be kept below the water level of the bath. The extract should never be allowed to become dry.

- r) Transfer the 0.5 ml concentrate to a 2 ml screw top vial and bring the volume to 1.0 ml using rinsates from the concentrator ampule, use quantitative transfer. Label the vial with sample ID.

Submit the sample for GC/FID screening, GC/MS screening or GC/MS analysis.

- s) Fill out the organic extraction bench sheet (PNL-SA-45) and submit it with the sample.

REFERENCES

USA EPA Contract Laboratory Program, Statement of Work for Organic Analysis, 2/88.

USA EPA Test Methods for Evaluating Solid Waste, SW-846, 3rd Edition.

APPENDIX B

PNL-ALO-122, CLEANUP OF HYDROCARBON CONTAINING SAMPLES FOR THE
ANALYSIS OF SEMIVOLATILE ORGANIC COMPOUNDS, Revision 1.0

PNL TECHNICAL PROCEDURE

TITLE: PNL-ALO-122, CLEANUP OF HYDROCARBON CONTAINING SAMPLES FOR THE ANALYSIS OF SEMIVOLATILE ORGANIC COMPOUNDS, REVISION 1.0

APPLICABILITY

This procedure is useful for the analysis of semivolatile compounds in samples containing interfering quantities of normal hydrocarbons. This procedure was developed to remove normal paraffin hydrocarbons (NPH), typically C_{12} through C_{14} , often used as a non-miscible hydrostatic fluid for obtaining radioactive tank waste samples on the Hanford site. Previously, diluting the sample to reduce the quantity injected into the analytical system has been the only available method of dealing with high concentration levels of hydrocarbons. Often the dilution required may be several orders of magnitude which also dilutes the analytes of interest and affects detection limits substantially. Use of this procedure in conjunction with most GC/MS methods does not require dilution and allows the stated detection limits to remain unchanged. Any sample matrix containing excessive quantities of normal hydrocarbons, such as waters, sorbents, or soils, are applicable to this procedure.

DEFINITIONS

NPH (Normal Paraffin Hydrocarbons)
CRQL (Contract Required Quantitation Limits)

RESPONSIBLE STAFF

Cognizant Scientist
Analyst

PROCEDURE

1.0 SCOPE AND APPLICATION

- 1.1 This method is employed following extraction of a waste or soil using other procedures such as PNL-ALO-120 or -344. In this procedure, residues from PNL-ALO-120 or -344 are partitioned into their base-neutral and acidic components. The base-neutral fraction is cleaned using a silica gel column. Analysis of the cleaned fraction is per other methods such as PNL-ALO-345. Unless specifically mentioned within this procedure, there are no further deviations of the methods used to perform extractions or to analyze residues from this preparation.
- 1.2 This method may be used for the analysis of radioactive storage tank wastes which contain hydrocarbons in excessive quantity. This procedure does not allow for distinction between hydrocarbons introduced during the sampling process as contamination and those which are native to the tank wastes. An attempt to quantify any

PNL TECHNICAL PROCEDURE

compound not included in the Semivolatiles Target Compound List (TCL) following use of this procedure without further validation is inappropriate.

- 1.3 This procedure may be used to remove oil from various matrices, such as soil. Before using this procedure, validation employing matrix spikes and establishing control limits for each distinct matrix should be considered.

2.0 SUMMARY OF METHOD

An aliquot of the sample residue produced from a preceding extraction procedure is added to a volume of reagent water. The pH is adjusted to >11 and the base-neutral fraction is extracted into methylene chloride. The pH of the aqueous phase is again adjusted to <2 and the acid fraction is extracted into methylene chloride. Following concentration and exchange into cyclohexane, the base-neutral fraction is eluted through a silica gel (an amorphous silica with weakly acidic properties) column. The hydrocarbon fraction is discarded, followed by collection of the analyte fraction of interest. After concentration, the residue is now ready for GC/MS analysis along with the acid fraction.

3.0 INTERFERENCES

- 3.1 Method interferences may be caused by contaminated solvents, reagents, or glassware. Particular care should be taken when plastic labware of any kind is in the laboratory. Substantial quantities of various phthalate esters, used as plasticisers, can contaminate the sample even from a small amount of solvent exposed to plastic surfaces.
- 3.2 If pre-packed silica cartridges are substituted for glass columns, it is imperative that the manufacturers lot is checked for contamination. Phthalate contamination is a primary obstacle using pre-packed commercial columns.

4.0 APPARATUS AND MATERIALS

- 4.1 Balance: Analytical, capable of accurately weighing 0.01 gram.
- 4.2 Vials: Pre-cleaned 1.5 mL amber screw top with teflon lined closures.
- 4.3 Pipets: Pasteur type or equivalent.

PNL TECHNICAL PROCEDURE

4.4 Kuderna-Danish (KD) Apparatus:

- 4.4.1 Concentrator tube: 10 or 25 mL with ground glass fitting and stopper.
- 4.4.2 Evaporation flask: KD type, 500 mL.
- 4.4.3 Snyder columns: Three ball macro and two ball micro or equivalent.
- 4.7 Oven: Capable of maintaining 150°C.
- 4.8 Water bath: Capable of temperature control and boiling water.
- 4.9 Boiling chips: Cleaned silicon carbide or equivalent.
- 4.10 Erlenmeyer flasks: 250 mL and as needed.
- 4.11 Beakers: 500 mL and as needed.
- 4.12 Chromatographic column: 250 mm long x 10 mm I.D. with pyrex glass wool plug and Teflon stopcock or equivalent, including commercially available pre-packed cartridges.
- 4.13 Muffle furnace: Capable of maintaining 400°C for several hours.
- 4.14 Separatory funnel: 125 mL with teflon stopcock.
- 4.15 Culture tube: Disposable 16 x 150 mm with screw top teflon lined closures.
- 4.16 Centrifuge: Bench top with openings capable of holding 16 x 150 mm culture tubes and achieving 2000 rpm.

5.0 REAGENTS

- 5.1 Sodium Sulfate: Granular, anhydrous purified by heating at 400°C for at least 4 hours.
- 5.2 Water: In which an interferant is not observed at or above the CRQL for each parameter of interest.
- 5.3 Sodium hydroxide solution: 10 N, dissolve 40 grams sodium hydroxide into reagent water and dilute to 100 mL.
- 5.4 Sulfuric acid solution: A.C.S grade or equivalent sulfuric acid and reagent grade water 1/1 (v/v).

PNL TECHNICAL PROCEDURE

- 5.5 Solvents: Cyclohexane, pentane, methylene chloride (pesticide quality or equivalent).
- 5.6 Silica gel: 100/200 mesh activated at 140°C for at least 16 hours in a beaker covered with foil. The silica gel should not exceed a 5 cm depth.

6.0 SAMPLE COLLECTION, PRESERVATION, AND HANDLING

- 6.1 Samples to be used for this procedure shall follow guidelines provided by the appropriate procedure and matrix, for example PNL-ALO-344.

7.0 PROCEDURE

- 7.1 Acid/Base-neutral partitioning: Proceed to 7.2 when cleaning water extracted residues.
 - 7.1.1 Quantitatively transfer the sample extract obtained from the appropriate extraction procedure such as PNL-ALO-344 or -120 into a clean 16 x 150 mm culture tube. Add 10 mL of reagent water.
 - 7.1.2 Check the pH of the sample with wide range pH paper and adjust to pH >11 with sodium hydroxide solution.
 - 7.1.3 Add 5 mL of methylene chloride to the culture tube. Extract the sample by shaking the culture tube for two minutes with periodic venting to release excess pressure.
 - 7.1.4 Allow the organic layer to separate from the water phase for a minimum of 10 minutes. If the emulsion interface between layers is more than one-third the volume of the solvent layer, then centrifuging should be employed to mechanically aid the phase separation. If centrifuging is unsuccessful, the contents of the culture tube may be transferred to a 125 mL separatory funnel followed by an addition of 50 mL reagent water. All subsequent extractions are then performed using 20 mL of methylene chloride.
 - 7.1.5 Collect the lower methylene chloride layer containing the base-neutral fraction using a pasteur pipet and transfer to a concentrator tube.

PNL TECHNICAL PROCEDURE

Repeat steps 7.1.3 through 7.1.5 two additional times. Save the base-neutral fraction for solvent exchange, Section 7.2.

- 7.1.6 Adjust the aqueous phase to pH <2 using the sulfuric acid solution. Perform three methylene chloride extractions as described in steps 7.1.3 through 7.1.5 collecting the acid fraction of the partition.
- 7.1.7 Add a boiling chip and concentrate the collected acid fractions using a micro KD apparatus. For analysis using PNL-ALO-345, when the apparent volume of the residue reaches 0.5 mL, about 5-20 minutes, remove the KD apparatus and allow it to drain and cool for at least 10 minutes.
- 7.1.8 Remove the micro Snyder column and rinse its lower joint into the concentrator tube with a minimum amount of methylene chloride. Further, centrifuging may be required to remove small particulates in the residue if the radioactivity level exceeds desired levels. Adjust the volume to 1.0 mL. Transfer the residue to a 1.5 mL amber screw top vial. Cap with a teflon lined closure. Proceed with the appropriate analysis on the acid residue.

7.2 Solvent exchange:

- 7.2.1 Transfer the sample extract obtained from the base-neutral partitioning procedure to a KD concentrator tube. Add a boiling chip.
- 7.2.2 Add 4 mL of cyclohexane and attach a micro Snyder column to the concentrator tube.
- 7.2.3 Place the assembled micro KD apparatus in boiling water so that the concentrator tube is partially immersed in the water.
- 7.2.4 When the apparent volume of the cyclohexane residue reaches 0.5 mL, about 5-20 minutes, remove the KD apparatus and allow it to drain and cool for at least 10 minutes.
- 7.2.5 Remove the micro Snyder column and rinse its lower joint into the concentrator tube with a minimum amount of cyclohexane. Adjust the volume to about 2 mL.

PNL TECHNICAL PROCEDURE

7.3 Column cleanup:

- 7.3.1 Prepare a slurry of 10 grams of activated silica gel in methylene chloride and place this into a 10 mm I.D. chromatographic column. Tap the column several times to settle the silica gel. Allow 3-4 cm of unpacked space to remain in the column.
- 7.3.2 Elute the methylene chloride to 1-2 cm above the silica gel. Add 1-2 cm of anhydrous sodium sulfate to the top of the silica gel.
- 7.3.3 If using commercial pre-packed columns, begin their use here. Pre-elute the column with 40 mL of pentane. The rate for all elutions should be about 2 mL/minute. Discard the pentane eluate.
- 7.3.4 Just prior to exposure of the sodium sulfate layer to the air, transfer the 2 mL of cyclohexane base-neutral sample residue onto the column. Use an additional 2 mL of cyclohexane to rinse and complete the transfer. Continue to discard the column eluent.
- 7.3.5 Just prior to exposure of the sodium sulfate layer to the air, add 25 mL of pentane and continue elution of the column. Discard the pentane eluate.
- 7.3.6 Just prior to exposure of the sodium sulfate layer to the air, add 25 mL of 2/3 (v/v) methylene chloride/pentane. Collect this eluent into a 250 mL erlenmeyer flask or directly into a 500 mL KD flask equipped with a concentrator tube.
- 7.3.7 Just prior to exposure of the sodium sulfate layer to the air, add 25 mL of methylene chloride to the column. Continue to collect the column eluent.
- 7.3.8 Just prior to exposure of the sodium sulfate layer to the air, add 25 mL 1/1 (v/v) of methylene chloride/acetone to the column.
- 7.3.9 Just prior to exposure of the sodium sulfate layer to the air, add 25 mL of acetone to the column. Collect the eluent until the column runs dry.
- 7.3.10 Concentrate the column eluent using the macro KD apparatus, followed by the micro KD apparatus if necessary, to whatever volume is required by the analysis procedure, for

PNL TECHNICAL PROCEDURE

PNL-ALO-345 discontinue concentration when the apparent volume is 0.5 mL.

- 7.3.11 Remove the Snyder column and rinse the lower joint and KD flask into the concentrator tube. Adjust the final volume to 1.0 mL with methylene chloride. Transfer the residue to a 1.5 mL amber screw top vial. Cap with a teflon lined closure. Proceed with the appropriate analysis on the residue.

8.0 QUALITY CONTROL

- 8.1 Before processing samples, the analyst shall demonstrate through the analysis of a method blank that all glassware and reagents are interference free when used in the quantity as specified in this method. With each batch of samples, and whenever there is a change in reagents or the location of performing this procedure, a method blank shall be processed as a safeguard against chronic laboratory contamination. The blank samples shall be carried out through the all stages of the sample preparation including matching handling and storage times and location.
- 8.2 Standard quality assurance practices consistent with the analysis method shall be employed while using this method. Fortified samples shall be carried through all stages of sample preparation adhering to those parameters previously specified with the blank preparation.
- 8.3 Limits of uncertainty for all measurements shall be in accordance with PNL QA Plan MCS-033 Rev. 0. Briefly, the limit is ± 5 in the next digit beyond the last one stated. For example, 5.0 mL means 5.0 ± 0.05 mL.
- 8.4 Control Limits:
 - 8.4.1 Surrogate standard determinations shall be performed on all samples and blanks. All samples and blanks are fortified with surrogate spiking compounds before extraction and this cleanup procedure in order to monitor the preparation and analysis of samples.
 - 8.4.2 Matrix spikes shall be performed at a frequency specified by the appropriate method such as PNL-ALO-344 or -120. Briefly, matrix spike and matrix spike duplicate analyses are required for each matrix and every twenty samples, the same as specified by the EPA-CLP SOW dated 2/88.

PNL TECHNICAL PROCEDURE

- 8.4.3 Control limits for this method were calculated in accordance with EPA and PNL QA Plan MCS-033 Rev 0 procedures. Table 1 provides surrogate recovery control limits and Table 2 provides spiking compound recovery limits obtained from analyses using PNL-ALO-345. Due to the limited quantity of data provided from validation study samples and sample matrices, all control limits for this method shall be considered advisory only. Currently, no action will be required if these control limits can not be met, however, these instances shall be noted in the case narrative. Additionally, these control limits were obtained from calculations which were estimated based on only two representative samples.

PNL TECHNICAL PROCEDURE

TABLE 1. SURROGATE RECOVERY CONTROL LIMITS

<u>Surrogate Compound</u>	<u>Level*</u>	<u>Recovery Limit(%)</u>	<u>RPD Limit</u>
Nitrobenzene-d ₅	100 ug	34-85	50
2-Fluorobiphenyl	100 ug	34-84	50
p-Terphenyl-d ₁₄	100 ug	51-94	50
Phenol-d ₅	100 ug	20-89	50
2-Fluorophenol	100 ug	12-81	50
2,4,6-Tribromophenol	100 ug	35-99	50

All limits are for advisory purposes only

* At time of injection in the validation study. Actual levels used in the procedure are typically 100-200 for acids and 100-50 ug for base-neutrals.

TABLE 2. MATRIX SPIKE RECOVERY LIMITS

<u>Matrix Spike Compound</u>	<u>Level*</u>	<u>Recovery Limit(%)</u>	<u>RPD Limit</u>
1,2,4-Trichlorobenzene	100 ug	25-81	50
Acenaphthene	100 ug	58-92	50
2,4-Dinitrotoluene	100 ug	56-91	50
Pyrene	100 ug	64-101	50
N-Nitroso-Di-n-Propylamine	100 ug	49-114	50
1,4-Dichlorobenzene	100 ug	15-76	50
Pentachlorophenol	100 ug	39-138	50
Phenol	100 ug	29-79	50
2-Chlorophenol	100 ug	31-70	50
4-Chloro-3-Methylphenol	100 ug	37-96	50
4-Nitrophenol	100 ug	30-123	50

All limits are for advisory purposes only

* At time of injection in the validation study. Actual levels used in the procedure are typically 100-200 for acids and 100-50 ug for base-neutrals.

PNL TECHNICAL PROCEDURE

TABLE 3. METHOD ACCURACY AND PRECISION

Compound Name	-----Percent Recovery-----	
	Mean	Standard Deviation (n-1)
1) *1,4-Dichlorobenzene-d4	100.0	0.0
2) #2-Fluorophenol	46.6	11.4
3) #Phenol-d5	54.5	11.6
4) Phenol	54.2	8.4
5) bis(2-Chloroethyl)ether	56.3	6.8
6) 2-Chlorophenol	50.5	6.6
7) 1,3-Dichlorobenzene	53.3	10.9
8) 1,4-Dichlorobenzene	45.6	10.2
9) Benzyl alcohol	71.0	7.4
10) 1,2-Dichlorobenzene	56.2	8.4
11) 2-Methylphenol	50.7	7.4
12) bis(2-Chloroisopropyl)ether	92.5	10.8
13) 4-Methylphenol	52.3	8.4
14) N-Nitroso-di-n-propylamine	81.8	10.8
15) Hexachloroethane	30.8	11.6
16) *Naphthalene-d8	100.0	0.0
17) #Nitrobenzene-d5	59.5	8.6
18) Nitrobenzene	66.2	5.4
19) Isophorone	55.8	23.7
20) 2-Nitrophenol	62.9	8.7
21) 2,4-Dimethylphenol	27.5	5.7
22) Benzoic acid	37.1	16.8
23) bis(2-Chloroethoxy)methane	76.6	5.9
24) 2,4-Dichlorophenol	64.5	8.4
25) 1,2,4-Trichlorobenzene	52.9	9.2
26) Naphthalene	68.6	5.9
27) 4-Chloroaniline	68.2	14.4
28) Hexachlorobutadiene	29.7	10.1
29) 4-Chloro-3-methylphenol	66.7	9.9
30) 2-Methylnaphthalene	65.9	4.6
31) *Acenaphthene-d10	100.0	0.0
32) Hexachlorocyclopentadiene	14.5	8.4
33) 2,4,6-Trichlorophenol	66.6	10.2
34) 2,4,5-Trichlorophenol	64.9	9.9
35) 2-Chloronaphthalene	56.7	5.3
36) #2-Fluorobiphenyl	59.3	8.4
37) 2-Nitroaniline	84.9	7.9
38) Dimethylphthalate	82.3	6.6
39) Acenaphthylene	62.2	5.5
40) 2,6-Dinitrotoluene	68.1	5.3
41) 3-Nitroaniline	73.9	6.2

PNL TECHNICAL PROCEDURE

TABLE 3. METHOD ACCURACY AND PRECISION (continued)

Compound Name	-----Percent Recovery-----	
	Mean	Standard Deviation (n-1)
42) Acenaphthene	75.2	5.8
43) 2,4-Dinitrophenol	64.3	8.7
44) 4-Nitrophenol	76.6	15.4
45) Dibenzofuran	68.7	6.1
46) 2,4-Dinitrotoluene	73.5	5.8
47) Diethylphthalate	61.7	28.3
48) 4-Chlorophenyl-phenylether	129.5	6.1
49) Fluorene	82.8	6.3
50) 4-Nitroaniline	38.1	15.2
51) *Phenanthrene-d10	100.0	0.0
52) 4,6-Dinitro-2-methylphenol	71.9	10.9
53) N-Nitrosodiphenylamine	74.8	5.9
54) #2,4,6-Tribromophenol	67.3	10.6
55) 4-Bromophenyl-phenylether	79.5	7.0
56) Hexachlorobenzene	50.8	13.5
57) Pentachlorophenol	88.7	16.4
58) Phenanthrene	81.2	5.8
59) Anthracene	65.3	6.4
60) Di-n-butylphthalate	71.7	27.8
61) Fluoranthene	78.8	6.5
62) *Chrysene-d12	100.0	0.0
63) Pyrene	82.4	6.3
64) #Terphenyl-d14	72.5	7.2
65) Butylbenzylphthalate	83.0	5.6
66) 3,3'-Dichlorobenzidine	37.0	8.3
67) Benzo(a)anthracene	82.0	7.0
68) Chrysene	83.5	6.5
69) Bis(2-Ethylhexyl)phthalate	66.9	29.3
70) *Perylene-d12	100.0	0.0
71) Di-n-octylphthalate	67.2	27.9
72) Benzo(b)fluoranthene	88.7	6.1
73) Benzo(k)fluoranthene	67.8	3.9
74) Benzo(a)pyrene	63.3	5.8
75) Indeno(1,2,3-cd)pyrene	75.5	6.5
76) Dibenz(a,h)anthracene	73.2	5.1
77) Benzo(g,h,i)perylene	74.0	6.6

* Compound is an Internal Standard

Compound is a Surrogate Spike

PNL TECHNICAL PROCEDURE

9.0 METHOD PERFORMANCE AND SPECIFIC QUALIFICATION

9.1 Table 3 provides method accuracy and precision based on analysis of:

- 22 acid surrogate spiked samples
- 12 base-neutral surrogate spiked samples
- 17 acid TCL spiked samples
- 9 base-neutral TCL spiked samples

These data were obtained from three different matrices. As seen from Tables 1, 2, and 3, the base-neutral ranges are quite wide. The low number of determinations contributes substantially to variability. It is anticipated that the range will narrow when more data is available.

9.2 The validation of this procedure was performed using spikes of the complete Semivolatiles Target Compound List (TCL) from the USEPA-CLP, SOW dated 2/88, on a limited number of authentic single shell tank wastes and blank samples. See reference 1.

REFERENCES

1. Hoppe, E. W.; Stromatt, R. W.; Campbell, J. A.; Steele, M. J.; Jones, T. E., Development and Validation of a Cleanup Method for Hydrocarbon Containing Samples for the Analysis of Semivolatile Organic Compounds, Pending Battelle-PNL Technical Report, 4/92.
2. PNL-ALO-120, Semivolatile Organic Extractions of Single Shell Tank Samples, Rev. 0, Battelle-PNL Technical Procedure, 8-3-89.
3. PNL-ALO-344, Procedure for Sample Preparation of Samples for GC/MS Analysis for Semivolatile Organic Compounds, Rev. 0, Battelle-PNL Technical Procedure, 4-26-91.
4. PNL-ALO-345, GC/MS Analysis of Extractable Semivolatile Organic Compounds, Rev. 0, Battelle-PNL Technical Procedure, 8-3-89.
5. QA Plan for Activities Conducted by the Analytical Chemistry Laboratory (ACL), QA Plan MCS-033, Rev. 0, 9-4-91
6. USEPA Contract Laboratory Program, Statement of Work for Organic Analysis, Multi-Media, Multi-Concentration, SOW No. 2-88.

PNL TECHNICAL PROCEDURE

7. USEPA Contract Laboratory Program, Statement of Work for Organic Analysis, Multi-Media, Multi-Concentration, Document Number OLM01.0, March 1990.
8. USEPA Office of Solid Waste, Test Methods for Evaluating Solid Waste, SW-846, Third Edition, 11-86.

DISTRIBUTION

No. of
Copies

No. of
Copies

OFFSITE

12 DOE Office of Scientific and
Technical Information

ONSITE

2 DOE Richland Field Office

J. M. Clark, A4-02
J. R. Noble-Dial, A4-02

11 Westinghouse Hanford Company

D. A. Dodd, T6-50
S. G. McKinney, T6-50
S. G. Metcalf, T6-50
A. F. Noonan, R2-12
L. M. Sasaki (5), R2-12
K. L. Silvers, T6-08
W. I. Winters, T6-50

15 Pacific Northwest Laboratory

J. A. Campbell, P8-08
E. W. Hoppe (5), P7-22
T. E. Jones, P7-27
M. J. Steele, P7-53
R. W. Stromatt, P7-22
Publishing Coordination
Technical Report Files (5)

END

**DATE
FILMED**

6 / 30 / 92