PROGRESS REPORT

Grant No. DE-FG05-87ER13751
(July 1, 1991 through June 30, 1992)

DOE/ER/13751--5

"Theoretical Atomic Collision Physics"

DE92 013571

March 9, 1992

N.F. Lane
Rice University
Houston, Texas 77251

The theoretical atomic collision physics program at Rice includes studies of a variety of low-to-intermediate energy atomic collision processes of possible importance to advanced energy research. Semi-classical, classical and quantum mechanical methods are used to investigate: excitation and electron-capture in ion-atom/molecule collisions; state-changing and ionization in collisions of excited atoms with atoms and molecules; and Rydberg atom collisions. Our research program continues to benefit significantly from collaborations with colleagues at Rice and other institutions, including many researchers supported by the DOE atomic physics program.

The current focus of the research is low-energy (collision $v \ll$ bound v_e) inelastic collisions for selected systems that possess the characteristic that many intermediate states are strongly coupled, such as can occur in classes of excited-atom (including Rydberg atom) collisions with atoms, molecules and positive and negative ions: (1) We are interested in the dependence of various differential and total cross sections on the angular momentum of the initial excited state and on the alignment of the initial electron charge distribution (for non-spherical initially excited states). (2) We wish to understand how characteristics of the classical trajectories (in CTMC calculations), *e.g.* multiple encounters, quasi-periodicity, chaos, relate to characteristics of the probability (scattering) amplitudes obtained from semiclassical (quantum mechanical) treatments. (3) In particular, in order to investigate a range of "interaction regimes," we have proposed to study low-Rydberg-atom collisions with: ions and polar molecules (long range interaction); non-polar molecules and atoms (short-range interaction); as well as electron-attaching atoms/molecules (transient electron capture possible). (4) We plan to look for observable signatures of possibly novel intracollisional interference effects and quasi-vibrational resonance effects that may occur in low-Rydberg collisions.

In the first grant year, which ends on June 30, 1992, we proposed to complete a number of projects that were begun under the previous grant and initiate new studies addressing the research objectives outlined above. We have been able to make considerable progress on all the projects proposed for this first year; very brief descriptions follow. Research is continuing in all these areas.

Ionization of low-Rydberg atoms in collisions with polar molecules:

Classical (CTMC) calculations of ionization (and other state-changing) cross sections have been performed for $\text{Na}(nl)$ on HCl molecules to study the relationship between the collision velocity dependence of the cross sections and the onset of multiple electron-molecule encounters in the collision.

MASTER

MAR 12 1992

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

We find that at thermal velocities, ionization results from a transfer of rotational energy from the molecule and usually involves many electron-molecule encounters. The resulting velocity dependence, as expected, is not a simple $1/v$ form; and it appears to agree, at least qualitatively with the experimental observations of Pesnelle's group (Phys. Rev. A 38, 4552, 4560 (1988)).

Dependence of amplitudes and cross sections for low-Rydberg-atom collisions with He on alignment of initial (np) state : Semi-classical (impact-parameter) calculations have been performed to study the dependence of $n,l \rightarrow n',l'$ state-changing cross sections on the alignment (direction of principal axis of the p orbital, in this case) of the initial electron charge distribution. For Na(7p)/He collisions at low velocities, the dominant final channels are Na(6f, 6d, 6p and 7s). The alignment dependence is found to be significant and is not entirely washed out, therefore observable, in the integrated cross sections.

Intracollisional effects and quasi-vibrational resonances in low-Rydberg collisions with atoms: Only preliminary studies have so far been carried out to see if these effects will be experimentally observable, and at this point we are not ready to say. The resonances are due to quasi-vibrational (pre-dissociation) states in the shallow wells that occur in some of the oscillatory adiabatic potential curves of the Na(nl)/He system. Our tentative conclusion is that several resonance states exist with lifetimes that are likely to be sufficiently long to produce structure in the cross sections. However, we have not yet performed the quantum mechanical collision calculations. The intracollisional interference effects refer to the possibility of an observable situation in which an incident He atom passing through different lobes of the wavefunction of the aligned Rydberg atom "scatters" the electron twice (or more times), with amplitudes which interfere with one another. We have seen the effect in semi-classical calculations for a selected narrow range of impact parameters. The phenomena should be present in differential cross sections, which, unfortunately, would be very difficult to measure; is not clear if the features wash out in a sum over impact parameters.

Charge-transfer cross sections and alignment/orientation parameters for Na(3p)/He+ collisions--comparison with Na(3s)/He+ collisions: Semi-classical (MO multi-state) calculations have been performed for collision energies in the low keV/amu range. The cross sections for single charge-transfer are found to be comparable for Na(3s) and Na(3p) collisions up to 8-10 keV/amu. For Na(3p) collisions, the final-state orientations of He($2^{1,3}P$) are relatively independent of energy in the range 3-10 keV/amu but show large variations with respect to impact parameter.

Other projects: In addition to the projects described above, a few from the earlier grant are nearing completion and many are completed and papers either submitted or in various states of preparation. These include: charge-transfer in collisions between stripped ions and H and H₂, quenching in collisions between Rydberg Na and Li atoms and rare-gas atoms, and a few small projects related to experimental programs at Rice.

PAPERS PUBLISHED IN 1991, OR LATER, AND PAPERS IN PRESS

Many of the following papers report research begun under the previous grant, when Dr. Kimura was a particularly active collaborator. Some of the following papers are also co-authored by scientists who visited Rice for short periods during the summers and received very modest support from the grant. All papers acknowledge the DOE OBES Division of Chemical Sciences.

R.S. Gao, L.K. Johnson, G.J. Smith, C.L. Hakes, K.A. Smith, N.F. Lane, R.B. Stebbings and M. Kimura, "Collisions between H^+ and H_2 at kilo-electron-volt energies: absolute differential cross sections for small-angle direct, single-, and double-charge-transfer scattering," Phys. Rev. A 44, 5599 (1991)

A. Kumar, M. Kimura and N.F. Lane, "Selective-state charge transfer in a collision between Li^{3+} ion and a ground-state Li atom: A molecular-state approach," Phys. Rev. A 43, 217 (1991).

B.C. Saha, N.F. Lane and M. Kimura, "Molecular-state treatment of $He^+(2p)$ excitation through electron capture in He^{2+} - H_2 collisions at low energies," Phys. Rev. A 44, R1 (1991).

M. Kimura and N.F. Lane, "Theoretical study of charge transfer in $He^+ + H_2$ collisions in the meV regime," Phys. Rev. A 44, 259 (1991)

R.S. Gao, C.M. Dutta, N.F. Lane, K.A. Smith, R.F. Stebbings and M. Kimura, "Experimental and theoretical studies of the $He^{2+} - He$ system: Differential cross sections for direct, single-, and double-charge-transfer scattering at keV energies," Phys. Rev. (to be published 1992)).

N. Shimakura* and M. Kimura, "Electron capture in collisions of N^{5+} ions with H atoms from the meV to keV energy regions," Phys. Rev. A 44, 1659 (1991)

N. Shimakura*, M. Itoh and M. Kimura, "Molecular treatment of electron capture in collisions of N^{4+} ions with H atoms," Phys. Rev. 45, 267 (1992).

*(Dr. Shimakura was a summer visitor at Rice U. while a portion of this work was completed and received a small amount of support from the grant.)

C.M. Dutta, N.F. Lane and M. Kimura, "theoretical study of nonresonant $^3He^+ + ^4He \rightarrow ^3He + ^4He^+$ charge-transfer in the threshold region," (submitted for publication).

END

**DATE
FILMED**

6 / 29 / 92