WSRC-RP--91-1244

DE92 013118

**THE LIMNOLOGY OF L LAKE:
RESULTS OF THE L-LAKE MONITORING PROGRAM
1986-1989 (U)**

by

James A. Bowers

Westinghouse Savannah River Company
Savannah River Site
Aiken, South Carolina 29808

DISCLAIMER

This is a Technical Report

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This report was prepared in connection with work done under Contract No. DE-AC09-89SR18035 with the U.S. Department of Energy. By acceptance of this report the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this report, along with the right to reproduce and to authorize others to reproduce all or part of the copyrighted report.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

EB

D. B. Moore-Shedrow, Section Manager
Authorized Derivative Classifier

**THE LIMNOLOGY OF L LAKE:
RESULTS OF THE L-LAKE MONITORING PROGRAM 1986-1989 (U)**

James A. Bowers

Approved by D. B. Moore-Shedrow, Section Manager

Publication Date: December 15, 1991

Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808

THIS DOCUMENT WAS PREPARED IN CONNECTION WITH WORK UNDER THE U.S. DEPARTMENT OF ENERGY (CONTRACT DE-AC09-88SR18035). By acceptance of this paper, the publisher and /or recipient acknowledges the U. S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper, along with the right to reproduce and to authorize others to reproduce all or part of the copyrighted paper.

WSRC-RP-91-1244

**THE LIMNOLOGY OF L LAKE:
RESULTS OF THE L-LAKE MONITORING PROGRAM 1986-1989 (U)**

James A. Bowers

Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808

CONTENTS

EXECUTIVE SUMMARY	i
ABSTRACT	1
INTRODUCTION	1
DISCUSSION	4
Water Quality, Macronutrients, and Phytoplankton	4
Aquatic Macrophytes and Habitat Formers	9
Zooplankton	10
Benthic Macroinvertebrates	12
Fish	14
Community Succession	19
CONCLUSIONS	20
REFERENCES	21

LIST OF FIGURES

		<u>Page</u>
Figure 1.	Aerial view of L Reactor and L Lake.	1
Figure 2.	NPDES Permit Region of L Lake.	2
Figure 3.	Offshore Sampling Stations in L Lake.	3
Figure 4.	L-Lake Program History.	3
Figure 5.	Mean Annual Temperature ($^{\circ}\text{C}$) in Mixed Layer of L Lake, 1986-1989.	4
Figure 6.	Depth Integrated Primary Production in L Lake, 1986-1989.	5
Figure 7.	Dissolved Oxygen Concentrations in the Mixed Layer of L Lake, 1986-1989.	6
Figure 8.	pH Concentrations in the Mixed Layer of L Lake, 1986-1989.	6
Figure 9.	Mean Number of Algal Species in L Lake.	7
Figure 10.	Mean Integrated Primary Productivity in L Lake.	7
Figure 11.	Mean Integrated Chlorophyll <i>a</i> in the Mixed Layer of L Lake.	7
Figure 12.	Mean Concentrations of Total Phosphorus and Chlorophyll <i>a</i> in the Mixed Layer of L Lake.	8
Figure 13.	Percent Abundance of Blue-greens/Greens, and Diatoms in L Lake.	8
Figure 14.	Percent Cover of Aquatic Macrophytes in L Lake.	9
Figure 15.	Submerged Surface Area of L Lake Macrophytes.	10
Figure 16.	Mean Total Zooplankton Densities in L Lake.	11
Figure 17.	Mean Cladoceran and Copepod Densities in L Lake.	11
Figure 18.	Mean Species Richness for Total Zooplankton in L Lake.	12
Figure 19.	Mean Taxa Richness for Macroinvertebrates in L Lake.	13
Figure 20.	Macroinvertebrate Densities in L Lake.	13
Figure 21.	Macroinvertebrate Biomass (Ash Free Dry Mass/AFDM) in L Lake.	13
Figure 22.	<i>Chaoborus</i> Densities in L Lake.	13
Figure 23.	Species Richness of Fish in L Lake.	14
Figure 24.	Abundance of Threadfin Shad, Redbreast Sunfish, Bluegill Sunfish, and Largemouth Bass Estimated by Electrofishing in L Lake.	15
Figure 25.	Estimated Pelagic Fish Abundance in L Lake from SONAR.	16
Figure 26.	Bluegill Sunfish Condition Factor in L Lake.	16
Figure 27.	Redbreast Sunfish Condition Factor in L Lake.	16
Figure 28.	Largemouth Bass Condition Factor in L Lake.	17
Figure 29.	Relative Abundance of Prey Groups for Redbreast Sunfish in L Lake.	17
Figure 30.	Relative Abundance of Prey Groups for Bluegill Sunfish in L Lake.	17
Figure 31.	Relative Abundance of Prey Groups for Largemouth Bass in L Lake.	17
Figure 32.	Relative Abundance of Dominant Fish Compared to Regional Lakes.	18
Figure 33.	Fish Kill Events in L Lake.	19

EXECUTIVE SUMMARY

L Lake was constructed in 1985 on the upper regions of Steel Creek, SRS to mitigate the heated effluents from L Reactor. In addition to the NPDES permit specifications (Outfall L-007) for the L-Reactor outfall, DOE-SR executed an agreement with the South Carolina Department of Health and Environmental Control (SCDHEC), that thermal effluents from L-Reactor will not substantially alter ecosystem components in the approximate lower half of L Lake. This region should be inhabited by Balanced (Indigenous) Biological Communities (BBCs) in accordance with Section 316(a) of the Pollution Control (Clean Water) Act (Public Law 92-500). In response to this requirement the Environmental Sciences Section/Ecology Group initiated a comprehensive biomonitoring program which documented the development of BBCs in L Lake from January, 1986 through December, 1989. This report summarizes the principal results of the program with regards to BBC compliance issues and community succession in L Lake. The results are divided into six sections, water quality, macronutrients, and phytoplankton, aquatic macrophytes, zooplankton, benthic macroinvertebrates, fish, and community succession. One of the prime goals of the program was to detect potential reactor impacts on L-Lake. During 1987 and 1988, L-Reactor power levels were reduced approximately 50% compared to 1986 operating levels, lessening the amount of heat to be dissipated by the lake. Consequently, lake temperatures for the November through June reactor operating periods were generally lower during 1987 and 1988 than those observed during 1986. L-Reactor has not operated since June 1988; therefore data for the last half of 1988 and all of 1989 was collected during an ambient thermal regime for this latitude.

Water quality in L Lake was generally comparable to other regional basins. Compared to other reservoirs, dissolved oxygen (DO), total suspended solids (TSS), total organic carbon (TOC), alkalinity, nitrate-nitrogen, cations, sulfate, chloride, and hydrogen sulfide concentrations were similar in magnitude. L Lake had relatively high values for pH, phosphorus, hypolimnetic ammonia, iron, and manganese. L Reactor had only a slight impact on seasonal temperature cycling during fall and winter in 1986 and 1987 by delaying fall overturn and sustaining thermal stratification for a few weeks longer than would normally occur. Hypolimnetic discharge was performed during 1987 which resulted in increased bottom temperatures, higher oxidation-reduction potentials, increased oxygen concentrations, and lower dissolved materials. Cooling water from the Savannah River caused high rates of phosphorus loading into L Lake which in turn promoted algal production. There were no consistent gradients or trends along the long axis of the lake that potentially would impact the biota.

The phytoplankton community in L Lake was similar to other eutrophic basins in the region. *Microcystis aeruginosa*, a pollution but not thermally tolerant species, dominated the community during 1986 and 1987. *M. incerta*, another closely related blue-green species, and *Cryptomonas erosa* dominated the algal assemblage during 1988 and 1989. Phytoplankton population dynamics were not correlated to L-Reactor operational cycles with regard to biomass, chlorophyll *a*, or primary production. An excess of phosphorus from the Savannah River created conditions of nitrogen limitation for the algae which gave blue-green species a competitive advantage to the point where light probably limited algal growth in

L Lake. Species diversity compared well with other basins. Phytoplankton densities were generally higher than in most reservoirs with chlorophyll *a*, primary production rates, and ash free dry mass within ranges for other regional lakes.

Rotifera and protozoans dominated the zooplankton assemblage in L Lake. While taxa richness was similar to other regional basins, protozoan densities were generally greater than in other basins. Planktivorous size-selective predation by threadfin shad progressively removed larger and more vulnerable species of copepods and cladocerans during 1988 and 1989.

Relative to BBC criteria L-Lake's macrophyte community is still at an early stage of successional development. Most of the plant diversity in the littoral regions of the lake were due to the SREL planting program performed in 1987. Though percent cover and plant biomass increased steadily during the four years of monitoring, both macrophyte attributes were partially a function of water levels. Generally, submergent macrophyte development remains low, most likely due to light inhibition which resulted from the high densities of algal populations. During the next few years with the increasing water transparency submergent growth is expected to develop at a faster rate.

The benthic macroinvertebrates in L Lake doubled in density from 1986 to 1988 and stabilized during 1989. Chironomids, especially the Tribe Chironomini, constituted 40.3 to 58.3% of the benthic community which is greater than most benthic communities in the region. However, chironomid diversity continually increased during the four years. *Chaoborus punctipennis*, common to other reservoirs, dominated meroplankton collections. When classified according to foraging behaviors, the collector-gatherers dominated functional group densities. There were significant increases of the clam *Corbicula fluminea*

in 1988 and 1989. Though taxa richness increased and reached a maximum in 1988 of 110 species, richness is slightly lower than other southeastern reservoirs. There were no results indicating that thermally tolerant species dominated L Lake or that L-Reactor operations affected benthic structure and function. This rich productive bottom fauna provided benthic feeding fish species with a suitable and substantial food resource.

Development of fish populations are best described in the succession construct where relative abundance shifts from flowing water species found in rivers and streams to a lacustrine and currentless habitat where lake species dominate the community. Most historical records of fish succession in reservoirs suggest that populations may require 5 to 10 years before stable populations are realized. Typical of this region in the United States, largemouth bass, bluegill, sunfish species, and shad dominate. Juvenile and adult bluegill and redbreast sunfish populations peaked in 1988 and declined slightly in 1989 for the bluegill, while production for both species remained high. General physical condition of bluegill, redbreast sunfish, and largemouth bass in 1989 had slightly declined when compared to 1987 and 1988. Such trends are not unusual in reservoirs and possibly reflect natural successional changes. Threadfin shad, which increased continually during the first three years, declined slightly in 1989 and may have reached its carrying capacity in the lake. Voracious planktivores, threadfin shad exerted a profound influence on the size structure of zooplankton species in L Lake, a process common to virtually all lake and ocean communities. Red sore disease infected L-Lake populations during the springs of 1988 and 1989. Outbreaks of this disease were concurrent with other regional events. When integrating all of the parameters of fish dynamics in the lake, results indicate BBC criteria are met.

Direct L-Reactor impacts were most identifiable during restart events which resulted in fish kills in the outfall canal and uppermost regions of the lake. The greatest loss occurred during December, 1986 when approximately 3.5 million sunfish, < 80 mm in length, were killed. All subsequent fishkills during restart episodes were at least two orders of magnitude less than the December 1986 event. Due to the reproductive strategies and the age classes of fish killed, fishkill events in L Lake did not have any significant ecological impacts on the fish populations themselves or the L-Lake ecosystem.

Several features of community succession in L Lake can be interpreted within current models of lake ecosystem development. Mechanistic hypotheses are proposed to explain nitrogen and light limitation of the phytoplankton, zooplankton size distributions and relative abundance, macrophyte growth, planktivore predation effects, and species succession in fish populations.

THE LIMNOLOGY OF L LAKE: RESULTS OF THE L-LAKE MONITORING PROGRAM 1986-1989 (U)

By James A. Bowers

Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808

ABSTRACT

L Lake is a comparatively young reservoir experiencing a successional pattern typical of most regional reservoirs. L-Reactor impacts on the lake ecosystem and the predicted successional trends are confounded by their cooccurrence. Reactor operations have not created permanent physical or chemical gradients which would impact the structure, size, or distribution of the biota. Relative to BBC mandates L Lake compares favorably with other regional basins.

INTRODUCTION

The Department of Energy's (DOE) nuclear production reactors for defense materials are located at the Savannah River Site (SRS). L Reactor releases heated effluents into the cooling reservoir, L Lake, built in 1985 on the upper

regions of Steel Creek. After exiting L Lake, water flows down the impounded portion of Steel Creek and discharges into the Savannah River (Figure 1).

Figure 1. Aerial view of L Reactor and L Lake.

The NPDES permit specifications, (Outfall L-007) for the thermal discharge into L Lake, mandate for the approximate lower one-half of the basin (Figure 2) that (1) the integrated daily mean temperature $\geq 50\%$ of the

L-Lake basin not exceed 35°C , (2) at least 50% of the lake be maintained at 32.2°C or below based upon a consecutive 5-day integrated mean temperature at 0.6 m (2 ft.) depth, (3) water temperature leaving the L-Lake dam into Steel Creek should not exceed 32.2°C .

Figure 2. NPDES Permit Region of L Lake.

In addition to the NPDES Permit compliance the DOE achieved an accord with the South Carolina Department of Health and Environmental Control (SCDHEC), that the effluent from L-Reactor will not significantly impact ecosystem components in the approximate lower half of L Lake. This region must be populated by Biological Balanced (Indigenous) Communities (BBCs) in accordance with Section 316(a) of the Pollution Control (Clean Water) Act (Public Law 92-500). The establishment of BBCs in L Lake would in terms of this law, "... assure the protection and propagation of a balanced, indigenous population of shellfish, fish, and wildlife in and on the body of water into which the discharge is to be made...". Preparation of a L-Lake 316(a) Demonstration (Gladden et al. 1989) followed the United States

Figure 3. Offshore Sampling Stations in L Lake.

Environmental Protection Agency's (USEPA) guidelines (EPA 1977) that specify four generally accepted criteria for concurrence. Communities satisfying the law:

- 1) are not dominated by pollution (thermally) tolerant species,

- 2) have biotic diversity and productivity characteristic of that found in other lakes and streams of the region,
- 3) have populations that are reproducing and are not maintained by repeated stocking and reseeded, and
- 4) have all trophic groups that would be expected in an ecosystem of that type.

A comprehensive biomonitoring program, which documented the trophic development of BBCs in the lake, was initiated in January, 1986 and continued through December, 1989 in its original form. Figure 3 indicates sampling regions and the offshore sampling stations in L Lake, while Figure 4 indicates the monitoring program's history with respect to L-Reactor power levels. Reactor operating cycles will be further discussed in the sections below.

Figure 4. L-Lake Program History

The purpose of this report is to summarize the main findings of the original monitoring program with regards to BBC compliance issues and community succession within L Lake. The results are divided into six components, water quality, macronutrients, and primary producers (phytoplankton), habitat formers (macrophytes), zooplankton, macroinvertebrates (benthic community), fish, and community succession.

All recently created impoundments follow an often observed sequence of events termed reservoir aging which is equivalent to the more universally applied term, ecological succession (Kimmel and Groeger 1986). Reservoir aging is divided into two phases, an initial trophic disequilibrium phase, lasting up to 5 years and a stabilization and maturation phase which may require

another 5 to 10 years. During the first year or two years of the disequilibrium phase impoundments experience what is termed a "trophic upsurge" (Baranov 1961) where, nutrient loading from the inlet and basin into the water column, stimulates algal blooms throughout the basin. Obviously this upsurge in lake productivity cannot continue for a long period of time with finite supplies of nutrients, mainly nitrogen and phosphorus. As will be seen shortly, L Lake experienced this upsurge for approximately four to five months. Following decreasing internal loadings of nitrogen and phosphorus, a decline in labile organic detritus, a cessation of habitat expansions throughout the basin, and a reduction of favorable habitats for niches already "occupied", impoundments enter a period of "trophic depression" (Kimmel and Groeger 1986) where productivity significantly decreases for at least one to three years. Afterwards a stabilization and maturation condition exists with fluvial nutrient inputs being the most important factor in determining lake metabolism for reservoirs. Limnetic community structure, most importantly the fish, may require five to ten years to stabilize. However, reservoirs are rarely isolated completely, because cultural inputs from point source industrial effluents and nonpoint source agricultural runoffs often dominate. Reservoirs, like all lakes, reflect their drainage basins (Hutchinson 1957). It is within this construct of reservoir aging and potential L-Reactor impacts, that this program was performed. Data interpretations and synthesis will be performed mostly on an annual basis because of the long-term nature of the results which will eventually cover the years from 1986 through 1992.

DISCUSSION

Water Quality, Macronutrients, and Phytoplankton

Physical and chemical conditions in L Lake reflected the successional trends expected for a newly built reservoir at this latitude. Compared to other regional reservoirs, dissolved oxygen (DO), total suspended solids (TSS), total organic carbon (TOC), alkalinity, nitrate nitrogen, Si, cations, sulfate, chloride, hydrogen sulfide were similar. Phosphorus concentrations, pH, hypolimnetic ammonia, iron, and manganese were comparatively higher than other southeastern lakes (Applied Biology 1981; Carr and Assoc. 1982; Kimsey et al. 1982; Harmon 1983). There were no consistent gradients or trends for water quality variables along the long axis of the lake. There were however downlake gradients in the hypolimnion for phosphorus, iron, manganese, ammonia-nitrogen, and TSS due simply to basin depth (redox potentials). Hypolimnetic discharge during 1987 resulted in increased bottom temperatures, higher oxidation-reduction potentials, increased oxygen concentrations, and lower dissolved materials. Chemical clines over depth for DO, ammonia-nitrogen, and hydrogen sulfide were reduced during this period. The most important water quality variables, relative to L Lake, are the seasonal temperature cycle and the significant input of phosphorus from the Savannah River during periods of L-Reactor operation which accelerates production processes in the basin (EPA 1976, 1980).

Figure 5. Mean Annual Temperature (°C) in Mixed Layer of L Lake, 1986-1989.

In L Lake production rates typically reached maximum rates (during summer) from April through September. Midsummer declines indicate nutrient limitation, in this case a paucity of nitrogen due to excess phosphorus supplies. These declines are characteristic of temperate nutrient limiting conditions. Winter minimum rates reached lowest levels during December, January, and February. Because photosynthesis produces significant amounts of oxygen and consumes carbon dioxide, dissolved oxygen are proportional to primary production rates and pH (see Wetzel 1983). This effect is seen in Figures 7 and 8 which indicate the seasonal cycles of DO and pH during the four-year study period. Both DO and pH emulate the temperature cycle in L Lake.

On an annual frequency several aspects of algal succession and production become apparent for this newly created reservoir. Species richness (Fig. 9) increased at an approximate linear rate from 1986 through 1988 reaching a mean maximum of 50 species. This declined slightly during 1989, but is representative of other basins in the southeastern United States. After the predicted trophic upsurge during the first six months' of the lake's

Figure 7. Dissolved Oxygen Concentrations in the Mixed Layer of L Lake, 1986-1989.

Figure 8. pH Concentrations in the Mixed Layer of L Lake, 1986-1989.

existence, primary production has continually declined during the study period (Fig. 10). Concurrently, chlorophyll *a*, a generally used indicator of algal biomass, decreased approximately 30% (Fig. 11). This decrease was loosely correlated to integrated estimates of total phosphorus (Fig. 12), an often used predictor of eutrophic status (Wollenweider 1976). Productivity estimates are similar to other reservoirs. Examples are Clarks Hill, 900-11,400 mg C/m²/h (Applied Biology 1982), Hartwell Lake, 800-35,100 mg C/m²/h (Carr and Assoc. 1982), and Lake Seminole, 200-60,500 mg C/m²/h (Water and Air Res. 1982). Similarly, chlorophyll *a* concentrations in these basins encompass estimates from L Lake (see Applied Biology 1982; Carr and Assoc. 1982; Water and Air Res. 1982).

Figure 9. Mean Number ($\pm$ SD) of Algal Species in L Lake.

Figure 10. Mean Integrated Primary Productivity ($\pm$ SD) in L Lake.

Figure 11. Mean Integrated Chlorophyll *a* ($\pm$ SD) in the Mixed Layer of L Lake.

Figure 12. Mean Concentrations ($\pm$ SD) of Total Phosphorus and Chlorophyll *a* in the Mixed Layer of L Lake.

Figure 13. Percent Abundance of Blue-greens, Greens, and Diatoms in L Lake.

The most important taxonomic groups in L-Lake's phytoplankton community were the Cyanophyta (blue-greens), Chlorophyta (greens), and Bacillariophyta (diatoms). Due to the high rates of nitrogen and phosphorus inputs into L Lake from the Savannah River, blue-green species have dominated the L-Lake algal community, particularly the species *Microcystis aeruginosa* and *M. incerta*. Figure 13 indicates the four-year history of blue-green, green, and diatom densities in the pelagic region of L Lake. During 1986 and 1987 *Microcystis aeruginosa* dominated that algal assemblage. Blue-green densities declined during 1988, while in 1989, *M. incerta* became the dominant blue-green species which accounted for the resurgence of blue-greens in the total percentage. Blue-green dominance in L Lake is typical of all basins having an excess of phosphorous. Newman et al. (1987) performed nutrient-addition experiments in L Lake to test whether the phytoplankton populations were either phosphorus and or nitrogen limited. In every experiment only the nitrogen additions produced significant increases in growth. Although field or laboratory experiments have not been performed to prove light inhibition, the light environment in L Lake strongly suggests that during 1986 and 1987, when the 1% PAR level was approxi-

mately one meter, a significant fraction of the algal assemblage was shaded out by surface mats of *Microcystis*. This combination of nitrogen and light limitation of phytoplankton growth is common to eutrophic reservoirs (Humphries and Lyne 1988). Chlorophyte species increased during 1988 and 1989. Small coccoid nanoplankton species (< 5 mm) dominated most of this increase. Concurrent to the relative increase of Chlorophyte species were increases in the flagellated chrysophytes, chiefly *Cryptomonas erosa* and *Rhodomonas minuta* during 1988 and 1989. Diatom populations were sporadically abundant during the winters of 1986-1987, spring of 1988, and April and July of 1989. Several species were commonly collected including *Melosira granulata*, *Skeletonema potamos*, *Cyclotella pseudostelligera*, and *Nitzschia holsetica*.

The above trends in L Lake duplicate successional progression experienced by newly constructed basins. Because L-Reactor operations were suspended during 1988, events in 1989 cannot be specifically correlated to this cessation of operations. Successional events will always be confounded with reactor operations. Intense monitoring just prior and during future L-Reactor restarts will significantly aid in deciphering reactor

induced impacts.

Relative to BBC criteria phytoplankton population dynamics were not correlated to L-Reactor operational cycles with regard to biomass, chlorophyll *a*, or primary production. During 1986 and 1987 the pollution (not thermally) tolerant *Microcystis aeruginosa* dominated the lake's algal assemblage. Algal diversity compared well with other regional basins, while phytoplankton densities were higher than in most reservoirs. Primary production rates, chlorophyll *a* and algal dry mass were well within the ranges for regional lakes.

Aquatic Macrophytes and Habitat Formers

Provided with the appropriate physical habitat, macrophytes become important structural and functional components of the littoral zone in reservoirs (Shireman et al. 1982). They indirectly structure the whole lentic ecosystem by affecting basin nutrient dynamics (Carpenter 1983). Additionally, aquatic plants provide habitat for phytoplankton, zooplankton, epiphytes, macroinvertebrates, and fish (Wiley et al. 1984; see Duarte et al. 1986). Macrophytes are an important food source for grazers and detritivores; they trap and recycle nutrients, stabilize the substrate, and are sources of oxygen for the water column (Raschke and Rusanowski 1984) and sediments (Chen and Barko 1988). Littoral zone flora produce significant amounts of organic material, most of which are placed into the nearshore sediments (Wetzel 1983).

Several habitat features affect the establishment and growth of macrophytes in the littoral zone. These include: sediment composition (Keddy and Constabel 1985), solar radiation (King 1975), nutrients (Wetzel 1983), shoreline slope (Duarte et al. 1986), and water level fluctuations (Moore and Keddy 1988). Slope is highly correlated to littoral zone macrophyte development (Duarte and Kalff 1986). Normally, a lake with a gentle slope will have a wider littoral zone and is more productive than a lake with a steep slope (Reid and Wood 1976). Gradually sloping shores tend to support greater vascular plant cover than do stepped shorelines (Sculthorpe 1967). Extended periods of elevated lake levels gradually eliminate emergent plant taxa (van der Valk and Davis 1978), whereas persistent low lake levels eliminate taxa intolerant of dry conditions. Growth is greater on nutrient-rich organic sediments than on sandy nutrient-poor substrates (Wetzel 1983). Solar radiation is a limiting factor in growth controlling photosynthetic rates (Wetzel 1983). Often in eutrophic basins, similar to L Lake, high light extinction coefficients limit macrophyte development.

The University of Georgia's Savannah River Ecology Laboratory (SREL) initiated an extensive transplanting program to accelerate colonization of the L-Lake littoral zone (Wein et al. 1987). Approximately 52 species of aquatic and wetland plants from Par Pond, SRS were planted throughout the L-Lake shoreline in 1987. During 1988 per cent cover (Fig. 14) ranged from < 5% in the winter to approximately 25% during the summer and changed little in 1989. No submerged plants were found in the sampling belt plots which were not located in the SREL-planted areas. Submerged surface area reached to approximately 300 m²/hectare in 1988, and decreased to 150 m²/hectare in 1989 (Fig. 15). This decrease reflects changes in water levels rather than true changes in plant growth and development. Generally submergent macrophyte development remains low due to a combination of slope, fetch length, and light inhibition. The emergents, *Typha*, *Eleocharis*, *Pontedaria*, and *Panicum* were often mapped in the eulittoral zone. Based on dominant species the results suggest that the current macrophyte diversity in L Lake is mainly due to the SREL planting program.

Figure 14. Percent Cover of Aquatic Macrophytes in L Lake.

Figure 15. Submerged Surface Area of L-Lake Macrophytes.

Abiotic or nonphotosynthetic components of the littoral zone increased through the four years. Standing timber left in shallow regions of the constructed basin and artificial structures placed there for increased habitat provided macroinvertebrates and fish an initial resource for habitat and cover. Because there is no import of such material from upper Steel Creek, these structures will continually decrease. Macrophyte growth will ultimately provide most of the lake's habitat structure.

Relative to BBC criteria L-Lake's macrophyte community is still at an early stage of successional development. The SREL planting program has been very successful with significant growth in the planted areas (Wein et al. 1987). Continued spread of these areas will contribute to a littoral community similar to older reservoirs. The rate of spread will partially depend upon continuing trends in algal production which limits water clarity and thus available light for macrophyte growth. Because light extinction coefficients have decreased in L Lake this growth limitation would appear to be decreasing with time. Physical inhibition due to those shorelines exposed to a long fetch will continue. None of the dominant species are pollution or thermally tolerant.

Zooplankton

Zooplankton populations, comprised of small protozoans, rotifers, and microcrustaceans couple a lake's primary production to the basin's fisheries (Hutchinson 1957; Gliwicz, 1969). Secondary consumer zooplankters "graze" upon phytoplankton in very selective modes based on size, shape, taste, and abundance (Haney 1973). This selective predation partially controls algal abundance and productivity. In turn zooplankton themselves are the principle prey of planktivorous fish. This predation is also size selective and has profound effects on the whole reservoir ecosystem through an process termed "cascading effects" (Carpenter et al. 1985). This process has been observed in L Lake through its first four years of existence with regards to zooplankton and fish.

Mean total zooplankton densities ranged from approximately 10,000 individuals/L to approximately 150,000 individuals/L, where in excess of 75% were ciliate protozoans (Fig. 16). Peaks in abundance were observed during all seasons which is characteristic of temperate lakes (Hutchinson 1967; Wetzel 1983). The large abundance peak during 1988 was comprised of an unidentified ciliate and the genus *Uronema* protozoans. During 1989 loricate rotifers dominated most of the large summer pulse. Cladoceran abundances (Fig. 17) declined steadily throughout the four-year period in response to intense predation from planktivorous fish, principally threadfin shad which continually increased in abundance during the study period. The mechanism for this decline is the mode of planktivore feeding which is visual and thus selects for large, visually vulnerable targets such as cladocerans, primarily the genus *Daphnia* (Brooks and Dodson 1965). Additionally, *Daphnia* are known to avoid or receive little food value from blue-green phytoplankters which were always present in L-Lake's algal assemblage (Porter and Orcutt 1980). Thus the history of zooplankton population dynamics in L Lake are partially described by planktivore abundance, typical of all freshwater and marine pelagic communities (Carpenter and Kitchell 1984).

With respect to BBC criteria zooplankton populations are very similar in structure to reservoirs in the southeastern United States. Taxa richness declined slightly during the four years (Fig. 18), but still is comparable to other reservoirs in the region (Bowers, 1990). Only the dominance of protozoans and rotifers in L Lake separates the basin from others which possibly is due to a large resource of bacteria and small picoplankton that are their principle foods (Porter et al. 1979).

Figure 16. Mean Total Zooplankton Densities in L Lake.

Figure 17. Mean Cladoceran and Copepod Densities in L Lake.

There were no periods during the four years when pollution (thermally) tolerant species dominated the zooplankton assemblage, abundance reflect food resources and fish predation. All trophic levels expected in a temperate reservoir were present. Macrozooplankton were represented by known grazers such as *Daphnia* and *Ceriodaphnia*, while predators

such as *Cyclops* were always in abundance. Furthermore, all populations exhibited successful reproduction and were not supported by continuous importation from Savannah River. Evidence for this lies in considering their life spans, which ranges from weeks for microzooplankton to months for the large crustacean macrozooplankton. Additionally, immature life stages

of crustacean fauna present, nauplii and copepodids, and egg-bearing rotifers. Population structure and foodweb functionality comply with BBC criteria.

Figure 18. Mean Species Richness for Total Zooplankton in L Lake.

Benthic Macroinvertebrates

Benthic macroinvertebrates are commonly partitioned into several arbitrarily defined components based primarily on taxa and size. They include nematodes, flatworms, bryozoans, annelid oligochaetes, leaches, ostracods, crustaceans, mollusks, and particularly the insects. This diverse fauna plays many roles in the ecosystem through their foraging behaviors (Cummins 1973; Wetzel 1983). Macroinvertebrates are also good indicators of ecological stress because of their sedentary habits and relatively long life spans (Wetzel 1983). Unfortunately, these animals have been difficult to study because of their complicated taxonomy and problems with quantitative sampling (Cummins 1973).

When stream beds are dammed to create impoundments the shift from the lotic to lentic habitat causes equally significant changes in the biota (Voshell and Simmons 1984; McLachlan 1977; Petr 1972; McDonald and Bernhard 1978). Lentic species, not requiring flowing water habitats replace the endemic-obligate rheophite species (Paterson and Fernando 1970; Nursall 1952; Voshell and Simmons 1984; McDonald and Bernhard 1978). Those species having aerial adults or high dispersal rates are the first groups to colonize a newly created reservoir, especially those carried into the basin

from upstream habitats (Paterson and Fernando 1969a, 1969b, 1970; Paterson and Fernando 1970; Petr 1972; Voshell and Simmons 1984).

Subsequently, benthic community succession reflects lotic to lentic changes from terrestrial-type sediments (leaf litter, dead terrestrial plants, other decaying organic matter) to silt and mud substrates. In turn this change alters the energy source for the benthic community from outside inputs of detritus to organic deposition from the primary production of phytoplankton and macrophytes (Voshell and Simmons 1984; McLachlan 1977; Paterson and Fernando 1970; Nursall 1952). Macroinvertebrate species composition and density are associated with this shift. Chironomids often dominate the macroinvertebrate community, if a large amount of terrestrial organic material was left in place when the reservoir was created (Krzyzanek et al. 1986; Baxter 1977; Paterson and Fernando 1969a). Excepting restricted areas left unaltered for fish habitat, most of the terrestrial organic material was removed prior to the filling of L Lake.

Potential thermal impacts from the L-Reactor discharge on the L-Lake macroinvertebrate community include the elimination of thermally-sensitive species, the subsequent domination by thermally tolerant species (Parkin and Stahl 1981), and accelerated insect production (Sweeney and Vannote 1978). However, most macroinvertebrate species in L Lake are endemic to this latitude and genetically warm-adapted.

Macroinvertebrate succession exhibited patterns consistent with many newly built reservoirs with regards to species richness, densities, and biomass. Cumulative mean taxa richness, based on quarterly PONAR and dip net sampling, remained relatively constant during the study with an annual mean low of approximately 24 species to a maximum of 33 species (Fig. 19). Including all types of sampling as many as 110 species were observed during 1988. These numbers are slightly less than values reported for many other regional reservoirs (Duke Power, Inc. 1985). Early dominance by collector-gatherer guilds was succeeded by collector-filterer species indicating a more stable community is developing.

Densities of benthic macroinvertebrates increased substantially during the four-year period where densities ranged from 3,955/m² to 11,762/m² (Fig. 20). The chironomid Tribe Chironomini dominated all taxa comprising no less than 40% at any time during the study with the genus *Glyptotendipes* being the most commonly sampled group. Correspondingly, biomass (Ash Free Dry Mass, AFDM) also significantly increased from 1986 to 1989 (Fig. 21).

Figure 19. Mean Taxa Richness for Macroinvertebrates in L Lake.

Figure 20. Macroinvertebrate Densities in L Lake.

Collector-gatherers comprised most of this biomass during 1986 through 1988, and collector-filterers dominated in 1989. Much of these increases resulted from large increases of the Asiatic clam *Corbicula fluminea* in 1988 and 1989. Densities and biomass were

inversely correlated to depth reflecting anoxic hypolimnia during prolonged periods of thermal stratification.

The predatory dipteran, *Chaoborus punctipennis*, dominated the meroplankton portion of the macroinvertebrate community reaching a maximum density of approximately 155,000/m³ in 1987 (Fig. 22). A voracious predator of zooplankton crustacea, *Chaoborus* probably accounts for a significant portion of zooplankton mortality in L Lake (Fedorenko 1973, 1975).

Figure 21. Macroinvertebrate Biomass (Ash Free Dry Mass/AFDM) in L Lake.

Figure 22. *Chaoborus* Densities in L Lake.

Relative to BBC considerations the L-Lake macroinvertebrate community is still experiencing succession characteristic to recently created basins. Species richness is at a level lower than most regional basins. These levels will increase as littoral macrophyte beds grow and provide additional and more diverse habitats (Voshell and Simmons 1984). Thermally-tolerant species did not dominate any region in the lake. Throughout the study period densities and biomass in sampling region 4 were always comparable to regions further downstream. Small thermal gradients during 1986 and 1987 were not correlated to absolute or relative abundances. No rare or endangered species were collected (SCWMRD 1976). Emergence trap results indicate that repeating generations of aquatic insects are reproducing and sustaining growth. All functional feeding groups (Merritt and Cummins 1984) were present during the study providing the necessary trophic couplings needed for interactions with the rest of the lake's ecosystem, particularly as a resource for benthic feeding bluegills and sunfish.

Fish

Fish populations are an important component of L-Lake because of their known impact on the ecosystem through size-selective predation (Hickman and Congdon 1974; Carpenter 1983). Assessment requires consideration of several interrelated factors including thermal effects, nutrient loading, and successional changes. The L-Lake fish community is unique in that no fishing pressure is imposed, either commercial or sport. Thermal effects can be lethal, destroy acceptable habitat, influence processes such as growth, activity, and oxygen consumption, and change fish movement and orientation in response to temperature gradients and relatively rapid changes in temperature (Brown 1974). Additionally, successional changes complicate the assessment of thermal effects by the confounding of natural changes in species composition, richness, and abundance with thermally-induced disturbances. The fish communities of new reservoirs undergo successional changes as lotic species are replaced by species better adapted to a lentic environment (Turner 1971; Kimmel and Groeger 1986). Fishing can strongly influence the size, age structure, and abundance of predator fishes, with subsequent effects on forage species (Hickman and Congdon 1974). The thermal influences of reactor operation and associated nutrient loading may also influence succession by altering the length of the growing season and the extent of algal productivity (Carlander 1977; Taylor 1971). The assessment presented here addresses species richness, absolute and relative abundances, condition indices, foraging patterns, regional comparisons, and fish kills.

Taxa richness in L Lake decreased from 1986 to 1987 and was basically unchanged from 1987 through 1988 (Fig. 23; Heuer et al. 1989). This trend continued in 1989 with only a slight increase. The same dominant taxa were found during 1989, despite a general decline in abundance from 1988 levels. Rare species were occasionally collected and this may explain minor changes in taxa richness from year to year. The elimination of thermally sensitive species from L Lake was observed. Species richness declined precipitously when temperatures exceeded 35 °C in Four Mile Creek (Paller and Saul 1986). It is unlikely that the illustrated initial decline in species richness in L Lake, from 1986 through 1987, was caused by thermal stress because the lake region never experienced water temperatures greater than 35 °C (Heuer et al. 1989). Limited habitat diversity is the more probable cause for the decline in taxa richness especially with the lack of a well developed littoral zone.

Figure 23. Species Richness of Fish in L Lake.

Figure 24 indicates the average monthly catch per unit effort (CPUE) for the four dominant species in L Lake. The principle successional trends, observed from electrofishing, included the steady decline in bluegill, the relatively constant abundance of the redbreast sunfish and largemouth bass, and the significant increase in threadfin shad during 1987 and 1988. These changes were most significant during 1986 and 1987 reflecting the expected successional changes in the whole basin.

Mechanisms for these changes are numerous and complex. They ultimately include all abiotic and biotic factors affecting age-specific birth and death rates which are very difficult to estimate and beyond the scope of this effort. Most importantly, the observed changes are similar to other successional patterns for reservoirs at this latitude (Carlander 1977). Increases in threadfin shad abundances were also indirectly observed by lakewide SONAR transects which estimated nighttime abundances of fish in offshore watercolumn (Fig. 25). Note the steady increase in target density magnitudes through 1988 and the slight declines during 1989. This trend corresponds directly to the concurrent declines in the larger species of calanoid copepods and cladoceran macrozooplankton. Parallel with these trends in the dominant species were the reduction in lotic species such as dollar sunfish, spotted sunfish, spotted sucker, redfin pickerel, and shiners which were endemic to the upper Steel Creek system prior to basin formation (Paller et al. 1988). These patterns have often been observed in other impoundments (Beckman and Elrod 1971; Turner 1971; Walburg et al. 1971). The proposed mechanism for this shift is the change from a diverse habitat like that in the former Steel Creek (Firth et al. 1986) to the more uniform lacustrine habitat in L Lake.

lower for the juvenile age groups, and that the highest condition factors occurred during 1986. The condition index decreased through 1988 for both adult and young bluegills and sunfish, and remained approximately the same for 1989. Condition indices for three size classes of largemouth bass exhibited roughly the same pattern. Although thermal impacts from reactor operations were considered as possibly being responsible for these declines from 1986 through 1987, index declines continued through 1988 when maximum littoral temperatures in 1988 were lower than in previous years. More likely, increased competition, especially among juveniles for declining food resources, and differential predation by largemouth bass on redbreast and bluegill are responsible for these trends. All three species are planktivorous with juveniles preferring crustacean macrozooplankton, copepods and cladocerans which declined significantly during 1988 and 1989. Threadfin shad, which increased from 1987 to 1988, would also compete for these prey which markedly declined in abundance due to this size-selective predation. Therefore all four dominant species, adult threadfin shad and juvenile sunfish and bass, were competing for the same decreasing macrozooplankters. Redbreast, bluegill, and bass adults also compete to varying degrees (Figs. 29, 30, and 31). All three species prey on Diptera,

Figure 24. Abundance of Threadfin Shad, Redbreast Sunfish, Bluegill Sunfish, and Largemouth Bass Estimated by Electrofishing in L Lake.

Average condition factors, using Lagler's (1956) equation of weight and length, were calculated for bluegill, redbreast sunfish, and largemouth bass (Figs. 26, 27, and 28). Though not indicated here, average factors were similar among different regions of the lake for all size classes. Note that condition values were

especially the sunfish. Redbreast and bluegill overlap extensively on Diptera, phytoplankton, and macrozooplankton which could account for the decreasing condition factors. Added to this competition is the largemouth bass predation on both species. Only a quantitative study, focusing on these trophic web

relationships, would conclusively demonstrate the hypothesized scenario above.

Figure 25. Estimated Pelagic Fish Abundance in L Lake from SONAR.

Figure 26. Bluegill Sunfish Condition Factor in L Lake.

Figure 27. Redbreast Sunfish Condition Factor in L Lake.

Figure 28. Largemouth Bass Condition Factor in L Lake.

Figure 30. Relative Abundance of Prey Groups for Bluegill Sunfish in L Lake.

Figure 29. Relative Abundance of Prey Groups for Redbreast Sunfish in L Lake.

Figure 31. Relative Abundance of Prey Groups for Largemouth Bass in L Lake.

When compared to other fish communities in southeastern reservoirs, the fish populations in L Lake are generally comparable to reservoirs of this size. Shad, bluegill, redbreast sunfish, and largemouth bass dominate the fish assemblage (Fig. 32). The successional decline in the proportion of bluegill, redbreast sunfish, and largemouth bass, coupled with the increase in the proportion of shad has evolved into a community more similar to other regional basins. The paucity of minnows, suckers, catfishes, and silversides appears to be the most significant difference between L Lake and other reservoirs which accounted for the other species not shown in Figure 32. However, the difference is not atypical for the southeastern United States. Dominance by either clupeids or centrarchids is common among reservoirs of the region as evidenced by Lakes Greenwood, Russell, and Robinson. Given a sufficient length of time, the fish assemblage of L Lake can probably be expected to resemble those of Pond B and Par Pond.

The L-Lake fish community satisfy BBC criteria for a reservoir of this age. Pollution (thermally) tolerant species do not dominate the lake. Bluegill, redbreast sunfish, largemouth bass, and threadfin shad are common to the climate of the southern United States (Clugston 1973; Beitingner 1974). The continued relative abundance pattern in L-Lake would not be unusual or indicative of thermal influences relating to operation of L Reactor. Some impoundments become overpopulated by bluegill (Carlander 1977), and the increased abundance of larval bluegills in 1988 and 1989 as well as the increase in the ratio of bluegill to redbreast sunfish, suggests this may be occurring in L Lake. From 1987 through the summer 1989, the ratio of redbreast sunfish to bluegill has remained near 0.5, but late in 1989 it increased to 1.5. Thus, trends observed during 1989 indicate that successional changes in L Lake are typical of the development of a warm-water reservoir fish assemblage.

Successful recruitment of each species requires collection of juveniles or adults as well as eggs and larvae (Nikolsky 1963; Summerfelt 1975; Paller et al. 1988). From 1987 through 1989, similar patterns of occurrence of adults and larvae were noted for the dominant taxa, reflecting continued successful reproduction of these taxa.

All representative trophic components in L-Lake food webs were present by 1987. Threadfin and gizzard shad function as primary consumers (Mosher 1984). Based on gut contents studies in L Lake threadfin shad have exhibited two feeding patterns. Some feed on benthos or detritus, others filter-feed on plankton indicating that energy pathways in the limnetic zone are important. The continued use of threadfin shad as prey by largemouth bass completes a grazing-based food web from

phytoplankton to secondary consumer (Paller et al. 1988). There is an extensive literature describing the planktivorous habits of small bluegills and redbreast sunfish (see O'Brien 1987; Threlkeld 1987). A detritus-based food web is initiated by the intense bluegill and redbreast sunfish predation on chironomid larvae and pupae. These bluegills and sunfish themselves are in turn prey for largemouth bass which terminates the food web. Bass also link themselves directly to detritus by preying on threadfin shad that occasionally feed directly on detritus. Detritus and grazing based food chains are typical of lakes and reservoirs (Odum 1971; Wetzel 1983). The presence and diversity of these two energy pathways, as well as the presence of all trophic groups, supports the designation of L Lake as a balanced biological community.

Figure 32. Relative Abundance of Dominant Fish in L Lake Compared to Regional Lakes.

Comprehensive assessments of fish communities must encompass any events in the basin which result in lethal impacts. In L Lake fishkill episodes have resulted from the naturally occurring redsore disease and L-Reactor restarts. The aquatic bacterium, *Aeromonas hydrophila* causes redsore disease that infects many freshwater fish, amphibians, reptiles, and man. The bacterium is common to all lakes and rivers throughout North America (Hazen et al. 1978a), and occasionally causes significant fish kills in North and South Carolina (Hazen et al. 1978a; Hazen et al. 1978b; Miller and Chapman 1976). Studies in Par Pond (Hazen 1979) and Albemarle Sound, NC (Hazen and Esch 1983) have indicated that the abundance and thus the rate of infection of fish, are positively correlated to total Kjeldahl nitrogen, orthophosphate, and total phosphorus, and chlorophyll *a*. During 1988 red sore disease occurred in L Lake from approximately the end of March through the end of May. A total of 9495 dead fish were counted along the L-Lake shoreline. Most of the fish were sunfish, but significant numbers of largemouth bass also died during the outbreak. The red-sore disease during 1989 began in mid-April and continued through the third week of June. Taxa experiencing the highest mortality were redbreast sunfish (1201), black crappie (967), bluegill sunfish (661), *Lepomis* spp. (380), and largemouth bass (348). The total count for the 1989 epizootic census was 4018 fish. Neither disease outbreaks during 1988 and 1989 resulted in mortality levels that would impact community level structure and function.

Direct reactor impacts have been experienced in L Lake as thermal kills during restart episodes. During a restart event flows and water temperature increase over a period of hours. Fish in the upstream region of the lake and especially in the outfall canal are exposed to lethal temperatures ($> 35^{\circ}\text{C}$) without the time or routes of escape for fish. The worst event occurred in December, 1986 when 3.5 million small ($< 80\text{ mm}$) sunfish were killed (Fig. 33). Two events, orders of magnitude less in scale, occurred during the January and March 1987. Again these kills involved small bluegills and redbreast sunfish. During November of the same year another episode occurred killing 27,000 fish. The last event occurred in April 1988 when only 12,000 fish were killed. Because L Reactor was shutdown in the fall of 1988, there were no fishkill events during 1989. The most important consideration relative to BBC criteria is the relative impact of these kills on the fish populations and the L-Lake ecosystem. Studies on the population dynamics of all types of freshwater fishes (Lagler 1956; Nikolsky 1963; Clugston, J.P. 1973; Mayhew 1974) have indicated that juvenile mortality is very high through the first year of life. Because of these high rates fish reproductive strategy is to produce large numbers of fry to guarantee enough adults to sustain the species.

The fishkills in L Lake primarily affect that age class that experiences high rates of mortality. Because the magnitudes of losses due to fishkill events are small relative to natural mortality rates for this age class, the impact has been negligible and virtually undetectable in the adult age group.

Figure 33. Fish Kill Events in L Lake.

Community Succession

After summarizing the four-year history of ecological events in L Lake it is now appropriate for a further synthesis addressing ecosystem level limnology. All natural lakes and reservoirs, experience the process called eutrophication. Eutrophication is defined here as a natural or culturally accelerated process leading from oligotrophy (nutrient poor) through mesotrophy (moderate nutrient conditions) to eutrophy (heavy nutrient enrichment), and ending in a dystrophic lake, a basin having large amounts of humic organic matter (see NAS 1969; Wetzel 1983). The process is modified for reservoirs as detailed earlier in the introduction. Superimposed on this natural and culturally accelerated process are current proposals regarding how lake ecosystems function as a whole. They are termed the "bottom up" model (Odum 1957), the "top down" model (Hrbacek 1962), and the "cascading effects" model (Carpenter and Kitchell 1984). The "bottom up" model states that primary productivity, regulated by nutrient loading, sets limits on lake metabolism, while the hypothesis in the "top down" construct states that size-selective fish predation controls the relative abundance and size distributions of their zooplankton

prey assemblages. The "cascading effects" model can be thought of as a corollary to the "top down" model. When top consumers dominate a community, it benefits those prey two trophic links lower in the food chain, because it reduces the abundance of the intermediate consumer (Abrams 1984). Reviews of these concepts are found in Bowers (1988).

Interpreting the trophic history of L Lake within these constructs indicate how L Lake has evolved when compared to other freshwater lacustrine systems. All lakes are reflections of their drainage basins with L-Lake's drainage basin being extended, in that the drainage basin includes its own basin plus those regions draining into the Savannah River (Hutchinson 1957). Boundary conditions for the lake's primary production were set by high rates of phosphorus and nitrogen inputs from the Savannah River which concurred with "bottom up" predictions. Nitrogen and light inhibition of the algal assemblage occurred during 1986 and 1987, due to an excess of phosphorus input from the Savannah River (Newman et al. 1987). The predicted "trophic upsurge" event occurred immediately during the spring of 1986 within two months after filling. Internal release of phosphorus from the newly inundated sediments assisted this short-term explosion of production. During this period *Microcystis* and other blue-green species had a competitive advantage over other phytoplankton groups. Herbivorous zooplankters, including the larger Copepoda and Cladocera, were provided an abundance of food without suffering substantial mortality to planktivorous fish when L-Lake fish populations were shifting from a riverine assemblage to a lacustrine group. Although macrophytes had been planted on a limited basis, inhibition of light, current disturbances from long wind fetches, and limited favorable shoreline slopes, retarded littoral development. This limited littoral habitat could have restricted sunfish and largemouth bass populations. Benthic macroinvertebrates experienced dominance by collector-gatherer guilds, especially the chironomid Tribe Chironomini. Accelerated algal production provided the energy for this benthic development through high sedimentation rates of organic resources.

"Top down" and "cascading effects" model impacts were observed in the shift from a riverine fish community to one comprised of lacustrine species. The transition resulted in a significant increase of the threadfin shad, an effective size-selective planktivore, resulting in the removal of all large zooplankton in L Lake, especially *Daphnia*. The removal of larger herbivorous zooplankton probably lessened predation on larger algal species suitable as food, although experimental proof of this process is lacking. Pelagic Rotifera and Protozoa dominated the zooplankton community. Accompanying these changes was an increase in green

and flagellate phytoplankters during 1988 and 1989. Water transparency increased. However, nitrogen and phosphorus concentrations during 1989 indicated continuing favorable conditions for eutrophic algal species, similar to other Southeast reservoirs. Macrophyte cover in the littoral zone increased steadily within the restrictions of slope and wind fetch. Collector-filterer species of benthos, especially, the Asiatic clam *Corbicula fluminea*, replaced much of the collector-gatherer guilds of benthic fauna by 1989. The mechanism for this change cannot be proposed with current information. After four years of life L Lake had reached a stage where it strongly resembled other regional reservoirs.

CONCLUSIONS

- L Lake is a comparatively young reservoir experiencing a successional pattern typical of most regional reservoirs. High rates of nitrogen and phosphorus inputs from the Savannah River have resulted in eutrophic conditions similar to other regional basins which receive an excess of nutrients from their drainage basins. Successional patterns, when interpreted within successional models, are typical of most other reservoirs.
- L-Reactor impacts on the lake ecosystem and the predicted successional trends are confounded, although reactor operations have not created permanent physical or chemical gradients which would impact the structure, size, or distribution of the biota.
- Relative to BBC criteria L Lake compares favorably with other regional basins. Only the dominance of *Microcystis aeruginosa*, a pollution tolerant blue-green algal species, inhabited the lake. No thermally tolerant species have dominated L Lake. Productivity and diversity of all components of the lake ecosystem were within the variation of regional basins, although primary production during the first two years of the lake's existence, were relatively high for basins of this size and latitude. All of the lake's flora and fauna were self-sustaining and not maintained by repeated stockings or reseeds from other communities. Food webs in L Lake provide all trophic groups expected in a temperate reservoir.

REFERENCES

- Abrams, P. A. 1984. Foraging time optimization and interactions in food webs. *Amer. Naturalist* 124:80-96.
- Applied Biology, Inc. 1981. Clarks Hill Lake water quality study. February - October 1981. Prepared under contract no. DACW21-81-C-0011 for U.S. Army Corps of Engineers, Savannah District.
- Baranov, I. V. 1961. Biohydrochemical classification of reservoirs in the European U. S. S. R. pp.139-183. In P. V. Tyurin (ed.), *The storage lakes of the U. S. S. R. and their importance for fishery*. Israil Program for Scientific Translations, Tel Aviv, Israel.
- Baxter, R. M. 1977. Environmental effects of dams and impoundments. *Ann. Rev. Ecol. Sys.* 8:255-283.
- Beckman, L.G., and J.H. Elrod. 1971. Apparent abundance and distribution of young-of-year fish in Lake Oahe, 1965-69. In: Hall, E. (ed.), *Reservoir Fisheries and Limnology*. Spec. Publ. No. 8. Am. Fish. Soc. Washington, DC.
- Beitinger, T.L. 1974. Thermoregulatory behavior and diel activity patterns of bluegill, *Lepomis macrochirus*, following thermal shock. *U.S. Fish and Wild. Serv. Bull.* 72:1087-1093.
- Bowers, J. A. 1988. A Review of Abiotic and Biotic Interactions in Pelagic Communities: Processes Relevant to Lake. DPST 88-707, E. I. du Pont de Nemours & Co., Savannah River Laboratory, Environmental Sciences Division, Aiken, South Carolina 29808-0001
- Bowers, J. A. 1990. DRAFT, WSRC-90-RP-000, Regional Lakes Study in South Carolina, 1988-1989 (U), Westinghouse Savannah River Company, Savannah River Site, Aiken, SC 29808-0001
- Brooks, J. L., and S. I. Dodson. 1965. Predation, body size, and the composition of the plankton. *Science* 150:28-35.
- Brown, H.W. 1974. Handbook on the effects of temperature on some North American fishes. Am. Elec. Power Ser. Corp. Canton, OH. ~
- Carlander, K.D. 1977. *Handbook of Freshwater Fisheries Biology*. Vol.II. The Iowa State Univ. Press. Ames, IA.
- Carpenter, S. R., and J. K. Kitchell. 1984. Plankton community structure and limnetic primary production. *Amer. Naturalist* 124:159-172.
- Carpenter, S. R., J. K. Kitchell, and J. R. Hodgson. 1985. Cascading trophic interaction and lake ecosystem productivity. *Bioscience* 35:635-639.
- Carpenter, S.R. 1983. Submersed macrophyte community structure and internal loading: Relationship to lake ecosystem productivity and succession, pp. 105-111. In: J. Taggart (ed.), *Lake Restoration, Protection and Management*. U.S. E.P.A., Washington, D.C.
- Carr, James H., and Assoc. 1982. Hartwell Lake water quality study. February 1981 - November 1981. Prepared under contract no. DACW 21-81-C-008 for U.S. Army Corps of Engineers, Savannah District.
- Chen, R.L. and J.W. Barko. 1988. Effects of freshwater macrophytes on sediment chemistry. *J. Freshw. Ecol.* 4:279 - 289.
- Clugston, J.P. 1973. The effects of heated effluents from a nuclear reactor on species diversity, abundance, reproduction and movement of fish. Ph.D. Dissertation. Univ. Georgia. Athens, GA.
- Cummins, K. W. 1973. Trophic relations of Aquatic Insects. *Ann. Rev. Entomol.* 18:183-206.
- Duarte, C.M. and J. Kalff. 1986. Littoral slope as a predictor of the maximum biomass of submerged macrophyte communities. *Limnol. Oceanogr.* 31:1072 - 1080.
- Duarte, C.M., J. Kalff, and R.H. Peters. 1986. Patterns in biomass and cover of aquatic macrophytes in lakes. *Can. J. Fish. Aquatic Sci.* 43:1900-1908.
- Duke Power Co. 1985. Catawba Nuclear Station 316(a) demonstration preoperational report. 1973-1974 and 1983, 1984. Lake Wylie chemical and biological characteristics.
- EPA. 1976. Quality Criteria for Water. United States Environmental Protection Agency.

- EPA. 1980. Clean lakes program guidance manual. US EPA. Office of Water Regulations and Standards. EPA/440/581-003.
- Fedorenko, A. Y. 1973. Predation interactions between zooplankton and two species of *Chaoborus* (Diptera, Chaoboridae) in a small coastal lake. M. S. Thesis, Univ. British Columbia. 123p.
- Federenko, A. Y. 1975. Feeding characteristics and predation impact of *Chaoborus* (Diptera, Chaoboridae) larvae in a small lake. *Limnol. Oceanogr.* 20:250-259.
- Firth, P., J.R. O'Hop, B. Coler, and R.A. Green. 1986. Lotic aquatic ecosystems of the Savannah River Plant: impact evaluation, habitat analyses and the lower food chain communities. ECS-SR-26. Prepared by Environmental & Chemical Sciences, Inc. for E.I. du Pont de Nemours & Company. Aiken, SC.
- Gladden, J. B., W. L. Specht, J. A. Bowers, N. V. Halverson, H. E. Mackey, and E. W. Wilde. *Compliance of the Savannah River Plant L-Reactor Cooling System with Environmental Regulations (U)*. USDOE Report DP-1776 Rev. 1 (UC-702), Savannah River Plant, Aiken SC 29808 (1989).
- Gliwicz, Z. M. 1969. Studies on the feeding of pelagic zooplankton in lakes with varying trophy. *Ekologia Polska Series A.* 17:663-708.
- Haney, J. F. 1973. An in situ examination of the grazing activities of natural zooplankton communities. *Archiv fur Hydrobiologie.* 72:87-132.
- Harmon Engineering and Testing Company, Inc. 1983. Water quality management studies. Middle Black Warrior and Lower Tombigbee Rivers. Warrior and Demopolis Lakes. July 1978 - October 1979. Prepared under contract no. DACW-78-C-0181 for U.S. Army Corps of Engineers, Environmental Quality Section. Mobile, AL.
- Hazen, T. C. 1979. Ecology of *Aeromonas hydrophila* in a South Carolina cooling reservoir. *Microbial Ecology* 5:179-195.
- Hazen, T. C. 1983. A model for the density of *Aeromonas hydrophila* in Albemarle Sound, North Carolina. *Microb. Ecology* 9:137-153.
- Hazen, T. C., and G. W. Esch. 1983. Effect of effluent from a nitrogen fertilizer factory and a pulp mill on the distribution and abundance of *Aeromonas hydrophila* in Albemarle Sound, North Carolina. *Appl. and Environmental Microbiol.* 45:31-42.
- Hazen, T. C., C. B. Fliermans, R. P. Hirsch, and G. W. Esch. 1978a. Prevalence and distribution of *Aeromonas hydrophila* in the United States. *Appl. & Environ. Microbiol.* 36:731-738.
- Hazen, T. C., G. W. Esch, A. B. Glassman, and J. W. Gibbons. 1978b. Relationship of season, thermal loading and red-sore disease with various haematological parameters in *Micropterus salmoides*. *J. Fish Biol.* 12:491-498.
- Heuer, J.H., L.A. Kissick, H.G. Mcaling, D.W. Hughes, and B.P. Lindsey. 1989. L-Lake Fish: L-Lake/Steel Creek Biological Monitoring Program. January 1986 - December 1988. NAI-SR-80. Prepared by Normandeau Associates, Inc. Southeast Regional Office of E.I. du Pont de Nemours & Co., Aiken, SC.
- Hickman, G.O., and J.C. Congdon. 1974. Effects of length limits on the fish population of five North Missouri lakes. In: Funk, J.L. (ed.). Symposium on overharvest and management of largemouth bass in small impoundments. N. Central Division, Spec. Publ. No. 3. Am. Fish. Soc.
- Hrbacek, J. 1962. Species composition and the amount of zooplankton in relation to the fish stock. *Rozpr. Cesk. Akad. Ved.* 72:1-116.
- Humphries, S. E., and V. D. Lyne. 1988. Cyanophyte blooms: The role of cell buoyancy. *Limnol. Oceanogr.* 33:79-91.
- Hutchinson, G. E. 1957. A treatise on limnology. Vol. I, Wiley and Sons, London.
- Hutchinson, G. E. 1967. A treatise on limnology. Vol. II, Wiley and Sons, London.
- Keddy, P.A., and P. Constabel. 1985. Germination of ten shoreline plants in relation to seed size, soil particle size and water level: an experimental study. *J. Ecol.* 74:133-141.

- Kimmel, B. L., and A. W. Groeger. 1986. Limnological and ecological changes associated with reservoir aging. In G. E. Hall and M. J. Van Den Avyle, (eds.), *Reservoir Fisheries management: strategies for the '80's*. Amer. Fish. Soc. pp. 103-109.
- Kimsey, C.D., Jr., A.C. Boozer, J.N. Knox, L.E. Turner, K.K. Cain, and G.W. Long. 1982. South Carolina clean lakes classification survey. Tech. Rept. No. 019 - 82, Div. Water Qual. Assess. Enforce. SC DHEC, Columbia, SC.
- King, T.J. 1975. Inhibition of seed germination under leaf canopies in *Arenaria serpyllifolia*, *Veronica arvensis* and *Cerastium holosteoides*. *New Phytol.* 75:87-90.
- Krzyzanek, E., H. Kasza, W. Krzanowski, T. Kuflikowski, and G. Pajak. 1986. Succession of communities in the Goczalkowice Dam Reservoir in the period 1955 - 1982. *Arch. Hydrobiol.* 106:21-43.
- Lagler, K.F. 1956. *Freshwater Fishery Biology*. Wm. C. Brown Company Dubuque, IA.
- Mayhew, J. 1974. Abundance, distribution, mortality and production of O-age fish. Iowa Fed. Aid Proj. F-88R-1 (performance report for jobs 1 and 3). Iowa Conservation Commission. 83 pp.
- McDonald, D.B., and H.F. Bernhard. 1978. Ecological aspects of artificial cooling ponds and lakes. In: Cairns, J., E.F. Benfield, and J.R. Webster (eds.), *Current Perspectives on River-Reservoir Ecosystems*. North Am. Benthol. Soc.
- McLachlan, A.J. 1977. The changing roles of terrestrial and allochthonous organic matter in newly flooded lakes. *Hydrobiologia* 54:215-217.
- Merritt, R.W., and K.W. Cummins. 1984. *An Introduction to the Aquatic Insects of North America*. 2nd ed. Kendall/Hunt. Dubuque, IA.
- Miller, R. W., and W. R. Chapman. 1976. *Epistylis* and *Aeromonas hydrophila* infections in fishes from North Carolina reservoirs. *Prog. Fish Cult.* 38:165-168.
- Moore, D.R.J., and P.A. Keddy. 1988. Effects of a waterdepth gradient on the germination of lakeshore plants. *Can. J. Bot.* 66:548-552.
- Mosher, T. D. 1984. An evaluation of threadfin shad *Dorosoma petenense* introduction in Kansas. Kansas Fish and Game Comm. Project Report FW-9-2, 54 pp.
- NAS (National Academy of Sciences). 1969. Eutrophication: Causes, consequences, correctives. Proceedings of a symposium, Washington, D. C.
- Newman, M., K. Anderson, R. Bayer, and R. Lew. 1987. Comprehensive Cooling Water Report: Vol. 2: Water Quality, Savannah River Ecology Laboratory, Divisions of Wetlands Ecology, Aiken, SC 29808
- Nikolsky, G.V. 1963. *The Ecology of Fishes* Academic Press, New York, NY.
- Nursall, J.R. 1952. The early development of a bottom fauna in a new power reservoir in the Rocky Mountains of Alberta. *Can. J. Zool.* 30:387-409.
- O'Brien, W. J. 1987. Planktivory by freshwater fish: Thrust and parry in the pelagia. pp. 3-16. In W. C. Kerfoot and A. Sih. (eds.), *Predation, Direct and indirect impacts on aquatic communities*, Univ. Press of New England, Hanover, NH.
- Odum, H. T. 1957. Trophic structure and productivity of Silver Springs, Florida. *Ecol. Monogr.* 27:55-112.
- Odum, E.P. 1971. *Fundamentals of Ecology*. 3rd ed. W.B. Saunders Co. Philadelphia, PA.
- Paller, M.H., and B.M. Saul. 1986. Effects of thermal discharges on the distribution and abundance of adult fishes in the Savannah River and selected tributaries. Annual Report. November 1984 - August 1985. ECS-SR-28. Prepared by Environmental & Chemical Sciences, Inc. for E.I. du Pont de Nemours & Company. DPST-8. Aiken, SC.
- Paller, M.H., J.K. Heuer, L.A. Kissick, H.G. Mealing, and D.W. Hughes. 1988. L-Lake Fish: L-Lake/Steel Creek Biological Monitoring Program. January 1986 - December 1987. ECS-SR-65. Prepared by Environmental & Chemical Sciences, Inc. for E.I. du Pont de Nemours & Co., Aiken, SC.

- Parkin, R.B., and J.B. Stahl. 1981. Chironomidae (Diptera) of Baldwin Lake, Illinois, a cooling reservoir. *Hydrobiologia* 76:119-128.
- Paterson, C.G., and C.H. Fernando. 1969a. The macroinvertebrate colonization of a small reservoir in Eastern Canada. *Verh. Internat. Verein. Limnol.* 17:126-136.
- Paterson, C.G., and C.H. Fernando. 1969b. Macroinvertebrate colonization of the marginal zone of a small impoundment in Eastern Canada. *Can. J. Zool.* 47:1229-1238.
- Paterson, C.G., and C.H. Fernando. 1970. Benthic fauna colonization of a new reservoir with particular reference to the Chironomidae. *J. Fish. Res. Board. Can.* 27:213-232.
- Petr, T. 1972. Benthic fauna of a tropical man-made lake (Volta Lake, Ghana 1965 - 1968). *Arch. Hydrobiol.* 70:484-533.
- Porter, K. G., and J. D. Orcutt Jr. 1980. Nutritional adequacy, manageability, and toxicity as factors that determine food quality of green and blue-green algae for *Daphnia*. pp.268-281, *In* Kerfoot, W. C. (ed.), *Evolution and ecology of zooplankton communities*. Univ. Press New England, Hanover, NH.
- Porter, K. G., M. L. Pace, and J. F. Battey. 1979. Ciliate protozoans as a link in freshwater planktonic food chains. *Nature* 227:563-564.
- Raschke, R.L., and P.C. Rusanowski. 1984. Aquatic Macrophyton Field Collection Methods and Laboratory Analyses. *In*: Dennis, W.M. and B.G. Isom (eds.), *Ecological Assessment of Macrophyton: Collection, Use, and Meaning of Data*. Am. Soc. for Testing and Materials.
- Reid, G.K., and R.D. Wood. 1976. *Ecology of Inland Waters and Estuaries*. D. Van Nostrand Company New York, NY.
- Sculthorpe, C.D. 1967. *The Biology of Aquatic Vascular Plants*. Edward Arnold, Ltd. London.
- SCWMRD. 1976. Proceedings of the First South Carolina Endangered Species Symposium. November 11 - 12, 1976. Charleston, SC.
- Shireman, J.V., W.T. Haller, D.E. Canfield, Jr., and V.T. Vandiver. 1982. The impact of aquatic plants and their management techniques on the aquatic resources of the United States: an overview. US EPA/600/4-81/007.
- immerfelt C. 1975. Relationship between weather and year-class strength of largemouth bass. *In*: Clipper, H. (ed.), *Black Bass Biology and Management*. Sport Fishing Institute, Washington, D.C.
- Sweeney, B.W., and R.L. Vannote. 1978. Size variation and the distribution of hemimetabolous aquatic insects: two thermal equilibrium hypotheses. *Science* 200:444-446.
- Taylor, M.P. 1971. Phytoplankton productivity response to nutrients correlated with certain environmental factors in six TVA reservoirs. *In*: Hall, G.E. (ed.), *Reservoir Fisheries and Limnology*. Spec. Publ. No. 8. Am. Fish. Soc. Washington, DC.
- Threlkeld, S. T. 1987. Experimental evaluation of trophic-cascade and nutrient-mediated effects of planktivorous fish on plankton community structure. pp. 161-174, *In* W. C. Kerfoot and A. Sih. (eds.), *Predation, Direct and indirect impacts on aquatic communities*. Univ. Press of New England, Hanover, NH.
- Turner, W.R. 1971. Sport fish harvest from the Rough River, Kentucky, before and after impoundment. *In*: G.E. Hall (ed.), *Reservoir Fisheries and Limnology*. Spec. Publ. No. 8. Am. Fish. Soc. Washington, D.C.
- Van der Valk, A.G. and B.C. Davis. 1978. Primary production of prairie glacial marshes. *In*: Good, R.E., D.F. Whigham, and R.L. Simpson (Eds.), *Freshwater Wetlands: Production Processes and Management Potential*. Academic Press. New York, NY.
- Voshell, J.R., and G.M. Simmons, Jr. 1984. Colonization and succession of benthic macroinvertebrates in a new reservoir. *Hydrobiologia* 112:27-39.
- Walburg, C.H., G.L. Kaiser, and P.L. Hudson. 1971. Lewis and Clark Lake Tailwater Biota and Some Relations of the Tailwater and Reservoir Fish Populations. pp 449-467 *In* G.E. Hall, ed. *Reservoir Fisheries and Limnology*. Am. Fish. Soc. Spec. Publ. No. 8 Washington, DC.

Water and Air Research, Inc. 1982. Water quality management studies Lake Seminole. February - December 1979. Phase II. Tech. Pub. ACF 80-11. Contract #DACW01-78-C-O101.

Wein, G.R., S. Kroeger, and G.J. Pierce. 1987. Lacustrine vegetation establishment within a cooling reservoir. Proc. 14th Annual Conf. on Wetlands Restoration and Creation. Hillsborough Community College.

Wetzel, R.G. 1983. *Limnology*. 2nd ed. Saunders College Publ. Philadelphia, PA.

Wiley, J.M., R.W. Gorden, S.W. Waite, and T. Powless. 1984. The relationship between aquatic macrophytes and fish production in Illinois ponds: a simple approach. N. Am. J. Fish. Manage. 4: 119.

Wollenweider, R.A. 1976. Advances in defining critical loading levels for phosphorus in lake eutrophication. Mem. Inst. Ital. Idrobiol. 33:53-83.

END

DATE
FILMED

6 / 24 / 92