THEORETICAL PARTICLE PHYSICS
Technical Progress Report

Steven A. Gottlieb, Archibald W. Hendry, V. Alan Kostelecký and Don B. Lichtenberg
Physics Department, Indiana University
Bloomington, Indiana 47405

Prepared for the U.S. Department of Energy

~~Contract No. DE-AC02-84ER40125, Task B,~~ May 1, 1991-March 31, 1992

Grant No. DE-FG02-91ER40661, Task B, January 1, 1992-April 30, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ABSTRACT

We have carried out research in lattice gauge theory, superstring theory, supersymmetry, the solar neutrino puzzle, QCD perturbation theory, and phenomenological models of hadrons. In this report, we summarize our work in each of these areas.

CONTENTS

Introduction	2
Research work	3
Personnel	22
Short-term visitors	22
Invited talks, 1989-92	23
Research papers supported in part by DOE, 1989-92	25

INTRODUCTION

In this Technical Progress Report we summarize the principal research of the Task B High Energy Theory Group at Indiana University during the period from May 1, 1991 to April 30, 1992. We have obtained important results in lattice gauge theory, the solar neutrino puzzle, supersymmetry, superstrings, gravitation, applications of QCD perturbation theory, and hadron properties.

The principal investigators are S. Gottlieb, A. Hendry, A. Kostecký, and D. Lichtenberg. Other contributing researchers supported by the contract and grant include postdoctoral research associates A. Krasnitz and R. Potting. One graduate student, J. Tsai, supervised by A. Hendry, received his Ph.D. in August, 1991. Another graduate student, R. Roncaglia, has worked with D. Lichtenberg, but has been supported from other sources. Three graduate students, Ross Hyman, Jason Smith, and Bogdan Tudose, are going to work with A. Kostecký beginning in May, 1992.

We include near the end of this report a list of personnel supported in whole or in part by Task B and a list of names of short-term visitors during the last twelve months. This is followed by a list of recent invited talks and research papers by members of our group.

RESEARCH WORK

STAGGERED SPECTRUM ON THE IPSC/860

(Gottlieb, Krasnitz *et al.*)

In a QCD simulation, as the lattice spacing and quark mass are decreased, one must increase the spatial volume of the lattice in order to avoid finite size effects. If the spatial size of the lattice is too small, the hadrons become squeezed, generating systematic errors in the masses. We have seen some of these effects in the HEMCGC spectrum calculations [1], and this point was spectacularly brought home by Ukawa at Lattice '90 [2,3] with some calculations on very small lattices.

Our desire was to demonstrate that we actually have control of the finite size effects. This requires going to a moderate to large volume and getting very good statistics to show that the effects can be observed where they are still fairly small. Doing this at moderately strong coupling, we can set the appropriate volume for more challenging, weaker coupling calculations, thus insuring from the beginning that finite size effects there will be small. It would also be of value to make contact with the analytic studies of Lüscher [4], and this can be done best if we have a range of pion masses with which to work.

We have used Intel machines at SDSC, the SSC lab and NASA Ames Laboratory during the course of this project. (The DOE has generously supported our use of the San Diego Supercomputer Center iPSC/860 with a \$125,000 supplement to the grant of J. Kuti at UCSD. An additional \$125,000 has come from the NSF and \$120,000 in matching funds from the six universities with researchers in the MILC collaboration.) We now have results for $6/g^2 = 5.445$ and $am_q = 0.025$ on $8^3 \times 24$, $10^3 \times 24$, $12^3 \times 24$ and $16^3 \times 24$ lattices. In addition, we have extensive results for the larger two lattices with a lighter quark mass 0.0125. The larger volume calculation is not quite completed. Below we show our results for the rho and nucleon masses as a function of the spatial length of the lattice. Using the rho mass to set the lattice spacing, we find that we studied boxes with sizes from 1.8 to 3.6 fermis. A box size of 2.7 fm seems comfortable to control the finite size effects at this value of the pion mass.

Some of these results have been presented in an invited talk at the Workshop on Fermion Algorithms [5] and at Lattice '91 [6]. It will be very interesting to complete the analysis with the lighter mass since the finite size effects are supposed to fall exponentially with the pion mass [4].

We have also attempted to study the decay of the rho by looking at the correlation between rho momentum and polarization in the above spectrum studies. Understanding rho decay is clearly of great importance as we decrease the quark mass toward the physical value. Unfortunately, we have not yet seen any splitting between the different rho polarizations.

References

1. Quantum Chromodynamics at $6/g^2 = 5.6$, K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U. M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang, Phys. Rev. Lett. **65**, 2106 (1990); Hadron Spectrum in QCD at $6/g^2 = 5.6$, Phys. Rev. D **42**, 3794 (1990).

Nucleon (a) and rho (b) masses as a function of spatial lattice size N_s . In addition to the data on each of the lattice sizes, the average of all sizes is plotted. Finally, there is a line depicting the size of a 1% error.

2. M. Fukugita, H. Mino, M. Okawa and A. Ukawa, Nucl. Physics B (Proc. Suppl.), 20, 376, 1991.
3. M. Fukugita, H. Mino, M. Okawa and A. Ukawa, Phys. Rev. Lett. **68**, 761, 1992.
4. M. Lüscher, Communications in Mathematical Physics, **104**, 177, 1986; and in *Fields, Strings and Critical Phenomena*, edited by E. Brèzin and J. Zinn-Justin (North-Holland, Amsterdam, 1990).
5. QCD on the iPSC/860, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, in *Workshop on Fermion Algorithms*, edited by H. J. Hermann and F. Karsch, (World Scientific, Singapore, 1991).
6. Finite-Size Effects and T_c for $\beta = 5.445$ with Two Staggered Flavors, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, in Proceedings of Lattice '91, Tsukuba, Japan, Nucl. Phys. B(Proc. Suppl.) to appear.

QCD THERMODYNAMICS WITH STAGGERED FERMIONS

(Gottlieb, Krasnitz *et al.*)

For some time, we have been studying QCD at finite temperature with an eye toward predicting the temperature for deconfinement [1]. This number is quite interesting for heavy-ion collision

experiments, such as those that will be conducted at RHIC, because it indicates how hard it will be to produce the quark-gluon plasma. So far, we have seen that with dynamical fermions the transition temperature is markedly reduced as compared with pure gluon calculations. The temperature for two flavors of dynamical fermions is about 150 MeV, compared with about 230 in the pure glue theory. Calculations done so far have been at relatively strong coupling. It is important to extend these calculations to weaker coupling because results are not entirely trustworthy until the scaling regime has been reached [2].

In the quenched approximation, the transition couplings are known up to $N_t = 14$, where N_t is the number of time slices. (The continuum limit corresponds to increasing N_t .) However, with dynamical fermions, only up to 8 time slices have been studied. We have refined our calculations of the crossover couplings for $N_t = 6$ and 8, using the Intel iPSC/860 at the San Diego Supercomputer Center and the CM2 at the Pittsburgh Supercomputer Center, respectively. These results have been published in the International Journal of Supercomputer Applications in an issue highlighting work at the NSF Supercomputer Centers [3,4], at various conferences [5-9] and in Physical Review [10]. We find that the best current estimate of the crossover temperature for restoration of chiral symmetry is 147 ± 7 MeV, where the rho mass is used to set the scale. The quality of the dynamical fermion results is now such that their statistical error bars are as small as or smaller than for the quenched approximation.

One of the results that we are quite excited about involves looking at the spatial correlation of the quarks in the plasma phase. We have compared the analogous quantities at zero and finite temperature and find very little difference in the correlations. The falloff of the correlation as a function of distance in the zero temperature phase is indicative of the confinement scale. We expected to see a slower falloff in the high temperature phase indicating the onset of deconfinement. This result has been published in Physical Review Letters [11] and presented at Quark Matter '91 [12] and Lattice '91 [7].

References

1. S. Gottlieb, W. Liu, D. Toussaint, R.L. Renken and R.L. Sugar, Phys. Rev. Lett. **59**, 1513 (1987).
2. For a review of the current status of QCD thermodynamics, see S. Gottlieb, Nucl. Phys. B(Proc. Suppl.)**20**, 247, 1991.
3. Studying Quarks and Gluons on MIMD Parallel Computers, C. Bernard, M. C. Ogilvie, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, R. L. Sugar and D. Toussaint, Internat. J. Supercomput. Appl. **5**, 61, 1991.
4. Hadron Thermodynamics on the Connection Machine, S. Gottlieb, A. Krasnitz, U. M. Heller, A. D. Kennedy, W. Liu, J. B. Kogut, R. L. Renken, D. K. Sinclair, K. C. Wang, R. L. Sugar and D. Toussaint, Internat. J. Supercomput. Appl. **5**, 50, 1991.
5. QCD on the iPSC/860, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, in *Workshop on Fermion Algorithms*, edited by

- H. J. Hermann and F. Karsch, (World Scientific, Singapore, 1991).
6. Finite Temperature Lattice QCD Simulations with Staggered and Wilson Fermions, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, U. M. Heller, A. D. Kennedy, J. B. Kogut, A. Krasnitz, W. Liu, M. C. Ogilvie, R. L. Renken, D. K. Sinclair, R. L. Sugar, D. Toussaint, and K. C. Wang, in Proceedings of Hot Summer Daze, a workshop at Brookhaven National Laboratory, August, 1991, to be published by World Scientific.
 7. QCD Thermodynamics with Two Flavors of Quarks, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, in Proceedings of Lattice '91, Tsukuba, Japan, Nucl. Phys. B(Proc. Suppl.) to appear.
 8. Thermodynamics of Lattice QCD with 2 Light Dynamical (Staggered) Quark Flavours on a $16^3 \times 8$ Lattice, S. Gottlieb, U. M. Heller, A. D. Kennedy, J. B. Kogut, A. Krasnitz, W. Liu, R. L. Renken, D. K. Sinclair, R. L. Sugar, D. Toussaint and K. C. Wang, in Proceedings of Lattice '91, Tsukuba, Japan, Nucl. Phys. B(Proc. Suppl.) to appear.
 9. Nuclear physics at 1,700,000,000,000 degrees Kelvin, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, to be published in the proceedings of the *1991 Intel Supercomputer Users Group*, Dallas, Texas, 1991.
 10. QCD Thermodynamics with Two Flavors at $N_t = 6$, (with C. Bernard, T. A. DeGrand, C. DeTar, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), Phys. Rev., to appear.
 11. The Spatial Structure of Screening Propagators in Hot QCD, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, Phys. Rev. Lett., **68**, 2125, 1992.
 12. Quark Structure of Static Correlators in High Temperature QCD, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint, in Proceedings of Quark Matter '91, to be published.

GRAND CHALLENGE SPECTRUM CALCULATION

(Gottlieb, Krasnitz *et al.*)

The High Energy Monte Carlo Grand Challenge Collaboration has been working for some three years on state-of-the-art simulations of QCD [1] with dynamical fermions, first on the Supercomputer Computations Research Institute (SCRI) ETA10, and more recently on the Connection Machine installed to replace the ETA10. During that period, we have published ten papers [2-11].

During the past year, we have completed a QCD simulation on a $16^3 \times 32$ lattice with gauge coupling $6/g^2 = 5.6$ using staggered fermions. We have used two masses $am_q = 0.025$ and 0.01 . These are the same masses that we used previously. For the heavier mass, our previous calculations were done on 12^4 lattices, doubled in time for the spectrum analysis, and our desire in the current calculation is to check finite size effects. For the lighter mass, we had done a simulation on a 16^4 lattice replicated in time for the spectrum analysis. We had seen a very strange behavior in the

effective pion mass that we attributed to replication in time. Because the CM5 is both faster than the ETA10 and has much more memory, we could avoid time replication in the current calculation. We find, as shown below, that the pion effective mass now has a smooth plateau. Additional theoretical evidence for replication in time as a source of the strange oscillations comes from a study of a model system.[12]

We are still working on the mass fitting for the other particles. Our preliminary findings are that there are not big finite size effects, though there may be some for the lighter quark mass. The latter effect had already been seen with the 16^4 lattice; however, we now have improved statistics for the rho mass. For the nucleon, we have a new source not previously studied, and we are still trying to complete the analysis.

We have also used the gauge configurations generated here to look at the spectrum of Wilson valence quarks [13]. Here we find some evidence for differences between quenched calculations and this hybrid calculation. The nucleon to rho mass ratio is smaller in the quenched approximation than with our lighter dynamical quark mass.

Some preliminary results from both types of valence quarks were presented at Tsukuba [14].

Plot of the effective pion mass versus distance from the source. Note the dip near distance 10 in the results for the 16^4 lattice doubled in time (◇). We believe that this is an artifact of the doubling in time, as it is not present in the current run (squares and crosses).

References

1. For recent reviews, see A. Ukawa, Nucl. Phys. B (Proc. Suppl.), **10A**, 66, 1989, T. DeGrand, Nucl. Phys. B (Proc. Suppl.), **20**, 353, 1991, or the one by D. Toussaint at Lattice '91.
2. Hadron Spectrum with Staggered Dynamical Quarks, K. M. Bitar, R. Edwards, U. M. Heller, A. D. Kennedy, T. A. DeGrand, S. Gottlieb, A. Krasnitz, J. B. Kogut, R. L. Renken, M. C. Ogilvie, P. Rossi, D. K. Sinclair, K. C. Wang, R. L. Sugar, M. Teper and D. Toussaint, Nucl. Phys. B(Proc. Suppl.)**17**, 404 (1990).
3. Hadronic Spectroscopy with Wilson Valence Quarks, K. M. Bitar *et al.*, Nucl. Phys. B(Proc. Suppl.)**17**, 400 (1990).
4. QCD with Dynamical Staggered Quarks, K. M. Bitar *et al.*, in *Proceedings of The Rice Meeting*, edited by B. Bonner and H. Miettinen (World Scientific, Singapore, 1990).
5. Hadron Spectrum in QCD at $6/g^2 = 5.6$, K. M. Bitar *et al.*, Phys. Rev. D **42**, 3794 (1990).
6. The High Energy Monte Carlo Grand Challenge: Simulating Quarks and Gluons, K. M. Bitar *et al.*, Internat. J. Supercomput. Appl. **4**, 48 (1990).
7. Quantum Chromodynamics at $6/g^2 = 5.6$, K. M. Bitar *et al.*, Phys. Rev. Lett. **65**, 2106 (1990).
8. Lattice Spectrum with Dynamical Staggered Quarks at $6/g^2 = 5.6$, K. M. Bitar *et al.*, presented at PANIC XII, to appear in Proceedings of the Twelfth International Conference on Particle and Nuclei, Massachusetts Institute of Technology, June 24–29, 1990, edited by J. I. Matthews *et al.*
9. Hadronic Spectroscopy at $a \approx 0.15$ fm, K. M. Bitar *et al.*, Nucl. Phys. B(Proc. Suppl.)**20**, 362, 1991.
10. Glueball and Topological Charge in the Presence of Dynamical Quarks, K. M. Bitar *et al.*, Nucl. Phys. B(Proc. Suppl.)**20**, 390, 1991.
11. On Glueballs and Topology in Lattice QCD with Two Light Flavors, K. M. Bitar *et al.*, Phys. Rev. D **44**, 2090, 1991.
12. Spectroscopy with Multiple Field Configurations, A. Krasnitz, Phys. Rev. D **43**, 2694, 1991.
13. Hadron Spectrum in QCD with Valence Wilson Fermions and Dynamical Staggered Fermions at $6/g^2 = 5.6$, K. M. Bitar *et al.*, Submitted to Physical Review.
14. QCD Hadron Spectroscopy with Staggered Dynamical Quarks at $\beta = 5.6$, K. M. Bitar *et al.*, to appear in Nucl. Phys. B(Proc. Suppl.), Proceedings of Lattice '91.

QCD THERMODYNAMICS WITH WILSON FERMIONS

(Gottlieb, Krasnitz *et al.*)

The study of high temperature QCD with Wilson quarks has received much less attention than the study with staggered quarks. The staggered quarks have the advantage of retaining a remnant of chiral symmetry on the lattice, and the chiral limit can be approached by simply decreasing the quark mass to zero. However, for finite lattice spacing, the flavor symmetry among the pion states is broken, and full chiral symmetry is achieved only in the continuum limit. By contrast, the Wilson quark action explicitly breaks chiral symmetry for finite lattice spacing. In order to identify the chiral limit, one must search in the two-dimensional space of the gauge coupling and hopping parameter. Given the very different behaviors of the two formulations of quarks for finite lattice spacing, it is essential to study both and to verify that they do yield the same continuum results.

Some studies with Wilson fermions have suggested that for $N_t = 4$, it might not even be possible to have a light pion and be in the low temperature phase [1]. This is certainly not the case for staggered fermions. For staggered fermions, we have studied hadron screening lengths [2] to shed light on the high temperature phase. Previously, we have done similar studies with Wilson quarks using the ETA10 at the John von Neumann Center [3,4]. During the past year, we wanted to push further toward the continuum limit which requires increasing the value of N_t , and a concomitant increase in the computing resources required.

At the San Diego Supercomputer Center, we have completed a study of high temperature QCD with Wilson quarks using the iPSC/860. This work used $12^3 \times 6$ lattices with two flavors of quarks, using hopping parameters of $\kappa = 0.16, 0.17$ and 0.18 . We have determined the location of the crossover between the high and low temperature states for these hopping parameters, and have carried out spectrum calculations for the same hopping parameters on $12^3 \times 24$ lattices in order to determine the critical value of the gauge coupling, that is, the value for which the pion and quark masses vanish. One unexpected result is that for the larger hopping parameters (smaller quark masses), there is evidence for coexistence between the high and low temperature states. Whether this result is indicative of a first order phase transition or is an artifact of the rather large lattice spacing at which we are working is not clear, but the possibility that the two formulations of quarks give very different thermodynamics deserves further study. At all of these κ values the zero temperature pion mass remained large at the values of κ and $6/g^2$ corresponding to the high temperature transition with six time slices. This indicates that we have not approached the point in the phase diagram corresponding to light quarks, where the zero temperature pion mass should vanish. (Equivalently, κ_c , where the zero temperature squared pion mass extrapolates to zero, should equal κ_t , where we see the crossover to high temperature behavior.) Some preliminary results of this work were presented in an invited talk [5] at the Brookhaven workshop Hot Summer Daze, organized by the theory group in support of the RHIC program. Additional results were presented at the Lattice '91 conference [6], and a full writeup is being prepared for publication.

The results for Wilson fermions are quite tantalizing, and it is clear that this program should

be pursued with weaker coupling. We intend to do so in the coming year.

References

1. M. Fukugita, S. Ohta and A. Ukawa, Phys. Rev. Lett. **57**, 1974 (1986).
2. S. Gottlieb, W. Liu, D. Toussaint, R.L. Renken and R.L. Sugar, Phys. Rev. Lett. **59**, 1881 (1987); C.E. DeTar and J.B. Kogut, Phys. Rev. Lett. **59**, 399 (1987).
3. Hadron Thermodynamics with Wilson Quarks, K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U.M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint and K.C. Wang, Phys. Rev. D **43**, 2396 (1991).
4. QCD Thermodynamics with Wilson Quarks, K.M. Bitar *et al.*, Nucl. Phys. B(Proc. Suppl.)**20**, 272, 1991.
5. Finite Temperature Lattice QCD Simulations with Staggered and Wilson Fermions, C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, U. M. Heller, A. D. Kennedy, J. B. Kogut, A. Krasnitz, W. Liu, M. C. Ogilvie, R. L. Renken, D. K. Sinclair, R. L. Sugar, D. Toussaint, and K. C. Wang, in Proceedings of Hot Summer Daze, a workshop at Brookhaven National Laboratory, August, 1991, to be published by World Scientific.
6. The MIMD Lattice Computations (MILC) Collaboration, QCD Thermodynamics with Two Flavors of Quarks, to be published in Nuc. Phys. B(Proc. Suppl.), proceedings of Lattice '91.

QCD TERAFLUP PROJECT

(Gottlieb, and many others)

I am a member of the steering committee of the QCD Teraflop project. We have proposed that DOE fund the construction of a \$40 million computer capable of sustaining a speed of 1 trillion floating point operations a second on lattice QCD calculations. We have published the physics goals of our proposal [1] and have suggested that the site for the computer be Brookhaven National Laboratory. I plan to spend my sabbatical there next year in order to make maximal contributions to the project.

References

1. Physics Goals of the QCD Teraflop Project, the QCD Teraflop Collaboration, Intl. J. Mod. Phys. C, **2**, 829, 1991.

NEUTRINOS FROM THE SUN

(Hendry and Tsai)

The study of neutrinos from the Sun seems to be about the only way in the foreseeable future of getting an estimate of neutrino masses and mixings, should they be as small as they are presently thought to be. These properties of neutrinos are tantalizingly important since they may yield the first clues to what lies beyond the present standard model of particle physics.

There are however two aspects to resolving the neutrino problem. One is exactly how they are produced in the Sun, the other is related to how they are transported through the solar material as they escape from the hot core. So far, most attention has been fixed on the latter aspect. If at least one variety of neutrinos has mass, neutrino oscillations are possible [1,2]; the existing measurements [3-5] of the electron neutrino flux from the Sun can then be used to pin down an allowed region in Δm^2 , $\sin^2 2\theta$ space. Unfortunately much less scrutiny has been given to the production of neutrinos in the Sun. The reason is that astrophysicists [6,7] have developed a very detailed model of the Sun, the so-called standard solar model (SSM), which seems to describe rather well the general characteristics of the Sun. However, as pointed out Bahcall [8], the SSM is not infallible since it rests on a number of assumptions that are hard to prove, as well as on a great deal of input nuclear and atomic data.

Clearly one must consider both neutrino production and neutrino oscillations to pin down the fundamental properties of neutrinos. The two aspects are intimately intertwined. Neutrino oscillations for example depend on the density of electrons that the neutrinos meet as they make their way out through the Sun; density of electrons however is something that comes from solar models. During the past year therefore, we have taken a closer look at solar models.

Our first step has been to re-examine the SSM itself. We have shown that the SSM exhibits several interesting characteristics, not previously noticed. One is that the Sun seems to behave as a set of three distinct regions. In each region, the local pressure varies with the local density according to the relationship $P = K\rho^\gamma$. A gas satisfying a relationship of this kind is called a polytrope. In the Sun, the convective zone of the SSM has this property, while the radiative zone of the SSM has two such regions. This observation therefore allows us to derive a novel phenomenological model of the Sun in which the Sun is represented by a set of three matching polytropes. This work is described in ref. [9]. The model approximates rather well the pressure, density and temperature profiles of the SSM. One can also calculate the neutrino fluxes (in particular, the boron-8 and pp neutrinos) in such a model, and again good agreement with the SSM is obtained.

Having developed a phenomenological model to reproduce the results of the SSM, we can now generate variations of this model. These represent perturbations away from the SSM. (However each variant must still satisfy the observed constraints of radius $R_\odot$, mass $M_\odot$, surface temperature T_s and surface chemical composition μ_s .) They can therefore be used to probe the sensitivity of various quantities to small perturbations from the SSM. Some of this work constitutes the Ph.D. thesis of J. Tsai [10], and is also described in ref. [11]. So far, we have concentrated on a set of model-suns whose core temperatures are lower than those of the SSM and have shown quantitatively

how the neutrino fluxes change. One can for example account for all or part of the boron-8 neutrino deficit in this way.

Different model-suns yield not only different neutrino fluxes, but also different electron densities. Thus the allowed regions in Δm^2 , $\sin^2 2\theta$ are different from the region determined from the SSM. We are at present calculating these changes, as well as the associated changes in the energy spectrum of the solar neutrino output. With a new result from GALLEX [12] imminent, and further results forthcoming in the next few years, neutrino physics in the 1990's may well provide the much-needed clues for future progress in particle theory.

References

1. S.P. Mikheyev and A. Yu. Smirnov, Sov. J. Nucl. Phys. 42 (1985) 913;
L. Wolfenstein, Phys. Rev. D17 (1978) 2369 and D20 (1979) 2634.
2. H.A. Bethe, Phys. Rev. Lett. 56 (1986) 1305 and 63 (1989) 837;
S.P. Rosen and J.M. Gelb, Phys. Rev. D34 (1986) 969; *ibid.* D39 (1989) 3190;
E.W. Kolb, M.S. Turner and T.P. Walker, Phys. Lett. B175 (1986) 478;
T.K. Kuo and J. Pantaleone, Phys. Rev. D39 (1989) 1930; D41 (1990) 3842; and Rev. Mod. Phys. 61 (1989) 937.
3. R. Davis, Jr. in *Proc. of the Seventh Workshop on Grand Unification (ICOBAN '86)*, Toyama (1986), ed. by J. Arafune (World Scientific, Singapore 1986).
4. K.S. Hirata et al., Phys. Rev. Lett. 65 (1990) 1297; *ibid.* 63 (1989) 16; Phys. Rev. D44 (1991) 2241.
5. A.I. Abazov et al. (SAGE collaboration), Phys. Rev. Letts. 67 (1991) 3332.
6. J.N. Bahcall and R.K. Ulrich, Rev. Mod. Phys. 60 (1988) 297.
7. S. Turke-Chièze, S. Cahen, M. Cassé and C. Doom, Astrophysics J. 335 (1988) 415.
8. J.N. Bahcall, *Neutrino Astrophysics* (Camb. Univ. Press, Cambridge 1989).
9. A.W. Hendry, Indiana preprint IUHET 214 (1992).
10. J. Tsai, Ph.D. thesis (Indiana University, August, 1991).
11. A.W. Hendry and J. Tsai, Indiana preprint IUHET 203 (1992).
12. GALLEX collaboration, see T. Kirsten, *Proc. 14th International Conference on Neutrino Physics and Astrophysics (Neutrino'90)*, Nucl. Phys. B (Proc. Suppl.) 19 (1991), ed. J. Panman and K. Winter, North Holland 1991.

POSSIBLE STRING SIGNATURE FROM CPT VIOLATION

(Kostelecký and Potting)

The CPT theorem offers a profound connection among discrete symmetries in particle theory. Experimental tests [1] with kaons have placed bounds on any potential CPT violations of less

than five parts in 10^{18} . It is thought that the theorem is exact for local relativistic quantum field theories. It has been demonstrated both by direct construction [2-5] and via axiomatic field theory [6,7].

Covariant field theories of strings [8] are known to contain stringy features. For instance, the string field action contains infinitely many fields and interactions and has arbitrary powers of derivative interactions. For string theory, the presence of an exact and rigorous result for CPT valid for local particle theories together with the existence of precision experiments means the discrete symmetry CPT is particularly interesting. The idea is that the extended nature of strings implies that they need not necessarily preserve spacetime CPT .

In a recent publication [9], we studied the spacetime CPT properties of the basic critical open strings and open superstrings, for which technical aspects of gravity are avoidable and covariant string field theories exist. We extended the approach of ref. [10] for particle theories to the string case, to sidestep difficulties with the standard Pauli-type proof that would necessitate further developments of the formalism of string field theory. This approach enabled us to extract the behavior of the string fields under CPT transformations for both the theories considered. We were thus able to examine the question of violation of CPT in an explicit framework.

It turns out that in string field theory a natural mechanism exists that can break CPT spontaneously. In usual particle field theories, this mechanism does not appear. It can therefore be viewed as a stringy feature. As the four-dimensional CPT symmetry contained in the higher-dimensional string theory can be violated along with the full CPT symmetry, we have argued in [9] that in a low-energy effective field theory in four dimensions arising from a realistic string theory might contain some breaking of CPT .

More recently, we have extended our investigation into the mechanism of CPT breaking in string theory, with the goal of identifying a particular signature in the kaon system that would be a consequence of an underlying stringy physics. We have therefore examined effective lagrangians with terms of the string CPT -violating type for physical systems. In a favorable scenario, the results we have obtained recently may offer the possibility of identifying an experimental signature for string theory that would be accessible to direct test within the next decade.

References

1. R. Carosi et al., Phys. Lett. B **237**, 303 (1990).
2. J. Schwinger, Phys. Rev. **82**, 914 (1951).
3. G. Lüders, Det. Kong. Danske Videnskabernes Selskab Mat.-fysiske Meddelelser **28**, no. 5 (1954); Ann. Phys. (N.Y.) **2**, 1 (1957).
4. W. Pauli, p. 30 in W. Pauli, ed., *Niels Bohr and the Development of Physics*, McGraw-Hill, New York, 1955.
5. G. Lüders and B. Zumino, Phys. Rev. **106**, 345 (1957).
6. R.F. Streater and A.S. Wightman, *PCT, Spin and Statistics, and All That*, Benjamin Cummings, Reading, 1964.

7. R. Jost, *The General Theory of Quantized Fields*, AMS, Providence, 1965.
8. E. Witten, Nucl. Phys. **B268**, 253 (1986); Nucl. Phys. **B276**, 291 (1986); C. Preitschopf, C.B. Thorn and S.A. Yost, Nucl. Phys. **B337**, 363 (1990); I.Ya. Aref'eva, P. Medvedev and A. Zubarev, Nucl. Phys. **B341**, 464 (1990).
9. V.A. Kostelecký and R. Potting, Nucl. Phys. **B359**, 545 (1991).
10. R.G. Sachs, Prog. Theor. Phys. Suppl. **86**, 336 (1986).

TIME-DEPENDENT SUPERCOHERENT STATES

(Kostelecký, Man'ko, Nieto and Truax)

Superalgebras are mathematical constructs formally consisting of a set of generators satisfying commutation and anticommutation relations. Extensively investigated in the last twenty years, they play an important role in that they contain information on infinitesimal supersymmetry transformations. There are now many subfields of physics in which supersymmetries arise [1]. In many cases, it suffices to consider infinitesimal transformations. Partly for this reason and partly due to inherent technical difficulties, the mathematical structure involved with *finite* supersymmetry [2] transformations has been less extensively studied. Over several years our group has been involved in an ongoing investigation of these finite transformations, called supergroups, paying especial attention to their use in physical situations.

We initiated this program with a study of the basic features of supergroups via their one-parameter subgroups [3]. This provided us with a set of canonical coordinates and parametrizations on the supergroup manifold. The next step was to develop a general approach to the construction of formulae connecting the various schemes. These analytical formulae, called Baker-Campbell-Hausdorff (BCH) relations, can be found by setting up and solving a system of simultaneous nonlinear differential equations with Grassmann-valued dependent variables [4]. We thereby established [5,6] BCH relations for the simple orthosymplectic supergroup $OSP(1|2)$ and for the super-Poincaré group $IOSP(1|4)$. (Both these supergroups are widely applied in physical situations.) In a separate development arising from our resulting increased understanding of Grassmann valued differential equations, we were able to reformulate fluid dynamics in Grassmann variables and find solution sets for compressible fluid flow [7].

In more recent work, we have addressed one large class of physical situations in which the group structure is important: coherent states. These are of great value in a broad set of situations, from descriptions of laser physics to the mathematics of path integrals [8]. An open problem in coherent state theory during the past decade was the issue of constructing natural extensions of coherent states for supersymmetric systems. A number of distinct definitions exist for the usual coherent states. For instance, for the simple harmonic oscillator definitions that have been adopted include: minimum uncertainty states that preserve the uncertainty in time, eigenstates of the annihilation

operator, and the displacement operator. All these result in coherent states that agree, and they have several desirable aspects. For the supersymmetric version of the harmonic oscillator, however, the natural extension of these definitions was lacking. We were able to solve this difficulty by extending to supergroups the generalized displacement operator approach used with Lie groups [9]. One result [10] was our 'supercoherent' states that for the supersymmetric harmonic oscillator have extensions of all the usual attractive properties. Another consequence was the possibility of constructing natural supercoherent states for *any* supersymmetric model. We used our method to develop explicit supercoherent states for a theory with an electron moving in a constant magnetic field, and for a theory describing an electron in the field of a monopole.

In the past year, we have succeeded in combining our methods with developments introduced by a group of Soviet physicists to handle time-dependent situations [11]. In particular, we have been investigating the issue of constructing supercoherent states for the quantum motion of an electron in a *time-dependent* magnetic field. This technically involved calculation demonstrates that the quantum motion is inherently supersymmetric and thereby provides a solution to the question of constructing closest-to-classical behavior of an electron subjected to a varying field.

References

1. An overview of the applications of supersymmetry to physics may be found in: *Supersymmetry in Physics*, eds. V.A. Kostecký and D.K. Campbell (North-Holland, Amsterdam, 1985). These must now be supplemented by a number of more recent developments in nuclear physics, atomic physics, condensed matter physics, and string theory.
2. A. Rogers, J. Math. Phys. **22**, 939 (1981).
3. D.R. Truax, V.A. Kostecký and M.M. Nieto, J. Math. Phys. **27**, 754 (1986).
4. V.A. Kostecký, M.M. Nieto and D.R. Truax, J. Math. Phys. **27**, 1419 (1986).
5. V.A. Kostecký and D.R. Truax, J. Math. Phys. **28**, 2480 (1987).
6. B.W. Fatyga, V.A. Kostecký and D.R. Truax, J. Math. Phys., **30**, 291 (1989).
7. B.W. Fatyga, V.A. Kostecký and D.R. Truax, J. Math. Phys., **30**, 1464 (1989).
8. See, for example, J.R. Klauder and B.-S. Skagerstam, *Coherent States* (World Scientific, Singapore, 1985).
9. See, for example, A.M. Perelomov, *Generalized Coherent States and Their Applications* (Springer-Verlag, Berlin, 1986).
10. B.W. Fatyga, V.A. Kostecký, M.M. Nieto and D.R. Truax, Phys. Rev. D **43**, 1403 (1991).
11. V.A. Kostecký, V. Man'ko, M.M. Nieto, D.R. Truax, in preparation.

FERMIONS WITH SPACELIKE DISPERSION

(Chodos, Kostelecký, Potting, and Gates)

The issue of constructing consistent quantum theories for fields with spacelike dispersion has occupied the attention of many researchers over the years [1]. For important technical reasons involving the representation structure of the Poincaré group, most of the literature has been devoted to the spin-0 case. In 1985, two of us authored a paper in which the possibility of spin-1/2 fields with spacelike dispersion was considered [2]. In that work, we presented a modified hermitian lagrangian for the free-fermion case that is Lorentz invariant and that has spacelike solutions. The straightforward quantization of this theory contains negative-norm states, as might be anticipated on the basis of the general unitarity theorems of representation theory. However, we noted that interactions for such fermions written with left-chirality projections are just what is needed to ensure zeroth-order decoupling of the negative-norm states.

On the basis of this observation and some other technical arguments, we suggested the possibility that fermions with spacelike dispersion might exist in nature and pointed out that the natural candidates for these would be neutrinos, whose dispersion at that time was not well measured. We also examined a number of experimental signatures in an effort to identify candidate tests.

In the ensuing years, further experimental data relevant to this issue have been obtained. The results remain ambiguous but are suggestive, and certainly do not rule out the possibility of an effect. In a recent investigation [3], we have therefore extended our analysis to propose an additional experimental test and to incorporate several alternative suggestions for handling the negative-norm states. The test requires a precision measurement that is likely to be feasible. Among the suggestions is the possibility of attempting to use an infinite-component theory to avoid the difficulties. Some evidence to support this approach was adduced. We also considered the possibility of applying an appropriate constraint to our original lagrangian in order to define a physical subspace free of negative norms, much as occurs in the covariant quantization of gauge field theories. This may lead to a phenomenologically satisfactory model but a completely adequate theory has not yet been shown to exist.

References

1. For some early literature on the subject, see, e.g., G. Feinberg, Phys. Rev. **162**, 1274 (1967), Phys. Rev. D **17**, 1651 (1978); M.E. Arons and E.C.G. Sudarshan, Phys. Rev. **173**, 1622 (1968).
2. A. Chodos, V.A. Kostelecký, Phys. Lett. B **150**, 431 (1985).
3. A. Chodos, V.A. Kostelecký, R. Potting, E. Gates, Mod. Phys. Lett. **7**, 467 (1992).

PARABOLIC SUPERSYMMETRIES

(Bluhm and Kostelecký)

Supersymmetry is a mathematical construct that is widely used in various subfields of physics [1]. One of the few known examples of supersymmetry realized in nature is one that interrelates the eigenvalues [2] and eigenfunctions of distinct elements [3]. Using a supersymmetry for a radial equation in the context of a quantum defect theory, it has been possible to construct *analytical* wavefunctions [4]. Fine-structure effects can also be incorporated [5]. These wavefunctions can be used to obtain oscillator strengths, which have been shown to be in good agreement with experimental data and with numerical computations [4,6]. They can also be used as trial wavefunctions for detailed simulations [7].

In a recent investigation [8], we have made a detailed study of supersymmetry in the Schrodinger equation in a large class of coordinate systems. Focussing on the directly separable case, we have performed an exhaustive search for alternate supersymmetries that might in principle also have physical import. We have discovered that the radial supersymmetry also manifests itself in conical coordinates. We also identified a novel pair of entwined supersymmetries in parabolic coordinates. The eigenvalues and eigenfunctions can be obtained explicitly, as before, and the physical interpretation is different. It is possible that these supersymmetries could have a useful role to play in the calculation of matrix elements for the Stark effect.

References

1. An overview of the applications of supersymmetry to physics may be found in: *Supersymmetry in Physics*, eds. V.A. Kostelecký and D.K. Campbell (North-Holland, Amsterdam, 1985). These must now be supplemented by a number of more recent developments in nuclear physics, atomic physics, condensed matter physics, and string theory.
2. V.A. Kostelecký and M.M. Nieto, Phys. Rev. Lett. **53**, 2285 (1984).
3. V.A. Kostelecký and M.M. Nieto, Phys. Rev. A **32**, 1293 (1985).
4. V.A. Kostelecký and M. M. Nieto, Phys. Rev. A **32**, 3243 (1985).
5. V.A. Kostelecký, M.M. Nieto and D.R. Truax, Phys. Rev. A **38**, 4413 (1988).
6. M.T. Djerad, J. Phys. II **1**, 1 (1991).
7. See, for example, R.E.H. Clark and A.L. Merts, J. Quant. Sp. Rad. Tr. **38**, 287 (1987).
8. R. Bluhm and V.A. Kostelecký, in preparation.

COLOR-HYPERFINE SPLITTING FROM PERTURBATIVE QCD

(Lichtenberg, Potting, Predazzi, Roncaglia)

Much attention has been focused recently on the P -wave color-hyperfine splitting in charmonium [1-4]. The reason is that a measurement of this splitting has been undertaken in E760 at

Fermilab. Recently, a preliminary result has been reported [5].

In one paper on this subject [1], we pointed out that in the generalized Fermi-Breit (GFB) approximation, the center of gravity of the three 3P_J states of charmonium lies a few MeV higher than the 1P_1 state, whereas in one-loop perturbative QCD calculations, the reverse is true [6–8]. The preliminary measurement [5] favors the QCD calculations. In a second paper, we extended the one-loop QCD calculations to include the effects of a running coupling constant. We were able to obtain approximate upper and lower bounds on the splitting. The measurement [5] lies between those bounds.

Our considerations [1,4] also apply to the P -wave color-hyperfine splitting in bottomonium. However, no experiment is presently underway to test the theory in the case of bottomonium.

We are presently looking at the effects on charmonium and bottomonium P states of tensor and spin-orbit forces in addition to the color-hyperfine interaction. The tensor and color-hyperfine interactions are short range, and can be calculated reasonably well in QCD perturbation theory, but the spin-orbit forces arise in part from nonperturbative effects of QCD, and are less well known.

References

1. D. B. Lichtenberg, E. Predazzi, R. Roncaglia, Phys. Rev. D 45, 3265 (1992).
2. J. Stubbe and A. Martin, Phys. Lett B 271, 208 (1991)
3. K. T. Chao, Y. B. Ding, and D. H. Qin, Peking U. preprint PUTP-91-05 (1991).
4. D. B. Lichtenberg and R. Potting, IUHET 208, MAD/PH/668 (March, 1992).
5. R. Cester, seminar at Fermilab (Jan, 1992).
6. S. N. Gupta, S. F. Radford and W. W. Repko, Phys. Rev. D 26, 3305 (1982); 34, 201 (1986).
7. J. Pantaleone, S.-H. H. Tye, and Y. J. Ng, Phys. Rev. D 33, 777 (1986).
8. J. Pantaleone and S.-H. H. Tye, Phys. Rev. D 37, 3337 (1988).

DIQUARKS

(Lichtenberg, Anselmino, Ekelin, Fredriksson, Predazzi)

We have finished a comprehensive review of diquarks [1] which we began last year. We have been in communication with the editors of *Reviews of Modern Physics*, and have submitted the paper for publication to that journal. To our knowledge, ours is the first comprehensive review of diquarks, although a a prior review [2] has appeared on some aspects of diquarks and a conference has been held on the subject [3].

Our general definition of a diquark is that it is a two-quark state considered collectively. A diquark can also be regarded as a two-quark correlation in a hadron containing more than two

quarks. Dynamical diquark correlations arise in QCD (among other reasons) because of short-range spin-dependent forces between two quarks. Kinematical diquark correlations also arise from mass differences among quarks of different flavors. of a spin-1 pair. Taking into account these correlations affects the calculated values of many observables in hadron physics.

Our review includes the effects of diquarks on the static properties of hadrons, on deep-inelastic scattering, on nuclei and nuclear matter, on electron-positron annihilation, on hadron-hadron collisions, and on exclusive processes.

We have written this review, not only because we believe it is needed, but also to familiarize ourselves with the present state of the art so as to do further research on the role of diquark correlations in hadrons.

References

1. M. Anselmino, S. Ekelin, S. Fredriksson, D.B. Lichtenberg, and E. Predazzi, DFTT 7/92, IUHET-215, TULEA 1992:05 (Feb, 1992).
2. M. Szczekowski, Int. J. Mod. Phys. A 4, 3985 (1989).
3. Workshop on Diquarks, Turin, Italy, October, 1988, eds. M. Anselmino and E. Predazzi (World Scientific, Singapore, 1989).

DYNAMICAL SUPERSYMMETRY OF MESONS AND BARYONS

(Lichtenberg)

We have continued to work on the question of the consequences of an approximate dynamical supersymmetry between mesons and baryons. This supersymmetry arises in QCD because an antiquark in a meson and a diquark in a baryon each belongs to the same color multiplet ($\mathbf{\bar{3}}$) and each interacts with a quark to form a color singlet.

This idea has been considered previously [1,2], but only for the case of light quarks, where the supersymmetry is so badly broken that it is not useful. We were the first to extend the idea to the case of hadrons containing at least one heavy quark [3,4], where the situation is more promising.

References

1. H. Miyazawa, Prog. Theor. Phys. 36, 1266 (1966); 1968 Phys. Rev. 170, 1586 (1968).
2. S. Catto and F. Gürsey, 1985 Nuovo Cimento 86, 201 (1988); 99, 685 (1988).
3. D.B. Lichtenberg, Proc Rice Meeting, DPF90, B. Bonner and H. Miettinen, eds., World Scientific, Singapore (1990), p. 643.
4. D.B. Lichtenberg, J. Phys. G **16**, 1599 (1990).

GLUEBALLS, HYBRIDS, AND EXOTIC MESONS

(Lichtenberg)

The existence of glueballs follows very plausibly from QCD, but so far it has not been possible to make reliable predictions of the masses and decay modes of these states. Furthermore, glueballs should not be considered in isolation because in principle they can mix with ordinary quark-antiquark states, exotics, and hybrids.

We are studying these problems using phenomenological models in the hope that we can make a significant contribution in this difficult but important area. It is especially appropriate to do this theoretical research at Indiana University because the experimental group at Indiana University (Task A of this grant) is participating in an experiment at Brookhaven National Lab to search for glueballs and hybrid mesons [1]. Alex Dzierba of Task A is co-spokesman for this experiment.

References

1. S.U. Cheng et al., Search for mesons with unusual quantum numbers (E852), proposal (1989).

LECTURE NOTES ON THE STANDARD MODEL

(Lichtenberg)

We have written up a set of notes on the standard model based on lectures given at Shanxi University (Taiyuan, China) and at the University of Naples (Naples, Italy). The notes were published as a book in July, 1991, by Bibliopolis publishing company, Naples, in its series, *Lectures at the University of Naples* [1].

The book contains a description of the elementary particles of the standard model, an introduction to ideas in group theory and gauge theory, including spontaneous symmetry breaking, a discussion of the Lagrangian of the standard model, and selected applications.

A unique feature of the book is its emphasis on bound-state problems which cannot be solved with perturbative QCD. We have discussed nonperturbative methods, including our own research on potential models which incorporate some important features of lattice QCD.

References

1. D.B. Lichtenberg, *The Standard Model of Elementary Particles*, Bibliopolis, Naples (1991), xiv + 152 pp.

SPHALERON TRANSITION RATE: DETERMINATION FROM A CANONICAL ENSEMBLE

(A. Krasnitz, R. Potting)

A possibility that the observed baryon asymmetry of the universe may be attributed to the electroweak phase transition has attracted much attention recently¹ (it was also suggested that baryon-number violating processes may be observed in the SSC range of energies). It turns out that for temperatures exceeding the gauge-boson mass scale the baryon number evolves approximately according to the classical equations of motion. Numerical determination of the evolution rate presents computational physics with a new challenge. To date, the only approach to the problem has consisted of integrating the equations of motion with a finite time step and finding the autocorrelation time for a quantity of interest directly from its time history²⁻⁵. This dependency between the way the ensemble is generated and the measurement is inconvenient and makes generalization to the canonical case difficult and time-consuming.

We are currently working on extracting the rate from a canonical ensemble. The latter may be generated in the most efficient way available and independently of the measurement. We chose to apply the new method first to determine the kink-antikink pair creation rate in 1+1-dimensional ϕ^4 theory (a popular test case). Essentially, the evolution of the kink-antikink pair number is modelled by an effective stochastic equation of motion. So far we have been able to show that the equilibrium distribution of the kink-antikink number agrees well with the stationary solution of the corresponding Fokker-Planck equation.

If this exploratory study is met with success, the method can be applied to study the sphaleron transitions in the abelian Higgs model in 1+1 dimensions, and, finally, to determine the baryon number violation rate in the physically interesting case of the $SU(2)$ gauge-Higgs system in 3+1 dimensions.

References

1. E. Mottola, S. Raby, and G. Strikman, in *Baryon number violation at the SSC*, M. Mattis and E. Mottola (eds.), World Scientific (1990) 188; M. Dine, O. Lechtenfeld, B. Sakita, W. Fischler, and J. Polchinski, Nucl. Phys. **B342** (1990) 381; L. McLerran, Phys. Rev. Lett. **62** (1989) 1075.
2. J. Ambjoern, T. Askgaard, H. Porter, M.E. Shaposhnikov, Phys. Lett. **B244** (1990) 479; Nucl. Phys. **B353** (1991) 346.
3. D.Yu. Grigoriev, V.A. Rubakov, M.E. Shaposhnikov, Nucl. Phys. **B326** (1989) 737.
4. D.Yu. Grigoriev, V.A. Rubakov, Nucl. Phys. **B299** (1988) 67.
5. A.I. Bochkarev, Ph. de Forcrand, Phys. Rev. **D44** (1991) 519.

PERSONNEL

Principal Investigators

S. Gottlieb, professor
A. W. Hendry, professor
V. A. Kostelecký, professor
D. B. Lichtenberg, professor

Postdoctoral research associates

A. Krasnitz
R. Potting

Graduate Students

S. Mustafa (on leave since Dec. 31, 1990)
J. Tsai (PhD, August, 1991) with Hendry.

SHORT-TERM VISITORS

During the last year, we have had several short-term visitors who consulted on topics of interest to our research effort. The cost of some of these visitors was supported in whole or in part by the contract. Additional support came from Indiana University and other sources. These visitors and their home institutions are listed below, together with the approximate dates of their visits.

C. S. Kim, University of Wisconsin (Sep. 16, 1991)
Malcolm Perry, University of Cambridge (Sep. 23, 1991)
Vladimir Eletsky, Institute for Theoretical and Experimental Physics, Moscow (Sep. 30, 1991)
W. Zakrzewski, University of Durham, UK (Oct. 7, 1991)
Jochen Fingberg, HLRZ, KFA Jülich (Dec. 9, 1991)
James Pantaleone, Fermilab (Jan. 13, 1992)
Al Shapere, Cornell (April 13, 1992)
Roberto Vega, SLAC (April 20, 1992)
Ikaros Bigi, Notre Dame (April 27, 1992)

INVITED TALKS AT CONFERENCES

1989-92

Finite Size Scaling and the QCD High Temperature Phase Transition, S. Gottlieb, International Symposium Lattice 89, Capri, Italy, September 18, 1989.

Gravity and Lorentz Breakdown in Higher-Dimensional Theories and Strings, A. Kostecký, XVIIIth International Conference on Differential Geometric Methods in Theoretical Physics, Granlibakken, California, July 7, 1989.

Atomic Supersymmetry, A. Kostecký, 10th Canadian Theoretical Chemistry Conference, Banff, Canada, August 26, 1989.

A Nonperturbative Vacuum for the Open Bosonic String, A. Kostecký, Conference on Superstrings and Particle Theory, Tuscaloosa, Alabama, November 12, 1989.

QCD with Dynamical Staggered Fermions, S. Gottlieb, Division of Particles and Fields, Houston, TX, January 5, 1990.

Thermodynamics, S. Gottlieb, QCD Teraflop Workshop, SCRI, Florida State University, January 17, 1990.

Quantum Chromodynamics at $6/g^2 = 5.6$, S. Gottlieb, PANIC XII (International Conference on Particles and Nuclei), Massachusetts Institute of Technology, Cambridge, MA, June 28, 1990.

Finite Temperature QCD with Dynamical Fermions, S. Gottlieb, Lattice '90, SCRI, Florida State University, October 11, 1990.

A Nonperturbative String Vacuum, A. Kostecký, 1990 Annual Meeting of the Division of Particles and Fields of the American Physical Society, Houston, Texas, January 4, 1990.

Collective Effects in Bosonic Strings, A. Kostecký, Strings '90 Conference, College Station, Texas, March 16, 1990.

Gauge Masses in String Field Theory, A. Kostecký, Conference Beyond the Standard Model II/Supersymposium, Norman, Oklahoma, November 3, 1990.

Flavor and Higher Symmetries in Hadrons Containing both Heavy and Light Quarks, D.B. Lichtenberg, Rice Meeting, DPF90 (Jan. 1990)

Effect of Color-Hyperfine Interactions on Baryon Masses, D.B. Lichtenberg (with M. Anselmino and E. Predazzi), 25th Int.Conf. on High Energy Physics, Singapore (Aug. 2-8, 1990)

Lattice QCD—How We Expect it to Develop in the Future, S. Gottlieb, LISS workshop, IUCF, February 9, 1991.

QCD on the iPSC/860, S. Gottlieb, Workshop on Fermion Algorithms, Juelich, April 10-12, 1991.

What Lattice Calculations Can Tell Us About QCD at High Temperature, S. Gottlieb, QCD in Nuclear Physics, Gordon Conference, July 22-26, 1991.

Recent Finite Temperature Lattice QCD Simulations with Wilson and Staggered Fermions, S. Gottlieb, Brookhaven Workshop, August 1991.

Gauge Boson Masses in String Field Theories, A. Kostelecký, Meeting of the Southern Association of High Energy Theorists, Gulf Shores, Alabama, January 4, 1991.

Supercoherent States for Physical Systems, A. Kostelecký, Conference on Squeezed States and Uncertainty Relations, College Park, Maryland, March 29, 1991.

String Signatures, A. Kostelecký, Joint International Lepton-Photon Symposium/Europhysics Conference in High Energy Physics, Geneva, Switzerland, July 25, 1991.

Neutrino Mass Bounds, A. Kostelecký, Meeting of the Southern Association of High Energy Theorists, Gulf Shores, Alabama, January 3, 1992.

Parabolic Atomic Supersymmetry, A. Kostelecký, Conference on Harmonic Oscillators, College Park, Maryland, March 1992.

Dynamical Supersymmetry for Mesons and Baryons, D.B. Lichtenberg, to be presented at the International Workshop on Quark Cluster Dynamics, Bad Honnef, Germany, June 29-July 1, 1992.

Diquarks in Exotic Hadrons, D.B. Lichtenberg, to be presented at the International Workshop on Diquarks, Turin, Italy, Nov. 2-4, 1992.

During 1989-1992, Gottlieb presented 12 colloquia and seminars at universities, laboratories, computer companies and conferences; Hendry presented 6 colloquia and seminar at universities and laboratories, Kostelecký presented 30, and Lichtenberg presented 27.

RESEARCH PAPERS SUPPORTED IN PART BY DOE
 CONTRACT DE-AC02-84ER40125 and GRANT DE-FG02-91ER40661, TASK B
 1989-92

Since 1970, we have written more than 300 papers supported in part by DOE (including prior contracts with AEC and ERDA). We include below only a list of papers for the period January, 1989–March, 1992.

1. CP Violation in Supergravity Models (V.A. Kostelecký, G.C. Branco), *Phys. Rev. D* **39**, 2075 (1989).
2. Baker–Campbell–Hausdorff Relations for the Orthosymplectic Group $Osp(1|2)$ (A. Kostelecký, B.W. Fatyga and D.R. Truax), *J. Math. Phys.* **30**, 291 (1989).
3. QCD Simulations with Dynamical Fermions, (S. Gottlieb, W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), in *Proceedings of the Storrs Meeting*, edited by K. Haller *et al.* (World Scientific, Singapore, 1989).
4. More on the Spectrum with Kogut-Susskind Fermions, (S. Gottlieb, W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), (with W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), presented at Lattice '88, Fermilab, September, 1988, *Nucl. Phys. B(Proc. Suppl.)***9**, 259, 1989.
5. Phase Diagram of Four Flavor QCD, (S. Gottlieb, W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), presented at Lattice '88, Fermilab, September, 1988, *Nucl. Phys. B(Proc. Suppl.)***9**, 325, 1989.
6. Hadronic Screening Lengths and Quark Number Susceptibility from Lattice QCD, (S. Gottlieb, W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), presented at Quark Matter '88, Lenox, Massachusetts, September 1988, *Nucl. Phys. A***498**, 435c, 1989.
7. Reply to Comment on 'Fine Structure and Analytical Quantum-Defect Wavefunctions' (V.A. Kostelecký, M.M. Nieto, D.R. Truax), *Phys. Rev. A* **41**, 1732 (1989).
8. Spontaneous Breaking of Lorentz Symmetry in String Theory, (V.A. Kostelecký, S. Samuel), *Phys. Rev. D* **39**, 683 (1989).
9. Toy Superstrings, (V.A. Kostelecký, M. Corvi and P. Moxhay), *Phys. Rev. D* **39**, 1611 (1989).
10. Paton-Chan Factors in the $N=2$ Open Superstring (M. Corvi) *Phys. Lett. B*, **231**, 240 (1989).
11. The Tachyon Potential in String Theory (with S. Samuel), in K. Haller *et al.*, eds., *Proceedings of the Storrs Meeting*, World Scientific, Singapore, 1989.
12. Grassmann-Valued Fluid Dynamics, (V.A. Kostelecký, B.W. Fatyga and D.R. Truax), *J. Math. Phys.* **30**, 1464 (1989).
13. Phenomenological Constraints on Strings and Higher-Dimensional Theories, (V. A. Kostelecký, S. Samuel), *Phys. Rev. Lett.* **63**, 224 (1989).
14. Gravitational Phenomenology of Higher-Dimensional Theories and Strings, (V.A. Kostelecký, S. Samuel), *Phys. Rev. D* **40**, 1886 (1989).
15. Testing Static Quark-Antiquark Potentials with Bottomonium (D.B. Lichtenberg, E. Predazzi, R. Roncaglia, M. Rosso, and J.G. Wills), *Z. Phys. C* **41**, 615 (1989).

16. Why it is Necessary to Consider Diquarks (D.B. Lichtenberg), Proc. Workshop on Diquarks, Turin, Italy (Oct. 24–26, 1988), M. Anselmino and E. Predazzi, eds. World Scientific, Singapore (1989).
17. Effects of Diquarks and Mesons on Baryon Magnetic Moments (D.B. Lichtenberg and W. Namgung), Phys. Rev. D **40**, 1606 (1989).
18. Lower Bounds on the Constituent Quark Mass Differences (D.B. Lichtenberg), Phys. Rev. D **40**, 3675 (1989).
19. Application of a Generalized Feynman–Hellmann Theorem to Bound-State Energy Levels (D.B. Lichtenberg), Phys. Rev. D **40**, 4196 (1989).
20. Flavor Dependence of the Static Quark-Antiquark Potential (D.B. Lichtenberg, E. Predazzi, R. Roncaglia, M. Rosso, and J.G. Wills), Z. Phys. C **46**, 75 (1990).
21. Masses of $c\bar{s}$, $b\bar{s}$, and $c\bar{b}$ States in Potential Models, (D.B. Lichtenberg, R. Roncaglia, J.G. Wills, and E. Predazzi), Z. Phys. C **47**, 83 (1990).
22. Gravity and Lorentz Breakdown in Higher-Dimensional Theories and Strings, (V.A. Kostelecký, S. Samuel), *Proceedings of the XVIIIth Conference on Differential Geometric Methods in Theoretical Physics*, Plenum, New York, 1991.
23. On a Nonperturbative Vacuum for the Open Bosonic String (V.A. Kostelecký, S. Samuel), Nucl. Phys. **B336**, 263 (1990).
24. Nonperturbative Vacuum Structure and the Open Bosonic String (V.A. Kostelecký, S. Samuel), in L. Clavelli and B. Harms, eds., *Superstrings and Particle Theory*, World Scientific, Singapore, 1990.
25. A Nonperturbative String Vacuum (V.A. Kostelecký, S. Samuel), accepted for publication in B. Bonner and H. Miettinen, eds., *Proceedings of the Rice Meeting*, World Scientific, Singapore, 1990.
26. Nonperturbative Many-Body Physics for the Open Bosonic String (V.A. Kostelecký, S. Samuel), Phys. Rev. Lett. **64**, 2238 (1990).
27. Collective Effects in Bosonic Strings (V.A. Kostelecký, S. Samuel), in R. Arnowitt et al., eds., *Strings '90*, World Scientific, Singapore, 1991.
28. Collective Effects in the Closed Bosonic String (V.A. Kostelecký, S. Samuel), Phys. Rev. D **42**, 1289 (1990).
29. A Quantum-Mechanical Central Limit Theorem (M. Corvi, A. Lenard), IUHET 163, (June 1989).
30. The Off-Shell Koba-Nielsen Formula (R. Bluhm, S. Samuel), Nucl. Phys. **B323**, 337 (1989).
31. Off-Shell Conformal Field Theory at the One-Loop Level (R. Bluhm, S. Samuel), Nucl. Phys. **B325**, 325 (1989).
32. On the Modified Superstring Field Theory (R. Bluhm, S. Samuel), Nuclear Physics **B338**, 38 (1990).
33. Conformal Methods in String Field Theory Amplitudes (R. Bluhm, S. Samuel), in L. Clavelli et al., eds., *Superstrings and Particle Theory*, World Scientific, Singapore, 1990.

34. Four Flavor QCD with Intermediate and Light Mass Quarks, (S. Gottlieb, W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), Phys. Rev. D **40**, 2389 (1989).
35. QCD Thermodynamics with Eight Time Slices, (with W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), Phys. Rev. D **41**, 622, 1990.
36. Finite Size Scaling and the QCD High Temperature Phase Transition, (with W. Liu, R.L. Renken, R.L. Sugar and D. Toussaint), Nucl. Phys. B(Proc. Suppl.)**17**, 173, 1990.
37. Hadron Spectrum with Staggered Dynamical Quarks, (with K. M. Bitar, A. D. Kennedy, W. Liu, T. A. DeGrand, J. B. Kogut, R.L. Renken, M.C. Ogilvie, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper and D. Toussaint), Nucl. Phys. B(Proc. Suppl.)**17**, 404, 1990.
38. Hadronic Spectroscopy with Wilson Valence Quarks, (with K. M. Bitar, A. D. Kennedy, W. Liu, T. A. DeGrand, J.B. Kogut, R.L. Renken, M.C. Ogilvie, P. Rossi, D.K. Sinclair, K.C. Wang, R.L. Sugar, M. Teper and D. Toussaint), Nucl. Phys. B(Proc. Suppl.)**17**, 400, 1990.
39. QCD with Dynamical Staggered Quarks, (with K. M. Bitar, T. A. DeGrand, R. Edwards, U. Heller, A. D. Kennedy, J. B. Kogut, A. Krasnitz, C. Liu, M. C. Ogilvie, R. L. Renken, P. Rossi, D. K. Sinclair, R. L. Sugar, M. Teper, D. Toussaint and K. C. Wang), in *Proceedings of The Rice Meeting*, edited by B. Bonner and H. Miettinen (World Scientific, Singapore, 1990).
40. Lattice Gauge Theory on the Intel Parallel Scientific Computer, in *Computing for High Luminosity and High Intensity Facilities*, edited by J. Lillberg and M. Oothoudt (American Institute of Physics, New York, 1990).
41. On a Fermion Correlation Function in Multiple Field Configurations (A. Krasnitz), Phys. Rev. Letts. **68**, (1992) 2125.
42. Block-conjugate-Gradient Method (J.F. McCarthy), Phys. Rev. D **40**, 2149 (1989).
43. Determination of the Order of Phase Transitions in Numerical Simulations (J.F. McCarthy), Phys. Rev. D **41**, 9530 (1990).
44. Flavor and Higher Symmetries in Hadrons Containing both Heavy and Light Quarks (D.B. Lichtenberg) Proc. Rice Meeting, DPF90, Billy Bonner and Hannu Miettinen, eds. World Scientific, Singapore, 1990, pp. 643-647.
45. Nonstatic Spin-Independent Term in the Quarkonium Potential (J.G. Wills and D.B. Lichtenberg) Phys. Rev. D **42**, 2408-11 (1990).
46. Semiempirical Formulas for the Color-Hyperfine Mass Splitting in hadrons (Yong Wang and D.B. Lichtenberg) Phys. Rev. D **42**, 2404-7 (1990).
47. Dynamical Supersymmetry in Hadrons Containing Both Heavy and Light Quarks (D.B. Lichtenberg) J. Phys. G **16**, 1599-06 (1990).
48. Quark Color-hyperfine Interactions in Baryons (M. Anselmino, D.B. Lichtenberg, and E. Predazzi) Z. Phys. C. **45**, 605-12 (1990).
49. Hadron Spectrum in QCD at $6/g^2 = 5.6$, (K. M. Bitar, R. Edwards, U. M. Heller, A. D. Kennedy, W. Liu, T. A. DeGrand, S. Gottlieb, A. Krasnitz, J. B. Kogut, R. L. Renken, M. C. Ogilvie, P. Rossi, D.K. Sinclair, K.C. Wang, R.L. Sugar, M. Teper and D. Toussaint), Phys. Rev. D **42**, 3794, 1990.

50. The High Energy Monte Carlo Grand Challenge: Simulating Quarks and Gluons, (K. M. Bitar, A. D. Kennedy, W. Liu, T. A. DeGrand, S. A. Gottlieb, U. Heller, A. Krasnitz, J. B. Kogut, R.L. Renken, M.C. Ogilvie, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper and D. Toussaint), *Internat. J. Supercomput. Appl.* **4**, 48, 1990.
51. Quantum Chromodynamics at $6/g^2 = 5.6$, (K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U. M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), *Phys. Rev. Lett.* **65**, 2106, 1990.
52. Lattice Spectrum with Dynamical Staggered Quarks at $6/g^2 = 5.6$, (K.M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U.M. Heller, A.D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), presented at PANIC XII, Particles and Nuclei, Proceedings of the Twelfth International Conference on Particle and Nuclei, Massachusetts Institute of Technology, June 24–29, 1990, edited by J. L. Matthews *et al.*, page 527c.
53. Hadron Thermodynamics with Wilson Quarks, (K.M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U.M. Heller, A.D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), *Phys. Rev. D* **43**, 2396, 1991.
54. Finite Temperature QCD with Dynamical Fermions, S. Gottlieb, *Nucl. Phys. B(Proc. Suppl.)* **20**, 247, 1991.
55. Hadronic Spectroscopy at $a \approx 0.15$ fm, (K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U. M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), *Nucl. Phys. B(Proc. Suppl.)* **20**, 362, 1991.
56. QCD Thermodynamics with Wilson Quarks, (K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U. M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), *Nucl. Phys. B(Proc. Suppl.)* **20**, 272, 1991.
57. Glueball and Topological Charge in the Presence of Dynamical Quarks, (K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U. M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), *Nucl. Phys. B(Proc. Suppl.)* **20**, 390, 1991.
58. On Glueballs and Topology in Lattice QCD with Two Light Flavors, (K. M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U. M. Heller, A. D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, P. Rossi, D.K. Sinclair, R.L. Sugar, M. Teper, D. Toussaint, K.C. Wang), *Phys. Rev. D* **44**, 2090, 1991.
59. Studying Quarks and Gluons on MIMD Parallel Computers, (C. Bernard, S. Gottlieb, M. C. Ogilvie, T. A. DeGrand, C. DeTar, A. Krasnitz, R. L. Sugar and D. Toussaint), *Internat. J. Supercomput. Appl.* **5**, 61, 1991.

60. Hadron Thermodynamics on the Connection Machine, (S. Gottlieb, A. Krasnitz, U. M. Heller, A. D. Kennedy, W. Liu, J. B. Kogut, R. L. Renken, D. K. Sinclair, K. C. Wang, R. L. Sugar and D. Toussaint), *Internat. J. Supercomput. Appl.* **5**, 50, 1991.
61. QCD on the iPSC/860, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), in *Workshop on Fermion Algorithms*, edited by H. J. Hermann and F. Karsch, (World Scientific, Singapore, 1991).
62. QCD Thermodynamics with Two Flavors at $N_t = 6$, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), IUHET-210 (October 1991), submitted to *Physical Review*.
63. Finite Temperature Lattice QCD Simulations with Staggered and Wilson Fermions, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, U. M. Heller, A. D. Kennedy, J. B. Kogut, A. Krasnitz, W. Liu, M. C. Ogilvie, R. L. Renken, D. K. Sinclair, R. L. Sugar, D. Toussaint, and K. C. Wang), in *Proceedings of Hot Summer Drive*, a workshop at Brookhaven National Laboratory, August, 1991, to be published by World Scientific.
64. The Spatial Structure of Screening Propagators in Hot QCD, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), IUHET-211 (November 1991), submitted to *Physical Review Letters*.
65. Quark Structure of Static Correlators in High Temperature QCD, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), in *Proceedings of Quark Matter '91*, to be published.
66. Neutrinos in a Phenomenological Solar Model (A. Hendry, J. Tsai), IUHET-203 (1991).
67. Reply to Comment on 'Fine Structure and Analytical Quantum-Defect Wavefunctions' (A. Kostelecký, M.M. Nieto and D.R. Truax), *Phys. Rev. A* **41**, 1732 (1990).
68. Reply to Comment on 'Analytical Wavefunctions for Atomic Quantum-Defect Theory' (A. Kostelecký and M.M. Nieto), *Phys. Rev. A* **43**, 4063 (1991).
69. Supercoherent States (A. Kostelecký, B.W. Fatyga, M.M. Nieto and D.R. Truax), *Phys. Rev. D* **43**, 1403 (1991).
70. Photon and Graviton Masses in String Theories (A. Kostelecký and S. Samuel), *Phys. Rev. Lett.* **66**, 1811 (1991).
71. Gauge Masses in String Field Theory (A. Kostelecký and S. Samuel), in K. Milton et al., eds., *Beyond the Standard Model II*, World Scientific, Singapore, 1991.
72. Supercoherent States and Physical Systems, (B.W. Fatyga, A. Kostelecký, M.M. Nieto and D.R. Truax), in D. Han et al., eds., *Squeezed States and Uncertainty Relations*, NASA, Washington, D.C., 1992.
73. CPT and Strings (A. Kostelecký and R. Potting), *Nucl. Phys.* **B359**, 545 (1991).
74. Experimental Constraints on Extra Dimensions (A. Kostelecký and S. Samuel), *Phys. Lett.* **B270**, 21 (1991).
75. Hidden Weak Ferromagnetism in Slightly Doped Antiferromagnets, (A. Krasnitz and A. Kovner), *Phys. Rev.* **B41** (1990) 547.

76. Fermion Correlation Function in Multiple Field Configurations, (A. Krasnitz), Phys. Rev. **D42** (1990) 1301.
77. Spectroscopy with Multiple Field Configurations, (A. Krasnitz), Phys. Rev. **D43** (1991).
78. Irreducible Representations of Lie groups from the quantization of constrained dynamical systems, (E. Gates, R. Pottling, C. Taylor, B. Velikson), IUHET-195 (1991).
79. The Standard Model of Elementary Particles (D.B. Lichtenberg), Bibliopolis, Naples (1991).
80. Effect of Color-Hyperfine Interactions on Baryon Masses (D.B. Lichtenberg, M. Anselmino, and E. Predazzi), *Proc. 25th Int. Conf. on High Energy Physics*, Singapore, Aug. 2-8, 1990, K.K. Phua and Y. Yamaguchi, eds., World Scientific, Singapore (1991), pp. 577-81.
81. Sign of the Color-Hyperfine Splitting in Charmonium as a Test of Perturbative QCD (D.B. Lichtenberg, E. Predazzi, and R. Roncaglia), Phys. Rev. D **45**, 3265 (1992).
82. Color-Hyperfine Splitting in Heavy Quarkonium P States from Perturbative QCD with a Running Coupling Strength (D.B. Lichtenberg and R. Potting), IUHET 208 (Revised Mar. 1992)
83. Diquarks (M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson, and D.B. Lichtenberg), IUHET 215 (Feb. 1992)
84. Null Experiments for Neutrino Masses (A. Kostelecký, A. Chodos, R. Potting, and E. Gates), Mod. Phys. Lett. A **7**, 467 (1992).
85. Space Time CPT and String Theory (A. Kostelecký and R. Potting), to appear in *Proceedings of the XXth Conference on Differential Geometric Methods in Theoretical Physics*.
86. String Signatures (A. Kostelecký and S. Samuel), to appear in *Proceedings of the 1991 Joint International Lepton-Photon Symposium and Geophysics Conference on High-Energy Physics*.
87. Neutrinos in a Phenomenological Solar Model (A. Hendry and J. Tsai). IUHET 203 (1991).
88. A Polytropic Solar Model (A. Hendry), IUHET 214 (1992).
89. QCD Thermodynamics with Two Flavors of Quarks, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), in Proceedings of Lattice '91, Tsukuba, Japan, Nucl. Phys. B(Proc. Suppl.) to appear.
90. Finite-Size Effects and T_c for $\beta = 5.445$ with Two Staggered Flavors, (C. Bernard, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, M. C. Ogilvie, R. L. Sugar and D. Toussaint), in Proceedings of Lattice '91, Tsukuba, Japan, Nucl. Phys. B(Proc. Suppl.) to appear.
91. Thermodynamics of Lattice QCD with 2 Light Dynamical (Staggered) Quark Flavours on a $16^3 \times 8$ Lattice, (S. Gottlieb, U. M. Heller, A. D. Kennedy, J. B. Kogut, A. Krasnitz, W. Liu, R. L. Renken, D. K. Sinclair, R. L. Sugar, D. Toussaint and K. C. Wang), in Proceedings of Lattice '91, Tsukuba, Japan, Nucl. Phys. B(Proc. Suppl.) to appear.
92. QCD Hadron Spectroscopy with Staggered Dynamical Quarks at $\beta = 5.6$, (K.M. Bitar, T. DeGrand, R. Edwards, S. Gottlieb, U.M. Heller, A.D. Kennedy, J.B. Kogut, A. Krasnitz, W. Liu, M.C. Ogilvie, R.L. Renken, D.K. Sinclair, R.L. Sugar, D. Toussaint, K.C. Wang), Nucl. Phys. B(Proc. Suppl.) to appear.
93. Physics Goals of the QCD Teraflop Project, S. Gottlieb with the QCD Teraflop Collaboration), Intl. J. Mod. Phys. C, **2**, 829, 1991.

END

**DATE
FILMED**

7 / 9 / 92