NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

DE-FG03-91ER61198 DOE/ER/
A001

DOE/ER/61198--T1

DE92 013772

**Arm Tropical Pacific Experiment (ATPEX) : Role of Cloud, Water Vapor and
Convection Feedbacks in the Coupled Ocean/Atmosphere System**

V. Ramanathan & T.P. Barnett

California Space Institute & Climate Research Division

Scripps Institution of Oceanography, UCSD, La Jolla, CA 92093

Project Funded by Department of Energy, Office of Energy Research

Progress Report for the Period: September 1, 1991 - August 31, 1992

Report Date: March 5, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

<u>Table of Contents</u>	<u>Page Number</u>
1. Research Progress	1
(a) First Six Months	1
(b) Planned Studies for the Next Six Months	4
2. Papers Submitted	5
3. Personnel & Status of Appointments	5
4. Equipment Purchases	6

MASTER

1. RESEARCH PROGRESS

(a) First Six Months:

During the first six months, we have created most of the infrastructure needed to integrate the ARM data with radiation models, AVHRR/EPBE satellite data, and GCMs. We have initiated studies that include radiation model validation, improved treatment of the three-dimensional structure of cloud-radiation interactions, and GCM sensitivity runs that will unravel the role of cloud-convection-radiation interactions in the Pacific SSTs and the overlying Walker and Hadley circulation.

As mentioned in the proposal (see p.10), the research program is divided into three phases. We are in Phase 1 which extends from Year 1 to Year 2. The specific scientific issues are subdivided into three categories (p.10): (i) Radiation; (ii) Cloud Parameterization Issues; (iii) Feedback and Ocean-Atmosphere Interactions. We will list the progress under each of these three categories.

(i) Radiation (J. del Corral, A. Inamdar & V.Ramanathan):

The LOWTRAN 7 radiation code has been integrated with radiosonde soundings taken from ships over the tropical Pacific, Indian, and Atlantic oceans. A five-year data set from 1984 to 1989 has been assembled with a total of about 1500 soundings. This period was chosen because of the availability of the ERBE data set. We have computed the surface and top-of-the atmosphere fluxes, as well as flux divergences within the troposphere.

Role of Convection in the Atmospheric Greenhouse Effect: The computed atmospheric greenhouse effect (G_a) from the soundings is in very good agreement with the ERBE data (Figure 1). In particular, the computed greenhouse effect simulates the super-greenhouse effect identified from ERBE data by Ramanathan & Collins (1991, Nature, 351, p.27). Encouraged by this agreement, we are now analyzing the surface radiation flux and its relationship with sea surface temperature (SST) and the greenhouse effect. In particular, we are interested in the dependence of these processes on deep convection. Since one necessary condition for deep convection (Graham & Barnett, 1987, Science, 238, p.657) is that SST should exceed about 300 K, we separate the computed fluxes into two categories: (i) SST > 300 K to denote regions of convection; (ii) SST < 298 K to denote regions of no-convection. The averages of the greenhouse effect and the surface fluxes for the two regions (Figure 2) reveal several interesting features. The greenhouse effect and the downward emission are much larger in the convection regions. The entire troposphere contributes to the increase in G, but roughly 70% of the increase comes from the increase in water vapor and lapse rate from surface to about 500 mb. It is also

seen that water vapor increases rapidly in the boundary layer (1000 to 700 mb) up to a SST of about 300 K. For SST > 300 K, the boundary layer humidity is subject to considerable scatter and does not reveal a discernible trend with SST. For SST > 300 K, the increase in vapor concentration is found mainly above the boundary layer.

The above results set the stage very nicely for our studies with the ARM site data over the western tropical Pacific. Since the radiosonde data are unreliable above pressure levels of about 500 mb, we are very keen on examining the link between convection, upper tropospheric water vapor and the greenhouse effect.

A paper on these results has been submitted for presentation at the AGU Spring 1992 Meeting.

(ii) Cloud Parameterization Issues (I. Boer, JDC, & VR):

Three-Dimensional Imaging of Clouds: We have made substantial progress in the development of this tool. As proposed (p.13 of the proposal), we have incorporated the following features in a three-dimensional cloud imager, which employs the full resolution (1 km x 1 km) AVHRR data :

- Clear-sky Identification: As a first guess, we employ the Coakley/Bretherton spatial coherence scheme on the IR window channel. We then examine the ratio of the radiances of the two visible channels. One of these channels is in the blue end of the spectrum (Channel 1), while the other is near the red end (Channel 2). Since the clear sky is blue and cloudy sky is white, the ratio of these two channels should be much larger for the clear-sky than for the cloudy sky. We use this property to filter out the clear scenes from the pixels chosen by the spatial coherence method.

- Optical Depth Determination: We then employ a delta-eddington multiple scattering code to solve for the optical depth of cloudy pixels, assuming that they fill the field of view.

- Edge Detection: A Detect and Spread (DAS) method has been developed to label each cloud element within the scene. DAS first selects a pixel with large optical depth, and then spreads this cloud all around until it encounters a clear scene. An optically thin pixel attached to a thick cloud is assumed to have the same altitude as the thick cloud.

Many details of this cloud retrieval and imaging algorithms are currently being documented in a report. An image of a cloud system in the tropical Pacific, as reconstructed by the cloud imager is attached herewith (Figure 3). The imager reveals realistic structure of deep convective clouds and

anvils. However, many serious problems remain which can only be resolved with in-situ data from the ARM sites.

(iii) Feedback and Ocean-Atmosphere Interactions: (T.P. Barnett, M. Meyer, U. Schlese (MPI), JDC, & VR):

Atmosphere GCM :

The Max-Planck Institute has given their latest version of the GCM, the T42 ECHAM3 version, for our studies. MM & TPB have started a 14 year integration with observed SSTs for the period 1970-83; the ten years before AMIP, plus four overlap years. Lack of computer time will slow the progress of this run. This eventually will give us 20 years of T42 results, enough to do a decent statistical comparison with observations.

Greenhouse Effect and Cloud Forcing Results: Tom Peterson (a visitor from CSU) and TPB compared a reduced resolution version of ECHAM3 simulated cloud and radiation variability over 1979-85 with satellite data. The model did remarkably well, and a paper describing the results has been submitted. MM, JDC, and VR have begun to compare the T42 model run for the 1980s with observed ERBE data. We are focusing on the tropical Pacific oceans. The model is able to capture the gross features of the observed data. For example, it simulates the super-greenhouse effect over the convective regions and captures the near cancellation of the longwave and shortwave cloud forcing. However, the extreme values of the cloud-radiative forcing parameters are much larger in the model.

Radiative Effects of the 1987 El-Nino (MM, JDC & VR): A more critical test of the model is the effect of the El-Nino on the radiative fluxes as identified in the Ramanathan & Collins (1991) paper. The model is quite successful in simulating the increases in the greenhouse effect, the longwave and shortwave cloud forcing over the warmer central Pacific. The monthly mean differences shown in Figure 4 shows a remarkable similarity with the observed changes (compare Figures 4a & 4b). The spatial distribution of radiation and greenhouse effects is also surprisingly well reproduced (compare Figures 5a & 5b). However, the magnitude of the cloud forcing changes are larger in the model (note the scale difference of the extreme in Figure 5). What is the significance of this agreement? First, note that the warming in the central Pacific is accompanied by deep convection, which in turn modifies the water vapor distribution and cirrus clouds. Since the altered water vapor distribution changes the greenhouse effect, the agreement with the observed changes in the greenhouse effect and cloud forcing suggests that the model may have a realistic treatment of the coupling between convection, water vapor distribution, and cloudiness.

This is certainly an encouraging beginning and comparison with the ARM surface data is the next step.

Cloud-Radiation Experiments (MM, VR, US): Motivated by the success of the model in simulating the El-Nino changes, we have begun cloud-radiation sensitivity experiments. A preliminary 60-day run has been completed, in which we are examining the importance of cirrus-radiative forcing to the atmospheric dynamics and convection. By the end of Year-1, we would have completed several sensitivity studies, which will improve our understanding of the feedback effects of western Pacific cirrus clouds on the Walker and Hadley cell circulation.

Tropical Ocean GCM (TPB, A. Miller, J. Ritchie, & N. Schneider (from May 1)): Jack Ritchie has spun up a new version of the OPYC (MPI OGCM), and is now testing it to see how well it did the seasonal cycle. Results to data are moderately encouraging, although the model does have some flaws. In addition, a 16-year run with observed winds has just been completed. We are just starting to study this run to see how well OPYC simulates natural variability. First indications are that the responses are too weak, but this may be due to a stronger than necessary empirical damping, which could be easily corrected. We have received a new MPI ocean model, which is a more conventional z-coordinate model with better numerics and a somewhat improved vertical mixing scheme. We are just starting to test it as an alternative to OPYC.

(b) Planned Studies for the Second Six Months:

Further work on the infrastructure (i) John del Corral will contact the ARM data group to acquire the data that will soon be collected at the first site in Oklahoma. The data that we need include surface and satellite (AVHRR) data. We will automate the process of acquiring the data, computation of radiation flux, imaging of the three-dimensional cloud structure and validation, and tuning of the computed three-dimensional cloud and radiation fields; (ii) JDC will work with M. Meyer (GCM programmer) to develop an interactive image processing system to analyze the three-dimensional time evolution of ARM and GCM fields.

Scientific Studies: (i) We will undertake comparison studies of ARM data with model fields. (ii) Preliminary estimates will be made of the non-linear relationship between sub-grid scale cloud properties and the averaged GCM scale radiation fields. (iii) We will complete our studies on the interaction between convection, water vapor, and the greenhouse effect with the ship data. (iv) We will complete about 6 AGCM sensitivity studies in which we systematically strip the model of cirrus clouds in various regions of the tropics to understand the planetary scale waves and motion

fields forced by the cirrus radiative forcing. (v) The two ocean models will be inter-compared, and the best one retained for production work. (vi) N. Schneider will work to improve the vertical mixing process in the selected model, concentrating on performance in the warm pool. (vii) Various ENSO/non-ENSO simulations will be carried out to determine the interaction of radiation and ocean mixing processes.

2. Papers Submitted

E. Boer & V. Ramanathan, 1992: Three-dimensional imaging of cloud structure using satellite radiances: a new tool for cloud studies. Submitted to: All-Union Session on Scientific Visualization, AGU Joint Spring Meeting, Montreal, Canada, May 12th-15th.

A.K. Inamdar & V. Ramanathan, 1992: More on the water vapor greenhouse effect over the tropical oceans. AGU Joint Spring Meeting, Montreal, Canada, May 12th-15th.

T.C. Peterson, T.P. Barnett, E. Roeckner, & T.H. Vonder Haar, 1992: An analysis of the relationship between cloud anomalies and sea surface temperature anomalies in a global circulation model. Submitted to: Journal of Geophysical Research - Atmospheres.

3. Personnel & Status of Appointment

We list below the research personnel whose salaries are currently paid by this proposal:

(i) Programmers:

M. Meyer:	100%
J. del Corral:	50%
J. Ritchie:	50%

(ii) Scientists:

A. Miller	25%
A.K. Inamdar:	50%

(iii) New Appointments: We proposed to hire three new research scientists by the end of three months after getting the grant (i.e., by December 1st). Because of changes in appointment policies, there has been some delays. A post-doctoral Oceanographer, Dr. N. Schneider will join

us by May 1st to work on problems dealing with the vertical mixing parameterization in ocean models. We have selected a short list of candidates for the two atmospheric science positions, and hope to have them on-board by June 1st. In the meantime, Dr. Inamdar (a post-doc) will work on radiation model studies in the western Pacific region. E. Boer, an imaging specialist, will complete work on the three-dimensional cloud imager. We have received approval from DOE to use a portion of the unused funds for buying urgently and badly needed computing equipment, the details of which are given next.

4. Equipment Purchases

ARM & Satellite Data Studies: We have ordered a high capacity erasable optical automated removable media mass storage system. This system includes disk drives and media for 60 Gigabytes of storage, a high performance workstation front-end, and a 5 Gigabyte 8mm tape drive. The workstation (SPARC 2) will run database software to manage and retrieve the data stored on optical disks and also provide network access to the data. The data stored on this system will include the ARM data set, AVHRR, and ERBE satellite images, field measurements, and GCM model results.

The total cost of the system is \$102,780, and was cost-shared by \$35,000 from DOE funds, with the balance from ERBE and CERES grants and \$40,000 seed funds from the NSF Science and Technology Center for Clouds, Chemistry, and Climate. As soon as this system arrives (by mid-March), JDC will integrate the system into our facility for analysis of satellite data and model results (Figure 6).

GCM Studies: A disk farm system has been ordered, consisting of a server and 14 Gigabytes of on-line storage. This will accommodate the AGCM history files and sensitivity runs, as well as to provide rapid data access for real-time graphics work. We plan to mate this system with the optical jukebox system during Year 2.

SHIP DATA LOCATIONS 1985 - 89

FIGURE 1: Top panel shows the location of ship sondes used in the radiation model study. Bottom panels shows a comparison of observed atmospheric greenhouse effect with computed values for co-located regions and times. The computed values are obtained from the radiosonde soundings obtained from the ship observations.

"Non-Convective Region"
(292 - 298 K)

"Convective Region"
(> 300 K)

FIGURE 2:

Global longwave energy budgets for the non-convective and convective regimes. G is the total greenhouse trapping in the atmosphere, Q is the radiative cooling of the entire atmospheric column. $G1$, $G2$ and $G3$ in the lower half of the figure refer to the greenhouse trapping in the lower, middle and upper tropospheres respectively. The downward fluxes at the boundaries of these layers are shown to the right. All energy units are in W/m^2 .

FIGURE 3: Three-dimensional reconstruction of clouds from AVHRR image of a convective-cirrus scene in the Pacific. Cloud properties are obtained from the two visible and two IR window channels. The top and bottom images show two different 150 km x 75 km regions. The ruler on the bottom image shows the height scale. The top of the red marker is 20 km in altitude. The scene is viewed from an altitude of about 16 km for the top image and from 32 km for the bottom image. Both reveal deep convective cloud clusters with extended anvil cirrus clouds with the base at about 10 km. Source: E. Boer & V. Ramanathan (1992).

u8638 @ sdsc.edu

Feb 28 10:09

twocld.rgb

-J: Box B11

FIGURE 4: Comparison of the simulated greenhouse effect and cloud forcing with ERBE results.

Figure 4a: ERBE results for the tropical Pacific as shown in Ramanathan and Collins (1991).

Top left panel shows, G , the greenhouse effect of the atmosphere and clouds plotted against SST. Top right panel shows G_a , the greenhouse effect of just the atmosphere plotted against SST. Bottom left panel shows a scatter plot of shortwave cloud forcing against longwave cloud forcing.

Figure 4b: Same as Figure 4a, but for computed fields by ECHAM.

FIG. Monthly mean greenhouse effect and cloud-forcing terms for the Pacific extending from 140° E to 90° W. Each point in this and other figures represents a monthly averaged value for a 2.5° (latitude) × 2.5° (longitude) region. *a*, Total greenhouse effect, G , plotted as a function of SST for April 1987 from 85° N to 85° S. Regions with SST < 274 K are excluded. *b*, Atmospheric greenhouse effect G_a against SST for the tropical Pacific from 25° N to 25° S for April 1987. The least-squares linear slope is $dG_a/dT = 6.5 \text{ W m}^{-2} \text{ K}^{-1}$ with a correlation coefficient of $r = 0.96$. *c*, Cloud short-wave forcing against cloud long-wave forcing for the region 10° N to 10° S for 4/85 (•), 2/87 (○), 2/88 (×), and 4/87 (·). The linear correlation is $dC_s/dC_l = -0.951$ ($r = -0.9$).

FIGURE 4a: ERBE Data

Source: Ramanathan & Collins (1991).

ECHAM

FIGURE 4b

FIGURE 5: Changes in SST and radiation fluxes during El-Niño of 1987.

Figure 5a (first image): Observed changes between April 1985 and April 1987. Top left is change in SST. Top right is change in atmospheric greenhouse effect. Bottom left is change in shortwave cloud forcing. Bottom right is change in longwave cloud forcing.

Figure 5b (second image): Same as Figure 5a, but for the computed changes by the model ECHAM.

SST (K)

4/87 - 4/85

-2. -1. 0. 1. 2. 3. 4. 5.

Shortwave Cloud Forcing (W/m^2)

-100. -50. -15. 0. 15. 40. 60.

Atmospheric Greenhouse Effect (W/m^2)

-25. -15. -5. 5. 15. 25.

Longwave Cloud Forcing (W/m^2)

-45. -25. -5. 5. 15. 35. 55. 75.

Computing and Network Facilities for the joint Satellite/ARM modeling studies at the California Space Institute/Climate Research Division at Scripps Institution of Oceanography

Figure 6
Source: J. del Corral

END

**DATE
FILMED**

7 / 10 / 92