DOE/ER/13911--T2

DOE/ER/13911--T2

DE92 012574

ANNUAL PROGRESS REPORT FOR
"TRANSPORT PROPERTIES OF MULTI-COMPONENT
FLUIDS AND OF SUSPENSIONS"

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Prepared By

Irwin Oppenheim

and

John McBride

September 15, 1989

DOE Grant No. DE-FG03-88ER13911

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

I.	SUMMARY OF PROGRESS	1
II.	FOKKER-PLANCK DESCRIPTION OF SUSPENSIONS.....	2
	A. Hamiltonian and the Equations of Motion	2
	B. Liouville Operator	6
	C. Equilibrium Properties	7
	D. Derivation of the Fokker-Planck Equation	8
	E. Distribution Function for the Bath Particles.....	21
III.	SMOLUCHOWSKI EQUATION FOR THE REDUCED DISTRIBUTION FUNCTION.....	21
IV.	ELLIPSOIDAL BROWNIAN PARTICLE	29
V.	FUTURE WORK	34
	REFERENCES.....	36

I. SUMMARY OF PROGRESS

This report describes work performed under grant No. DE-FG03-88ER13911 for the period June 15, 1988 through June 14, 1989. During this time, significant progress was made in the derivation of the fundamental equations describing suspensions and multicomponent fluid flow.

We first considered a system consisting of n spherical heavy (Brownian) particles of mass M immersed in a bath of N spherical particles of mass m (suspensions). The deviations of the bath from equilibrium are due to the nonequilibrium motions of the Brownian particles. The densities of the bath and of a Brownian particle are similar. An expansion in powers of the mass ratio, $\epsilon = (m/M)^{1/2}$, yields a Fokker-Planck equation for the distribution function of the Brownian particles, $W(R^n, P^n, t)$, including the effects of direct and hydrodynamic interactions amongst these particles. The effect of the Brownian particle motion on the bath properties has also been described. The conditions under which a closed equation for the coordinate space distribution function, $R(R^n, t)$, can be obtained have been investigated and a Smoluchowski equation for this quantity has been derived.

The techniques utilized above were then applied to a system containing n Brownian particles of arbitrary shape in a bath of N spherical bath particles. In this more general case, we have also obtained a Fokker-Planck equation [Equation (44)] for the distribution function $W(Q^n, P^n, t)$, where Q^n and P^n denote the generalized coordinates and momenta of the Brownian particles. The treatment includes translation, rotation and internal motions (vibrations). The effects of the Brownian particle motion on the bath have been described [see Equation (66)] and a Smoluchowski equation [Equation (96)] for the distribution function $R(Q^n, t)$ has been derived. The conditions under which the translational and rotational motions can be separated have been studied. The derivation of these fundamental equations for nonspherical Brownian particles are

described in Sections II and III. A specific example for an ellipsoidal Brownian particle is given in Section IV, which illustrates some of the simplifications due to symmetries in certain problems.

The equations of hydrodynamics for multicomponent systems have been derived. The special case in which a solute molecule is much larger than a solvent molecule has been treated and new expressions for the chemical potentials in this system have been obtained. Multicomponent hydrodynamics will be considered further during the second year of the grant and results will be documented in future reports.

II. FOKKER-PLANCK DESCRIPTION OF SUSPENSIONS

We wish to study the conditions under which a Fokker-Planck (or generalized Fokker-Planck) equation can be used to study the time-dependence of a reduced distribution function. In particular, is the Fokker-Planck equation a suitable starting point for studying the reduced distribution function for heavy particles suspended in a fluid of light particles (the bath)?

A. Hamiltonian and the Equations of Motion

We consider an overall system consisting of n identical, non-spherical, heavy particles of mass M with internal degrees of freedom immersed in a bath of N identical particles of mass m . The deviations of the bath from equilibrium are due to the non-equilibrium motions of the heavy particles. We assume that all interaction potentials are short-range and that the densities of the heavy particles and of the bath are similar.

The generalized coordinates and momenta of the α th heavy particle are denoted by Q_α and P_α , respectively. These quantities are of two types: those describing the translational and rotational motions of the particle as a whole will be denoted by $Q_{\alpha B}$, $P_{\alpha B} \equiv X_{\alpha B}$; those describing

the internal motions (vibrations) of the particle will be denoted by $Q_{\alpha I}$, $P_{\alpha I} \equiv X_{\alpha I}$. The motions described by $X_{\alpha B}$ are slow due to the heavy mass of the particle and the subscript B stands for Brownian. The motions described by $X_{\alpha I}$ need not be slow but we shall assume that they are weakly coupled to the bath. The n heavy particles will be called the system.

The Hamiltonian $H_s(X_s)$ for the n interacting heavy particles can be written:

$$H_s(X_s) = H_B(X_B) + H_I(X_I) + V(Q), \quad (1)$$

where $X_s \equiv Q, P$; $X_B \equiv Q_B, P_B$ and $X_I \equiv Q_I, P_I$. Here, Q stands for all the generalized coordinates of the n heavy particles, Q_B stands for all the generalized translational and rotational coordinates for the particles as a whole, etc. The first term on the RHS of Equation (1) is a sum of one particle terms describing the kinetic energies of translation and rotation. The second term on the RHS of Equation (1) is a sum of one particle terms describing the kinetic and potential energies of the internal motions of the heavy particles. Within each particle, there will be a sum of terms for each mode of vibration. At a later stage, these modes will be assumed to be harmonic, for simplicity. The third term on the RHS of Equation (1) describes the interactions among the heavy particles; it is a sum of two-body terms each of which goes to zero when the center of mass separation of the two particles is larger than ξ_s .

The generalized coordinates and momenta of the i th bath particle will be denoted by q_i and p_i , respectively. The Hamiltonian for the N bath particles is given by

$$H_v(X_v) = \sum_{i=1}^N H_i(q_i, p_i) + U(q) \quad (2)$$

where $X_v \equiv \mathbf{q}, \mathbf{p}$ and $\mathbf{q}, \mathbf{p}$ denotes all of the generalized coordinates, momenta of the N bath particles. The quantity H_i is the Hamiltonian for the i th isolated particle. The interaction among the bath particles is given by U which is a sum of two particle terms each of which goes to zero when the center of mass separation of the two bath particles is greater than ξ_v .

The Hamiltonian for the overall system consisting of the N bath particles and the n heavy particles is given by:

$$H(X_s, X_v) = H_s(X_s) + H_v(X_v) + \Phi(\mathbf{q}, \mathbf{Q}), \quad (3)$$

where Φ describes the interaction between the bath particles and the heavy particles. It is again a sum of two-body terms

$$\Phi = \sum_{\alpha=1}^n \sum_{i=1}^N \phi(\mathbf{q}_i, \mathbf{Q}_\alpha) \quad (4)$$

where ϕ goes to zero when the center of mass separation between the i th bath particle and the α th heavy particle is larger than ξ .

The Hamilton equations of motion for the overall system are:

$$\dot{\mathbf{q}}_i = \partial H_v / \partial \mathbf{p}_i, \quad \dot{\mathbf{p}}_i = -\partial H_v / \partial \mathbf{q}_i - \partial \Phi / \partial \mathbf{q}_i \quad (5a)$$

$$\dot{\mathbf{Q}}_{\alpha B} = \partial H_B / \partial \mathbf{P}_{\alpha B}, \quad \dot{\mathbf{P}}_{\alpha B} = -\frac{\partial H_B}{\partial \mathbf{Q}_{\alpha B}} - \frac{\partial V}{\partial \mathbf{Q}_{\alpha B}} - \frac{\partial \Phi}{\partial \mathbf{Q}_{\alpha B}} \quad (5b)$$

$$\dot{\mathbf{Q}}_{\alpha I} = \frac{\partial H_i}{\partial \mathbf{P}_{\alpha I}}, \quad \dot{\mathbf{P}}_{\alpha I} = -\frac{\partial H_i}{\partial \mathbf{Q}_{\alpha I}} - \frac{\partial V}{\partial \mathbf{Q}_{\alpha I}} - \frac{\partial \Phi}{\partial \mathbf{Q}_{\alpha I}} \quad (5c)$$

There are two smallness parameters in this system: one of them, $\varepsilon \equiv (\frac{m}{M})^{1/2} \ll 1$, is due to the difference in masses of a heavy particle and a bath particle; the other, $\lambda \ll 1$, is associated with the weak coupling of the internal modes of the heavy particles with the bath.

We recall^{1,2} that we obtain simple Fokker-Planck or Langevin descriptions of Brownian motion when the momentum of the Brownian particle obeys the inequalities:

$$(M/m) |p| > |P_B| > |p|. \quad (6)$$

It is then convenient to define a reduced momentum

$$P_B^* \equiv \varepsilon P_B \quad (7)$$

which is assumed to be of order $\varepsilon^0 = 1$. In terms of P_B^* and Q_B , Equations (5b) yield $\dot{Q}_{\alpha B} \sim \varepsilon$, $\dot{P}_{\alpha B}^* \sim \varepsilon$.

Finally we rewrite Equation (5c) to make the weak coupling explicit

$$\dot{P}_{\alpha I} = -\partial H_s / \partial Q_{\alpha I} - \lambda \frac{\partial \Phi}{\partial Q_{\alpha I}}. \quad (8)$$

The overall system now has 5 characteristic times: (1) $\tau_v \sim 10^{-12}$ sec, the relaxation time of the isolated bath; (2) τ_I , the characteristic time of the internal motions which are arbitrary; (3) $\tau_B \sim \varepsilon^{-1} \tau_v$, the characteristic time for the isolated Brownian motions; (4) $\tau_{RB} \sim \varepsilon^{-2} \tau_v$, the relaxation time of the Brownian motions to equilibrium due to interactions with the bath; and (5) $\tau_{RI} \sim \lambda^{-2} \tau_v$, the relaxation time of the internal motions of the heavy particles to equilibrium due to interactions with the bath. We are assuming that the number of degrees of freedom in the isolated bath is sufficiently large that its macroscopic variables decay to their equilibrium values in the time τ_v . However, the isolated heavy particles do not come to equilibrium because of their

small number of degrees of freedom. They approach equilibrium due to their interactions with the bath. It is the separation in time scales given above that enables us to obtain a simplified equation of motion for the reduced heavy particle distribution function.

B. Liouville Operator

The Liouville operator for the overall system in $X^* \equiv X_v + X_I + X_B^* (\equiv Q_B, P_B^*)$ space is:

$$L(X^*) = L_0(X_v, Q) + L_I(X_I, q, Q_B) + L_B(X_B^*, q, Q_I) \quad (9)$$

where

$$L_0 = -\frac{\partial H_v}{\partial p} \cdot \frac{\partial}{\partial q} + \frac{\partial H_v}{\partial q} \cdot \frac{\partial}{\partial p} + \frac{\partial \Phi}{\partial q} \cdot \frac{\partial}{\partial p}, \quad (10)$$

$$\begin{aligned} L_I &= -\frac{\partial H_I}{\partial P_I} \cdot \frac{\partial}{\partial Q_I} + \frac{\partial H_I}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \frac{\partial V}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \lambda \frac{\partial \Phi}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} \\ &\equiv L_I^0 + \lambda \frac{\partial \Phi}{\partial Q_I} \cdot \frac{\partial}{\partial P_I}, \end{aligned} \quad (11)$$

and

$$L_B = \epsilon \left[-\frac{\partial H_B}{\partial P_B^*} \cdot \frac{\partial}{\partial Q_B} + \frac{\partial H_B}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} + \frac{\partial V}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} + \frac{\partial \Phi}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right] \quad (12)$$

The Liouville operator L_0 corresponds to a system with Hamiltonian

$$H_0 = H_v + \Phi \quad (13)$$

in which all of the motions of the heavy particle are fixed.

C. Equilibrium Properties

Before proceeding with the derivation for the equation of motion for the reduced distribution function of the heavy particles, it is convenient to study the equilibrium properties of the overall system.

The equilibrium distribution function for the overall system is given by:

$$\begin{aligned} \rho_e(X^*) &= e^{-\beta H} / \int e^{-\beta H} dX^* \\ &= \frac{e^{-\beta H_v} e^{-\beta \Phi} e^{-\beta H_s}}{\int dX_s^* e^{-\beta H_s} \int dX_v e^{-\beta H_v} e^{-\beta \Phi}}. \end{aligned} \quad (14)$$

The potential of mean force for the heavy particles in the presence of the bath is defined by

$$e^{-\beta w(Q)} \equiv \int \rho_v e^{-\beta \Phi} dX_v, \quad (15)$$

where ρ_v is the equilibrium distribution function for the isolated bath, i.e.

$$\rho_v \equiv e^{-\beta H_v} / \int e^{-\beta H_v} dX_v. \quad (16)$$

Thus, Equation (14) becomes:

$$\rho_e = \frac{\rho_v e^{-\beta \Phi} e^{-\beta H_s}}{\int dX_s^* e^{-\beta H_s} e^{-\beta w}}. \quad (17)$$

The equilibrium reduced distribution function for the heavy particles, $W_e(X_s^*)$, is given by:

$$W_e(X_s^*) \equiv \int \rho_e(X^*) dX_v = \frac{e^{-\beta H_s} e^{-\beta w}}{\int e^{-\beta H_s} e^{-\beta w} dX_s^*} \quad (18)$$

The equilibrium conditional distribution function for the bath, $\tilde{\rho}_e(X_v, Q)$, is given by

$$\tilde{\rho}_e(X_v, Q) \equiv \rho_e / W_e = \rho_v e^{-\beta \Phi} e^{\beta w} \quad (19)$$

which is normalized, i.e.

$$\int \tilde{\rho}_e dX_v = 1. \quad (20)$$

The distribution function $\tilde{\rho}_e$ plays an essential role in our future developments. The bath relaxes to this distribution function in the presence of fixed heavy particles on the time scale τ_v .

Averages with this function are given by:

$$\bar{B} \equiv \int \tilde{\rho}_e B dX_v \quad (21)$$

$$\hat{B} \equiv B - \bar{B} \quad (22)$$

where B is an arbitrary dynamical variable.

D. Derivation of the Fokker-Planck Equation

The equation of motion for the distribution function of the overall system, $\rho(X^*, t)$, is the Liouville equation

$$\frac{\partial \rho}{\partial t} \equiv \dot{\rho}(t) = L \rho(t). \quad (23)$$

In order to obtain an equation for the reduced non-equilibrium distribution function, $W(X_s^*, t)$, it is convenient to introduce the projection operator, P ,¹⁻⁴ which is defined by

$$PB \equiv \tilde{\rho}_e \int B dX_v \quad (24)$$

In particular:

$$P\rho(t) = \tilde{\rho}_e W(t) \equiv y(t) \quad (25a)$$

$$(1-P) \rho(t) \equiv z(t), \quad (25b)$$

where $W(X_s^*, t)$ is the time dependent reduced distribution function for the heavy particles. The quantities $y(t)$ and $z(t)$ have the properties

$$\int y(t) dX_v = W(t) \quad (26a)$$

$$\int z(t) dX_v = 0 \quad (26b)$$

The equations for the time derivatives of $y(t)$ and $z(t)$ follow from Equation (23) in the form:

$$\dot{y}(t) = \tilde{\rho}_e \dot{W}(t) = \tilde{\rho}_e \int dX_v L(y(t) + z(t)) \quad (27a)$$

$$\dot{z}(t) = (1-P) L (1-P) z(t) + (1-P) L y(t) \quad (27b)$$

We make use of the properties:

$$\int dX_v L_0 B = 0 \quad (28)$$

$$\int dX_v L_I^0 B = L_I^0 \int dX_v B \quad (29)$$

$$\begin{aligned} \int dX_v L_B B = \epsilon \left[-\frac{\partial H_B}{\partial P_B^*} \cdot \frac{\partial}{\partial Q_B} + \frac{\partial (H_B + V)}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right] \int dX_v B \\ + \epsilon \int \frac{\partial \Phi}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} B dX_v \end{aligned} \quad (30)$$

and Equations (26) to write Equation (27a) as:

$$\begin{aligned} \dot{W}(X_s^*, t) = L_I^0 W(t) + \lambda \frac{\partial w}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} W(t) \\ + \epsilon \left[-\frac{\partial H_B}{\partial P_B^*} \cdot \frac{\partial}{\partial Q_B} + \frac{\partial (H_B + V)}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} + \frac{\partial w}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right] W(t) \\ + \lambda \frac{\partial}{\partial P_I} \cdot \int \frac{\partial \Phi}{\partial Q_I} z(t) dX_v + \epsilon \frac{\partial}{\partial P_B^*} \cdot \int \frac{\partial \Phi}{\partial Q_B} z(t) dX_v \end{aligned} \quad (31)$$

where

$$\frac{\partial w}{\partial Q} = \overline{\frac{\partial \Phi}{\partial Q}}. \quad (32)$$

Equation (31) is exact and contains terms of order 1, λ , ϵ , λz and ϵz on the RHS.

We now turn our attention to Equation (27b) for $\dot{z}(t)$. The homogeneous term on the RHS can be written

$$\begin{aligned} (1-P) L z(t) &= (1-P) (L_o + L_I^o) \dot{z}(t) + (1-P) \left(L_B + \lambda \frac{\partial \Phi}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} \right) z(t) \\ &= (L_o + L_I^o) \dot{z}(t) + (1-P) \left(L_B + \lambda \frac{\partial \Phi}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} \right) z(t). \end{aligned} \quad (33)$$

The first term on the RHS of Equation (33) is of order 1 and the second term of order λ or ϵ . The inhomogeneous term is given by:

$$(1-P) L y(t) \equiv (1-P) L (\tilde{\rho}_e W(t)) = (1-P) [W(t) L \tilde{\rho}_e + \tilde{\rho}_e L W(t)]. \quad (34)$$

Using Equations (10)-(12) for L and the definition of $\tilde{\rho}_e$ in Equation (19), we find

$$L \tilde{\rho}_e = \tilde{\rho}_e \left[\beta \lambda \frac{\partial H_I}{\partial P_I} \cdot \frac{\partial \Phi}{\partial Q_I} + \beta \epsilon \frac{\partial H_B}{\partial P_B} \cdot \frac{\partial \Phi}{\partial Q_B} \right] \quad (35)$$

$$P L \tilde{\rho}_e = 0. \quad (36)$$

Since

$$W(t) = W(X_s^*, t), \quad (37)$$

$$L W(t) = (L_I + L_B) W(t) \quad (38)$$

and

$$(1-P) \tilde{\rho}_e L W(t) = \tilde{\rho}_e \left[\lambda \frac{\widehat{\partial \Phi}}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \varepsilon \frac{\widehat{\partial \Phi}}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right] W(t) . \quad (39)$$

Thus, the inhomogeneous term becomes:

$$(1-P) L y(t) = \tilde{\rho}_e \left[\lambda \frac{\widehat{\partial \Phi}}{\partial Q_I} \cdot \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) + \varepsilon \frac{\widehat{\partial \Phi}}{\partial Q_B} \cdot \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_I}{\partial P_B^*} \right) \right] W(t) . \quad (40)$$

Note that, as in the equation for $\dot{W}$, the inhomogeneous term is of order λ or ε . Substitution of Equations (33) and (40) into Equation (27b) yields:

$$\begin{aligned} \dot{z}(t) &= (L_o + L_I^o) z(t) + (1-P) \left(L_B + \lambda \frac{\widehat{\partial \Phi}}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} \right) z(t) \\ &+ \tilde{\rho}_e \left[\lambda \frac{\widehat{\partial \Phi}}{\partial Q_I} \cdot \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) + \varepsilon \frac{\widehat{\partial \Phi}}{\partial Q_B} \cdot \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_I}{\partial P_B^*} \right) \right] W(t) . \end{aligned} \quad (41)$$

Equations (31) and (41) are exact.

The exact formal solution of Equation (27b) is given by

$$z(t) = e^{(1-P)Lt} z(0) + \int_0^t e^{(1-P)L\tau} (1-P) L \tilde{\rho}_e W(t-\tau) d\tau \quad (42)$$

which can be substituted into Equation (31) to yield

$$\begin{aligned}
\dot{W}(t) = & \left[L_I^0 + \lambda \frac{\partial w}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \varepsilon \left(-\frac{\partial H_B}{\partial P_B^*} \cdot \frac{\partial}{\partial Q_B} + \frac{\partial (H_B + V + w)}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right) \right] W(t) \\
& + \int dX_v \left[\lambda \frac{\partial \Phi}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \varepsilon \frac{\partial \Phi}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right] e^{(1-P)Lt} z(0) \\
& + \int dX_v \left(\lambda \frac{\partial \Phi}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \varepsilon \frac{\partial \Phi}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right) \int_0^t d\tau e^{(1-P)L\tau} \\
& \times \tilde{\rho}_e \left[\lambda \frac{\partial \Phi}{\partial Q_I} \cdot \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) + \varepsilon \frac{\partial \Phi}{\partial Q_B} \cdot \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_B}{\partial P_B^*} \right) \right] W(t-\tau) \quad (43)
\end{aligned}$$

Equation (43) is exact. It contains Euler or streaming terms of order 1, ε , or λ ; an initial value term which is at least of order ε or λ ; and time integrals over correlation functions which are at least of quadratic order in ε and/or λ .

In order to reduce Equation (43) to manageable form we must be able to:

- (a) Eliminate the initial value term containing $z(0)$;
- (b) Simplify the propagator $e^{(1-P)L\tau}$;
- (c) Obtain a simple expression for $W(t-\tau)$ in terms of $W(t)$.

All of these desiderata can be accomplished for small λ and ε and for time $t > \tau_v$ for the following reasons:

- (1) Correlation functions of the form $\overline{Ae^{Lot}B} \rightarrow \overline{A}\overline{B}$ for $t > \tau_v$ where A and B are dynamical variables involving bath coordinates and momenta.
- (2) $e^{Lot} z(0)$ decays to zero for $t > \tau_v$ since the conditional bath distribution function $\tilde{\rho}(t) \rightarrow \tilde{\rho}_e$ on this time scale.
- (3) The Liouville operations L^0 and L^0_I commute to order 1.

To quadratic order in the smallness parameters ε and λ and for $t > \tau_v$ we find:^{2,3}

- (1) The initial value term is negligible.
- (2) The limit of integration in the time integral in the last term on the RHS of Equation (43) can be changed from t to ∞ .

Since this term is explicitly of quadratic order in ε and/or λ , we need evaluate the propagator and the time dependence of $W(t-\tau)$ to order one only.

$$(3) \quad \text{Thus: } e^{(1-P)L\tau} \cong e^{(L_o + L_I^o)\tau} \cong e^{L_o\tau} e^{L_I^o\tau}$$

and

$$(4) \quad W(t-\tau) \cong e^{-L_I^o\tau} W(t)$$

Results (3) and (4) depend on the fact that the correlation function decays to zero for $t > \tau_v$. Further simplification arises because $L_I^o \tilde{\rho}_e \sim \lambda$.

Under these circumstances Equation (43) becomes:

$$\begin{aligned}
\dot{W}(t) = & \left[L_I^0 + \lambda \frac{\partial w}{\partial Q_I} \cdot \frac{\partial}{\partial P_I} + \varepsilon \left(-\frac{\partial H_B}{\partial P_B^*} \cdot \frac{\partial}{\partial Q_B} + \frac{\partial (H_B + V + w)}{\partial Q_B} \cdot \frac{\partial}{\partial P_B^*} \right) \right] W(t) \\
& + \varepsilon^2 \Gamma_{BB}(Q) : \frac{\partial}{\partial P_B^*} \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_B}{\partial P_B^*} \right) W(t) \\
& + \varepsilon \lambda \Gamma_{BI}(Q) : \frac{\partial}{\partial P_I} \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_B}{\partial P_B^*} \right) W(t) \\
& + \varepsilon \lambda \int_0^\infty d\tau \overline{\frac{\partial \Phi}{\partial Q_B} e^{L_o \tau} e^{L_I^0 \tau} \frac{\partial \Phi}{\partial Q_I}} : \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) \frac{\partial}{\partial P_B^*} e^{-L_I^0 \tau} W(t) \\
& + \lambda^2 \int_0^\infty d\tau \overline{\frac{\partial \Phi}{\partial Q_I} e^{L_o \tau} \frac{\partial}{\partial P_I} e^{L_I^0 \tau} \frac{\partial \Phi}{\partial Q_I}} : \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) e^{-L_I^0 \tau} W(t) . \quad (44)
\end{aligned}$$

where

$$\Gamma = \int_0^\infty d\tau \overline{\frac{\partial \Phi}{\partial Q} e^{L_o \tau} \frac{\partial \Phi}{\partial Q}} . \quad (45)$$

Equation (45) is the desired generalized Fokker-Planck equation. It contains short range interactions among the heavy particles in the V and w terms and long range hydrodynamic interactions due to the dependence of the correlation functions in the quadratic terms on Q . It contains Euler or streaming terms of order 1, λ or ε . When the Euler operator acts on W_e , we get zero. The first two quadratic terms are straight forward and involve Fokker-Planck operators of the usual type. These terms also have the property that when operating on W_e they yield zero to all orders.

The last two quadratic terms are more complicated and demand that we can obtain explicit expressions for the dynamics included in L_I^0 . [They yield zero when acting on the equilibrium distribution to lowest order.] The following approximations are valid in almost all cases:

(1) The potential of interaction between the bath and the heavy particles can be written as $\Phi(q, Q) = \Phi_0(q, Q_B) + \lambda \Phi_1(q, Q_B, Q_I)$

(2) The dependence of Φ , on Q_I is linear. In this case, the Γ 's are functions of Q_B only, the correlation functions in the last two terms become $\frac{\widehat{\partial \Phi}}{\partial Q} e^{L_0 \tau} \frac{\widehat{\partial \Phi}}{\partial Q}$, and we have to evaluate the

operator $e^{L_I^0 \tau} \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) e^{-L_I^0 \tau}$ acting on $W(t)$.

These operators are easily evaluated if the internal modes are harmonic and we can neglect the dependence of V on Q_I . For one oscillator:

$$H_I = P_I^2 / 2\mu + \mu \Omega^2 Q_I^2 / 2 \quad (46)$$

and

$$L_I^0 = - (P_I / \mu) \frac{\partial}{\partial Q_I} + \mu \Omega^2 Q_I \frac{\partial}{\partial P_I} \quad (47)$$

Then:

$$e^{-L_I^0 \tau} W(t) = W(X_I(\tau), X_B^*, t) \quad (48)$$

where

$$\begin{aligned} Q_I(\tau) &= Q_I \cos \Omega \tau + (P_I / \mu \Omega) \sin \tau \\ P_I(\tau) &= P_I \cos \Omega \tau - \mu \Omega Q_I \sin \Omega \tau. \end{aligned} \quad (49)$$

Thus

$$\begin{aligned}
\frac{\partial}{\partial P_I} e^{-L_I^0 \tau} W(\tau) &= \frac{\partial W(\tau, t)}{\partial Q_I(\tau)} \frac{\partial Q_I(\tau)}{\partial P_I} + \frac{\partial W(\tau, t)}{\partial P_I(\tau)} \frac{\partial P_I(\tau)}{\partial P_I} \\
&= \frac{\partial W(\tau, t)}{\partial Q_I(\tau)} \frac{1}{\mu \Omega} \sin \Omega \tau + \frac{\partial W(\tau, t)}{\partial P_I(\tau)} \cos \Omega \tau
\end{aligned} \tag{50}$$

where $W(\tau, t) \equiv W(X_I(\tau), X_B^*, t)$ and

$$e^{L_I^0 \tau} \frac{\partial}{\partial P_I} e^{-L_I^0 \tau} W(t) = \frac{1}{\mu \Omega} \frac{\partial W(t)}{\partial Q_I} \sin \Omega \tau + \frac{\partial W(t)}{\partial P_I} \cos \Omega \tau. \tag{51}$$

Further,

$$\begin{aligned}
e^{L_I^0 \tau} \beta \frac{\partial H_I}{\partial P_I} e^{-L_I^0 \tau} W(t) &= \beta P_I(-\tau) / \mu \\
&= (\beta / \mu) (P_I \cos \Omega \tau + \mu \Omega Q_I \sin \Omega \tau)
\end{aligned} \tag{52}$$

and

$$\begin{aligned}
e^{L_I^0 \tau} \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial H_I}{\partial P_I} \right) e^{-L_I^0 \tau} W(t) &= \\
\left[\cos \Omega \tau \left(\frac{\partial}{\partial P_I} + \beta \frac{P_I}{\mu} \right) + \frac{\sin \Omega \tau}{\mu \Omega} \left(\frac{\partial}{\partial Q_I} + \beta \mu \Omega^2 Q_I \right) \right] W(t)
\end{aligned} \tag{53}$$

Substitution of Equation (53) for each internal mode simplifies the last two terms in Equation (44). It is remarkable that the $\cos \Omega \tau$ in Equation (53) contributes to the dissipation while the $\sin \Omega \tau$ term affects the dynamical oscillatory frequency.

For a general system, with linear interactions with the bath, we can write a compact form for the operator

$$O \equiv e^{L_I^0 \tau} \left(\frac{\partial}{\partial \mathbf{P}_I} + \beta \frac{\partial H_I}{\partial \mathbf{P}_I} \right) e^{-L_I^0 \tau} \quad (54)$$

acting on the distribution function $W(t)$. We use the following facts:

$$e^{-L_I^0 \tau} W(\mathbf{Q}, \mathbf{P}, t) = W(\mathbf{Q}_o(\tau), \mathbf{P}_o(\tau), t) \equiv W(\tau, t) \quad (55)$$

where the subscript o implies that the τ time dependence is due to the Liouville operator L_I^0 . It then follows that

$$\frac{\partial}{\partial \mathbf{P}_I} e^{-L_I^0 \tau} W(\mathbf{Q}, \mathbf{P}, t) = \frac{\partial W(\tau, t)}{\partial \mathbf{Q}_o(\tau)} \cdot \frac{\partial \mathbf{Q}_o(\tau)}{\partial \mathbf{P}_I} + \frac{\partial W(\tau, t)}{\partial \mathbf{P}_o(\tau)} \cdot \frac{\partial \mathbf{P}_o(\tau)}{\partial \mathbf{P}_I} \quad (56)$$

and

$$\begin{aligned} e^{L_I^0 \tau} \frac{\partial}{\partial \mathbf{P}_I} e^{-L_I^0 \tau} W(\mathbf{Q}, \mathbf{P}, t) &= \left[e^{L_I^0 \tau} \frac{\partial \mathbf{Q}_o(\tau)}{\partial \mathbf{P}_I} \right] \cdot \frac{\partial W(\mathbf{Q}, \mathbf{P}, t)}{\partial \mathbf{Q}} \\ &+ \left[e^{L_I^0 \tau} \frac{\partial \mathbf{P}_o(\tau)}{\partial \mathbf{P}_I} \right] \cdot \frac{\partial W(\mathbf{Q}, \mathbf{P}, t)}{\partial \mathbf{P}} \end{aligned} \quad (57)$$

The term

$$e^{L_I^0 \tau} \beta \frac{\partial H_I}{\partial \mathbf{P}_I} e^{-L_I^0 \tau} W(\mathbf{Q}, \mathbf{P}, t) = \beta \left[\frac{\partial H_I}{\partial \mathbf{P}_I} \right] (-\tau) W(\mathbf{Q}, \mathbf{P}, t) \quad (58)$$

where again the τ time dependence is due to the Liouville operator L_I^0 . A more convenient form for this quantity can be obtained by noting that

$$\frac{\partial H_I}{\partial \mathbf{P}_I} = \frac{\partial (H_I + V)}{\partial \mathbf{P}_I} \quad (59)$$

and that the Hamiltonian, $H_I + V$, is invariant to the time dependence generated by L_I^0 . Thus

$$\begin{aligned} e^{L_I^0 \tau} \frac{\partial H_I}{\partial \mathbf{P}_I} &= e^{L_I^0 \tau} \frac{\partial}{\partial \mathbf{P}_I} (H_I + V)(\mathbf{Q}_0(\tau), \mathbf{P}_0(\tau)) \\ &= \left[e^{L_I^0 \tau} \frac{\partial \mathbf{Q}_0(\tau)}{\partial \mathbf{P}_I} \right] \cdot \frac{\partial (H_I + V)}{\partial \mathbf{Q}}(\mathbf{Q}, \mathbf{P}) \\ &\quad + \left[e^{L_I^0 \tau} \frac{\partial \mathbf{P}_0(\tau)}{\partial \mathbf{P}_I} \right] \cdot \frac{\partial (H_I + V)}{\partial \mathbf{P}}(\mathbf{Q}, \mathbf{P}). \end{aligned} \quad (60)$$

Combining Equations (51) and (58), we obtain:

$$\begin{aligned} OW(\mathbf{Q}, \mathbf{P}, t) &= \left\{ \left[e^{L_I^0 \tau} \frac{\partial \mathbf{Q}_0(\tau)}{\partial \mathbf{P}_I} \right] \cdot \left(\frac{\partial}{\partial \mathbf{Q}_I} + \beta \frac{\partial (H_I + V)}{\partial \mathbf{Q}_I} \right) \right. \\ &\quad \left. + \left[e^{L_I^0 \tau} \frac{\partial \mathbf{P}_0(\tau)}{\partial \mathbf{P}_I} \right] \cdot \left(\frac{\partial}{\partial \mathbf{P}_I} + \beta \frac{\partial (H_I + V)}{\partial \mathbf{P}_I} \right) \right\} W(\mathbf{Q}, \mathbf{P}, t) \end{aligned} \quad (61)$$

where we have used the facts that

$$\frac{\partial Q_o(\tau)}{\partial P_I} \left(\frac{\partial P_o(\tau)}{\partial P_I} \right) \text{ are zero unless } Q = Q_I (P = P_I).$$

Substitution of Equation (61) into Equation (44) yields the following expression for the quadratic terms:

$$\begin{aligned} & \left\{ \varepsilon^2 \Gamma_{BB} : \frac{\partial}{\partial P_B^*} \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_B}{\partial P_B^*} \right) \right. \\ & + \varepsilon \lambda \Gamma_{IB} : \frac{\partial}{\partial P_I^*} \left(\frac{\partial}{\partial P_B^*} + \beta \frac{\partial H_B}{\partial P_B^*} \right) \\ & + \left(\varepsilon \lambda \chi_{BI} : \frac{\partial}{\partial P_B^*} + \lambda^2 \psi_{II} \right) \cdot \left(\frac{\partial}{\partial Q_I} + \beta \frac{\partial (H_I + V)}{\partial Q_I} \right) \\ & \left. + \left(\varepsilon \lambda \chi_{BI} : \frac{\partial}{\partial P_B^*} + \lambda^2 \hat{\psi}_{II} \right) \cdot \left(\frac{\partial}{\partial P_I} + \beta \frac{\partial (H_I + V)}{\partial P_I} \right) \right\} W(t) \end{aligned} \quad (62)$$

where

$$\chi_{BI} = \int_0^\infty d\tau \frac{\widehat{\partial \Phi}}{\partial Q_B} e^{L_o \tau} \frac{\widehat{\partial \Phi}}{\partial Q_I} \cdot \left[e^{L_I^o \tau} \frac{\partial Q_o(\tau)}{\partial P_I} \right] \quad (63)$$

$$\psi_{II} = \int_0^\infty d\tau \frac{\widehat{\partial \Phi}}{\partial Q_I} e^{L_o \tau} \frac{\widehat{\partial \Phi}}{\partial Q_I} : \frac{\partial}{\partial P_I} \left[e^{L_I^o \tau} \frac{\partial Q_o(\tau)}{\partial P_I} \right] \quad (64)$$

and $\hat{\chi}_{BI} (\hat{\psi}_{II})$ is obtained from $\chi_{BI} (\psi_{II})$ by replacing $\left[e^{L_I^o \tau} \frac{\partial Q_o(\tau)}{\partial P_I} \right]$ by $\left[e^{L_I^o \tau} \frac{\partial P_o(\tau)}{\partial P_I} \right]$.

E. Distribution Function for the Bath Particles

The technique that we have used also yields the first order terms in the expansion of

$$z(t) \equiv \rho(t) - \tilde{\rho}_e W(t) \quad (65)$$

in powers of ε and λ . Using the approximations described above, we find that the non-equilibrium conditional distribution function for the bath is given by

$$\begin{aligned} \tilde{\rho}(t) \equiv \rho(t)/W(t) &= \tilde{\rho}_e + \int_0^\infty d\tau \tilde{\rho}_e e^{L_0 \tau} \\ &\times \left[\varepsilon \frac{\widehat{\partial \Phi}}{\partial \mathbf{Q}_B} \cdot \left(\frac{\partial}{\partial \mathbf{P}_B^*} + \beta \frac{\partial H_B}{\partial \mathbf{P}_B^*} \right) + \lambda \frac{\widehat{\partial \Phi}}{\partial \mathbf{Q}_I} \cdot e^{L_I^0 \tau} \left(\frac{\partial}{\partial \mathbf{P}_I} + \beta \frac{\partial H_I}{\partial \mathbf{P}_I} \right) e^{-L_I^0 \tau} \right] \ln W(t) \end{aligned} \quad (66)$$

which is valid for $t > \tau_D$ and to first order in the smallness parameters. The correction to $\tilde{\rho}_e$ is due to the fact that the bath does not adjust instantaneously to the state of the heavy particles. Note that the correction term is zero for $W(t) = W_e$.

III. SMOLUCHOWSKI EQUATION FOR THE REDUCED DISTRIBUTION FUNCTION

We next study under what circumstances we can obtain a relatively simple equation for the time dependence of the reduced distribution function in coordinate space, $R(Q, t)$. This will occur if the momentum relaxation of the heavy particles is faster than their relaxation in coordinate space. The resulting equation - the Smoluchowski equation - is valid only for times longer than the momentum relaxation time which is proportional to $[\beta \varepsilon^2 \Gamma/m]^{-1}$ and only for situations in which $\frac{\partial R}{\partial Q}$ is small in some sense.

In order to simplify the algebra we shall consider the same overall system treated above in the absence of internal degrees of freedom. Again, we first examine the equilibrium forms of the reduced distribution functions and then introduce a suitable projection operator formalism.

The equilibrium form of W is given in Equation (18) by

$$W_e(Q, P) = \frac{e^{-\beta H_B(Q, P)} e^{-\beta w(Q)} e^{-\beta V(Q)}}{\int e^{-\beta H_B(Q, P)} e^{-\beta w(Q)} e^{-\beta V(Q)} dP dQ} \quad (67)$$

where we have dropped the subscript B on Q and P . The only motions of the heavy particles now are Brownian. The equilibrium coordinate space distribution function is given by

$$R_e(Q) \equiv \int W_e dP = \frac{e^{-\beta W} e^{-\beta V} \int e^{-\beta H_B} dP}{\int e^{-\beta w} e^{-\beta V} dQ \int e^{-\beta H_B} dP} \quad (68)$$

The equilibrium conditional distribution function for the momenta is given by:

$$\begin{aligned} \tilde{\Pi}_e(P, Q) &\equiv W_e(Q, P) / R_e(Q) \\ &= e^{-\beta H_B(Q, P)} / \int e^{-\beta H_B(Q, P)} dP. \end{aligned} \quad (69)$$

We recall that H_B contains the kinetic energy terms for translation and rotation. The distribution $\tilde{\Pi}_e$ is the form the momentum distribution function relaxes to for fixed Q .

We now define the projection operator $\hat{P}$ by⁴:

$$\hat{P} B = \tilde{\Pi}_e \int B dP. \quad (70)$$

In particular,

$$\hat{P} W(Q, P, t) = \tilde{P}_e R(Q, t) = \hat{y}(t) \quad (71a)$$

and

$$(1 - \hat{P}) W(Q, P, t) = \hat{z}(t) \quad (71b)$$

where

$$\int dP \hat{y}(t) = R(Q, t) \quad (72a)$$

$$\int dP \hat{z}(t) = 0. \quad (72b)$$

The equation for $\dot{R}(t)$ is obtained by applying the projection operator $\hat{P}$ to Equation (44), in the absence of the internal modes, and integrating the results over P . The equation for $\dot{\hat{z}}$ is obtained by applying the projection operator $(1 - \hat{P})$ to this equation. We rewrite equation (44) in the form:

$$\dot{W}(Q, P, t) = O_E W(t) + O_D W(t) \quad (73)$$

where the Euler operator, O_E , is given by

$$O_E = - \frac{\partial H_B}{\partial P} \cdot \frac{\partial}{\partial Q} + \frac{\partial (H_B + V + w)}{\partial Q} \cdot \frac{\partial}{\partial P} \quad (74)$$

and the dissipative operator, O_D , is

$$O_D = \Gamma(Q) : \frac{\partial}{\partial P} \left(\frac{\partial}{\partial P} + \beta \frac{\partial H_B}{\partial P} \right). \quad (75)$$

Here,

$$\Gamma(Q) = \int_0^\infty \frac{\partial \Phi}{\partial Q} e^{L \circ \tau} \frac{\partial \Phi}{\partial Q} d\tau \quad (76)$$

is a symmetric tensor.

We make use of the properties:

$$\int dP O_D B = 0 \quad (77)$$

$$\int dP O_E B = \frac{\partial}{\partial Q} \cdot \int dP \frac{\partial H_B}{\partial P} B \quad (78)$$

$$\int dP \frac{\partial H_B}{\partial P} \tilde{\Pi}_e = - (1/\beta) \int dP \frac{\partial \tilde{\Pi}_e}{\partial P} = 0 \quad (79)$$

to write

$$\dot{R}(Q, t) = - \frac{\partial}{\partial Q} \cdot \int dP \frac{\partial H_B}{\partial P} \hat{z}(t). \quad (80)$$

The equation for $\hat{z}(t)$ is

$$\begin{aligned} \dot{\hat{z}}(t) &= (1 - \hat{P})(O_E + O_D) \hat{z}(t) \\ &+ (1 - \hat{P})(O_E + O_D) \hat{y}(t). \end{aligned} \quad (81)$$

In addition to the properties in Equations (77)-(79) we use the fact that

$$O_D \hat{y}(t) = (O_D \tilde{\Pi}_e) R(t) = 0 \quad (82)$$

to write the inhomogeneous term in Equation (81) as

$$\begin{aligned} (1-\hat{P})(O_E + O_D) \hat{y}(t) &= O_E (\tilde{\Pi}_e R(t)) \\ &= R(t) (O_E \tilde{\Pi}_e) + \tilde{\Pi}_e O_E R(t) \\ &= R(t) O_E \tilde{\Pi}_e - \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \cdot \frac{\partial}{\partial \mathbf{Q}} R(t). \end{aligned} \quad (83)$$

The effect of O_E on $\tilde{\Pi}_e$ is most easily obtained by using the fact that

$$\begin{aligned} O_E W_e &= 0 = O_E (\tilde{\Pi}_e R_e) \\ &= R_e O_E \tilde{\Pi}_e + \tilde{\Pi}_e O_E R_e. \end{aligned} \quad (84)$$

Thus

$$O_E \tilde{\Pi}_e = \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \cdot \frac{\partial}{\partial \mathbf{Q}} \ln R_e \quad (85)$$

and Equation (83) becomes

$$(1-\hat{P})(O_E + O_D) \hat{y}(t) = -\tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \cdot \left[\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right] R(t) \quad (86)$$

Substitution of Equation (86) into Equation (81) yields

$$\begin{aligned} \dot{\hat{z}}(t) = & O_D \hat{z}(t) + (1 - \hat{P}) O_E \hat{z}(t) \\ & - \tilde{\Pi}_e \left[\frac{\partial H_B}{\partial \mathbf{P}} \right] \cdot \left[\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right] R(t) \end{aligned} \quad (87)$$

which is an exact equation. It is important to note that

$$\int \tilde{\Pi}_e \dot{\mathbf{Q}} d\mathbf{P} = \int \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} d\mathbf{P} = 0 \quad (88)$$

so that the average time dependence of $\mathbf{Q}$ goes to zero once the momentum distribution function has relaxed. This implies that $R(\mathbf{Q}, t)$ relaxes more slowly than the momentum function.

The formal solution of Equation (87) is easily written down and substituted into Equation (80) to obtain:

$$\begin{aligned} \dot{R}(t) = & -\frac{\partial}{\partial \mathbf{Q}} \cdot \int d\mathbf{P} \frac{\partial H_B}{\partial \mathbf{P}} e^{(1 - \hat{P})(O_D + O_E)t} \hat{z}(0) \\ & + \frac{\partial}{\partial \mathbf{Q}} \cdot \int d\mathbf{P} \int_0^t d\tau \frac{\partial H_B}{\partial \mathbf{P}} e^{(1 - \hat{P})(O_D + O_E)\tau} \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \\ & \cdot \left[\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right] R(t) \end{aligned} \quad (89)$$

Equation (89) is exact. It contains an initial value term and a time integral of a correlation function.

In order to simplify this equation, we make the assertion that all derivatives with respect to $\mathbf{Q}$ have a smallness parameter δ associated with them. Thus the initial value term is at least of

order δ , the time integral is at least of order δ^2 , O_D is of order 1, and O_E is of order δ . We wish to obtain an equation which is valid to order δ^2 . The propagator e^{ODt} has the properties:

$$(1) \quad e^{ODt} \hat{z}(0) = 0 \quad \text{for } t > \tau_p$$

$$(2) \quad \int \tilde{\Pi}_e A e^{ODt} B d\mathbf{P} \rightarrow \int \tilde{\Pi}_e A d\mathbf{P} \int \tilde{\Pi}_e B d\mathbf{P} \quad \text{for } t > \tau_p$$

where τ_p is the momentum relaxation time. Thus, to order δ^2 , and for $t > \tau_p$, Equation (89) becomes:

$$\dot{R}(\mathbf{Q}, t) = \frac{\partial}{\partial \mathbf{Q}} \cdot \int_0^\infty d\tau \left[\int d\mathbf{P} \frac{\partial H_B}{\partial \mathbf{P}} e^{OD\tau} \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \right] \cdot \left(\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right) R(t) \quad (90)$$

In order to complete the derivation of the Smoluchowski equation we must study the effects of $e^{OD\tau}$ on $\tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}}$. We shall look at the effects of various powers of O_D on $\tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}}$ and look at the results in terms of components in the multidimensional generalized momentum space of the n heavy particles. We shall indicate components by Greek subscripts, use the summation convention for repeated indices, and use the facts that the tensor $\underline{\Gamma}$ is symmetric and that H_B is a quadratic function of the momenta. We obtain:

$$O_D^0 \tilde{\Pi}_e \frac{\partial H_B}{\partial P_\alpha} = \tilde{\Pi}_e \frac{\partial H_B}{\partial P_\alpha} \quad (91a)$$

$$\begin{aligned}
O_D \tilde{\Pi}_e \frac{\partial H_B}{\partial P_\alpha} &= \Gamma_{\gamma\gamma'} \frac{\partial}{\partial P_{\gamma'}} \left(\frac{\partial}{\partial P_\gamma} + \beta \frac{\partial H_B}{\partial P_\gamma} \right) \left(\tilde{\Pi}_e \frac{\partial H_B}{\partial P_\alpha} \right) \\
&= \Gamma_{\gamma\gamma'} \frac{\partial}{\partial P_{\gamma'}} \left(\tilde{\Pi}_e \frac{\partial^2 H_B}{\partial P_\gamma \partial P_\alpha} \right) = -\beta \tilde{\Pi}_e \frac{\partial H_B}{\partial P_{\gamma'}} \Gamma_{\gamma\gamma'} \frac{\partial^2 H_B}{\partial P_\gamma \partial P_\alpha} \\
&= \tilde{\Pi}_e \frac{\partial H_B}{\partial P_{\gamma'}} \left(-\beta \Gamma_{\gamma\gamma'} \frac{\partial^2 H_B}{\partial P_\gamma \partial P_\alpha} \right)_{\gamma'\alpha}
\end{aligned} \tag{91b}$$

$$\begin{aligned}
O_D^2 \left(\tilde{\Pi}_e \frac{\partial H_B}{\partial P_{\gamma'}} \right) &= \tilde{\Pi}_e \frac{\partial H_B}{\partial P_{\theta'}} \left(-\beta \Gamma_{\gamma\gamma'} \frac{\partial^2 H_B}{\partial P_\gamma \partial P_\alpha} \right)_{\theta'\gamma'} \left(-\beta \Gamma_{\gamma\gamma'} \frac{\partial^2 H_B}{\partial P_\gamma \partial P_\alpha} \right)_{\gamma'\alpha} \\
&\vdots \\
&\vdots
\end{aligned} \tag{91c}$$

We define the tensor

$$T_{\gamma\gamma'} = \Gamma_{\gamma\gamma'} \frac{\partial^2 H_B}{\partial P_\gamma \partial P_{\gamma'}} \tag{92}$$

and write

$$\begin{aligned}
e^{O_D \tau} \tilde{\Pi}_e \frac{\partial H_B}{\partial P} &= \tilde{\Pi}_e \frac{\partial H_B}{\partial P} \cdot \left[1 - \beta T_{\gamma\gamma'} \tau + \frac{\beta^2 \tau^2}{2} T_{\gamma\gamma'} \cdot T_{\gamma\gamma'} \dots \right] \\
&= \tilde{\Pi}_e \frac{\partial H_B}{\partial P} \cdot e^{-\beta T \tau}
\end{aligned} \tag{93}$$

Substitution of Equation (93) into Equation (90) and integration over τ yields:

$$\begin{aligned}
\dot{R}(\mathbf{Q}, t) &= \frac{\partial}{\partial \mathbf{Q}} \cdot \left[\int \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \frac{\partial H_B}{\partial \mathbf{P}} d\mathbf{P} \right] \cdot \beta^{-1} T^{-1} \cdot \left[\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right] R(t) \\
&= k_B T \frac{\partial}{\partial \mathbf{Q}} \cdot \underline{K}(\mathbf{Q}) \cdot \underline{T}^{-1}(\mathbf{Q}) \cdot \left(\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right) R(t)
\end{aligned} \tag{94}$$

where

$$\begin{aligned}
\underline{K}(\mathbf{Q}) &\equiv \int \tilde{\Pi}_e \frac{\partial H_B}{\partial \mathbf{P}} \frac{\partial H_B}{\partial \mathbf{P}} d\mathbf{P} \\
&= \frac{1}{\beta} \frac{\partial^2 H_B}{\partial \mathbf{P} \partial \mathbf{P}}
\end{aligned} \tag{95}$$

Thus:

$$\begin{aligned}
\dot{R}(\mathbf{Q}, t) &= (k_B T)^2 \frac{\partial}{\partial \mathbf{Q}} \cdot \underline{\Gamma}^{-1}(\mathbf{Q}) \cdot \left(\frac{\partial}{\partial \mathbf{Q}} - \frac{\partial \ln R_e}{\partial \mathbf{Q}} \right) R(t) \\
&= (k_B T)^2 \frac{\partial}{\partial \mathbf{Q}} \cdot \underline{\Gamma}^{-1}(\mathbf{Q}) \cdot \left[\frac{\partial}{\partial \mathbf{Q}} + \beta \left(\frac{\partial (W + V)}{\partial \mathbf{Q}} + \frac{\int \frac{\partial H_B}{\partial \mathbf{Q}} e^{-\beta H_B} d\mathbf{P}}{\int e^{-\beta H_B} d\mathbf{P}} \right) \right] R(t)
\end{aligned} \tag{96}$$

Equation (96) is the desired result. The direct interactions are included in the w , V and $\partial H_B / \partial \mathbf{Q}$ terms; the hydrodynamic interactions are included in $\underline{\Gamma}(\mathbf{Q})$.

IV. ELLIPSOIDAL BROWNIAN PARTICLE

In the special case of a single ellipsoidal Brownian particle, the Hamiltonian may be written as

$$\begin{aligned}
H &= P^2/2M + \frac{1}{2I_x} \left[\frac{(P_\phi - P_\psi \cos \theta)^2}{\sin^2 \theta} + P_\theta^2 \right] + \frac{P_\psi^2}{2I_z} \\
&+ H_v + \Phi(r^N, R, \theta, \phi) \\
&= \hat{H} + P_\psi^2/2I_z.
\end{aligned} \tag{97}$$

In Equation (97), R , and P refer to the center of mass position and momentum of the Brownian particle. The angles ϕ, θ, ψ are the Eulerian angles described in Reference 5, and P_ϕ, P_θ, P_ψ are the corresponding angular momenta. $I_x = I_y$ and I_z are the moments of inertia of the Brownian particle [See Figure 4-6 of Reference 5 for the topology]. We must consider P_ψ to be an externally fixed quantity since $\dot{P}_\psi = 0$, and ψ to be an auxiliary quantity determined by

$$\dot{\psi}(t) = \left[\frac{1}{I_x} \left(\frac{\cos^2 \theta}{\sin^2 \theta} \right) (t) + \frac{1}{I_z} \right] P_\psi - \frac{1}{I_x} \left[\frac{P_\phi \cos \theta}{\sin^2 \theta} \right] (t). \tag{98}$$

We write the Fokker-Planck equation as an equation for the distribution function $W(R, P; \theta, P_\theta, \phi, P_\phi, t)$, which goes to Equation (44) with respect to the Hamiltonian $\hat{H}$. The Smoluchowski equation is then for the distribution function $R(R, \theta, \phi, t)$. From these equations, $\bar{\psi}(t)$ can be determined. The center of mass coordinates are special since the interaction terms go to zero when their relative separation is $> \xi$.

Due to the symmetries of this special case, certain simplifying relationships exist among the elements of the interaction matrix Γ defined by Equation (45). The potential ϕ describing the interaction of the ellipsoidal Brownian particle with a bath particle has the form

$$\phi = \phi(x'^2 + y'^2 + z'^2). \tag{99}$$

We consider the radius of the bath particle to be much smaller than the dimensions of the Brownian particle, and thus treat the bath particles as point particles for simplicity. The primed coordinates refer to the relative separation of the center of mass of the Brownian and bath particles, referred to a rotated coordinate system. They are related to those in a nonrotated Cartesian coordinate system by

$$\underline{X'} = \underline{A} \cdot \underline{X} . \quad (100)$$

where the transformation matrix is given by⁵

$$\underline{A} = \begin{pmatrix} \cos \psi \cos \phi - \cos \theta \sin \phi \sin \psi & , & \cos \psi \sin \phi + \cos \theta \cos \phi \sin \psi & , & \sin \psi \sin \theta \\ -\sin \psi \cos \phi - \cos \theta \sin \phi \cos \psi & , & -\sin \psi \sin \phi + \cos \theta \cos \phi \cos \psi & , & \cos \psi \sin \theta \\ \sin \theta \sin \phi & , & -\sin \theta \cos \phi & , & \cos \theta \end{pmatrix} \quad (101)$$

Note that the potential ϕ is indeed independent of ψ since

$$\begin{aligned} x'^2 + y'^2 &= x^2 (\cos^2 \phi + \cos^2 \theta \sin^2 \phi) + y^2 (\sin^2 \phi + \cos^2 \theta \cos^2 \phi) \\ &+ z^2 \sin^2 \theta + 2(yz \cos \phi - xz \sin \phi) \sin \theta \cos \theta + 2xy \sin \phi \cos \phi \sin^2 \theta \end{aligned} \quad (102)$$

and

$$z' = x \sin \theta \sin \phi - y \sin \theta \cos \phi + z \cos \theta \quad (103)$$

Straight forward calculation leads to the following form for the interaction matrix

$$\Gamma_{ij} = \Gamma_{ji}$$

$$\Gamma = \begin{pmatrix} \Gamma_{xx} & \Gamma_{xy} & \Gamma_{xz} & 0 & 0 \\ \Gamma_{yx} & \Gamma_{yy} & \Gamma_{yz} & 0 & 0 \\ \Gamma_{zx} & \Gamma_{zy} & \Gamma_{zz} & 0 & 0 \\ 0 & 0 & 0 & \Gamma_{\theta\theta} & 0 \\ 0 & 0 & 0 & 0 & \Gamma_{\phi\phi} \end{pmatrix},$$

where the non zero matrix elements are

$$\Gamma_{xx} = 4(\cos^2 \phi + \cos^2 \theta \sin^2 \phi) \langle \phi_1^2 x' x' \rangle + 4 \sin^2 \theta \sin^2 \phi \langle \phi_2^2 z' z' \rangle \quad (104a)$$

$$\Gamma_{yy} = 4(\sin^2 \phi + \cos^2 \theta \cos^2 \phi) \langle \phi_1^2 x' x' \rangle + 4 \sin^2 \theta \cos^2 \phi \langle \phi_2^2 z' z' \rangle \quad (104b)$$

$$\Gamma_{zz} = 4 \sin^2 \theta \langle \phi_1^2 x' x' \rangle + 4 \cos^2 \theta \langle \phi_2^2 z' z' \rangle \quad (104c)$$

$$\Gamma_{xy} = \Gamma_{yx} = 4 \sin^2 \theta \sin \phi \cos \phi (\langle \phi_1^2 x' x' \rangle - \langle \phi_2^2 z' z' \rangle) \quad (104d)$$

$$\Gamma_{xz} = \Gamma_{zx} = -4 \sin \theta \cos \theta \sin \phi (\langle \phi_1^2 x' x' \rangle - \langle \phi_2^2 z' z' \rangle) \quad (104e)$$

$$\Gamma_{yz} = \Gamma_{zy} = 4 \sin \theta \cos \theta \cos \phi (\langle \phi_1^2 x' x' \rangle - \langle \phi_2^2 z' z' \rangle) \quad (104f)$$

$$\Gamma_{\theta\theta} = \Gamma_{\phi\phi} / \sin^2 \theta = 4 \langle (\phi_1^2 - \phi_2^2)^2 \rangle (z' x') (z' x') \quad (104g)$$

Note that

$$\Gamma_{xz} = -\Gamma_{xy} \frac{\cos \theta}{\sin \theta \cos \phi}$$

$$\Gamma_{yz} = \Gamma_{xy} \frac{\cos \theta}{\sin \theta \sin \phi}$$

We have defined

$$\phi_1 = \frac{\partial \phi}{\partial (x'^2 + y'^2)} \quad (105a)$$

$$\phi_2 = \frac{\partial \phi}{\partial z'^2} \quad (105b)$$

$$\langle \phi_1^2 x' x' \rangle = \int_0^\infty d\tau \frac{\partial \phi_1}{\partial (x'^2 + y'^2)} x' e^{L \circ \tau} x' \frac{\partial \phi_1}{\partial (x'^2 + y'^2)} \quad (106)$$

et.sim. for $\langle \phi_2 x' x' \rangle$, etc., and sums over all bath particles are implied.

The inverse matrix Γ^{-1} which enters the Smoluchowski equation for R [Equation (96)] can now be obtained in terms of the matrix elements Γ_{ij} . The result is

$$\Gamma_{ij}^{-1} = \Gamma_{ji}^{-1}$$

$$\Gamma^{-1} = \begin{pmatrix} \Gamma_{xx}^{-1} & \Gamma_{xy}^{-1} & \Gamma_{xz}^{-1} & 0 & 0 \\ \Gamma_{yx}^{-1} & \Gamma_{yy}^{-1} & \Gamma_{yz}^{-1} & 0 & 0 \\ \Gamma_{zx}^{-1} & \Gamma_{zy}^{-1} & \Gamma_{zz}^{-1} & 0 & 0 \\ 0 & 0 & 0 & \Gamma_{\theta\theta}^{-1} & 0 \\ 0 & 0 & 0 & 0 & \Gamma_{\phi\phi}^{-1} \end{pmatrix},$$

The matrix elements are as follows:

$$\Gamma_{xx}^{-1} = (\Gamma_{yy} \Gamma_{zz} - \Gamma_{yz}^2) / D \quad (107a)$$

$$\Gamma_{yy}^{-1} = (\Gamma_{xx} \Gamma_{zz} - \Gamma_{xz}^2) / D \quad (107b)$$

$$\Gamma_{zz}^{-1} = (\Gamma_{xx} \Gamma_{yy} - \Gamma_{xy}^2) / D \quad (107c)$$

$$\Gamma_{xy}^{-1} = \Gamma_{yx}^{-1} = -(\Gamma_{xy} \Gamma_{zz} - \Gamma_{yz} \Gamma_{xz}) / D \quad (107d)$$

$$\Gamma_{xz}^{-1} = \Gamma_{zx}^{-1} = -(\Gamma_{xy} \Gamma_{yz} - \Gamma_{yy} \Gamma_{xz}) / D \quad (107e)$$

$$\Gamma_{yz}^{-1} = \Gamma_{zy}^{-1} = -(\Gamma_{xx} \Gamma_{yz} - \Gamma_{xz} \Gamma_{xy}) / D \quad (107f)$$

$$\Gamma_{\theta\theta}^{-1} = \Gamma_{\phi\phi}^{-1} \sin^2 \theta = 1 / \Gamma_{\theta\theta},$$

where

$$D = \Gamma_{xx} (\Gamma_{yy} \Gamma_{zz} - \Gamma_{yz}^2) - \Gamma_{xy} (\Gamma_{zz} \Gamma_{xy} - \Gamma_{xz} \Gamma_{yz}) + \Gamma_{xz} (\Gamma_{xy} \Gamma_{yz} - \Gamma_{xz} \Gamma_{yy})$$

V. FUTURE WORK

As mentioned in the introduction, our work on the hydrodynamics of multicomponent systems, begun in the first year, will be completed and documented.

The treatment of Brownian motion in nonequilibrium baths will be pursued using techniques similar to those used in equilibrium systems. The interaction between bath and Brownian particle motions will be studied. The relationship between the Fokker-Planck and Smoluchowski equations

in nonequilibrium baths and the equations of hydrodynamics will be investigated. Appropriate hydrodynamic boundary conditions on the surfaces of the Brownian particles will be obtained.

The dependence of the transport coefficients (viscosity, diffusion constant, etc.) on the concentration of the suspended particles will be studied. In particular, the origin of the surprising experimental result that the diffusion constants of some large molecules are independent of concentration as a result of cancellation between the effects of direct interaction and the hydrodynamic interaction among the particles will be investigated. The distinction between short-time and long-time diffusion constants will be pursued.

As a result of these investigations, the appropriate equations for the transport of suspended particles or solutes will be obtained together with appropriate approximations for the dissipative coefficients in these systems.

REFERENCES

1. P. Mazur and I. Oppenheim, *Physica* 50, 241 (1970); I. Oppenheim and J. M. Deutch, *J. Chem. Phys.* 54, 3547 (1971).
2. N. G. van Kampen and I. Oppenheim, *Physica* 138A, 231 (1986).
3. V. Romero-Rochin and I. Oppenheim, *J. Stat. Phys.* 53, 307 (1988).
4. J. M. Deutch and I. Oppenheim, *Faraday Discussions, Chem. Soc.* 83, 1 (1987).
5. H. Goldstein, in Classical Mechanics (Addison-Wesley, 1965) pp. 107-109.

END

DATE
FILMED
6/10/92