PROGRESS REPORT

for

Grant No. DE-FG02-90ER20009

"Post Transcriptional Regulation of Chloroplast Gene Expression by Nuclear Encoded Gene Products"

For the Budget Period

6/1/91 - 5/31/92

Principal Investigator

Michael R. Kuchka

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

PROGRESS REPORT FOR GRANT NO. DE-FG02-90ER20009 - "POST-TRANSCRIPTIONAL REGULATION OF CHLOROPLAST GENE EXPRESSION BY NUCLEAR ENCODED GENE PRODUCTS" - Principal Investigator - Michael R. Kuchka, Ph.D.

The following is a review of research accomplished in the first two years of funding for the above mentioned project. The work performed is a molecular characterization of nuclear mutants of Chlamydomonas reinhardtii which are deficient in different stages in the post-transcriptional expression of a single chloroplast encoded polypeptide, the D2 protein of Photosystem II. Our long-term goals are to understand the molecular mechanisms by which nuclear gene products affect the expression of chloroplast genes. Specifically, we wish to understand how specific nuclear gene products affect the turnover rate of the D2 encoding mRNA (psbD), how other nuclear encoded factors work to promote the translation of psbD mRNA and/or stabilize the D2 protein, and what the role of the D2 protein itself is in Photosystem II assembly and in the control of expression of other chloroplast genes. This progress report will be organized into four major sections concerning (I) The characterization of nuclear mutants affected in D2 translation/turnover, (II) The study of trans-acting factors which associate with the 5' end of the psbD mRNA, (III) In vitro mutagenesis of the psbD gene, and (IV) Additional studies.

I. D2 TRANSLATION/TURNOVER MUTANTS

This research is described in detail in the enclosed preprint and so will be discussed only briefly here. Four independently isolated nuclear mutants of Chlamydomonas reinhardtii were shown to have normal levels of psbD mRNA but lack D2 protein because of a block in D2 translation and/or an increase in D2 protein turnover. Genetic analysis showed that these four mutants identify at least two and possibly three different nuclear genes. Two genes were mapped genetically. The psbD mRNA was determined to be associated with large classes of polysomes in all mutants, suggesting that late stages in translation elongation and/or D2 protein turnover is affected in these strains. Phenotypic revertants have been generated from all mutant backgrounds. We are characterizing these revertants in an attempt to identify additional genes whose products interact with the original gene products to promote D2 translation and/or stabilization. We have focused on a particular suppressor derived from one of the revertants. This suppressor mutation suppresses the phenotypes of three mutants affected in two different genes. We would like to understand the nature of suppression in these strains and the exact function of the suppressor gene product. Our plans for the continued examination of these mutants and revertants are presented in the Continuation Request section of this proposal.

II. PSBD TRANSCRIPT STABILITY MUTANT

We have described a nuclear mutant of Chlamydomonas reinhardtii which transcribes but fails to accumulate stably the psbD mRNA (Kuchka et al., 1989). Previous experimentation suggested that the 5'

untranslated region of psbD mRNA is an important cis element in determining message half-life (Kuchka et al., 1989; Rochaix et al., 1990). We are interested in characterizing proteins which bind specifically to this part of the psbD transcript. Some of these proteins may confer psbD message stability by their binding. We have generated by polymerase chain reaction a 220 base pair sequence of DNA which encodes the full 5' untranslated region and the first several codons of psbD. This sequence was cloned into an in vitro transcription vector which is used to generate labeled transcripts. We have used these labeled RNAs in mobility shift assays and uv-crosslinking experiments with total chloroplast proteins derived from mutant and wild type cells. These studies are still in progress. We are now planning to compare proteins which bind to the 5' untranslated region of psbD with those which bind to the complete psbD message. These experiments are described in the Continuation Request section of this proposal.

III. IN VITRO MUTAGENESIS OF PSBD

In characterizing mutants of Chlamydomonas which are deficient in D2 synthesis, we have noticed that these strains differ significantly in their ability to assemble stably the Photosystem II (PS II) complex. Mutants affected in psbD mRNA stability are deficient in all PS II proteins, even peripheral membrane proteins involved in oxygen evolution (the OEE proteins). Other mutants, including one with an internal duplication within the coding region of the chloroplast psbD gene, accumulate these OEE proteins in wild type quantities. We have wondered to what extent these differences reflect variations in the levels of D2 protein in these mutants or to what extent they reflect additional roles of the mutant gene products. In part, to distinguish between these possibilities we have undertaken an approach to generate a true psbD null mutant by in vitro mutagenesis and transformation. We have constructed a plasmid which is deleted in Chlamydomonas chloroplast DNA and disrupted in the psbD gene by insertion of the bacterial aadA sequence which confers spectinomycin and streptomycin resistance to transformed cells (Goldschmidt-Clermont, 1991). This construct was introduced into Chlamydomonas cells by particle bombardment and putative transformants which are antibiotic resistant have been identified. These putative transformants show abnormal chlorophyll fluorescence kinetics, a characteristic of photosynthetic mutants. Plasmid construction was performed in our laboratory. Transformation and fluorescence kinetics characterizations were carried out at the University of Geneva by Dr. Michel Goldschmidt-Clermont. Subsequent analysis of putative transformants is currently underway in our lab (see Continuation Request section of proposal). This collaboration was arranged because our lab presently does not have a particle gun for transformation nor the proper equipment for fluorescence measurements. We plan to continue this productive collaboration. A true psbD null mutant will be useful in elucidating the role of the D2 protein in PS II complex assembly and will also help clarify D2's implied role in regulating the expression of other photosynthetic proteins, including the related D1 polypeptide of PS II (Erickson et al., 1988).

IV. ADDITIONAL STUDIES

In the year one progress report I submitted, I discussed our need for a D2 specific antibody. I described our attempts to clone psbD into an expression vector for the generation of sufficient quantities of the D2 protein for antibody production. These experiments are currently underway in the laboratory. Appropriate psbD restriction fragments have been made and constructs are being examined. In the interim period, we received a D2-specific antiserum raised against the D2 protein from barley chloroplasts (a kind gift of Dr. John Mullet). We have tested this antiserum by immunoblot analysis and it cross-reacts well and specifically with the D2 protein of Chlamydomonas. We plan to use this antiserum for immunoprecipitation and immunoblotting experiments (see Continuation Request section). Nevertheless, we have a limited quantity of this antiserum and so find it necessary to continue the pursuance of our initial approach of cloning psbD into an expression vector and isolating purified D2 protein. Antibodies resulting from this work will be an invaluable tool in our research.

REFERENCES

- Erickson, J. M., Rahire, M., Malnoe, P., Girard-Bascou, J., Pierre, Y., Bennoun, P., and Rochaix, J.-D. (1986) EMBO J., 5: 1745.
- Goldschmidt-Clermont, M. (1991) Nucl. Acids Res., 19: 4083.
- Kuchka, M. R., Goldschmidt-Clermont, M., van Dillewijn, J., and Rochaix, J.-D. (1989) Cell, 58: 869.
- Rochaix, J.-D., Goldschmidt-Clermont, M., Choquet, Y., Kuchka, M., and Girard-Bascou, J. (1990) in Plant Molecular Biology (ed. R. G. Hermann) Plenum Press, NY.

END

**DATE
FILMED**

6 / 29 / 92