YEAR 1 PROGRESS REPORT FOR DE-FG02-91ER61247

Kevin Harrison
Wally Broecker
5/21/92

THE IMPACT OF CHANGING LAND USE, NITRATE DEPOSITION AND CO₂
FERTILIZATION ON SOIL CARBON STORAGE

Outline

- Abstract
- Introduction
- Changing Land Use
 - Strategy
 - Materials and Methods
 - Results/Discussion
 - Future Research
 - Summary
- Nitrate deposition and C/N ratios
 - Strategy
 - Results/Discussion
 - Future Research
 - Summary
- CO₂ Fertilization
 - Strategy
 - Results/Discussion
 - Future Research
 - Summary
- Acknowledgements

ABSTRACT

This research strives to assess the impact of changing land use, nitrate deposition and CO₂ fertilization on soil carbon storage. Our motivation is that this reservoir is the most likely candidate for the so-called missing carbon sink. We are working on several aspects of this problem by measuring carbon content, nitrogen content and radiocarbon ratios in paired soil samples from neighboring sites, to determine the impact of land use on soil carbon inventories and turnover times. We are also gathering information on how the C/N ratios in soils vary with climate and changing land use, in an effort to estimate how much carbon has been sequestered as a result of atmospheric fallout of NH₄OH and HNO₃. Finally, we are developing a soil greening model that uses CO₂ growth-enhancement results and bomb radiocarbon-based estimates of soil carbon inventory response times.

MASTER*Wally*

INTRODUCTION

Houghton et al. (1990) estimate that 32 GtC have been released by changing land use between 1958 and 1978. In this same time period, Marland and Rotty (1984) estimate that burning fossil fuel has released 76 GtC. The oceans have absorbed 28 GtC (Broecker, et al., 1979); and the atmosphere has increased by 43 GtC (Bolin, 1986). This results in a "missing sink" of 37 GtC. Sarmiento and Sundquist (1992) cite annual values for the 1980's, with the following representative figures for the decade:

Sources: Fossil fuel (5.4 ± 0.5)

Deforestation (1.6 ± 1.0)

Sinks: Atmosphere (3.2 ± 0.1)

Oceans (2.0 ± 0.8)

Missing sink = (1.8 ± 0.4) GtC/yr.

In what ways could the biosphere be sequestering carbon in response to man?

Possibilities include changing land use, nitrate deposition and CO₂ fertilization. Carbon losses upon cultivation have been documented by Post and Mann (1990) and Schlesinger (1986). Houghton et al. (1980) estimate a 90 percent soil-carbon recovery from disturbances such as cultivation, based on a handful of references. The recovery of abandoned, managed ecosystems has not been well-documented (Schlesinger, 1986). Bazzaz and Fajer (1992) suggest a CO₂ fertilization factor of 0.35 for well-watered and fertilized plants grown in carbon-dioxide chambers. However, they doubt that this fertilization factor could apply in natural settings, because of limited nutrients, limited water and species competition. When fertilization factors for natural settings become available,

they can be used in conjunction with soil-carbon residence time spectra to calculate how much carbon gets sequestered because of carbon-dioxide fertilization.

Nitrate deposition can potentially lead to the sequestration of large amounts of carbon. Duce et al. (1991) suggest that 50×10^{12} gN/year of oxidized nitrogen and 54×10^{12} gN/year of reduced nitrogen get deposited in the land and oceans. The amount of carbon sequestered by this 104×10^{12} gmN/year depends on where it ends up. C/N ratios range from 160 in trees, to 15 in soil humus, to 8 in aquatic-marine organic matter. Since the soil humus and aquatic-marine organic sinks dominate, the carbon sequestered is likely to fall in the range of 0.8 to 1.5 GtC, but it could be as high as that required to explain the value of 1.8 GtC cited by Sarmiento and Sundquist (1992). Further, Suzanne Bailey (1992, personal communication) has found C/N ratios ranging from 50-100 for peat, and has shown that fertilizing peat bogs with N doesn't alter the C/N ratio. Kauppi et al. (1992) concur, citing nitrate deposition as the primary candidate for increasing European biomass by 25 percent from 1970 to 1990.

CHANGING LAND USE

Strategy

In this study, the effects of changing land use have been investigated by looking at paired plots of land having different uses (table 2). One aspect involved checking to see if carbon lost from shallow layers of soil was being re-deposited in deep soil. Hence, we sampled as deeply as possible (1-3 meters). We chose to work at the Duke Experimental Forest to compare a native forest and recovering forest ecosystem. About 32 percent of the US is forested, and about 80 percent of the Eastern forests have been cleared for cultivation, most of which has been abandoned. We wanted to document the rate of recovery.

The second site we selected was in Kansas: the Konza Long-Term Ecological Research Site. It consisted of native grassland (extremely rare, since most grassland has been cultivated), currently-farmed land, and a recovering site that had been seeded with brome, a non-native grass. Pasture and rangeland account for 26 percent of the US, with cropland representing 21 percent. Hence, the forested and grassland sites represent 79 percent of the US land area.

Italy was selected as a grassland site with land-use histories extending over 100 years and a markedly different climate than Kansas. The Canadian sites provided a platform to study forest and grassland ecosystems in a cooler climate than the previous sites, and to assess how slopes can impact carbon reservoirs. The Canadian sites consisted of arid grassland and sub-humid boreal forest.

Materials and Methods:

Soil samples were oven dried at 60 degrees to a constant dry weight, ground and sieved (<2mm) before analysis. Bulk densities were calculated from a fixed-volume sample and weight. Radiocarbon measurements and Cs-137 analyses were performed following Trumbore (1988). Nitrogen and carbon were measured using the procedures described by Verardo et al. (1990). The soil was acid-washed before being oxidized for graphite targets to remove any carbonates.

Results/Discussion

The results of the Duke Experimental Forest are plotted in figures 1. The recovering site had 15% more C and 8% more N than the native site, presumably because N-fixing scrub replenished the N that had been lost through prior cultivation. Once the N was fixed and added to the soil, the carbon was efficiently sequestered by the pines.

Prior to 1931, the recovering site was cultivated, abandoned and scrub was allowed to grow. In 1931, conifers were planted and soil samples were collected in 1991. The paired native

site was next to the recovering site, had similar parent material and topography and consisted of mature deciduous hardwoods.

Using the C and N loss estimates of Post and Mann (1990) one can estimate the pre-recovery carbon (58,700 kg C/ha) and nitrogen (4,300 kg N/ha). How did the recovering site gain 635 kg N/ha? Possible mechanisms include residual fertilizer, N-fixation and N deposition. The amount of residual fertilizer is unknown. A reasonable value for N deposition is 8.8 kg N/ha/yr (Cole and Rapp, 1981), but some of this would be lost through denitrification (2 kg N/ha/yr; Schlesinger, 1991) and leaching (3.5-4.3 kg/ha/yr; Cole and Rapp, 1981). The upper limit on the amount of N deposited by rain and dry fallout would be about 175 kg/ha since abandonment. Both the native and the recovering site would receive this nitrogen. N-fixation could deposit up to 650 kg N/ha. If symbiotic N fixation from scrub occurred at a rate of 100 kg N/ha/yr and asymbiotic fixation occurred at 2.5 kg N/ha/yr (Schlesinger, 1991), we estimate it would only take 5 years of scrub growth and 60 years of asymbiotic fixation to produce 650 kg N/ha.

Once the N has been deposited, the subsequent pines efficiently use N to build up soil carbon reservoirs. Although deciduous trees produce about 20% more litter than conifers (Cole and Rapp, 1981), conifers have a 100% higher C/N ratio (Cole and Rapp, 1981) and have 50% more lignin. The higher C/N ratio means that more C is sequestered per N and the higher lignin makes the litter more recalcitrant. The Cs-137 profiles were very similar for the two sites, with slightly more being present in the native site. The recovering site had a higher soil-radiocarbon value, indicating that it was sequestering more carbon during the bomb spike than the native site. If both forests had achieved steady state, we would have expected the conifer forest to have lower C-14 values, because of the longer residence time for its humus, a function of the higher lignin content of its litter.

The observed 115% recovery at the Duke Experimental Forest sharply contrasts the 90% recovery estimated by Houghton (1980). Clearly, more effort needs to be directed toward studying recovering ecosystems. Recovering ecosystems can be engineered to sequester more

carbon simply by allowing N-fixing scrub to grow for a few years. This contrasts current forestry practice, where every effort is made to eliminate the scrub.

The results from the Kansas study are plotted in figure 2. Although the abandoned site had the least carbon, it looked like carbon was starting to accumulate in the upper ten centimeters and its nitrogen content was about the same as the native site. Perhaps, the recovering vegetation has a lower C/N ratio than the native grasses. The farmed land had low surface carbon, but deep carbon wasn't as depleted, so its carbon inventory fell between the native and abandoned sites. The soybean crops helped maintain N levels in the farmed site because they are able to fix atmospheric N.

The results from Italy are listed in figure 3. Carbon lost from the shallow, disturbed soil appears to have been spread downward by plowing. The N profile shows a similar trend. The Cs-137 was significantly reduced in the cultivated site, possibly by the same mechanism as Kansas. These soil samples and supporting data were obtained from Dr. Costantini. More data should be forthcoming. Land use records extend back over 100 years.

The results from Canada are listed in figure 4. Canadian samples and supporting data were obtained from Jeff Schoenau of the University of Saskatchewan. They are described in his PhD. thesis. We have compared native and cultivated forest and grassland sites. The sub-arid grassland was ungrazed and undisturbed by man, while the disturbed land had been cultivated for 70 years, using a cereal fallow rotation. The lowest grassland site was frequently submerged and anoxic. The forest site consisted of an aspen forest with various legumes and grasses making up the lower story. The cultivated site had had the litter layer scraped off and had been planted for barley. The cultivated grassland had about 50% of the native carbon in the upper and mid-slope positions. However, the disturbed land had more carbon in the lower site. Carbon at the native sites increased downslope, but not as dramatically as the cultivated site. The anoxic conditions of the cultivated site may have further enhanced the buildup of carbon. The native forested site had carbon distribution similar to native grassland site from top slope to bottom slope, but the

cultivated site showed increasing levels of carbon downslope. Hence, roots seem to stabilize soil transport, while cultivation accelerates it.

Future research

We plan to confirm the Duke Experimental Forest results to make sure our observations are not due to lateral inhomogeneity. Then, we would like to look at currently cultivated land and recovering land where scrub growth has been inhibited. In Kansas, we would like to look at currently cultivated land with the non-N fixing crops and look at recovering grassland that has native vegetation instead of brome. In Canada, we would like to have looked at land that wasn't sloped and at some recovering sites, both with more closely-spaced sampling intervals. In Italy, we are waiting for bulk density data; we also plan to obtain samples at a recovering site and to sample at more closely spaced sampling intervals. The radiocarbon data for these sites will be discussed in greater detail in subsequent sections.

Summary:

The Duke site illustrates how N-fixing scrub growth enhances recovery. The Kansas site demonstrates how N-fixing crops can build up soil N and that non-native vegetation can hinder carbon recovery if it has lower C/N ratios than the native vegetation. In Canada, the sites showed the impact slope can have on carbon storage and provided a climatic contrast for the Duke and Kansas sites. These results can also be compared with the values of Post and Mann (1990) and Schlesinger (1986) for estimating soil carbon losses upon disturbance. Both the Kansas and Italy results fall in the range found by Post and Schlesinger. The Canadian results don't apply because of the slope. The Duke recovery result of 115% contrasts with the 90% recovery used by Houghton et al (1980).

NITRATE DEPOSITION

Strategy

Nitrate deposition can sequester between 16.6 and 0.8 GtC, annually depending on whether it's assimilated by trees, deposited in soils or incorporated into marine humus. The actual

value depends on where the nitrate is deposited, how much is assimilated into organics and the C/N ratio of these organics.

Results/Discussion

To estimate the impact of nitrate deposition, one needs to know the temporal input of N and how ecosystems respond to the additional N. The carbon cycling time determines the rate of carbon sequestration and the C/N ratio determines the amount.

We measured C/N ratios in Duke, Kansas, Italy and Canada to see how C/N ratios vary with climate and changing land use (figure 5). The C/N ratio was higher in the Duke forest than the Canadian forest; the Kansas grassland had higher C/N ratios than the Canadian grassland, while the Italian grassland had the lowest values. The forests had higher C/N ratios than the grasslands.

Future Research

We plan to estimate the impact of nitrate deposition on soil carbon storage by using the temporal input of N, by allocating the distribution of N between vegetation and humus, and by using the C/N ratios to determine the amount of carbon sequestered. We plan a similar approach for the marine environment.

Using acid rain maps, we will estimate the contemporary nitrate input, project N input over time by looking at temporal variation from existing stations, and scale the global results accordingly.

Allocating the N between vegetation and soil presents the next challenge. N's impact will be greater if kept in the vegetation (C/N=50-160) compared to soil (C/N=15). Some N deposited on land will be transported by rivers into the ocean, which, along with atmospheric deposition to the oceans, would sequester carbon at a ratio of 8 to 1.

Summary

Atmospheric deposition of N could sequester large amounts of C, depending on where it ends up. For example, trees would sequester the most carbon per nitrogen. We have measured

C/N ratios in soil and plan to model global nitrate deposition and concomitant carbon sequestering using available acid rain deposition distribution data, vegetation maps and observed C/N values.

CO₂ FERTILIZATION

Strategy

Soil carbon residence times control the humus' response to perturbations, such as changing land use and nutrient input. Here we discuss how increasing atmospheric CO₂, a nutrient increase, affects soil carbon storage (figure 6).

The 25% atmospheric carbon dioxide increase since 1850 may have increased plant growth and soil carbon input if other nutrients are not limiting. Bazzaz and Fajer (1992) suggest a 0.35 CO₂ fertilization factor based on chamber experiments. However, they doubt this factor applies to natural settings, where species competition and water and nutrients may limit growth.

Results/Discussion

Translating soil radiocarbon measurements into residence time provides an exciting challenge. We define R as the measured C-14/C ratio divided by the C-14/C ratio of 1850 wood. $R_{\text{atmosphere}}$ was diminished 2.5% between 1850 and 1950 by burning fossil fuel and greatly enhanced by atmospheric nuclear bombs between 1850 and the present. This atmospheric signal propagates through soil carbon reservoirs as a function of soil carbon residence times (figure 7). For example, we plotted R_{soil} vs residence time for 1985 (figure 8). If soil carbon consisted of one component, the residence time could be obtained directly. However, a slow cycling component lowers the observed ratio. We have estimated the size and residence time of these components by looking at pre- and post-bomb native soils.

The pre-bomb soil values range from 0.8 to 0.9 with an average value of 0.85, which translates into a 1300-year apparent residence time. This leads to expected R_{soil} of slightly above 0.9 in 1985, contrasting observations exceeding 1.1 (figure 9).

We estimated a range of possible residence times for the fast cycling component that brackets the observed native soil radiocarbon values. Ten-year and 100-year cycling times provide upper and lower limits with 30-year cycling times providing the best fit of the observations (figure

10). Determining the fraction of soil having the 30-year residence time is the next step in our calculations.

Having a pre-bomb value of 0.85 and knowing the residence time of the fast cycling component does provide broad limits for the fast-cycling and slow-cycling fractions. For example, if the slow component has a 1500-year cycling time, 12% of the soil carbon would be fast-cycling and 88% would be slow-cycling. If the slow part had a 3000-year cycling time, 49% would be fast and 51% slow. For "dead" radiocarbon, 85% would be fast and 15% slow. Deep radiocarbon measurements further constrain the slow cycling component. These values range from 0.4 to 0.7 and average 0.575, which corresponds to a 4,500 year cycling time. Hence, 65% would be fast and 35% would be slow. Therefore, the active soil carbon reservoir increased by up to 5% by 1985 assuming a 30-year residence time and a 0.35 fertilization factor (figure ?). What is the size of the active reservoir. The upper limit would be the total carbon soil reservoir (1500 GtC) times the portion of soil carbon with a fast-cycling time (0.65) or 975GtC. The lower limit would be 50 GtC, the estimated annual litter input (Schlesinger, 1991). Hence, we estimate that the fast cycling reservoir size is 500 GtC. If we assume a CO₂ fertilization factor of 0.35 and use a 30-year cycling time, the this reservoir would increase by 5% or 25GtC (figure 6).

Future Research

In the next year, we would like to improve our model by estimating fast cycling residence times for different types of vegetation and for different climates. Toward this end, we are compiling existing radiocarbon values for various climes and flora and continue to expand our own experimental measurements (figure 11 and table 1). Our model translates radiocarbon values, which cannot be compared from one year to the next, to residence times, which can be compared. We hope to improve our understanding of the slow-cycling part by looking at deep soils, where the fast-cycling soil carbon component has decayed. We hope to improve our understanding of the slow and fast cycling components' sizes and residence times to further validate our model by examining pre- and post- bomb archived soil time series in collaboration with Dan Richter and Mac Post.

Summary

Soil carbon cycling times control humus response to perturbations. We are using the infiltration of bomb radiocarbon to estimate these residence times in conjunction with an integrated modelling and experimental approach. Carbon cycling times could help predict future changes in the soil carbon inventory.

ACKNOWLEDGEMENTS:

Jim Simpson and Inez Fung helped this project evolve with their insightful questions, patient discussions and modelling insights. Eric Haney and Peter Harrell helped select the Duke Experimental Forest field sites and did considerable work in constructing the two meter deep soil pits. Bill Schlesinger, Dan Richter, and Norm Christenson hosted my visit to Duke, helped select field sites and shared their expertise in many discussions that helped me clarify my research objectives. Jeff Schoenau hosted my visit to the University of Saskatchewan, provided soil samples and supporting data, and made valuable suggestions for further research. Michel Ransom hosted my visit to Kansas State University, and in conjunction with Bill Weilmeuler, helped me select field sites and use the soil probe truck. We thank Dr. Edoardo A. C. Costantini (Istituto Sperimentale per lo Studio) for helping us in Italy by providing soil samples on short notice. Nancy Mager helped make numerous text, budget and figure revisions and ensured smooth and rapid communication between the various investigators associated with this project. Rachel Oxburgh helped considerably with the modelling. Elaine Mathews and Katy Prentice provided a seemingly endless supply of answers and references to the multitude of questions raised by our research. DOE funded this research. We thank them all.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

TABLE 1

location	C	N	C/N	Rsoil*	Cs-137	T °C	P(mm)
	<u>% by mass</u>		<u>molar</u>		<u>dpm/cm2</u>		
Italy native	.9%	.11%	9.6	—	31	14	1792
Italy cultivated	.6%	.08%	8.8	1.03(0-10cm)**	11.5	14	1792
	<u>kg/ha</u>	<u>kg/ha</u>					
Kansas native	48656	3948	14.4	1.15(0-2cm)**	10.73	13	732
Kansas cultivated	30617	3664	9.8	1.13(0-2cm)**	11.9	13	732
Kansas recovering	26100	4047	7.5	1.01(0-2cm)**	5.46	13	732
Duke native	76200	4580	19.4	1.12(0-10cm) 0.98(20-30cm)	22.32	15	1075
Duke recovering	88600	4940	20.9	1.15(0-10cm) 1.02(20-30cm)	20.16	15	1075
Canada grassland NT	81460	8629	11.0	1.22(0-15cm)**	5.8	7	300-350
NM	110340	11808	10.9	—	4.83		
NB	141240	14107	11.7	—	3.16		
Canada grassland CT	50850	6229	9.53	—	4.83		
CM	64570	7178	10.5	—	5.66		
CB	170310	17352	11.5	—	13.50		
Canada forest NT	112960	9247	14.2	—	5.16	6	400-450
NM	127480	10492	14.2	—	5.33		
NB	119270	9636	14.4	—	8.66		
CT	54590	5370	11.9	1.03(0-7cm)**	2.5		
CM	67190	6916	11.3	—	4.33		
CB	88300	8970	11.5	—	6.00		

*Rsoil = $\frac{C-14/C}{C-14/C \ 1850}$

** preliminary

TABLE 2

<u>site</u>	<u>crop type</u>	<u>recovering vegetation</u>	<u>native vegetation</u>
Duke	--	Loblolly pine	Oak
Kansas	soybeans	brome	grass
Canada forest	cereal/fallow	--	Aspen
Canada grassland	barley	--	grass
Italy	crop	--	grass

FIGURE CAPTIONS

Figure 1: Plots of carbon, nitrogen and Cs-137 for the Duke Experimental Forest are presented. The native site had slightly more carbon in the 0-10cm interval, but the recovering site had more carbon from 20-40cm. Slight differences below 40cm tend to cancel. The nitrogen profile resembles the carbon profile. Cs-137 data suggests that the two sites had similar erosional history since atmospheric atomic bomb testing.

Figure 2: Kansas carbon, nitrogen, and Cs-137 are plotted for native, farmed and recovering sites. The native site had the most carbon. The recovering site had the least carbon, but carbon appears to be building up in the shallowest layers of recovering soil. The farmed site had high N levels because of the N-fixing soybean crops. The recovering site has about the same N as the disturbed site. Cs-137 is depleted for the cultivated site.

Figure 3: Italy carbon, nitrogen and Cs-137 data are plotted for cultivated and native land. Carbon lost from the shallow layers of disturbed soil appears to be deposited deeper in the soil. Nitrogen shows a similar trend. Cs-137 is depleted in the cultivated soil.

Figure 4: Canadian carbon, nitrogen and Cs-137 are plotted for paired cultivated and native grassland and forested sites. These sites were selected to see how a slight slope would affect soil carbon storage. Low, middle and high refer to the slope position of each of the sites. Carbon migrated downhill and cultivation accelerated this process.

Figure 5: Variations in C/N ratios for Duke, Kansas and Italy are plotted against depth. At Duke, values tend to decrease with depth. In Kansas, the farmed site ratios are pretty constant, but the native and recovering sites tend to decrease with depth. In Italy, the ratios tend to slightly increase with depth.

Figure 6: We plotted residence time vs percent carbon reservoir increase for hypothetical soils to illustrate how the residence time affects carbon increase with a given CO₂ fertilization factor. We used a fertilization factor of 0.35 and multiplied this times the CO₂ increase from 1850 to 1985. The carbon pools having the shortest cycling time had an almost 9% increase in reservoir size, contrasting the 1000-year residence time carbon, which increased by less than 1%.

Figure 7: C-14 ratio is plotted against time, from 1950 to 1995. The ratio ranges from 0.8 to almost 1.9. The atmospheric ratio shows the greatest increase, which stems from atmospheric atom bomb tests. This bomb spike is propagated through the soil carbon having different residence times. The shortest soil carbon residence time plotted, 3 years, shows the greatest perturbation. However, even at this short residence time, the atmospheric signal is significantly attenuated. The longest residence time of 1000 years shows the least change.

Figure 8: The soil radiocarbon ratio is plotted against soil carbon residence time for the year of 1985. The ratio ranges from slightly above 0.9 to almost 1.35. The maximum corresponds to a 10 year residence time. Carbon having shorter residence times have lower ratios because they more closely follow the atmospheric bomb signal, which has rapidly attenuated. Carbon having longer residence times are still lagging the bomb signal.

Figure 9: Rsoil carbon experimental values are plotted against time from 1950 to 1991. The native values tend to be higher than the cultivated values, although considerable variability is present. The lowest values tended to be from cultivated soil, with wet cultivated soil being bracketed by the native and dry cultivated values. Native values for

soil collected before 1950 were assigned an arbitrary year of 1957 to expand the horizontal scale.

Figure 10: Rsoil carbon for native soil was plotted against years. We superimposed our soil carbon modelling results for 10, 30 and 100-year residence times over this data. Our hypothetical curves were corrected for slow cycling carbon (see text). The 10 and 100-year cycling times tend to bracket the experimental data, with the 30 year cycling time providing the best fit to the data.

Figure 11: Percent modern radiocarbon for Duke is plotted against depth. The recovering sight had the highest values, while both sites radiocarbon values decreased with increasing depth.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7

FIGURE 8

FIGURE 9

FIGURE 10

FIGURE 11

REFERENCES

- Aber, John D. and J.M. Melillo, "Litter Decomposition: measuring relative contributions of organic matter and nitrogen to forest soils." Canadian Journal of Botany, 58, 416-421, 1980.
- Anderson, D.W. and E.A. Paul. "Organo-mineral complexes and their study by radiocarbon dating." Soil Sci. Soc. Am. J., 48, 298-301, 1984.
- Bailey, J.P. R.K. Dixon, T.P. Droessler, S. Henderson, H. Gucinski, M.G. Johnson, G.A. King, R. Leemans, K.G. Mattson, R.P. Neilson, D.L. Phillips, P.E. Schroeder, D.P. Turner and J.K. Winjum, "Response and Feedbacks of Forest Systems to Global Climate Change." Corvallis: EPA, Ed. by G.A. King, J.K. Winjum, R.K. Dixon and L.Y. Arnaut, September, 1990.
- Bazzaz, F.A. and E.D. Fajer. "Plant life in a CO₂-rich world." Scientific American, January 1992, 68-74.
- Becker-Heidmann, P., Liu Liang-wu, H.W. Scharpenseel. "Radiocarbon dating of organic matter fractions of a Chinese mollisol." Z. Pflanzenernahr. Bodenk., 151, 37-39, 1988.
- Becker-Heidmann, P. and H.W. Scharpenseel. "Carbon isotope dynamics in some tropical soils." Radiocarbon, 31, 3, 672-679, 1989.
- Billings, W.D. "The Structure and Development of Old Field Shortleaf Pine Stands and Certain Associated Properties of the Soil." Ecological Monographs, 8, 437-499, 1938.
- Blank, R. R. "Cultivated and Adjacent Virgin Soils in Northcentral South Dakota: I. Chemical and Physical Comparisons." Soil Sci. Soc. Am. J., 53, 1484-1490, 1989.
- Bolin, B., "Changes in land biota and their importance for the carbon cycle." Science, 196, 613-615, 1977.
- Bolin, B., "How much CO₂ will remain in the atmosphere?" In The Greenhouse Effect, Climate Change, and Ecosystems, Scope 29, ed. by B. Bolin, B.R. Doos and J. Jager. New York: John Wiley, pp. 93-155, 1986
- Bopp, Richard, Michael Gross, Huazi Tong, H. James Simpson, Stephen J. Monson, Bruce L. Deck, and Frederick Moser. "A major incident of dioxin contamination: sedimentary New Jersey estuaries." ES & T, 25, 951-956, 1991.
- Botkin, Daniel B., and L.G. Simpson, "Biomass of North American Boreal Forest." Biogeochemistry, 9, 161-174, 1990.
- Bouwman, A.F. (ed.), Soils and the Greenhouse Effect. Proceedings of the International Conference on Soils and the Greenhouse Effect. Wageningen, The Netherlands, Wiley & Chichester.
- Broecker, W. S., T. Takahashi, H. J. Simpson, and T. H. Peng. "Fate of fossil fuel carbon dioxide and the global carbon budget." Science, 206, 409-418, 1979.

Buringh, P., "Organic Carbon in Soils of the World." In The role of the Terrestrial Vegetation in the Global Carbon Cycle, Scope 23, ed. by G.M. Woodell. New York: John Wiley, 1984, pp. 91-109.

Burke, I.C., C.M. Yonker, W.J. Parton, C.V. Cole, K. Flach and D.S. Schimel, "Texture, Climate and Cultivation Effects on Soil Organic Matter Content in U.S. Grassland Soils." Soil Science Society of America Journal, 53(3), 800-805, 1989.

Burke, Ingrid C., William A. Reiners and David S. Schimel, "Organic matter turnover in a sagebrush landscape." Biogeochemistry, 1:11-31, 1989.

Burke, Ingrid C., D.S. Schimel, C.M. Yonker, W.J. Parton, L.A. Joyce and W.K. Lauenroth, "Regional modeling of grassland biogeochemistry using GIS." Landscape Ecology, 4 (1), 45-54, 1990.

Campbell, C.A., E.A. Paul, D.A. Rennie and K.J. McCallum. "Applicability of the carbon-dating method of analysis to soil humus studies." Soil Science, 104, 3, 217-223, 1967.

Chan, Yip-Hoi and J.S. Olson, "Limits on the Organic Storage of Carbon from Burning Fossil Fuels." Journal of Environmental Management, 11, 147-163, 1980.

Dörr, Helmut and K.O. Münnich, "Annual Variations of the ^{14}C Content of Soil CO_2 ." Radiocarbon, 28 (2A), 338-345, 1986.

Duce, R.A., P.S. Liss, J.T. Merrill, et al. "Atmospheric input of trace species to the world oceans." Global Biogeochemical Cycles, 5, 3, 193-259, 1991.

Encyclopedia Britannica, "Rice," vol. 19, p. 283-284, Encyclopedia Britannica, Inc., Chicago, IL, 1956.

Gilbert-Blein, N., G. Marien and J. Evin. "Unreliability of C-14 dates from organic matter of soils." Radiocarbon, 22, 3, 919-929, 1980.

Goh, K.M., T.A. Rafter, J.D. Stout and T.W. Walker. "Accumulation of soil organic matter and its carbon isotope content in a chronosequence of soils developed on aeolian sand in New Zealand." J. Soil Science, 27, 89-100, 1976.

Goh, K.M., J.D. Stout and T.A. Rafter. "Radiocarbon enrichment of soil organic fractions in New Zealand soils." Soil Science, 123, 6, 385-390, 1977.

Harkness, DD, AF Harrison, and PJ Bacon. "Temporal distribution of bomb C-14 in a forest soil." Radiocarbon, 28, 2A, 328-337, 1986.

Hart, J.F. "Loss and Abandonment of Cleared Farmland in the Eastern United States." Annals, Association of American Geog., 58, 417-440, 1968.

Houghton, R.A. and D.L. Skole, "The Long-Term Flux of Carbon Between Terrestrial Ecosystems and the Atmosphere as a Result of Changes in Land Use." Oak Ridge: Carbon Dioxide Information Analysis Center, July, 1990.

Houghton, RA, JE Hobbie, JM Melillo, B Moore, BJ Peterson, GR Shaver and GM Woodwell. "Changes in carbon content of terrestrial biota and soils between 1860 and

1980: a net release of CO₂ to the atmosphere." Ecological Monographs, 53(3), 235-262, 1983.

Kauppi, PE, K Mielikainen and K Kuusela. "Biomass and carbon budget of European Forests, 1971-1990." Science, 256, 70-74, 1992

Keeling, Charles D., R.B. Bacastow, A.F. Carter, S.C. Piper, T.P. Whorf, M. Heimann, W.G. Mook and H. Roeloffzen, "A Three-Dimensional Model of Atmospheric CO₂ Transport , 1. Analysis of Observational Data," (ed. David H. Peterson), Geophysical Monograph 55, AGU, Aspects of Climate Variability in the Pacific and Western Americas, p. 165-236, 1989.

Keeling, Charles D., Stephen C. Piper and Martin Heimann, "A Three-Dimensional Model of Atmospheric CO₂ Transport Based on Observed Winds: 4. Mean Annual Gradients and Interannual Variations. In: Aspects of Climate Variability in the Pacific and the Western Americas. (ed. by David H. Peterson), Geophysical Monograph 55, AGU, 1989.

Keeling, Charles D., "The Carbon Dioxide Cycle: Reservoir Models to depict the exchange of atmospheric carbon dioxide with the oceans and land plants. In Chemistry of the Lower Atmosphere, ed. by S. Rasool, 251-329, 1973.

Kulp, J.L. The Photosystem as a Sink for Carbon Dioxide. EN-6786, Special Report, May, 1990. Palo Alto: Electric Power Research Inst., 1990.

SJ Ladyman and DD Harkness. "Carbon isotope measurement as an index of soil development." Radiocarbon, 22,3 885-891, 1980.

Mann, L. K. "Changes in soil carbon storage after cultivation." Soil Science, 142, 5, 279-288, 1986.

Marland, GR; RM Rotty. (1984) "Carbon dioxide emissions from fossil fuels: a procedure for estimation and results for 1950-1982, Tellus, 36B, 232-262.

Martel, Y. A. and A. F. MacKenzie. "Longer-term effects of cultivation and land use on soil quality in Quebec." Can. J. Soil Sci., 60, 411-420, 1980.

Martel, Y. A. and E. A. Paul. "Effects of cultivation on the organic matter of grassland soils as determined by fractionation and radiocarbon dating." Can. J. Soil Sci., 54, 419-426, 1974.

Martel, YA and EA Paul. "Use of radiocarbon dating of organic matter in the study of soil genesis." Soil Sci. Soc. Amer. Proc., 38, 501-506, 1974.

Meints, V. W. and G. A. Peterson. "The influence of cultivation on the distribution of nitrogen in soils of the ustoll suborder." Soil Science, 124, 6, 334-342, 1977.

McGraw-Hill Encyclopedia of Science & Technology, "Rice," vol. 11, p. 662 - 667, McGraw-Hill, NY, 1982.

Oades, J.M., "The retention of organic matter in soils," Biogeochemistry 5:35-70, 1988.

O' Brien, BJ and JD Stout. "Movement and turnover of soil organic matter as indicated by carbon isotope measurements." Soil Biol. Biochem., 10, 309-317, 1978.

- O'Brien, B.J. "Soil organic carbon fluxes and turnover rates estimated from radiocarbon enrichments." Soil Biol. Biochem., 16, 2, 115-120, 1984.
- O'Brien, B.J. "The use of natural and anthropogenic C-14 to investigate the dynamics of soil organic carbon." Radiocarbon, 28, 2A, 358-362, 1986.
- Oeschger, H., U. Siegenthaler, U. Schotterer and A. Gugelmann, "A box diffusion model to study the carbon dioxide change in nature," Tellus, 27:168-192, 1975.
- Parton, W.J., C.V. Cole, J.W.B. Stewart, D.S. Ojima and D.S. Schimel, "Stimulating regional patterns of soil, C,N and P dynamics in the U.S. central grasslands region," Ecology of Arable Land, 99-108, 1989, (eds. M. Clarholm and L. Bergström) Lkuwer Academic Publishers, 1989.
- Parton, W.J., D.S. Schimel, C.V. COle, and D.S. Ojima, "Analysis of factors controlling soil organic matter levels in Great Plains Grasslands," Soil Science America Journal, 51:1173-1179, 1988.
- Pastor, John and W.M. Post, "Influence of climate, soil moisture and succession on forest carbon and nitrogen cycles," Biogeochemistry, 2, 3-27, 1986.
- Post, W.M., P.J. Zinke, A.G. Strangenberger, W.R. Emanuel, H. Jenny, and J.S. Olson, "Summaries of soil carbon storage in world life zones," in Global Dynamics of Biospheric Carbon. Ed. by S. Brown, p. 131-139, 1982.
- Post, WM and LK Mann. "Changes in soil organic carbon and nitrogen as a result of cultivation." Soils and the Greenhouse Effect, ed by AF Bouwman. New york: John Wiley and Sons, 1990, 401-407.
- Prentice, Katharine C. & I. Y. Fung, "The Sensitivity of Terrestrial Carbon Storage to Climate Change," Nature, 346, 48-51, 1990.
- Rice, Charles W. and Kirby L. Rogers. "Denitrification in the Vadose Zone: The Microbial Factor on Nitrate Persistance." Report #G1563, Kansas Water Resources Research Institute, 1991.
- Revelle, R. and W. Munk, "The Carbon Dioxide Cycle and the Biosphere," in Energy and Climate, Studies in Geophysics, p. 140-158, National Research Council, National Academy of Sciences, Washington, DC, 1977.
- Ritchey, Jerry C. and J. Roger McHenry. "Application of radioactive fallout Cs-137 for measuring soil erosion and sediment accumulation rates and patterns: a review." J. Environ. Qual., 19, 215-233, 1990.
- Sarmiento, JL and ET Sundquist. "Revised budget for the oceanic uptake of anthropogenic carbon dioxide." Nature, 356, 589-593, 1992.
- Scharpenseel, HW and P Becker-Heidmann, HU Neue and K. Tsutsuke. "Bomb-carbon, C-14 dating and C-13 measurements as tracers of organic matter dynamics." The Science of the total environment, 81/81, 99-110, 1989.
- Scharpenseel, H-W and P Becker-Heidmann. "Shifts in C-14 patterns of soil profiles due to bomb carbon." Radiocarbon, 31, 3, 627-636, 1989.

Schiff, S. L., R. Aravena, S. E. Trumbore, and P. J. Dillon. "Dissolved organic carbon cycling in forested water sheds: A carbon isotope approach." Water Resources Research, 26, 2949-2957, 1990.

Schiffman, P.M., and W.C. Johnson. "Phytomass and detrital carbon storage during forest regrowth in the Southeastern US Piedmont." Canadian Journal of Forestry Research, 19, 69-78, 1989.

Schimel, David S., "Carbon and Nitrogen Turnover in Adjacent Grassland and Cropland Ecosystems," Biogeochemistry, 2:345-357, 1986.

Schimel, David S., D.C. Coleman, and K.A. Horton, "Soil Organic Matter Dynamics in Paired Rangeland and Cropland Toposequences in North Dakota," Geoderma, 36, 201-214, 1985.

Schimel, David S., M.A. Stillwell, and R.G. Woodmansee, "Biogeochemistry of C, N and P in a Soil Catena of the Shortgrass Steppe", Ecology, 66(1), p. 276-282, 1985.

Schlesinger, W. H. Biogeochemistry: An Analysis of Global Change. New York: Academic Press, 1991.

Schlesinger, W.H., "Soil organic matter: a source for atmospheric CO₂." In Woodell, G.W. (ed.), The Role of Terrestrial Vegetation in the Global Carbon Cycle, Scope vol. 23, New York: Wiley, 1984, pp.111-127.

Schlesinger, W.H., "Changes in soil carbon storage and associated properties with disturbance and recovery," in Trabalka, J.R. and D.E. Reichle, (eds.), The Changing Carbon Cycle - A global Analysis. Springer-Verlag, 1986, pp. 194-220.

Schlesinger, W. H. "Evidence from chronosequence studies for a low carbon storage potential of soils." Nature, 348, 232-234, 1990.

Schoenau, J. J. "Nutrient Cycling Processes in Prairie and Boreal Forest Soils." University of Saskatchewan, Ph. D. Thesis, 1988.

Siegenthaler, U and H Oeschger. "Biospheric CO₂ emissions during the past 200 years reconstructed by deconvolution of ice core data." Tellus, 39B, 140-154, 1987.

Singh, J.S., and S.R. Gupta, "Plant Decomposition and Soil Respiration in Terrestrial Ecosystems." The Botanical Review, 43, 449-528, 1977.

Stewart, J.W.B., and C.V. Cole, "Influence of elemental interactions and pedogenic processes in organic matter dynamics." In Clarholm, L and L. Bergstrim (eds.), Ecology of arable land -- Perspectives and Challenges, *Development in Plant and Soil Sciences*, vol. 39, Kluwer Academic Publishers, 1989.

Sundquist, Eric T., "Geological Perspectives on Carbon Dioxide and the Carbon Cycle," in The Carbon Cycle and Atmospheric CO₂: Natural Variations Archean to Present, *Geophysical Monograph* 32, 1985.

Tans, Pieter P., I.Y. Fung and T. Takahashi, "Observational Constraints on the Global Atmospheric CO₂ Budget." Science, 247, 1431-1438, 1990.

Trexler, MC. "Minding the carbon store: weighing US Forestry strategies to slow global warming." World resources institute, 1990.

Trumbore, Susan E., Carbon Cycling and Gas Exchange in Soils, Columbia University, Ph.D. thesis, 1988.

Trumbore, SE, G. Bonani and W Wolfi. "The rates of carbon cycling in several soils from AMS C-14 measurements of fractionated soil organic matter." *Soils and the Greenhouse Effect*. ed by AF Bouwman. New York: John Wiley and Sons, 1990, 407-414.

Trumbore, SE, JS Vogel and JR Southon. "AMS C-14 measurements of fractionated soil organic matter." Radiocarbon, 31, 3, 644-654, 1989.

Tsutsuki, K, C Suzuki, S. Kuwatsuka, P Becker-Heidmann and H-W Scharpenseel. "Investigation on the Stabilization of the humus in mollisols." S. Pflanzenernahr. Bodenk., 151, 87-90, 1988.

Verardo, David J., P.N. Froelich and A. McIntyre, "Determination of organic carbon and nitrogen in marine sediments using the Carlo Erba NA-1500 Analyzer," Deep-Sea Research, 37:157-165, 1990.

Wassenaar, Leonard, Ramon Aravena and Peter Fritz. "The Geochemistry and evolution of natural organic solutes in groundwater." Radiocarbon, 32,3, 865-876, 1989.

Woodwell, G.M., R.H. Whittaker, W.A. Reiners, G.E. Likens, C.C. Delwiche and D. B. Botkin, "The biota and the world carbon budget," Science, 199:141-146, 1978.

World Almanac and Book of Facts 1992. ed by Mark S. Hoffman. New York: Pharos Books, 1991.

The World Almanac Book of Facts, ed. Mark S. Hoffman, Pharos Books, NY, 1990.

Yonker, C.M., D.S. Schimel, E. Paroussis and R.D. Hiel, "Patterns of Organic Carbon Accumulation in a Semiarid Shortgrass Steppe, Colorado." Soil Science Society of America Journal, 52 (2), 478-483, 1988.

Zielke, R. C. and D. R. Christenson. "Organic carbon and nitrogen changes in soil under selected cropping systems." Soil Sci. Soc. Am. J., 50, 363-367, 1986.

Zinke, P.J., A.G. Strangenburger, W.M. Post, W.R. Emanuel and J.S. Olson, "Worldwide Organic Soil Carbon and Nitrogen Data," Report ORNL/TM-8857, Oak Ridge National Laboratory, Oak Ridge, TN, 1984.

**DATE
FILMED**
8 105192