DOE/ER/13657--T1

DOE/ER/13657--T1

DE92 011427

**FINAL TECHNICAL REPORT: EXPERIMENTAL AND
COMPUTER MODELING STUDIES OF ISOTOPICALLY SELECTIVE,
TWO-STEP, LASER PHOTODISSOCIATION OF SMALL MOLECULES**

Prepared by

PAUL F. ZITTEL
Space and Environment Technology Center
Technology Operations

25 March 1992

Engineering and Technology Group
THE AEROSPACE CORPORATION
El Segundo, CA 90245-4691

Prepared for

DEPARTMENT OF ENERGY
Washington, DC 20585

Grant No. DE-FG03-87ER13657

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

PREFACE

This document is a final report on research conducted under Department of Energy Grant No. DE-FG03-87ER13657, which covered the period 12/1/86 - 12/31/91. The DOE grant was a follow-on to a previous DOE contract. By agreement this document is also a final report on activities under the aforementioned Department of Energy Contract No. DE-AC03-76ER01017 (originally AEC No. AT(04-3)-1017 and later ERDA No. EY-76-C-03-1017), which covered the period 6/1/74 - 11/30/86.

The results of the research conducted under this contract/grant have been thoroughly documented in journal articles and publicly released technical reports, which are listed herein. It is not the intention of this report to review every aspect of the research program in detail.

After an introduction (Section I), a general overview section (Section II) provides a brief summary of the different problems studied during the program and provides references to more detailed reports, which are itemized in a list of publications (Appendix). Following the overview, a few topics are reviewed in more detail (Sections III-V). The latter topics are those which hold particular scientific interest, or which have particular relevance to practical isotope enrichment schemes. Finally, some general observations on the two-step photodissociative isotope separation scheme are presented in a conclusion section (Section VI).

CONTENTS

PREFACE	ii
I. INTRODUCTION	1
II. GENERAL OVERVIEW	3
III. PHOTODISSOCIATION CROSS SECTIONS OF VIBRATIONALLY EXCITED MOLECULES	6
IV. ISOTOPE SEPARATION IN THE LOW-TEMPERATURE GAS PHASE	10
V. ISOTOPE SEPARATION IN CRYOGENIC LIQUID SOLUTIONS	14
VI. CONCLUSIONS	20
REFERENCES	23
APPENDIX: LIST OF PUBLICATIONS	24

FIGURES

1.	Photodissociation of Vibrationally Excited HBr	7
2.	Ozone Photodissociation Cross Sections	8
3.	OCS Photodissociation Cross Sections	9
4.	Temperature Dependence of Oxygen Isotope Enrichment Factors	11
5.	Cost of Enriching ^{33}S by Two-Step Photodissociation of OCS	12
6.	Cryostat for Liquid Samples	15
7.	FTIR Spectrum of OCS in Liquid Ar	15
8.	Computer-Modeled, Liquid-phase, Carbon and Oxygen Enrichment Factors	19

TABLES

1.	O, C, and S Isotope Enrichment Factors at 295K	13
2.	$^{16}\text{O}^{12}\text{C}^{32}\text{S}$ Vibrational Band Centers, Half-Widths, and Strengths in Cryogenic Liquid Solutions	16
3.	OCS Vibrational Relaxation in Cryogenic Solutions	18

I. INTRODUCTION

Since their advent, lasers have been considered as potential tools for achieving efficient and highly selective isotope separation.¹ A variety of laser isotope separation methods have been demonstrated, including photoionization, photodissociation, photochemical, and momentum transfer techniques. Unlike most current separation processes, which are based on small isotopic differences in mass transport or thermodynamic properties, proposed laser processes are based on spectroscopic differences, which can be very large for the different isotopes of an atom, or isotopomers of a given molecule. Particularly in the case of middle isotopes, which are collected in the isotopically cluttered middle fraction of a typical mass transport or thermodynamic separation, the laser excitation approach can take advantage of clean differences in the absorption spectrum of a molecule containing a middle isotope (e.g., ³³S) from the spectra of isotopomers containing neighboring isotopes (e.g., ³²S and ³⁴S). The widths of absorption lines of small molecules in the gas phase can be much smaller than spectral shifts due to isotopic substitution (i.e., $\sim 10^{-2} \text{ cm}^{-1}$ versus $\sim 10 \text{ cm}^{-1}$ for rovibrational features) and smaller than the spacing of absorption lines (i.e., $0.1\text{-}1 \text{ cm}^{-1}$ for many linear triatomics). Even in condensed phases, isotopic spectral shifts can be much larger than the entire width of a collapsed rovibrational absorption band.

The approach to laser isotope separation taken in this study is based on isotopically selective, *two-step*, laser photodissociation of small molecules. The first, infrared, laser photon selectively excites molecules containing a particular atomic isotope to a vibrational level above those levels populated thermally in the bulk sample. The second, ultraviolet, laser photon is not by itself isotopically selective, but photodissociates vibrationally excited molecules more efficiently than ground state molecules, producing isotopic enrichment in the photodissociation fragments. Subsequent chemical scavenging reactions provide the means for collecting the enriched photodissociation fragments. The isotopic selectivity and efficiency of such a process depend on a number of fundamental spectroscopic, kinetic and chemical properties. Important questions include the extent to which vibrational populations can be changed by infrared laser pumping, the isotopic selectivity of the vibrational excitation process, the quantitative effect of vibrational excitation on photodissociation cross sections, and the rapidity of various isotope-scrambling energy transfer and chemical reaction processes.

A primary goal of this study is the *measurement of fundamental molecular processes* which control the two-step, photodissociative isotope enrichment process. This objective has led to experimental measurements of UV photodissociation cross sections for vibrationally excited states of several small molecules, including the first cross section reported for any neutral molecule in a specific, excited vibrational state. Another fundamental property is the rate constant for collisional relaxation of a vibrationally excited molecule. As part of this study,

vibrational relaxation rate constants have been measured for relevant molecules in both the gas phase and in cryogenic liquid solutions. Finally, our recent work with cryogenic liquids has resulted in fundamental spectroscopic measurements of vibrational band centers, widths and strengths for some small molecules in a variety of cryogenic liquid solvents.

A second goal of this study has been the *laboratory demonstration of isotope enrichment* for isotopes of practical interest and for processes with a potential for larger scale production. Where possible, efforts have focussed on the separation of middle isotopes, such as ^{17}O and ^{33}S , which are expensive and difficult to separate using other techniques. Considerable success has been achieved in demonstrating the enrichment of isotopes of bromine, carbon, oxygen and sulfur. Although the laboratory enrichment measurements are primarily used with measurements of fundamental properties and computer modeling to understand the enrichment process, crude economic estimates suggest that it is reasonable to consider the techniques studied here for larger scale, practical isotope enrichment.

A third goal of this study has been the *application of computer modeling* to the two-step enrichment process. Experimental measurements define as many as possible of the critical photophysical and chemical parameters required by an *ab initio* computer model of the enrichment process. A comparison of experimental enrichment measurements and computer modeling results can confirm the adequacy of the physical model of the process, or indicate that some important elements have not been considered. A particularly useful function of an experimentally verified model is the prediction of isotope enrichments for a wide range of conditions that for various reasons cannot be immediately investigated in the laboratory.

Progress toward these goals has been documented in detail in a variety of journal articles and technical reports. It is not the objective of this final report to discuss each aspect of the work done under this contract/grant in detail. Section II of this document is a general overview of the program, touching briefly on each of the problems addressed by the study and providing references to more detailed discussions. Sections III-V review in more detail a few topics that are of particular scientific interest, or which are relevant to practical isotope separation. The objective of the detailed review is to draw together into an organized discussion the new information that has been learned about a particular topic over the course of the program. In Section VI, some general conclusions are presented about the two-step, photodissociative isotope enrichment method. Finally, a list of publications under this contract/grant is given in an Appendix.

II. GENERAL OVERVIEW

This section presents an overview of the different problems addressed during the course of our study of two-step photodissociative isotope separation. The discussion is not detailed, and references are provided to full accounts in journal articles and technical reports.

During the course of this program a versatile experimental and computer modeling capability have been developed to study a variety of small molecules in different chemical environments. A preliminary modeling study used a modified version of the Aerospace Corporation NEST Chemistry Computer Program² to model the complex photoabsorption, energy transfer, and chemical reaction processes associated with isotopically selective, two-step photodissociation of HBr in mixtures with NO and rare gases.³ A subsequent computer study modeled the effect of rotational "bottlenecking" on the infrared absorption step of the two-step photodissociation process.⁴ The modeling results indicated the possibility of separating the ⁷⁹Br and ⁸¹Br isotopes of bromine by two-step photodissociation of HBr and chemical scavenging of Br atoms by reaction with NO. The use of an intracavity, isotopic, cold gas filter was experimentally investigated as an efficient means of generating isotopically specific emission on the rovibrational lines of hydrogen halide molecular lasers for isotopically specific vibrational pumping in enrichment experiments.⁵

The first isotope separation measurements of the program focussed on isotopically selective photodissociation of HBr. The critical UV photodissociation cross section of vibrationally excited HBr($v=1$) was measured at 258.9 nm in the red wing of the HBr bound-continuous absorption band.⁶ Vibrational excitation was found to produce a large increase in the photodissociation cross section, indicating the potential for substantial isotope enrichment through two-step photodissociation. An real-time mass spectrometric sampling technique was subsequently used to study the enrichment of the ⁸¹Br isotope with respect to ⁷⁹Br, generated by isotopically specific vibrational excitation of H⁸¹Br with an HBr molecular laser followed by UV photodissociation at 258.9 nm.⁷ Extensive computer modeling of the specific experimental conditions indicated that isotope exchange between Br and HBr limited enrichments.⁷ The real-time mass spectrometric sampling technique also made it possible to determine diffusion coefficients for highly reactive Br atoms in Ar gas mixtures and for Br₂ molecules in NO.⁸

The isotope enrichment that can be achieved in a two-step, IR/UV, photodissociation process is critically dependent on the UV photodissociation cross section of the vibrationally excited target molecule. Prior to the present study, no photodissociation cross sections had been reported for neutral molecules in specific vibrational levels. Several experimental studies were undertaken to obtain excited state cross section information. The effect of excitation of the vibrational stretching modes of O₃ on its UV photodissociation cross section was studied as a function of UV wavelength on the red side (i.e., $\lambda=259-315$ nm) of the intense Hartley band.⁹

An extremely large enhancement was observed for the dissociation channel producing $\text{O}_2 + \text{O}(^1\text{D})$ products. Intense IR laser pumping and vibrationally enhanced UV photodissociation of a more complicated polyatomic molecule were studied in the case of CH_3Br .¹⁰ The most detailed study of vibrationally state specific cross sections was undertaken for the OCS molecule.¹¹⁻¹³

Two-step photodissociation of carbonyl sulfide (OCS) has been studied comprehensively, providing detailed information about the vibrational state dependence of photodissociation cross sections, measurements of vibrational relaxation rate constants, and new spectroscopic data in the condensed phase. Extensive measurements of isotope enrichment factors have been made in the gas phase as a function of temperature and in the cryogenic liquid phase. Initial experiments demonstrated two-step enrichment of the middle isotope ^{33}S and investigated basic chemical systems for scavenging photogenerated sulfur atoms.¹¹ A detailed study in the room temperature gas phase, using a CO_2 laser to excite the $2\nu_2$ (bending overtone) vibrational level of OCS and a UV excimer laser for photodissociation, determined the UV photodissociation cross section of $\text{OCS}(2\nu_2)$.¹² All of the stable isotopes of carbon, oxygen, and sulfur were enriched by employing various CO_2 laser lines to vibrationally excite selected OCS isotopomers prior to photodissociation.

Two-step photodissociation of OCS through the ν_1 (OC-S stretching) vibrational level has also been investigated.¹³ The effort followed development of a CO_2 laser-pumped, NH_3 molecular laser for exciting the ν_1 vibrational mode of OCS. The experiments determined a cross section for UV photodissociation of $\text{OCS}(\nu_1)$. Additional analysis of temperature-dependent UV absorption spectra also inferred a cross section for the $\text{OCS}(\nu_2)$ level. Various lines of the tunable NH_3 laser were used in two-step photodissociation experiments to achieve enrichment of each of the stable isotopes of sulfur.

It is evident from accumulated cross section measurements that a very sharp increase in the UV photodissociation cross section of OCS occurs with increasing vibrational energy. The results suggest that two-step photodissociation through the highest energy fundamental vibrational mode of OCS (i.e., the ν_3 O-CS stretch) should substantially increase enrichment factors over the values achieved by excitation of lower energy modes. A frequency-doubled CO_2 laser provided $4.8\text{-}\mu\text{m}$ radiation for direct excitation to the ν_3 vibrational level of OCS in a study of the rate constants for collisional relaxation of the ν_3 level.¹⁴ The $4.8\text{-}\mu\text{m}$ laser power levels were inadequate for two-step photodissociation measurements; however, the measured vibrational relaxation rate constants indicate that isotopically selective, two-step photodissociation through the $\text{OCS}(\nu_3)$ level is possible and is probably the most promising of the approaches considered for selective photodissociation of OCS.

Thermal population of the low-lying vibrational levels of polyatomic molecules can reduce enrichment factors. The enrichment of oxygen and sulfur isotopes by two-step photodissociation of OCS was experimentally investigated in the gas phase between room

temperature and a low temperature of 150K, which is slightly above the freezing point of OCS.¹⁵ A computer model that explicitly included all low lying vibrational levels of OCS reproduced the observed, large improvement in enrichment factors with decreasing temperature.

Solution in a cryogenic liquid drastically alters the IR absorption spectrum, energy transfer rates, and chemistry of small molecules. The cryogenic liquid environment presents some advantages for a photodissociative isotope separation scheme, including the simplification of the vibrational absorption spectrum and photofragment scavenging processes. A study of OCS dissolved in liquid Ar and liquid O₂ determined band centers, widths and strengths for vibrational absorption bands in the 500-4200 cm⁻¹ range, as well as isotope shifts for the singly substituted isotopomers of OCS.¹⁶ Isotopically selective, two-step photodissociation of OCS was also investigated, focussing on the dependence of ¹⁶O, ¹⁷O, and ¹⁸O enrichment factors on laser characteristics and liquid sample composition. A subsequent study defined the spectral parameters of the vibrational absorption bands of OCS dissolved in liquid krypton and liquid xenon.¹⁷ The study also determined critical rates for vibrational relaxation of the low-lying vibrational levels of OCS by the four cryogenic solvents. Infrared spectral parameters and vibrational relaxation rates have also been determined for O₃ in the four cryogenic liquid solvents.¹⁷

A comprehensive computer model has been developed for two-step photodissociation of OCS (or other small molecules) in cryogenic, liquid solutions.¹⁸ The model is based on our measured spectroscopic and energy transfer data and has been used to predict isotope enrichment factors for optimum short-pulse and cw laser sources not typically available in the laboratory.

III. PHOTODISSOCIATION CROSS SECTIONS OF VIBRATIONALLY EXCITED MOLECULES

This section reviews studies of the effect of vibrational excitation on the UV photodissociation cross sections of small molecules. The vibrational enhancement of bound-continuous photodissociation cross sections is the basis for isotope separation by two-step photodissociation of molecules. Prior to the present effort, there were essentially no experimental measurements of vibrationally state-specific photodissociation cross sections for neutral molecules.

The isotope enrichment factor achieved by a two-step, molecular photodissociation process is critically dependent on the effect of vibrational excitation on the molecular photodissociation cross section. The enrichment factor for atomic isotope x with respect to atomic isotope y is defined by Eq. (1),

$$\beta_y^0(x)_{\text{observed}} = \frac{([x]/[y])_{\text{product}}}{([x]/[y])_{\text{original}}} \quad (1)$$

where "product" indicates isotopic abundance in the photodissociation products and "original" indicates abundance in the original material prior to photodissociation. For a simple system with two vibrational levels connected by an IR laser-pumped optical transition, the isotope enrichment factor in the absence of isotope scrambling by spectral, energy transfer, and chemical scavenging processes is given theoretically by Eq. (2),

$$\beta_y^0(x)_{\text{theoretical}} = f_v \left(\frac{\sigma_{UV,v}}{\sigma_{UV,0}} - 1 \right) + 1 \quad (2)$$

where f_v is the fraction of molecules containing isotope x that are pumped to the excited vibrational level v by the IR laser, and $\sigma_{UV,v}$ and $\sigma_{UV,0}$ are the UV photodissociation cross sections for the excited and ground vibrational levels, respectively. The IR pumping step can typically be saturated and rotational bottlenecking avoided, so that for non-degenerate vibrational levels the factor f_v approaches the value of ~ 0.5 .

The effect of vibrational excitation on the bound-continuous photodissociation cross section of a simple diatomic is illustrated for HBr in Fig. 1. In the red wing of the UV transition, vibrational excitation increases the Franck-Condon overlap factor of the transition by decreasing the internuclear separation at which the UV transition takes place and by expanding the vibrational wave function to larger internuclear separations. For HBr at 258.9 nm, the photodissociation cross section ratio, $\sigma_{UV,v=1}/\sigma_{UV,v=0}$, is found to be 34 ± 7 .⁶ The enhancement effect is several times larger than what one would estimate from the UV absorption

Figure 1. Photodissociation of Vibrationally Excited HBr. $\lambda_{UV}=258.9$ nm. The potential surfaces and vibrational wave functions are drawn to scale. From Ref. 6.

spectrum by simply increasing the UV photon energy by a vibrational quantum.

In principle, photodissociation cross sections for vibrationally excited levels can be calculated from highly accurate *ab initio* potential energy surfaces that fully accommodate vibrational displacements from equilibrium geometry. In reality, such surfaces are generally unavailable for any but the simplest of molecules. Simple models, such as those based on a "reflection approximation" derivation of potential energy surfaces, are inadequate.⁶ In the future, improvements in computational speed and theory may make *ab initio* calculations a useful source of excited level photodissociation cross sections for polyatomic molecules. At present, experimental measurements are necessary.

The UV wavelength dependence of excited state photodissociation cross sections is an important factor in two-step photodissociative isotope separation schemes. Qualitatively, one expects the cross section ratio $\sigma_{UV,v}/\sigma_{UV,0}$ to increase with increasing UV wavelength, although the absolute cross sections and photodissociation yield decrease. The effect was studied for photodissociation of vibrationally excited O_3 .⁹ Figure 2 shows the photodissociation cross section for the channel, $O_3+h\nu \rightarrow O_2+O(^1D)$, for vibrationally unexcited, room temperature O_3 and for O_3 that is excited with an average of 0.6 quanta of stretching vibration per molecule. The effect of vibrational excitation ranges from negligible at the UV cross section maximum near 260 nm to a factor of 100 increase in the cross section per vibrational stretching quantum

Figure 2. Ozone Photodissociation Cross Sections. From Ref. 9

at 315 nm in the red wing.

Although the isotope enrichment factors determined in our study are typically measured at a single UV wavelength, it is clear that enrichment is a function of UV wavelength. Higher cross section ratios and isotope enrichment factors are generally expected for longer UV wavelengths as one approaches the energy threshold for photodissociation. Conversely, the absolute magnitudes of the cross sections decrease with increasing wavelength. Thus, in a practical isotope enrichment scheme a trade-off exists between enrichment and photon efficiency.

The qualitative picture of vibrationally enhanced photodissociation, displayed in Fig. 1 for a diatomic molecule, is complicated for polyatomic molecules by the three-dimensional nature of the vibrational motions and potential energy surfaces. In a simple Franck-Condon approximation, the ratio of excited vibrational state to ground vibrational state photodissociation cross sections is the product of two factors. One factor depends primarily on the energy of the vibrational quantum. The second factor is a ratio of Franck-Condon overlap integrals and thus depends to an extent on the geometrical nature of the vibrational mode. For example, one might expect vibrational motions in the ground electronic state that involve large excursions along the dissociation coordinate of the upper electronic state to enhance the photodissociation cross section most strongly.

The vibrational mode dependence of the enhancement of photodissociation cross sections was investigated for the OCS molecule.^{12,13} Figure 3 shows UV (i.e., $\lambda=249$ nm)

Figure 3. OCS Photodissociation Cross Sections. $\lambda_{UV}=249$ nm. Vibrational quantum numbers label data points. $\sigma_{295K}=1.4 \times 10^{-20}$ cm². From Ref. 13.

photodissociation cross sections for specific vibrational levels of OCS relative to the cross section for OCS that is in complete thermal equilibrium at 295K. The data points are labeled by the quantum numbers ($n_{\nu_1}, n_{\nu_2}, n_{\nu_3}$) of the ν_1 (OC-S stretch), ν_2 (O-C-S bend), and ν_3 (O-CS stretch) vibrational modes of the linear triatomic molecule. The photodissociation cross sections of the vibrationally excited levels increase exponentially with the energy of vibrational excitation in the ground electronic state. The increase is much steeper than a simple shift in the UV absorption spectrum by the vibrational energy would imply. Perhaps coincidentally, the enhancements by bending and sulfur-stretching vibrational modes follow the same correlation. If the high energy, ν_3 ($E_{vib}=2062$ cm⁻¹), vibrational mode follows the same correlation, the cross section ratio $\sigma_{UV, \nu_3} / \sigma_{UV, 0}$ is predicated to be ~ 400 .

For several reasons it appears that two-step photodissociation through the ν_3 intermediate vibrational level of OCS *in the gas phase* may be a highly selective, efficient means of isotope enrichment. Although the vibrational enhancement effect for the ν_3 mode may not be as large as the simple correlation of Fig. 3 suggests, the cross section ratio is probably much larger than the ratios for lower energy vibrational levels. Vibrational relaxation experiments¹⁴ have determined gas-phase relaxation times for the ν_3 level that are compatible with a two-step enrichment process. Finally, the band strength of the $\nu_3=0 \rightarrow 1$ infrared absorption band is very large, allowing saturation of the vibrational excitation step with a relatively low-power IR laser.

IV. ISOTOPE SEPARATION IN THE LOW-TEMPERATURE GAS PHASE

This section reviews a study of the effect of temperature on the enrichment factors achieved by two-step photodissociation of molecules in the gas phase. For polyatomic target molecules, the simple expedient of lowering the gas temperature leads to a large enhancement of isotope enrichments.

Polyatomic molecules typically have low energy vibrational levels, which can be significantly populated at thermal equilibrium at room temperature. To the extent that the photodissociation cross sections from such levels are significantly larger than the photodissociation cross section of the vibrationless ground level, thermal population of the levels reduces two-step isotope enrichment factors. Thus, the enrichment factors can have a substantial dependence on gas temperature.

The simple theoretical expression for the enrichment factor given for a two level system in Eq. (2) can be modified for a polyatomic molecule to include the effect of thermally populated low energy vibrations. The result is given by Eq. (3),

$$\beta_v^0(x)_{\text{theoretical}} = f_{v*} \left(\frac{\sigma_{UV,v*} - \sigma_{UV,0}}{\sigma_{UV}(T)} \right) + 1 \quad (3)$$

where f_{v*} is the fraction of molecules containing isotope x that are pumped to the excited vibrational level v^* by the IR laser, $\sigma_{UV,v*}$ and $\sigma_{UV,0}$ are the UV photodissociation cross sections for the excited and ground vibrational levels, respectively, and $\sigma_{UV}(T)$ is the effective photodissociation cross section for a molecule at thermal equilibrium. For polyatomic molecules with degenerate vibrational levels, the factor f_{v*} approaches $d_{v*}/(d_{v*} + d_0)$, where d_{v*} and d_0 are the vibrational degeneracies of the excited and ground vibrational levels, respectively. The effective photodissociation cross section for a molecule at thermal equilibrium is given by Eq. (4),

$$\sigma_{UV}(T) = \sum_v n_v(T) \sigma_{UV,v} \quad (4)$$

where $n_v(T)$ is the fractional population of vibrational level v at thermal equilibrium and $\sigma_{UV,v}$ is the photodissociation cross section for the vibrational level. Knowledge of the photodissociation cross sections for the laser-pumped vibrational level and the low energy levels that have thermal populations of a few percent or more are required for modeling the temperature dependence of the enrichment factor.

The temperature dependence of enrichment factors has been studied for two-step photodissociation of OCS.¹⁵ Figure 4 shows the dependence of the enrichment of ^{18}O with respect to ^{17}O as a function of sample temperature (i.e., $\beta_{17}^0(18)$). The temperature dependence

Figure 4. Temperature Dependence of Oxygen Isotope Enrichment Factors. The solid curves are model calculations. From Ref. 15.

of the measured enrichment factors is fit well by the theoretical curve calculated from Eqs. (3) and (4) with the photodissociation cross sections for the low energy vibrational levels of OCS.¹³ Similar results were found for sulfur isotope enrichments. Thermal population of low energy vibrations can also reduce enrichments by allowing hot band absorptions of the IR laser pulse by isotopomers other than the isotopomer that is intended to be excited. As sample temperatures are lowered, the hot band absorptions are reduced. The effect is seen in Fig. 4, where an undesired enrichment of ^{16}O with respect to ^{17}O is reduced with decreasing temperature (i.e., $\beta_{17}^0(16)$). The effect is modeled well by a modified version of Eq. (3).

The reduction of gas-phase sample temperatures to 140-150K, which is just above the melting point of OCS, presents no obvious difficulties in gas handling procedures, or product throughput, for a large scale enrichment process. The advantage of almost doubling enrichment factors over room temperature values is important, since the cost of producing a high degree of isotope enrichment in a cycled process is strongly dependent on the single-cycle enrichment factor $\beta_y^0(x)$. A rough estimate of the cost of producing ^{33}S by cycling a two-step photodissociation process is shown in Fig. 5 as a function of the single-cycle enrichment factor.

Enrichments of all of the stable isotopes of C, O, and S by two-step photodissociation of OCS have been studied at room temperature.¹² The experimental enrichment factors at 295K are listed in Table 1. By analogy with the low-temperature measurements and computer

Figure 5. Cost of Enriching ^{33}S by Two-Step Photodissociation of OCS. From a previous progress report to DOE.

modeling results for ^{18}O and ^{33}S enrichment, it is expected that all of the enrichment factors will increase to values of 4-5 as sample temperatures are decreased to 140-150K. The low-temperature enrichment factors are a factor of 2-3 below the theoretical maximum predicted by Eq. (3) for photodissociation through the $2\nu_2$ vibrational level of OCS with perfectly selective saturation of the IR absorption step. The reason for the low experimental enrichments is not completely understood. The main factor is probably isotopically non-selective absorption of the somewhat broad CO_2 infrared laser lines, which are estimated to be $\sim 0.1 \text{ cm}^{-1}$ fwhm compared to OCS rovibrational line spacings of a few tenths of a cm^{-1} . Other factors may contribute to a lesser extent, including less than complete saturation of the IR absorption step and some relaxation of the OCS $2\nu_2$ vibrational level in the time interval between IR and UV laser pulses.

Although some improvement is possible in the enrichments obtained by photodissociation through the OCS $2\nu_2$ and ν_1 vibrational levels, an optimum gas-phase enrichment scheme will clearly involve photodissociation through the higher energy ν_3 vibrational level. Relevant energy transfer and photodissociation cross section measurements from the present study suggest the following experimental conditions. A narrow-band ($\leq 0.02 \text{ cm}^{-1}$ fwhm), short-pulse ($\leq 100 \text{ ns}$ fwhm), tunable, $4.8\text{-}\mu\text{m}$ laser source with pulse energies of only a few tens of millijoules is required to saturate the $\nu_3=0\rightarrow 1$ vibrational transition with high isotopic selectivity. Photodissociation by a 249 nm KrF excimer laser would follow the IR laser pulse by a delay of

Table 1. O, C, and S Isotope Enrichment Factors at 295K. ^a

Enriched Isotope	Isotope Mixture	CO ₂ Laser Line	Scavenging Systems	Enrichment Factors ^b	
¹² C	O ^{12,13} CS	P(22)	C ₂ F ₄ , NO ₂ , NO ₂ /O ₂	$\beta_{13}(12)$	3.4
¹³ C	O ^{12,13} CS	P(34)	C ₂ F ₄ , NO ₂	$\beta_{12}(13)$	1.7
¹⁶ O	^{16,17,18} OCS	P(22)	C ₂ F ₄ , NO ₂ , NO ₂ /O ₂	$\beta_{17}(16)$ $\beta_{18}(16)$	2.6 2.6
¹⁶ O	^{16,17,18} OCS	R(4)	C ₂ F ₄	$\beta_{17}(16)$ $\beta_{18}(16)$	3.1 3.1
¹⁷ O	^{16,17,18} OCS	P(16)	C ₂ F ₄ , NO ₂	$\beta_{16}(17)$ $\beta_{18}(17)$	1.5 1.5
¹⁶ O, ¹⁷ O	^{16,17,18} OCS	R(8)	C ₂ F ₄	$\beta_{18}(16)$ $\beta_{18}(17)$	1.8 1.5
¹⁸ O	^{16,17,18} OCS	P(26)	C ₂ F ₄ , NO ₂	$\beta_{16}(18)$ $\beta_{17}(18)$	1.8 3.0
³² S	OC ^{32,33,34} S	P(22)	NO ₂ /O ₂	$\beta_{33}(32)$ $\beta_{34}(32)$	2.9 2.9
³³ S	OC ^{32,33,34} S	P(10)	NO ₂ , NO ₂ /O ₂	$\beta_{32}(33)$ $\beta_{34}(33)$	3.1 1.8
³³ S	OC ^{32,33,34} S	P(12)	NO ₂ /O ₂	$\beta_{32}(33)$ $\beta_{34}(33)$	2.8 2.4
³³ S	OC ^{32,33,34} S	P(14)	NO ₂ /O ₂	$\beta_{32}(33)$ $\beta_{34}(33)$	3.0 3.2
³⁴ S	OC ^{32,33,34} S	P(16)	NO ₂ /O ₂	$\beta_{32}(34)$ $\beta_{33}(34)$	1.9 1.9

^a From Ref. 12.

^b Enrichments are not corrected for sample depletion; single-shot enrichments $\beta_{\nu}^0(x)$ are 5-10% larger. Enrichments were the same for different scavenging systems.

≤ 100 ns. Samples would be cooled to 140-150K. Gas compositions would be 1-2 Torr of OCS, 10-20 Torr of ethylene or propylene scavenger for S atoms, and 100-200 Torr of Ar buffer gas. The enrichments that could be achieved with this scheme depend on the UV photodissociation cross section of OCS(ν_3). Reasonable extrapolations from lower energy vibrational levels suggest a cross section that would generate enrichments roughly an order of magnitude greater than those achieved thus far.

V. ISOTOPE SEPARATION IN CRYOGENIC LIQUID SOLUTIONS

This section reviews the important elements of two-step photodissociative isotope separation in the cryogenic liquid phase. In addition to demonstrating isotope enrichment, the study has produced basic spectroscopic and kinetic data for the cryogenic liquid environment.

The infrared vibrational spectra of small, polyatomic molecules dissolved in cryogenic liquids are qualitatively different from gas-phase spectra. The rotational structure of a vibrational absorption band, which consists of many narrow lines spread over 50-100 cm^{-1} in the room temperature gas phase, can collapse to a single strong feature a few tenths to a few cm^{-1} wide in cryogenic solution. The cryogenic bandwidth can be much narrower than the shift in band center due to isotopic substitution. Thus, high isotopic specificity may be achieved in the vibrational excitation step, even with low resolution lasers. At cryogenic temperatures the thermal population of low energy vibrational levels is negligible, eliminating the complicating effects of hot band absorptions. In the cryogenic liquid environment, the kinetics of vibrational energy transfer and chemical reactions are expected to be significantly different from the gas phase. This may lead to some advantages over gas-phase isotope enrichment schemes. The solute densities that can be achieved in cryogenic solution are comparable to, or greater than, densities that would be employed in typical gas-phase laser isotope separation systems.

A potential advantage of the liquid environment is the simplification of a molecule's infrared absorption spectrum. A detailed study has been made of the spectra of OCS and O_3 dissolved in liquid rare gases and liquid O_2 .^{16,17} Figure 6 shows the cryogenic cell developed to conduct spectroscopic, kinetic, and isotope enrichment experiments in the cryogenic liquid phase (and low-temperature gas phase). The cryostat generates an 8.8 cm^3 sample of liquid at a selectable temperature down to 78K and holds the temperature to $\pm 0.2\text{K}$. The liquid cell is a cross, sealed with CsI or BaF_2 windows, permitting a variety of spectroscopic and kinetic experiments. Figure 7 shows a portion of a Fourier transform infrared spectrum of 15 ppm OCS dissolved in liquid Ar at a temperature of 84.4K. The spectrum illustrates the magnitude of the isotopic band center shifts relative to the bandwidth (i.e., 0.73 cm^{-1} hwhm) for the OCS ν_3 vibrational band. The band shapes and positions determine, for example, that the absorption cross section of an $^{18}\text{O}^{12}\text{C}^{32}\text{S}$ molecule at the center of its absorption band is a factor of ~ 2400 larger than the absorption cross section of a $^{16}\text{O}^{12}\text{C}^{32}\text{S}$ molecule at the identical IR wavelength. Similarly, for $^{17}\text{O}^{12}\text{C}^{32}\text{S}$ and $^{16}\text{O}^{13}\text{C}^{32}\text{S}$ the selectivity factors with respect to $^{16}\text{O}^{12}\text{C}^{32}\text{S}$ are ~ 680 and ~ 5200 , respectively. Sulfur isotope shifts are very small in the ν_3 band. Spectroscopic parameters measured in this study for a number of OCS vibrational bands and various solvents are given in Table 2. Isotope band center shifts have also been measured in liquid Ar and O_2 ¹⁶ and can be estimated accurately from gas-phase data.¹⁸

Figure 6. Cryostat for Liquid Samples.

Figure 7. FTIR Spectrum of OCS in Liquid Ar.

TABLE 2. $^{16}\text{O}^{12}\text{C}^{32}\text{S}$ Vibrational Band Centers, Half-Widths, and Strengths in Cryogenic Liquid Solutions^{a,b}

	upper vibrational level							
	(01 ¹ 0)	(10 ⁰ 0)	(02 ⁰ 0)	(20 ⁰ 0)	(12 ⁰ 0)	(00 ⁰ 1)	(04 ⁰ 0)	(10 ⁰ 1)
Band Center $\bar{\nu}_0$, cm ⁻¹								
Ar(l)	519.77	858.93	1045.42	1710.86	1890.57	2052.50	2101.72	2908.52
Kr(l)	518.73	858.42	1043.42	1709.79	1887.93	2049.66	2097.69	2905.21
Xe(l)	517.18	857.69	1040.40	1708.33	1884.11	2046.29	2091.73	2901.10
O ₂ (l)	520.01	859.14	1045.88	1711.25	1891.28	2052.64	2102.66	2908.93
gas	520.42	858.97	1047.04	1710.98	1892.23	2062.20	2104.83	2918.11
Band Half-Width γ , cm ⁻¹								
Ar(l)	0.34	0.65	0.71	0.92	0.76	0.73	0.94	0.84
Kr(l)	0.40	0.80	0.82	1.14	0.82	0.86	1.07	1.01
Xe(l)	0.63	1.14	1.32	1.61	1.20	1.22	1.73	1.35
O ₂ (l)	0.36	0.67	0.76	0.94	0.82	0.75	1.13	0.83
Band Strength S_b , 10 ⁻¹⁹ cm/molecule								
Ar(l)	7.0	12.0	4.8	3.0	4.5	1170	5.1	13.8
Kr(l)	7.0	12.0	4.8	2.9	4.6	1180	5.6	13.9
Xe(l)	6.7	10.5	4.6	2.6	4.4	1050	5.5	12.7
O ₂ (l)	7.5	12.6	4.9	2.9	4.7	1220	5.2	14.4
gas	4.8	14.7	4.8	2.7	4.4	1120	7.9	13.5

^a From Ref. 17.

^b Temperatures of Ar, Kr, Xe and O₂ solutions are 84.4, 116.8, 162.0 and 78.6 K, respectively.

The spectroscopic measurements show that OCS can be pumped with high isotopic selectivity in any of several vibrational bands. Different bands are optimum for O, C, and S isotopes. Relatively modest laser pulse energies are required to saturate the IR pumping step (e.g., ~ 0.1 J/cm²/pulse for the $2\nu_2$ band pumped by the CO₂ laser). Similar results have been obtained for other triatomic molecules. Analogous spectroscopic measurements have been made for O₃ in liquid rare gas solvents.¹⁷ Selectivity factors for oxygen isotopes in the ozone spectrum are found to be large, although not as large as for OCS. Spectral measurements have also been made for N₂O dissolved in liquid O₂ and liquid Ar.¹⁹ Narrow bandwidths (typically 1 cm⁻¹ hwhm) and large isotope shifts lead to large selectivity factors for oxygen and nitrogen isotopes.

The narrowest liquid-phase absorption bandwidths and consequently the largest isotopic selectivity factors were found for small *linear* molecules (e.g., OCS and N₂O). The bands are presumably narrowed by strong rotational hindering of the non-spherical molecules. Somewhat

broadband widths were found for non-linear molecules, such as O_3 and in preliminary spectra¹⁹ of SO_2 . In general, linear molecules are probably the best targets for a two-step enrichment process. In order to pump the vibrational bands that are optimum for separating isotopes of a particular atom, an IR laser source that is continuously tunable over a broad wavelength range would be ideal. Such sources will undoubtedly become available as laser technology progresses. Some bands can, however, be pumped with currently available, discretely tunable lasers.

A critical factor in liquid-phase, two-step, isotope enrichment is the vibrational relaxation of small polyatomic molecules by the cryogenic solvent. A very small amount of data on the relaxation process is available for a few polyatomics. Theories relating liquid-phase vibrational relaxation to relaxation in the higher temperature gas phase have been developed. An important part of the present study has been the experimental measurement of relaxation rates for OCS and O_3 by cryogenic liquids.¹⁷ Table 3 lists the measured relaxation rates that are relevant to two-step photodissociation of OCS through the $2\nu_2$ vibrational level. The rates measured for OCS (and O_3) in cryogenic solution are found to be in rough agreement with rates estimated by theoretically extrapolating room temperature gas-phase rates to the low temperature and high density of the cryogenic solution.¹⁷ The extrapolation procedure can provide a preliminary estimate of unmeasured, critical, liquid-phase relaxation rates, although more precise values are preferable for detailed modeling of the two-step enrichment process. The vibrational relaxation rates determine the laser time scale necessary for completing the two-step photodissociation process. In the case of OCS, for example, the relaxation rates in Table 3 show that a photodissociation time scale of 1-10 ns is required to take full advantage of photodissociation through the $2\nu_2$ intermediate vibrational level, whereas a time scale of a few hundred nanoseconds is adequate for the lower energy ν_1 and ν_2 levels.

Enrichment of the mass 16, 17 and 18 isotopes of oxygen by two-step photodissociation of OCS in liquid Ar has been investigated experimentally, using a CO_2 laser to vibrationally excite the $2\nu_2$ vibrational level and a KrF excimer laser for photodissociation. A detailed study of the dependence of enrichment on various experimental parameters was made, and results were compared with computer modeling calculations. Details of the study are given elsewhere.¹⁶ The experimentally measured oxygen isotope enrichment factors of 3-4 in the CO photodissociation product were a factor of 2-3 below theoretically optimum values. Modeling calculations showed that enrichments were reduced by rapid vibrational relaxation of the $2\nu_2$ vibrational level.

Our experimental studies have established that the main isotope enrichment limiting factor in the two-step, liquid-phase process is the rapid collisional relaxation of the IR laser-pumped vibrational level prior to UV photodissociation. The limitation is expected to become more serious if higher-energy, more rapidly relaxed, vibrational levels are used as the intermediate level in the two-step process. The rapidity of vibrational relaxation can in principle be

Table 3. OCS Vibrational Relaxation in Cryogenic Solutions. ^a

vibr level	solvent	T	1/K _{i,M} ^b	k _{i,M} ^c	P _{IBC,hs} ^d
i	M	(K)	(ns)	(10 ⁻¹⁶ cm ³ /s)	
2ν ₂ ^e	Ar	84.4	9	52	8.5 × 10 ⁻⁶
	Kr	116.8	17	34	6.0 × 10 ⁻⁶
	Xe	162.0	6	120	1.2 × 10 ⁻⁵
	O ₂	84.4	-	-	-
ν ₁	Ar	84.4	145 ± ³⁵ ₄₀	3.3 ± ^{1.2} _{0.7}	5.3 ± ^{2.0} _{1.1} × 10 ⁻⁷
	Kr	116.8	235 ± ⁸⁰ ₆₅	2.4 ± ^{1.0} _{0.5}	4.3 ± ^{1.7} _{0.8} × 10 ⁻⁷
	Xe	162.0	250 ± ¹⁵⁰ ₉₅	2.8 ± ^{1.6} _{1.1}	2.9 ± ^{1.8} _{1.1} × 10 ⁻⁷
	O ₂	84.4	22 ± ⁸ ₇	21. ± ⁸ ₆	2.6 ± ^{1.1} _{0.7} × 10 ⁻⁶
ν ₂	Ar	84.4	355 ± 35	1.33 ± 0.13	2.2 ± 0.2 × 10 ⁻⁷
	Kr	116.8	660 ± 60	0.87 ± 0.08	1.6 ± 0.2 × 10 ⁻⁷
	Xe	162.0	625 ± 75	1.10 ± 0.13	1.2 ± 0.1 × 10 ⁻⁷
	O ₂	84.4	53 ± 9	8.6 ± 1.5	1.1 ± 0.2 × 10 ⁻⁶

^a From Refs. 17 and 18.^b Reciprocal of relaxation rate.^c Rate constant, k_{i,M} = K_{i,M}/N_M, where N_M is the solvent number density.^d Relaxation probability, P_{IBC,hs} = K_{i,M}/K_{hs,M}, where K_{hs,M} is the hard sphere collision rate with the solvent.^e Values for the 2ν₂ level are estimated from theory and gas phase data. An upper limit of 70 ns was determined experimentally for 1/K_{2ν₂,M} for all four cryogenic solvents.

counterbalanced by using lasers with very short pulse widths.

A computer model of the liquid-phase isotope enrichment process has been developed to evaluate the effects of laser characteristics (and other parameters) on enrichment factors.¹⁸ Lasers with superior features, such as very short pulse widths, may not be immediately available in the laboratory, but may be available commercially, or may be developed in the near future. With experimental knowledge of the basic spectroscopic and kinetic parameters of the enrichment process, the enrichments achievable with various laser characteristics and other sample conditions can be computer modeled with some confidence. An extensive modeling study has been reported for two-step, liquid-phase photodissociation of OCS.¹⁸ As an example of the results, Figure 8 shows the enrichment factors expected for ¹³C, ¹⁷O and ¹⁸O, using 1-2 ns pulse-width lasers matched to the vibrational relaxation time of the laser-pumped, 2ν₂ intermediate vibrational level. The calculation indicates the IR laser fluence required for

Figure 8. Computer-Modeled, Liquid-Phase, Carbon and Oxygen Enrichment Factors. Legend "x/y" indicates enrichment of isotope x with respect to y. CO₂ laser lines are indicated. From Ref. 18.

maximum isotope enrichment. The computer modeling study also evaluates isotopically selective, two-step photodissociation through the ν_1 vibrational intermediate level of OCS, using pulsed lasers. A continuous, two-step process that employs cw lasers is also modeled.

VI. CONCLUSIONS

The research conducted under this contract/grant presents a comprehensive picture of the two-step, photodissociative, isotope enrichment process for selected molecules in the gas and cryogenic liquid phases. Basic photophysical, kinetic, and spectroscopic data have been acquired, and isotope enrichments have been demonstrated. Based on the collective results, some general observations about the isotopically selective, two-step photodissociation of small molecules are given here.

Vibrational enhancement of the molecular photodissociation cross section is the basis of the two-step isotope enrichment process. Our measurements of the enhanced cross sections for several small molecules have quantified the important effect of vibrational excitation in the red wing of bound-continuous photodissociation bands. The enhancement of photodissociation cross sections by vibrational excitation is much larger than one would estimate by simply shifting the photodissociation spectrum by a quantum of vibrational energy. The results indicate that isotope enrichment factors of approximately 10-100 should be achieved by two-step photodissociation of typical small molecules, while maintaining reasonable photodissociation yields. The steep increase in cross section enhancement with vibrational energy suggests that photodissociation through the highest energy, infrared active, fundamental vibrational level of a molecule should be particularly effective. Thus, the asymmetric stretching vibrations of linear triatomic molecules should be ideal intermediate levels for a two-step photodissociation process in gas-phase systems.

High isotopic selectivity can be achieved in the vibrational excitation step of the two-step photodissociation process. IR lasers with bandwidths of a few hundredths of a cm^{-1} are required in the gas phase to discriminate among the interlaced rovibrational lines of different isotopomers of linear triatomic molecules. Gas-phase enrichment factors in the present study have probably been reduced by broad laser bandwidth. Rotational bottlenecking is not found to be a serious problem in the gas phase. Experimental IR absorption measurements and computer modeling calculations show that rapid rotational relaxation by an inert bath gas allows near-saturation of the vibrational transition. A significantly larger IR laser bandwidth of several tenths of a cm^{-1} is adequate in the liquid phase to isotopically discriminate among collapsed vibrational features. Rotational relaxation is irrelevant in the cryogenic liquid phase.

Vibrational relaxation is not an enrichment limiting factor in the gas phase. Although high pressures of rare gas may be required to provide rapid rotational relaxation during the IR laser-pumping pulse, the rate constants for vibrational relaxation of the relevant levels of triatomic molecules by rare gases are not large enough to limit two-step photodissociation. Vibrational relaxation can be a serious limiting factor in the cryogenic liquid phase. Sub-nanosecond relaxation times are likely for vibrational levels above perhaps $1500\text{-}2000\text{ cm}^{-1}$

in energy, although experimental data for the higher energy levels is sparse. The vibrational relaxation times determine the maximum usable pulse widths of IR and UV laser sources. Since practical laser pulse widths are probably limited to the range of a few nanoseconds, two-step photodissociation in liquids may be limited to the use of intermediate vibrational levels of low energy, which produce only moderate vibrational enhancements of the photodissociation cross section. More experimental data are needed on vibrational relaxation times in cryogenic liquids.

Temperature is an important factor in the two-step enrichment process in the gas phase. Temperatures of 150-200K are necessary to prevent significant thermal population and isotopically non-selective photodissociation of low energy bending vibrational levels of typical triatomic molecules. For most molecular targets, an improvement of roughly a factor of two in the enrichment factor can be probably be achieved by the simple expedient of lowering sample temperature from room temperature to 150-200K.

Rapid chemical scavenging of active photofragments is essential to the two-step photodissociation process. In the case of sulfur atoms generated by photodissociation of OCS, highly efficient scavengers, such as NO_2 and ethylene, have been defined for the gas phase. Other active fragments, such as O atoms, may require different scavenging systems. In the liquid phase, three-body recombination of atomic fragments can be considered as an efficient scavenging technique. The low temperature in a cryogenic liquid retards bimolecular scavenging reactions, but together with the high solvent density enhances recombination reactions. In the case of sulfur atoms, scavenging by recombination is found experimentally to have the added advantage of producing a convenient, solid, sulfur precipitate.

Computer modeling of the two-step enrichment process is an important complement to experimental measurements. In some complicated systems, modeling has quantitatively identified the primary isotope scrambling processes. A particularly useful aspect of a good computer model is the ability to predict enrichments for experimental conditions that are achievable, but for various reasons not immediately accessible in the laboratory. For example, the beneficial effects of improving the temporal and spectral characteristics of a laser source can be assessed. Reliable modeling requires the input of basic spectral, photophysical, and kinetic parameters. It is important that the measurement of fundamental optical cross sections and kinetic rate constants be a part of an isotope enrichment study.

The results of the present study suggest an optimum system for achieving practical, high enrichments of C, N, O, S, and possibly other light isotopes. The optimum system is a fairly simple extension of the techniques already applied to OCS. From a practical standpoint, the simplest environment for an enrichment system is the low-temperature gas phase (i.e., 150-200K). Isotope enrichment would be based on two-step photodissociation of OCS, N_2O , or other linear triatomic molecules. A relatively low-power, tunable, 4.5- to 5.0- μm laser source would pump a selected isotopomer of the target molecule to the ν_3 (asymmetric stretching)

vibrational level. Since the absorption band strength of the ν_3 vibration is typically extremely large, the pulse energy requirement on the IR laser is small (a few tens of millijoules per pulse). Such tunable lasers exist for somewhat shorter IR wavelength regions and can undoubtedly be developed for the 5- μm region. Subsequent UV photodissociation would be by an excimer laser, or frequency-mixed dye laser, at the red edge of the molecule's UV photodissociation band. The vibrational enhancement of the photodissociation cross section and consequently the enrichment factor are expected to be very large for excitation of the ν_3 vibrational level. For a system based on OCS/Ar/scavenger mixtures, scavenging of S atoms would be by ethylene or higher alkenes. In $\text{N}_2\text{O}/\text{Ar}$ mixtures, simple recombination of electronically relaxed $\text{O}(^3\text{P})$ atoms would result in N_2 and O_2 products with no isotope scrambling. It is likely that enrichment factors on the order of ~ 50 could be achieved in such a system and that enrichment of the rarer isotopes could be economically practical.

REFERENCES

1. For early reviews see J. P. Aldridge, J. H. Birely, C. D. Cantrell, and D. C. Cartwright, "Experimental and Theoretical Studies of Laser Isotope Separation," Report No. LA-UR-75-2368, Los Alamos Scientific Laboratory, Los Alamos, NM, 1975; *Chemical and Biochemical Applications of Lasers, Vol. III*, edited by C. B. Moore, Academic, New York, 1977; V. S. Letokhov, *Nature* **277**, 605 (1979).
2. E. B. Turner, G. Emanuel, and R. L. Wilkins, "The NEST Chemistry Computer Program, Volume 1," Report No. TR-0059(6240-20)-1, Vol. 1, The Aerospace Corporation, El Segundo, CA, 30 July 1970.
3. W. C. Hwang, R. F. Kamada, and C. C. Badcock, *Int. J. Chem. Kinet.* **9**, 969 (1977).
4. W. C. Hwang and R. F. Kamada, "Modeling Study of Rotational Relaxation and Absorption," Report No. ATR-77(8127)-3, The Aerospace Corporation, El Segundo, CA, 20 January 1977.
5. C. C. Badcock, W. C. Hwang, J. F. Kalsch, and R. F. Kamada, *Appl. Phys. Lett.* **32**, 363 (1978).
6. P. F. Zittel and D. D. Little, *J. Chem. Phys.* **71**, 713 (1979).
7. P. F. Zittel and D. D. Little, *J. Chem. Phys.* **71**, 2748 (1979).
8. P. F. Zittel and D. D. Little, *J. Chem. Phys.* **71**, 723 (1979).
9. P. F. Zittel and D. D. Little, *J. Chem. Phys.* **72**, 5900 (1980).
10. P. F. Zittel and D. D. Little, *Chem. Phys.* **63**, 227 (1981).
11. P. F. Zittel and L. A. Darnton, *J. Chem. Phys.* **77**, 3464 (1982).
12. P. F. Zittel, L. A. Darnton, and D. D. Little, *J. Chem. Phys.* **79**, 5991 (1983).
13. P. F. Zittel and D. E. Masturzo, *J. Chem. Phys.* **85**, 4362 (1986).
14. P. F. Zittel and M. A. Sedam, *Chem. Phys. Lett.* **148**, 486 (1988).
15. P. F. Zittel and V. I. Lang, *J. Photochem. Photobiol. A: Chem.* **56**, 149 (1991).
16. P. F. Zittel and M. A. Sedam, *J. Phys. Chem.* **94**, 5801 (1990).
17. P. F. Zittel, *J. Phys. Chem.* **95**, 6802 (1991).
18. P. F. Zittel, "A Computer Modeling Study of Isotopically Selective, Laser Photodissociation of OCS in Cryogenic Solutions," Report No. ATR-91(7171)-2, The Aerospace Corporation, El Segundo, CA, 23 December 1991.
19. V. I. Lang and P. F. Zittel, unpublished data.

APPENDIX: LIST OF PUBLICATIONS

W. C. Hwang, R. F. Kamada, and C. C. Badcock, "Modeling Study of Two-Step Photodissociation Method of Isotope Enrichment," *Int. J. Chem. Kinet.* **9**, 969 (1977).

W. C. Hwang and R. F. Kamada, "Modeling Study of Rotational Relaxation and Absorption," Report No. ATR-77(8127)-3, The Aerospace Corporation, El Segundo, CA, 20 January 1977.

C. C. Badcock, W. C. Hwang, J. F. Kalsch, and R. F. Kamada, "Isotopic HCl Transfer Laser," *Appl. Phys. Lett.* **32**, 363 (1978).

P. F. Zittel and D. D. Little, "Photodissociation of Vibrationally Excited Hydrogen Bromide," *J. Chem. Phys.* **71**, 713 (1979).

P. F. Zittel and D. D. Little, "Diffusion Coefficients for Br₂ in NO and Br in Ar at 298°K," *J. Chem. Phys.* **71**, 723 (1979).

P. F. Zittel and D. D. Little, "Experimental and Computer Modeling Study of ⁸¹Br Isotope Enrichment by Isotopically Selective, Two-Step Laser Photodissociation of HBr," *J. Chem. Phys.* **71**, 2748 (1979).

P. F. Zittel and D. D. Little, "Photodissociation of Vibrationally Excited Ozone," *J. Chem. Phys.* **72**, 5900 (1980).

P. F. Zittel and D. D. Little, "Photodissociation of Vibrationally Excited CH₃Br," *Chem. Phys.* **63**, 227 (1981).

P. F. Zittel and L. A. Darnton, "Sulfur-33 Isotope Enrichment by Isotopically Selective, Two-Step, Laser Photodissociation of OCS," *J. Chem. Phys.* **77**, 3464 (1982).

P. F. Zittel, L. A. Darnton, and D. D. Little, "Separation of O, C, and S Isotopes by Two-Step, Laser Photodissociation of OCS," *J. Chem. Phys.* **79**, 5991 (1983).

P. F. Zittel and D. E. Masturzo, "Photodissociation of Vibrationally Excited OCS," *J. Chem. Phys.* **85**, 4362 (1986).

P. F. Zittel and M. A. Sedam, "Vibrational Relaxation of OCS(ν_3)," *Chem. Phys. Lett.* **148**, 486 (1988).

P. F. Zittel and M. A. Sedam, "Infrared Spectra, Vibrational Relaxation, and Isotopically Selective Photodissociation of OCS in Ar(l) and O₂(l) Cryogenic Solutions," *J. Phys. Chem.* **94**, 5801 (1990).

P. F. Zittel and V. I. Lang, "Isotopically Selective Photodissociation of Gas Phase OCS at Low Temperatures," *J. Photochem. Photobiol. A: Chem.* **56**, 149 (1991).

P. F. Zittel, "Vibrational Relaxation and Infrared Spectra of O₃ and OCS in Cryogenic Liquid Solutions," *J. Phys. Chem.* **95**, 6802 (1991).

P. F. Zittel, "A Computer Modeling Study of Isotopically Selective, Laser Photodissociation of OCS in Cryogenic Solutions," Report No. ATR-91(7171)-2, The Aerospace Corporation, El Segundo, CA, 23 December 1991.

END

**DATE
FILMED**

6/18/92