1678

179
03/4/80 MR.

JULY 1980

PPPL-1678

UC-20g

16. 1592

SOLVING THE CHIRIKOV TAYLOR MODEL
BY THE METHOD OF PATHS IN
FOURIER SPACE

BY

A. B. RECHESTER, M. N. ROSENBLUTH,
AND R. B. WHITE

MASTER

**PLASMA PHYSICS
LABORATORY**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

**PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY**

This work was supported by the U.S. Department of Energy
Contract No. DE-AC02-76-CHO 3073. Reproduction, translation,
publication, use and disposal, in whole or in part,
by or for the United States government is permitted.

Solving the Chirikov Taylor Model by the Method of Paths in Fourier Space

A. B. Rechester
Plasma Fusion Center
Cambridge, MA 02139

M. N. Rosenbluth
Institute for Advanced Study
Princeton, NJ 08540

R. B. White
Plasma Physics Laboratory
Princeton University
Princeton, NJ 08544

A compact path-diagram method has been introduced for the calculation of velocity moments of a probability function. This method is complementary to the approach developed in Ref. 1. It is applied to the Chirikov-Taylor model. A number of new results have been derived and verified by numerical computations.

DISCLAIMER

This document contains neither recommendations nor conclusions of the United States Government. It is the property of the United States Government and is loaned to your agency; it and its contents are not to be distributed outside your agency. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

I. A PROBABILITY DESCRIPTION FOR EVOLUTION IN PHASE SPACE

The purpose of this work is a further development of a probabilistic method outlined in Ref. 1. As before, we will consider the simple case of charged particles moving in a field of electrostatic plane waves. The equations of motion for the Chirikov-Taylor model are¹⁻³

$$x_{t+1} = x_t + v_t \quad (1)$$

$$v_{t+1} = v_t + \epsilon \sin x_{t+1} . \quad (2)$$

All variables here are dimensionless, x represents the coordinate, and v the velocity, and time t takes on integer values $t = 0, 1, 2, \dots$. The parameter ϵ is related to the electric field amplitude. Without loss of generality, we may assume it to be a positive number. We will also assume periodic boundary conditions in x , $0 \leq x \leq 2\pi$. Following Ref. 1, we substitute the deterministic description given by Eqs. (1), (2) with a probabilistic one by introducing an additional random step δx_t in Eq. (1). This variable has a probability distribution given by:

$$G(\delta x_t, v_t) = (2\pi\sigma)^{-1/2} \sum_{n=-\infty}^{\infty} \exp \left[- \frac{(\delta x_t - v_t + 2\pi n)^2}{2\sigma} \right] . \quad (3)$$

Equation (3) can be derived by introducing a diffusion term ($\sigma/2$ the diffusion coefficient) into the Vlasov equation.¹ Let

$P(x, v; t)$ be a time dependent probability function of phase space points x, v with the initial condition given by

$$P(x, v, 0) = \frac{1}{2\pi} \delta(v - v_0) \quad (4)$$

Then according to Eqs. (1) - (3) the time evolution of P in the interval $t - 1, t$ is given by

$$P(x, v, t) = \int_0^{2\pi} G(x - x', v) P(x', v + \varepsilon \sin x', t - 1) dx'. \quad (5)$$

The Fourier transformation of this function in x and v can be written as

$$P(x, v, t) = \frac{1}{(2\pi)^2} \sum_{m=-\infty}^{\infty} \int_{-\infty}^{\infty} dk a_m^t(k) \exp[i(mx + kv)]. \quad (6)$$

From (4) it follows that

$$a_m^0(k) = \delta_{m0} \exp(-ikv_0) \quad , \quad (7)$$

where $\delta_{m,n}$ is a Kronecker function. The relation between the Fourier amplitudes $a_m^t(k)$ at times t and $t - 1$ can be easily derived from Eqs. (5) and (6):

$$a_m^t(k) = \sum_{\ell=-\infty}^{\infty} J_{\ell}(|k'| \varepsilon) \exp[-(\sigma/2)m^2] a_{m'}^{t-1}(k'). \quad (8)$$

Here J_{ℓ} are Bessel functions and:

$$k' = k + m , \quad (9)$$

$$m' = m - \ell \operatorname{sign} k' , \quad (10)$$

and

$$\operatorname{sign} k = \begin{cases} +1 & k > 0 \\ -1 & k < 0 . \end{cases}$$

Introduce now velocity moments

$$\langle v^n \rangle_t = \int_0^{2\pi} dx \int_{-\infty}^{\infty} dv v^n P(x, v, t) . \quad (11)$$

With the use of (6) they can be expressed as

$$\langle v^n \rangle_t = \lim_{k \rightarrow 0+} \left(i \frac{\partial}{\partial k} \right)^n a_0^t(k) . \quad (12)$$

We will consider in this paper the second moment which is related to the diffusion coefficient D by¹

$$D = \frac{\langle (v - v_0)^2 \rangle_T}{2T} . \quad (13)$$

Throughout this paper we assume the final time T to be a large integer, $T \gg 1$.

II. PATHS IN FOURIER SPACE - ANALYTICAL CALCULATIONS

Starting from the initial Fourier amplitude a_m^0 we can find the final amplitude a_0^T as a product of T transitions. Every transition can be considered as a path in m, k space. In fact, we will go backward in time. Since we are interested in $a_0^T(k \rightarrow 0)$ (see Eq. 13), a path must start at the origin $m = 0, k = +0$. The simplest path is one which never leaves the origin. It contributes

$$a_0^T(k) = [J_0(k\varepsilon)]^T a_0^0(0) \quad (14)$$

Then from (13) we find

$$D_{QL} = \frac{\varepsilon^2}{4} \quad (15)$$

where we have used the expansion of J_0 for small argument

$$J_0(k\varepsilon) \approx 1 - \left(\frac{k\varepsilon}{2}\right)^2 \quad (16)$$

This simplest path corresponds to a value of diffusion given by quasilinear theory.^{1,3}

Consider now paths which do leave the origin. According to Eqs. (8) - (10) an amplitude $a_m^t(k)$ may be connected by a single path step to $a_{m'}^{t-1}(k')$ where (m', k') must lie on the vertical line $k' = k + m$. As a result of such a transition, the amplitude accumulates a factor of $J_0(|k'| \varepsilon) \exp[-(\sigma/2)m^2]$. This simple geometrical rule, illustrated in Fig. 1, allows

construction of all possible transition paths. For the case depicted in Fig. 1, the transition factor is equal to $J_1(3\epsilon) \exp(-2\sigma)$.

Note further that in leaving or returning to the origin, we obtain factors $J_\ell(k\epsilon)$. Since we are interested in the limit $k \rightarrow +0$, only $\ell = \pm 1$ are of low enough order in k to contribute to the diffusion. It is also clear that we do not need to consider any paths which leave or return to the origin more than once (too high order in k).

We will consider primarily the case $\epsilon \ll 1$, since the large ϵ case has been discussed already.¹ We utilize the fact that the Bessel functions $J_\ell(\epsilon)$ are proportional to ϵ^ℓ when $\epsilon \ll 1$. As we will see later, only paths involving $\ell = 0, \pm 1$ contribute to lowest order in ϵ .

Consider a path represented on Fig. 1. It starts at the origin, stays there for s transitions, then moves to an ending point $m = 0$, $k = p$ (the case $p = 3$ is depicted on Fig. 1) and then remains $T-s-p$ transitions at this point. The contribution from such a path for the case $p > 0$ is equal to

$$a_0^T(k) = J_0^S(k\epsilon) J_{-1}(k\epsilon) \prod_{i=1}^p J_0[(i+k)\epsilon] \exp[-(\sigma/2)p] \\ J_1[(p+k)\epsilon] J_0^{T-s-p}[(p+k)\epsilon] a_0^O(p+k). \quad (17)$$

Path points on the k axis can remain stationary an arbitrary number of transitions, and we indicate these points with a cross.

Factors $J_0^S(k\varepsilon)$ give 1 in the limit $k \rightarrow 0$, and we will omit them in the following.

The diffusion is readily calculated to lowest order in ε for two limits, characterizing early and late times. For the early time limit assume that $p\varepsilon \ll 1$ and $1 \ll T \ll (2/p\varepsilon)^2$ so that $J_0^{T-S-P}(p\varepsilon) \approx 1$. Then with the use of the identity $J_{-1}(\varepsilon) = -J_1(\varepsilon)$ and $J_1(\varepsilon) = \varepsilon/2$, $\varepsilon \ll 1$, we find

$$a_0^T(k) = - \frac{Tk(p+k)\varepsilon^2}{4} \exp[-(\sigma/2)p - ipv_0] .$$

The factor of T in front of this equation comes from the summation over s .

The case $p < 0$ can be considered in a similar way. Using Eq. (13) and summing over all $p = +1, +2, \dots$ we can get the contribution to D which we combine with expression (15) to find

$$D = \frac{\varepsilon^2}{4} \frac{[1 - \exp(-\sigma)]}{(1 - 2 \cos v_0 \exp[-(\sigma/2)] + e^{-\sigma})} . \quad (19)$$

The approximate region of applicability of this formula is

$$1 \ll T \ll (2/\varepsilon)^2, \quad \sigma \gg \varepsilon . \quad (20)$$

It is easy to show that contributions from other types of paths which end up outside of the origin are small.

Consider now the limit $T \gg (2/\epsilon)^2$. Then the factors $J_0^{T-S-P}(p\epsilon)$ in Eq. (17) are small. Due to this, only paths which come back to the origin contribute. The simplest path of this type is shown in Fig. 2a. Its contribution is equal to

$$a_0^T(k) = T J_{-1}(k\epsilon) J_1(\epsilon) \exp[-(\sigma/2)] J_0^n(\epsilon) J_1(\epsilon) \exp(-\sigma/2) J_{-1}(k\epsilon) a_0^0. \quad (21)$$

Summing over all $n = 0, 1, 2, \dots$ we find

$$a_0^T(k) = T \left(\frac{k\epsilon}{2} \right)^2 \frac{J_1^2(\epsilon)}{1 - J_0(\epsilon)} e^{-\sigma}. \quad (22)$$

In the limit of $\epsilon \ll 1$ its contribution to the diffusion is equal to $-(\epsilon^2/4) \exp(-\sigma)$. Exactly the same value arises from a symmetric path leaving the origin in a downward direction.

Let us now turn to a more complicated path, depicted on Fig. 2b. Every crossing of a path with the abscissa at k_c gives a factor similar to that of Eq. (22)

$$\pm \frac{J_1^2(|k_c|\epsilon)}{1 - J_0(|k_c|\epsilon)} \approx \pm 1, \quad k_c \epsilon \ll 1.$$

If we designate by k_R the last cross to the right different from the origin, then the number of crosses between the origin and that cross is $n_R = k_R - 1 \geq 0$. The same applies for the number

of crosses between the origin and the last one to the left,
 $n_L = -k_L - 1 \geq 0$. The total number of such crosses is

$$n = n_R + n_L. \quad (23)$$

In the case $n > 0$, the contribution from any given path of the type 2b is equal to

$$a_0^T(k) = (-1)^{d-2} T\left(\frac{k\varepsilon}{2}\right)^2 \exp[-\sigma(n+2)]. \quad (24)$$

Here d is the total number of diagonal lines on the path. It is easy to see that for a given n_R and n_L the total number of different paths which have a positive sign in Eq. (24) is equal to $\sum_{i=0}^n C_{2n}^{2i}$, while the number of paths with a negative sign is $\sum_{i=1}^n C_{2n}^{2i-1}$, where C_n^i is a binominal coefficient. Due to the identity

$$\sum_{i=0}^{2n} C_{2n}^i (-1)^i = (1-1)^{2n} = 0, \quad (25)$$

we have exact cancellation of contributions from these paths. Thus only paths with $n = 0$ contribute. They are drawn on Fig. 2c. It is straightforward to show that their contribution to the diffusion is given by

$$\frac{\varepsilon^2}{4} (-1)^p e^{-p\sigma} \quad p = 1, 2, 3 \dots \quad (26)$$

Here the integer p corresponds to the number of half circles

around the origin. Also the contributions from paths in clockwise and anticlockwise directions are equal.

Summing over P and combining with Eq. (15), we can finally obtain, to lowest order in ϵ , for $T \gg (2/\epsilon)^2$,

$$\frac{D}{D_{QL}} = \tanh(\sigma/2) \quad . \quad (27)$$

The main result of Ref. 1, Eq. (20), can be easily rederived using the method described here.

III. NUMERICAL SUMMATION OF PATHS IN FOURIER SPACE

The analytical calculation of D/D_{QL} to higher order in ϵ involves the summation of a rather large set of paths in Fourier space, and we have carried out this summation numerically. From Eq. (8), we find that symmetry or antisymmetry of $a_m^t(k)$ under $m, k \rightarrow -m, -k$ is propagated from $t-1$ to t . Taking $a_m^0(k)$ as given by Eq. (7) with $V_0 = 0$, we thus have that $a_m^t(k)$ is symmetric under this reflection. We are interested in long time behavior so the value of V_0 is irrelevant. We then need keep only $k \geq 0$. Now truncate in Fourier space through

$$\begin{aligned} -M &\leq m \leq M \\ 0 &\leq k \leq K \end{aligned} \quad (28)$$

and replace (m, k) with the single index $n = (2M+1)k+m$ where $1 \leq n \leq N$, $N = (2M+1)K+M$. From the preceding analysis regarding paths in Fig. 2, we can take the path to originate by leaving

the origin through $m = -1, k = 1$, i.e., $a_n^1 = \delta_{n,2M} \exp(-\sigma/2)$. (We are now following the path forward in time, i.e., backwards along the arrows of Fig. 2.) The term $a_0^t(0)$ is excluded since the paths considered return to the origin only once, passing through $(m,k) = (1,0)$. The diffusion is given by

$$\frac{D}{D_{QL}} = 1 - 2A_1, \quad (29)$$

where $A_1 = \lim_{t \rightarrow \infty} a_1^t$, since the path must terminate by entering the origin through $(m,k) = (1,0)$. From Eq. (8), the vector A satisfies the matrix equation

$$A = (1 + M + M^2 + \dots) a^1 = (1 - M)^{-1} a^1, \quad (30)$$

where

$$M_{ij} = \exp[-(m^2\sigma/2)] [J_{m-m'}(k'\epsilon) \delta_{k',m+k} - J_{-(m+m')}(k'\epsilon) \delta_{k',-m-k}] \quad (31)$$

with

$$M = \begin{cases} i \bmod (2M+1) & m \leq M \\ i \bmod (2M+1) - (2M+1) & i \bmod (2M+1) > M \end{cases} \quad (32)$$

$$k = \frac{i-m}{2M+1} \quad (33)$$

and similar equations relating m', k' to j . Gaussian elimination is then used to find the diffusion through Eq. (29). As numerical checks on this method, we used the trivial case $M = 1, K = 1$, for which M_{ij} is 4×4 , as well as confirming that for

$\epsilon \ll 1$, $\sigma \ll 1$ the case $M = 1$, $K \gg 1$ converges for large K to the partial result

$$\frac{D}{D_{QL}} = \tanh \frac{\sigma}{2} + \frac{3\epsilon^2}{16\pi^2} \quad , \quad (34)$$

obtained by summing all paths restricted within $-1 \leq m \leq 1$. Equation (34) is not a correct expression for D/D_{QL} to order ϵ^2 ; paths with excursions out to $m = \pm 2$ cancel this ϵ^2 term. In addition, we have compared this method to results obtained by numerically advancing the Chirikov-Taylor mapping for values of ϵ and σ for which this is practical. We have not been able to analytically derive the lowest order corrections (in ϵ) to Eq. (27). The numerical results indicate that these corrections are not analytic in σ . These results are discussed in the following section.

IV. NUMERICAL RESULTS FOR SMALL ϵ

We turn now to the results of numerical computations. Two methods have been used. First, where practical, we have calculated the diffusion directly through Eq. (13) by advancing Eqs. (1,2) with the introduction of a random step in x having a normal distribution of mean σ . A large number of particles N_p is used, due to the presence of fluctuations in D proportional to $N_p^{-1/2}$. Full advantage of the vector processing capabilities of the MFE CRAY 1 have been used, but it is still not practical to obtain values for the diffusion in this manner when both ϵ and σ are quite small. Approximately, 2 μ sec of cpu time is

necessary per particle step. In Figure 3, is shown the ratio of D to the quasilinear value as a function of σ for $\epsilon = 0.06$, and two values of V_0 . Each point is obtained by advancing 1000 particles initially distributed uniformly in x at $V = V_0$. For values of T such that $1 \ll T \ll (2/\epsilon)^2$, the observed values agree with the analytical result given by Eq. (19), shown with a solid curve. Also shown is the result for $T \gg (2/\epsilon)^2$, when the particles have diffused sufficiently far in phase space that the initial correlations associated with the value of V_0 have been lost. The diffusion is then seen to be described by Eq. (27). We have also investigated the small ϵ , small σ regime. Results obtained by advancing Eqs. (4,5) to $T = 29,000$ are shown in Fig. 4. We have used this method for $0.6 < \epsilon < 1.5$. At small ϵ large times are necessary to obtain convergent results due to the very small rates of diffusion, and for $\epsilon < 0.6$, it is prohibitive to find the correction to Eq. (27) in this manner. At $\epsilon = 0.7$ for example, the diffusion approaches its asymptotic limit as

$$\frac{D}{D_{QL}} = \left(\frac{D}{D_{QL}} \right)_{\text{asym}} + \frac{40}{T} \quad , \quad (35)$$

for the initial conditions used. Here we have used 2000 particles initially distributed uniformly in x for $0 \leq x \leq 2\pi$, and randomly in V for $0 \leq V \leq 2\pi$. A detailed investigation of the dependence of D on σ and ϵ is not feasible, and $\sigma = 10^{-2}$, $\epsilon = 0.5$ is practically the lower bound.

To complement these results, we have used the numerical path summation technique discussed in Section IV. In Figure 5, is shown the dependence of the numerical results on the Fourier space truncation parameters M, K , for ϵ below the Chirikov threshold.² For fixed M convergence is seen to proceed stepwise in $1/k$, indicating the existence of critical scale lengths in phase space for contributions to a particular order in ϵ . The values $K = 30, M = 10$ result in a 640×640 dimensional matrix M_{ij} , which is approximately the computational upper bound. A determination of the diffusion requires approximately two minutes of CRAY time with M, K this large, and the practical ranges of ϵ, σ are significantly extended over what is possible by directly using the Chirikov-Taylor mapping. For $\epsilon > 1$, except in the vicinity of $\epsilon = 2\pi$, much smaller values of K, M suffice to give convergent results. The large extent in Fourier space of the contributing paths undoubtedly indicates that for $\epsilon < 1$ the diffusion is spatially localized, perhaps into filaments attached to separatrices. In Figure 4, are shown the results obtained in the small ϵ regime, for different values of σ . The $\sigma = 0$ results are consistent with the vanishing of the diffusion below the Chirikov limit. These results are seen to agree quite well with those obtained by stepping Eqs (4,5).

In Figure 6, is shown the correction to the lowest order expression for D , given by Eq. (27). Determination of this correction below $\epsilon = 0.5$ is not numerically feasible with the present schemes. Assuming the corrections to be of the form of even powers in ϵ , a result expected by the nature of terms

appearing in the path summations, we find that these results are reasonably well fit by

$$\frac{D}{D_{QL}} = a\sigma^{2/3} \epsilon^2 + b\epsilon^4 \quad (36)$$

with $a \approx 1/8$ and $b \approx 2 \times 10^{-3}$. The dependence of b on σ cannot be determined from the data available, except that it must be quite weak. Of course, b must vanish as $\sigma \rightarrow 0$.

In Figure 7, is shown D/D_{QL} for $0 < \epsilon < 20$, and $\sigma = 10^{-3}$ obtained with the numerical Fourier path summation technique. We find that large M, K is necessary to observe the peak at $\epsilon \approx 2\pi$, where D becomes more than three times the quasilinear value. This indicates that this peak is associated with streaming localized in phase space. Chirikov² has pointed out the existence of domains with regular motion for $\epsilon \approx 2\pi n$, n integer. Also shown is the analytic expression for large ϵ derived in Ref. 1,

$$\frac{D}{D_{QL}} = 1 - 2J_2(\epsilon) e^{-\sigma} - 2J_1^2(\epsilon) e^{-\sigma} + 2J_3^2(\epsilon) e^{-3\sigma} . \quad (37)$$

V. CONCLUSION

Determination of the time evolution of the solution of the Vlasov equation through the method of paths in Fourier space greatly extends the results obtainable through direct stepping of the Chirikov-Taylor mapping. The method provides lowest order closed form analytic expressions for the diffusion, and the means of calculating to higher order numerically. We believe that it

will prove useful in the examination of other problems involving chaotic or turbulent dynamics.

ACKNOWLEDGMENTS

We would like to acknowledge valuable conversations with J. Greene, and C. Oberman, and valuable numerical assistance from M. Petravic.

This work was supported by United States Department of Energy Contract No. DE-AC02-76-CHO3073 and DE-AS02-78ET53074.

REFERENCES

1. A. B. Rechester and R. B. White
Phys. Rev. Lett. 44, 1586 (1980).
2. B. V. Chirikov
Phys. Reports 52, 263 (1979).
3. G. M. Zaslavsky and B. V. Chirikov
Usp. Fiz. Nauk 14, 195 (1972),
(Sov. Phys. Usp. 14, 549 (1972)).

(PPPL-802109)

Fig. 1. Examples of paths in Fourier space contributing to the probability amplitude. A single path step can connect the point (m, k) to any point (m', k') with $k' = k + m$. The crosses indicate points at which the path can remain for several time steps.

(PPPL-802111)

Fig. 2. Paths in Fourier space which contribute to the long time $(T \gg (2/\epsilon)^2)$ diffusion. They must leave the origin through the point $(m, k) = \pm(1, 0)$ and return through $\pm(-1, 1)$.

(PPPL-802110)

Fig. 3. The ratio of the diffusion to the quasilinear value as a function of σ obtained using the Chirikov-Taylor mapping. The particles were initially uniformly distributed in x and located at $V = V_0$. For $1 \ll T \ll 2/\epsilon^2$ the results agree with Eq. (19), whereas, for $T \gg 2/\epsilon^2$, the diffusion is given by Eq. (27). Here $\epsilon = 0.06$.

(PPPL-802139)

Fig. 4. The diffusion as a function of ϵ for small ϵ , and various values of σ . The points were obtained using the Chirikov-Taylor mapping directly, and the curves by the path summation technique of Section IV. For $\sigma = 0$, the diffusion vanishes for ϵ less than the Chirikov threshold value.

(PPPL-802143)

Fig. 5. The convergence of the Fourier path summation method. Shown are values of $D/D_{0L} - \tanh \sigma/2$ as a function of $1/K$ for various M . Here $\epsilon = 0.5$ and $\sigma = 10^{-2}$. As $\epsilon \rightarrow 0$ and $\sigma \rightarrow 0$, the $M = 1$ curve is found to converge to the value $(3\epsilon^2/16\pi^2)$.

(PPPL-802140)

Fig. 6. Corrections to Eq. (27) as a function of ϵ for various σ , obtained using the Fourier path summation method.

(PPPL-802141)
Fig. 7. The ratio of D/D_{QL} as a function of ϵ for $0 < \epsilon < 20$, as obtained using Fourier path summation. The peak at $\epsilon \approx 2\pi$ is associated with paths with large extent in (m,k) space. Also shown is the large ϵ expression given in Eq. (37). Here $\sigma = 10^{-3}$.