Received by OSTI

The Behavior of Matter under Non-equilibrium Conditions MAY 20 1992

Fundamental Aspects and Applications

Progress Report

July 15, 1991 - July 14, 1992

Ilya Prigogine

Center for Studies in Statistical Mechanics

The University of Texas at Austin

Austin, Texas 78712

April 1992

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY

UNDER GRANT NUMBER DE-FG05-88ER13897

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

PROGRESS REPORT
The Behavior of Matter under Non-equilibrium Conditions:
Fundamental Aspects and Applications

Our report contains a brief summary of what has been achieved over the period of the contract. We believe that we have formulated the laws of unstable "chaotic" systems in classical and quantum mechanics, using a probabilistic approach. Basically, the scheme that emerges is the following:

1. Dynamical instability leading to a probabilistic description
2. Establishment of spectral representation of the evolution operator for probabilities (This cannot be done by conventional methods.)
3. Proof that these spectral representations involve time symmetry breaking and therefore, solve the time paradox

The results obtained in this portion of the project are due to I. Prigogine, T. Petrosky, I. Antoniou, D. Driebe, H. Hasegawa, J. Lee, J. Levitan, R. Passante W. Saphir and S. Tasaki.

Maps:

Various maps are the prototypes of systems which are governed by deterministic laws but exhibit chaotic behaviors. One of the simplest maps is the Bernoulli map

$$x_{n+1} = 2x_n \pmod{1} . \quad (1)$$

Corresponding to the phase space description of the usual Hamiltonian systems, the evolution can be described by a distribution function ρ defined on $[0,1]$. Their time evolution is governed by the Perron-Frobenius operator U defined by

$$\rho_{n+1}(x) = U\rho_n(x) \equiv \int dy \delta(x - f(y)) \rho_n(y) = \frac{1}{2} \left\{ \rho_n\left(\frac{x}{2}\right) + \rho_n\left(\frac{x+1}{2}\right) \right\} . \quad (2)$$

It is well-known that the Bernoulli map is mixing and that, for any integrable initial distribution ρ_0 , $\rho_n = U^n \rho_0$ tends to a uniform distribution when n goes to infinity. In the usual Hilbert space formalism, U does not admit a spectral decomposition. Indeed, while the adjoint $U^\dagger$ is an isometry (which implies that the modulus of the eigenvalues should be one), U has eigenvalues $1/2^n$ as can be easily verified associated with polynomial eigenfunctions. On the contrary, as shown in Refs. 1 and 2, in our complex spectral theory, we can obtain a spectral decomposition of U

$$U\rho(x) = \varphi_0(x) \langle \varphi_0 | \rho \rangle + \sum_{n=1}^{\infty} \varphi_n(x) \frac{1}{2^n} \langle \tilde{\varphi}_n | \rho \rangle , \quad (3)$$

where Lyapounov times $1/2^n$ appear in the spectrum. Here $\varphi_n(x)$ is the n th Bernoulli polynomial defined through the generating function

$$\frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} \varphi_n(x) t^n , \quad (4)$$

and $\tilde{\varphi}_n$ is the distribution defined through the $n-1$ th derivative of $\rho(x)$ by

$$\langle \tilde{\varphi}_n | \rho \rangle \equiv \rho^{(n-1)}(1) - \rho^{(n-1)}(0) . \quad (5)$$

These formal results can be justified in an extended framework called rigged Hilbert spaces³. To define the generalized states $\tilde{\varphi}_n$ and ensure the convergence of the sum (3), the class of admissible densities ("test functions") should be restricted. Here we use the space $\mathcal{D}$ of all polynomials. This space together with its topological dual and the space L^2 of square integrable functions form a rigged Hilbert space $\mathcal{D} \subset L^2 \subset \mathcal{D}^\dagger$.

Then the spectral decomposition (3) is valid for any $\rho \in \mathcal{D}$ and the generalized eigenfunctions $\tilde{\varphi}_n$ live in the topological dual $\mathcal{D}^\dagger$.

One of the interesting features^{1,2,3} of the complex spectral decomposition is that the eigenvalues coincide with Ruelle's resonances [D. Ruelle, Phys. Rev. Lett. **56**, (1986), 405; Commun. Math. Phys. **125**, (1989), 239] determined by the zeros of the Fredholm determinant $d(z) \equiv \det(1 - z^{-1}U)$, which is usually calculated through the symbolic dynamics. This is also true for other maps like the baker transformation^{2,6}.

Besides the spectral decomposition containing polynomial eigenfunctions, the Perron-Frobenius operator U admits infinitely many spectral decompositions, each of which corresponds to an arbitrary closed path in the unit disk in the complex plane. These eigenstates are the generalization of coherent states of quantum harmonic oscillators and their graphs are fractals. The existence of these coherent states is the result of non-exponential decay of functions such as $\sin 2\pi x$ ($U \sin 2\pi x = 0$). This result will appear soon⁴.

The same method has been applied to the multi-Bernoulli map, an array of coupled Bernoulli maps. This map exhibits a diffusion process and much closer to realistic situations. For this system, we found eigenvalues corresponding to the diffusion process, piecewise polynomial eigenfunctions and generalized left eigenfunctions⁵.

The baker transformation, a typical example of unstable dynamical systems called Kolmogorov-systems, is closer to conservative Hamiltonian systems as it is invertible. The Perron-Frobenius operator U of the Baker transformation is unitary and has an absolutely continuous spectrum of countable uniform multiplicity. Apart from the usual spectral decomposition of the unitary operator U in Hilberts space, through the complex spectral theory^{2,6}, we can introduce a second complex spectral decomposition. The new spectrum representation corresponds to Lyapounov times $1/2^n$ and the evolution operator reduces to Jordan blocks. As a result of the appearance of Jordan blocks, the time evolution of densities deviates from exponential decay.

Our complex spectral theory has also applied to the standard map⁷, a typical non-linear map which shows various aspects of chaos. For this map, the appearance of diffusion process and the corrections to diffusion coefficient have been discussed by a perturbation method where the fully chaotic regime is chosen as an unperturbed state. Through this method, we have recovered the oscillatory corrections to the diffusion coefficient.

Other maps such as the cat map⁸ can be studied by similar methods.

Mathematical Aspects of the Complex Spectral Representations:

The integration of Chaotic Dynamical Systems is an unsolvable problem. Due to the instability or resonances, numerical or analytic techniques fail to provide the solution. As shown recently, these difficulties have been related to Gödel's undecidability⁹. Koopman's suggestion to study densities instead of trajectories does not resolve the problem either. In fact, as the generator of the densities evolution has continuous spectrum, it is an unbounded operator in the Hilbert Space of square integrable densities. It has also been shown recently that the eigenvalue problem for unbounded operators is not computable in Hilbert or Banach spaces⁹. This is typical not only for Classical, but also for Quantum Large Poincaré Systems¹⁰.

On the contrary, the theory of subdynamics provides a constructive method^{11,12} to obtain complex spectral representations of dynamical systems with continuous spectra like Large Poincaré Systems, Kolmogorov systems, or chaotic maps, invertible or non-invertible. The complex eigenvalues correspond to the relevant kinetic properties like Lyapunov exponents, diffusion coefficients or lifetimes. The constructions have, however, no meaning in the conventional Hilbert space framework, but acquire a natural meaning in "Rigged" Hilbert Spaces, which consist of linear functionals over suitable test subspaces.

In the case of Large Poincaré Systems like the Friedrichs Model, the eigenvectors associated with the resonances^{13,14} live in Rigged Hilbert Spaces of Hardy functions^{10,15}. Furthermore, the unitary group solution of the Schrödinger equation, when extended to the Rigged Hilbert Space, splits into two semigroups. This split shows explicitly the intrinsically irreversible character of the complex spectral representation^{10,15}. The

construction resolves therefore, not only the problem of irreversibility, but also the problem of integration of Large Poincaré Systems.

In the case of chaotic maps^{1,2,3,5,6} like the Renyi maps on the unit interval, the spectral decomposition of the Koopman (or Perron-Frobenius) operator acquires meaning in the Rigged Hilbert Space of linear functionals over the polynomial test functions^{3,16}. Trajectories lose their meaning in this representation, as the Dirac δ functions are not in the domain of the spectral decomposition. In this sense test functions play the role of the traditional coarse graining. This implies the impossibility of deterministic predictions for chaotic systems. Only probabilistic predictions are possible.

Large Poincaré Systems:

Among various types of unstable dynamical systems, of special interest for us are Hamiltonian systems with a continuous spectrum classified as Poincaré "non-intergrable system". We have called these systems Large Poincaré systems (LPS). Most systems in nature are indeed LPS. Our *complex spectral analysis* provides us with a systematic perturbational method for the construction of eigenfunctions and complex eigenvalues of the Liouville-von Neumann operator (which describes the time evolution of the statistical distribution function ρ)^{11,13,17,28} in both classical and quantum systems. This permits us to study in detail the approach to equilibrium.

These results are of special importance for quantum mechanics^{11,18,19} as our theory leads to an irreducible representation for broken time symmetry with the density matrix. That means that the time evolution of the density matrix for unstable systems *cannot* be expressed in terms of wave functions. Even if we start at the initial time with a wave function, we have a "collapse" of the wave function time going on. The reason for this collapse is the manifestation of "quantum chaos"¹⁸. As well known, classical chaos originates from the sensitivity to initial conditions. Quantum chaos has a quite different origin. It comes from the fact that wave functions become ill-defined as time going on due to the Poincaré resonances and acquire meaning only with suitable test functions. As a result, we obtain a density matrix which is not any more be "reducible" with respect to wave functions.

We have discussed simple examples corresponding to scattering¹⁸. The standard example of the S -matrix approach used in quantum theory is to two-body scattering. Both "in-states" and "out-states" are described by free-moving wave packets. In this case, there exists no irreducible representation.

However, the situation is different when we consider *persistent* interactions. For example, starting from a plane wave, the asymptotic limit of the wave function becomes ill-defined. The usual relation between density matrices and wave functions is lost¹⁸. An interesting example is three-body scattering. The conventional theory leads then to divergences (the so-called "rescattering anomaly") but our theory predicts a finite value for the three-body scattering cross section. We hope to develop these questions further, in more detail and to verify our predictions by computer experiments.

Of course, the main interest lies in the study of interactions of fields. This is the basic problem in modern field theory. There is no difficulty to find the spectral representation of free fields. However, a spectral representation of interacting fields has not been constructed. Our method suggests that the difficulty originates from the neglect of dissipative properties leading to Poincaré's divergences. We need a complex spectral theory whatever the value of the coupling constant¹⁹.

Moreover, the complex spectral theory avoids the well-known epistemological problems of the quantum measurement theory^{9,20}. As well known, Bohr suggested that the measurement apparatus had to be described in classical terms, and that the measurement should be associated to an irreversible macroscopic process. Our approach gives a precise meaning to Bohr's suggestions. In our view, the apparatus must be an unstable dynamical system leading to an irreducible representation in the Liouville space. As a result, we deal directly with probabilities and not with wave functions. While wave functions have no classical analogue quantum probabilities have. Our solution of the measurement paradox satisfies as well the second Bohr requirement: measurement must lead to irreversible effects. Of course these two requirements are not independent as dissipation is the result of resonances which lead to instability.

Selected Applications:

Let us indicate a few selected applications of the techniques we have discussed in the preceding sections:

For decay process of unstable states, "non-Markovian" processes (manifested as deviations from the exponential decay) have attracted much interest in the context of quantum measurement theory. Unfortunately, the time scales associated with the non-Markovian effects do not lend themselves readily to measurement, because of its too-short time scale ($\sim 10^{-15}$ sec). Applying the complex spectral analysis, we suggest a mechanism which renders an experimental investigation possible. We propose the injection into the interacting system of an additional time-dependent signal. It has been shown that the line shape of emitted photons in far off resonance condition could then give access to the non-Markovian of the system²¹.

Another application refers to dissipative processes in quantum optics. Usually theoretical works are restricted to the "Born-Markov" approximations. Now we can go systematically to higher order calculations²². An interesting effect which has emerged from our calculations is the following: In the case of a multi-level system, there exists the possibility of "indirect spectroscopy". In addition to the usual Weisskopf-Wigner "direct" decay from one state to a lower one, first a virtual transition brings the system to a higher excited level, from which it subsequently decays by resonance transition. This new phenomenon leads to a correction to Einstein's A and B coefficients. Experimental verification of this effect is under consideration.

As the result of the work by Misra and Sudarshan [J. Math. Phys. **18** (1977) 756], the short-time Zeno effect has been attributed to the quantum measurement process. However, in a recent paper we have shown that a Zeno period (short time deviation from exponential decay) exists also in classical systems²⁵. This question has recently attracted much interest as the result of a paper by Itano and others [Phys. Rev. **A41** (1990) 2259] reporting that they have succeeded in observing the quantum Zeno effect. However, we have shown that these results can be recovered through conventional quantum mechanics and do not involve a repeated reduction (or collapse) of the wave function^{23,24}. Our result has since been confirmed independently by other investigators [A. Peres and A. Ron, Phys. Rev. **A42** (1990) 5720]. Recently, Itano and others have admitted ambiguity of their interpretation [Phys. Rev. **A 43** (1991) 5168].

We have studied the approach to equilibrium of a classical gas with initial long-range spatial correlations²⁶. We have obtained "anti-Boltzmann" behavior in the evolution of the velocity distribution as the Boltzmann entropy decreases from the non-equilibrium to the equilibrium state. We also have non-trivial behavior on a short time scale due to the non-Markovian effects. The results obtained show the interplay between irreversible processes leading to disorder and to order in classical gas.

For driven systems, there may exist resonances between internal and external frequencies in addition to those among internal frequencies. For these systems, we have constructed projection operators which decompose the Liouville equation²⁷. This new approach recovers the previous formulations of time-dependent subdynamics but is based on a simple recursion formula, which significantly simplifies the explicit construction. These projectors are non-Hermitian and provide a description with broken time symmetry.

List of Publications (References)

- 1) H.H. Hasegawa and W.C. Saphir : "Decaying Eigenstates for Simple Chaotic Systems," *Physics Letters A* **161** (1992) 471.
- 2) H.H. Hasegawa and W.C. Saphir : "Unitarity and Irreversibility in Chaotic Systems," to be presented *Physical Review A* (1992).
- 3) S. Tasaki and I. Antoniou : "Generalized Spectral Decomposition for r -adic Renyi Map," "The 3rd International Workshop-Conference on Evolution Equations, Control Theory and Biomathematics," Hans-sur Lesse, Belgium, October, (1991).
- 4) I. Antoniou and S. Tasaki, "Coherent States of Shifts" and "Coherent States Decomposition of Chaotic Dynamical Systems," to appear.
- 5) H.H. Hasegawa and D.J. Driebe : "Transport as a Dynamical Property of a Simple Map," submitted to *Physics Letters A*.
- 6) H.H. Hasegawa and W.C. Saphir : "Non-equilibrium Statistical Mechanics of the Baker Map: Ruelle Resonances and Subdynamics," *Physics Letters A* **161** (1992) 477.
- 7) H. Hasegawa and W. Saphir : "Kinetic Theory for the Standard Map," in *Solitons and Chaos* pp.192 - 200, eds. I. Antoniou and F. Lambert, Springer, Berlin (1991).
- 8) H.H. Hasegawa and W.C. Saphir, to appear.
- 9) I. Prigogine : "Dissipative Processes in Quantum Theory," the XXth Solvay Conference, November 5 - 9 (1991), eds. P. Mandel., see also Antoniou's remarks.
- 10) I. Antoniou and I. Prigogine : "Dynamics and Intrinsic Irreversibility," *Nuovo Cimento*, (1992) in press.
- 11) T. Petrosky and I. Prigogine : "Alternative Formulation of Classical and Quantum Dynamics for Non-integrable Systems," *Physica A* **175** (1991), 146.
- 12) I. Antoniou and S. Tasaki : "Generalized Spectral Decompositions of Mixing Dynamical Systems," to appear in *Journal of Quantum Chemistry Symposia*, (1992).
- 13) H. Hasegawa, T. Petrosky, I. Prigogine and S. Tasaki : "Quantum Mechanics and the Direction of Time," *Foundation of Physics*, **21** (1991), 263.
- 14) T. Petrosky, I. Prigogine and S. Tasaki : "Quantum Theory of Non-integrable Systems," *Physica A* **173** (1991), 175.
- 15) I. Antoniou : "Intrinsic Irreversibility of Quantum Systems, Rigged Hilbert Space for the Lee-Friedrichs Model," the 2nd Int. Wigner Symposium, July 15 - 20 (1991), eds. H. Doebner, F. Schroeck.
- 16) I. Antoniou and S. Tasaki : "Generalized Spectral Decompositions of the r -adic Renyi Transformations," to appear (1992).
- 17) I. Prigogine, T. Petrosky, H. Hasegawa and S. Tasaki : "Integrability and Chaos in Classical and Quantum Mechanics," *Solitons and Chaos* pp.3 - 24, eds. I. Antoniou and F. Lambert, Springer, Berlin (1991).
- 18) I. Prigogine and T. Petrosky : "Quantum Chaos - Toward the Formulation of an Alternative Quantum Theory," the 2nd Int. Wigner Symposium, July 15 - 20 (1991), eds. H. Doebner, F. Schroeck.
- 19) S. Tasaki, P. Nardone and I. Prigogine : "Negative Energy Instability in a Toy Model of Quantum Gravity," to appear (1992).
- 20) I. Prigogine : "From Poincaré's Divergences to Quantum Mechanics with Broken Time Symmetry," in honor of Prof. Sudarshan, to appear (1992).
- 21) T. Petrosky, Lee Jing-Yee and J. Levitan: "Non-Markovian Effects in an Exactly Soluble Matter-Field Interaction System," in press, *Physics Letters* (1992).
- 22) R. Passante, T. Petrosky and I. Prigogine : "Virtual Transitions, Self-Dressing and Indirect Spectroscopy," submitted to *Europhysics Letters* (1992).
- 23) T. Petrosky, S. Tasaki and I. Prigogine : "Quantum Zeno Effect," *Physics Letters A*, **151** (1990), 109.
- 24) T. Petrosky, S. Tasaki and I. Prigogine : "Quantum Zeno Effect," *Physica A* **170** (1991), 306.
- 25) T. Petrosky and I. Prigogine, "Laws and Events: The dynamical basis of self-organization," *Can. J. Phys.* **68** (1990), 670.
- 26) D. Driebe and T. Petrosky : "Order and Disorder in the Approach to Equilibrium of a Classical Gas," in press, *J. Stat. Phys.*, (1992).

List of Publications (References) Continued

- 27) Lee Jing-Yee and S. Tasaki : "Poincaré's Theorem and Subdynamics for Driven Systems," *Physica A*, **182** (1992), 59.
- 28) T. Petrosky and I. Prigogine: "Integration of Poincaré's Non-integrable Systems --New Formulation of Classical and Quantum Dynamics--," *Proceedings of Ninth Symposium on Energy Engineering Sciences (Research on Fluid and Dynamical Systems) at the Argonne National Laboratory*, edited by D. Frederick, (1991).

END

**DATE
FILMED**

7 / 2 / 92