MEASUREMENTS OF IONIZATION BALANCE PARAMETERS
IN ATOMIC IONS

Grant DE-FG02-86ER13634

Final Report

For the period 1 May 1982 through 31 May 1989

Principal Investigator

Dr. John L. Kohl

June 1991

Prepared for
U. S. Department of Energy
Washington, D. C. 20545

Received by OST
MAR 30 1992

Smithsonian Institution
Astrophysical Observatory
Cambridge, MA 02138

The Smithsonian Astrophysical Observatory
is a member of the
Harvard-Smithsonian Center for Astrophysics

NOTICE

This work was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed or represents that its use would not infringe privately-owned rights.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MASTER

~~CONFIDENTIAL~~ 402

2.0 Publications and Presentations

Scientific publications and presentations resulting from this research program are as follows:

Papers:

- An Experimental Study of the Electron Impact Excitation of Positive Ions, G.P. Lafyatis, Ph.D. Thesis, Harvard University (1982).
- Photon-Ion Concidence Measurements of Dielectronic Recombination in C^{3+} , J.L. Kohl, L.D. Gardner, G.P. Lafyatis, A.R. Young, and A. Chutjian, in Atomic Excitations and Recombination in External Fields, M.H. Nayfeh and C. Clark, eds., (New York: Gordon & Breach Science Publishers, 1985) pp. 421-437.
- An Experimental Apparatus for Photon/Ion Coincidence Measurements of Dielectronic Recombination, L.D. Gardner, J.L. Kohl, G.P. Lafyatis, A.R. Young and A. Chutjian, *Rev. Sci. Instrum.* 57, 2254 (1986).
- Experiments on Electron Impact Excitation and Dielectronic Recombination at Harvard-Smithsonian, L.D. Gardner, J.L. Kohl, D.W. Savin and A.R. Young, *Journal de Physique* 50, C1-405 (1989).
- An Experiment to Measure Dielectronic Recombination in a Known External Field, A.R. Young, Ph.D. Thesis, Harvard University (1990).
- A Measurement of Dielectronic Recombination in a Known External Field, A.R. Young, L.D. Gardner, J.L. Kohl, D.W. Savin, G.P. Lafyatis, and A. Chutjian, submitted to *Physical Review A* (1991).

Abstracts:

- A Photon-Ion Coincidence Method for the Measurement of Dielectronic Recombination, G.P. Lafyatis, presented at the NATO Advanced Summer Institute on Atomic and Molecular Processes in Controlled Thermonuclear Fusion, Palermo, Italy (1982).
- Progress toward Measurement of Dielectronic Recombination in C^{3+} , J.L. Kohl and G.P. Lafyatis, Atomic Physics Program Contractors' Workshop, April, 1984.
- Progress Towards an Inclined Beams Measurement of Dielectronic Recombination in C^{3+} , G.P. Lafyatis and J.L. Kohl, *Bull. Am. Phys. Soc.* 29, 805 (1984).
- Measurements of Absolute Collisional Cross Sections at Harvard-Smithsonian, CfA, J.L. Kohl, L.K. Deutsch, L.D. Gardner, G.P. Lafyatis, and A. Young, contributed paper to the IAU Colloquium No. 86, the Eighth International Colloquium on EUV and X-ray Spectroscopy of Astrophysical and Laboratory Plasmas, Washington, D.C., Aug. 27-29, 1984.

- Photon-Ion Coincidence Measurements of Dielectronic Recombination, J.L. Kohl, L.D. Gardner and A.R. Young, DOE Atomic Physics Program Contractor's Workshop, Chapel Hill, NC, May 13-14, 1985.
- Coincidence Method Results for C^{3+} Dielectronic Recombination, L.D. Gardner, J.L. Kohl, G.P. Lafyatis and A.R. Young, Bull. Am. Phys. Soc. 31, 989 (1986).
- Measurements of Dielectronic Recombination in C^{3+} , L.D. Gardner, J.L. Kohl, D.W. Savin and A.R. Young, Bull. Am. Phys. Soc. 32, 1274 (1987).
- Measurements of Dielectronic Recombination in a Known Electric Field, J.L. Kohl, L.D. Gardner, A.R. Young and D.W. Savin, VIIIth Atomic Physics Program Workshop (DOE Atomic Physics Program Contractor's Workshop), Cambridge, MA, May 21-22, 1987.
- Cross Beams Determination of the Dielectronic Recombination Cross Section in C^{3+} , L.D. Gardner, J.L. Kohl, D.W. Savin and A.R. Young, Abstracts of Contributed Papers to the XVth International Conference on the Physics of Electronic and Atomic Collisions, Brighton, U.K., 1987, p. 384.
- Experimental Cross Section for Dielectronic Recombination in C^{3+} , A.R. Young, L.D. Gardner, J.L. Kohl, and D.W. Savin, Post-Deadline Paper to the XVIth International Conference on the Physics of Electronic and Atomic Collisions, New York, NY, 1989.
- Measurements of Dielectronic Recombination in C IV in an External Electric Field, A.R. Young, L.D. Gardner, J.L. Kohl, J.C. Raymond, D.W. Savin, and D.B. Reisenfeld, Bull. Am. Astron. Soc. 22, 799 (1990).

3.0 Experimental Results

A diagram of the experimental arrangement is shown in Figure 1. A 38 keV beam of C^{3+} is purified with a charge state analyzer and crossed with an electron beam at 45 degrees. A controllable magnetic field along the electron beam provides a known electric field in the ion rest frame. Photons are collected by a mirror and detected with a photomultiplier (PMT). Recombined ions are selected by a charge state analyzer and detected with a channel electron multiplier (CEM). The DR event rate is determined by counting delayed coincidences between stabilizing photons and recombined ions. The electron energy is switched off and on resonance at a rate of 0.1 Hz, and plots of coincidences versus delay time are recorded for each energy. In order to determine the DR rate coefficient, one must measure or otherwise determine the DR event rate, the current density in each beam in the region of intersection, the electron-ion relative velocity, the detection efficiency of the CEM, the efficiency of the photon collection system as a function of position in the emitting region (essentially the same as the beams' intersection region), and the detection efficiency of the PMT as a function of position on its photocathode. The reader is referred to the attached preprint and to Gardner, et al. (1986) for detailed discussion of the experimental techniques and analysis procedures.

Figure 1: Schematic of the experimental apparatus.

Measurements of the DR rate coefficient for the C^{3+} ion were made for the case where the process occurs in an external electric field of 12 volts/cm and a 155 nm photon is emitted. The measured value of the rate coefficient $3.6 \pm 1.5 \times 10^{-10} \text{ cm}^3/\text{sec}$ and the energy averaged cross section is $2.1 \pm 0.9 \times 10^{-18} \text{ cm}^2$. This measurement is dominated by statistical uncertainties (± 30 percent, quoted here at the 68 percent confidence level). It is the first measurement of DR for a multiply charged ion in a directly measured external electric field. All uncertainties and systematics affecting the result have been very carefully considered. The experiment has been verified through measurements of other atomic processes that are better understood.

Measurements of the electron impact excitation of C^{3+} have been made near threshold in order to check our data analysis procedures, to calibrate the ion rest frame collision energy, and to determine the energy distribution of the electrons in the beam. This measurement and theoretical calculations of the rate coefficient (Gau and Henry,

1975) were used to set the absolute scale in the experiment. The data are presented in Figure 2. The error bars shown are statistical at 90% confidence. The electron energy distribution is found to be consistent with a gaussian with full width at half maximum of 1.8 ± 0.2 volts.

Figure 2: The electron impact excitation rate coefficient near threshold for the $1s^2 2s \ ^2S$ to $1s^2 2p \ ^2P^o$ transition in C^{3+} . The solid curve is the best fit of the experimental data to theory (Gau and Henry, 1977). The error bars on the data are the statistical uncertainties at 90 percent confidence.

Since DR experiments require an unambiguous determination of the conditions under which DR is measured, several auxiliary measurements were made. The energy distribution derived from the excitation data, the electric fields in the interaction region (shown in Figure 3), and the charge state analyzer fields that determine the cutoff in the principal quantum number, n , of the captured electron due to field-ionization of the recombined ions were measured.

Figure 3: The electric fields in the ion rest frame.

Figure 4: Comparison of the measured DR rate coefficient to theory. The error bars shown are total uncertainties at the 68 percent confidence level. The lower solid curve uses the zero field calculations of McLaughlin and Hahn (1983). The upper solid curve uses calculations of LaGattuta (1985)/McLaughlin and Hahn (1983) for 12 V/cm (see the text). The lower dotted curve is a zero field calculation and the upper dotted curve a 12 V/cm calculation, both by Griffin, et al. (1986,1989).

Figure 4 shows the predicted DR rate coefficient for two values of the external electric field strength, where the field dependence is taken from the work of LaGattuta (1985), and the zero field cross sections are taken from McLaughlin and Hahn (1983). Theoretical curves generated by Griffin et al. (1986, 1989) are also shown. There are several points to note concerning the comparison of the theoretical curves to the measurement.

The first is that this measurement, including the 90 percent confidence uncertainties, differs from the theoretical predictions for a zero external field. The field effect is required in the theory to produce agreement with the present result.

The second point to note here is that when the effects of the external field are included in the theory the measured value of the rate coefficient is still a factor of 3 larger than that predicted by LaGattuta/McLaughlin and Hahn, and 70 percent larger than that predicted by Griffin et al., although the experimental uncertainties may account for the differences. This is consistent with the work of Dittner et al. (1988) which indicated a rate coefficient that is 50 percent larger than rate coefficients computed with a saturated field effect (where no further field enhancement is expected), and the recent work of Andersen and Bolko (1990) (for fields estimated to be 2 to 3 V/cm) which indicates a possible factor of two discrepancy between experiment and theory. It should be noted that the fields were not precisely known in either of these latter experiments.

The rate coefficients for a given value of the electric field strength as calculated by Griffin et al. are generally larger than those of LaGattuta. One of the principal differences between the two calculations is that the work of Griffin includes fine structure interactions: i.e., an intermediate coupling calculation. The predicted size of the field enhancement tends to be smaller when intermediate coupling calculations are performed.

A further point is that neither of these calculations take into account rotational effects due to changes in direction of the external electric field on the measured recombination rate coefficient or the possibility of overlapping-interacting resonances. The rotation effects arise because the rest frame electric fields in the collision volume and in the post-analyzer are at 90° to one another in the present experiment. The recombined ions therefore experience an electric field that is changing in orientation and magnitude, a situation which is known to affect the relative populations of the Stark states of the captured electron. This, in turn, can affect the probability that the recombined ion is field-ionized in the post-analyzer. This has been studied in some detail for the case of Mg^+ and resulted in a maximum reduction of 15 percent in the size of the predicted rate coefficient (LaGattuta, 1986). A similar effect may be present in C^{3+} . Analytical

treatments using quantum defect theory indicate there may be a reduction in the size of the field enhancement at large values of the captured electron's principal quantum number due to overlapping-interacting resonances (Harmin, 1986). This, too, would result in a small decrease in the size of the predicted rate coefficient. Both effects tend to push the experimentally determined rates out of agreement with theory, but the effects are expected to be small enough that there is no dramatic change in the conclusions drawn from the experimental data.

Further measurements are planned at the same and other external electric field values. In order to facilitate the additional measurements, improvements have been made to the apparatus. Additional vacuum pumping has been added to the interaction chamber with the goal of reducing the recombined ion background from charge transfer. We now achieve pressures below 1.0×10^{-10} torr with the experiment operating. The ion background was further reduced by modifying the final charge state analyzer and primary beam dump.

Computer simulations of the "leakage" fields from the electrostatic charge state analysers have been carried out. Results indicate that stray fields from this source are negligible (< 0.1 V/cm) in our experiment. The total experimental uncertainty in the external field strength is 1.6 V/cm, a level that is adequate for comparisons of the experimental DR results to theory.

4.0 Future Directions

At present, the program is continuing with two Harvard graduate students pursuing their Ph.D. research. One of them has made considerable progress on a measurement to determine the variation in the C^{3+} DR rate with the strength of the external electric fields. Measurements of the C^{3+} DR rates at three values of the external electric field are in progress. The other student has initiated a measurement of electron impact excitation in C^+ . The work is expected to lead to DR measurements in that atomic system.

5.0 References

- Anderson, L.H., Bolko, J., and Kvistgaard, P., 1990, Phys. Rev. A **41**, 1293.
Burgess, A., and Summers, H.P., 1969, Ap. J. **157**, 1007.
Dittner, P.F., 1988, Phys. Scr. **T22**, 65.
Griffin, D.C., Pindzola, M.S., and Bottcher, C., 1986, Phys. Rev. A **33**, p. 3124.
Griffin, D.C., Pindzola, M.S., and Krylstedt, P., 1989, in press.
Gau, J.N. and Henry, R.J.W., 1977, Phys. Rev. A **16**, 986.
Harmin, D., 1987, Phys. Rev. Lett. **57**, 1570.
Jacobs, V.L., Davis, J., and Kepple, P.C., 1976, Phys. Rev. Lett. **37**, 1390.
LaGattuta, K.J., 1985, J. Phys. B **18**, L467.
LaGattuta, K.J., Nasser, I., and Hahn, Y., 1986, Phys. Rev. A **33**, p. 2782.
McLaughlin, D.J., and Hahn, Y., 1983, Phys. Rev. A **27**, 1389.

*Preprint removed
& cited separately*

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DATE
FILMED
5/18/92**