MASTER

There is no objection from the patent point of view to the publication or dissemination of the document(s) listed in this letter.

BROOKHAVEN PATENT GROUP

12/16 19 81 By CRK

NEW YORK UNIVERSITY
FACULTY OF ARTS AND SCIENCE
DEPARTMENT OF APPLIED SCIENCE

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

DOE/EV/10610--01

DE82 005484

GEOCHEMICAL AND GEOPHYSICAL MODELS OF
THE FOSSIL-FUEL CO₂ CLIMATE PROBLEM

Semiannual Progress Report
for Period February 2, 1981 - August 31, 1981

Martin Hoffert and Warren Wiscombe

Prepared for

THE U. S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-ACO2-81EV10610

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

DISCLAIMER
This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

EP

NEW YORK UNIVERSITY

FACULTY OF ARTS AND SCIENCE

DEPARTMENT OF APPLIED SCIENCE

26-36 Stuyvesant Street
New York, New York 10003

GEOCHEMICAL AND GEOPHYSICAL MODELS OF
THE FOSSIL FUEL CO₂ CLIMATE PROBLEM

Semi - Annual Progress Report

2/2/81 - 8/31/81

Our initial efforts have been directed towards development of the coupled ocean-atmosphere climate model described in our original proposal. In the course of setting up a prototype of this model, we made an important discovery with far-reaching implications for the CO₂ problem.

Basically, our discovery was this: the evaporative constraint on tropical ocean temperature described by Newell and Dopplnick (1979), when put into a climate model, yields much more realistic simulations than the current generation of energy-balance climate models have been able to achieve. That this constraint is correct is strongly supported by paleoclimatic data showing near-constancy (to $\pm 1-2K$) of the tropical sea-surface temperature over the past 700 million years.

As is well known, Newell and Dopplnick's paper roused a storm of protest (Ramanathan, 1981 etc.) and indeed could legitimately be faulted for ignoring a number of feedback processes which other models include. But lost in the hubbub was the germ of a very good idea. By incorporating this idea properly into a new kind of climate model, which Newell and Dopplnick never did, we have, we believe, made an important stride in our ability to predict future climates and explain past ones.

Let us briefly review the deficiencies of Budyko-Sellers-North (BSN) climate models, which all current energy-balance climate models are either copies of or slight variations upon. These models, by tuning several empirical formulas, have been successful in predicting the present surface air temperature distribution as a function of latitude. They also "predict" the latitudinal location of the ice-line, however this is illusory since when they recapture the present latitudinal temperature distribution, which they are tuned to do, they automatically get the present average ice-line location.

Where these models flounder is in predicting climatic change, which of course is a stiff test. For climates near that of the present, they predict much less sensitivity than we actually observe. They simply cannot explain the tremendous variability we see in our current climate, which must be due to relatively small forcings. They are unable to recover the last ice age or even come close to it, in spite of numerous forcing mechanisms which have been tried (including the famous Milankovitch mechanism, related to changes in the earth's orbit). This again is because of too much insensitivity to forcing functions. They are also unable to reproduce the climate of the warm Cretaceous period (140-65 million years ago) when the poles were ice-free.

Yet for the larger excursions of ice toward the equator, BSN models are far too sensitive. They would have the ice advancing uncontrollably once it reached about 40° N, a latitude which was actually reached (repeatedly) by the Laurentide sheet in North America during the last ice age. Much study has been devoted to this instability even though it appears patently ridiculous in light of past climates.

The above deficiencies are all remedied in our new type of climate model, which we have dubbed a "tropical evaporative constraint" (TEC) model.

This model manifests a quite different and, we believe, more realistic response to CO_2 forcing than BSN models. In particular:

- high-latitude amplification (the prediction that high-latitude temperatures change much more than the global average) is much greater than in BSN models; hence the icecaps will melt even earlier due to CO_2 forcing
- present-day climate is much more sensitive than in BSN models
- there are no unstable climates, no "iceball earth" solutions; it requires greater and greater forcing to move the ice-line closer and closer to the equator
- the ice-line in the last ice age can be reproduced using the lower measured concentrations of CO_2 at that time (200 ppm, Oeschger, 1981); the BSN models are unable to do this
- the warm Cretaceous climate can be successfully reproduced
- the ice-age phasing problem found in BSN-style approaches to Milankovitch forcing (Suarez and Held, 1979) goes away; that is, rather than being in the midst of an ice age at present, as the Suarez-Held model would have it, we are properly predicted to be in an interglacial.

We are in the process of preparing a formal publication documenting these results. To facilitate review of this work by the DOE CO₂ Research Division, Office of Energy Research, a summary of our model and major results in graphical form are given below.

TECHNICAL DETAILS

The equator-to-pole distribution of longitudinally-averaged surface air temperature $t_a(x)$ where $x \equiv \sin\phi$ and ϕ is latitude, is known to depend critically on the poleward heat flux F . Without this flux (carried about equally by the atmosphere and oceans) the equator would be much warmer and the poles much colder than they are. In terms of an effective thermal diffusion coefficient D with the units W/m^2-K , energy balance climate models generally express F by a kind of Fourier heat conduction law,

$$F = -2\pi R^2 D (1 - x^2) dT_a / dx \quad (1)$$

where $R \approx 6.37 \times 10^6$ m is the Earth's radius and the $(1 - x^2)$ term comes in due to spherical geometry. At any latitude the energy absorbed by the Earth per unit surface area is Sa , where $S(x)$ is the incident solar flux and $a(x)$ is the local planetary absorptance (equal to one minus the local albedo). The longwave flux to space I is a function of (among other things) the surface temperature T_a and the CO₂ content of the atmosphere. It is through "I" that direct CO₂ forcing enters the picture. Both a and I depend on cloudiness in ways which may (Cess, 1976) or may not (Ohring and Clapp, 1980; Hartmann and Short, 1980) cancel out. In any event, the cloudiness feedback is suppressed initially in order to focus on evaporation damping of tropical temperature per se. Temperature-ice albedo feedback is retained, however, by

switching the absorptance "a" from a higher to a lower value (signifying increased surface albedo) whenever T_a drops below some critical temperature T^* associated with the deposition of snow or the freezing of ice on the underlying land or water surface. This point is called the iceline latitude ϕ^* or $x^* = \sin\phi^*$.

Global conservation of energy requires

$$\frac{dF}{2\pi R^2 dx} = S_a - I, \quad (2)$$

and we assume the following functional forms for annual mean quantities on an equatorially-symmetric planet:

$$\begin{aligned} T_a(x) &= T_0 - T_1 x^2, \\ S(x) &= Q(s_0 - s_1 x^2), \\ a(x) &= \begin{cases} a_0 - a_1 x^2; & \text{when } T_a > T^* \\ b_0 & ; \text{when } T_a < T^* \end{cases} \\ I(x) &= A + B T_a = A + B T_0 - B T_1 x^2 \end{aligned} \quad (3)$$

where T_0 is the temperature at the equator, T_1 the equator-to-pole temperature difference, Q is 1/4 of the solar constant, s_0 and s_1 are coefficients in the latitude distribution of insolation which depend on the earth's spin axis tilt or obliquity (see Hoffert et al., 1981, for equations), a_0 and a_1 are coefficients in the distribution of absorptance equatorward of the iceline, b_0 is the absorptance over snow and ice and A and B are coefficients in the linear I - T_a relation. These functions are shown in Figure 1 along with the satellite data to which they are fitted. Current values of these

parameters used in our model are $T_0 = 302$ K, $T_1 = 44$ K, $s_0 = 1.246$, $s_1 = 0.738$, $a_0 = 0.785$, $a_1 = 0.263$, $b_0 = 0.380$, $x^* = 0.941$ ($\phi^* \sim 70^\circ$), and $B = 2.2$ W/m²-K, corresponding to the present solar output $Q = Q_0 = 340$ W/m² and present CO₂ level c_0 embodied in $A = A_0 = -395$ W/m². The CO₂ content of the atmosphere is related to this coefficient approximately by $A = -395 - 7.93 \ln(c/c_0)$ W/m² (see Hoffert, et. al. 1980, for derivation). Thus Climatic changes can be generated by varying solar output (Q), obliquity (s_0 and s_1), absorptance (a_0 , a_1 , b_0) or CO₂ content (c). The numerical values were tuned to satisfy the integral constraint on (2) that meridional flux vanishes at the equator and poles:

$$\int_0^1 (Sa - I) dx = \int_0^{x^*} (Sa - I) dx + \int_{x^*}^1 (Sa - I) dx = 0 \quad (4)$$

Actually, equation (4) must always be satisfied when climate changes, but it is only necessary to carry the integral to the iceline x^* when the earth is partly glaciated. Under these conditions the iceline condition

$$T(x^*) \equiv T^* = 263 \text{ K} = \text{constant} \quad (5)$$

carried over from the models of Budyko (1969), Sellers (1969), North (1975), etc., is applied. One more condition is needed which traditionally relates to fixing the thermal conduction D during climate change. But " D " is essentially an empirical quantity that results from the complex interplay of transport processes in the poleward direction as well as exchange between surface and atmospheric heat reservoirs. Integrating equation (2) for $F(x)$ using (3) and the present-day values for T_0 , etc., leads to the dashed curve in Figure 2. Substituting this $F(x)$ back into (1) gives the dashed $D(x)$ curve

of Figure 2; as expected, it is a weak function of latitude. The present global mean value $\bar{D} \equiv \int_0^1 D(x)dx = 0.641 \text{ W/m}^2\text{-K}$. The standard BSN assumption is to keep $\bar{D}(T_0, T_1, x^*) = 0.641 \text{ W/m}^2\text{-K}$ constant during climate change, which together with (4) and (5) closes the system. This assumption controls the equator-to-pole temperature gradient in energy-balance climate models and is responsible, we find, for the unrealistic predictions of paleoclimatic equator-to-pole temperature distributions calculated by North *et al.* (1981), Barron *et al.* (1981), Suarez and Held (1979) and many others. This suggests the substitution of a different constraint better grounded in the paleo-temperature data of earlier periods.

The great stability of wet, tropical regions to changes in insolation was attributed by Priestly (1966) to the dominance of evaporation cooling in the surface energy balance. More recently Newell and Doplick (1979) argued that evaporation would dramatically damp the tropical ocean surface temperature response to the enhanced downward IR radiation produced by increasing CO_2 . They find 1/10 of the current consensus value, i.e., 0.25K instead of 2.5 K global warming for a CO_2 doubling.

To study this evaporative constraint in the context of our energy-balance climate model, we started from the surface energy balance of the ocean under steady-state conditions, which can be written

$$\hat{S}_w + \Delta S_w + \Delta I_{aw} = LH_{wa} + SH_{wa} + \hat{I}_{wa} + \frac{dF_w}{2\pi R^2 dx} \quad (6)$$

All terms are per unit surface area of ocean. Those on the left-hand-side are, respectively, the solar flux absorbed under current conditions, the increment in absorbed solar flux, and the increment in downward longwave

radiation from the CO₂ greenhouse effect. On the right are the latent heat flux from the water into the atmosphere, the sensible heat flux, the net longwave flux from water to air under current conditions and the poleward flux carried by the oceans. We now assume that, on the right-hand side of (6), all but the latent heat term can be neglected (this is the "tropical evaporative constraint", TEC), and that it is given by the usual Dalton formula (Geiger, 1960)

$$LH_{wa} = \gamma(e_{sat} - e) \quad (7)$$

where γ is related to average wind speed (assumed invariant), e_{sat} is saturation vapor pressure at the ocean temperature, and e is vapor pressure in the surface air. In the tropics, the ocean and surface air temperatures are always less than a degree apart, so, upon assuming RH_a = surface air relative humidity is invariant, we arrive at

$$LH_{wa} = \gamma e_{sat}(T_o) (1 - RH_a) = \gamma' \exp(-\theta/T_o) \quad (8)$$

γ' is invariant, $\theta_L = 5420K$, and the approximation to e_{sat} results from an approximate integration of the Clausius-Clapeyron equation (Hess, 1959).

Putting (8) back into (6) for present $(\hat{T}_o = 302K)$ and then for perturbed conditions yields a temperature change

$$\Delta T_o \equiv T_o / -\hat{T}_o \approx (\hat{T}_o^2 / \theta_L) \ln \left[\frac{\hat{S}_w + \Delta S_w + \Delta I_{aw}}{\hat{S}_w} \right] \approx 16.8 \ln(Q/Q_o) \quad (9)$$

where the approximation on the far right applies when $\Delta I_{aw} = 0$ and only the solar constant changes. From (9) the sensitivity parameter at the equator is $\beta_o \equiv (1/Q_o) dT_o/dQ \approx 17 K$, roughly an order-of-magnitude below the 110-170K values commonly cited for the global mean sensitivity parameter $\bar{\beta} \equiv (1/Q_o) d\bar{T}/dQ$

($\bar{T} = \frac{1}{\int_0} T(x)dx$). We believe the order-of-magnitude discrepancies between the Newell-Dopplnick results and the global modeler's is due to this evaporation damping effect, not to Newell & Dopplnick's neglect of water vapor and other feedbacks as claimed in a recent paper by Ramanathan (1981). Indeed, Newell and Dopplnick's (1981) most recent calculations with water vapor feedback confirm their earlier predictions.

Evidence for the near-constancy of T_0 is summarized in Figure 3, showing that it has remained stable over the past 700Myr of geologic time to within a few degrees K, while bottom water temperatures due to downwelling at polar latitudes fluctuated over a 15-20 K range. Bottom waters near freezing correspond to glacial ice at the pole while warm bottom waters correspond to ice-free poles.

Donn and Shaw (1977) used a constant-D BSN model to "successfully" reproduce the evolution of climate from the early Triassic (200Myr ago) to the present, using continental drift as the only forcing function. But the more recent study of the Cretaceous period (65-140 Myr ago) carried out by Barron et al. (1981) resulted in different conclusions. They found it was necessary to have some additional forcing either from the sun or a greenhouse gas and that the equator-to-pole temperature profile in the record could not be fit by their constant-D BSN model. Moving on to more recent geological times, we see in Figure 3 the formation of the Antarctic ice sheet, and later the Northern Hemisphere ice sheets led to the cold bottom waters of the present Glacial Age including the most recent (Wisconsin) Ice Age whose surface temperatures were reconstructed by CLIMAP (1976). Typical water temperatures versus latitude for this period (18 kyr ago) as well as those of the current and Cretaceous climates are shown in Figure 4. Using the

intersection of the zonal-annual-average sea surface temperature with the freezing point of water as a working definition of the global mean iceline, CLIMAP data predict $\phi^* = 56^\circ$ ($x^* = 0.83$). A number of authors (North et al., 1981; Hartmann and Short, 1979; etc.) have commented that constant-D BSN models are too insensitive to recover this CLIMAP iceline for plausible amounts of climatic forcing.

To overcome the aforementioned problems we replaced the constant-D constraint with the "TEC" constraint derived above.

Figure 5 illustrates our results for the location of the iceline x^* vs. solar output Q/Q_0 for both the constant-D and TEC models. For comparison, the results of Hartmann and Short (1979) who used a 3-mode Legendre polynomial expansion a la North (1975) and a similar $a(x)$ to ours is also shown. Clearly, our parabolic $T(x)$ and constant- $\bar{D}$ model is equivalent to the traditional BSN models and exhibits the usual multiple solutions at the present solar constant. As discussed by Held and Suarez (1974) and Lindzen and Farrell (1980) the slope $|dx^*/dQ|$ is a general measure of climate stability and sensitivity when the planet is glaciated, with $dx^*/dQ > 0$ implying stability, $dx^*/dQ < 0$, instability. In general, smaller positive values of dx^*/dQ imply lesser climate sensitivity.

We find the iceline location is almost four times as sensitive to solar forcing with the TEC model than with the constant-D model, as shown in Figure 5. This is because the weak (logarithmic) variation of tropical temperature with solar output forces temperature changes to be greater at high latitudes, where they are further amplified by ice-albedo feedback. For example, the condition for polar cap melting in the long-term equilibrium is a 0.9% increase in solar

output for the TEC model, 3.5% for the constant-D. These results should not be taken too literally at this point, since other feedbacks may come into play on the real earth to preserve polar ice. (especially the strong polar-night cooling which is not accounted for in our present annually-averaged model). But they dramatically illustrate that evaporation-stabilized tropical temperatures have precisely the opposite effect on global sensitivity than they do on tropical sensitivity.

Obviously, the TEC model will break down for sufficiently large changes in climatic forcing, here represented by solar output as a proxy for other types of forcing like CO_2 . To study the regions of validity and probable breakdown we computed the global mean temperature, equator-to-pole temperature difference and peak poleward flux for the TEC and constant-D models. They are shown in Figure 6. If Q decreases sufficiently far, it is likely that other terms in the surface energy balance will become comparable to the latent heat and thus (9) will no longer be valid; but it is interesting that if (9) is maintained while Q is decreased, the poleward flux keeps decreasing because the tropics can only maintain their high T_0 by suppressing poleward heat transport, until a zonal radiative equilibrium ($F=0$) condition is created at about 83% of the current solar output. It follows that even with the TEC model, which is more stable than the constant-D model for large decreases in Q , it is necessary to invoke some additional warming mechanism, like highly abundant CO_2 as a greenhouse gas, to account for habitable temperatures on earth during the first billions of years after solar system formation when the solar output was probably 30% less than at present. On the warm side, the TEC model can reasonably be considered to break down when it forces surface temperatures to be uniform over the earth, which occurs at about 114% of the present solar output. Nevertheless, for

the perturbations in climate forcing which have actually been experienced on earth, Figures 3 and 4 suggest the tropical evaporative constraint has actually been in operation.

Another check on the realism of the TEC model is given in Figure 7 where its equator-to-pole temperature distribution is compared to the constant-D result having the same global mean temperature $\bar{T} = 298.1 \text{ K}$ typical of the Cretaceous. The TEC model agrees much better with the actual latitudinal distribution of Cretaceous sea surface temperature, as cited by Baron et al. (1981).

Finally, we considered the effect of the evaporative constraint on the response of the Wisconsin Ice Age to astronomical and CO_2 forcing. According to Milankovitch (1941) and his current interpreters, the periodic advances and retreats of high-latitude ice sheets through the Pleistocene (2 Myr-10 kyr ago) were driven by quasi-cyclic variations in the earth's orbital eccentricity, precession and obliquity with periods of 100, 19 and 23, and 41 kyr, respectively (Figure 8). Hays et al. (1976) found evidence of this periodicity by spectrally analyzing oxygen isotope proxy temperatures in two Indian Ocean sediment cores. Suarez and Held (1979) applied a constant-D model forced by orbital variations over the past 150 kyr, and found a response qualitatively similar to the geologic record of that period, except for a phase lag predicting present-day temperatures colder than those 18 kyr ago (the peak of the Ice Age). Besides the astronomical explanation, the recent finding by Delmas et al. (1980) and Oeschger (1981) that CO_2 levels in gas samples trapped in polar icecores were roughly 1/2 of present-day values, suggests a greenhouse effect explanation of the last Ice Age.

We tested these theories by computing the iceline variation with solar absorptance poleward of the iceline, with obliquity (over the present range), with annual mean insolation and with atmospheric CO_2 levels (Figure 9). We found: (a) the TEC model is much more effective at moving the iceline toward the equator when forced by ice-albedo changes; (b) obliquity-driven changes in the iceline are too small to explain the Wisconsin Ice Age but significantly are reversed in phase from those with the constant-D assumption; (c) a 2% decrease in solar output is needed to move the iceline to the Wisconsin value, compared to 5% with the constant-D model (there is no reason to believe Q actually varied this much in the past 18 kyr) and (d) the lowered CO_2 levels of Oeschger are capable all by themselves of explaining the Wisconsin iceline in the TEC model (but, significantly, not in the constant-D model). Newell and Dopplick (1981) give further evidence of evaporation-limited temperatures during the CLIMAP period. Apparently, S. Manabe (personal communication) has been able to reproduce CLIMAP-like conditions using a GCM driven with such reduced CO_2 levels.

FIGURE 1: (a) zonal-annual-mean surface air temperature $T_a(x)$ and planetary absorptance of solar radiation $a(x)$ versus latitude, and (b) longwave radiation $I(T_a)$ to space versus surface air temperature. Surface and satellite data from Ellis and Vonder Haar (1976) and Warren and Schneider (1979). The crosses (+) and circles (o) are Northern and Southern Hemisphere data points, respectively. The water temperature curve $T_w(x)$ is also assumed parabolic in x , equal to the air temperature at the equator and to 273 K at the ice margin (see text).

FIGURE 2: Effective thermal diffusion coefficient and poleward heat flux across latitudes for the present climate; for the parabolic $T(x)$ distributions assumed here D is weakly dependent on x and is approximated by $\bar{D}$ which gives the $F(x)$ distribution of the solid curve. The atmospheric part of the total flux F_a is estimated by averaging northern and southern hemisphere data points for all ambient transports; the implied oceanic part $F_w = F - F_a$ is obtained as a residual.

FIGURE 3. Summary of changes in solar luminosity predicted by a standard solar model and changes in tropical ocean surface and bottom water temperature deduced from paleoclimate proxy records versus the logarithm of geologic time (adapted from Sears, 1964; Imbrie and Imbrie, 1979; Schopf, 1980 and Barron *et al.*, 1981).

FIGURE 4: Sea surface temperature versus latitude during Cretaceous, Present and Wisconsin Ice Age climates. Data points are annual and latitude band means generated by averaging values at given latitudes with equal weighting to Northern and Southern Hemispheres. Global mean icelines are estimated by intersection of sea surface temperature curve with water freezing point (see text). Polar regions were apparently ice-free in Cretaceous times.

FIGURE 5:

Equilibrium iceline latitude versus solar output for TEC and constant-D models.

Both limits recover the present $T(x)$ distribution of Figure 3(a) with $x^* \approx 0.94$ @ $Q/Q_0 = 1.00$; but the constant-D model has additional solutions at the current solar output, an unstable root at $x^* \approx 0.17$ and a stable one at $x^* = 0$ corresponding to an ice-covered planet.

FIGURE 6 Equilibrium values of (a) global mean air temperature $\bar{T}_a$, (b) equator-to-pole temperature difference T_1 , and (c) peak poleward heat flux normalized to present solar power incident on a hemisphere versus solar output for TEC and constant-D model.

FIGURE 7: Comparison of surface air temperature distributions in TEC and constant-D models having the same planetary mean temperature of 298.1 K with the paleotemperatures of the Cretaceous sea surface of Figure 2. This warmer mean temperature compared to the present 287.3 K corresponds to an increase in solar output or planetary absorptance of $\approx 9\%$; it would also have resulted from the greenhouse effect of an atmosphere whose CO_2 content was ≈ 15 times greater than today's.

FIGURE 8 Changes in the earth's orbital eccentricity, spin axis tilt (obliquity) and precession computed by Berger (1977). These are the changes driving advances and retreats of ice sheets during glacial ages according to the astronomical theory of Milankovitch (1941) and its current interpretations (Imbrie and Imbrie, 1979).

FIGURE 9: Annual mean iceline location with TEC and constant-D models for 4 different types of climatic forcing: (a) solar absorptance poleward of the iceline, (b) obliquity (axial tilt), (c) solar output and (d) atmospheric carbon dioxide levels. There are a number of possible causes for the Wisconsin glaciation, but the iceline location at that time seems most consistent with the diminished CO₂ levels inferred for that time from trapped gas samples and an evaporation limit on tropical ocean temperatures.

REFERENCES

- Barron, E.J., S.L. Thompson and S.H. Schneider 1981. An ice-free cretaceous results from climate model simulations. Science, 212, 501-508.
- Berger, A. 1977. Long-term variations of the earth's orbital elements Celestial Mechanics, 15, 53-74.
- Budyko, M. 1969: The effect of solar radiation variations on the climate of the earth, Tellus, 21, 5-13.
- Cess, R., 1976: Climate change: an appraisal of atmospheric feedback mechanisms employing zonal climatology, J. Atmos. Sci., 33, 1831-1843.
- CLIMAP Project Members 1976. The surface of the ice-age earth, Science, 191, 1131-1137.
- Delmas, R.J., J.M. Ascencio and M. Legrand 1980. Polar ice evidence that atmospheric CO₂ 20,000 P was 50% of present. Nature, 284, 155-157.
- Donn, W.J. and D.M. Shaw 1977. Model of climate evolution based on continental drift and polar wandering. Geol. Soc. America Bull., 88, 390-396.
- Geiger, R. 1960. The Climate Near the Ground, Academic Press, New York
- Hartmann, D.L. and D.A. Short 1979. On the role of zonal asymmetries in climate change, J. Atmos. Sci., 36, 519-528
- Hartmann, D. and D. Short, 1980. On the use of Earth radiation budget statistics for studies of clouds and climate, J. Atmos. Sci., 37, 1233-1250.
- Hays, J.D., J. Imbrie and N. Shackleton, 1976. Variations in the Earth's orbit: pacemaker of the Ice Ages, Science, 194, 1121-1131.
- Held, I. and M. Suarez, 1974. Simple albedo-feedback models of ice caps, Tellus, 26, 613-629.
- Hoffert, M., A. J. Callegari, C.T. Hsieh and W. Ziegler, 1981. Liquid water on Mars: an energy balance climate Model for CO₂/H₂O Atmosphere. Icarus (in press)
- Hoffert, M., A.J. Callegari and C.T. Hsieh 1980. The role of deep sea forcing in the secular response to climate forcing. J. Geophys. Res., 85, 6667-6679.
- Imbrie, J. and K.P. Imbrie, 1979, Ice Ages: Solving the Mystery, Enslow Publishers, Short Hills, NJ , 224 pp.

- Lindzen, R.S., and B. Farrell, 1977. Some realistic modifications of simple climate models J. Atmos. Sci., 34, 1487-1501.
- Milankovitch, M. 1941. Canon of Insolation and the Ice Age Problem. (Israel Program for Scientific Translations, Jerusalem, 1969, for U.S. Department of Commerce and National Science Foundation, Washington, D.C.)
- Newell, R. and T. Doplick, 1979. Questions concerning the possible influence of anthropogenic CO₂ on atmospheric temperature, J. Appl. Met., 18, 822-825.
- North, G., 1975. Analytical solution to a simple climate model with diffusive heat transport, J. Atmos. Sci., 32, 1301-1307.
- North, G. R., R. F. Cahalan, and J.A. Coakley, 1981. Energy balance climate models. Rev. Geophys. Space Phys., 19, 91-121.
- Oeschger, H. 1980. The history of atmospheric CO₂ as revealed from ice core studies. WMO project on research and monitoring of atmospheric CO₂. Rep. No. 1, Appendix G.
- Ohring, G. and P. Clapp, 1980. The effect of changes in cloud amount on the net radiation at the top of the atmosphere. J. Atmos. Sci., 37, 447-454.
- Oort, A.H. and Vonder Haar, T.H. 1976. On the observed annual cycle in the ocean-atmosphere heat-balance over the northern hemisphere. J. Phys. Oceanogr., 6, 781-800.
- Priestly, C.H.B. 1966. The limitation of temperature by evaporation in hot climates. Agric. Meteorol., 3, 241-246.
- Ramanathan, V. 1981. The role of ocean-atmosphere interactions in the CO₂-climate problem, J. Atmos. Sci.
- Schopf, T.J. M. 1980. Paleoceanography, Harvard University Press, Cambridge, Massachusetts, 341. pp.
- Sears, R.L. 1964. Helium content and neutrino fluxes in solar flux models. Astrophys. J. 146, 477-484.
- Sellers, W. 1969. A global climatic model based on the energy balance climate of the Earth-atmosphere system, J. Appl. Met., 8, 392-400.
- Suarez, M.J. and Held, I.M. 1979. The sensitivity of an energy balance climate model to variations in the orbital parameters. J. Phys. Res., 84, No. C8, 4825-4836.
- Trenberth, I.E. 1979. Mean annual poleward energy transports by the oceans in the southern hemisphere. Dyn. Atmos. Oceans., 4, 57-64.