APR 1 1987

DOE/PC/89785--9 9

DE92 011548

THE SINGLE ELECTRON CHEMISTRY OF COALS

Report for the period
October 1 to December 31 1991

John W. Larsen
and Thomas P. Eskay

Department of Chemistry
Lehigh University
Bethlehem, PA 18105

RECEIVED BY USN
APR 13 1992

PREPARED FOR THE UNITED STATES DEPARTMENT OF ENERGY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

OBJECTIVES AND SCOPE OF WORK

Objectives

To investigate and characterize the single electron reactions of alkyl and alkoxy aromatic compounds in order to determine the role these reactions play in the chemistry of coal.

Scope of Work

1. Attempts will be made to demonstrate that the radicals from inertinite maceral group will initiate the polymerization of 4-vinylpyridine.
2. The molecule, N,N-diphenyl-phenylenediamine, will be deposited in coals to characterize their native free radicals.
3. Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) will be used to characterize the numbers and strengths of single electron donors in coals.

Introduction

Depolymerization of coals at low temperatures may offer advantages over thermal bond cleavage. Because bond cleavage energies of radical cations are lower than the corresponding homolytic bond cleavage energies of the same bond^{1,2}, generation of radical cations in coal may make possible depolymerization at lower temperatures.

We seek to investigate the above possibility using single molecules containing functional groups common in coals. Since the generation of a radical cation requires the removal of an electron from a neutral molecule, a primary focus of the study will be finding oxidants that will remove an electron from compounds with structural similarity to those typically found in coals. The study must also be concerned with the decomposition of radical cations and the products formed as a result of the decomposition.

Results and Discussions

The single electron oxidation of benzyl phenyl ether with tris-(1,10-phenanthroline)-iron(III) perchlorate in refluxing acetonitrile produced the products phenol (P), benzyl acetamide (BAC), 2-benzylphenol (2BP), and 4-benzylphenol (4BP)³ (see reaction 1). Identification of these products was achieved by gas chromatography (GC) analysis. Previously we reported that

difficulty with the analysis made it impossible to obtain accurate mass balances⁴. Different response factors were found for the compounds produced in the reaction 1 in acetonitrile and methylene chloride, the solvent used for the reaction and the work-up extraction respectively. Because an accurate mass balance is necessary to ensure that other products are not being formed, the analysis must be perfected before proceeding with additional reactions.

Accurate mass balance determination requires that the response factors of the compounds in reaction 1, relative to an internal standard, are the same in both acetonitrile and methylene chloride, GC conditions were sought to give this consistency. Using a J&W DB-FFAP (15 m. x 0.25 mm.) column in the analysis, both methylene chloride and acetonitrile solutions containing the compounds in reaction 1, were analyzed using various GC conditions and injection sizes. Injections of 0.4 μl and an injection port temperature of 300°C gave the best results with phenol having the largest response factor variation: 1.16 ± 0.01 in methylene chloride and 1.24 ± 0.01 in acetonitrile (see tables I and II for results). To test these conditions a synthetic acetonitrile

reaction mixture was diluted with methylene chloride and a known amount of the internal standard, anthracene, was weighed into the mixture. GC analysis of the synthetic reaction mixture, using response factors obtained in methylene chloride failed to give quantitative agreement with the known amount weighed into the mixture (see table III). Furthermore, a portion of the synthetic reaction mixture was carried through the workup procedure (see experimental section for procedure) and GC analysis produced low recovery of compounds especially phenol (see table IV).

Because of the analytical difficulties described above, a new reaction workup procedure was developed. A small portion of aqueous sodium bicarbonate was added to the reaction mixture. The solution was dried with magnesium sulfate followed by GC analysis of the reaction mixture. This procedure was tested using synthetic reaction mixtures and was satisfactory.

Because anthracene was not soluble in acetonitrile, it was replaced as the internal standard by phenanthrene. Using a synthetic reaction mixture, two portions were carried through the workup procedure and one portion was used as a control. The weight of each compound in each portion determined from GC analysis agreed well with the known weight in the portion except for 4-benzylphenol (see table V). This compound gave GC analysis results 6 to 8 % too high. Since this is expected to be a minor product, this is not a significant error and studies of the reaction have begun.

Experimental

Materials

All compounds except benzyl acetamide (TCI), methylene chloride (Fisher A.C.S Grade), phenanthrene (Baker), and benzyl phenyl ether (see synthesis below) were purchased from Aldrich. 4-benzylphenol was recrystallized from n-hexane and phenanthrene was recrystallized from ethanol. All other chemicals were used from the bottle without further purification. Except for phenanthrene, which gave two small impurity peaks, all compounds used showed only one peak other than solvent in GC analysis.

NMR spectra were obtained on a Jeol FX90Q FT NMR spectrometer.

Benzyl Phenyl Ether

Benzyl phenyl ether was synthesized by a modification of a procedure by Pirkle⁵. In an oven dried, nitrogen purged, three neck 50 ml. flask fitted with a condenser and equipped with a magnetic stirrbar was placed 5.29 g. (38.3 mmol.) of anhydrous K_2CO_3 , 2.7 ml. (30 mmol.) of phenol, and 35 ml. of dried acetone (dried by shaking over $CaSO_4$ and distilling under nitrogen⁶). The resulting mixture was stirred for two hours prior to the addition of 3.8 ml. (34 mmol.) of benzyl chloride. The mixture was then heated to reflux for three days. The reaction solution was then filtered repeatedly washing the solid with acetone. The filtrate was then rotary evaporated to produce a yellow oil product $rf=0.58$ on Baxter MK6F/F254 TLC plates 1:4 CH_2Cl_2 :Petroleum ether b.p. 20-40°C. Flash chromatography of the oil with grade 60 (230-400 mesh) silica gel

using the same solvent system given for TLC produced a white crystalline powder which was recrystallized from ethanol twice. ^1H NMR 6.8-7.5 (m, 10H), 5.0 (s, 2H). The spectrum agreed well with previous spectra of benzyl phenyl ether.

Chromatography Analysis and Methods

Gas chromatography was carried out using a (15 m. x 0.25 mm. ID) J&W DB-FFAP column in a Hewlett-Packard 5880A gas chromatograph equipped with a FID detector. The carrier gas was He at a flow rate of 0.5 cm³/min. and the split ratio was 115:1. The column flow rate was determined by a methane injection at 180°C. The analysis was carried out using the following program: 180°C for 4.5 minutes, a 10°C/min. temperature increase to 240°C, and 9 minutes at 240°C with the injector temperature at 300°C and the detector at 250°C. The above program and conditions gave the following retention times in minutes: phenol 3.66, benzyl phenyl ether 6.89, benzylacetamide 9.52, phenanthrene 11.33, 2-benzylphenol 14.12, 4-benzylphenol 18.01.

The following is representative of how response factor standards and synthetic reaction mixtures were prepared. 0.20040 g. of BPE, 0.16245 g. of BAC, 0.29944 g. of phenanthrene, 0.06252 g. of 2BP, and 0.02697 g. of 4BP were weighed into a crimp top vial. The crimp top with a teflon lined septum was then crimped onto the vial and 0.09632 g. of phenol was then injected via syringe into the vial through the septum. This was followed by 15.95016 g. of acetonitrile also injected through the septum.

Two workup procedures were used and are described as follows

(see text). In the first procedure one ml. of a prepared synthetic reaction solution was removed from its crimp top vial by syringe. The solution was then injected into a preweighed 5 ml. conical vial capped and weighed. One ml. of saturated aqueous NaHCO_3 and one ml. of CH_2Cl_2 was added to the solution along with excess sodium chloride. The CH_2Cl_2 layer was removed after shaking the vial and placed into a vial containing a known amount of anthracene. The remaining aqueous layer was then extracted with 0.6 and 0.5 ml. of CH_2Cl_2 , and the organic layers from each extraction were added to the original organic layer removed. In the second workup procedure a known amount of anthracene was weighed into a 5 ml. conical vial. One ml. of the synthetic reaction mixture was injected into the vial and the weight was obtained. Three drops of saturated aqueous NaHCO_3 were then added to the solution and after shaking the vial magnesium sulfate was added. Compound weights present in each portion were determined by the weight percent of each compound in the master solution.

Response factors (RF) reported were calculated using the following equation and all data reported below are based on response factors relative to the internal standard phenanthrene except where otherwise noted.

$$RF = \frac{(\text{area of standard}) (\text{weight of compound})}{(\text{area of compound}) (\text{weight of standard})}$$

Data and Results

Table I. Response factors of compounds in reaction 1. determined in methylene chloride using 2-benzylphenol as the internal standard.

Compound	RF $\pm$ standard deviation
P	1.16 $\pm$ 0.01
BPE	0.98 $\pm$ 0.01
BAC	1.37 $\pm$ 0.02
4BP	1.17 $\pm$ 0.03

Table II. Response factors of compounds in reaction 1. determined in acetonitrile using 2-benzylphenol as the internal standard.

Compound	RF $\pm$ standard deviation
P	1.24 $\pm$ 0.01
BPE	1.00 $\pm$ 0.01
BAC	1.34 $\pm$ 0.01
4BP	1.14 $\pm$ 0.01

Table III. Weight percent of compound calculated from GC analysis divided by weight of compound weighed into solution. Anthracene internal standard.

Compound	Weight percent
P	94 $\pm$ 1 %
BPE	96 $\pm$ 1 %
BAC	98 $\pm$ 1 %
2BP	96 $\pm$ 1 %
4BP	99 $\pm$ 1 %

Table IV. Workup test. Weight percent of compound calculated from GC analysis divided by weight of compound weighed into solution. Anthracene internal standard.

Compound	Weight percent
P	86 %
BPE	88 %
BAC	90 %
2BP	94 %
4BP	96 %

Table V. Weight of compound determined by GC analysis divided by the weight of compound weighed into the solution after workup. Data given for two portions carried through the workup procedure and the control not carried through the procedure.

Compound	Portion 1	Portion 2	Control
P	100 %	99 %	98 %
BPE	99 %	101 %	99 %
BAC	99 %	98 %	100 %
2BP	99 %	100 %	100 %
4BP	106 %	108 %	108 %

References

1. D. M. Camaioni, J. A. Franz, J. Org. Chem., 1984, 49, 1601.
2. A. Okamoto, D. R. Arnold, Can. J. Chem., 1985, 63, 2340.
3. J. W. Larsen and T. Eskay, The Single Electron Chemistry of Coals, A paper prepared for the Department of Energy, # DE-FG22-89PC89785-7, August 1991
4. J. W. Larsen and T. Eskay, The Single Electron Chemistry of Coals, A paper prepared for the Department of Energy, # DE-FG22-89PC89785-8, October 1991
5. W. H. Pirckle and J. L. Schreiner, J. Org. Chem., 1981, 46, 4988.
6. D. D. Perrin and W. L. F. Armarego, Purification of Organic Compounds, Pergamon Press; New York, 1988, 3rd. edition.

END

**DATE
FILMED**

5/18/92