145
7-9-86 J5 (2)

38

LBL-21180
UC-4

Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Materials & Molecular Research Division

STRUCTURE OF
TRIS(TRIMETHYLSILYLCYCLOPENTADIENYL)
URANIUM(III), $[(CH_3)_3SiC_5H_4]_3U$

J. Brennan, R.A. Andersen, and A. Zalkin

February 1986

Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

LEGAL NOTICE

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Printed in the United States of America
Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161
Price Code: A02

LBL--21180

DE86 012828

Contribution from the Materials and
Molecular Research Division, Lawrence
Berkeley Laboratory and Department of
Chemistry, University of California,
Berkeley, California 94720 USA

Structure of Tris(trimethylsilylcyclopentadienyl)uranium(III),

John Brennan, Richard A. Andersen, and Allan Zalkin

Crystals of $[(CH_3)_3SiC_5H_4]_3U$ are orthorhombic, $Pbca$, with $a = 22.630(8)$, $b = 29.177(10)$ and $c = 8.428(3)$ Å at 23 °C. For $Z = 8$ the calculated density is 1.551 g/cm^3 . The structure was refined by full-matrix least-squares to a conventional R factor of 0.041 [2251 data, $F^2 > 2\sigma(F^2)$]. The uranium atom is bonded to the three cyclopentadienyl rings in a pentahapto fashion and is in the plane of the ring centroids. The U to ring distances are 2.54, 2.47 and 2.51 Å, and the average U-C distance is 2.78 ± 0.04 Å.

MASTER

643

Introduction. The title compound is one of several triscyclopentadienyl complexes of uranium(III) prepared for a study of their synthetic chemistry, structures and relative basicities.¹

Experimental. UCl_4 reacts with $\text{K}(\text{Me}_3\text{SiC}_5\text{H}_4)$ in tetrahydrofuran to give the red compound $(\text{Me}_3\text{SiC}_5\text{H}_4)_3\text{UCl}$ which when dissolved in hexane and stirred over Na/Hg or Na/K-yields the green compound $(\text{Me}_3\text{SiC}_5\text{H}_4)_3\text{U(III)}$.⁸ The air-sensitive crystals were sealed inside quartz capillaries in an argon filled drybox. X-ray diffraction intensities (0-20 scans) were obtained using a modified Picker FACS-I automatic diffractometer. The data were corrected for absorption (analytical method), crystal decay and Lorentz and polarization effects. Experimental details of the data collection are tabulated in Table I. The uranium atom position was deduced from three-dimensional Patterson maps, and subsequent least-squares refinements and electron density maps revealed the rest of the non-hydrogen atom positions. The structure was refined by full-matrix least-squares using anisotropic thermal parameters on all of the non-hydrogen atoms. Non-methyl hydrogen atoms were included in their estimated positions with fixed isotropic thermal parameters, but not refined; the methyl hydrogen atom positions could not be unambiguously estimated and were excluded. Atomic scattering factors and anomalous dispersion terms were taken from International Tables². Statistical results and other details of the least-squares refinements are tabulated in Table I.

Results Atomic parameters are listed in Table II. A list of selected bonds and angles is given in Table III. Tables of anisotropic thermal parameters, hydrogen positions, least-square planes, and additional bond distances and bond angles are given in the supplementary material.

The titled compound is a monomolecular complex of uranium bonded to three trimethylsilylcyclopentadienyl ligands in a pentahapto fashion. The uranium atom lies in the plane of the three cyclopentadienyl centroids with Cp-U-Cp angles all close to 120° (Table III); the average U-C distance is 2.78 ± 0.04 Å (Table III). An ORTEP³ schematic drawing of the structure is shown in Figure 1.

The U-Cp distances range from 2.48 to 2.54 Å and are comparable to the values found in the trimethylphosphine and thiophene coordination complexes of (MeCp)₃U(III); for (MeC₅H₄)₃U•PMe₃⁴ and (MeC₅H₄)₃U•SC₄H₈⁵ the U-Cp distances range from 2.51-2.54 Å and 2.54-2.55 Å, respectively. These U(III)-Cp distances are remarkably similar to those in some U(V)-Cp and U(IV)-Cp complexes; i.e. in (C₅H₅)₃U(V)=CHP(CH₃)₂(C₆H₅)⁶ the U-Cp distance range is 2.49 to 2.55 Å; in (C₅H₅)₃U(IV)(n-C₄H₉) and (C₅H₅)₃U(IV)(CH₂C₆H₄Me)⁷ the U-Cp distance range 2.45 to 2.52 Å.

Acknowledgment. This work was supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U. S. Department of Energy under Contract No. DE-AC03-76SF-00098.

References

1. Brennan, J. G. Ph.D. Thesis, Univ. of Calif. Berkeley, 1985, Lawrence Berkeley Laboratory Report No. 20845.
2. International Tables for X-ray Crstallography (1974). Vol. IV; Table 2.2, pp 71-102. Birmingham: Kynoch Press.
3. Johnson, C. K. (1965). ORTEP. Report ORNL-3794. Oak Ridge National Laboratory, Tennessee.
4. Brennan, J. G.; Zalkin, A. Acta Crystallogr., 1985, C41, 1038-1040.
5. Zalkin, A.; Brennan, J. G. Acta Crystallogr., 1985, C41, 1295-1297.
6. Cramer, R. E.; Maynard, R. B.; Paw, J. C.; Gilje, J. W. Organometallics, 1983, 2, 1336-1340.
7. Perego, G.; Cesari, M.; Farina, F.; Lugli, G. Gazz. Chim. Ital., 1985, 115, 643-646.

Table I. Crystallographic Summary and Data Processing for $[(CH_3)_3SiC_5H_4]_3U$

a, Å ^a	22.630(8)
b, Å	29.177(10)
c, Å	8.428(3)
cryst syst	orthorhombic
space group	Pbca
volume, Å ³	5564.8
d(calcd), g/cm ³	1.551
Z	8
temp (°C)	23
empirical formula	C ₂₄ H ₃₉ Si ₃ U
f(000)	2536
mol wt	649.87
x-ray	MoKα (graphite monochromated)
wave-length (Kα ₁ , Kα ₂), Å	0.70930, 0.71359
crystal size (mm)	0.21 x 0.30 x 0.48
abs coeff, cm ⁻¹	56.64
abs corr range	2.99-4.82
cryst decay corr range	0.60-1.27
2θ limits, deg	4.6-50.1
hkl limits	h 0, 26; k 0, 34; l -10, 10
scan type	θ-2θ
scan width, °2θ	1.20 + 0.693 × tanθ
no. of standards	3
no. reflections betwn stds	250
no. scan data	7595
no. unique reflections	4909

$R_{\text{int}}^{\text{b}}$	0.054
no. non-zero weighted data	2251 ($F^2 > 2\sigma$)
p^{c}	0.04
no. parameters	253
R (non-zero wtd dat) $^{\text{d}}$	0.041
R_{w}^{e}	0.041
R (all data)	0.147
Goodness of fit $^{\text{f}}$	1.22
max shift/esd in least-square	0.03
max,min in diff map ($e/\text{\AA}^3$)	1.78, -2.33

$^{\text{a}}$ Unit cell parameters were derived by a least-squares fit to the setting angles of the unresolved MoK α components of 29 reflections ($24^\circ < 2\theta < 34^\circ$).

$^{\text{b}}$ R_{int} = agreement factor between equivalent or multiply measured reflections = $\Sigma[I_{\text{hkl}} - \langle I_{\text{hkl}} \rangle] / \Sigma \langle I_{\text{hkl}} \rangle$

$^{\text{c}}$ In the least-squares, the assigned weights to the data are $1.0/[\sigma(F)]^2$ were derived from $\sigma(F^2) = [S^2 + (pF^2)^2]$, where S^2 is the variance due to counting statistics and p is assigned a value that adjusts the weighted residuals of the strong reflections to be comparable to the weak ones.

$^{\text{d}}$ $R = \Sigma(|F_{\text{obs}}| - |F_{\text{cal}}|) / \Sigma |F_{\text{obs}}|$

$^{\text{e}}$ $R_{\text{w}} = \sqrt{[\Sigma w(|F_{\text{obs}}| - |F_{\text{cal}}|)^2 / \Sigma w F_{\text{obs}}^2]}$

$^{\text{f}}$ σ_1 = error in observation of unit weight = $\sqrt{[\Sigma(w(|F_{\text{obs}}| - |F_{\text{cal}}|)^2) / (n_{\text{o}} - n_{\text{v}})]}$, where n_{o} is the number of observations and n_{v} is the number of variables.

Table II. Positional Parameters with Estimated Standard Deviations^afor $[(CH_3)_3SiC_5H_4]_3U$

atom	x	y	z
U	0.22514(2)	0.12380(2)	0.08037(6)
Si(1)	0.10228(21)	0.02486(15)	0.2233(7)
Si(2)	0.37799(21)	0.17855(16)	0.2461(8)
Si(3)	0.08792(20)	0.14995(15)	-0.2207(7)
C(1)	0.1792(7)	0.0451(4)	0.2281(20)
C(2)	0.2263(10)	0.0286(5)	0.1363(19)
C(3)	0.2812(10)	0.0449(6)	0.1966(27)
C(4)	0.2641(12)	0.0707(7)	0.343(3)
C(5)	0.2042(10)	0.0718(6)	0.3510(22)
C(6)	0.2976(7)	0.1887(5)	0.2182(21)
C(7)	0.2505(9)	0.1824(7)	0.3197(25)
C(8)	0.1979(8)	0.2001(5)	0.2570(29)
C(9)	0.2125(9)	0.2171(5)	0.1063(23)
C(10)	0.2725(10)	0.2113(4)	0.0833(21)
C(11)	0.1650(6)	0.1312(5)	-0.2127(15)
C(12)	0.1860(9)	0.0845(5)	-0.2028(19)
C(13)	0.2479(9)	0.0854(7)	-0.2063(23)
C(14)	0.2694(7)	0.1310(6)	-0.2292(18)
C(15)	0.2162(7)	0.1582(5)	-0.2296(18)
C(16)	0.0477(8)	0.0747(6)	0.229(3)
C(17)	0.0902(9)	-0.0116(8)	0.408(3)
C(18)	0.0899(8)	-0.0123(6)	0.0435(28)
C(19)	0.4193(9)	0.2355(6)	0.237(3)
C(20)	0.4058(8)	0.1407(7)	0.078(3)
C(21)	0.3937(8)	0.1506(6)	0.4463(23)
C(22)	0.0549(7)	0.0989(5)	-0.242(3)
C(23)	0.0798(9)	0.1874(7)	-0.401(3)
C(24)	0.0642(9)	0.1863(7)	-0.044(3)

^a Estimated standard deviations in this and subsequent tables are indicated in parentheses.

Table III: Bond Lengths and Angles for $[(CH_3)_3SiC_5H_4]_3U$

atoms			dist, Å	atoms			dist, Å
U	-	C(1)	2.811(14)	Si(1)	-	C(1)	1.839(16)
U	-	C(2)	2.816(14)	Si(1)	-	C(18)	1.885(21)
U	-	C(3)	2.806(17)	Si(1)	-	C(17)	1.905(24)
U	-	C(4)	2.843(21)	Si(1)	-	C(16)	1.909(15)
U	-	C(5)	2.780(17)	Si(2)	-	C(6)	1.860(16)
U	-	C(6)	2.761(14)	Si(2)	-	C(20)	1.900(23)
U	-	C(7)	2.706(20)	Si(2)	-	C(21)	1.905(19)
U	-	C(8)	2.748(17)	Si(2)	-	C(19)	1.914(15)
U	-	C(9)	2.746(16)	Si(3)	-	C(11)	1.829(14)
U	-	C(10)	2.768(13)	Si(3)	-	C(23)	1.882(21)
U	-	C(11)	2.828(13)	Si(3)	-	C(24)	1.903(23)
U	-	C(12)	2.792(15)	Si(3)	-	C(22)	1.920(14)
U	-	C(13)	2.713(18)	U	-	Cp(1) ^a	2.54
U	-	C(14)	2.802(16)	U	-	Cp(2)	2.48
U	-	C(15)	2.806(14)	U	-	Cp(3)	2.51

atoms			angle, °	atoms			angle, °
C(1)	-	Si(1) - C(16)	111.6(8)	C(20)	-	Si(2) - C(21)	110.7(9)
C(1)	-	Si(1) - C(17)	107.3(9)	C(11)	-	Si(3) - C(22)	111.6(7)
C(1)	-	Si(1) - C(18)	110.1(8)	C(11)	-	Si(3) - C(23)	107.2(8)
C(16)	-	Si(1) - C(17)	108.11(12)	C(11)	-	Si(3) - C(24)	114.0(9)
C(16)	-	Si(1) - C(18)	111.24(10)	C(22)	-	Si(3) - C(23)	108.38(10)
C(17)	-	Si(1) - C(18)	108.4(9)	C(22)	-	Si(3) - C(24)	109.10(10)
C(6)	-	Si(2) - C(19)	109.4(8)	C(23)	-	Si(3) - C(24)	106.27(10)
C(6)	-	Si(2) - C(20)	108.9(8)	Cp(1) - U	-	Cp(2)	120.2
C(6)	-	Si(2) - C(21)	111.3(9)	Cp(1) - U	-	Cp(3)	120.9
C(19)	-	Si(2) - C(20)	108.21(10)	Cp(2) - U	-	Cp(3)	118.0
C(19)	-	Si(2) - C(21)	108.25(10)				

^a Cp(1), Cp(2) and Cp(3) are the centroids of atoms C(1)-C(5), C(6)-C(10) and C(11)-C(15) respectively.

Figure Caption

Fig. 1. ORTEP drawing of tris(trimethylsilylcyclopentadienyl)uranium(III); thermal ellipsoids at 50% probability level.

Fig. 1

Supplemental Material

Structure of Tris(trimethylsilylcyclopentadienyl)uranium(III),

John Brennan, Richard A. Andersen, and Allan Zalkin

Crystals of $[(CH_3)_3SiC_5H_4]_3U$ are orthorhombic, $Pbca$, with $a = 22.630(8)$, $b = 29.177(10)$ and $c = 8.428(3)$ Å at 23 °C. For $Z = 8$ the calculated density is 1.551 g/cm^3 . The structure was refined by full-matrix least squares to a conventional R factor of 0.041 [2251 data, $F^2 > 2\sigma(F^2)$]. The uranium atom is bonded to the three cyclopentadienyl rings in a pentahapto fashion and is in the plane of the ring centroids. The U to ring distances are 2.54, 2.47 and 2.51 Å, and the average U-C distance is 2.78 ± 0.04 Å.

Table S1. Thermal Parameters^a (Å²) with Estimated Standard Deviations for
[(CH₃)₃SiC₅H₄]₃ U

Atom	B ₁₁	B ₂₂	B ₃₃	B ₁₂	B ₁₃	B ₂₃
U	4.713(23)	2.641(17)	3.046(19)	0.053(29)	0.08(3)	-0.044(27)
SI(1)	5.27(26)	5.39(23)	7.1(3)	0.04(19)	0.60(24)	1.66(24)
SI(2)	5.80(28)	6.18(25)	7.2(3)	-1.82(20)	-1.14(26)	0.40(26)
SI(3)	5.54(27)	4.95(21)	7.3(3)	-1.11(18)	-1.25(25)	1.14(23)
C(1)	5.6(9)	2.8(6)	4.0(8)	-1.0(6)	-1.0(8)	0.1(6)
C(2)	7.0(10)	3.0(6)	5.2(9)	0.3(9)	-1.5(11)	0.5(6)
C(3)	6.2(12)	6.6(10)	5.5(12)	-1.4(10)	-0.5(12)	3.3(9)
C(4)	7.9(15)	5.8(11)	9.3(17)	-2.0(12)	-4.4(15)	4.4(11)
C(5)	8.7(13)	4.9(8)	4.9(9)	-2.7(8)	1.4(9)	0.8(7)
C(6)	5.9(10)	4.0(7)	3.1(8)	-0.7(6)	-0.8(7)	-0.4(6)
C(7)	9.9(18)	5.7(12)	2.4(9)	-0.5(8)	1.0(8)	-1.6(8)
C(8)	7.2(12)	4.0(8)	6.5(12)	0.9(7)	0.6(11)	-0.2(8)
C(9)	6.5(12)	4.4(7)	5.6(11)	0.7(7)	-1.7(9)	-1.5(7)
C(10)	7.9(10)	3.0(6)	4.7(8)	-0.5(8)	-0.2(13)	-0.9(7)
C(11)	4.7(8)	4.9(8)	3.4(7)	-0.9(7)	-0.6(6)	0.0(6)
C(12)	7.9(12)	5.0(9)	4.8(10)	0.5(9)	-2.1(10)	-1.0(7)
C(13)	10.4(15)	5.9(11)	3.5(10)	2.6(9)	-1.0(8)	-1.8(8)
C(14)	6.1(9)	7.0(11)	3.2(6)	-0.7(10)	-1.8(8)	0.4(8)
C(15)	5.2(10)	5.2(8)	3.1(7)	0.3(7)	-0.6(7)	1.4(6)
C(16)	8.2(12)	10.2(12)	12.0(16)	5.5(10)	-2.7(13)	-3.0(14)
C(17)	8.1(12)	12.8(16)	11.0(16)	-3.8(11)	0.0(14)	5.9(15)
C(18)	6.7(11)	6.7(10)	11.3(18)	-2.9(8)	-0.5(11)	-2.9(11)
C(19)	12.9(16)	7.9(11)	9.3(13)	-6.5(10)	-0.4(14)	1.4(12)
C(20)	6.8(11)	11.3(14)	11.1(15)	1.1(10)	-0.9(13)	-1.5(14)
C(21)	8.3(11)	9.3(11)	6.5(12)	-2.2(9)	-2.6(10)	3.3(10)
C(22)	7.0(11)	6.1(8)	11.1(13)	-2.8(8)	-1.2(12)	0.7(11)
C(23)	7.6(12)	11.1(14)	17.9(23)	-1.5(10)	-3.2(14)	11.3(16)
C(24)	7.6(12)	9.5(13)	14.4(21)	-0.8(10)	3.2(13)	-4.5(14)

H(1) - H(12) B = 10.0

^a The anisotropic temperature factor has the form:

$$-0.25\exp(h^2 a^{*2} B_{11} + 2hka^* b^* B_{12} + \dots).$$

The isotropic temperature factor has the form: $\exp-B(\sin\theta/\lambda)^2$.

Table S2. Estimated Hydrogen Positional Parameters in

$$[(CH_3)_3SiC_5H_4]_3 U$$

Atom	x	y	z
H(1) C(2)	0.2219	0.0088	0.0433
H(2) C(3)	0.3207	0.0399	0.1529
H(3) C(4)	0.2917	0.0850	0.4180
H(4) C(5)	0.1816	0.0885	0.4306
H(5) C(7)	0.2537	0.1680	0.4243
H(6) C(8)	0.1588	0.2000	0.3053
H(7) C(9)	0.1852	0.2312	0.0303
H(8) C(10)	0.2945	0.2213	-0.0115
H(9) C(12)	0.1614	0.0568	-0.1947
H(10) C(13)	0.2734	0.0586	-0.1939
H(11) C(14)	0.3100	0.1415	-0.2406
H(12) C(15)	0.2159	0.1915	-0.2410

Table S3. Additional Distance and Angles in $[(CH_3)_3SiC_5H_4]_3U$

atoms	d(Å)	atoms	d(Å)
C(1) - C(2)	1.402(23)	H(1) - C(2)	0.979
C(1) - C(5)	1.415(21)	H(2) - C(3)	0.978
C(2) - C(3)	1.425(26)	H(3) - C(4)	0.982
C(3) - C(4)	1.497(27)	H(4) - C(5)	0.972
C(4) - C(5)	1.358(28)	H(5) - C(7)	0.979
C(6) - C(7)	1.378(21)	H(6) - C(8)	0.974
C(6) - C(10)	1.431(21)	H(7) - C(9)	0.979
C(7) - C(8)	1.401(23)	H(8) - C(10)	0.986
C(8) - C(9)	1.403(26)	H(9) - C(12)	0.983
C(9) - C(10)	1.382(28)	H(10) - C(13)	0.978
C(11) - C(15)	1.409(17)	H(11) - C(14)	0.974
C(11) - C(12)	1.447(18)	H(12) - C(15)	0.976
C(12) - C(13)	1.401(21)		
C(13) - C(14)	1.429(23)		
C(14) - C(15)	1.441(19)		

atoms	angle(°)	atoms	angle(°)
SI(1) - C(1) - C(2)	126.7(13)	C(6) - C(7) - C(8)	112.0(19)
SI(1) - C(1) - C(5)	124.9(16)	C(7) - C(8) - C(9)	105.8(18)
C(1) - C(2) - C(3)	110.6(17)	C(8) - C(9) - C(10)	108.3(16)
C(2) - C(3) - C(4)	103.6(21)	C(6) - C(10) - C(9)	109.6(17)
C(3) - C(4) - C(5)	108.2(24)	C(12) - C(11) - C(15)	105.2(13)
C(1) - C(5) - C(4)	110.4(20)	C(11) - C(12) - C(13)	108.0(15)
SI(2) - C(6) - C(7)	131.0(14)	C(12) - C(13) - C(14)	111.2(15)
SI(2) - C(6) - C(10)	124.3(15)	C(13) - C(14) - C(15)	103.3(13)
C(7) - C(6) - C(10)	104.3(16)	C(11) - C(15) - C(14)	112.3(13)

Table S4. Least-squares Planes and Deviations from Least-squares planes
in $[(CH_3)_3SiC_5H_4]_3U$

Equation of Plane #1. Atoms C(1) - C(5)
 $-1.17127 a + 23.95055 b - 4.79354 c = -0.22019$

Deviations from Plane

atom	d(Å)	atom	d(Å)
C(1)	-0.003(22)	U	2.536(10)
C(2)	-0.012(28)	Si(1)	-0.375(33)
C(3)	0.023(30)		
C(4)	-0.041(35)		
C(5)	0.019(28)		

Equation of Plane #2. Atoms C(6) - C(10)
 $4.17598 a + 25.98811 b + 3.50068 c = 6.91270$

Deviations from Plane

atom	d(Å)	atom	d(Å)
C(6)	-0.001(23)	U	-2.473(09)
C(7)	-0.006(26)	Si(2)	0.164(33)
C(8)	0.014(30)		
C(9)	-0.010(25)		
C(10)	0.009(27)		

Equation of Plane #3. Atoms C(11) - C(15)
 $1.15662 a + 2.86370 b + 8.37624 c = -1.22051$

Deviations from Plane

atom	d(Å)	atom	d(Å)
C(11)	0.006(18)	U	2.509(08)
C(12)	-0.021(24)	Si(3)	-0.097(27)
C(13)	0.024(25)		
C(14)	-0.012(23)		
C(15)	0.001(20)		

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.