48
3/10/92 JS @

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2821
UC-420,427

PPPL-2821

THEORY OF MINIMUM DISSIPATION OF ENERGY
FOR THE STEADY STATE

BY

T.K. CHU

February 1992

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
703-487-4650

Theory of minimum dissipation of energy for the steady state

by

PPPL--2821

DE92 008650

T. K. Chu

Plasma Physics Laboratory, Princeton University

Princeton, New Jersey 08543

The magnetic configuration of an inductively driven steady-state plasma bounded by a surface (or two adjacent surfaces) on which $\mathbf{B} \cdot \mathbf{n} = 0$ is force-free: $\nabla \times \mathbf{B} = 2\alpha \mathbf{B}$, where α is a constant, in time and in space. α is the ratio of the Poynting flux to the magnetic helicity flux at the boundary. It is also the ratio of the dissipative rates of the magnetic energy to the magnetic helicity in the plasma. The spatial extent of the configuration is noninfinitesimal. This global constraint is a result of the requirement that, for a steady-state plasma, the rate of change of the vector potential, $\partial \mathbf{A} / \partial t$, is constant in time and uniform in space.

PACS numbers: 52.53Fa, 41.10.-j, 47.25.-c, 52.55 Hc

MASTER

EP

To maintain the steady state, a plasma needs a constant supply of energy since, as it is commonly understood, energy is dissipated continuously. In a current-carrying plasma, the replenishing energy that sustains the current is often drawn from an external electromagnetic field through the Poynting flux. When magnetic energy is supplied to a plasma, magnetic helicity is also supplied, the latter characterizing the ordered topological structure of the field - the linking of flux tubes and the twisting, kinking and knotting of a flux tube. In a hypothetical plasma in which no dissipation occurs, the topological structure of the field cannot change and magnetic helicity is an invariant. In a real plasma, magnetic helicity, like magnetic energy, is dissipated Ohmically. What is the mutual constraint, in the steady state, between the dissipative rates of energy and helicity? This question is addressed here. But first it is useful to recount historical background on the subjects of the force-free state and the minimum-energy principle.

Lundquist¹ first introduced the concept of force-free field as a candidate characterizing the interstellar magnetic field $\mathbf{B}$ embedded in a conducting fluid. A force-free field is described by

$$\nabla \times \mathbf{B} = 2\alpha \mathbf{B}. \quad (1)$$

Lüst and Schlüter² also pointed out that cosmic magnetic fields often satisfy this condition. Chandrasekhar³ gave general solutions to Eq. (1) for fields with spherical symmetry and a constant α . The constant- α condition, meaning α being discontinuous at the boundary S ($\mathbf{B} \cdot \nabla \alpha = 0$) whose unit normal vector is $\mathbf{n}$, requires

$$\mathbf{B} \cdot \mathbf{n} = 0 \text{ on } S, \quad (2)$$

that is, the surface is a magnetic surface (or a conductor) held at static equilibrium. In a further inquiry, Chandrasekhar and Woltjer⁴ showed that a plasma in a force-free field can be interpreted as being in either a state of maximum magnetic energy for a given mean-square current density (maximum $\int_V B^2 dV$ for a given $\int_V j^2 dV$, with the volume V defined by S) or a state of minimum energy dissipation for a given magnetic energy (minimum $\int_V j^2 dV$ for a given $\int_V B^2 dV$), with the resistivity assumed to be uniform in V .

Soon afterwards, Woltjer, not satisfied with these two interpretations, showed in a paper⁵ communicated by Chandrasekhar that helicity⁶ of the field in a fluid with infinite conductivity is an invariant:

$$K = \int_V \mathbf{A} \cdot \mathbf{B} dV = \text{const.}, \quad (3)$$

where $\mathbf{A}$ is the vector potential, $\mathbf{B} = \nabla \times \mathbf{A}$, and the volume V is finite or the whole space - not infinitesimal. (For an inductively sustained toroidal plasma, the toroidal component of $\mathbf{A}$ on the surface of the torus and of $\mathbf{B}$ in the torus determine the linking, with V being the volume of all space. Let $V = V_t + V_h$, where V_t is the volume of the torus and V_h is the remaining volume that includes the torus hole where the inductive core is placed. For the flux linking in V_t , let C_t be a toroidal loop on the surface of V_t whose poloidal cross-sectional area is a_p , then $\int_{C_t} \mathbf{A} \cdot d\mathbf{l} = \psi_\theta$, the total poloidal flux from the inductive core, and $\int_{a_p} \mathbf{B} \cdot d\mathbf{a}_p = \psi_\phi$, the total toroidal flux in the plasma. Thus, $K = 2\psi_\phi\psi_\theta$, where the factor two comes from an equal amount of linking in V_h). Woltjer then derived the force-free condition by a variational method applied to the magnetic energy, subject to the constraint of constant helicity - hence the dissipation needed for the configuration to evolve was ignored. Reversing the earlier argument, Woltjer stated that this stationary value of energy is intuitively obvious being the minimum magnetic energy, since when the plasma is at that state the field can produce no motion and thus the Lorentz force vanishes. The Woltjer principle, that the force-

free field corresponds to the state of minimum magnetic energy, gained popularity after it was applied - relying on an argument that energy is dissipated much faster than helicity during "relaxation" and hence the total helicity is conserved - to laboratory plasmas in reversed field pinches (RFP).⁷ In recent years, the principle and its modified forms have also been used to explain solar corona heating.⁸ In the last few years, it has been applied to tokamak plasmas, but with the plasma current as the substituted constraint.⁹⁻¹¹ The glaring contradiction between the concept of minimum magnetic energy and the earlier conclusion of maximum magnetic energy (or minimum energy dissipation) has not been resolved. The incompatibility between the notion of minimum magnetic energy and a centrally peaked profile of current density j and the observed abundance of instabilities has been ignored. The meaning and determination of the proportionality constant between j and B , 2α , have also evaded a solution.

The present work shows that the magnetic configuration of an inductively driven steady-state plasma is force-free and α is the ratio of the Poynting flux to the helicity flux at the boundary as well as the ratio of the dissipative rates of magnetic energy to helicity in the enclosed, finite-volume plasma. The minimum dissipation rate of energy corresponds to

the lowest eigenvalue of the solution of the force-free field. These effects are results of the requirement that, in the steady state, $\partial \mathbf{A} / \partial t$ is constant in time and uniform in space.

Let W be the energy in a dissipative plasma in a finite volume V ,

$$W = \int_V [B^2/2 + p/(\gamma - 1)] dV, \quad (4)$$

where p is the plasma pressure and γ is the ratio of specific heats. The boundary conditions are $\mathbf{B} \cdot \mathbf{n} = 0$ and $\mathbf{j} \cdot \mathbf{n} = 0$. To allow energy and helicity flow across the boundary, a surface voltage, and therefore an electric field, is set up on the surface (or across some gaps on the surface). Calculate the rate of change of W and obtain

$$\partial W / \partial t = - \int_S \mathbf{E} \times \mathbf{B} \cdot \mathbf{n} dS + \int_V [(\gamma - 1)^{-1} \partial p / \partial t - \mathbf{E} \cdot \mathbf{j}] dV, \quad (5)$$

which, when the energy is at the steady state, reduces to

$$\int_S \mathbf{E} \times \mathbf{B} \cdot \mathbf{n} dS = - \int_V \mathbf{E} \cdot \mathbf{j} dV, \quad (6)$$

equating the Poynting flux (energy is supplied when $\mathbf{B}$ has a component perpendicular to $\mathbf{E}$) to the dissipation of magnetic energy in the plasma. In Eq. (6), $\mathbf{E}$ (that is, $\partial \mathbf{A} / \partial t$) is

constant in time. This integral form of power balance is more general than its differential-form counterpart since it allows singularities on the surface as well as discrete surfaces within the volume [$V = \Delta V_1 + \Delta V_2 + \dots$, with ΔV replacing V and the difference of two (adjacent) surface integrals replacing the surface integral in Eq. (6); if each noninfinitesimal ΔV is separately force-free, V is piecewise force-free].

The magnetic helicity in V can be expressed by a gauge-invariant helicity,¹²

$$K_I = \int_V (\mathbf{A} + \mathbf{A}') \cdot (\mathbf{B} - \mathbf{B}') dV, \quad (7)$$

where all fields extend to the whole space of which V is a part. $\mathbf{A}$ and $\mathbf{B}$ are fields of interest. $\mathbf{A}'$ and $\mathbf{B}'$ are fixed reference fields (vacuum fields) differing from $\mathbf{A}$ and $\mathbf{B}$, respectively, only in V . The tangential components of $\mathbf{A}$ and $\mathbf{A}'$ are equal on the boundary and $\partial \mathbf{A}' / \partial t$ is zero in V . Calculate the rate of change of helicity, use Eq. (2) and obtain

$$\partial K_I / \partial t = -2 \int_S \mathbf{E} \times \mathbf{A} \cdot \mathbf{n} dS - 2 \int_V \mathbf{E} \cdot \mathbf{B} dV, \quad (8)$$

in which the surface integral is the helicity inflow^{13,14} and the vector $2\mathbf{E} \times \mathbf{A}$ can be regarded as the helicity flux vector

(helicity is supplied when $\mathbf{B}$ has a component parallel to $\mathbf{E}$).

When helicity is at the steady state, the helicity flux balances its dissipation:

$$2 \int_S \mathbf{E} \times \mathbf{A} \cdot \mathbf{n} dS = -2 \int_V \mathbf{E} \cdot \mathbf{B} dV. \quad (9)$$

For a steady-state plasma, Eqs. (6) and (9) must be satisfied simultaneously. This requires

$$\frac{\int_V \mathbf{E} \cdot \mathbf{j} dV}{2 \int_V \mathbf{E} \cdot \mathbf{B} dV} \equiv \frac{\dot{W}_B}{K} = \alpha \quad (10)$$

$$\text{and} \quad \frac{\int_S \mathbf{E} \times \mathbf{B} \cdot \mathbf{n} dS}{2 \int_S \mathbf{E} \times \mathbf{A} \cdot \mathbf{n} dS} \equiv \frac{\Gamma_B}{\Gamma_K} = \alpha, \quad (11)$$

where α , equal to the ratio of the dissipative rates of energy to helicity in the volume and the ratio of the fluxes of energy to helicity at the boundary, must be a constant.

The magnetic configuration corresponding to an α constant in time can be obtained from Eq. (10) by varying $\dot{W}_B$, subject to the constraint of constant K :

$$\begin{aligned}
\delta(\dot{W}_B - \alpha \dot{K}) &= \delta \int_V (\mathbf{E} \cdot \mathbf{j} - 2\alpha \mathbf{E} \cdot \mathbf{B}) dV \\
&= \delta \int_V (-\partial \mathbf{A} / \partial t - \nabla \phi) \cdot (\mathbf{j} - 2\alpha \mathbf{B}) dV = 0.
\end{aligned}
\tag{12}$$

The terms involving the electrostatic potential ϕ vanishes when the integral is converted to a surface integral because of the boundary conditions. In a steady-state plasma, in V ,

$$\partial \mathbf{A} / \partial t = \text{const.}, \tag{13}$$

as a nonuniform $\partial \mathbf{A} / \partial t$ would result in a time-dependent $\mathbf{B}$ field

$[\mathbf{B} = \nabla \times (\mathbf{A}_i + \delta t \partial \mathbf{A}(\mathbf{r}) / \partial t) = \mathbf{B}_i + \delta \mathbf{B}(t)$, where the subscript

i denotes the initial value]. Equation (12) becomes

$\partial \mathbf{A} / \partial t \cdot \int_V [\partial (\mathbf{j} - 2\alpha \mathbf{B}) / \partial \mathbf{B}] \delta \mathbf{B} dV = 0$. For an arbitrary $\delta \mathbf{B}$,

$\partial (\mathbf{j} - 2\alpha \mathbf{B}) / \partial \mathbf{B} = 0$. Integrate once, use the condition that

$\mathbf{B} = 0$ when $\mathbf{j} = 0$ and obtain Eq. (1) in which α is constant, in

space. The lowest eigenvalue of the solution of Eq. (1)

corresponds to the minimum rate of dissipation of magnetic

energy for a given rate of dissipation of magnetic helicity; it

also gives the Poynting flux needed to dissipate the supplied

helicity influx.

The following are some consequences of the theory.

1. Pinch velocity. For a toroidal plasma, the pinch velocity of the magnetic energy and of the quantity $\mathbf{A} \cdot \mathbf{B}$ is

$$\mathbf{v}_p = 2\mathbf{E} \times \mathbf{B} / B_\theta^2, \quad (14)$$

where B_θ is the poloidal magnetic field. In a dissipative plasma, the particle pinch velocity across a magnetic surface^{15,16} (on which $\mathbf{B} \cdot \mathbf{n} = 0$) is, rightly, the same as that of magnetic energy and helicity. (In a hypothetical plasma in which there is a constant $\mathbf{E}$ field but no dissipation the inward radial velocity of a flux surface¹⁵ or particles¹⁷ is only one half of $\mathbf{v}_p$.)

2. Magnetic configuration. Evaluate Eq. (11) for a toroidal plasma and obtain

$$\alpha = V_L I_p / 2V_L \Psi_\phi = I_p / 2\Psi_\phi, \quad (15)$$

where V_L is the loop voltage and I_p is the plasma current. Thus,

$$\mathbf{j} = (I_p / \Psi_\phi) \mathbf{B}. \quad (16)$$

Note that although α is the ratio of the dissipative rates, it does not depend on the magnitude of the loop voltage (the configuration does not depend on the actual mechanism of dissipation as long as the dissipative process can be described by Ohm's law). Also, since $\partial \mathbf{A} / \partial t = \mathbf{v} \times \mathbf{B} - \eta \mathbf{j} - \nabla \phi$, where

$\mathbf{v}$ is the fluid velocity, uniform $\partial \mathbf{A} / \partial t$ results in

$\partial \mathbf{A} / \partial t \cdot \int_V \mathbf{B} dV = \int_V \eta \mathbf{j} \cdot \mathbf{B} dV$ and η cannot be assumed, in general, to be uniform in V .

3. Time scales of the dissipation rates. Form dot product of Eq. (1) with $\mathbf{A}$, integrate over V and obtain¹⁸

$$\alpha = \frac{W_B - W_S}{K}, \quad (17)$$

where $W_B = \int_V (B^2/2) dV$ is the total magnetic energy,

$W_S = (1/2) \int_S (\mathbf{A} \times \mathbf{B}) \cdot \mathbf{n} dS$ is the sum of the magnetic energy due to an induced current in the boundary⁴ and the external currents, and αK is the magnetic energy required to assemble the current in V .¹⁹ W_S is nonzero for a toroidal plasma. Insert this α in Eq. (10) and obtain

$$\frac{\dot{W}_B}{W_B - W_S} = \frac{\dot{K}}{K}. \quad (18)$$

The time scales of the two volume dissipation rates are, rightly, equal.

4. Scale length of a force-free configuration. α determines the domain of the force-free field, L_B :

$$\alpha = |\nabla \times \mathbf{B}| / 2|\mathbf{B}| \equiv 1/2L_B \quad (19)$$

(the minor radius of a toroidal vessel, for example, generally does not). As long as α is finite, the spatial extent of a force-free configuration is finite ($L_B \rightarrow 0$ only if $\alpha \rightarrow \infty$, that is, only if $|\mathbf{j}| = |\nabla \times \mathbf{B}| \rightarrow \infty$), as demanded by Ampere's law.

5. The bases for a force-free configuration. In a steady-state plasma, the force-free condition is a result of dissipating the supplied magnetic helicity. Berger⁸ and Boozer²⁰ showed that if there exists a process that dissipates energy and helicity at rates faster than the corresponding rates at the force-free condition, the enhanced rate of dissipation of magnetic energy due to this process exceeds that of magnetic helicity. Such a process, not associated with a force-free configuration, is also not in the steady state. Also, a force-free configuration requires static equilibrium of its boundary. Given a boundary at static equilibrium, the enclosed finite-volume plasma, at the steady state, is force-free. The present theorem does not specify how the boundary (static equilibrium) condition, $\mathbf{B} \cdot \mathbf{n} = 0$, can be maintained: it does not guarantee that a steady state exists; nor does the theorem indicate how the current density and the plasma pressure can be kept constant in time when heating occurs as a result of dissipation.

6. Pressure containment and magnetic surfaces. The global constraint results in zero $\mathbf{j} \times \mathbf{B}$ force in a force-free region. Hence, in this region there is no diamagnetic current and, in a toroidal plasma, no secondary current (Pfirsch-Schlüter current).^{21,22} Equation (2) shows that a discontinuity on a surface distinguishes the surface as a magnetic surface.²³ Known analytic solutions^{1,24} of force-free configurations often have internal surfaces with $B_n = 0$ but without a discontinuity in α . Such a configuration, when realized by inductive-drive means, is in fact made of piecewise force-free regions²⁵ all having the same time-averaged α and hence unsuitable for confinement purposes.

7. Concept of energy dissipation and of Poynting flux. To maintain the steady throughput of the magnetic flux (in a torus, the flux generated by the transformer), the ordered topological structure of the field associated with the flux can be conceptually regarded as entering the plasma from the boundary and to be dismantled in the interior. The required rate of dissipation of magnetic energy simply equals the rate at which work is done to dissipate that structure. The difference between a conservative field and a field that is no longer conservative can be viewed as that the topological structure of the latter has undergone dissipation while that of the former has remained intact. Alternatively, as work is done to pare

down the order of the structure, the supplied magnetic energy takes on a measurable form as "the energy that is dissipated." After the ordered structure of the field is taken apart, no more expenditure of magnetic energy is needed, nor there is any left to increase the stored energy. The cause of energy dissipation is the need to dismantle the topological structure of the field.

The concept of Poynting flux requires that the energy of an electromagnetic field flows into a system from its boundary in a direction along which $\mathbf{E}$, and hence $\mathbf{j}$, have no component. Physical descriptions of the concept have not been completely satisfactory.²⁶ When helicity flux and dissipation are included in the consideration, the concept of Poynting flux - the concept of energy dissipation borne by Poynting flux - becomes quite natural: it is the topological structure of the field that enters from the boundary; and energy dissipation, that is, the dissipation of the field topology, begins at the boundary. Hence, both Poynting flux and helicity flux are properly defined electrodynamic concepts in a real (dissipative) plasma.

In summary, in an inductively driven steady-state plasma, the ratio of the supply rates of energy and helicity through a closed surface held at static equilibrium equals the ratio of the corresponding dissipation rates in the finite volume enclosed by the surface, and the rate of change of the vector potential, $\partial \mathbf{A} / \partial t$, is constant in time and uniform in space.

These requirements result in the global force-free constraint.

ACKNOWLEDGEMENT. I benefited from discussions with Wolfgang Stodiek, John Johnson, Cris Barnes, Allan Reiman and Schweickhard von Goeler. This work is supported by U. S. Department of Energy Contract No. DE-AC02-76-CHO-3073.

Reference

- 1 S. Lundquist, Arkiv. fysik, Bd. 2, 35 (1950); Phys. Rev., **83**, 307 (1951).
- 2 R. Lüst and A. Schlüter, Z. Astrophys. **34**, 263 (1954).
- 3 S. Chandrasekhar, Proc. of Nat. Acad. of Sci. **42**, 1 (1956).
- 4 S. Chandrasekhar and L. Woltjer, Proc. of Nat. Acad. of Sci., **44**, 285 (1958).
- 5 L. Woltjer, Proc. of Nat. Acad. of Sci., **44**, 489 (1958).
- 6 H. K. Moffatt, *Magnetic Field Generation in Electrically Conducting Fluids*, (Cambridge U. P., Cambridge, 1978).
- 7 J. B. Taylor, Phys. Rev. Lett. **33**, 1139 (1974); Rev. Modern Phys. **58**, 741 (1986).
- 8 M. A. Berger, Geophys. Astrophys. Fluid Dynamics, **30**, 79 (1984); the references in this paper include a partial listing of the work on solar corona heating.
- 9 D. Biskamp, Comments Plasma Phys. Controlled Fusion, **10**, 165 (1986).
- 10 B. B. Kadomtsev, Comments Plasma Phys. Controlled Fusion, **11**, 153 (1987).
- 11 J. Y. Hsu and M. S. Chu, Phys. Fluids **30**, 1221 (1987).
- 12 J. M. Finn and T. M. Antonsen, Jr., Comments Plasma Phys. Controlled Fusion **9**, 111 (1985).
- 13 T. H. Jensen and M. S. Chu, Phys. Fluids, **27**, 2881 (1984).
- 14 M. A. Berger and G. B. Field, J. Fluid Mechanics, **147**, 133

- (1984).
- 15 F. L. Hinton and R. D. Hazeltine, *Rev. of Modern Phys.* **48**, 239 (1976).
 - 16 I. B. Bernstein and K. Molvig, *Phys. Fluids*, **26**, 1488(1983).
 - 17 A. A. Ware, *Phys. Rev. Lett.* **25**, 15 (1970).
 - 18 A. Reiman, *Phys. Fluids* **23**, 230 (1980).
 - 19 A. Reiman, *Phys. Fluids* **24**, 956 (1981); J. A. Stratton, *Electromagnetic Theory*, (McGraw-Hill, New York, 1941), p.122.
 - 20 A. H. Boozer, *J. Plasma Phys.* **35**, 133 (1986).
 - 21 L. Spitzer, Jr., *Phys. Fluids*, **1**, 253 (1958).
 - 22 M. D. Kruskal and R. M. Kulsrud, *Phys. Fluids*, **1**, 265, (1958).
 - 23 H. Grad and H. Rubin, in *Proc. of 2nd U.N. Int. Conf. on the Peaceful Uses of Atomic Energy* 1958, Geneva (Columbia U. P., New York, 1959), Vol. 31, p. 190.
 - 24 J. Hastie, *Nuclear Fusion* **29**, 96 (1989).
 - 25 T. K. Chu, "Theory and application of maximum magnetic energy in toroidal plasmas," Princeton Plasma Physics Laboratory Report PPPL-2822, December 1991 (submitted to *Phys. Fluids*).
 - 26 R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman Lectures on Physics*, (Addison-Wesley, Reading, 1964), Sec. 27-5, Vol. 2.

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Prof. I. Kawakami, Hiroshima Univ., JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Research Info. Ctr., National Inst. for Fusion Science, JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. S. Tanaka, Kyoto Univ., JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	Library, Kyoto Univ., JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Prof. R. Boucqué, Rijksuniversiteit Gent, BELGIUM	S. Mori, Technical Advisor, JAERI, JAPAN
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL	J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
Documents Office, Atomic Energy of Canada Ltd., CANADA	D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Prof. B.S. Uley, Univ. of Waikato, NEW ZEALAND
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Inst. of Physics, Chinese Acad. Sci. PEOPLE'S REP. OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Prof. T.W. Johnston, INRS-Energie, CANADA	Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Dr. C.R. James, Univ. of Alberta, CANADA	Dr. O. Petrus, AL I CUZA Univ., ROMANIA
Dr. P. Lukáč, Komenského Univerzita, CZECHO-SLOVAKIA	Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
The Librarian, Culham Laboratory, ENGLAND	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS	Dr. L. Stenflo, Univ. of UMEA, SWEDEN
P. Mähönen, Univ. of Helsinki, FINLAND	Library, Royal Inst. of Technology, SWEDEN
Prof. M.N. Bussac, Ecole Polytechnique, FRANCE	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
C. Moutat, Lab. de Physique des Milieux Ionisés, FRANCE	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Bibliothek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
Prof. E. Economou, Univ. of Crete, GREECE	Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
Ms. C. Rinni, Univ. of Ioannina, GREECE	Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
Dr. T. Mui, Academy Bibliographic Ser., HONG KONG	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA	Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Librarian, International Center for Theo. Physics, ITALY	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN	Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

END

**DATE
FILMED
416192**

7