577
2-25-82

I-1610

(1)

(40)

LB. 286
LBL-13116
UC-66a

Lawrence Berkeley Laboratory
UNIVERSITY OF CALIFORNIA

EARTH SCIENCES DIVISION

MASTER

ON A FUNCTION, $J(x,y)$, OCCURRING IN PROBLEMS OF
SOLUTE TRANSPORT WITH NON-EQUILIBRIUM INTERPHASE
MASS TRANSFER

Chalon L. Carnahan

July 1981

Prepared for the U.S. Department of Energy under Contract W-7405-ENG-48

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

LEGAL NOTICE

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Printed in the United States of America
Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161
Price Code: A03

LBL--13116

DE82 011098

ON A FUNCTION, $J(x,y)$, OCCURRING IN PROBLEMS
OF SOLUTE TRANSPORT WITH NON-EQUILIBRIUM
INTERPHASE MASS TRANSFER

Chalon L. Carnahan

Earth Sciences Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

July, 1981

DISCLAIMER

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Prepared for the Director, Office of Energy Research, Office of Basic Energy Sciences, Division of Engineering, Mathematics and Geosciences, of the U. S. Department of Energy under contract W-7405-ENG-48.

TABLE OF CONTENTS

List of Figures	ii
Abstract	iii
1. Introduction	1
2. Definitions and Properties of $J(x,y)$	2
3. Applications	4
4. Tabulations	9
5. Computational Methods	11
6. Summary and Conclusions	22
Acknowledgements	23
References	24
Figures	27

LIST OF FIGURES

- Figure 1. $J(x,y)$ on probability scale versus x on logarithmic scale with y as a parameter.
- Figure 2. $J(x,y)$ on probability scale versus y on logarithmic scale with x as a parameter.
- Figure 3. Curves of constant $J(x,y)$ as functions of arguments x and y .
- Figure 4. $J(x,x)$ versus x .

ON A FUNCTION, $J(x,y)$, OCCURRING IN PROBLEMS
OF SOLUTE TRANSPORT WITH NON-EQUILIBRIUM
INTERPHASE MASS TRANSFER

Chalon L. Carnahan
Earth Sciences Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

ABSTRACT

The function, $J(x,y)$, which has appeared frequently in analytical solutions of a variety of technical problems, is described and its applications briefly reviewed. Two detailed examples of applications are given. Tabulations of functions related to $J(x,y)$ are listed, and relationships of $J(x,y)$ to these functions are stated. Methods of computation of $J(x,y)$, suitable for use with digital computers, are described.

1. INTRODUCTION

Analytical solutions of the partial differential equations of mass or heat transport in porous media with time-dependent (non-equilibrium) transfer of mass or heat between fluid and solid phases may include certain irreducible integrals in which the integrands are, in general, products of exponential functions and Bessel functions. These integrals have appeared frequently in the literature on heat and mass transport, as well as in that describing other phenomena, and in recent years have been abbreviated by use of the functional designation $J(x,y)$. Aside from purely mathematical interest, knowledge of the properties of the function $J(x,y)$ is required for its numerical evaluation in a variety of technical applications. Here we briefly review properties and applications of the function $J(x,y)$, describe published (and unpublished) tabulations of related functions, and discuss methods for numerical evaluation of $J(x,y)$.

2. DEFINITIONS AND PROPERTIES OF $J(x,y)$

The function $J(x,y)$ can be defined (Goldstein, 1953) by:

$$J(x, y) = 1 - e^{-y} \int_0^x e^{-\xi} I_0(2\sqrt{y\xi}) d\xi, \quad x \geq 0, y \geq 0, \quad (1)$$

where $I_0(z)$ is the modified Bessel function of the first kind, of order zero, with argument z . The form (1) appears to have been used most frequently in technical applications. Other, equivalent forms have been derived by Anzelius (1926):

$$J(x, y) = 1 - e^{-y} \int_0^x e^{-\xi} J_0(2\sqrt{y\xi}) d\xi, \quad (2)$$

and by Terazawa (1922) and Goldstein (1932, 1953):

$$J(x, y) = 1 - x^{1/2} \int_0^\infty e^{-\xi} J_0(2\sqrt{y\xi}) J_1(2\sqrt{x\xi}) \xi^{-1/2} d\xi, \quad (3)$$

where $J_0(z)$ and $J_1(z)$ are the Bessel functions of the first kind, of orders zero and one respectively, with argument z . Vermeulen and Hiester (1952) appear to have been the first to use the symbol $J(x,y)$ for this function.

Mathematical properties of $J(x,y)$ have been summarized by Goldstein (1953) and, more extensively, by Luke (1962). Some of the more useful properties derivable from (1) are listed below, where subscripts x and y indicate partial differentiation.

$$J(x, y) + J(y, x) = 1 + e^{-x-y} I_0(2\sqrt{xy}), \quad (4)$$

$$J(x, x) = \frac{1}{2} + \frac{1}{2}e^{-2x} I_0(2x) \quad (5)$$

$$J_x(x, y) = -e^{-x-y} I_0(2\sqrt{xy})$$

$$J_y(x, y) = e^{-x-y} \left(\frac{x}{y}\right)^{1/2} I_1(2\sqrt{xy})$$

$$J(x, 0) = e^{-x}$$

$$J(0, y) = 1$$

$$\lim_{x \rightarrow \infty} J(x, y) = 0$$

$$\lim_{y \rightarrow \infty} J(x, y) = 1$$

$$\lim_{x \rightarrow \infty} J(x, x) = \frac{1}{2}$$

Figures 1-4 show the variations of $J(x, y)$ over limited ranges of the arguments x and y .

3. APPLICATIONS

The function $J(x,y)$ and related functions occur in the analytical solutions of a variety of technical problems. A sampling of such problems is given below with two detailed examples; the reader is referred to the original literature for detailed statements of other problems, including governing differential equations, initial conditions, and boundary conditions.

(1) heat exchange and heat transport:

Anzelius, 1926; Brinkley, 1947; Hougen and Marshall, 1948; Klinkenberg, 1948; Carslaw and Jaeger, 1959 (pp. 259-260, 393-396).

(2) hydrodynamics:

Terazawa, 1922 (p. 99); Goldstein, 1932 (pp. 66-72).

(3) probability and statistics:

Bose, 1947; Brinkley and Brinkley, 1947; Masters, 1955.

(4) fluid flow in pipes, electric current in cables:

Binnie and Miller, 1955.

(5) mass transport in porous media with interphase mass transfer:

(a) convective transport only:

Thomas, 1944, 1948, 1949; Walter, 1945; Hiester and Vermeulen, 1952; Vermeulen and Hiester, 1952; Goldstein, 1953; Opler and Hiester, 1954; Gardner and Brooks, 1957 .

(a) convective and dispersive transport:

Lapidus and Amundson, 1952; Ogata, 1961, 1964, 1969, 1970, 1976; Banks and Iqbal, 1964; Eldor and Dagan, 1972; Lindstrom and Narasimhan, 1973; Lindstrom, 1976; Cameron and Klute, 1977; deSmedt and Wierenga, 1979a, 1979b; Carnahan and Remer, 1981.

Two detailed examples of applications of the function $J(x,y)$ in problems

concerning mass transport with interphase mass transfer are set forth below. The first example considers purely convective transport with a nonlinear expression for the rate of mass transfer. The second example considers convective and dispersive transport with a linear rate expression.

Hiester and Vermeulen (1952) considered a fixed-bed ion exchange process in which an ionic solute, A, in the feed solution exchanges with another solute previously sorbed on the solid phase (resin) occupying a packed column. The material balance relation for solute A at a given cross section of the column is (with minor changes of notation):

$$-\left(\frac{\partial c}{\partial v}\right)_V = \rho \left(\frac{\partial q}{\partial V}\right)_V + f \left(\frac{\partial c}{\partial V}\right)_V \quad (6)$$

where c is the number of equivalents of A per unit volume of fluid phase, q is the number of equivalents of A per unit mass of dry resin, ρ is the bulk density of dry resin, f is the porosity of the column, v is the total column volume from inlet to cross section, and V is the volume of fluid fed to the column. The rate expression for ion exchange is:

$$\frac{dq}{d\tau} = k \left[c(q_\infty - q) - \frac{1}{K} q(c_0 - c) \right] \quad (7)$$

where τ is time, k is a rate constant, K is the selectivity coefficient for the ion-exchange reaction, q_∞ is the exchange capacity of the resin, and c_0 is the total solute concentration in the fluid phase. The problem is recast by use of the dimensionless variables, r , s , and t ,

where $r = K^{-1}$

$$s = kv\rho q_\infty R^{-1}$$

$$t = k(V-fv)c_0 R^{-1} = k(\tau-fvR^{-1})c_0$$

and R is the volumetric rate of flow of feed solution. With this notation (6) and (7) become, respectively,

$$- \left[\frac{\partial(c/c_o)}{\partial s} \right]_t = \left[\frac{\partial(q/q_\infty)}{\partial t} \right]_s$$

$$\left[\frac{d(q/q_\infty)}{dt} \right]_v = \frac{c}{c_o} \left(1 - \frac{q}{q_\infty} \right) - r \frac{q}{q_\infty} \left(1 - \frac{c}{c_o} \right)$$

with the boundary conditions:

$$c/c_o = 1 \text{ at } s = 0$$

$$q/q_\infty = 0 \text{ at } t = 0$$

The resulting solutions are

$$\frac{c}{c_o} = \frac{J(rs, t)}{J(rs, t) + e^{(r-1)(t-s)} [1 - J(s, rt)]} \quad (8)$$

and

$$\frac{q}{q_\infty} = \frac{1 - J(t, rs)}{J(rs, t) + e^{(r-1)(t-s)} [1 - J(s, rt)]} \quad (9)$$

where $J(x, y)$ is defined by (1).

Carnahan and Remer (1981) considered a three-dimensional porous medium with porosity ϵ supporting a flow of fluid with velocity v parallel to the z -axis and moving toward the positive direction. A solute is introduced at a constant rate, m_o mass units per unit time, at the origin of coordinates and is subjected to convective transport with hydrodynamic dispersion, the dispersion being characterized by coefficients D_L and D_T in directions parallel and perpendicular, respectively, to the direction of fluid flow. If c is the solute mass per

unit volume of fluid and q is the sorbed solute mass per unit volume of solid phase, the mass balance equation at a point (x,y,z) at time $t > 0$ is

$$\frac{\partial c}{\partial t} + \alpha \frac{\partial q}{\partial t} = D_T \left(\frac{\partial^2 c}{\partial x^2} + \frac{\partial^2 c}{\partial y^2} \right) + D_L \frac{\partial^2 c}{\partial z^2} - v \frac{\partial c}{\partial z} \quad (10)$$

where

$$\alpha = \frac{1-\epsilon}{\epsilon}$$

The rate expression for sorption is assumed to be linear and takes the form

$$\frac{\partial q}{\partial t} = k_1 c - k_2 q \quad (11)$$

where k_1 and k_2 are forward and backward rate constants. The linear rate expression used here corresponds to "Langmuir" sorption in which q is much smaller than the sorptive capacity of the solid. Initial conditions are:

$$c = 0, q = 0 \text{ for all } (x,y,z) \text{ at } t=0 \quad (12)$$

and the boundary conditions are

$$\lim_{\substack{c=0, t>0 \\ x,y,z \rightarrow \infty}} \quad (13)$$

$$\int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dz [\epsilon c + (1-\epsilon)q] = m_0 t, \quad t > 0 \quad (14)$$

The solutions to (10) and (11) satisfying (12) - (14) were found to be:

$$c = Q \int_0^t e^{-\frac{x^2+y^2}{4D_T\tau} - \frac{(z-v\tau)^2}{4D_L\tau}} J \left[\alpha k_1 \tau, k_2 (t-\tau) \right] \frac{d\tau}{\tau^{3/2}} \quad (15)$$

$$q = \frac{k_1}{k_2} Q \int_0^t e^{-\frac{x^2+y^2}{4D_T\tau} - \frac{(z-v\tau)^2}{4D_L\tau}} \left\{ 1 - J \left[k_2 (t-\tau), \alpha k_1 \tau \right] \right\} \frac{d\tau}{\tau^{3/2}} \quad (16)$$

where

$$Q = \frac{m_0}{8\pi^{3/2} D_T D_L^{1/2} \epsilon} .$$

Although solutions (8) and (9) could be evaluated by use of tables of the function $J(x,y)$, it is evident that numerical evaluation of the irreducible integrals in (15) and (16) requires an efficient computational scheme for $J(x,y)$.

4. TABULATIONS

Although the function $J(x,y)$ as defined by (1)-(3) does not appear to have been tabulated in the published literature, tabulations of several related functions have been published or reported.

- (1) Bose (1947) has tabulated the function $L(P,p,\lambda)$ where

$$L(P,p,\lambda) = \int_0^P \eta^{p/2} \lambda^{1-p/2} e^{-(\eta^2 + \lambda^2)/2} I_{(p-2)/2}(\lambda\eta) d\eta$$

For the case $p = 2$,

$$J(x,y) = 1 - L(\sqrt{2x}, 2, \sqrt{2y})$$

- (2) Brinkley and Brinkley (1947) reported an unpublished table of the probability, $p(r,R)$, that the point of impact of a missile aimed at the origin of coordinates will lie within a circle of radius r whose center is distance R from the origin; $p(r,R)$ is sometimes called the offset circular probability function, and was defined by

$$p(r,R) = e^{-R^2} \int_0^{r^2} e^{-t} I_0(2R\sqrt{t}) dt$$

Then,

$$J(x,y) = 1 - p(\sqrt{x}, \sqrt{y})$$

- (3) The Rand Corporation (1951) has tabulated the offset circular probability function $q(R,r)$, where

$$q(R,r) = \int_R^\infty t e^{-(t^2 + r^2)/2} I_0(rt) dt \quad (17)$$

Then,

$$J(x,y) = q(\sqrt{2x}, \sqrt{2y})$$

- (4) Wilson (1951) and the Admiralty Research Laboratory (1953) have published tabulations of the Function $F(\beta, \rho)$ where

$$F(\beta, \rho) = \frac{2e^{-\rho^2}}{\sqrt{\pi}\beta^2} \int_0^\beta e^{-\eta^2} I_0(2\rho\eta) \eta d\eta$$

Then,

$$J(x,y) = 1 - \sqrt{\pi x} F(\sqrt{x}, \sqrt{y})$$

- (5) Brinkley, Edwards, and Smith (1952) reported an unpublished table of the function $\phi_0(x,y)$ applicable to exchange of heat between a fluid and a porous solid, where

$$\phi_0(x,y) = e^x \int_0^x e^{-t} I_0(2\sqrt{yt}) dt$$

Then,

$$J(x,y) = 1 - e^{-x-y} \phi_0(x,y)$$

- (6) Masters (1955) tabulated the offset circular probability function, $P(z/\sigma, R/\sigma)$, and indicated the utility of this function in problems of nuclear particle scattering and heat conduction. $P(z/\sigma, R/\sigma)$ was defined by:

$$P(z/\sigma, R/\sigma) = \frac{e^{-R^2/2\sigma^2}}{\sigma^2} \int_0^z e^{-r^2/2\sigma^2} I_0(rR/\sigma^2) r dr$$

and is related to $J(x,y)$ by:

$$J(x,y) = 1 - P(\sqrt{2x}, \sqrt{2y})$$

5. COMPUTATIONAL METHODS

The tabulations of functions related to the function $J(x,y)$ are not suitable for repeated, numerical evaluations of $J(x,y)$ required for many practical applications. However, several computational methods are available.

Walter (1945) and Klinkenberg (1948) have given an analytical approximation to $J(x,y)$, based on the error function and applicable for large values of the arguments (x,y) . Hastings and Wong (1953) have provided twenty-two analytical approximations to the function $q(R,r)$ defined by (17); their approximations are valid for a variety of ranges of argument values and exhibit a variety of accuracies.

An asymptotic expansion in a series of Bessel functions was attributed to L. Onsager and was described by Thomas (1944, 1948). Asymptotic expansions which are more easily implemented numerically have been derived by Goldstein (1953).

The available analytical approximations are not accurate enough for evaluation of expressions such as (15) and (16), where $J(x,y)$ appears as part of the integrand in integrals which must be evaluated by numerical methods. The remainder of this section describes methods which have been implemented at the Lawrence Berkeley Laboratory for computation of $J(x,y)$ as part of a study of mass transport in porous media (Carnahan and Remer, 1981).

One method uses an iterative Simpson's rule routine to evaluate the integral $S(x,y)$, where

$$S(x,y) = \int_0^1 e^{-x\xi} I_0(2\sqrt{xy\xi}) d\xi$$

Approximations to $S(x,y)$ are made on successively smaller subintervals of $\Delta \xi$

until the absolute fractional error between successive approximations is less than an assigned value. The modified Bessel function is computed using polynomial approximations given by Olver (1965). If argument $x \leq 10$, or if $x > 10$ while $x \leq y$, $J(x,y)$ is formed from:

$$J(x,y) = 1 - xe^{-y}S(x,y)$$

If $x > 10$ while $x > y$, the routine computes $S(y,x)$, and $J(x,y)$ is formed from:

$$J(x,y) = ye^{-x}S(y,x) + e^{-x-y}I_0(2\sqrt{xy})$$

This method produces results which have relative errors of about 1×10^{-6} .

Another method uses infinite series expansions of $J(x,y)$. If the modified Bessel function in (1) is expanded in the series (Olver, 1965)

$$I_0(z) = \sum_{k=0}^{\infty} \frac{(z^{2/4})^k}{(k!)^2} \quad (18)$$

and the integration is performed term by term, the following result is obtained:

$$J(x,y) = e^{-x-y} \sum_{n=0}^{\infty} \frac{y^n}{n!} \sum_{m=0}^n \frac{x^m}{m!} \quad (19)$$

This or equivalent results have been stated by Thomas (1944, 1948), Walter (1945), and deSmedt and Wierenga (1979a, 1979b). An alternative form, stated by deSmedt and Wierenga (1979a, 1979b), can be derived by using (19) to write $J(y,x)$ and finding $J(x,y)$ by use of (4) and (18). The result is:

$$J(x,y) = 1 - e^{-x-y} \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{m=0}^{n-1} \frac{y^m}{m!} \quad (20)$$

In principle, (19) and (20) can be used to compute $J(x,y)$ to any desired degree of accuracy. We have found that, for a given degree of accuracy, (19) requires fewer computational steps if $x > y$, while (20) is faster in this sense if $y > x$. In practice, large values of x and y require a large number of computations because of the slow rate of convergence of the double sum in the two expressions. Also, the numerical value of the double sum may exceed the range of the computing device for very large argument values. For example, assume that the computation is carried to a number of terms so that the contribution of the next term in the summation over n , relative to the double sum computed thus far, is less than 1×10^{-6} . Then if both x and y equal 116, the double sum equals about 10^{100} , and if both x and y equal 372, the double sum equals about 10^{322} , which is the upper limit of the range of real constants allowed by the CDC-7600 computer in use at this laboratory. In both examples, the value of $J(x,x)$ is about 0.5.

The problems of slow convergence and out-of-range computations encountered in use of (19) and (20) can be minimized by application of available asymptotic expansions of the function $J(x,y)$. We have adapted several expansions given by Goldstein (1953) to computations of $J(x,y)$ with large values of the arguments, our principal modification being the substitution of an asymptotic expansion of the modified Bessel function of the first kind of order zero wherever the Bessel function occurs in Goldstein's equations. We have also derived recursion relations for the coefficients occurring in the expansions.

The asymptotic expansion of the modified Bessel function of the first kind of order zero with argument z is (Goldstein, 1953):

$$I_0(z) \sim \frac{e^z}{\sqrt{2\pi z}} \sum_{m=0}^{\infty} \frac{A_m}{(2z)^m} \quad (21)$$

where $A_0 = 1$, $A_1 = \frac{1}{4}$, ..., $A_m = \frac{1^2 \cdot 3^2 \dots (2m-1)^2}{4^m m!}$

The recurrence relation for the coefficient A_m is

$$A_m = \frac{(2m-1)^2}{4m} A_{m-1} \quad (22)$$

The expansion (21) can be used in (5) to derive an asymptotic expansion of $J(x, x)$:

$$J(x, x) \sim \frac{1}{2} + \frac{1}{4\sqrt{\pi x}} \sum_{m=0}^{\infty} \frac{A_m}{(4x)^m} \quad (23)$$

The following computational scheme provides $J^{(n)}(x, x)$, the approximation of $J(x, x)$ including terms through $m = n$:

$$Q_0(x) = \frac{1}{4\sqrt{\pi x}}, \dots, Q_m(x) = \frac{(2m-1)^2}{16mx} Q_{m-1}(x)$$

$$J^{(n)}(x, x) = \frac{1}{2} + \sum_{m=0}^n Q_m(x) = J^{(n-1)}(x, x) + Q_n(x)$$

Define the relative difference, R_n , between successive approximations $J^{(n)}(x, y)$ by

$$R_n = \left| \frac{J^{(n)}(x, y) - J^{(n-1)}(x, y)}{J^{(n-1)}(x, y)} \right|$$

In this case,

$$R_n = \left| \frac{Q_n(x)}{J^{(n-1)}(x, x)} \right|$$

The computational scheme for $J(x,x)$ gives $R_n = 5 \times 10^{-8}$ for $x = 100$ and $n = 2$, and $R_n = 8 \times 10^{-7}$ for $x = 10$ and $n = 3$. In contrast, use of (19) requires 152 terms in the outer summation to achieve $R_n = 8 \times 10^{-7}$ with $x = y = 100$. It is noted that (23) begins to diverge ($Q_m > Q_{m-1}$) at a value of m approximately equal to $4x + 1$, and if a desired R_n value has not been achieved by this step, it never will be achieved. For this reason, use of (23) is not advised for argument values less than $x = 4$ if $R_n < 10^{-7}$.

If the arguments of $J(x,y)$ are unequal, the asymptotic expansions given in the following paragraphs can be used. Using Goldstein's notation, the following variables are defined:

$$\xi = 2\sqrt{xy}, \quad \eta = \sqrt{\frac{y}{x}}, \quad z = (\sqrt{y} - \sqrt{x})^2$$

If ξ is large while $\xi/z < 1/2$, we have

$$J(x,y) \sim \frac{e^{-z}}{2\sqrt{2\pi\xi}} S_\infty(x,y) \quad \text{for } \eta < 1$$

$$J(x,y) \sim 1 + \frac{e^{-z}}{2\sqrt{2\pi\xi}} S_\infty(x,y) \quad \text{for } \eta > 1$$

where

$$S_\infty(x,y) = \sum_{m=0}^{\infty} \frac{A_m}{(2\xi)^m} \left[1 + \frac{1+\eta}{1-\eta} \beta_m(\xi,z) \right]$$

A_m is defined by (22), and $\beta_m(\xi,z)$ is defined by

$$\beta_0(\xi,z) = 1, \quad \beta_m(\xi,z) = 1 + \sum_{k=1}^m B_{m,k}(\xi,z) \quad (24)$$

$$B_{m,k}(\xi, z) = (-)^k \frac{2m(2m-2)\dots(2m-2k+2)}{(2m-1)(2m-3)\dots(2m-2k+1)} \left(\frac{2\xi}{z}\right)^k$$

The following computational scheme provides $J^{(n)}(x, y)$:

$$B_{m,1}(\xi, z) = -\frac{2m}{2m-1}\left(\frac{2\xi}{z}\right), \quad B_{m,k}(\xi, z) = -\frac{2m+2-2k}{2m+1-2k}\left(\frac{2\xi}{z}\right) B_{m,k-1}(\xi, z)$$

Compute $\beta_m(\xi, z)$ according to (24).

$$C_0 = 1 + \frac{1+\eta}{1-\eta}, \quad C_m = 1 + \frac{1+\eta}{1-\eta} \beta_m(\xi, z)$$

$$D_0 = C_0, \quad D_m = \frac{(2m-1)^2}{8m\xi} \cdot \frac{C_m}{C_{m-1}} \cdot D_{m-1}$$

$$S_n = \sum_{m=0}^n D_m = S_{n-1} + D_n$$

$$J^{(n)}(x, y) = \frac{e^{-z}}{2\sqrt{2\pi\xi}} S_n \quad \text{for } \eta < 1$$

$$J^{(n)}(x, y) = 1 + \frac{e^{-z}}{2\sqrt{2\pi\xi}} S_n \quad \text{for } \eta > 1$$

$$R_n = \left| \frac{D_n}{S_{n-1}} \right| \quad \text{for } \eta < 1$$

$$R_n = \frac{e^{-z}}{2\sqrt{2\pi\xi}} \left| \frac{D_n}{J^{(n-1)}(x, y)} \right| \quad \text{for } \eta > 1$$

Using arguments $x = 100$, $y = 0.1$, this scheme produces $J^{(10)}(x, y) = 3.13601 \times 10^{-42}$ with $R_{10} = 0.99 \times 10^{-6}$.

If ξ is large while $\xi/z > 1$, we have

$$J(x, y) \sim F(\xi, z) + \frac{1}{2} e^{-z} \sum_{m=1}^{\infty} [G_m(\xi) + H_m(\xi, z)]$$

where

$$F(\xi, z) = \frac{e^{-z}}{2\sqrt{2\pi\xi}} + \frac{1}{2} (1 \mp \operatorname{erf} \sqrt{z}), \operatorname{erf} \sqrt{z} = \frac{2}{\sqrt{\pi}} \int_0^{\sqrt{z}} e^{-u^2} du \quad (25)$$

$$G_m(\xi) = \frac{A_m}{\sqrt{2\pi\xi} (2\xi)^m}$$

$$H_m(\xi, z) = \pm \frac{\alpha_m T_m(z)}{(2\xi+z)^m} \sqrt{\frac{z}{\pi}} \quad (26)$$

and A_m is defined in (22).

In (25) and (26) the upper signs are used if $\eta < 1$, and the lower signs are used if $\eta > 1$. In (26), note that $2\xi + z = (\sqrt{x} + \sqrt{y})^2$. α_m and $T_m(z)$ are defined by:

$$\alpha_m = \frac{1 \cdot 3 \cdot 5 \dots (2m-1)}{2 \cdot 4 \cdot 6 \dots (2m)}$$

$$T_1(z) = 1, T_2(z) = z + \frac{1}{2}, T_3(z) = z^2 + z + \frac{3}{4},$$

$$T_m(z) = z^{m-1} + \frac{1}{2} (m-1) z^{m-2} + \frac{1}{2} \cdot \frac{3}{2} \cdot \frac{(m-1)(m-2)}{2!} z^{m-3} + \dots$$

$$+ \frac{1}{2} \cdot \frac{3}{2} \dots (k - \frac{1}{2}) \cdot \frac{(m-1)(m-2) \dots (m-k)}{k!} z^{m-k-1} + \dots + \frac{1}{2} \cdot \frac{3}{2} \dots (m - \frac{3}{2})$$

The following computational scheme provides $J^{(n)}(x, y)$:

$$G_1(\xi) = (8\xi\sqrt{2\pi\xi})^{-1}, \quad G_m(\xi) = \frac{(2m-1)^2}{8m\xi} G_{m-1}(\xi).$$

$$T_m(z) = \sum_{k=0}^{m-1} C_{m,k} z^{m-k-1}$$

$$\text{where } C_{m,0} = 1, \quad C_{m,k} = \frac{(k-\frac{1}{2})(m-k)}{k} \cdot C_{m,k-1}$$

$$H_1(\xi, z) = \pm \frac{\sqrt{z}}{2(2\xi+z)\sqrt{\pi}}, \quad H_m(\xi, z) = \frac{2m-1}{2m} \cdot \frac{T_m(z)}{T_{m-1}(z)} (2\xi+z)^{-1} H_{m-1}(\xi, z)$$

$$S_n = \sum_{m=1}^n \left[G_m(\xi) + H_m(\xi, z) \right] = S_{n-1} + G_n(\xi) + H_n(\xi, z)$$

$$J^{(n)}(x, y) = F(\xi, z) + \frac{1}{2} e^{-z} S_n$$

$$R_n = \frac{e^{-z}}{2} \left| \frac{G_n(\xi) + H_n(\xi, z)}{J^{(n-1)}(x, y)} \right|$$

Using arguments $x = 100$, $y = 10$, this scheme produces $J^{(10)}(x, y) = 3.61548 \times 10^{-22}$ with $R_{10} = 2.3 \times 10^{-7}$.

The following asymptotic expansion is useful when ξ is large and z is neither very large nor very small relative to ξ , in particular when $\xi \leq z \leq 2\xi$:

$$J(x, y) \sim M(\xi, z) + \frac{e^{-z}}{2\sqrt{2\pi\xi}} \sum_{m=1}^{\infty} \left[L_m(\xi) + K_m(\xi, z) \right] \text{ for } \eta < 1$$

$$J(x, y) \sim 1 + M(\xi, z) + \frac{e^{-z}}{2\sqrt{2\pi\xi}} \sum_{m=1}^{\infty} \left[L_m(\xi) + K_m(\xi, z) \right] \text{ for } \eta > 1$$

where

$$L_m(\xi) = \frac{A_m}{(2\xi)^m}$$

$$M(\xi, z) = \frac{e^{-z}}{2\sqrt{2\pi\xi}} \pm \frac{\sqrt{x} + \sqrt{y}}{2\sqrt{2\xi}} \operatorname{erfc}\sqrt{z} \quad , \quad \operatorname{erfc}\sqrt{z} = \frac{2}{\sqrt{\pi}} \int_{\sqrt{z}}^{\infty} e^{-u^2} du \quad (27)$$

$$K_m(\xi, z) = \frac{(x-y)\alpha_m S_m(z)}{(2\xi)^m}$$

and A_m and α_m remain as defined previously.

In (27) the plus sign is to be used if $n < 1$, and the minus sign if $n > 1$. Goldstein (1953) defines the function $S_m(z)$ by

$$S_m(z) = \frac{\Gamma(m - \frac{1}{2})}{\Gamma(\frac{1}{2})} - \frac{\Gamma(m - \frac{3}{2})}{\Gamma(\frac{1}{2})} z + \frac{\Gamma(m - \frac{5}{2})}{\Gamma(\frac{1}{2})} z^2 - \dots$$

$$+ (-)^m \frac{\Gamma(\frac{3}{2})}{\Gamma(\frac{1}{2})} z^{m-2} + (-)^{m-1} z^{m-1} + (-)^m z^{m-1} \sqrt{\pi z} e^z \operatorname{erfc}\sqrt{z}$$

$$S_1(z) = 1 - \sqrt{\pi z} e^z \operatorname{erfc}\sqrt{z}$$

and gives a recurrence relation which we write as

$$S_m(z) + z S_{m-1}(z) = \frac{\Gamma(m - \frac{1}{2})}{\Gamma(\frac{1}{2})} \quad (m > 1)$$

$\Gamma(\zeta)$ is the gamma function with argument ζ and has the property (Davis, 1965):

$$\frac{\Gamma(m - \frac{1}{2})}{\Gamma(\frac{1}{2})} = \frac{1 \cdot 3 \cdot 5 \cdot 7 \dots (2m-3)}{2^{m-1}} \quad (m > 1)$$

The following computational scheme provides $J^{(n)}(x,y)$:

$$L_1(\xi) = \frac{1}{8\xi}, \quad L_m(\xi) = \frac{(2m-1)^2}{8m\xi} L_{m-1}(\xi)$$

$$S_1(z) = 1 - \sqrt{\pi z} e^z \operatorname{erfc}\sqrt{z}$$

$$S_m(z) = N_m - z S_{m-1}(z)$$

$$\text{where } N_1 = 1, \quad N_m = \frac{2m-3}{2} N_{m-1}$$

$$K_1(\xi, z) = \frac{(x-y)S_1(z)}{4\xi}$$

$$K_m(\xi, z) = \frac{2m-1}{4m\xi} \cdot \frac{S_m(z)}{S_{m-1}(z)} K_{m-1}(\xi, z)$$

$$J^{(n)}(x,y) = M(\xi, z) + \frac{e^{-z}}{2\sqrt{2\pi\xi}} \sum_{m=1}^n \left[L_m(\xi) + K_m(\xi, z) \right], \quad \eta < 1$$

$$J^{(n)}(x,y) = 1 + M(\xi, z) + \frac{e^{-z}}{2\sqrt{2\pi\xi}} \sum_{m=1}^n \left[L_m(\xi) + K_m(\xi, z) \right], \quad \eta > 1$$

$$R_n = \frac{e^{-z}}{2\sqrt{2\pi\xi}} \left| \frac{L_n(\xi) + K_n(\xi, z)}{J^{(n-1)}(x,y)} \right|$$

Using arguments $x = 100$, $y = 4$, this scheme produces $J^{(5)}(x,y) = 1.26273 \times 10^{-29}$ with $R_5 = 8.8 \times 10^{-8}$.

The computational routines for the asymptotic expansions given here can be performed on a desk-top electronic calculator, provided certain functions such as $\operatorname{erf}\sqrt{z}$, $\operatorname{erfc}\sqrt{z}$, and $e^{-z}/(2\sqrt{2\pi\xi})$ are calculated or obtained from tables and stored beforehand. For implementation with a digital computer, software routines are usually available for calculation of the error function or its

complement; alternatively, rational approximations and asymptotic expansions are available and are easily coded (see, for example, Gautschi, 1965).

6. SUMMARY AND CONCLUSIONS

The function $J(x,y)$ is defined by an irreducible integral and appears both by itself and as part of the integrands of other irreducible integrals in a number of applications. Available methods for numerical computation of $J(x,y)$ over a wide range of argument values have been examined. Numerical integration of the defining integral, (1), is limited in accuracy by the accuracy of existing rational approximations of the Bessel function in the integrand. Infinite series expansions of $J(x,y)$ are capable, in principle, of greater accuracy, but for large values of the arguments they exhibit slow convergence and can produce numerical values for partial sums which are beyond the range of computing devices. These problems can be overcome by use of asymptotic expansions. Three expansions are applicable individually to the cases $\xi z < 1/2$, $1/2 \leq \xi/z \leq 1$, and $\xi/z > 1$ (ξ and z being the functions of x and y defined previously), when arguments x and y are unequal. A fourth expansion applies to the case of large, equal arguments. The expansions can be programmed easily on electronic calculators and digital computers by use of the recurrence relations presented here.

ACKNOWLEDGEMENTS

Janet Remer provided valuable assistance in computer programming and in reviewing this paper. This work was supported by the Office of Energy Research; Office of Basic Energy Sciences, Division of Engineering, Mathematics and Geosciences, of the U.S. Department of Energy under Contract W-7405-ENG-48.

REFERENCES

- Admiralty Research Laboratory, Table of $F(\beta, \rho)$, Teddington, Middlesex, England, 1953.
- Anzelius, A., Über Erwärmung vermittelt durchströmender Medien, Z. angew. Math. Mech., 6, 291-294, 1926.
- Banks, Robert B., and Iqbal Ali, Dispersion and adsorption in porous media flow, J. Hydraulics Div. ASCE, 90 (HY5), 13-31, 1964.
- Binnie, A.M., and J.C.P. Miller, Tables of two functions required in certain attenuation problems, Quart. J. Mech. Appl. Math., 8, 468-479, 1955.
- Bose, P.K., On recursion formulae, tables and Bessel function populations associated with the distribution of classical D^2 -statistic, Sankhya, 8, 235-248, 1947.
- Brinkley, Stuart R., Jr., Heat transfer between a fluid and a porous solid generating heat, J. Appl. Phys., 18, 582-585, 1947.
- Brinkley, S.R., Jr., and Ruth F. Brinkley, Unpublished mathematical tables, note 52(k), Math. Tables Aids Comp., 2, 221, 1947.
- Brinkley, S.R., Jr., H.E. Edwards, and R.W. Smith, Jr., Unpublished mathematical tables, note 141(I), Math. Tables Aids Comp., 6, 40, 1952.
- Cameron, D.R., and A. Klute, Convective-dispersive solute transport with a combined equilibrium and kinetic adsorption model, Water Resour. Res., 13, 183-188, 1977.
- Carnahan, Chalon L., and Janet S. Remer, Non-equilibrium and equilibrium sorption with a linear sorption isotherm during mass transport through an infinite, porous medium: some analytical solutions, Lawrence Berkeley Laboratory, Report LBL-13005, Berkeley, California, 1981.
- Carslaw, H.S., and J.C. Jaeger, Conduction of Heat in Solids, Second Edition, Oxford University Press, London, 1959.
- Davis, Philip J., Gamma function and related functions, in National Bureau of Standards Applied Mathematics Series, No. 55, Handbook of Mathematical Functions, edited by M. Abramowitz and I. Stegun, U.S. Govt. Printing Office, Washington, D.C., 1965.
- Eldor, M., and G. Dagan, Solutions of hydrodynamic dispersion in porous media, Water Resour. Res., 8, 1316-1331, 1972.
- Gardner, W.R., and R.H. Brooks, A descriptive theory of leaching, Soil Sci., 83, 295-304, 1957.
- Gautschi, Walter, Error function and Fresnel integrals, in National Bureau of

Standards Applied Mathematics Series, No. 55, Handbook of Mathematical Functions, edited by M. Abramowitz and I. Stegun, U.S. Govt. Printing Office, Washington, D.C., 1965.

Goldstein, S., Some two-dimensional diffusion problems with circular symmetry, Proc. London Math. Soc., Ser.2, 34, 51-88, 1932.

-----, On the mathematics of exchange processes in fixed columns. I. Mathematical solutions and asymptotic expansions, Proc. Roy. Soc.(London), A219, 151-171, 1953.

Hastings, Cecil, Jr., and James P. Wong, Jr., Note 153. Analytical approximations, Math. Tables Aids Comp., 7, 212-213, 1953.

Hiester, N.K., and T. Vermeulen, Saturation performance of ion-exchange and adsorption columns, Chem. Eng. Prog., 48, 505-516, 1952.

Hougen, O.A., and W.R. Marshall, Adsorption from a fluid stream flowing through a stationary granular bed, Chem. Eng. Prog., 43, 197-208, 1947.

Klinkenberg, Adriaan, Numerical evaluation of equations describing transient heat and mass transfer in packed solids, Ind. Eng. Chem., 40, 1992-1994, 1948.

Lapidus, L. and N.R. Amundson, Mathematics of adsorption in beds. VI. The effect of longitudinal diffusion in ion exchange and chromatographic columns, J. Phys. Chem., 56, 984-988, 1952.

Lindstrom, F.T., Pulsed dispersion of trace chemical concentrations in a saturated sorbing porous medium, Water Resour. Res., 12, 229-238, 1976.

Lindstrom, F.T., and M.N.L. Narasimhan, Mathematical theory of a kinetic model for dispersion of previously distributed chemicals in a sorbing porous medium, SIAM J. Appl. Math., 24, 496-510, 1973.

Luke, Yudell L., Integrals of Bessel's Functions, McGraw-Hill Book Co., Inc., New York, 1962.

Masters, J.I., Some applications in physics of the P function, J. Chem. Phys., 23, 1865-1874, 1955.

Ogata, Akio, Transverse diffusion in saturated isotropic granular media, U.S. Geological Survey, Professional Paper 411-B, U.S. Govt. Printing Office, Washington, D. C., 1961.

-----, Mathematics of dispersion with linear adsorption isotherm, U.S. Geological Survey, Professional Paper 411-H, U.S. Govt. Printing Office, Washington, D.C. 1964.

-----, Two-dimensional dispersion in a granular-medium disk source emitting at a constant rate, U.S. Geological Survey, Professional Paper 650-D, U.S. Govt. Printing Office, Washington, D.C., 1969.

- , Theory of dispersion in a granular medium, U.S. Geological Survey, Professional Paper 411-I, U.S. Govt. Printing Office, Washington, D.C., 1970.
- , Two-dimensional steady-state dispersion in a saturated porous medium, J. Res. U.S. Geol. Survey, 4, 277-284, 1976.
- Olver, F.W.J., Bessel functions of integer order, in National Bureau of Standards Applied Mathematics Series, No. 55, Handbook of Mathematical Functions, edited by M. Abramowitz and I. Stegun, U.S. Govt. Printing Office, Washington, D.C., 1965.
- Opler, Ascher, and Nevin K. Hiester, Tables for predicting the performance of fixed bed ion exchange and similar mass transfer processes, Stanford Research Institute, Stanford, California, 1954.
- Rand Corporation, Table of offset circle probabilities, Rand Corporation, Santa Monica, California, 1951.
- Smedt, F. de, and P.J. Wierenga, Mass transfer in porous media with immobile water, J. Hydrology, 41, 59-67, 1979a.
- , A generalized solution for solute flow in soils with mobile and immobile water, Water Resour. Res., 15, 1137-1141, 1979b.
- Terazawa, K., On the decay of vortical motion in a viscous fluid, Tokyo Imperial University, Report of the Aeronautical Research Institute, 1(4), 87-134, 1922.
- Thomas, Henry C., Heterogeneous ion exchange in a flowing system, J. Am. Chem. Soc., 66, 1664-1666, 1944.
- , Chromatography: a problem in kinetics, Ann. New York Acad. Sci. 49, 161-182, 1948.
- , The kinetics of fixed-bed ion exchange, in Ion Exchange Theory and Application, edited by F.C. Nachod, Academic Press, New York, 1949.
- Vermeulen, T., and N.K. Hiester, Ion-exchange chromatography of trace components, Ind. Eng. Chem., 44, 636-651, 1952.
- Walter, John E., Rate-dependent chromatographic adsorption, J. Chem. Phys., 13, 332-336, 1945.
- Wilson, E.M., Solutions of the equation $(y'')^2 = yy'$ and two other equations, Admiralty Research Laboratory, Teddington, Middlesex, England, 1951.

Figure 1. $J(x,y)$ on probability scale versus x on logarithmic scale with y as a parameter. XBL 819-11611

XBL 819-11612

Figure 2. $J(x,y)$ on probability scale versus y on logarithmic scale with x as a parameter.

XBL 819-11613

Figure 3. Curves of constant $J(x,y)$ as functions of arguments x and y .

Figure 4. $J(x, x)$ versus x .

XBL 819-11614

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720