PPPL-2034

UC20-G

PPPL-2034

Dr. 1747-7

I-11079

166
9-9-83 QS (1)

PPPL--2034

DE83 017720

STABILITY OF $n = 1$ KINK MODES IN BEAM-SHAPED TOKAMAKS

By

J. Manickam, R.C. Grimm, and M. Okabayashi

AUGUST 1983

PLASMA
PHYSICS
LABORATORY

PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY

MASTER

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-OR0-2073

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

STABILITY OF $n = 1$ KINK MODES IN BEAN-SHAPED TOKAMAKS

J. Manickam, R. C. Grimm, and M. Okabayashi

Plasma Physics Laboratory,

Princeton University

Princeton, New Jersey, 08544

ABSTRACT

Numerical studies show that by indenting the small-major-radius side of conventional finite-aspect-ratio tokamaks, significant improvements to the stability of pressure-driven ideal MHD modes can be achieved. The internal $n = 1$ kink mode can be stabilized completely with quite modest indentation. Kink-ballooning mode stability is also improved, and, in the presence of a nearby conducting wall, accessibility to a second stable region at high plasma β is possible.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

229

Detailed theoretical studies have established that the ideal MHD stability properties of axisymmetric tokamak equilibria can depend strongly on the shape of the poloidal cross section of the plasma. Small aspect ratio, modest elongation, and triangularity are all favorable in the sense that the critical value of $\beta \equiv 2\langle p \rangle / \langle B^2 \rangle$ for the onset of internal, pressure-driven ballooning modes is larger than for the comparable circular cross-sectional shape.¹ The stabilizing effect of triangularity can be enhanced by indenting the small major radius side of the cross section, forming a "bean-shaped" plasma.^{2,3} In fact, with certain plasma pressure and safety factor profiles, it has been shown recently that for sufficiently large indentations complete ballooning stability can be achieved, providing an accessible path to the second stability region at smaller indentations.⁴

While stability in the short wavelength ballooning limit can usually be extrapolated to longer wavelength modes, the effects of which one would expect to be most directly observable experimentally, it is necessary to carry out a separate study for the low toroidal mode numbers (such as $n = 1$). This is the subject of this paper. For the internal $n = 1$ kink, we find that even modest indentation can completely stabilize the mode, thus connecting the favorable effect of finite aspect ratio at low β ⁵ to the stabilization associated with the second stable region at very high β .⁶ For the external kinks we find that in the presence of a nearby conducting shell the stability properties with large indentation are quantitatively similar to previous parameter studies at low β , but that, at larger β , a second stable region to external kinks appears in a similar way to the high- β ballooning results.

Equilibrium solutions employed for the study were prepared following now standard procedures. A flux-coordinate, Grad-Shafranov equilibrium code was used with pressure and safety-factor profiles given as

$$p(\psi) = p_0 (1 - \tilde{\psi}^\alpha)^\beta ; \quad q(\psi) = \sum_{i=1}^3 q_i \tilde{\psi}^i , \quad (1)$$

with $\tilde{\psi} = (\psi - \psi_a) / (\psi_\ell - \psi_a)$, (ψ_a , ψ_ℓ being the flux measured at the magnetic axis and plasma surface, respectively), and p_0 , α, β , q_i parameters chosen to fix β and the required profiles. In the cases reported here the q_i were fixed so that $q(0) = 1.03$, $q(1) = 4.2$, $q'(0) = 0.84$, and $q'(1) = 9.0$. The equilibria generated were obtained as fixed boundary solutions with the plasma surface determined from the formulae⁴

$$X(\theta) = \tilde{X} + \rho \cos \theta ; \quad Z(\theta) = E \rho \sin \theta , \quad (2)$$

with $\rho = A + B \cos \theta$, $t = C \sin \theta$, and $0 < \theta < 2\pi$. A typical example, showing the various parameters of interest, is given in Fig. 1. In the study reported here $R/a = 4$, $b/a = 1.386$ and the indentation, $i \equiv d/2a$, is varied by changing the angle C in Eq. (2). Sequences of flux conserved equilibria at different values of the indentation were then obtained by increasing the parameter, p_0 , holding all other profile parameters fixed. For the external kink studies, this gives a matrix of equilibria in the $\langle \beta \rangle - i$ plane, for each of which the stability must be examined. For the $n = 1$ internal kink we are interested in the stability properties for equilibria with $0.5 < q(0) \leq 1$. To study this, each of the equilibria described above was scaled by varying the strength of the external toroidal field.⁷ This procedure generates additional sequences of equilibria at constant $\beta_p \equiv 2\langle p \rangle / I_\phi^2$. At fixed indentation, the stability can then be determined in the $\epsilon \beta_p - \text{vs} - q(0)$ plane.

The stability calculations were carried out with PEST 2.⁸ Typically, 30-50 Fourier harmonics in PEST coordinates were kept with 100 radial finite elements. Convergence studies were performed to ensure that this resolution

was adequate.

Stability boundaries for the $n = 1$ internal kink mode are presented in Figs. 2 and 3 for two different choices of the pressure profile. In Fig. 2 $\alpha = \beta = 2$ so that the profile is flat near the magnetic axis. This provides some stabilization when the $q = 1$ surface lies close to the magnetic axis, even without indentation. The $n = 1$ mode is unstable inside the curves, which shrink rapidly with indentation. In the inset we have plotted the maximum height and width of the unstable regions as a function of indentation. The critical indentation is seen to be approximately 0.1. Detailed calculations with $i > 0.11$ show complete stability to the internal kink even with $q(0)$ as low as 0.5.

In Fig. 3 we have taken $\alpha = 1$, $\beta = 2$, so that $p'(0) \neq 0$. For given indentation the instability region is larger than for the flatter pressure profile, but indentation is seen to have a similar, strongly stabilizing effect. Extrapolation of these results gives a critical indentation of approximately 0.3, which corresponds closely to the critical indentation for stability to internal ballooning modes for these configurations with $q(0) = 1.03$.⁴

It is useful to examine the nature of the $n = 1$ eigenmodes as a function of $\epsilon\beta_p$. At low $\epsilon\beta_p$ the structure, illustrated in Fig. 4, is consistent with the conventional picture of a current driven perturbation with driving forces localized around the $q = 1$ surface. At larger β_p , however, the mode becomes primarily pressure driven and shows a radial structure of poloidal harmonics typical of ballooning modes.

External kink modes are usually difficult to avoid in MHD models of an infinitely conducting plasma surrounded by a vacuum region, and set β -limits below internal ballooning results.¹ Theoretical studies show a sensitivity to

the shape of the current profile near the plasma surface. Stability depends on the presence of a nearby conducting shell and we adopt the usual approach of presenting the stability boundaries as a function of wall radius. To do this we have examined each of the equilibria in the $\langle \beta \rangle - i$ plane as a function of A_w/A , and plot contours of marginal stability in Fig. 5.

At low β the external modes are localized near the plasma surface and can be described as surface kinks. Indentation is destabilizing in the sense that the wall must be closer to achieve stability as indentation is increased. At modest indentations, $i < 0.2$, the modes take on a strong ballooning character as β is increased (see Fig. 6). Eventually, as the second stable region to internal ballooning modes is approached, the advantages associated with favorable local shear and the magnetic well arising from the Shafranov shift appear, and the configuration becomes stable. At strong indentations, finite β effects are seen to stabilize the kink modes, an effect previously observed in a circular cross section, quasi-uniform-current, $\beta \sim a/R$ model.⁹ When the results of Fig. 5 are considered simultaneously with the infinite- n ballooning mode boundary⁴ (broken line), we see that, with a perfectly conducting wall at 1.3 plasma radius, accessibility to the second stable region without encountering current driven kinks should be achievable.

While the experimental verification of external kink modes in modern tokamaks remains an open issue, operation with $q(0) < 1$ is usual and measurements indicate significant activity associated with the $m = 1$, $n = 1$ internal mode. It is usually believed that the mode occurs as a resistive kink, whose nonlinear evolution is associated with the sawtooth phenomenon; but, in high temperature discharges the driving forces will have much in common with the ideal MHD internal kink. Recently this mode has been associated with the onset of fishbone MHD oscillations in PDX,¹⁰ resulting in

poor fast ion confinement. Since, to date, this is the only clearly identifiable detrimental ideal MHD experimental effect to come with increasing β , it is obviously of considerable interest to study configurations which are stable to this mode.

Currently, preparations are underway to modify the PDX tokamak to enable a study of the effect of indentation on the achievable $\langle\beta\rangle$ values. Considerable analysis of equilibria with profiles consistent with data taken from actual high- β PDX discharges has been carried out. While these profiles do not display the second stable region to ballooning modes, they do indicate ballooning $\langle\beta\rangle$ limits ($\sim 10\%$) considerably above previous studies, and with $q(0) = 0.8$, stability to the internal kink up to the maximum $\langle\beta\rangle$ studied of 20%. These results give a quantitative measure to the basic idea that the stabilizing properties of shear and minimum-B can be enhanced in tokamaks by strong cross-sectional shaping.

ACKNOWLEDGMENTS

We are indebted to Dr. M. S. Chance whose results inspired this work, and who together with Dr. D. A. Monticello and Dr. M. Reusch helped in the preparation of the equilibria used here. We are grateful for many useful discussions with Dr. P. H. Rutherford and the PPL PDX and Theory groups.

This work was supported by U. S. Department of Energy Contract No. DE-AC02-76-CHO-3073.

REFERENCES

- ¹A. M. M. Todd, J. Manickam, M. Okabayashi, M. S. Chance, R. C. Grimm, J. M. Greene, and J. L. Johnson, Nucl. Fusion 19, 743 (1979).
- ²C. Mercier, in Lectures in Plasma Physics, EURATOM-CEA/CEN/EUR 5/27e, Luxembourg (1974).
- ³R. Miller and R. Moore, Phys. Rev. Lett. 43, 765 (1979).
- ⁴M. S. Chance, S. C. Jardin, and T. H. Stix (to be published).
- ⁵M. N. Bussac, R. Pellat, D. Edery, and J. L. Soule, Phys. Rev. Lett. 35, 1638 (1975).
- ⁶B. Coppi, G. B. Crew, and J. J. Ramos, MIT Report PTP-82/6, 1982.
- ⁷R. L. Dewar, J. Manickam, R. C. Grimm, and M. S. Chance, Nucl. Fusion 21, 493 (1981).
- ⁸R. C. Grimm, R. L. Dewar, and J. Manickam, J. Comput. Phys. 49, 94 (1982).
- ⁹J. P. Freidberg, J. P. Goodbloed, W. Grossman, and F. A. Haas, in Plasma Physics and Controlled Thermonuclear Fusion Research, (IAEA, Tokyo, 1974), Vol. I, pp. 505.
- ¹⁰D. Johnson et al., in Proceedings of the Ninth International Conference on Plasma Physics and Controlled Nuclear Fusion Research, (IAEA, Baltimore, 1983) Vol. 1, p. 9.

FIGURE CAPTIONS

- FIG. 1. Parameters used to describe the bean-shaped plasmas studied.
- FIG. 2. Stability boundaries for the internal kink with $\alpha = \beta = 2$ and various indentations.
- FIG. 3. Stability boundaries for the internal kink with $\alpha = 1$, $\beta = 2$, and various indentations.
- FIG. 4. Internal kink displacement vector ξ , for $\epsilon\beta_p = 0.16$, $q(0) = 0.9$.
- FIG. 5. Stability boundaries to the external $n = 1$ kink mode for different wall positions. Shaded regions are unstable.
- FIG. 6. Strongly ballooning driven kink mode with $i = 0.112$, $\langle\beta\rangle = 8.2\%$, $A_w/A = 0.5$.

Fig. 1

#83T0293

Fig. 2

#83T0292

Fig. 3

Fig. 4

Fig. 5

Fig. 6

EXTERNAL DISTRIBUTION IN ADDITION TO TIC UC-20

Plasma Res Lab, Austri Nat'l Univ, AUSTRALIA
 Dr. Frank J. Paoloni, Univ of Wollongong, AUSTRALIA
 Prof. I.R. Jones, Flinders Univ., AUSTRALIA
 Prof. M.H. Brennan, Univ Sydney, AUSTRALIA
 Prof. F. Cap, Inst Theo Phys, AUSTRIA
 Prof. Frank Verheest, Inst theoretische, BELGIUM
 Dr. D. Palumbo, Dg XII Fusion Prog, BELGIUM
 Ecole Royale Militaire, Lab de Phys Plasmas, BELGIUM
 Dr. P.H. Sakanaka, Univ Estadual, BRAZIL
 Dr. L.R. James, Univ of Alberta, CANADA
 Prof. J. Teichmann, Univ of Montreal, CANADA
 Dr. H.M. Skarsgaard, Univ of Saskatchewan, CANADA
 Prof. S.R. Sreenivasan, University of Calgary, CANADA
 Prof. Tudor W. Johnston, INRS-Enargie, CANADA
 Dr. Hannes Barnard, Univ British Columbia, CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Zhengwu Li, SW Inst Physics, CHINA
 Library, Tsing Hua University, CHINA
 Librarian, Institute of Physics, CHINA
 Inst Plasma Phys, SW Inst Physics, CHINA
 Dr. Peter Lukac, Komenského Univ, CZECHOSLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Prof. Schatzman, Observatoire de Nice, FRANCE
 J. Redet, CEN-BP6, FRANCE
 AM Dupes Library, AM Dupes Library, FRANCE
 Dr. Tom Mual, Academy Bibliographic, HONG KONG
 Preprint Library, Cent Res Inst Phys, HUNGARY
 Dr. A.K. Sunderan, Physical Research Lab, INDIA
 Dr. S.K. Trehan, Panjab University, INDIA
 Dr. Indra, Mohan Lal Das, Benares Hindu Univ, INDIA
 Dr. L.K. Chavda, South Gujarat Univ, INDIA
 Dr. R.K. Chhajlani, Var Ruchi Marg, INDIA
 P. Kew, Physical Research Lab, INDIA
 Dr. Phillip Rosenau, Israel Inst Tech, ISRAEL
 Prof. S. Cupernan, Tel Aviv University, ISRAEL
 Prof. G. Rostaqni, Univ DI Padova, ITALY
 Librarian, Int'l Ctr Theo Phys, ITALY
 Miss Ciella De Palo, Assoc EURATOM-CIEN, ITALY
 Biblioteca, del CNR EURATOM, ITALY
 Dr. H. Yamato, Toshiba Res & Dev, JAPAN
 Prof. M. Yoshikawa, JAERI, Tokai Res Est, JAPAN
 Prof. T. Uchida, University of Tokyo, JAPAN
 Research Info Center, Nagoya University, JAPAN
 Prof. Kyoji Nishikawa, Univ of Hiroshima, JAPAN
 Prof. Sigeru Mori, JAERI, JAPAN
 Library, Kyoto University, JAPAN
 Prof. Ichiro Kawakami, Nihon Univ, JAPAN
 Prof. Satoshi Itoh, Kyushu University, JAPAN
 Tech Info Division, Korea Atomic Energy, KOREA
 R. England, Ciudad Universitaria, MEXICO
 Bibliotheek, Fom-Inst Voor Plasma, NETHERLANDS
 Prof. B.S. Lillev, University of Waikato, NEW ZEALAND
 Dr. Suresh C. Sharma, Univ of Calabar, NIGERIA
 Prof. J.A.C. Cabral, Inst Superior Tech, PORTUGAL
 Dr. Octavian Petrus, ALI CUZA University, ROMANIA
 Dr. R. Jones, Nat'l Univ Singapore, SINGAPORE
 Prof. M.A. Heilberg, University of Natal, SO AFRICA
 Dr. Johan de Villiers, Atomic Energy Bd, SO AFRICA
 Dr. J.A. Togle, JEN, SPAIN
 Prof. Hans Wilhelmson, Chalmers Univ Tech, SWEDEN
 Dr. Lennart Stenflo, University of UMEA, SWEDEN
 Library, Royal Inst Tech, SWEDEN
 Dr. Erik T. Karlson, Uppsala Universitet, SWEDEN
 Centre de Recherches, Ecole Polytech Fed, SWITZERLAND
 Dr. W.L. Weise, Nat'l Bur Stand, USA
 Dr. W.M. Stacey, Georgia Inst Tech, USA
 Dr. S.T. Wu, Univ Alabama, USA
 Prof. Norman L. Gleson, Univ S Florida, USA
 Dr. Benjamin Ma, Iowa State Univ, USA
 Prof. Magne Kristiansen, Texas Tech Univ, USA
 Dr. Raymond Askew, Auburn Univ, USA
 Dr. V.T. Tolok, Kharkov Phys Tech Ins, USSR
 Dr. D.D. Ryutov, Siberian Acad Sci, USSR
 Dr. M.S. Rabinovich, Lebedev Physical Inst, USSR
 Dr. G.A. Eliseev, Kurchatov Institute, USSR
 Dr. V.A. Glukhikh, Inst Electro-Physical, USSR
 Prof. T.J. Boyd, Univ College N Wales, WALES
 Dr. K. Schindler, Ruhr Universität, W. GERMANY
 Nuclear Res Estab, Jülich Ltd, W. GERMANY
 Librarian, Max-Planck Institut, W. GERMANY
 Dr. H.J. Kaeppler, University Stuttgart, W. GERMANY
 Bibliothek, Inst Plasmaforschung, W. GERMANY