DOE/ER/13754--T1

DOE/ER/13754--T1

DE92 010607

DOE Grant No: DE-FG03-87ER-13754
PI: Christopher J. Lamb
INSTITUTION: Salk Institute for Biological Studies
TITLE: Biosynthesis of Plant Plasmamembrane Polypeptides

In this project we examined molecular aspects of the functional organization of plant plasma membranes with an initial focus on plasma membrane antigens that reacted with a previously characterized monoclonal antibody, designated MAb 16.4B4, identified by immunofluorescent staining of the surface *Nicotiana glutinosa* protoplasts (1). The epitope reactive with MAb 16.4B4 was shown to be present in a glycan moiety and defined a family of surface glycoproteins with molecular masses in the range 135-180 kilodalton (kDa) (2). These glycoproteins were exclusively associated with the plasma membrane as demonstrated by immunostaining of highly purified plasma membrane vesicles obtained by aqueous two-phase partitioning of microsomal fractions. The bulk of these glycoproteins were not released by high-salt washing, sonication or hypotonic shock treatment of plasma membrane vesicles, demonstrating a tight association with the membrane. Triton X-114 partitioning of plasma membrane vesicles indicates that these antigens were hydrophilic, peripheral membrane glycoproteins. The glycoproteins were purified by immunoaffinity chromatography following solubilization in sodium dodecyl sulfate and shown to contain glycan moieties abundant in arabinose and galactose linked to a 50-kDa polypeptide rich in alanine, glycine, serine and threonine.

To investigate the biosynthesis of these plasma membrane glycoproteins in relation to the ontogeny of plant cell membranes, we investigated the subcellular distribution of antigenic sites immunoreactive to MAb 16.4B4 in tobacco (*Nicotiana tabacum*) leaf cells (3). We demonstrated by immunogold electron microscopy that the epitope reactive with MAb 16.4B4 was localized on the cell surface in the leaf parenchyma cell periplast. The 16.4B4 antigen was also localized in multivesicular invaginations of the plasma membrane also known as plasmalemmasomes, implying a

MASTER
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

biochemical and, hence, functional interrelationship between these structures. MAb 16.4B4 also labels intracellular multivesicular bodies that appear to represent internalized plasmalemmasomes. Antibody reactivity was also observed in partially degraded multivesicular bodies sequestered within the central vacuole. We propose that the subcellular distribution of the epitope reactive with MAb 16.4B4 defines a plasmalemmasome (or multivesicular body)-mediated pathway for the internalization of the periplasmic matrix for vacuolar-mediated disposal. The multivesicular bodies appear to be equivalent to the well-characterized endosomes and multivesicular bodies of animal cells involved in the internalization and lysosome-mediated degradation of extracellular materials.

The plasma membrane arabinogalactan protein family defined by MAb 16.4B4 has now been re-purified by immunoaffinity chromatography for the generation of antibodies to the apoprotein and peptide sequencing prior to the molecular cloning of cDNAs encoding the 50-kDa apoprotein. This will allow further studies of the structure, expression and function of these abundant plasma membrane glycoproteins.

In view of the involvement of these arabinogalactan proteins in membrane and vesicle internalization from the cell surface, we started to examine the perturbation of plasma membrane and cell surface functions during plant : pathogen interactions. In the initial stages of these studies we showed that treatment of bean or soybean cells with fungal elicitor or glutathione causes a rapid insolubilization of pre-existing (hydroxy)proline-rich structural proteins in the cell wall (4). This insolubilization, which involves H_2O_2 -mediated oxidative cross-linking, is initiated within 2 minutes and complete within 10 minutes under optimal conditions, and hence precedes the expression of transcription-dependent defense responses such as the synthesis of phytoalexin antibiotics and lytic enzymes. Cross-linking is also under developmental control associated with cellular maturation during hypocotyl growth and in tissues subject to mechanical stress such as the stem : petiole junction. Stimulus-dependent oxidative cross-linking of plant cell wall structural proteins is a novel site of cellular regulation with potentially important functions in the flexible realization of the final functional architecture of cell walls during development and rapid toughening of cell walls in the initial stages of plant defense.

In addition, we have examined the molecular basis for two key plasma membrane functions: synthesis of callose and translocation of sucrose. There is a strong correlation between induced callose synthesis and resistance of barley to all races of the fungal pathogen powdery mildew. Upon pathogen attack callose (1,3- β -polyglucoase) is deposited as papillae between the plasma membrane and the cell wall. We have purified the polypeptides constituting the host plasma membrane-localized callose synthase. Five putative polypeptides are being sequenced. We intend to clone the genes for these polypeptides by screening cDNA libraries with complementary oligonucleotide probes, or alternatively by cloning using PCR amplification. The cDNA clones will be used to study gene regulation in response to infection and wounding. Antisera to synthetic peptides or purified polypeptides will be used to monitor the constitutive/induced levels of the proteins in the plasma membrane. Both the cellulose and the callose synthesizing activities are suggested to be localized in the same high molecular weight protein complex, the "rosettes" of the plant plasma membrane. Antisera will allow us to investigate the presumed identity of the callose synthase and the cellulose synthases.

We have identified and purified a sucrose translocator of the plant plasma membrane which transports sucrose into cells and seems to be involved in phloem loading. With antisera raised against the purified sucrose translocator we have isolated and sequenced a cDNA clone from an *Arabidopsis thaliana* expressed library. Some sequence similarities to eukaryotic hexose carriers can be discerned. We are currently characterizing and verifying this clone. By immunogold labeling we are investigating the role of the sucrose translocator in phloem loading.

REFERENCES

- (1) Norman, P.M., Wingate, V.P.M., Fitter, M.S. and Lamb, C.J. (1986) Monoclonal antibodies to plant plasmamembrane antigens. *Planta* **167**, 452-459.
- (2) Norman, P.M., Kjellbom, P., Bradley, D.J., Hahn, M.G. and Lamb, C.J. (1990) Immunoaffinity purification and biochemical characterization of plasma membrane arabinogalactan-rich glycoproteins. *Planta* **181**, 365-374.

- (3) Herman, E.M. and Lamb, C.J. (1992) Arabinogalactan-rich glycoproteins are localized on the cell surface and in intravacuolar multivesicular bodies. *Plant Physiol.* **98**, 264-272.
- (4) Bradley, D.J., Kjellbom, P. and Lamb, C.J. (1992) Stimulus-dependent oxidative cross-linking of a proline-rich plant cell wall protein: A novel, rapid defense response and control point in cell maturation. *Cell*, submitted.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor an agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

preprint removed.
ds

END

**DATE
FILMED
5/13/92**