DOE/PC/89772--T6

**APPLICATIONS OF MICELLAR ENZYMOLOGY TO CLEAN COAL
TECHNOLOGY**

DOE/PC/89772--T6

DE92 010939

Ninth Quarterly Report

DOE Grant No. DE-FG22-89PC89772

MAR 30 1992

Submitted to:

**U.S. Department of Energy
Pittsburgh Energy Technology Center
P.O. Box 10940
Pittsburgh, PA 15236**

Prepared by:

**Carol T. Walsh, Ph.D.
Principal Investigator
Boston University School of Medicine
80 E. Concord Street, L-603
Boston, MA 02118
617-638-4326**

January 30, 1991

**US/DOE Patent Clearance is not required prior to the publication
of this document.**

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

FEB 03 1992

TABLE OF CONTENTS

Section		Page
1	INTRODUCTION	1
1.1	Program Overview	1
1.2	Summary of Results	4
2	MATERIALS AND METHODS	5
2.1	Materials	5
2.2	Methods	5
2.2.1	Enzyme Activity in Reverse Micelle Solutions with MC as Substrate	5
2.2.2	Enzyme Activity in Reverse Micelle Solution with EPS as Substrate	6
2.2.2a	Quantitation of EPS Sulfur Oxidation Products	6
2.2.2b	Reverse Micelle Solutions	7
3	RESULTS	8
3.1	Comparison of Chloroperoxidase Activity in CTAB/1-Hexanol and AOT/Isooctane Reverse Micelle Solutions with MC as Substrate	8
3.2	Relationship of H ₂ O ₂ to Chloroperoxidase Activity with MC as Substrate	11
3.2.1	Dependence on H ₂ O ₂ Concentration in Reverse Micelle Solution	11
3.2.2.	Inhibition by Preincubation with H ₂ O ₂	15
3.3	EPS Sulfoxidation by Chloroperoxidase in Reverse Micelle Solution	18
4	PLANS FOR THE TENTH QUARTER	24
5	BIBLIOGRAPHY	26

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
Figure 1: Enzymatic Sulfoxidation In Reverse Micelles	2
Figure 2: DBT, EPS and Metabolites in Sulfur Oxidation Pathway	3
Figure 3: Concentration vs Absorbance of MC in CTAB/1-Hexanol and AOT/Issooctane Reverse Micelle Solutions	9
Figure 4: Comparison of Chloroperoxidase Activity in CTAB/1-Hexanol and AOT/Issooctane Reverse Micelle Solutions	10
Figure 5: Dependence of Chloroperoxidase Activity in CTAB/1-Hexanol and AOT/Issooctane Reverse Micelle Solutions on MC Concentration	12
Figure 6: Effect of Preincubation Time in Reverse Micelle Solution on Chloroperoxidase Activity	13
Figure 7: Dependence of Chloroperoxidase Activity in Reverse Micelle Solution on H ₂ O ₂ Concentration	14
Figure 8: Inhibition of Chloroperoxidase by Preincubation with H ₂ O ₂ in Aqueous Solution	16
Figure 9: Effect of Chloroperoxidase Concentration (11-44 μ M) on EPS Sulfoxidation in Reverse Micelle Solution	19
Figure 10: Effect of Chloroperoxidase Concentration (1-11 μ M) on EPS Sulfoxidation in Reverse Micelle Solution	20
Figure 11: Time Dependence of EPS Sulfoxidation by Chloroperoxidase in Reverse Micelle Solution	21
Figure 12: Effect of H ₂ O ₂ Concentration on EPS Sulfoxidation by Chloroperoxidase in Reverse Micelle Solution	23
Figure 13: Effect of Sequential H ₂ O ₂ Addition on EPS Sulfoxidation in Reverse Micelle Solution	24

LIST OF TABLES

<u>Table</u>	<u>Page</u>
Table 1: Effect of Preincubation with H ₂ O ₂ on Chloroperoxidase Activity in Reverse Micelle Solution	17

Section 1

INTRODUCTION

1.1 Program Overview

Full implementation of coal fuel sources will require more effective methods of providing "clean coal" as a fuel source. Methods must be developed to reduce the sulfur content of coal which significantly contributes to environmental pollution. This project is designed to develop methods for pre-combustion coal remediation by implementing recent advances in enzyme biochemistry. The novel approach of this study is incorporation of hydrophilic oxidative enzymes in reverse micelles in an organic solvent (Figure 1).

Enzymes from commercial sources or microbial extracts are being investigated for their capacity to remove organic sulfur from coal by oxidation of the sulfur groups, splitting of C-S bonds and loss of sulfur as sulfuric acid. Dibenzothiophene (DBT) and ethylphenylsulfide (EPS) are serving as models of organic sulfur-containing components of coal in initial studies (Figure 2).

A goal of this project is to define a reverse micelle system that optimizes the catalytic activity of enzymes toward desulfurization of model compounds and ultimately coal samples. Among the variables which will be examined are the surfactant, the solvent, the water:surfactant ratio and the pH and ionic strength of the aqueous phase. Studies by several groups (Martinek et al., 1981; Kabanov et al., 1988; Martinek, 1989; Verhaert et al., 1990) have shown that the surfactant AOT over a broad concentration range in organic solvents produces micelles, comparatively uniform in

Figure 1: Enzymatic Sulfoxidation in Reverse Micelles

Figure 2: DBT, EPS and Metabolites in Sulfur Oxidation Pathway

diameter, which incorporate hydrophilic enzymes. The activity (kcat) of certain enzymes in this system is higher than in aqueous solution. This surfactant is therefore being examined first, although the potential disadvantages of an SO_3 containing molecule have been recognized. Other surfactants to be tested include Tritons, Tweens, CTAB and Brij 35.

1.2 Summary of Results

Chloroperoxidase was active in both AOT/isooctane and CTAB/1-hexanol reverse micelle solutions, as evidenced by chlorination of MC. The latter reverse micelle system produced more variable results with a lower reaction velocity than observed with the former solution. The activity of the enzyme decreased with incubation time at room temperature but the drop was not substantially different from that observed in aqueous solution. The enzyme was inhibited by brief preincubation with H_2O_2 , which almost certainly explained the lack of EPS sulfoxidation in reverse micelle solutions, reported earlier. Definitive enzyme mediated sulfoxidation of EPS in an AOT/isooctane reverse micelle solution was demonstrated, when peroxide was added already diluted in AOT/isooctane as in MC studies. However, the sulfoxidation of EPS was incomplete, considerably less than that observed under aqueous conditions, and not enhanced by an increase in enzyme concentration. The sequential addition of peroxide during the incubation was found to markedly enhance sulfoxidation. This observation is currently being further explored. In addition, the CTAB/1-hexanol reverse micelle solution is being evaluated for its ability to support chloroperoxidase catalysed EPS sulfoxidation.

Section 2

MATERIALS AND METHODS

2.1 Materials

Enzyme studies were performed with chloroperoxidase from Caldariomyces fumago. Chloroperoxidase, 2.7 mg/ml in 0.1 M sodium phosphate buffer (pH 4.0), was obtained from Sigma Chemical Co. and refrigerated at 0 °C. EPS (Aldrich Chemical Co.) and MC were evaluated as enzyme substrates. Two reverse micelle solutions were studied: 1) surfactant AOT (0.1 M, di (2-ethylhexyl) sodium sulfosuccinate from Sigma Chemical Co.), organic solvent isooctane (OmniSolv, EM Scientific), and 2.3% (v/v) aqueous phase (potassium phosphate, 0.1 M, pH 2.75); 2) surfactant CTAB (0.1 M, hexadecyltrimethylammonium bromide, Sigma Chemical Co.), solvent 1-hexanol (Fisher Scientific) and 2.3% (v/v) aqueous phase (potassium phosphate, 0.1 M, pH 2.75). Hexane, chloroform, acetone and isopropanol, all from Fisher Scientific, were used in sample analysis. Solid phase extractions were performed on 20H (diol) Bond Elut (500 mg) Bond Elut columns from AnalytiChem Inc. (Van Horne, 1990). A Scientific Industries rotator was used for agitation of incubation samples.

2.2 Methods

2.2.1 Enzyme Activity in Reverse Micelle Solutions with MC as Substrate

The assay method for measuring chloroperoxidase activity is based on procedures developed by Morris and Hager (1966) and Hager et al. (1966) and involves monitoring the enzymatic chlorination of

monochlorodimedon to form the dichloro derivative, 1,1-dimethyl-4,4-dichloro-3,5-cyclohexanedione. To test the activity of chloroperoxidase in reverse micelles, two solutions were made. The first contained 0.1 M AOT/isooctane or CTAB/1-hexanol with 0.1 mM monochlorodimedon, to which was added an aqueous phase (2.3%) containing enzyme (4.4 μ M) and KCl (0.04 M) in phosphate buffer (pH 2.75). The second solution contained the same organic phase but the aqueous phase (2.3%) consisted of 20 mM H₂O₂ in phosphate buffer. These solutions were combined in a 1:1 ratio and vortexed for 10 sec. After an additional 10-sec delay, the absorbance of the solution was measured for 20-30 sec at 278 nm on an LKB Biochrom 4050 spectrophotometer using "Reaction Rate" software to collect absorbance data and calculate reaction velocity. Nonenzyme-containing solutions were also tested. All conditions were tested in triplicate.

2.2.2 Enzyme Activity in Reverse Micelle Solution with EPS as Substrate

2.2.2a Quantitation of EPS Sulfur Oxidation Products

Isopropanol (1 column volume, 2.8 ml) was used to solvate a 20H (diol) Bond Elut column (500 mg). Excess isopropanol was removed with two column-volume washes of isooctane at a flow rate of 15 ml/min. A 0.5 ml aliquot of standards or experimental samples in reverse micelle solutions was applied and run to dryness at a flow rate of 4.8 ml/min. The sample was eluted five times with 0.25 ml aliquots of 80:20 chloroform:acetone. Eluates 2-5 were collected, solvent evaporated under N₂, and the residue redissolved in 85:15 hexane:isopropanol (0.1 ml) for HPLC analysis. Recovery of 0.009 mM

or 0.125 mM EPSx from reverse micelle solution, based on comparison of area values to that of EPSx in hexane:isopropanol (85:15), consistently averaged about 100%.

EPS (0.5 mM), EPSx (0.009 or 0.125 mM), and EPSn (0.5 mM) were added alone to hexane:isopropanol (85:15, v/v). These standards and aliquots of eluates from extraction columns (20 μ l) were analyzed with a Shimadzu HPLC system (LC-600 analytical pumps) using a Waters μ Porasil column (10 μ m, 3.9x300 mm). This normal phase chromatographic system was based on the method used by Light *et al.* (1982) and Waxman *et al.* (1982) to separate sulfones and sulfoxides. The mobile phase consisted of 85:15 hexane:isopropanol at a flow rate of 1.2 ml/min for 15.0 min. The detector was a SPD-6AV UV-VIS spectrophotometer set at 254 nm. Chromatograms were produced on a Chromatopac CR 501 data processor.

2.2.2b Reverse Micelle Solutions

EPS (0.5 mM) was incubated with chloroperoxidase and H₂O₂ for 0.5-4.0 hour at 23°C in reverse micelle solutions. These solutions were made by: 1) combining one aqueous phase solution, containing both enzyme and peroxide in 0.1 M phosphate buffer, and the organic phase, 0.1 M AOT-isooctane, containing 0.5 mM EPS; 2) making separate enzyme-containing and peroxide-containing reverse micelles solutions and then combining them. The solutions were vortexed for 1 min and then incubated. In each experiment controls included EPS without enzyme or peroxide and EPS with peroxide and no enzyme. All conditions were tested with at least duplicate incubations; replicate aliquots of each (0.5 ml) were analyzed by 20H (diol) extraction and HPLC.

Section 3

RESULTS

3.1 Comparison of Chloroperoxidase Activity in CTAB/1-Hexanol and AOT/Isooctane Reverse Micelle Solutions with MC as Substrate

The activity of chloroperoxidase, measured as the chlorination rate of MC, was tested in 0.1 M AOT-isooctane and 0.1 M CTAB/1-hexanol reverse micelle solutions with 2.3 % aqueous phase. The concentration-absorbance relationship of MC at 278 nm in these two solutions was similar (Figure 3). Enzyme activity data are therefore presented as absorbance units per min since correction for differences in extinction coefficient appeared unnecessary. Peroxide-containing controls (without enzyme) exhibited insignificant levels of activity.

The enzymatic reaction velocity was significantly higher in AOT/isooctane as compared to CTAB/1-hexanol reverse micelle solutions (Figure 4). The rate was significantly increased in AOT/isooctane solutions when the enzyme concentration was increased ($p < 0.0001$, Student's t-test). In contrast, the rate was not significantly greater when the same increase in enzyme level was tested in CTAB/1-hexanol solutions; greater variability was also noted. Subsequently, a summary of five experiments indicated that for identical experimental conditions the coefficient of variation for the reaction rate in AOT/isooctane was 15.6% whereas that for CTAB/1-hexanol was 62.2%. The mean rate in CTAB/1-hexanol averaged 57.8% of that in AOT/isooctane. In both reverse micelle solutions the 0.1 M concentration of MC appeared to generate

Figure 3: Concentration vs Absorbance of MC in CTAB/1-Hexanol and AOT/Isooctane Reverse Micelle Solutions

Figure 4: Comparison of Chloroperoxidase Activity in CTAB/1-Hexanol and AOT/Isooctane Reverse Micelle Solutions

maximal velocities; 50 and 100% increases in MC concentrations did not significantly increase reaction rates (Figures 5).

The stability of the enzyme in these two reverse micelle solutions was also compared. In this study the enzyme-containing reverse micelle solution was formed and the rate of MC chlorination then tested immediately or after a 2 or 4 hour incubation at room temperature. New peroxide-containing reverse micelle solutions were generated just prior to enzyme testing to eliminate a potential time-dependent change in this component of the assay mixture. The results, depicted in Figure 6, indicated that the rate of loss of enzyme activity was similar in the two types of reverse micelle solutions. After 4 hours the per cent of activity in the AOT/isooctane solution compared to 0 time, 45.5%, was similar to that observed when the enzyme stability was tested in aqueous solution (p. 10, Seventh Quarterly Report).

A partitioning experiment determined that 0.1 mM MC in isooctane was entirely extracted into an equal volume of 0.1 M phosphate buffer (pH 2.75).

3.2 Relationship of H₂O₂ to Chloroperoxidase Activity with MC as Substrate

3.2.1 Dependence on H₂O₂ Concentration in Reverse Micelle Solution

The effect of varying the H₂O₂ concentration on chloroperoxidase activity in the AOT/isooctane reverse micelle solution was determined. As indicated in Figure 7, maximal activity was observed with peroxide concentrations of 10-40 mM in the

Figure 5: Dependence of Chloroperoxidase Activity in CTAB/1-Hexanol and AOT/Isooctane Reverse Micelle Solutions on MC Concentration

Figure 6: Effect of Preincubation Time in Reverse Micelle Solution on Chloroperoxidase Activity

Figure 7: Dependence of Chloroperoxidase Activity in Reverse Micelle Solution on H₂O₂ Concentration

final aqueous phase of the reverse micelle solutions.

3.2.2 Inhibition by Preincubation with H₂O₂

Addition of peroxide to chloroperoxidase (4.0 nM) in aqueous solution without substrate present was found to markedly inhibit the chlorination of MC on subsequent addition of the peroxide-enzyme combination (within 5-15 min) to MC in phosphate buffer (Figure 8). This study was designed such that the final peroxide concentration in all cases was 3.3 mM. In one experiment where enzyme was preincubated with 10 mM peroxide, after combination with the substrate solution, additional peroxide was added equivalent to the amount used in control studies. However, activity was even further decreased. An inhibitory effect was also observed when the peroxide-enzyme solution was tested by addition to MC in AOT/isooctane (Table 1). These observations indicated that the most recent protocol for testing EPS sulfoxidation by this enzyme in reverse micelle solutions was inappropriate. The procedure of combining peroxide and enzyme in aqueous solution prior to addition to surfactant/solvent had been adopted in June of 1991 (p. 8-9, Seventh Report), when initial studies with peroxide and enzyme separately added to EPS-containing reverse micelle solutions had proved negative. Subsequent work described below modified the procedure to attempt to avoid the pronounced inhibitory effect of peroxide.

Figure 8: Inhibition of Chloroperoxidase by Preincubation with H_2O_2 in Aqueous Solution

Table 1: Effect of Preincubation with H₂O₂ on Chloroperoxidase Activity in Reverse Micelle Solution

REACTION VELOCITY^a (abs units/min)		
<u>CONDITION</u>	<u>ENZYME^b</u>	<u>BLANK</u>
Control ^c	0.508 ± 0.075 (3)	0.032 ± 0.006 (2)
Preincubation: 10 mM H ₂ O ₂ ^d	0.039 ± 0.025 (3)	0.026 ± 0.006 (2)
Control: 1 hr later	0.478 ± 0.083 (2)	-

a Change in absorbance units of 0.1 mM MC contained in all 0.1M AOT:isooctane solutions, measured at 278 nm for 1 minute.

b Chloroperoxidase concentration in aqueous phase: 2.2 μ M; concentration based on total volume:50 nM.

c Enzyme in 0.04 M KCl-phosphate (pH 2.75) added to 0.1 M AOT:isooctane, combined with an equal volume of 20 mM H₂O₂ in phosphate added to 0.1 M AOT:isooctane.

d Enzyme and 10 mM H₂O₂ in phosphate added after 5 min to 0.1 M AOT:isooctane, combined with 0.04 M KCl-10 mM H₂O₂ added to 0.1 M AOT:isooctane.

3.3 EPS Sulfoxidation by Chloroperoxidase in Reverse Micelle Solution

EPS as a substrate for chloroperoxidase in reverse micelle solutions was tested under several conditions. These studies were carried out with 20 mM peroxide in phosphate buffer added to AOT/isooctane, equivalent to the optimum for MC chlorination in the comparable experimental system. Concentrations of the enzyme in the aqueous phase added to the organic solution were initially tested at 44 μ M which exceeded the level of enzyme active in MC chlorination in reverse micelle solution by twenty fold and in EPS sulfoxidation in aqueous solution by two-hundred fold. In this initial study the production of EPSx after 2 hours was five-fold greater in enzyme-containing reverse micelle solutions than in controls. This encouraging observation led to further studies with a range of enzyme concentrations. The results of the two studies, shown in Figures 9 and 10, demonstrated significant enzyme-dependent sulfoxidation of EPS in the AOT/isooctane reverse micelle solution. The magnitude of EPSx production in two hours did not show a positive correlation with enzyme concentration. Furthermore, only about 10% of the substrate was oxidized to EPSx with no other metabolites detectable.

Additional studies were carried out to explore the factors limiting the extent of EPS sulfoxidation, since increasing the amount of enzyme in the reverse micelle solution did not enhance the conversion. The next experiment, summarized in Figure 11, revealed little difference in the extent of product formation after 2.0 hours as compared to a much shorter incubation time, 0.5 hours. This

Figure 9: Effect of Chloroperoxidase Concentration (11-44 μM) on EPS Sulfoxidation in Reverse Micelle Solution

Figure 10: Effect of Chloroperoxidase Concentration (1-11 μM) on EPS Sulfoxidation in Reverse Micelle Solution

Figure 11: Time Dependence of EPS Sulfoxidation by Chloroperoxidase in Reverse Micelle Solution

finding was inconsistent with earlier studies of enzyme activity as a function of incubation time, based on MC as substrate, and suggested either product inhibition or depletion of peroxide might be limiting further EPS sulfoxidation. EPSx had previously been shown to be a substrate for chloroperoxidase in aqueous solution so that the former hypothesis seemed less likely. Therefore, in the next experiment, a four-fold higher peroxide concentration was tested. The magnitude of EPSx formation after a 0.5 hr incubation was increased by the higher peroxide level but only by 34% (Figure 12). The effect of sequential peroxide addition to the incubation mixture was examined next. In this experiment aliquots of the reverse micelle solutions were removed for EPSx determination after 0.5 hr incubations, and at that time additional phosphate buffer (2.3% of the total volume), either alone or containing 10 mM peroxide, was added. Marked enhancement of EPS sulfoxidation was produced by the second addition of peroxide (Figure 13, "samples" 1=just before and 0.5 hr after second peroxide addition, 2=just before and 0.5 hr after addition of phosphate control, dark bars are "before" condition). The extent to which this effect was dependent on enzyme catalysis cannot be determined from this experiment, since a nonenzyme-containing control was not included. Subsequent work will explore this question.

Figure 12: Effect of H_2O_2 Concentration on EPS Sulfoxidation by Chloroperoxidase in Reverse Micelle Solution

Figure 13: Effect of Sequential H₂O₂ Addition on EPS Sulfoxidation in Reverse Micelle Solution

Section 4

PLANS FOR THE TENTH QUARTER

1. Further study of chloroperoxidase and bacterial and mammalian enzymes (Holland, 1988, Kobayashi et al., 1986, and Rettie et al., 1990) for sulfur oxidizing activity.
2. Methodology for peroxide addition to maximize EPS sulfoxidation by chloroperoxidase in reverse micelle solution.
3. Development of analytical assay for EPSx in a CTAB/1-hexanol reverse micelle solution and determination of EPS sulfoxidation by chloroperoxidase in this solution.

Section 5

BIBLIOGRAPHY

Hager, L.P., Morris, D.R., Brown, F.S., and Eberwein, H. Chloroperoxidase: utilization of halogen anions. Journal of Biological Chemistry **241**: 1769-1777, 1966.

Holland, H.E. Chiral sulfoxidation by biotransformation of organic sulfides. Chem. Rev. **88**: 473-485, 1988.

Kabanov, A.V., Levashov, A.V., Klyachko, N.L., Namyotkin, S.N., Pshezhetsky, A.V. and Martinek, K. Enzymes entrapped in reversed micelles of surfactants in organic solvents: a theoretical treatment of the catalytic activity regulation. J. Theor. Biol. **133**: 327-343, 1988.

Kobayashi, S., Nakano, M., Goto, T., Kimura, T., and Schaap, A.P. Evidence of the peroxidase-dependent oxygen transfer from hydrogen peroxide to sulfides. Biochemical and Biophysical Research Communications **135**: 166-171, 1986.

Light, D. R., Waxman, D. J., and Walsh, C. T. Studies on the chirality of sulfoxidation catalyzed by bacterial flavoenzyme cyclohexanone monooxygenase and hog liver flavin adenine dinucleotide containing monooxygenase. Biochemistry **21**: 2490-2498, 1982.

Martinek, K. Micellar enzymology: potentialities in fundamental and applied areas. Biochemistry International **18**: 871-893, 1989.

Martinek, K., Levashov, A.V., Klyachko, N.L., Pantin, V.I., and Berezin, I.V. The principles of enzyme stabilization vs. catalysis by water-soluble enzymes entrapped into reversed micelles of

surfactants in organic solvents. Biochim. Biophys. Acta 657: 277-294, 1981.

Morris, D.R., and Lowell, L.P. Chloroperoxidase I. Isolation and properties of the crystalline glycoprotein. J. Biol. Chem. 241: 1763-1768, 1966.

Rettie, A.E., Bogucki, B.D., Lim, I., and Meier, G.P. Stereoselective sulfoxidation of a series of alkyl p-tolyl sulfides by microsomal and purified flavin-containing monooxygenases. Molecular Pharmacology 37: 643-651, 1990.

Verhaert, R.M.D., Hilhorst, R., Vermue, M., Schaafsma, J.J., and Veeger, C. Description of enzyme kinetics in reversed micelles. 1. Theory. Eur. J. Biochem. 187: 59-72, 1990.

Van Horne, K.C. Sorbent Extraction Technology. Analytichem International, Inc., 1990.

Waxman, D. J., Light, D. R., and Walsh, C. T. Chiral sulfoxidations catalyzed by rat liver cytochromes P-450. Biochemistry 21: 2499-2507, 1982.

END

**DATE
FILMED**

5 / 20 / 92