9:
201223

1 OF 1

DOE/ER/13705--T2

Progress report for DOE grant #DE-FG03-87ER13705

Title: Characterization and Expression of Clostridium Cellulase Genes

Period of support: May 1, 1987 to April 30, 1989

Principal Investigator: Roy H. Doi

DOE/ER/13705--T2

I. Personnel

DE92 012231

A. Principal Investigator

Roy H. Doi, Department of Biochemistry & Biophysics, University of California, Davis, CA 95616
Tel. (916)752-3191
Fax (916)752-3085

B. Postdoctoral Fellows

Dr. Ida K. Yu
Dr. Oded Shoseyov

II. Major goals of research

The overall goal of the project is to understand the genetics and molecular genetics of the gram positive, anaerobic, spore-forming Clostridium cellulovorans. We have chosen the cellulase enzyme and the genes that code for the subunits of the cellulase enzyme as a model system for our studies.

The major goals for the past two years have been as follows:

- A. The isolation and characterization of the cellulase enzyme complex from Clostridium sp.
- B. Cloning and characterization of the genes that encode the proteins of the cellulase enzyme complex.

We have made significant progress in both areas and have laid a sound foundation for our future studies.

III. Research accomplishments

- A. Purification and characterization of the cellulase enzyme complex from Clostridium cellulovorans.

- 1. Ammonium sulfate and cellulose affinity purification

We have been able to purify the cellulase enzyme from C. cellulovorans by growing the cells in medium containing Avicel (cellulose) as the carbon source. The supernatant portion of the

growth medium was precipitated with 80% ammonium sulfate and the enzyme activity in the precipitate included cellulase and cellobiase activity. The extracellular cellulase complex was isolated from the crude protein by taking advantage of the extremely high affinity of the complex to cellulose. The cellulase could not be washed off the cellulose by 1 M sodium chloride nor by phosphate-citrate buffer (50 and 12.5 mM, respectively, pH 7.0). Repeated washings of the cellulase-cellulase complex with deionized water resulted in the elution of the cellulase complex from cellulose.

2. SDS-PAGE analysis of the cellulase complex

This complex resolved by SDS-PAGE revealed three major protein bands having molecular masses of 170 kDa (P170), 100 kDa (P100), and 70 kDa (P70) and at least six minor protein bands of 80 kDa (P80), 75 kDa (P75), 65 kDa (P65), 58 kDa (P58), 43 kDa (P43), and 40 kDa (P40) (Fig. 1A). The cellulase fraction eluted by deionized water migrated as a single high molecular weight band after native PAGE (Fig. 1B). These results indicated that the C. cellulovorans cellulase is a multiple protein complex composed of different subunits having a heterologous stoichiometry. Among the three major bands the P170 eluted only at the last distilled water wash, whereas the P100 and P70 continuously "leaked off" at all the washing steps (Fig. 1A).

3. Analysis for CMCase activity of the cellulase subunits

CMCase activity was tested on the SDS-PAGE separated protein subunits of cellulase (Fig. 2). The major CMCase activity was associated with P100 with lower amounts of CMCase activity present in P80, P75, P65, and P58. Among the three major proteins, P70 was less active based on the intensities of the Commassie blue band compared to its activity band and our current feeling is that P70, in fact, does not contain any CMCase activity. The P170, P43, and P40 proteins showed no CMCase activity. The significance of these results with respect to the P170 is that this major protein, although it had no hydrolytic activity (that could be detected) possessed the highest affinity to cellulose. A similar protein was found recently in Clostridium thermocellum. That protein is believed to play a role as an anchor protein, an essential component that is required for the degradation of crystalline cellulose fibers.

4. Analysis of the native molecular weight of cellulase by FPLC

In order to determine the molecular weight of the cellulase complex the enzyme was injected into an FPLC system equipped with a molecular weight sizing column. Total protein, CMCase and cellulase activities were monitored (Fig. 3). Two major CMCase activities were detected that coincided with two protein peaks at 900 kDa and 100 kDa (Figs. 3A and 3B). The major cellulase activity appeared at the 900 kDa peak, but there was considerable

cellulase activity even at the fractions with masses less than the 12 kDa protein standard (cytochrome c) (Figs. 3A and 3C). SDS-PAGE followed by silver staining (Fig. 4) of the 900 kDa and 100 kDa peaks (arrows marked on Fig. 3A) revealed that 11 fractions containing cellulase activity had P170, P100 and P70. The most surprising phenomenon was the detection of relatively high amounts of P170 in the last low molecular weight fractions indicating a strong affinity of the lower molecular weight components to P170 that managed to retard its elution from the column.

This latter observation raises questions concerning the validity of several works that claimed the presence of low molecular weight cellulases as estimated by gel filtration chromatography. This technique can cause artifactual results as demonstrated in Figs. 3 and 4.

The nine proteins of the cellulase complex have been purified to homogeneity except for P80 and P75 which could not be resolved (Fig. 5). CMCase activity assay of all the isolated proteins confirmed our previous findings by the CMC-SDS-PAGE system (Fig. 2) from which we concluded that the major CMCase activity could be attributed to P100 and that P170, P43 and P40 had no detectable CMCase activity.

5. Specific inactivation of cellulase by antibodies against P170.

Specific polyclonal antibodies that had been prepared against P170 (Fig. 6) were mixed separately with the isolated cellulase complex and these mixtures were tested for cellulase, CMCase, and cellobiohydrolase (CBHase) activities. Anti-P170 inhibited cellulase activity, but not CMCase or CBHase activities (Fig. 7).

B. Construction of a gene bank and cloning of the cellulase subunit genes

1. A method was developed to isolate C. cellulovorans DNA rapidly and in high quantity. This DNA was used to construct a lambda gtl1 bank of the C. cellulovorans genome.
2. By use of selective substrates in the medium, four different CMCase clones and one cellobiase clone were isolated. The CMCase clones differ in the size and restriction map of the fragments containing the CMCase activity. Out of 5 cellobiase positive clones selected, it was shown that all 5 clones contained the identical fragment. Thus it is likely that only one cellobiase gene may be present. The EcoRI fragments that carry the CMCase genes are 3.2, 3.1, 2.8, and 2.2 kb and 2.5 kb for the cellobiase gene.
3. Since the CMCase and the cellobiase activities were shown to be present in the presence or absence of the inducer (IPTG), it is possible that these genes are being expressed from their own promoters and that C. cellulovorans promoters are active in E. coli.

C. Several other data concerning C. cellulovorans

1. The organism is sensitive to kanamycin. Thus if pUB110 can be transformed into this organism, we can use kanamycin as the selective marker.
2. The organism contains a beta-galactosidase gene. Thus if we are to attempt to use beta-galactosidase as a reporter gene, e.g. in gene fusion studies, we will have to select a beta-galactosidase negative mutant of C. cellulovorans.

IV. Summary of research accomplishments

- A. The cellulase enzyme complex of C. cellulovorans has been purified and its subunits identified. The complex has a relatively small molecular weight of about 900,000.
- B. The MWs of the subunits are as follows: 170,000 (P170), 100,000 (P100), 80,000 (P80), 75,000 (P75), 70,000 (P70), 65,000 (P65), 58,000 (P58), 43,000 (P43) and 40,000 (P40).
- C. No activity could be found associated with P170, P70, P43 and P41.
- D. CMCase activity was found to be associated with P100, P80, P75, P65, and P58.
- E. Antibodies have been prepared against the major proteins P170, P100 and P70.
- F. The antibody against P170 inactivates cellulase, but not CMCase or cellobiohydrolase (CBHase).
- G. A quick and efficient method for extracting DNA from C. cellulovorans was developed.
- H. A genomic bank of C. cellulovorans was constructed in lambda gt11.
- I. Four different CMCase clones and one cellobiase clone has been selected from the gene bank.

V. Manuscripts in preparation

- A. Shoseyov, O. and Doi, R.H. Properties of the Clostridium cellulovorans cellulase complex and its subunits.
- B. Shoseyov, O. and Doi, R.H. A 170 kDa subunit anchors Clostridium cellulovorans cellulase complex to cellulose.

Figure 1. PAGE analysis of the affinity purified cellulase complex. A. (SDS-PAGE). Lane 1, 0-80% $(\text{NH}_4)_2\text{SO}_4$ fraction; Lane 2, proteins that did not bind to the cellulose; Lane 3, 4 ml 1 M NaCl wash; Lane 4, 4 ml phosphate buffer wash; Lanes 5 and 6, 4 ml deionized water washes; Lane 7, 400 ml deionized water wash; Lane 8, molecular weight markers. B. (native-PAGE). Lane 1, 0-80% $(\text{NH}_4)_2\text{SO}_4$ fraction; Lane 2, the purified cellulase (A-7), Lane 3, 272,000 dalton urease marker.

Figure 2. CMC-SDS-PAGE analysis of the cellulase. Lane 1, Commassie blue staining; Lane 2, Activity staining.

Figure 3. FPLC analysis of the cellulase complex. A. 280 nm monitoring of the eluted protein. B. CMCase analysis of the fractions. C. Cellulase analysis of the fractions.

Figure 4. SDS-PAGE analysis of the FPLC fractions. Lane 1, the cellulase complex; Lane 2, Fraction no. 33; Lane 3, Fraction no. 51; Lane 4, Fraction no. 58; Lane 5, Fraction no. 66.

Figure 5. SDS-PAGE analysis of the purified cellulase subunits. A. Lane 1, the cellulase complex; Lane 2, P170; Lane 3, P100; Lane 4, P70. B. Lane 1, P40; Lane 2, P43; Lane 3, P58; Lane 4, P65; Lane 5, P75, P80; Lane 6, the cellulase complex; Lane 7, molecular weight markers.

Figure 6. Western blot analysis of P170. Lane 1, immunostaining of rabbit polyclonal anti-P170-IgG by alkaline phosphatase conjugated to goat anti-rabbit IgG. Lane 2, Coomassie blue staining of cellulase blotted on Immobilon-P. Lane 3, Molecular weight markers (200, 97, 68, and 43 kDa).

Figure 7. The effect of P170-IgG on cellulase (A), CMCase (B), and CBHase (C) activities. No. 1, cellulase. No. 2, cellulase + P170-IgG. No. 3, cellulase + preserum. No. 4, boiled cellulase.

END

**DATE
FILMED
6/10/92**