

BNL-NCS--43160

Prepared for
Pure Applied Chem.

DE90 001017

Total Half-lives for Selected Nuclides*

NORMAN E. HOLDEN
National Nuclear Data Center
Brookhaven National Laboratory

JULY 1 1980

Abstract — Measurements of the half-lives of ^3H , ^{10}Be , ^{14}C , ^{26}Al , ^{40}K , ^{38}Ar , ^{53}Mn , ^{87}Rb , ^{82}Nb , ^{129}I , ^{138}La , ^{147}Sm , ^{176}Lu , ^{174}Hf , ^{180}Ta , ^{187}Re , ^{186}Os , ^{190}Pt , ^{204}Pb , ^{210}Pb , ^{210}Po , ^{222}Rn , ^{224}Th , ^{226}Ra , ^{227}Ac , ^{228}Ra , ^{228}Th , ^{230}Th , ^{232}Th , ^{231}Pa have been compiled and evaluated. The effect of the ^{14}C half-life value on carbon dating ages is discussed as well as the stability of ^{204}Pb .

I. Introduction

In the past, many compilations and evaluations of half-lives have been made which have uncritically accepted authors' values and uncertainties. They have merely recommended weight-averaging reported results. This evaluation attempts to reanalyse each experiment in the literature including an estimate of the standard deviation utilizing, where possible, an estimate of the systematic error. The long-lived nuclides of light elements are of interest for their use in dating methods and for calculating cosmic-ray exposure ages of meteorites. The heavy mass nuclides are of interest in determining geological ages using the Re-Os or the Lu-Hf dating methods, in supplying information on the natural radioactive decay chains and in the case of tantalum because it represents the first long-lived state which is actually an isomer.

The impact of the recommended ^{14}C half-life of 5715 years on the carbon dating technique, which uses the Libby value of 5568 years, will be discussed. Also the possible primordial occurrence of ^{92}Nb is now definitely ruled out by the recommended half-life of 3.7×10^7 years. Based on the recommended ^{26}Al half-life value, the ^{21}Ne production rate for calculating cosmic-ray exposure ages remains too high, compared to rates using the ^{53}Mn and ^{10}Be half-life values. It is shown that ^{204}Pb , which was previously thought to be radioactive, is stable.

*Research carried out under the auspices of the U.S. Department of Energy, Contract No. DE-AC02-76CH00016.

MASTER

It will be noted that many of the uncertainties recommended here considerably exceed, by up to an order of magnitude, uncertainties quoted by individual authors in their publications; e.g. ^3H , ^{210}Po , ^{222}Rn , ^{227}Ac , and ^{228}Th .

The general procedure followed in this paper has been to review each experiment and to revise the published values for the latest estimates of various parameters originally reported by the authors; e.g. improved data on branching ratios assumed, on the half-lives of other nuclides involved, on the isotopic abundance in a natural sample, the nuclidic masses and the physical constants such as the Avogadro's constant. When detailed information on uncertainties was available in each experiment, the standard deviation was combined with one third of the systematic error to provide the uncertainty quoted in the table. The result of this procedure should be that the limit of error of the half-life would be obtained from the sum of the systematic error plus three standard deviations; i.e. 3σ . Where there was no adequate discussion of the systematic error and the total error was extremely small; e.g. 0.1 percent or less, a systematic error of 0.1 percent was estimated. One third of this amount, about 350 parts per million (ppm), was added to the published error to obtain the figure given in the various tables. The uncertainty listed for the recommended value in each table was calculated from a weighted average of the listed measurements using a variance weighting technique; either the reciprocal square of the author's reported uncertainty, or as revised according to the above scheme. Exceptions to the weighted average rule had to be made for some nuclides and will be discussed under the appropriate section for those nuclides. In such cases, recommendations were made using either a selected value considered superior to other listed measurements, or a weighted average was calculated for each of the different experimental techniques used and an unweighted average of these half-lives was recommended. All of the tables indicate the particular method chosen.

II. The Light Elements ($A < 100$)

For ^3H , a number of measurements have been reported for which the precision only is given. The reported values disagree by 20 to 30 standard deviations. The different techniques were weight-averaged and an unweighted average of these numbers was recommended.

For ^{14}C , Mann¹ discussed the problem of retention of a small amount of high specific activity ($\approx 0.02\%$) carbon dioxide during the gas dilution phase. This systematic effect could cause up to a 30% spread in the resulting half-life and was eliminated by substituting a clean flask during subsequent dilution phases. Earlier measurements, which varied from 4700–7200 years, were performed either with very low enrichment (a few percent) or with the above mentioned dilution process with large systematic error. These results were discarded. In Mann's revision² of his earlier measurement, he mentions a discrepancy between mass spectrometric determination of the amount of ^{14}C atoms. Samples which were run at NBS and Aldermaston showed a lower reading on one of the three machines at NBS. Mann noted that the result obtained on the mass spectrometer at AWRE agreed with the results on the two other NBS instruments but chose not to use this information. In my analysis, I have averaged the results on the

samples from all four instrument which has slightly lowered Mann's half-life. A weighted average gives 5692 ± 20 years, while an unweighted average gives 5715 ± 24 years. The unweighted average is recommended because the wide variation in authors estimates of systematic error sources tends to penalize those who do the best job of error analysis. The standard deviation is expanded to account for the variation in the weighted and unweighted averages and to allow for undisclosed systematic errors.

It should be noted that although the fifth (Godwin³) and sixth (Johnson⁴) International Carbon-14 Conferences recognized that the best available half-life at that time for the decay of radiocarbon was 5730 ± 40 years, the measurers of radiocarbon dates would continue to use 5568 years realizing that to obtain the correct dates, a factor of 1.03 must be used. The factor now becomes 1.026 with this recommended half-life.

For ^{39}Ar , the weighted average is 268 ± 8 years, where the 3% systematic error has been used rather than the 1% statistical error usually associated with this half-life.

For ^{40}K , the two significant decay branches of electron capture, ec, and negative beta decay, β , have been separately averaged and the total half-life has been calculated from the two decay constants. Most of the experiments have been reported at the 1% accuracy level. One similar experiment claims an accuracy of 0.1%. An unweighted average is recommended to treat all experiments on an equal level.

For ^{53}Mn , the early measurements assumed a constant cosmic ray flux over a period of 10 million years, which is a questionable assumption. Those early measurements have not been used.

For ^{92}Nb , Makino's result⁵ for the specific activity measurement as reported is in error. The half-life should be $3.98 \pm 0.76 \times 10^7$ years. In Nethaway's measurement⁶, he ignores all other measured ($n,2n$) cross section values for producing the m-state except his own⁷. The author notes a 10% effect is involved in treating the cross section for producing the long lived state. The author averages all total ($n,2n$) cross sections from 13 to 15 MeV, but selects the peak cross section for m-state production at 14.8 MeV. In this paper, I have renormalized the ^{238}U (n,f) flux monitor to the latest value of the Evaluated Nuclear Data File ENDF/B-V and I have recalculated the half life on the basis of 13-15 MeV average ($n,2n$) cross section difference for total and m-state production as well as 14.8 MeV differences. The former gives 3.79×10^7 years and the latter 4.02×10^7 years. An average is selected to represent this experiment.

III. The Medium Elements ($100 < A < 200$)

For ^{129}I , the most accurately quoted results are either unpublished or contain no details. An unweighted average of all data is recommended.

For ^{176}Lu , the two measurements which were performed with enriched samples do not agree. The difference is between four and seven standard deviations. An unweighted average is recommended.

For ^{174}Hf , ^{180}Ta , and ^{188}Os , the most recent measurement has been selected in each case. This corresponds to either the only value, a value which is far superior to other measurements or it is a higher upper limit to the total half-life.

For ^{187}Re , an unweighted average is recommended to take account of the measurement by Naldrett⁸, which is significantly lower than the other values.

II

IV. The Heavy Elements ($200 < A$)

For ^{204}Pb , Riezler⁹ used a nuclear emulsion technique to measure a sample of ^{204}Pb enriched to 27.0%. A peak was found between 8μ and 9μ in the emulsion, which from Faraggi's range energy curves¹⁰ was attributed to an alpha energy of 2.6 MeV. The latest mass data on ^{204}Pb , ^{200}Hg , and ^4He imply an available alpha energy of 1.93 MeV, i.e. a peak below 6.5μ . The peak has to be due to something other than the alpha decay of ^{204}Pb . There is no evidence that ^{204}Pb is radioactive. The most recent theoretical work¹¹ predicts a half-life value of 4.5×10^{35} years compared to Riezler's measurement of 1.4×10^{17} years.

For ^{210}Pb , the two most accurately quoted measurements do not agree. The difference is between seven and seventeen standard deviations. An unweighted average has been recommended.

For ^{210}Po , ^{222}Rn , ^{227}Ac and ^{228}Th , the recommended value is based on a weighted average of the measurements but the quoted value for the uncertainty has been increased to 0.1% (see discussion of results section).

For ^{230}Th , the results of Hyde¹² and Attree¹³ have been revised with the latest parameters as well as with the assumption that all the thorium in their samples, which was not ^{230}Th , was ^{232}Th . Meadows¹⁴ has recalculated all of the earlier measurements based on ^{226}Ra to the presently accepted half life of 1600 years.

For ^{232}Th , the recommended value is based on a weighted average of all measurements. The uncertainty has been increased from 0.5% to 0.7% to account for systematic errors.

V. Discussion of Results

In most cases, the recommended values and uncertainties in the tables are based on variance weighted averages. The recommended values listed are given in units of second (s), day (d), and year (a). Although it has been previously

discussed¹⁵, some words on the problem of error estimation can not be stated too often.

Various measurements in the tables below quote uncertainties that both disagree with and exclude many other good measurements from consideration. Undoubtedly, systematic errors have not been carefully considered in these publications. If one uses variance weighting indiscriminately in such cases, one penalizes the authors who attempt the difficult task of estimating the systematic error, while benefiting the authors who make no such attempt to determine all of their sources of error, (an admittedly difficult task). Some authors below have reported values and later revised their results of the same experiment by as much as twenty standard deviations. Statistically speaking, the two results could not have been estimating the same parameter.

In the review of nuclear data by the International Atomic Energy Agency¹⁶, their general comment on uncertainties included a statement questioning the validity of any presently stated uncertainties of less than 0.1% for half-lives. The same criteria has also been applied here in a few cases. No half-life has been recommended with an accuracy of better than 0.1%. The rationale for this rule is that systematic errors up to ten times smaller than the total statistical uncertainty quoted could have an appreciable effect on that total uncertainty, if there were a number of such errors. Recommending values at accuracy levels of a few hundred parts-per-million (ppm) would imply that all potential errors in the experiment at the level of ten ppm had been investigated, documented and their effect on the result taken into account. An experiment, in which such a thorough study has been both performed and documented, has yet to be reopored to my knowledge. In addition, many of these very precise results are based on the examination of only one sample.

The recommended data are given in the following tables.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Table 1. Total Half-life of ^3H

| Author | $T_{1/2}/(\text{a})$ | Comment |
|--------------------------|----------------------|--|
| Jenks ¹⁷ | 12.46 \pm 0.1 | He growth |
| Jones ¹⁸ | 12.41 +0.15 -0.25 | absolute counting |
| Jones ¹⁹ | 12.262 \pm 0.008 | He growth; precision only; rev. error |
| Popov ²⁰ | 12.57 \pm 0.18 | calorimetry |
| Merritt ²¹ | 12.31 \pm 0.13 | absolute counting |
| Jordan ²² | 12.346 \pm 0.007 | Calorim.; precision only; rev. error |
| Jones ²³ | 12.25 \pm 0.03 | He growth |
| Rudy ²⁴ | 12.323 \pm 0.008 | calorim.; precision only; rev. error |
| Unterweger ²⁵ | 12.43 \pm 0.05 | tritiated H_2O ; counting |
| Simpson ²⁶ | 12.32 \pm 0.03 | counting |
| Budick ²⁷ | 12.29 \pm 0.15 | ctg.; no details; $\sigma \times 1.5$ |
| Oliver ²⁸ | 12.38 \pm 0.03 | Neut. irradiated Li; He growth |
| Oliver ²⁹ | 12.38 \pm 0.04 | Tritiated H_2O ; He growth |

Recommended Value $12.32 \pm 0.03 \text{ a}$

Unweighted Average of Techniques

Table 2 Total Half-life of ^{10}Be

| Author | $T_{1/2}/(10^8 \text{ a})$ | Comment |
|-------------------------|----------------------------|---------------------------------|
| Hughes ³⁰ | 2.0 \pm n.u. | Revised from 2.9; not used |
| McMillian ³¹ | 2.5 \pm 0.5 | see ref. 33; not used |
| Yiou ³² | 1.55 \pm 0.3 | |
| McMillian ³³ | 1.71 \pm 0.34 | revision of 31 |
| Emery ³⁴ | 1.6 \pm 0.2 | no details; $\sigma \times 1.5$ |
| Makino ³⁵ | 1.48 \pm 0.15 | |
| Hofmann ³⁶ | 1.51 \pm 0.06 | accelerator mass spectrom. |

Weighted Average

 $1.52 \pm 0.05 \times 10^8 \text{ a}$

Recommended Value

Table 3 Total Half-life of ^{14}C

| Author | $T_{1/2}/(\text{a})$ | Comment |
|----------------------|----------------------|--|
| Libby ³⁷ | 5568 \pm 30 | Weighted Average of 3 (1949/50) values |
| Mann ¹ | 5760 \pm 50 | not used; revised; see ref. 2 |
| Watt ³⁸ | 5780 \pm 65 | mass spec., prop. ctg. |
| Olsson ³⁹ | 5680 \pm 40 | mass spec., prop. ctg. |
| Godwin ³ | 5730 \pm 40 | not used; avg. of 1, 38, 39 |
| Hughes ² | 5730 \pm 50 | revision of 1 |
| Bella ⁴⁰ | 5660 \pm 30 | |
| Emery ³⁴ | 5736 \pm 84 | no details; $\sigma \times 1.5$ |

Recommended Value $5715 \pm 30 \text{ a}$

Unweighted Average

Table 4 Total Half-life of ^{26}Al

| Author | $T_{\frac{1}{2}}/(10^3 \text{ a})$ | Comment |
|--------------------------|-------------------------------------|---------------------------------|
| Rightmire ⁺¹ | 7.1 ± 0.3 | Revised using ref. 42, 43 |
| Norris ⁺⁴ | 7.1 ± 0.2 | GeLi; Mass Spectrometry |
| Middleton ⁺⁵ | 7.0 ± 0.6 | Mass Spectrometry |
| Thomas ⁺⁸ | 7.8 ± 0.5 | not used; GeLi; verified others |
| <i>Recommended Value</i> | $7.1 \pm 0.2 \times 10^3 \text{ a}$ | <i>Weighted Average</i> |

Table 5 Total Half-life of ^{39}Ar

| Author | $T_{\frac{1}{2}}/(\text{a})$ | $\frac{\text{#}}{\text{#}}$ | Comment |
|--------------------------|------------------------------|-----------------------------|---|
| Zeldes ⁺⁷ | 265 ± 30 | | |
| Stoenner ⁺⁸ | 268 ± 8 | | Revised ^{37}Ar $T_{\frac{1}{2}}$ by Kishore ⁺⁹ |
| <i>Recommended Value</i> | $268. \pm 8. \text{ a}$ | | <i>Weighted Average</i> |

Table 6 Total Half-life of ^{40}K

| Author | $T_{1/2}/(10^9 \text{ a})$ | Comment |
|------------------------------|----------------------------|--------------------------------|
| Orban ⁵⁰ | 0.5 | not used; ec; cloud chamber |
| Gleditsch ⁵¹ | 11. ± 2. | not used; ec; GM ctr. |
| Ahrens ⁵² | 11.6 ± 0.2 | not used; ec; radiogenic |
| Graf ⁵³ | 1.48 ± 0.07 | not used; β ; GM ctr. |
| Stout ⁵⁴ | 1.29 ± 0.08 | not used; β ; GM ctr. |
| Floyd ⁵⁵ | 1.54 ± 0.39 | not used; total; GM ctr. |
| Sawyer ⁵⁶ | 12. ± 1. | not used; ec; stilbene crystal |
| Graf ⁵⁷ | 12. ± 2. | not used; ec; GM ctr. |
| Spiers ⁵⁸ | 1.18 | not used; total; ion cham., GM |
| Faust ⁵⁹ | 1.14 ± 0.16 | not used; total; |
| Sawyer ⁶⁰ | 1.27 ± 0.05 | not used; total; 4π ctr |
| Houtermans ⁶¹ | 1.31 ± 0.07 | not used; total; 4π ctr. |
| Smaller ⁶² | 1.76 ± 0.05 | not used; β ; KI crystal |
| Delaney ⁶³ | 1.24 ± 0.01 | not used; β |
| Good ⁶⁴ | 1.46 ± 0.03 | β ; KI crystal |
| Burch ⁶⁵ | 11.7 ± 0.5 | not used; ec; ion chamber |
| Suttle ⁶⁶ | 1.34 ± 0.03 | not used; β |
| | 13.4 ± 0.2 | not used; ec |
| Kono ⁶⁷ | 1.36 ± 0.05 | β ; KI crystal |
| Backenstoss ⁶⁸ | 11.3 ± 0.5 | not used; ec; NaI crystal |
| McNair ⁶⁹ | 1.44 ± 0.01 | β ; NaI crystal |
| Wetherill ⁷⁰ | 12.2 ± 0.6 | not used; ec; radiogenic |
| Wetherill ⁷¹ | 11.7 ± 0.4 | ec; radiogenic |
| Kelly ⁷² | 1.46 ± 0.03 | β ; KI crystal |
| Saha ⁷³ | 12.3 ± 0.6 | ec; NaI; stilbene |
| | 1.37 ± 0.04 | β |
| Glendenin ⁷⁴ | 1.40 ± 0.015 | β ; liquid scint. |
| Fleishman ⁷⁵ | 1.45 ± 0.4 | β ; scint. gel |
| Brinkman ⁷⁶ | 1.36 ± 0.02 | β ; liquid scint. |
| Leutz ⁷⁷ | 12.2 ± 0.3 | ec; NaI, CsI, KI |
| | 1.40 ± 0.002 | β |
| Feuerhake ⁷⁸ | 1.41 ± 0.02 | β ; scint. gel |
| DeRuytter ⁷⁹ | 12.2 ± 0.2 | ec; NaI |
| Egelkraut ⁸⁰ | 11.8 ± 0.5 | ec; KI, NaI |
| | 1.40 ± 0.07 | β |
| Venkataramaiah ⁸¹ | 1.31 ± 0.06 | not used; β |
| Gopal ⁸² | 1.13 ± 0.06 | not used; β |
| Cesana ⁸³ | 12.3 ± 0.04 | ec; GeLi |

Recommended Value

 $1.26 \pm 0.01 \times 10^9 \text{ a}$

Unweighted Average

Table 7 Total Half-life of ^{53}Mn

| Author | $T_{1/2}/(10^6 \text{ a})$ | Comment |
|--------------------------|-------------------------------------|--|
| Sheline ⁸⁴ | 2. | not used |
| Kaye ⁸⁵ | 1.9 ± 0.5 | not used |
| Hohlfelder ⁸⁶ | 10.8 ± 4.5 | not used |
| Matsuda ⁸⁷ | 2.9 ± 1.2 | not used |
| Hondo ⁸⁸ | 3.7 ± 0.2 | revised; mass spec., spec. act. |
| Woelfle ⁸⁹ | 3.8 ± 0.6 | revised; act. σ |
| Heimann ⁹⁰ | 3.7 ± 0.4 | revised; $^{53}\text{Mn}/^{26}\text{Al}$ in meteorites |
| Weighted Average | $3.7 \pm 0.2 \times 10^6 \text{ a}$ | Recommended Value |

Table 8 Total Half-life of ^{87}Rb

| Author | $T_{1/2}/(10^{10} \text{ a})$ | Comment |
|-------------------------------|--|---|
| Orban ⁹⁰ | 4.45 | not used; Cloud chamber |
| Strassmann ⁹¹ | 4.45 | not used; Pure ^{87}Sr in Rb mica |
| Eklund ⁹² | 5.8 ± 1.0 | not used; Geiger ctr. |
| Haxel ⁹³ | 6.9 ± 0.7 | not used; Geiger ctr. |
| Kemmerich ⁹⁴ | 6.0 ± 0.6 | not used; Geiger ctr. |
| Curran ⁹⁵ | 6.15 ± 0.3 | not used; Prop. ctr. |
| Lewis ⁹⁶ | 6.0 ± 0.3 | not used; Scint. ctr. |
| Flinta ⁹⁷ | 6.2 ± 0.2 | not used; |
| MacGregor ⁹⁸ | 6.2 ± 0.3 | not used; enriched ^{87}Rb |
| Geese-Baehnisch ⁹⁹ | $4.3 \pm 0.3 - 0.2$ | not used; |
| Fritze ¹⁰⁰ | 4.6 ± 0.5 | not used; Geolog. $^{87}\text{Sr}/^{87}\text{Rb}$ |
| Aldrich ¹⁰¹ | 5.0 ± 0.2 | not used; Geolog. $^{87}\text{Sr}/^{87}\text{Rb}$ |
| Libby ¹⁰² | 5.07 ± 0.2 | not used; Geiger ctr. |
| Flynn ¹⁰³ | 4.7 ± 0.1 | not used; Liq. scint. ctr. |
| Ovchinnikova ¹⁰⁴ | 5.0 ± 0.2 | not used; Geolog. $^{87}\text{Sr}/^{87}\text{Rb}$ |
| Rausch ¹⁰⁵ | 4.72 ± 0.08 | not used; 4π prop. ctg. |
| McNair ¹⁰⁶ | 5.25 ± 0.10 | not used; 4π ctg. |
| Egelkraut ¹⁰⁷ | 5.82 ± 0.1 | not used; Scint. ctr. |
| Beard ¹⁰⁸ | 5.53 ± 0.10 | not used; Scint. ctr. |
| Leutz ¹⁰⁹ | 5.80 ± 0.12 | not used; Scint. ctr. |
| Kovach ¹¹⁰ | 4.77 ± 0.10 | not used; Scint. ctr. |
| Thode ¹¹¹ | 4.60 ± 0.06 | not used; Mass spec. |
| Brinkman ⁷⁸ | 5.22 ± 0.15 | not used; |
| McMullen ¹¹² | 4.72 ± 0.04 | not used; Mass spec. |
| Neumann ¹¹³ | $4.88 \pm 0.06 - 0.10$ | 4π prop. ctg. |
| Davis ¹¹⁴ | 4.89 ± 0.04 | McMullen revised |
| Akatsu ¹¹⁵ | 5.56 ± 0.025 | not used |
| Recommended Value | $4.88 \pm 0.05 \times 10^{10} \text{ a}$ | Unweighted Average |

Table 9 Total Half-life of ^{92}Nb

| Author | $T_{\frac{1}{2}}/(10^7 \text{ a})$ | Comment |
|-------------------------|-------------------------------------|--------------------------|
| Apt ¹¹⁶ | 17 | not used |
| Makino ⁵ | 3.5 ± 0.4 | revised |
| Nethaway ⁸ | 3.9 ± 0.5 | revised |
| <i>Weighted Average</i> | $3.7 \pm 0.5 \times 10^7 \text{ a}$ | <i>Recommended Value</i> |

Table 10 Total Half-life of ^{129}I

| Author | $T_{\frac{1}{2}}/(10^7 \text{ a})$ | Comment |
|--------------------------|-------------------------------------|---------------------------------|
| Katcoff ¹¹⁷ | 1.72 ± 0.09 | prop. ctr., mass spec. |
| Russel ¹¹⁸ | 1.56 ± 0.06 | |
| Emery ¹¹⁹ | 1.57 ± 0.06 | No details; $\sigma \times 1.5$ |
| Kuhry ¹²⁰ | 1.97 ± 0.14 | Liq. scint. |
| <i>Recommended Value</i> | $1.7 \pm 0.1 \times 10^7 \text{ a}$ | <i>Unweighted Average</i> |

Table 11 Total Half-life of ^{138}La

| Author | $T_{1/2}/(10^{11} \text{ a})$ | Comment |
|---------------------------|--|----------------------------------|
| Turchinetz ¹²¹ | 1.15 ± 0.1 | not used; revised |
| Glover ¹²² | 1.13 ± 0.04 | not used; revised |
| DeRuytter ¹²³ | 1.04 ± 0.02 | not used |
| Ellis ¹²⁴ | 1.53 ± 0.3 | not used; revised; GeLi |
| Marsol ¹²⁵ | 1.23 ± 0.18 | revised; GeLi |
| Cesana ⁸³ | 1.25 ± 0.12 | not used; revised; GeLi |
| Taylor ¹²⁶ | 1.25 ± 0.12 | revised; GeLi |
| Sato ¹²⁷ | 1.03 ± 0.04 | GeLi |
| Norman ¹²⁸ | 1.05 ± 0.05 | revised; GeLi |
| Masuda ¹²⁹ | 2.5 ± 0.2 | not used; β^- ; radiogenic |
| <i>Weighted Average</i> | $1.06 \pm 0.04 \times 10^{11} \text{ a}$ | <i>Recommended Value</i> |

Table 12 Total Half-life of ^{147}Sm

| Author | $T_{1/2}/(10^{11} \text{ a})$ | Comment |
|----------------------------|--|-------------------------------------|
| Hevesy ¹³⁰ | 1.8 | not used; Geiger ctr. |
| Herzfinkiel ¹³¹ | 2.0 | not used; ion chamber |
| Mader ¹³² | 1.5 | not used; Ion chamber |
| Libby ¹³³ | 0.91 | not used |
| Hosemann ¹³⁴ | 1.5 ± 0.1 | not used; geiger ctr. |
| Cuer ¹³⁵ | 1.3 ± 0.1 | not used; nucl. emulsion |
| Picciotto ¹³⁶ | 0.99 ± 0.05 | not used; nucl. emulsion |
| Beard ¹³⁷ | 1.25 ± 0.06 | not used; 4π geiger ctr. |
| Leslie ¹³⁸ | 1.15 ± 0.03 | not used |
| Beard ¹³⁹ | 1.06 ± 0.04 | liq. scint.; corrected for wrong Sm |
| Karras ¹⁴⁰ | 1.13 ± 0.05 | not used; ion chamber |
| Mac Farlane ¹⁴¹ | 1.15 ± 0.05 | not used |
| Wright ¹⁴² | 1.05 ± 0.02 | not used; liquid scintillator |
| Donhofffer ¹⁴³ | 1.04 ± 0.03 | not used; Liquid scintillator |
| Valli ¹⁴⁴ | 1.08 ± 0.02 | not used; Ion.chamber,liquid scint. |
| Gupta ¹⁴⁵ | 1.06 ± 0.02 | 97% enriched |
| Al-Bataina ¹⁴⁶ | 1.05 ± 0.04 | 97.5% enriched |
| <i>Weighted Average</i> | $1.06 \pm 0.02 \times 10^{11} \text{ a}$ | <i>Recommended Value</i> |

Table 13 Total Half-life of ^{176}Lu

| Author | $T_{\frac{1}{2}}/(10^{10} \text{ a})$ | Comment |
|----------------------------|---------------------------------------|--------------------------------------|
| Heyden ¹⁴⁷ | 4. | not used; GM ctr. |
| Libby ¹⁴⁸ | $7.3 \pm 2.$ | not used; GM ctr. |
| Flammerfeld ¹⁴⁹ | 2.4 | not used; GM ctr. |
| Arnold ¹⁵⁰ | 2.15 ± 0.10 | not used; NaI |
| Dixon ¹⁵¹ | 4.56 ± 0.3 | not used; prop. ctr. |
| Glover ¹⁵² | 2.1 ± 0.2 | not used; NaI |
| Herr ¹⁵³ | 2.17 ± 0.35 | not used; radiogenic |
| Mc Nair ¹⁵⁴ | 3.6 ± 0.1 | not used; NaI |
| Brinkman ⁷⁸ | 3.59 ± 0.05 | not used; NaI, $\beta\gamma$ coin. |
| Donhoffer ¹⁴³ | 2.18 ± 0.06 | not used; liquid scint. |
| Sakamoto ¹⁵⁵ | 5.0 ± 0.3 | not used; NaI |
| Prodi ¹⁵⁶ | 3.27 ± 0.05 | not used; liquid scint. |
| Boudin ¹⁵⁷ | 3.3 ± 0.5 | not used; radiogenic |
| Komura ¹⁵⁸ | 3.79 ± 0.03 | 71% enriched; GeLi, NaI |
| Norman ¹⁵⁹ | 4.08 ± 0.24 | GeLi |
| Sguigna ¹⁶⁰ | 3.59 ± 0.05 | 54.4% enriched; $\gamma\gamma$ coin. |
| Patchett ¹⁶¹ | 3.57 ± 0.14 | radiogenic |
| Sato ¹⁶² | 3.78 ± 0.02 | GeLi |

Recommended Value $3.8 \pm 0.1 \times 10^{10} \text{ a};$ Unweighted Average

Table 14 Total Half-life of ^{174}Hf

| Author | $T_{\frac{1}{2}}/(10^{15} \text{ a})$ | Comment |
|----------------------------|---|-----------------|
| Riezler ¹⁶³ | 4.3. n.u. | natural sample |
| Mac Farlane ¹⁴¹ | 2.0 ± 0.4 | 10.14% enriched |
| Recommended Value | $2.0 \pm 0.4 \times 10^{15} \text{ a};$ | Selected Value |

Table 15 Total Half-life of ^{180}Ta

| Author | $T_{\frac{1}{2}}/(10^{15} \text{ a})$ | Comment |
|--------------------------|---------------------------------------|-------------------------|
| Eberhardt ¹⁶⁴ | > 0.00099 | β^- branch |
| Bauminger ¹⁶⁵ | $> 0.023 \pm 0.007$ | electron capture branch |
| | $> 0.017 \pm 0.006$ | β^- branch |
| Eberhardt ¹⁶⁶ | > 0.0000046 | K capture branch |
| Sakamoto ¹⁵⁵ | $> 0.015 \pm 0.005$ | electron capture branch |
| Ardisson ¹⁶⁷ | > 0.021 | electron capture branch |
| Norman ¹⁶⁸ | > 0.056 | elect. capt. branch |
| | > 0.056 | β^- branch |
| Cumming ¹⁶⁹ | > 1.2 | |
| Recommended Value | $> 1.2 \times 10^{15} \text{ a};$ | Selected Value |

Table 16 Total Half-life of ^{186}Os

| Author | $T_{1/2}/(10^{15} \text{ a})$ | Comment |
|----------------------|--|-----------------|
| Viola ¹⁷⁰ | 2.0 ± 1.1 | 61.27% enriched |
| Recommended Value | $2.0 \pm 1.1 \times 10^{15} \text{ a}$ | Selected Value |

Table 17 Total Half-life of ^{187}Re

| Author | $T_{1/2}/(10^{11} \text{ a})$ | Comment |
|---------------------------|--|-------------------------------------|
| Naldrett ¹⁷¹ | $40. \pm 10.$ | not used; geiger ctr. |
| Sugarman ¹⁷² | $40. \sim 70.$ | not used; geiger ctr. |
| Dixon ¹⁷³ | $> 1000.$ | not used; prop. ctr. |
| Suttle ¹⁷⁴ | $> 1.$ | not used; geiger ctr. |
| Herr ¹⁷⁵ | $0.05\text{--}2.5$ | not used; radiogenic |
| Herr ¹⁷⁶ | ~ 0.8 | not used; radiogenic |
| Walton ¹⁷⁷ | 2.1 ± 0.5 | not used; geiger ctr |
| Naldrett ¹⁷⁸ | 3.2 ± 0.7 | not used; geiger ctr. |
| Herr ¹⁷⁹ | 0.62 ± 0.07 | not used; radiogenic |
| Kocol ¹⁸⁰ | 0.79 | not used; geiger ctr., only 1 meas. |
| Wolf ¹⁸¹ | 1.2 ± 0.4 | not used; geiger ctr. |
| Hirt ¹⁸² | 0.43 ± 0.05 | radiogenic |
| Brodzinski ¹⁸³ | 0.66 ± 0.13 | not used; prop. ctr. |
| Watt ¹⁸⁴ | ~ 0.3 | not used; low backgd. meas. |
| Luck ¹⁸⁵ | 0.45 ± 0.02 | radiogenic |
| Naldrett ⁸ | 0.35 ± 0.04 | liquid scintillator |
| Lindner ¹⁸⁶ | 0.435 ± 0.013 | mass spectrometry |
| Recommended Value | $4.2 \pm 0.2 \times 10^{10} \text{ a}$ | Unweighted Average |

Table 18 Total Half-life of ^{190}Pt

| Author | $T_{1/2}/(10^{11} \text{ a})$ | Comment |
|----------------------------|--|-----------------------------|
| Hoffmann ¹⁸⁷ | 5. | not used |
| Porschen ¹⁸⁸ | 10. | not used; nuclear emulsion |
| Mac Farlane ¹⁴¹ | 6.9 ± 0.5 | 0.76% enriched |
| Petrzhak ¹⁸⁹ | 4.7 ± 1.7 | Natural Pt.; ion chamber |
| Graeffe ¹⁹⁰ | 5.4 ± 0.6 | natural + enriched Platinum |
| Al-Bataina ¹⁴⁸ | 6.65 ± 0.28 | natural Platinum |
| Weighted Average | $6.5 \pm 0.3 \times 10^{11} \text{ a}$ | Recommended Value |

Table 19 Total Half-life of ^{204}Pb

| Author | $T_{1/2}/(10^{17} \text{ a})$ | Comment |
|-----------------------|-------------------------------|---|
| Kohman ¹⁹¹ | ≥ 0.3 | Slight indication of activity |
| Riezler ⁹ | 1.4 | $E\alpha = 2.6 \text{ Mev} > \text{available energy}$ |

Recommended Value Stable

Table 20 Total Half-life of ^{210}Pb

| Author | $T_{1/2}/(\text{a})$ | Comment |
|-----------------------------|----------------------|-------------------------------------|
| Antonoff ¹⁹² | 16.5 | not used; ZnS ctg. |
| Albrecht ¹⁹³ | 22.5 ± 0.4 | not used |
| Curie ¹⁹⁴ | 22. | not used |
| Joliot-Curie ¹⁹⁵ | 23. | not used; α ctg. |
| Wagner ¹⁹⁶ | 25. | not used; ion chamb.; 0.7 year ctg. |
| Tobailem ¹⁹⁷ | 19.40 ± 0.35 | not used; 1/3 year ctg. |
| Merritt ¹⁹⁸ | 22.4 ± 0.4 | 4π prop. ctr., 5½ years ctg. |
| Harbottle ¹⁹⁹ | 20.4 ± 0.3 | ion chamber; 3/4 year ctg. |
| Pate ²⁰⁰ | 23.3 ± 0.5 | 4π prop. ctr., 5 years ctg. |
| Eckelmann ²⁰¹ | 21.4 ± 0.5 | geol. |
| Imre ²⁰² | 22.85 ± 0.70 | β counting |
| Ramthun ²⁰³ | 21.96 ± 0.51 | calorimetry |
| Von Gunten ²⁰⁴ | 22.2 ± 1.0 | prop. counter |
| Hoehndorf ²⁰⁵ | 22.26 ± 0.11 | α spectrometry |

Recommended Value $22.6 \pm 0.1 \text{ a};$ Unweighted AverageTable 21 Total Half-life of ^{210}Po

| Author | $T_{1/2}/(\text{d})$ | Comment |
|-----------------------------|----------------------|--------------------------------|
| Schweidler ²⁰⁶ | 136.5 | counted 6 years |
| Curie ²⁰⁷ | 140. | γ counted 4 year |
| Dorabialska ²⁰⁸ | 137.8 ± 0.6 | calorimetry, α counting |
| Sanielevici ²⁰⁹ | 138.7 ± 0.6 | calorimetry |
| Beamer ²¹⁰ | 138.3 ± 0.14 | calorimetry |
| Ginnings ²¹¹ | 138.39 ± 0.14 | calorimetry |
| Curtis ²¹² | 138.37 ± 0.098 | α ctg.; revised error |
| Eichelberger ²¹³ | 138.376 ± 0.05 | calorimetry; revised error |

Recommended Value $138.4 \pm 0.1 \text{ d};$ Weighted Average with Uncertainty Rule

Table 22 Total Half-life of ^{222}Rn

| Author | $T_{1/2}$ (d) | Comment |
|----------------------------|----------------------|---|
| Bothe ²¹⁴ | 3.825 \pm 0.004 | revised error |
| Curie ²¹⁵ | 3.823 \pm 0.003 | revised error |
| Tobailem ²¹⁶ | 3.825 \pm 0.006 | ion chamber; revised error |
| Marin ²¹⁷ | 3.8229 \pm 0.0017 | Counted 5½ $T_{1/2}$; revised error |
| Robert ²¹⁸ | 3.825 \pm 0.005 | revised error |
| Shimanskaja ²¹⁹ | 3.83 \pm 0.03 | Calorimetry |
| Butt ²²⁰ | 3.82351 \pm 0.0017 | Nal; counted 40 $T_{1/2}$; revised error |
| <i>Recommended Value</i> | 3.823 ± 0.004 d; | <i>Weighted Average with Uncertainty Rule</i> |

Table 23 Total Half-life of ^{224}Th

| Author | $T_{1/2}$ (s) | Comment |
|-------------------------|--------------------|--------------------------|
| Tove ²²¹ | 1.05 \pm 0.05 | Scintillation detector |
| Valli ²²² | 1.03 \pm 0.05 | semi-conductor |
| Ibowksi ²²³ | 1.05 \pm 0.02 | α spectrometry |
| <i>Weighted Average</i> | 1.05 ± 0.02 s; | <i>Recommended Value</i> |

Table 24 Total Half-life of ^{226}Ra

| Author | $T_{1/2}$ (a) | Comment |
|--------------------------|-------------------|-----------------------------------|
| Watson ²²⁴ | 1608. | not used; calorimetry |
| Braddick ²²⁵ | 1603. | not used; α current |
| Curie ²²⁶ | 1590. | not used; ion current |
| Ward ²²⁷ | 1599. | not used; no. α 's emitted |
| Meitner ²²⁸ | 1590. | not used; calorimetry |
| Gleditsch ²²⁹ | 1691. | not used; growth rate |
| Guenther ²³⁰ | 1603. | not used; He prod. |
| Kohman ²³¹ | 1622. \pm 13. | no. α 's emitted |
| Gorshkov ²³² | 1573. | not used; calorimetry |
| Sebaoun ²³³ | 1617. \pm 12. | no. α 's emitted |
| Gorshkov ²³⁴ | 1577. \pm 9. | calorimetry |
| Martin ²³⁵ | 1602. \pm 8. | calorimetry |
| Ramthun ²³⁶ | 1599. \pm 7. | calorimetry |
| <i>Weighted Average</i> | $1599. \pm 4.$ a; | <i>Recommended Value</i> |

Table 25 Total Half-life of ^{227}Ac

| Author | $T_{1/2}/(\text{a})$ | Comment |
|-----------------------------|----------------------|----------------------------|
| Joliot-Curie ²³⁷ | 21.7 | not used |
| Hollander ²³⁸ | 22.0 \pm 0.3 | ion chamber |
| Tobailem ²³⁹ | 21.8 \pm 0.4 | ion chamber |
| Shimanskaya ²⁴⁰ | 21.2 \pm 0.8 | calorimetry |
| Robert ²⁴¹ | 21.6 \pm 0.3 | calorimetry |
| Jordan ²⁴² | 21.773 \pm 0.012 | calorimetry; revised error |

Recommended Value $21.77 \pm 0.02 \text{ a};$ Weighted Average with Uncertainty Rule

Table 26 Total Half-life of ^{228}Ra

| Author | $T_{1/2}/(\text{a})$ | Comment |
|-----------------------|----------------------|----------|
| Curie ²⁴³ | 6.7 | not used |
| Dudley ²⁴⁴ | 5.7 \pm 0.2 | |
| Mays ²⁴⁵ | 5.75 \pm 0.03 | |

Weighted Average $5.75 \pm 0.03 \text{ a};$ Recommended Value

Table 27 Total Half-life of ^{228}Th

| Author | $T_{1/2}/(\text{a})$ | Comment |
|------------------------|----------------------|---------------------------------|
| Meitner ²⁴⁶ | 1.91 \pm 0.02 | |
| Kirby ²⁴⁷ | 1.910 \pm 0.002 | α counting, 2 years |
| Mays ²⁴⁸ | 1.908 \pm 0.004 | γ counting |
| Mays ²⁴⁸ | 1.924 \pm 0.020 | α counting |
| Jordan ²⁴⁹ | 1.9131 \pm 0.0020 | Calorim.; 9.3 years; rev. error |
| Hoppe ²⁵⁰ | 1.9113 \pm 0.0021 | revised error |

Recommended Value $1.912 \pm 0.002 \text{ a};$ Weighted Average with Uncertainty Rule

Table 28 Total Half-life of ^{230}Th

| Author | $T_{1/2}/(10^4 \text{ a})$ | Comment |
|-------------------------|----------------------------|--------------------------|
| Soddy ²⁵¹ | 7.42 | Meadows recalculation |
| Soddy ²⁵² | 7.14 \pm 0.36 | Meadows recalculation |
| Soddy ²⁵³ | 7.69 \pm 0.30 | Meadows recalculation |
| Curie ²⁵⁴ | 8.23 \pm 0.25 | not used |
| Soddy ²⁵⁵ | 7.32 \pm 0.37 | Meadows recalculation |
| Hyde ¹² | 7.99 \pm 0.34 | 26.4% enriched |
| Attree ¹³ | 7.61 \pm 0.14 | 12.11% enriched |
| Meadows ¹⁴ | 7.538 \pm 0.030 | 99.65% enriched |
| <i>Weighted Average</i> | | <i>Recommended Value</i> |

Table 29 Total Half-life of ^{232}Th

| Author | $T_{1/2}/(10^{10} \text{ a})$ | Comment |
|-------------------------|-------------------------------|-----------------------------------|
| Kovarik ²⁵⁶ | 1.39 \pm 0.03 | |
| Senftle ²⁵⁷ | 1.42 \pm 0.07 | Na(I) |
| Piciotto ²⁵⁸ | 1.39 \pm 0.03 | nuclear emulsion |
| Macklin ²⁵⁹ | 1.45 \pm 0.05 | incidental to cross section meas. |
| Farley ²⁶⁰ | 1.41 \pm 0.014 | ion-chamber α spectrometry |
| LeRoux ²⁶¹ | 1.40 \pm 0.007 | liquid scintillator |
| <i>Weighted Average</i> | | <i>Recommended Value</i> |

Table 30 Total Half-life of ^{231}Pa

| Author | $T_{1/2}/(10^4 \text{ a})$ | Comment |
|---------------------------|----------------------------|-----------------------------|
| Von Grosse ²⁶² | 3.2 \pm 0.3 | not used |
| Van Winkle ²⁶³ | 3.43 \pm 0.03 | not used; α counting |
| Kirby ²⁶⁴ | 3.248 \pm 0.026 | calorimetry |
| Brown ²⁶⁵ | 3.234 \pm 0.023 | α counting |
| Robert ²⁶⁶ | 3.278 \pm 0.011 | Calorimetry |
| <i>Recommended Value</i> | | <i>Unweighted Average</i> |

References

1. W. B. Mann, W. F. Marlow, E. E. Hughes, Appl. Radiat. Isot. **11**, 57 (1961).
2. E. E. Hughes, W. B. Mann, Appl. Radiat. Isot. **15**, 97 (1964).
3. H. Godwin, Nature **195**, 984 (1962).
4. F. Johnson, Science **149**, 1326 (1965).
5. T. Makino, M. Hondo, Geochim. Cosmochim. Acta **41**, 1521 (1977).
6. D. R. Nethaway, A. L. Prindle, R. A. Van Konynenberg, Phys. Rev./C **17**, 1409 (1978).
7. D. R. Nethaway, J. Inorg. Nucl. Chem. **40**, 1285 (1978).
8. S. N. Naldrett, Can. J. Phys. **62**, 15 (1984).
9. W. Riezler, G. Kauw, Z. Naturforschg. **13a**, 904 (1958).
10. H. Faraggi, Ann. Phys. (Paris) **6**, 325 (1951).
11. B. Al-Bataina, J. Jaenecke, Phys. Rev./C **37**, 1667 (1988).
12. E. K. Hyde, "The Transuranium Elements", National Nuclear Energy Series, Pt. II, 1435 (1949), McGraw-Hill Publ. - New York.
13. R. W. Attree, M. J. Cabell, R. L. Cushing, J. J. Pieroni, Can. J. Phys. **40**, 194 (1962).
14. J. W. Meadows, R. J. Armani, E. L. Callis, A. M. Essling, Phys. Rev./C **22**, 750 (1980).
15. N. E. Holden, Pure Appl. Chem. **61**, 1483 (1989).
16. C. W. Reich, R. Vaninbroukx, Int. Atomic Energy Agency Report, IAEA-TEC-DOC-336, 275 (1985).
17. G. H. Jenks, F. H. Sweeton, J. A. Ghormley, Phys. Rev. **80**, 990 (1950).
18. W. M. Jones, Phys. Rev. **83**, 537 (1951).
19. W. M. Jones, Phys. Rev. **100**, 124 (1955).
20. M. M. Popov, U. V. Gagarinskii, M. D. Sevin, I. P. Mikhaleko, Yu. M. Morozov, At. Energia **4**, 296 (1958).
21. J. S. Merritt, J. G. V. Taylor, Chalk River Report AECL-2510 (1966).
22. K. C. Jordan, B. C. Blanke, W. A. Dudley, J. Inorg. Nucl. Chem. **29**, 2129 (1967).
23. P. M. S. Jones, J. Nucl. Materials **21**, 239 (1967).
24. C. R. Rudy, K. C. Jordan, Mound Facility Report MLM-2458 (Dec. 22, 1977).
25. M. P. Unterweger, B. M. Coursey, F. J. Schima, W. B. Mann, Appl. Radiat. Isot. **31**, 611 (1980).
26. J. J. Simpson, Phys. Rev./C **35**, 752 (1987).
27. B. Budick, H. Lin, Bull. Am. Phys. Soc. **32**, 1063 (1987).
28. B. M. Oliver, H. Farrar IV, M. M. Bretscher, Appl. Radiat. Isot. **38**, 959 (1987).
29. B. M. Oliver, M. M. Bretscher, H. Farrar IV, Appl. Radiat. Isot. **40**, 199 (1989).
30. D. J. Hughes, C. Eggler, C. M. Huddleston, Phys. Rev. **71**, 269 (1947).
31. E. M. McMillian, Phys. Rev. **72**, 591 (1947).
32. F. Yiou, G. M. Raisbeck, Phys. Rev. Lett. **29**, 372 (1972).
33. E. M. McMillian, Phys. Rev./C **6**, 2296 (1972).
34. J. F. Emery, S. A. Reynolds, E. I. Wyatt, G. I. Gleason, Nucl. Sci. Eng. **48**, 319 (1972).
35. T. Makino, R. Gensho, M. Honda, Shitsuryo Bunseki (Mass Spectroscopy) **23**, 33 (1975).
36. H. J. Hofmann, J. Beer, G. Bonani, H. R. Von Gunten, S. Raman, M. Suter, R. L. Walker, W. Woelfli, D. Zimmermann, Nucl. Inst. Meth. **B29**, 32 (1987).

37. W. F. Libby, "Radiocarbon Dating", 2nd Ed., Univ. Chicago Press (1955).
38. D. E. Watt, D. Rameden, H. W. Wilson, Appl. Radiat. Isot. **11**, 68 (1961).
39. I. U. Olsson, I. Karlen, A. H. Turnbull, N. J. D. Prosser, Ark. f. Fys. **22**, 237 (1962).
40. F. Bella, M. Alessio, P. Fratelli, Nuovo Cim. **58B**, 232 (1968).
41. R. A. Rightmire, T. P. Kohman, H. Hintenberger, Z. Naturforsch. **13a**, 847 (1958).
42. E. A. Samworth, E. K. Warburton, G. A. P. Engelbertink, Phys. Rev./C **5** 138 (1972).
43. I. Sykora, P. Povinec, Nucl. Inst. Meth. **B27**, 467 (1986).
44. T. L. Norris, A. J. Gancarz, D. J. Rokop, K. W. Thomas, J. Geophys. Res. **88** (supplement) B333 (1983).
45. R. Middleton, J. Klein, G. M. Raisbeck, F. Yiou, Nucl. Inst. Meth. **218**, 430 (1983).
46. J. H. Thomas, R. L. Rau, R. T. Skelton, R. W. Kavanagh Phys. Rev./C **30** 385 (1984).
47. H. Zeldes, B. H. Ketelle, A. R. Brosi, C. R. Fultz, R. F. Hibbs, Phys. Rev. **86**, 811 (1952).
48. R. W. Stoermer, O. A. Schaeffer, S. Katcoff, Science **148**, 1325 (1965).
49. R. Kishore, R. Colle, S. Katcoff, J. B. Cumming, Phys. Rev./C **12**, 21 (1975).
50. G. Orban, Sitzb. Akad. Wiss. Wien Abt.IIA, **140**, 121 (1931).
51. E. Gleditsch, T. Graf, Phys. Rev. **72**, 640 (1947).
52. L. H. Ahrens, R. D. Evans, Phys. Rev. **74**, 279 (1948).
53. T. Graf, Phys. Rev. **74**, 831 (1948).
54. R. W. Stout, Phys. Rev. **75**, 1107 (1949).
55. J. J. Floyd, L. B. Borst, Phys. Rev. **75**, 1106 (1949).
56. G. A. Sawyer, M. L. Wiedenbeck, Phys. Rev. **76**, 1535 (1950).
57. T. Graf, Rev. Sci. Inst. **21**, 285 (1950).
58. F. W. Spiers, Nature **165**, 356 (1950).
59. W. R. Faust, Phys. Rev. **78**, 624 (1950).
60. G. A. Sawyer, M. L. Wiedenbeck, Phys. Rev. **79**, 490 (1950).
61. F. G. Houtermans, O. Haxel, J. Heintze, Z. Physik **128**, 657 (1950).
62. B. Smaller, J. May, M. Freedman, Phys. Rev. **79**, 940 (1950).
63. C. F. G. Delaney, Phys. Rev. **81**, 158 (1951).
64. M. L. Good, Phys. Rev. **81**, 891 (1951).
65. P. R. J. Burch, Nature **172**, 361 (1953).
66. A. D. Suttle, W.F. Libby, Anal. Chem. **27**, 921 (1955).
67. S. Kono, J. Phys. Soc. Japan **10**, 495 (1955).
68. G. Backenstoss, K. Goebel, Z. Naturforsch. **10a**, 920 (1955).
69. A. McNair, R. N. Grover, H. W. Wilson, Phil. Mag. **1**, 199 (1956).
70. G. W. Wetherill, G. J. Wasserberg, L. T. Aldrich, G. R. Tilton, R. J. Hayden, Phys. Rev. **103**, 987 (1956).
71. G. W. Wetherill, Science **126**, 545 (1957).
72. W. H. Kelly, G. B. Beard, R. A. Peters, Nucl. Phys. **11**, 492 (1959).
73. N. K. Saha, J. B. Gupta, Proc. Natl. Inst. Sci. India **26** No. 5 (1960).
74. L. E. Glendenin, Ann. N.Y. Acad. Sci. **91**, 166 (1961).
75. D. G. Fleishman, V. V. Glazunov, Sov. At. En. **12**, 338 (1962).
76. G. A. Brinkman, A. H. W. Aten, Jr., J. Th. Veenboer, Physica **31**, 1305 (1965).
77. H. Leutz, G. Schulz, H. Wenninger, Z. Phys. **187**, 151 (1965).

78. I. Feuerhake, A. Hinze, Naturwiss. **53**, 272 (1966).
79. A. W. DeRuytter, A. H. W. Aten, Jr., A. Van Dulmen, C. Krol-Konig, E. Zuidema, Physica **32**, 991 (1966).
80. K. Egelkraut, H. Leutz, Physik Verhandl. **11**, 67 (1966).
81. P. Venkataramiah, H. Sanjeevaiah, B. Sanjeevaiah, Ind. J. Pure Appl Phys. **9**, 133 (1971).
82. K. Gopal, H. Sanjeevaiah, B. Sanjeevaiah, Am. J. Phys. **40**, 721 (1972).
83. A. Cesana, M. Terrani, Anal. Chem. **49**, 1156 (1977).
84. R. K. Sheline, J. E. Hooper, Nature **179**, 85 (1957).
85. J. H. Kaye, P. J. Cressy, J. Inorg. Nucl. Chem. **27**, 1889 (1965).
86. J. J. Hohlfelder, Phys. Rev. **186**, 1126 (1969).
87. H. Matsuda, S. Umemoto, M. Hondo, Radiochimica Acta **15**, 51, (1971).
88. M. Hondo, M. Imamura, Phys. Rev./C **2**, 1182 (1971).
89. R. Woelfle, W. Herr, U. Herpers, Radiochimica Acta **18**, 207 (1972).
90. M. Heimann, P. P. Parekh, W. Herr, Geochim. Cosmochim. Acta **38**, 217 (1974).
91. F. Strassmann, E. Walling, Ber. d. dtach. Chem. Ges. **71B**, 1 (1938)
92. S. Eklund, Arkiv Mat. Astron. Fysik **A33**, #14 (1946)
93. O. Haxel, F. G. Houtermans, Z. Phys. **124**, 705 (1948); see also O. Haxel, F. G. Houtermans, M. Kemmerich, Phys. Rev. **74**, 1886 (1948).
94. M. Kemmerich, Z. Phys. **126**, 399 (1949).
95. S. C. Curran, D. Dixon, H. W. Wilson, Phys. Rev. **84**, 151 (1951).
96. G. M. Lewis, Phil. Mag. **43**, 1070 (1952).
97. J. Flinta, S. E. Klund, Arkiv Fysik **7**, 401 (1953).
98. M. H. MacGregor, M. L. Wiedenbeck, Phys. Rev. **94**, 138 (1954).
99. I. Geese-Baehnisch, E. Huster, Naturwiss. **21**, 495 (1954); see also I. Geese-Baehnisch, Z. Physik, **142**, 565 (1955).
100. K. Fritze, F. Strassmann, Z. Naturforschg. **11a**, 277 (1956).
101. L. T. Aldrich, G. W. Wetherill, G. R. Tilton, G. L. Davis, Phys. Rev., **103**, 1045 (1956).
102. W. F. Libby, Anal. Chem. **29**, 1566, (1957).
103. K. F. Flynn, L. E. Glendenin, Phys. Rev., **116**, 744 (1959)
104. G. V. Ovchinnikova, Geochemica, **5**, 392 (1959).
105. W. Rausch, W. Schmidt, Phys. Verhandl **11**, 66 (1960).
106. A. McNair, H. W. Wilson, Phil. Mag. **6**, 563 (1961).
107. K. Egelkraut, H. Leutz, Z. Physik, **161**, 13,(1961).
108. G. B. Beard, W. H. Kelly, Nucl. Phys. **28**, 570 (1961).
109. H. Leutz, H. Wenninger, K. Ziegler, Z. Physik, **169**, 409 (1962).
110. A. Kováč, Acta Phys. Hung. **17**, 341 (1964).
111. H. G. Thode, McMaster Univ. Reactor Research Report NP-15037, B27 (1965)
112. C. C. McMullen, K. Fritze, R. H. Tomlinson, Can. J. Phys. **44**, 3033 (1966).
113. W. Neumann, E. Huster, Z. Physik **270**, 121 (1971).
114. D. W. Davis, J. Gray, G. I. Cumming, H. Baadsgaard, Geochim. Cosmochim. Acta **41**, 1715 (1977).
115. E. Akatsu, Radioisotopes **30** 6-17 (1981).
116. K. E. Apt, J. D. Knight, D. C. Camp, R. W. Perkins, Geochim. Cosmochim. Acta **38**, 1185 (1974).
117. S. Katcoff, O. A. Schaeffer, J. M. Hastings, Phys. Rev. **82**, 688 (1951).

118. H. T. Russel, Oak Ridge National Laboratory Report ORNL-2293 (1957).
119. J. F. Emery, S. A. Reynolds, E. I. Wyatt, Nucl. Sci. Eng. 18, 319 (1972).
120. J. G. Kuhr, G. Bontems, Radiochem. Radioanal. Lett. 15, 29 (1973).
121. W. Turchinetz, R. W. Pringle, Phys. Rev. 103, 1000 (1956).
122. R. N. Glover, D. E. Watt, Phil Mag. 2, 49 (1957).
123. A. W. DeRuytter, A. H. W. Aten, Jr., A. Van Dulmen, C. Krol-Konig, E. Zuidema, Physica 32, 591 (1966).
124. J. L. Ellis, H. E. Hall Jr., Nucl. Phys., A179, 540 (1972).
125. C. Marsol, F. Armanet, G. Ardisson, Compt. Rend. 274, 904 (1972).
126. H. W. Taylor, R. J. Bauer, J. Phys. Soc. Japan 47, 1395 (1979).
127. J. Sato, J. Hirose, Radiochem. Radioanal. Lett. 46, 145 (1981).
128. E. B. Norman, M. A. Nelson, Phys. Rev./C 27, 1321 (1983).
129. A. Masuda, H. Shimizu, S. Nakai, A. Makishiwa, S. Rahti, Planet Sci. Lett. 89, 316 (1988).
130. G. Hevesy, M. Pahl, R. Hosemann, Z. Physik 83, 43 (1933).
131. M. Herzfinkiel, A. Wroneberg, Compt. Rend. 199, 133 (1934).
132. M. Mader, Z. Physik 88, 601 (1934).
133. W. F. Libby, Phys. Rev. 16, 196 (1934).
134. R. Hosemann, Z. Physik 99, 405 (1936).
135. P. Cuer, G. M. G. Lattes, Nature 158, 197 (1946).
136. E. Picciotto, Compt. Rend. 229, 117 (1949).
137. G. Beard, M. L. Wiedenbeck, Phys. Rev. 95, 1245 (1954).
138. G. E. Leslie, (M. S. Thesis North Carolina State College 1954. see also Nucl. Sci. Abst. 10, 1099 (1956).
139. G. Beard, W. H. Kelly, Nucl. Phys. 8, 207 (1958).
140. M. Karras, M. Nurmia, Nature 185, 601 (1960).
141. R. D. MacFarlane, T. P. Kohman, Phys. Rev. 121, 1758 (1961).
142. P. M. Wright, E. P. Steinberg, L. E. Glendenin, Phys. Rev. 123, 205 (1961).
143. D. Donhofer, Nucl. Phys. 50, 489 (1964).
144. K. Valli, J. Aaltonen, G. Graeffe, M. Nurmia, R. Poeyhuonen, Ann. Acad. Sci. Fenn. Ser. A, No. 177 (1965).
145. M. C. Gupta, R. D. MacFarlane, J. Inorg. Nucl. Chem. 32, 3425 (1970).
146. B. Al-Bataina, J. Jaenecke, Radiochim. Acta 42, 159 (1987).
147. M. Heyden, W. Wefelmeier, Naturwiss. 26, 612 (1938).
148. W. F. Libby, Phys. Rev. 56, 21 (1939).
149. A. Flammersfeld, Z. Naturforschg. 2a, 86 (1947); see also A. Flammersfeld, J. Mattauch, Naturwiss. 31, 66 (1943).
150. J. R. Arnold, Phys. Rev. 92, 7-13 (1954).
151. D. Dixon, A. McNair, S. C. Curran, Phil. Mag. 15, 683 (1954).
152. R. N. Glover, D. E. Watt, Phil. Mag. 2, 699 (1957).
153. W. Herr, E. Merz, Z. Nat. 13a, 268 (1958).
154. A. McNair, Phil. Mag., 6, 851 (1961).
155. K. Sakamoto, Nucl. Phys. A103, 134 (1967).
156. V. Prodi, K. F. Flynn, L. E. Glendenin, Phys. Rev. 188, 1930 (1969).
157. A. Boudin, S. Deutsch, Science, 168, 1219 (1970).
158. K. Komura, K. Sakamoto, S. Tanaka, Nucl. Phys. A198, 73 (1972).

159. E. B. Norman, Phys. Rev./C 21, 1109 (1980).
160. A. P. Sguigna, A. J. Larabee, J. C. Waddington, Can. J. Phys. 60, 361 (1982).
161. P. J. Patchett, Geochim. Cosmochim. Acta 47, 81 (1983).
162. J. Sato, Y. Ohoka, T. Hirose Radiochem. Radioanal. Lett. 58, 263 (1983).
163. W. Riezler, G. Kauw, Z. Naturforschg. 14a, 196 (1959).
164. P. Eberhardt, J. Geiss, C. Lang, W. Herr, E. Merz, Z. Naturforschg. 10a, 796 (1955).
165. E. R. Bauminger, S. G. Cohen, Phys. Rev. 110, 953 (1958).
166. P. Eberhardt, P. Signer, W. Herr, E. Merz, Z. Naturforschg. 13a, 1004 (1958).
167. G. Ardisson, Radiochem. Radioanal. Lett. 29, 7 (1977).
168. E. B. Norman, Phys. Rev./C 21, 2331 (1981).
169. J. B. Cumming, D. E. Alburger, Phys. Rev./C 31, 1494 (1985).
170. V. E. Viola Jr., C. T. Roche, M. M. Minor, J. Inorg. Nucl. Chem. 37, 11 (1975).
171. S. N. Naldrett, W. F. Libby, Phys. Rev. 73, 487 (1948).
172. N. Sugarman, H. Richter, Phys. Rev. 73, 1411 (1948).
173. D. Dixon, A. McNair, Phil. Mag. 45, 1099 (1954); see also D. Dixon, A. McNair, S. C. Curran, J. Phys. Rad. 16, 538 (1955).
174. A. D. Suttle, Jr., W. F. Libby, Phys. Rev. 95, 866 (1954).
175. W. Herr, H. Hintenberger, H. Voshage, Phys. Rev. 95, 1691 (1954).
176. W. Herr, E. Merz, Z. Naturforschg. 10a, 613 (1955).
177. J. R. Walton, Diss. Abst. 17, 2438 (1957).
178. S. N. Naldrett, Ann. N.Y. Acad. Sci. 72, 215 (1958).
179. W. Herr, E. Merz, Z. Naturforschg. 13a, 231 (1958).
180. H. Kocol, Nucl. Sci. Abst. 15, 3864 (1964).
181. C. J. Wolf, W. H. Johnston, Phys. Rev. 125, 307 (1962).
182. B. Hirt, G. R. Tilton, W. Herr, W. Hoffmeister, Helv. Phys. Acta 35, 320 (1962).
183. R. L. Brodzinski, D. C. Conway, Phys. Rev. 138, 1368 (1965).
184. D. E. Watt, R. N. Glover, Phil. Mag. 7, 105 (1962).
185. J. M. Luck, C. J. Allegre, Nature 302, 130 (1983).
186. M. Lindner, D.A. Leich, R.J. Borg, B.P. Russ, J.M. Bazan, D.S. Simons, A.R. Date, Nature 320 216 (1986).
187. G. Hoffmann, Z. Physik, 7, 254 (1921).
188. W. Porschen, W. Riezler, Z. Naturforschg. 11a, 143 (1956).
189. K. A. Petrzhak, M. I. Yakunin, Sov. Phys. JETP 14, 1265 (1962).
190. G. Graeffe, Ann. Acad. Sci. Fennicae Ser. A VI 128, 1 (1963).
191. T. P. Kohman, Proc. Conf. Nucl. Processes in Geological Settings, Williams Bay, Wisconsin, Sept. 21-23 (1953).
192. G. N. Antonoff, Phil. Mag. 19, 825 (1910).
193. E. Albrecht, Mitt. Ra. Inst. 123 Wien, Ber. IIIa 28 (1919).
194. M. Curie, I. Curie, J. Phys. Rad. 10, 385 (1929).
195. I. Curie, J. Phys. Rad. 10, 388 (1929).
196. F. Wagner, Argonne Nat'l Lab. Rpt. ANL-H90, 5 (1950).
197. J. Tobailem, J. Phys. Rad. 16, 235 (1955).
198. W. F. Merritt, P. J. Champion, R. C. Hawkings Can. J. Phys. 35, 16 (1957).

199. G. Harbottle, J. Inorg. Nucl. Chem. **12**, 6 (1959).
200. B. D. Pate, D. C. Santry, L. Yaffe, Can. J. Chem. **37**, 1000 (1959).
201. W. R. Eckelmann, W. S. Broecker, J. L. Kulp, Phys. Rev. **118**, 698 (1960)
202. L. Imre, G. Fabry, I. Dezsi Magy. Tud. Akad. Kem. Tud. Oszt. Kozlemen. **19**, 1 (1963).
203. H. Ramthun, Z. Naturforschg. **19a**, 1064 (1964).
204. H. R. von Gunten, A. Wyttensbach, H. Dulakas, J. Inorg. Nucl. Chem. **29**, 2826 (1976).
205. A. Hochndorf, Z. Naturforschg. **24a**, 612 (1969).
206. E. V. Schweidler, Verh. deutsch Phys. Ges. **13**, 539 (1912).
207. M. Curie, J. Phys. Rad. **1**, 12 (1920).
208. A. Dorabialska, Roczniki Chem. (Poland) **11**, 475 (1931).
209. A. S. Sanielevici, J. Chimie Phys. **33**, 779 (1936).
210. W. H. Beamer, W. E. Easton, J. Chem. Phys. **17**, 1298 (1949).
211. D. C. Ginnings, A. F. Ball, D. T. Vier, J. Res. NBS **50**, 75 (1953).
212. M.L. Curtis, Phys. Rev. **92**, 1489 (1953).
213. J.F. Eichelberger, K.C. Jordan, S.R. Orr, J.R. Parks, Phys. Rev. **96**, 719 (1954).
214. W. Bothe, Z. Phys. **16**, 266 (1923).
215. I. Curie, C. Chamie, J. Phys. Rad. **5**, 238 (1924).
216. J. Tobailem, Compt. Rend. (Paris) **233**, 1360 (1951).
217. J. Robert, J. Phys. Rad. **17**, 605 (1956).
218. P. C. Marin, Brit. J. Appl. Phys. **7**, 188 (1956).
219. N. S. Shimanskaja, Instr. Exptl. Techniques **2**, 283 (1956).
220. D. K. Butt, A. R. Wilson, J. Phys./A **5**, 1248 (1972).
221. P. A. Tove, Arkiv Fysik **13**, 549 (1958).
222. K. Valli, E. K. Hyde, J. Borggreen, Phys. Rev./C **1**, 2115 (1970).
223. R. Ibowksi, W. Scholz, J. Bisplinghoff, H. Ernst, J. Rama Rao, T. Mayer-Kuckuk, Juelich Report - JUEL-SPEZ-15 **19** (1978).
224. S. W. Watson, M. C. Henderson, Proc. Roy. Soc. A **118**, 318 (1928).
225. H. J. J. Braddick, H. M. Cave, Proc. Roy. Soc. A **121**, 367 (1928).
226. I. Curie, F. Joliot, Compt. Rend. (Paris) **187**, 43 (1928).
227. F. A. B. Ward, C. E. Wynn-Williams, H. M. Cave, Proc. Roy. Soc. A **125**, 713 (1929).
228. L. Meitner, W. Ortmann, Z. Phys. **60**, 143 (1930).
229. E. Gleditsch, E. Foeyn, Am. J. Sci. **29**, 253 (1935).
230. P. Guenther, Z. Phys. Chem. **A185**, 367 (1939).
231. T. P. Kohman, D. P. Ames, J. Sedlet, "The Transuranium Elements". National Nuclear Energy Series, Vol. IV-14B, 1675 (1949), McGraw-Hill Publ. - New York.
232. G. V. Gorshkov, N. S. Shimanskaya, Sov. At. En. **1** (5), 86 (1956).
233. W. Sebaoun, Ann. Phys. (Paris) **1**, 680 (1956).
234. G. V. Gorshkov, Z. G. Gretchenko T. A. Il'inskaya, B. S. Kuznetsov, N. S. Shimanskaya, Sov. At. En. **1**, 912 (1959).
235. G. R. Martin, D. G. Tuck, Int. J. Appl. Radiat. Isot. **5**, 141 (1959).
236. H. Ramthun, Nukleonik **8**, 244 (1966).
237. I. Joliet-Curie, G. Bouissieres, Cahiers Phys. **26**, 1 (1941).