H.H. Rossi and M. Zaider

College of Physicians & Surgeons of Columbia University
RARAF, Nevis Labs., P.O. Box 21, Irvington, NY 10533

Abstract

The development of the principles of the theory of dual radiation action is reviewed and a treatment of saturation is given. Although the relationship derived cannot be evaluated in general in closed form, approximations may be obtained that apply to a range of conditions.

INTRODUCTION

The theory of dual radiation action (TDRA) was developed with the aim of applying microdosimetry to radiobiology. It therefore can deal only with the first phases in a long chain of events that results in patent effects. It is, however, clear that the initial spatial and temporal pattern of energy deposition has a profound influence on the ultimate outcome. As often happens, the early formulation of the theory contained a number of simplifying assumptions. Although most of these were explicitly stated when the first version of the TDRA was published (1), experimental data obtained when the limitations are important were cited as contrary evidence causing considerable confusion. A more advanced version (2) eliminated some of the restrictions but there remain others, one of which relates to certain aspects of saturation which are addressed in this presentation.

There has, however, been no change in the fundamental tenet of the theory that is indicated by the adjective "dual." It is postulated that many radiation effects in eukaryotes are due to lesions that arise as the result of pairwise interaction of molecular products termed sublesions. However, the probability of cellular effects follows the same mathematical relation as that for lesion formation only if there is proportionality between these quantities. Exchange-type chromosome aberrations appear to satisfy this requirement. Other endpoints require further assumptions: the so-called "linear-quadratic" relation for radiation carcinogenesis can be supported by the TDRA only if radiogenic cancers arise from single autonomous cells (i.e., cells that are transformed by processes that are independent of any radiation received by the rest of the organism), in fact an unlikely assumption (3).

The point of departure of the theory was the recognition of regularities in the dependence of the RBE of neutrons (relative to electromagnetic radiations) on the level of effect (4). Since the RBE is merely a dose ratio for equal effect, the RBE-dose relation has the intrinsic advantage that it can be quantitatively applied even when numerical specification of the degree of effect (e.g., skin damage) must be arbitrary. A perhaps even more useful aspect is that the RBE reflects only differences in the effectiveness of lesion formation if the subsequent processes do not depend on radiation quality and cause complex dose-response relations when the effect is due to injury of several and/or non-autonomous cells. It is worth mentioning also that for clinical applications of the theory, RBE vs. dose relations permit extrapolation of the large body of empirical data accumulated with low-LET radiations to other modalities.

The RBE of neutrons is frequently observed to depend on the inverse square root of their dose when cellular damage is due to single particles (primarily protons). At higher doses, D , it tends to approach the value of 1, while data obtained at very low doses indicate a trend towards very high but dose-independent RBE. This is consistent with a dependence of effect probability on the square of specific energy as expressed in the relation

$$\bar{\epsilon} = (\bar{z}_D D + D^2) \quad (1a)$$

or in the nonspecific form

$$\bar{\epsilon} = \alpha D + \beta D^2, \quad (1b)$$

where $\bar{\epsilon}$ is the expected yield of lesions, and $\bar{z}_D$ (also symbolized by ζ) is the dose average of the specific energy in single events. In a logarithmic graph the principal variation of RBE appears as a line of slope $-1/2$ and its extrapolation to an RBE of 1 identifies both the value of $\bar{z}_D$ and (on the basis of microdosimetric data for the neutrons involved) an assumed spherical volume in which energy concentration is critical (the site). Typically its dimensions are of the order of 1 μm .

The line of slope $-1/2$ in the logarithmic RBE vs. dose relationship represents the situation where high-LET radiation (e.g., neutrons) acts in single events only, while the low-LET radiation (e.g., x rays) causes a quadratic dose-effect dependence typical for multi-event action. In terms of the spatial pattern of energy transfer points, the two radiations are characterized respectively by highly correlated (neutrons) or completely random (x rays) distribution. It is then only the latter radiations for which the mean interaction distance is comparable to the dimensions of the site.

The observation (5,6,7) that, within wide limits, the cross section for cell killing by high-LET particles is proportional to the square of the LET, indicates that in one-event inactivation dual radiation action obtains as well. However, the RBE-dose dependence does not readily disclose the scale of interactions for

this process. In principle, it could be obtained by comparing the RBE values at various neutron energies, but in the usual case when the range of the charged particles (primarily protons) exceeds a few micrometers, the ratio of the $\bar{z}_D$ values depends only weakly on site diameter.

The possibility of formulating dual radiation action in terms of the more general distance model was envisaged in the original publication (1) and it was made possible in the second publication (2) because of the development of the fundamental concept of the proximity function (8). In the generalized version

$$\bar{\epsilon} = k(\bar{D} + D^2) \quad (2)$$

and the linear-quadratic dependence on absorbed dose is retained, but the coefficient of D becomes

$$\xi = \int_0^{\infty} t(x)\gamma(x)dx \quad (3)$$

where $t(x)$ is the proximity function of energy transfer, and $\gamma(x)$ is the probability that two energy deposits initially separated by x result in the formation of a lesion. The molecular ion experiment (9), in which pairs of ions of various separations traverse cells, showed $\gamma(x)$ to be a very skewed function of x , indicating that one-event injury is primarily due to combinations of sublesions over distances substantially less than one micrometer. This is consistent with the findings of high RBE in the cell irradiations by very short-range electrons produced by soft x radiation (10,11).

The marked difference in the initial separations between combining sublesions have suggested the terms translesions (usually produced in two events) and cislesions (almost always produced in one event), and the possibility that this might be explained in terms of the structure of DNA (12).

While the distance model, Eq.2, represents the most general dependence of ϵ on D, it is in fact possible (and useful) to formally retain the site model provided the notion of simple spherical sites is replaced by the concept of more complex structures which need not be convex (13).

The function $\gamma(x)$ has been defined by

$$\gamma(x) = g(x)s(x)/4\pi x^2 \rho \int_0^{\infty} g(x)s(x)dx \quad (4)$$

where $g(x)$ is the probability that two sublesions separated by x result in a lesion, and ρ is the density of the sensitive matrix which has a point-pair distribution of distances (ppdd) equal to $s(x)/V$, where V is the volume of the matrix. The sensitive matrix is defined as the geometrical locus where energy transfers can result in sublesions. The gross sensitive volume (GSV) is the smallest convex body containing the matrix.

Although $\gamma(x)$ has been derived by analysis of certain experimental data (9,14,15), there is at this time no information concerning either of the two functions that determine $\gamma(x)$. It is therefore possible to postulate, as a

working hypothesis, that $g(x)$ is constant and therefore that

$$\gamma(x) = s(x)/4\pi x^2 \rho V \quad (5a)$$

or

$$s(x) = 4\pi x^2 \gamma(x) \rho V \quad (5b)$$

i.e., that $\gamma(x)$ is entirely determined by the shape of the matrix. If it should be found that $g(x)$ is not constant, and that therefore the ppdd defined by Eq.5b is different from that of the actual matrix, it may be necessary to abandon the site model as a realistic representation. One may, however, retain the concept of an equivalent site in which the $s(x)$ defined by Eq.5b is pertinent.

The fact that experimental microdosimetry is difficult, if not impossible for such a shape (that may even not have any physical existence), does not detract from the possibility of making useful deductions based on the postulate that the simple relation

$$\epsilon(z) = kz^2 \quad (6)$$

applies to it. This approach is discussed in a forthcoming publication (16).

SATURATION

It was evident at the outset that the biological effectiveness per unit of absorbed dose may not increase indefinitely and that the concept of saturation could be important at high doses and also at high LET. As used here, this refers to a condition in which an increment of absorbed energy is less effective when it follows one or more previous increments. In principle, saturation could occur at three levels.

The first of these is the production of sublesions where increasing density of transfer points might lead to excessive energy deposition in the site of the sublesion (the locus) if the average spacing between transfers is comparable to or less than the site diameter for the induction of sublesions. This would result in an LET dependence of k (or of β in Eq.1b). Present formulations of the TDRA assume that the yield of sublesions is simply proportional to specific energy, implying that sublesions are produced by single energy transfers and that the probability of multiple energy deposits in the locus can be neglected. There are indications that β does depend on LET (7) and on neutron energy (17) when the rate of cell killing is assumed to follow Eq.1b. However, the available information is fragmentary, and there is no theoretical guide for a quantitative treatment. Other mechanisms (to be discussed) modify the expression for cellular survival and lead to incorrect estimates of β when data are fitted to Eq.1b. There thus remains a possibility that the production of sublesions is independent of radiation quality, but at this time no firm conclusions can be drawn.

At the next step of biophysical action, the formation of lesions, there is the evident possibility of saturation because of inefficiently high concentrations of sublesions. Analyses have been given for both the linear component (1) and the quadratic component (18). It was recognized that these treatments can be only

approximate because they employ average values of ϵ to nonlinear relations. It should be noted that the modifications of dose-effect relations are different for high and low LET. Saturation affects the linear component when the LET is increased, which ultimately results in a decreased yield of cislesions and a reduction of α . On the other hand, saturation of the quadratic component, which is of primary importance for high doses of low-LET radiation, affects the value of the specific energy in the relatively large sites involved in translesion formation, and alters the shape of dose-effective relation at large doses which can no longer be expressed in terms of D^2 .

At a third level there is the possibility that at high absorbed doses more lesions are produced than are required for the effect (i.e., cell killing). There are in fact reasons for assuming that a single lesion is lethal (12).

Yield of Lesions and Survival

Adoption of the relation given by Eq.6 implies acceptance of Eq.1a with the understanding that $\bar{z}_D$ is the dose weighted mean specific energy produced by single events in the domain defined by Eq.5b. However, it should be stressed again that $\bar{\epsilon}$ is an inadequate index of cellular survival and that the relation

$$S(D) = \exp[-\bar{\epsilon}(D)] = \exp[-k(\bar{z}_D D + D^2)] \quad (7)$$

can not be generally valid.

A simple way of demonstrating this is based on the fact that, under condition of cell autonomy (which is assumed throughout this presentation), cells that experience no event must survive. The event frequency, n , at an absorbed dose D is $D/\bar{z}_r$. Since events are by definition statistically independent, there is, by Poisson statistics, a lower bound to survival given by

$$S(D) \geq \exp(-n) = \exp(-D/\bar{z}_r) . \quad (8)$$

Because of the quadratic term in Eq.7, $S(D)$ must, at sufficiently large doses, become less than the lower limit in Eq.(8). It may be noted that this constitutes a model-independent elimination of the general validity of the so-called α - β formulation for cell survival.

It will in the following be assumed that at a given specific energy, z , the yield of induced lesion is Poisson distributed with an average value kz^2 and that every lesion is lethal.* (Neither of these assumptions is essential for the formalism described below.) It follows that when averaged over all possible values of z the probability of survival is given by:

*The distribution of lesions at a given absorbed dose is not Poisson distributed, as can be shown by microdosimetric arguments.

$$S(D) = \int_0^\infty \exp(-kz^2) f(z, D) dz . \quad (9)$$

It should be noted that Eq.9 does not include saturation of sublesion or lesion formation which requires a formulation in which kz^2 would be replaced by $kz_0^2 [1 - \exp(-z^2/z_0^2)]$ (1).

Equation 9 can not be generally evaluated in closed form unless the geometry of the sensitive matrix and its corresponding distribution $f(z, D)$ are known. There are, however, particular cases in which Eq.9 can be further simplified and evaluated. They cover useful ranges of absorbed doses and/or radiation types.

A: Consider first the case where $n \ll 1$, i.e., when cell traversal by more than one charged particle is unlikely. Moderate absorbed doses of high-LET radiation fall into this category.

The general expression (1)

$$f(z, D) = \sum_{v=0}^{\infty} \exp(-n) \frac{n^v}{v!} f_v(z) \quad (10)$$

where $f_v(z)$ is the distribution of z in exactly v events, can then be approximated by

$$f(z, D) \approx \exp(-n) [\delta(z) + n f_1(z)] \quad (11)$$

Substitution into Eq.9 gives

$$S(D) \approx \exp(-n) [1 + \int_0^\infty f_1(z) \exp(-kz^2) dz] \quad (12)$$

or

$$S(D) \approx \exp \left\{ -n \int_0^\infty [1 - \exp(-kz^2)] f_1(z) dz \right\} . \quad (13)$$

This relation may be compared with the results of track-segment experiments in which inactivation cross sections are determined as a function of LET

$$S(D) = \exp(-\sigma \Phi) \quad (14)$$

where Φ is the particle fluence

$$\Phi = D/\bar{y}_p = 4D/S\bar{z}_p = 4n/S . \quad (15)$$

Here, $\bar{y}_p$ is the frequency average lineal energy, and the second equality follows from Cauchy's theorem, with S being the surface of the matrix (19).

Comparison with Eq.13 yields

$$\sigma = S/4 \int_0^\infty [1 - \exp(-kz^2)] f_1(z) dz . \quad (16)$$

This relation contains $f_1(z)$, which depends on the generally unknown geometry of the matrix. The import of Eq.16 can be illustrated by considering the case where the matrix is replaced by a spherical GSV of diameter d , which is traversed by rectilinear particle tracks which deposit energy at a rate that is proportional to their linear energy transfer, L (i.e., ignoring track structure). In this

case, $f_1(z)$ becomes a simple triangular distribution

$$f_1(z) = 2z/z_M^2 \quad (17)$$

with z varying between zero and z_M given by

$$z_M = 6L/\pi\phi d^2 .$$

Introducing Eq.17 into Eq.18, one obtains

$$\sigma = \pi \left(\frac{d}{z} \right)^2 \left\{ 1 - \frac{1}{kz_M^2} [1 - \exp(-kz_M^2)] \right\} \quad (18)$$

Despite the approximations employed, Eq.18 is in reasonable agreement with experimental results (5,6,7), and it should be noted that the equation was derived without invoking a "saturating mechanism" (20).

B: For doses such that in Eq.9 $kz^2 \ll 1$, one can write

$$S(D) \approx \int_0^\infty (1 - kz^2) f(z, D) dz = 1 - k\overline{z^2} \exp(-k\overline{z^2}) \quad (19)$$

In this range of doses and only then, Eq.7 (i.e., the usual α - β relation) is an adequate approximation of survival probability.

C: For high absorbed doses (i.e., when $n \gg 1$), the microdosimetric distribution $f(z, D)$ becomes normally distributed irrespective of the shape of the matrix or of the radiation:

$$f(z, D) \approx (2\pi\overline{z}_D^2)^{-1/2} \exp[-(z - D)^2 / 2\overline{z}_D^2] \quad (20)$$

and from Eq.9 one obtains

$$S(D) \approx (1 + 2k\overline{z}_D^2)^{-1/2} \exp[-kD^2 / (1 + 2k\overline{z}_D^2)] \quad (21)$$

This relation is of especial importance to low-LET radiation where n is large in the relatively big volume in which translesions are formed. It indicates that at doses of the order of tens of Gy the survival curve becomes quasi-exponential.

D: At even higher doses (such that $2k\overline{z}_D^2 \gg 1$), Eq.(23) evidently simplifies to

$$S(D) \approx (2k\overline{z}_D^2)^{-1/2} \exp(-D/2\overline{z}_D) \quad (22)$$

It should be noted that Eqs.21 and 22, although displaying a nonquadratic dependence of $S(D)$ on absorbed dose, still depend on radiation quality only through $\overline{z}_D$, a quantity which can be calculated directly from Eq.3, thus circumventing the need to know the actual geometry of the sensitive site.

CONCLUSION

The seemingly simple premises of the TDRA can lead to complex dependences of cell survival on absorbed dose. It is, however, possible to derive useful approximations in several instances. The need for further efforts is evident.

ACKNOWLEDGEMENT

We are greatly indebted to Prof. A.M. Kellerer who originated many of the concepts presented here and provided invaluable advice. This work was supported by Department of Energy Grants DE-FG02-86ER60449 and DE-FG02-86ER60402 to the Radiological Research Laboratory, Columbia University.

REFERENCES

- 1 Kellerer AM, Rossi HH (1972) Curr Topics Radiat Res Q 8:85
- 2 Kellerer AM, Rossi HH (1978) Radiat Res 75:471
- 3 Rossi HH (1985) Amer J Roentgen 144:1
- 4 Rossi HH (1970) Phys Med. Biol 15:2
- 5 Powers EL, Lyman JT, Tobias, CA (1968) Int J Radiat Biol 14:313
- 6 Barendsen GW (1967) In: Cole A (ed) Theoretical and Experimental Biophysics, p 167. Marcel Dekker, New York
- 7 Bird RP, Rohrig N, Colvett RC, Geard CR, Marino S (1980) Radiat Res 82:277
- 8 Kellerer AM, Chmelevsky D (1975) Rad and Environm Biophys 12:321
- 9 Kellerer AM, Lam YM, Rossi HH (1980) Radiat Res 83:511
- 10 Neary GJ, Simpson-Gildemeister, VFW, Peacocke AR (1970) Int J Radiat Biol 18:25
- 11 Goodhead DT, Thacker J, Cox R (1981) Int J Radiat Biol 36:101
- 12 Rossi HH (1985) Radiation Protection Dos 13(1-4):259
- 13 Zaider M, Rossi HH (1985) Radiat Res Suppl 104:S-68
- 14 Zaider M, Bird RP, Rossi HH, Marino S, Rohrig N (1983) Radiat Environm Biophys 22:239
- 15 Zaider M, Brenner DJ (1984) Radiat Res 99:484
- 16 Zaider M, Rossi HH (submitted 1987) Radiat Res
- 17 Kellerer AM, Hall EJ, Rossi HH, Teedla P (1976) Radiat Res 65:172
- 18 Zaider M, Rossi HH (1980) In: Radiation Physics, Biophysics and Radiation Biology, DOE Annual Report of Research Project, Columbia University, p 126
- 19 Zaider M, Rossi HH (1986) In: Orton CG (ed) Radiation Dosimetry, p 171. Plenum Publishing Corp, New York
- 20 Goodhead DT (1985) Radiat Res Suppl 104: S-58

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.