PROGRESS REPORT (5/86-1/91)

Title: ACTIVE AND PASSIVE CALCIUM TRANSPORT SYSTEMS IN PLANT CELLS
DOE/ER/13461-6

P.I. Heven Sze
Department of Botany, University of Maryland, College Park

The ability to change cytoplasmic Ca^{2+} levels ($[\text{Ca}^{2+}]$) by cells has made this cation a key regulator of many biological processes. Cytoplasmic $[\text{Ca}^{2+}]$ is determined by the coordination of passive Ca^{2+} fluxes which increase cytosolic $[\text{Ca}^{2+}]$ and active Ca^{2+} transport systems that lower cytosolic $[\text{Ca}^{2+}]$. The mechanisms by which plant cells achieve this is poorly understood. We have initially used isolated vesicles from the plasma membrane or organellar membranes to study Ca^{2+} transport systems in oat roots (a monocot) and carrot suspension cells (a dicot). The objectives of the proposal (submitted in 5/86) were to identify and characterize active (energy-dependent) and passive calcium transport systems that work together to regulate calcium levels in the cytoplasm of plant cells.

DETAILED PROGRESS REPORT

1. Calcium is accumulated into vacuoles via a $\text{H}^+/\text{Ca}^{2+}$ antiporter:
 - (a) Properties of the vacuolar $\text{Ca}^{2+}/\text{H}^+$ porter. [Schumaker & Sze (1986)]

Calcium is sequestered into vacuoles of oat root cells through a $\text{H}^+/\text{Ca}^{2+}$ antiporter system that is driven by the proton motive force of the tonoplast H^+ -translocating ATPase (Schumaker & Sze, 85). Unique properties of a transporter are needed in order to identify the protein. The characteristics of any H^+ -coupled transport system can be studied directly using an artificial pH gradient. We have used artificial pH gradients to study the properties of a $\text{Ca}^{2+}/\text{H}^+$ antiporter system from plant membranes. The pH gradient was imposed by diluting K^+ -loaded vesicles into a K^+ -free medium. Nigericin induced a K^+/H^+ exchange resulting in a pH gradient of 2 (acid inside). The pH gradient was capable of driving Ca^{2+} accumulation. Ca^{2+} uptake was tightly coupled to H^+ loss as increasing Ca^{2+} levels progressively dissipated the steady state pH gradient. Ca uptake displayed saturation kinetics with a K_m for Ca of 10 μM . The relative affinity of the antiporter for divalent cations was $\text{Ca}^{2+} > \text{Sr}^{2+} > \text{Ba}^{2+} > \text{Mg}^{2+}$. La^{3+} or Mn^{2+} blocked Ca^{2+} uptake possibly by occupying the Ca^{2+} binding site. Ruthenium red and N,N-dicyclohexylcarbodiimide specifically inhibited the $\text{H}^+/\text{Ca}^{2+}$ antiporter. A stoichiometry of the antiporter of $1\text{H}^+/1\text{Ca}^{2+}$ and a pH gradient of 2 would result in an accumulation ratio of 100:1; however a stoichiometry of 2 or 3H^+ per Ca^{2+} would result in a higher accumulation ratio. The antiporter thus plays an important role in Ca^{2+} accumulation into the vacuole.

- (b) Solubilization and reconstitution of the $\text{H}^+/\text{Ca}^{2+}$ antiporter. [Schumaker & Sze 1990]

Unlike transport ATPases, a cotransporter like the $\text{H}^+/\text{Ca}^{2+}$ antiporter, does not possess enzymatic activity and so it cannot be detected by enzymic activity after solubilization with detergents. One way to detect the porter is to functionally assay transport activity in a reconstituted system. Thus, the first

step towards purification and identification of the H^+/Ca^{2+} antiporter is to solubilize and reconstitute the transport activity in liposomes. The tonoplast proteins were solubilized with octylglucoside in the presence of soybean phospholipid and glycerol. After centrifugation, the soluble supernatant yielded a preparation of the transporter which could be reconstituted into liposomes by the detergent dilution method. A pH gradient (acid inside) was generated in proteoliposomes with an NH_4Cl gradient. The proteoliposomes catalyzed calcium uptake with a specific activity about 6 times higher than in native vesicles. Properties of the pH gradient-dependent calcium uptake such as the K_m for calcium (about 15 μM) and the sensitivity to inhibitors such as N,N' -dicyclohexylcarbodiimide, ruthenium red, and lanthanum were similar to those found in membrane vesicles, indicating that the H^+/Ca^{2+} antiporter has been reconstituted in active form.

The ability to reconstitute an active exchanger is an important and crucial step towards the identification of the polypeptide(s) responsible for calcium accumulation into higher plant vacuoles. This is one of our future goals.

2. Ca^{2+} in the vacuole can be released into the cytosol via Inositol 1,4,5-Trisphosphate ($InsP_3$)-sensitive Ca^{2+} channels: $InsP_3$ releases Ca^{2+} from tonoplast vesicles. [Schumaker & Sze (1987a)]

This is one of the most exciting findings related to signal transduction in plant cells. We demonstrated the first example of $InsP_3$ -induced Ca^{2+} release from vacuoles. Only $InsP_3$ caused a transient reduction of $^{45}Ca^{2+}$ levels in tonoplast vesicles, while inositol, 1,4-bisphosphate, myo-inositol 1-phosphate or fructose 2,6-bisphosphate did not. The decrease was dependent on $InsP_3$ concentration ($K_m = 0.6 \mu M$). The $InsP_3$ -induced Ca^{2+} release was blocked by the Ca^{2+} antagonist, 8-(N,N -diethylamino)-octyl 3,4,5-trimethoxybenzoate-HCl. These results suggest the vacuolar membrane is one target site for $InsP_3$ action and that $InsP_3$ may operate as a second messenger in the mobilization of intracellular Ca^{2+} in plant cells. We are now interested in identifying the $InsP_3$ receptor and the gated Ca^{2+} channels that respond to $InsP_3$ on the vacuolar membrane from oat roots.

3. H^+ -coupled anion transport on the vacuolar membrane: Decrease of pH gradients in tonoplast vesicles by NO_3^- and Cl^- . [Schumaker & Sze, 1987b]

Anions, like nitrate and phosphate, are essential nutrients; however, their transport and storage sites are poorly understood. We found that chloride or nitrate decreased a pH gradient (measured as ^{14}C -methylamine accumulation) in tonoplast vesicles. The pH gradient decrease was dependent on the anion concentration. These effects are independent of the anion-sensitive H^+ -ATPase of the tonoplast, since the pH gradient (acid inside) was imposed artificially using a K^+ gradient and nigericin. Two possible models to account for this anion-dependent decrease of the pH gradient are: (i) H^+ loss is accompanied by Cl^- or NO_3^- efflux via a H^+ /anion symport system and (ii) H^+ loss is accompanied by Cl^- or NO_3^- uptake via a H^+ /anion antiport system. These anion transport systems could serve to mobilize Cl^- or NO_3^- stored in the vacuole for use in the cytoplasm or to drive anions into the vacuole. The Cl^- transporter is sensitive to 4,4'-diisothiocyano-2,2-stilbene disulfonic acid which may be a good marker to identify this porter.

[From these studies, we found that the relative uptake of Ca into tonoplast vesicles directly reflected the relative magnitude of the pH gradient. This is a

useful property since Ca gradient is relatively stable ,so its uptake is often easier to measure than the pH gradient.]

4. Plasma membrane-type and Endoplasmic reticulum-type Ca^{2+} -pumping ATPases from carrot cells. [Pierce & Sze (1987); Hsieh, Pierce & Sze (in prep)]

A Ca^{2+} -ATPase on endoplasmic reticulum-enriched vesicles has been previously identified and characterized (Bush and Sze, 1986. Plant Physiol. 80: 549). This transport system is inhibited by vanadate and has a high affinity for Ca ($K_m[\text{Ca}] = 1 \mu\text{M}$). We concluded this transport activity was associated with the ER since it comigrated on sucrose density gradient with the ER marker, antimycin A-insensitive NADH-dependent cytochrome c reductase and Ca transport was also shifted by Mg in parallel with the ER marker. We later found that this Ca^{2+} transport activity is stimulated 2-4 fold by $1 \mu\text{M}$ calmodulin. To confirm the membrane identity of this transport activity, the vesicles were separated with a sucrose density gradient and then assayed for calmodulin-sensitive Ca^{2+} transport. A large part of the total calmodulin-stimulated activity was associated with ER as shown by its comigration with the NADH-cytochrome c reductase activity which peaked at 1.12 gm/ml . Calcium uptake into these vesicles were also stimulated by oxalate. However, there was also a small peak of calmodulin stimulated Ca^{2+} transport (expressed as specific activity) in vesicles at a density of 1.17 gm/ml , these vesicles were less responsive to oxalate (not shown). The results suggest that calmodulin-stimulated calcium transport was associated with ER vesicles as well as plasma membrane vesicles. Furthermore, an oxalate transport system appeared to be associated with the ER but not with the plasma membrane.

We were initially uncertain about the significance of our findings. In animal cells, calmodulin stimulates the plasma membrane-type Ca-ATPase directly and the SR/ER-type Ca-ATPase indirectly via protein phosphorylation of a regulatory polypeptide. Similar modes of regulation may exist in plant cells. One of our future goals is to identify the type of Ca^{2+} -pumping ATPase in carrot microsomes and determine the mode of regulation by calmodulin.

5. Calcium transport across the nuclear membrane.

The function of the nuclear membrane is very poorly understood and there are indications it plays a role in regulating nuclear and cytoplasmic Ca^{2+} levels and stages of mitosis. We first attempted to isolate intact nuclei according to a modified method of John Watson (in our department) and then to vesiculate the membranes. After repeated trials and errors, we have been unable to observe ATP-dependent Ca^{2+} uptake consistently in the preparations. We tried all the approaches known to our laboratory to enhance calcium transport with no success. Furthermore, examination of the isolated nuclei under light microscope revealed contamination by other membraneous material (probably the endoplasmic reticulum which can be contiguous with the nuclear membrane). For these reasons, we have decided not to pursue these studies for the present time.

6. Mode of Action of the *Helminthosporium maydis* toxin: Dissipation of the membrane potential and protection by dicyclohexylcarbodiimide.
(Holden & Sze, 1987; Holden & Sze, 1989)

[This study was initially supported by a preceding DOE grant. The time needed to complete the study overlapped partly with the present grant period.]

A toxin produced by the fungus, *Helminthosporium maydis* Race T (HmT), causes symptoms of leaf blight in corn with Texas male-sterile cytoplasm, while corn with N cytoplasm is insensitive to the toxin. Both toxin sensitivity and male sterility are maternally inherited, and mitochondrial gene products are thought to be responsible for these traits. We are interested in understanding the mode of toxin action and have shown that HmT toxin and various analogs induce H^+ leakage from T mitochondria. HmT toxin dissipated the steady state membrane potential (as measured by safranine absorbance change) similar to a protonophore. Synthetic toxin analogs (Cpd XIII: $C_{41}H_{86}O_{12}$ and Cpd IV: $C_{25}H_{44}O_6$) as well as reduced toxin (RT2C: $C_{41}H_{84}O_{13}$) and Pm toxin ($C_{33}H_{60}O_8$) produced by the fungus, *Phyllosticta maydis* were effective in dissipating the membrane potential or decreasing Ca uptake. In contrast, the toxins and analogs had no effect on membrane potential formation in normal (N) mitochondria. The striking similarities of the HmT toxin ($C_{41}H_{86}O_{12}$) and Cpd XIII on T mitochondrial activities provide strong evidence supporting the correctness of the polyketol structure assigned to the native toxin. Since the membrane potential in energized mitochondria is caused by the electrogenic extrusion of H^+ , the results support the idea that HmT toxin increases membrane permeability of T mitochondria to H^+ .

The effect of *Helminthosporium maydis* Race T toxin on electron transport in susceptible cytoplasmic male-sterile Texas corn mitochondria was investigated, using dichlorophenol indophenol and ferricyanide as electron acceptors. Succinate-dependent electron transport was stimulated by the toxin, consistent with the well-described increase in membrane permeability induced by the toxin. In contrast, malate-dependent electron transport was inhibited. Mitochondria from normal-fertile corn were not affected by the toxin. Both the inhibition of electron transport and the increase in ion permeability, such as dissipation of the membrane potential and Ca^{2+} gradients, induced by the toxin in T corn was prevented by N,N'-dicyclohexylcarbodiimide, a hydrophobic carbodiimide. A water-soluble carbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide was ineffective in preventing dissipation of membrane potential by the toxin. These results suggest that the various toxin actions are mediated via interaction of the toxin with one target molecule, most probably a 13 kDa polypeptide unique to T mitochondria. N,N'-dicyclohexylcarbodiimide (DCCD) may confer protection by modifying an amino-acid residue in a hydrophobic portion of the receptor.

These results were helpful in subsequent identification of the toxin receptor by C.S. Leving and colleagues. Our studies suggested the receptor binds to DCCD which was later shown to bind the 13 kDa polypeptide. However we still do not understand how the toxin induces an increase in membrane permeability in T corn mitochondria.

7. Molecular Cloning and Sequencing of the cDNAs encoding the 16kDa proteolipid of the vacuolar H^+ -ATPase

As a first step in understanding the molecular structure of transport proteins, we have begun to clone and sequence one subunit of the H^+ -pumping ATPase, specifically the 16 kda proteolipid. The techniques and procedures currently used to obtain cDNAs to the vacuolar ATPase will be later applied to cDNAs encoding calcium transport pumps or channels.

SUMMARY OF PROGRESS AND SIGNIFICANCE

Our studies show that plant vacuoles play a major role in regulating cytosolic $[Ca^{2+}]$. Ca^{2+} can be accumulated into the vacuole using the proton motive force generated by the tonoplast H^+ -pumping ATPase. The H^+/Ca^{2+} antiporter has been solubilized and reconstituted in active form. This is significant as the reconstituted activity will now allow us to identify the exchanger during purification. Unexpectedly, we discovered that the stored Ca can be released from the vacuole by inositol 1,4,5-trisphosphate, a compound which may be produced in response to environmental or hormonal signals. This was the first report that suggested the presence of an $InsP_3$ -sensitive Ca^{2+} channel on the vacuolar membrane from plant cells. Given the role of PI intermediates in signal transduction, our result has already caused much excitement among plant biologists and has stimulated other laboratories to detect, characterize and identify such Ca channels in other plant tissues.

We have also shown that two different Ca^{2+} -pumping ATPases, one at the plasma membrane and another on the endoplasmic reticulum, play important roles in removal of cytoplasmic Ca^{2+} . Surprisingly, both seem to be stimulated by calmodulin. However the mode of regulation is unclear and needs to be further investigated.

One of our significant findings to understand the mode of action of the Helminthosporium maydis T toxin, is demonstrating the prevention of toxin effects by DCCD. The implication that DCCD binds to the toxin receptor was used subsequently by others to identify the receptor.

PUBLICATIONS (supported in part by DOE)

Bush, D.R. & H. Sze. 1986. Calcium transport in tonoplast and endoplasmic reticulum vesicles from cultured carrot cells. *Plant Physiol.* 80, 549-55

Holden, M.J. & H. Sze. 1987. Dissipation of the membrane potential in susceptible corn mitochondria by the toxin of *Helminthosporium maydis* Race T, and toxin analogs. *Plant Physiol.* 84, 670-76

Holden, M.J. & H. Sze. 1989. Effects of *Helminthosporium maydis* Race T toxin on electron transport in susceptible corn mitochondria and prevention of toxin actions by dicyclohexylcarbodiimide. *Plant Physiol.* 91, 1296-302

Schumaker, K.S. & H. Sze. 1986. Calcium transport into the vacuoles of oat roots. Characterization of H^+/Ca^{2+} exchange activity. *J. Biol. Chem.* 261, 12172-78

Schumaker, K.S. & H. Sze. 1987a. Inositol 1,4,5-Trisphosphate releases Ca^{2+} from vacuolar membrane vesicles of oat roots. *J. Biol. Chem.* 262, 3944-46

Schumaker, K. S. & H. Sze. 1987b. Decrease of pH gradients in tonoplast vesicles by NO_3^- and Cl^- : evidence for H^+ -coupled anion transport. Plant Physiol. 83, 490-96

Schumaker, K. S. and H. Sze. 1990. Solubilization and reconstitution of the oat root vacuolar $\text{H}^+/\text{Ca}^{2+}$ exchanger. Plant Physiol. 92, 340-45

Hsieh, W-L, W. Pierce & H. Sze. Calcium pumping ATPases in vesicles from carrot cells: stimulation by calmodulin and phosphatidylserine and formation of a 120 kDa phosphoenzyme. (4/91 manuscript in preparation for Plant Physiology)

ABSTRACT

Pierce, W. & H. Sze. 1987. Calmodulin stimulation of calcium transport in carrot microsomal vesicles. Plant Physiol. 63: S-53

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DATE
FILMED**
418192

I