LA-8166-T

Thesis

DR. 960

MASTER

**Large Momentum Transfer Neutron Pickup
with the (π^+ , p) and (p, d) Reactions**

University of California

LOS ALAMOS SCIENTIFIC LABORATORY

Post Office Box 1663 Los Alamos, New Mexico 87545

Gregory Richard Smith*

- DISCLAIMER

[illegible]

CONTENTS

ABSTRACT	xix
Chapter	
I. INTRODUCTION	1
II. DWBA THEORY AND CALCULATIONS	8
2.1 Formalism	11
2.2 The PWBA Limit	25
2.3 Deuteron Wave Function	29
2.4 Neutron Bound State Potentials	36
2.5 Empirical Optical Potentials	42
2.6 KMT Optical Potentials	46
2.7 Well-Matched Optical Potentials	55
2.8 Core Coupling	57
2.9 Numerical Tests and Sensitivities	63
2.10 Angular Momentum Selectivity	73
III. THEORY OF THE (π^+ ,p) REACTION	75
3.1 One Nucleon Mechanism (π^+ ,p) Reaction Theory	79
3.2 General Features of the Theory	82
IV. EXPERIMENTAL METHODS FOR THE (p,d) STUDIES	95
4.1 Beam Preparation	95
4.2 HRS	101
4.3 Scintillation Detectors	105
4.4 Particle Identification	107

Chapter	Page
4.5 Wire Chambers	111
4.6 Targets	115
4.7 Monitors	118
4.8 Acceptance	120
4.9 Data Acquisition System	124
4.10 Correlations	128
4.11 Absolute Normalization	131
4.12 Cross Sections	133
4.13 Experimental Uncertainties	136
V. EXPERIMENTAL METHODS FOR THE (π^+ ,p) STUDIES	141
5.1 Pion Production	142
5.2 Pion Channel	144
5.3 Spectrometer	149
5.4 Scintillation Detectors	153
5.5 Wire Chambers	159
5.6 Targets	162
5.7 Monitors	163
5.8 Acceptance	165
5.9 Data Acquisition and Correlations	165
5.10 Absolute Normalization	167
5.11 Experimental Uncertainties	171
VI. FEATURES OF THE 800 MeV (p,d) REACTION	173
6.1 $^7\text{Li}(p,d)^6\text{Li}$	175

Chapter		Page
6.2	$^{13}\text{C}(\text{p},\text{d})^{12}\text{C}$	186
6.3	$^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$	204
6.4	$^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$	226
6.5	$^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}$	242
6.6	$^{28}\text{Si}(\text{p},\text{d})^{27}\text{Si}$	250
6.7	$^{40}\text{Ca}(\text{p},\text{d})^{39}\text{Ca}$	262
VII.	PRESENTATION OF THE (π^+, p) DATA AND CALCULATIONS	270
7.1	$^6\text{Li}(\pi^+, \text{p})^5\text{Li}$	271
7.2	$^7\text{Li}(\pi^+, \text{p})^6\text{Li}$	275
7.3	$^{12}\text{C}(\pi^+, \text{p})^{11}\text{C}$	281
7.4	$^{13}\text{C}(\pi^+, \text{p})^{12}\text{C}$	287
VIII.	COMPARISONS AND CONCLUSIONS	297
8.1	The 800 MeV (p,d) Reaction	297
8.2	The (π^+, p) Reaction Near 90 MeV and 180 MeV	301
8.3	Comparison of the (p,d) and (π^+, p) Reactions	303
	ACKNOWLEDGMENTS	307
	REFERENCES	309
	APPENDIX: TABLES OF THE π^+, p AND p,d DATA	318

LIST OF TABLES

Table	Page
2.1.1 $^{13}\text{C}(p,d)^{12}\text{C}$ Spectroscopic Factors . . .	23
2.4.1 Neutron Bound State Potentials	43
2.6.1 KMT Optical Potential Input Quantities (appropriate for protons of the specified kinetic energy)	51
2.6.2 800 MeV KMT Optical Potentials	54
2.7.1 800 MeV Well-Matched Optical Potentials	58
2.8.1 Optical Potential Parameter Sensitivities (Deuteron Imaginary Radius and Diffuseness) for 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}(\text{g.s.})$ ZR DWBA	69
3.2.1 Pion Optical Potentials for the $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ Reaction	84
3.2.2 Proton Optical Potentials for the $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ Reaction	87
4.11.1 Absolute Normalization Results for the 800 MeV (p,d) Data	134
5.10.1 $\pi^+d \rightarrow pp$ Differential Cross Section Parameterizations	169
6.3.1 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ Calibration	207
6.4.1 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ Calibration	228
6.5.1 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ Calibration	244
6.6.1 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ Calibration	253
6.7.1 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ Calibration	264

LIST OF FIGURES

Figure		Page
1.1.	Fourier transform of the $p_{3/2}$ neutron bound state wave function used in the analysis of the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction, plotted as a function of the momentum transfer q	4
2.1	Schematic representation of the One Nucleon Model, as explained in the text	10
2.1.1	Reaction coordinate definitions and labels for the constituents of the (p,d) reaction	13
2.3.1	Fourier transform of $V_{pn}(r_{pn})\phi_d(r_{pn})$ vs. momentum transfer q	33
2.4.1	Shown are 375 MeV $^{16}\text{O}(e,e)$ data (open circles) (SM 70) presented as the square of the charge form factor vs. q^2 (momentum transfer squared) . . .	41
2.8.1	Two step pathways for $p_{3/2}$ (solid lines) and $p_{1/2}$ (dashed lines) pickup to the 4.44 MeV 2^+ state of ^{12}C from the $^{13}\text{C}(p,d)^{12}\text{C}$ reaction . . .	60
2.8.2	Calculations of the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ (4.44 MeV 2^+) reaction . . .	62
2.9.1	Sensitivity of the DWBA calculations to the nuclear interior, demonstrated for 800 MeV $^{13}\text{C}(p,d)$ and $^{40}\text{Ca}(p,d)$. . .	66
2.9.2	Nuclear charge densities of ^{13}C (solid line) and ^{40}Ca (dashed line) vs. the nuclear radius, for purposes of comparison to Fig. 2.9.1 . . .	67

Figure		Page
2.9.3	800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}(\text{g.s.})$ ZR DWBA predictions using best fit (see Sect. 2.5) proton optical potentials derived from 800 MeV proton elastic scattering on targets of ^{12}C (dashed line) and ^{58}Ni (solid line)	70
2.9.4	Sensitivity of 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}(\text{g.s.})$ DWBA calculations to the choice of parameters for the neutron bound state: $r = 1.40$, $a = 0.45$ (dashed line) and $r = 1.44$, $a = 0.55$ (solid line)	72
3.1	Schematic (Feynmann-like) diagrams of the one nucleon model (ONM), the two nucleon model (TNM) and the one pion exchange model (OPEM)	77
3.2.1	Predicted energy dependence of the $^{12}\text{C}(\pi^+,p)^{11}\text{C}(3/2^- \text{ g.s.})$ reaction, at 90 MeV (solid curve) and 180 MeV (dashed curve)	90
3.2.2	A comparison of 162 MeV $^{12}\text{C}(\pi,\pi)^{12}\text{C}$ elastic scattering data (Pif+ 77) with the predictions for 180 MeV $^{12}\text{C}(\pi,\pi)^{12}\text{C}$ obtained from the pion optical potential of Table 3.2.1	92
4.1.1	General layout of the Los Alamos Meson Physics Facility, showing the relative locations of the various experimental areas	96
4.1.2	Experimental areas A, B, and C at LAMPF, including the switchyard. The HRS is housed in area C	98
4.1.3	Closeup of beam area C, showing the locations of the HRS and the various magnetic elements of the beam line	99
4.2.1	Side view of the HRS, showing the locations of the detectors, spectrometer dipoles, and scattering chamber	102

Figure		Page
4.2.2	Overhead view of the HRS scattering chamber, showing the locations of the various beam monitors such as the ion chambers (ICI-4), secondary emission monitor (SEM), and the elements (DI-4) of the monitor telescopes (MT1-2). Other devices shown are explained in the legend . . .	104
4.3.1	Detectors located atop the HRS, including jaw counters (J1-2), veto counters (V1-2), test counters (T1-3), trigger (scintillator) counters (S2-4, S6) and delay line proportional counters (DL1, DL6)	106
4.3.2	Schematic of the HRS trigger electronics. Shown are the signals from each end (up (U) and down (D)) of the trigger scintillation counters, and how these signals are combined to generate a flag used to identify a valid event. Also shown is the monitor telescope electronics . .	108
4.4.1	Particle identification spectra taken at the HRS on a ^{12}C target at spectrometer angles of 12.5° (A) and 29° (B). .	110
4.5.1	Shown are the signals from each end of the anode (A) and cathode (B) wire planes, as described in the text	113
4.8.1	The focal plane relative efficiency data taken for experiment 265 (A) and experiment 117 (B)	123
4.8.2	Software gates set on x angle (y-axis) vs. focal plane x	126
4.10.1	Focal plane θ vs. excitation energy with (A) and without (B) the polynomial correction terms discussed in the text	130
5.1.1	Experimental Area A showing the location of the EPICS, A-1 production target, and other meson lines	143

Figure	Page
5.2.1 Plot of the pion production cross section vs. production angle at the A-1 pion production target	146
5.2.2 Schematic of the EPICS channel and its associated optics in the horizontal plane (dashed line) and the vertical plane (solid line)	147
5.3.1 Schematic representation of the magnetic elements comprising the EPICS spectrometer	150
5.3.2 Schematic diagram of the EPICS scattering chamber, its various elements, and relevant angles	151
5.4.1 The EPICS detector arrangement showing the relative locations of the wire chambers, scintillators, and magnetic elements of the spectrometer	154
5.4.2 Particle identification dotplots of (^{12}C target) TOF vs. dE/dx as defined in Sect. 4.4, at spectrometer angles of 20° (A) and 100° (B) . .	157
5.4.3 Diagram of the fast electronics used at EPICS for this experiment	158
5.8.1 Acceptance curve for the EPICS spectrometer obtained using the 180 MeV $^2\text{H}(\pi^+, p)^1\text{H}$ reaction	166
6.1.1 A spectrum of the 800 MeV $^7\text{Li}(p, d)^6\text{Li}$ reaction at $\theta_L = 9.5^\circ$	176
6.1.2 Angular distributions for the 1^+ and 3^+ levels of ^6Li populated in the 800 MeV $^7\text{Li}(p, d)^6\text{Li}$ reaction	179
6.1.3 Two step pathways, deformations and spectroscopic amplitudes used as ingredients in the $^7\text{Li}(p, d)^6\text{Li}$ calculations (CCBA) including pickup of a $p_{3/2}$ (solid line) and $p_{1/2}$ (dashed line) neutron	180

Figure		Page
6.1.4	The 800 MeV ${}^7\text{Li}(p,d){}^6\text{Li}(1^+ \text{ g.s.})$ angular distribution is compared to EFR DWBA calculations employing KMT folded (solid line) and well-matched (dashed line) deuteron optical potentials with $\Sigma C^2S = 0.83$. .	181
6.1.5	Data for the 800 MeV ${}^7\text{Li}(p,d){}^6\text{Li}$ (2.185 MeV 3^+) level are shown, with four calculations: EFR DWBA predictions employing ($C^2S = 0.47$) KMT (solid line) and well-matched (dashed line) deuteron optical potentials, and 2R calculations ($D_0 = 80$ MeV $\sim \text{fm}^{3/2}$) utilizing one step (dotted line) and the two step (dot-dashed line) pathways of Fig. 6.1.3 . .	184
6.2.1	Sample spectrum of the 800 MeV ${}^{13}\text{C}(p,d){}^{12}\text{C}$ reaction at 12°	188
6.2.2	Schematic (Feynmann-like) diagram of the two nucleon model (TNM) representation of the (p,d) reaction mechanism	189
6.2.3	Shown are sample spectra of the 43 MeV ${}^{11}\text{C}({}^3\text{He},d){}^{12}\text{C}$ reaction (A) and the 43 MeV ${}^{13}\text{C}({}^3\text{He},\alpha){}^{12}\text{C}$ reaction (B) at $\theta_L = 30^\circ$	192
6.2.4	Angular distributions for some of the levels excited in the 800 MeV ${}^{13}\text{C}(p,d){}^{12}\text{C}$ reaction	194
6.2.5	The 800 MeV ${}^{13}\text{C}(p,d){}^{12}\text{C}(0^+ \text{ g.s.})$ angular distribution is compared to EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials	196

Figure		Page
6.2.6	Angular distributions of the 12.7 MeV $1^+ T = 0$, 15.1 MeV $1^+ T = 1$, and 16.1 MeV $2^+ T = 1$ levels excited in the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction are compared to EFR DWBA calculations with the KMT deuteron optical potential used in Fig. 6.2.4 . . .	198
6.2.7	The effect of the coupled channels pathways (shown in Fig. 2.8.1) is demonstrated for the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ (4.44 MeV 2^+) angular distribution, by comparing the results of EFR DWBA (dot-dash line), ZR one step (dashed line) and ZR two step (solid line) calculations ($D_0 = 80 \text{ MeV} \cdot \text{fm}^{3/2}$)	200
6.2.8	Shown is the deformation, spectroscopic amplitude, and pathway (A) for the two step calculation of $^{13}\text{C}(p,d)^{12}\text{C}$ (14.1 MeV 4^+)	203
6.3.1	Spectra collected at 5.2° (A) and 19° (B) of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction	205
6.3.2	Angular distributions of the differential cross sections for some of the levels excited in the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction	211
6.3.3	Angular distributions for some of the levels excited in the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction	212
6.3.4	The angular distribution of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ (6.48 MeV $7/2^-$) reaction is presented with an angular binning of only 0.46° , as opposed to the normal bin of 1.8° used in the rest of the figures	213

Figure	Page
6.3.5	The differential cross section of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction to the $3/2^-$ ground state of ^{11}C (open circles) are compared to those of the 700 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction to the same state (solid circles) (Bak+ 74) vs. momentum transfer q 215
6.3.6	Same as for Fig. 6.3.5 only for the 2.0 MeV $1/2^-$ state of ^{11}C 216
6.3.7	Same as for Fig. 6.3.6 only for the 6.48 MeV $7/2^-$ state of ^{11}C 217
6.3.8	The 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}(\text{g.s.})$ data are compared to EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials 218
6.3.9	Parentage of levels in ^{11}C populated by the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction, in the (extreme) weak coupling model 220
6.3.10	Two-step pathways, deformations, and spectroscopic amplitudes for $p_{3/2}$ (solid line) and $p_{1/2}$ (dashed line) neutron pickup to the $7/2^-$ member of a (suspected) multiplet located at 13.2 MeV in ^{11}C , populated by the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction 222
6.3.11	Shown is the angular distribution of the 13.2 MeV bump populated in the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction, and the results of two calculations 224
6.4.1	Spectra of the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction at 5° and 17° 227
6.4.2	Angular distributions of some of the levels excited in the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction 229

Figure	Page
6.4.3	Angular distributions of some of the levels excited in the 800 MeV $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$ reaction 230
6.4.4	The angular distribution of the $1/2^-$ ^{15}O ground state from the 800 MeV $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$ reaction is presented along with EFR DWBA calculations employing KMT (solid curve) and phenomenological (dashed curve) (Bru 78) deuteron optical potentials . . 235
6.4.5	Data from the 800 MeV $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$ (6.48 MeV $3/2^-$) reaction are presented with the results of EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials. 236
6.4.6	Saclay data (Bru 78) of the 700 MeV $^{16}\text{O}(\text{d},\text{d})^{16}\text{O}$ reaction are presented along with the elastic scattering predictions using a "best-fit" phenomenological (dashed line) (see Sect. 2.5), and a KMT (solid line) deuteron optical potential 238
6.4.7	Two step pathways, spectroscopic amplitudes, and deformation parameter for $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$ to the $5/2^+$ and $7/2^+$ levels at 5.2 MeV and 7.2 MeV 240
6.4.8	Angular distributions of the 800 MeV $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$ reaction to the $5/2^+$ (A) and $7/2^+$ (B) levels. Compared to the data are CCBA calculations employing the pathways of Fig. 6.4.7 241
6.5.1	Spectra of the 800 MeV $^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}$ reaction at 5.2° and 12.5° 243
6.5.2	Angular distributions for some of the levels populated in the 800 MeV $^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}$ reaction 245

Figure	Page
6.5.3 Angular distributions for some of the levels populated in the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction	246
6.5.4 Angular distributions for some of the levels populated in the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction	247
6.5.5 Data for the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction to the 0^+ ^{24}Mg ground state are presented with the results of an EFR DWBA calculation employing a KMT deuteron optical potential	249
6.6.1 Sample spectra of the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction at 5.2° and 12.5°	251
6.6.2 Correlation between the negative parity levels of ^{28}Si , and the strongest states excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction, suggesting the importance of weak coupling in the 800 MeV (p,d) reaction	256
6.6.3 Angular distributions for some of the levels excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction	258
6.6.4 Angular distributions for some of the levels excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction	259
6.6.5 Angular distributions for some of the levels excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction	260
6.6.6 Angular distribution for excitation of the ^{27}Si $5/2^+$ ground state, excited with the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction	261
6.7.1 Spectra of the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction at 5.2° and 12.5°	263

Figure		Page
6.7.2	Angular distributions for some of the levels excited in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction	266
6.7.3	Angular distributions for some of the levels excited in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction	267
6.7.4	Data for the excitation of the $3/2^+$ ^{39}Ca ground state, populated in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction, are presented with EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials	269
7.1.1	Sample spectrum of the 175 MeV $^6\text{Li}(\pi^+,p)^5\text{Li}$ reaction at $\theta_L = 20^\circ$. . .	272
7.1.2	Angular distributions of the 75 MeV and 175 MeV $^6\text{Li}(\pi^+,p)^5\text{Li}$ (g.s.) reaction	274
7.2.1	Sample spectrum of the 1975 MeV $^7\text{Li}(\pi^+,p)^6\text{Li}$ reaction at $\theta_L = 20^\circ$. . .	276
7.2.2	Angular distributions for the excitation of the 1^+ ground state and 2.185 MeV 3^+ levels of ^6Li , excited in the $^7\text{Li}(\pi^+,p)^6\text{Li}$ reaction at 75 MeV (A) and 1975 MeV (B)	278
7.2.3	Shown are the 75 MeV (A) and 195 MeV (B) angular distributions of the $^7\text{Li}(\pi^+,p)^6\text{Li}(1^+,3^+)$ reaction, with the results of calculations described in Chapter III for both the 1^+ (solid curve) and 3^+ (dashed curve) levels . .	280
7.3.1	Representative spectrum of the 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reaction collected at $\theta_L = 20^\circ$	282

Figure		Page
7.3.2	Angular distributions of some of the levels excited in the 130 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reaction	286
7.3.3	Energy dependence of the $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ (g.s.) reaction, sampled at 90 MeV (closed circles) and 180 MeV (open circles)	288
7.3.4	The 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ (g.s.) data are presented with the results of calculations for the 90 MeV (solid curve) and 180 MeV (dashed curve) $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ (g.s.) reaction	289
7.4.1	Typical spectrum of the 170 MeV $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ reaction at $\theta_L = 20^\circ$. . .	290
7.4.2	Angular distributions of some of the levels excited in the 170 MeV $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ reaction	293
7.4.3	Energy dependence of the $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ (4.44 MeV) reaction, sampled at 90 MeV (closed circles) and 170 MeV (open circles)	294
7.4.4	The 170 MeV $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ (g.s.) data are presented with the results of calculations for the 90 MeV (solid curve) and 170 MeV (dashed curve) $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ (g.s.) reaction	296
8.3.1	Spectra of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ and 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reactions are compared at a momentum transfer of 2.8 fm^{-1}	304
8.3.2	Spectra of the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ and 170 MeV $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ reactions are compared near a momentum transfer of 2.6 fm^{-1}	305

LARGE MOMENTUM TRANSFER NEUTRON PICKUP
WITH THE (π^+, p) AND (p, d) REACTIONS

by

Gregory Richard Smith

ABSTRACT

The (p, d) reaction has been studied for the first time at 800 MeV on seven targets ranging from ${}^7\text{Li}$ to ${}^{40}\text{Ca}$. The experimental resolution (~ 400 keV) attained was sufficient to observe many discrete levels in each of the residual nuclei. A modified version of the one-nucleon model successfully describes the magnitude and angular dependence of almost all of the transitions observed. A specific counter example to the two-nucleon model of the reaction mechanism is suggested. The calculations are also sensitive to the neutron single particle wavefunction, in accordance with the expectation that the high momentum components of this wavefunction are probed at higher bombarding energies. States which have never been seen before were strongly populated in the high excitation region (up to 25 MeV) of some of the residual nuclei. The relative intensities of the other levels observed suggest that coupled channels

mechanisms play an important role for some of these states. Explicit calculations have been performed to confirm this for several examples.

The first high resolution measurements of the (π^+, p) reaction have also been performed on ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^{12}\text{C}$ and ${}^{13}\text{C}$ at pion bombarding energies on and off the pion-nucleon resonance. Calculations employing a one-nucleon model of the reaction mechanism similar to the model successfully used for the (p, d) reaction are unable to account for transitions in the (π^+, p) reaction. It is, however, unclear whether this failure is due to a fundamental inadequacy of the model or improper treatment of details in the calculations. A striking similarity has been observed in the spectra of the (π^+, p) and 800 MeV (p, d) reactions on the same target, implying a similar mechanism for the two reactions.

CHAPTER I

INTRODUCTION

The advent of accelerators (such as LAMPF) which provide intermediate energy beams of protons and pions, coupled with the existence of spectrometers (such as HRS and EPICS) with sufficient resolution (≈ 100 keV) to resolve discrete nuclear levels, opens the door to an exciting new realm of nuclear physics. The shorter wavelengths of the proton probe (for example, an 800 MeV proton has a De Broglie wavelength of only 0.85 fm as opposed to 5.5 fm for a 30 MeV proton, in a medium where the average distance between nucleons is about 1.8 fm) has nourished the hope that new features of the nuclear system may be discovered and explored at intermediate energies. The bulk of the experimental data taken to date with intermediate energy projectiles has been focused on elastic and inelastic scattering to collective states—measurements which are mainly sensitive to the gross features of the nuclear density. More advantageous use of the short wavelengths of intermediate energy projectiles may be found in rearrangement reactions such as neutron pick-up, which sample the wavefunction of a single nucleon. In addition, reactions such as (p,d)

at intermediate energies require extremely large (linear) momentum transfers $q = |\vec{p}_p - \vec{p}_d|$, which must be supplied by the nucleus. These large momentum transfers therefore provide a means of sampling the high momentum components of the nucleus in the heretofore inaccessible region of momentum transfer $q \approx 2 \text{ fm}^{-1} - 5 \text{ fm}^{-1}$. Large momentum transfers may indeed be achieved in low energy neutron pick-up reactions such as $(^3\text{He}, \alpha)$ by increasing the scattering angle. However, the effects of distortion of the projectile and ejectile cloud the picture presented by these reactions, and these distortion effects increase in severity as the scattering angle is increased. Thus, in spite of the rapidly falling cross sections associated with increased projectile energy, the intermediate energy (800 MeV) (p, d) reaction may prove to be a most fruitful tool with which to study large momentum transfer neutron pickup.

One reason for studying the (p, d) reaction at intermediate energies is based on an assumption which seems to be valid at lower energies, namely, that the reaction probes neutron single particle wavefunctions. In particular, as the energy increases, higher momentum components of these wavefunctions assume increasing importance. Figure 1.1 shows the Fourier transform of the neutron bound state wavefunction, $\tilde{\phi}(q)$,

Fig. 1.1 Fourier transform of the $p_{3/2}$ neutron bound state wave function used in the analysis of the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction, plotted as a function of the momentum transfer q . Also shown are the regions of q sampled by the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction at various bombarding energies.

plotted as a function of momentum q . Also shown in this figure are the regions of momentum transfer sampled by the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction at several bombarding energies. The 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction clearly samples $\tilde{\phi}(q)$ at much higher q than the lower energy reactions. Any information one can obtain about these high q components is of considerable interest since that is where the details of the wavefunction are to be found. (The shell model quantum numbers and the nuclear radius are generally sufficient to determine $\tilde{\phi}(q)$ up to $\sim 1.5 \text{ fm}^{-1}$.)

The extraction of such information from 800 MeV (p,d) data depends on having a quantitative model of the reaction mechanism, whether it be the low energy model or something else.

Other features of interest may be accessible with the 800 MeV (p,d) reaction. For instance, new classes of final states might be excited at these larger momentum transfers. High spin states are expected to be preferentially populated because of the large momentum transfer, revealing, perhaps, new spectroscopic information. However, such spectroscopic information can only be obtained after the reaction mechanism is understood. This is another reason why major effort to understand the mechanism of neutron pick-up by the (p,d) reaction at 800 MeV is

warranted. This understanding is most easily gained by analysis of data for which the nuclear structure aspects of the reaction process are well known. These test cases should span a wide range in mass number, to reveal any systematic effects in the A-dependence of the reaction. Furthermore, they should have been extensively studied with the low energy (p,d) reaction. Many such low energy reaction studies are available in the literature.

The (π^+ ,p) reaction, in as much as it results in the removal of a single nucleon from a target nucleus, should be related to the intermediate energy (p,d) reaction. The comparable (large) asymptotic momentum transfers for the two reactions give some reason to expect considerable similarity between the two. If similarity can be established experimentally, for example by looking at results on the same targets, then it can possibly be exploited to help in understanding both the (p,d) and the (poorly understood) (π^+ ,p) reaction mechanisms. Once that is accomplished the two reactions might then be utilized as nuclear structure probes.

Data presented in this dissertation will establish a qualitative similarity between the (p,d) and (π^+ ,p) reactions. Both will be seen to excite nuclear levels unseen in low energy neutron pickup reactions.

The analysis of the 800 MeV (p,d) reaction presented here will show that nearly all the observed transitions to well known levels can be quantitatively explained using a somewhat refined version of the reaction model generally applied to the low energy (p,d) reaction. A similar model will be seen to be unable to account for similar transitions in the (π^+ ,p) reaction, although it is unclear whether the failure is due to a fundamental inappropriateness of the model or improper treatment of details in the calculations.

The thesis is organized so as to present the theoretical development of the reaction models first, followed by a description of the experimental techniques and a presentation of the data themselves. Finally the data are compared with the theoretical predictions and conclusions and a summary are presented.

CHAPTER II

DWBA THEORY AND CALCULATIONS

The purpose of this chapter is to develop the theoretical framework of the distorted wave Born approximation (DWBA) which will be tested in a new regime by comparison to 800 MeV (p,d) data on a variety of nuclei. At lower bombarding energies (≤ 100 MeV) the dominant mechanism for the (p,d) reaction is well understood: the incident proton encounters a target consisting of a core and a neutron bound in a shell-model orbital. The proton picks up the neutron, forming a deuteron, and leaves the residual (core) nucleus in one of its allowed states depending on the particular shell-model orbital of the picked-up neutron. Using this description (the one nucleon model or ONM) many features of nuclear structure have been investigated, such as single-particle quantum numbers and parentage of nuclear levels.

At higher bombarding energies like 800 MeV, however, this simple description may be questioned (Käl+ 78), (RSS 78), (CW 69). For example, real pion degrees of freedom are possible and may be of importance. Consequently, the ONM will be tested in this energy regime for cases where the nuclear structure is

well understood. In this way, the reaction mechanism and single nucleon DWBA treatment may be tested.

The ONM is typically represented schematically by a diagram such as that shown in Fig. 2.1. In this picture, the upper-left line represents the incoming proton, the lower-left lines represent the target nucleons. The outgoing deuteron is drawn as the two lines in the upper right, the residual nucleus is depicted by the lines on the lower right, and the picked-up neutron by the transverse line. The upper vertex connecting the proton, deuteron, and picked-up neutron contains information relevant to the reaction mechanism. The lower vertex contains the nuclear structure information. The wavy lines represent the general (non-rearrangement) interaction of the projectile (ejectile) with the target (residual) nucleus before (after) the neutron pick-up process. The consequence of such an interaction is referred to as distortion. The following section will develop the content of this diagram in detail.

This chapter, then, will first present the formalism of the DWBA and discuss the limits of its validity. Next, the distortion is accounted for using an optical potential (see Sects. 2.5, 2.6, and 2.7) which may (or may not) be constrained to fit elastic scattering data. Finally, relevant extensions and

Fig. 2.1 Schematic representation of the One Nucleon Model, as explained in the text.

modifications of the standard DWBA treatment will be discussed, and the sensitivity of the results to parameters of the theory will be investigated. The predictions of this theory will be compared to data in Chapter VI, and Chapter VIII will summarize the conclusions.

PART A: Theory of the DWBA

2.1 Formalism

The theory which will be discussed in this chapter will be relevant to the $A(p,d)B$ reaction at 800 MeV. In this notation, the target nucleus (A) consists of a neutron (n) bound to a core (B). The deuteron (d) consists of a neutron (n) and proton (p) bound by 2.22 MeV. The Schroedinger equation for the complete system is

$$H\Psi_{\alpha}^{(+)} = E\Psi_{\alpha}^{(+)} \quad (2.1.1)$$

where $\Psi_{\alpha}^{(+)}$ is the total wavefunction for the system with outgoing (+) wave boundary conditions. These boundary conditions specify that an incoming plane-wave is present in the $p + A$ (or α) partition and that outgoing spherical waves are present asymptotically in all other partitions. The Hamiltonian is

$$H = (H_i = H_{\alpha} + T_{\alpha} + V_{\alpha}) = (H_f = H_{\beta} + T_{\beta} + V_{\beta}) \quad (2.1.2)$$

Here H_β represents the internal Hamiltonian in partition β , and has associated with it a product eigenfunction

$$\phi_\beta(\vec{\xi}_\beta) = \phi_d(\vec{\xi}_d) \phi_B(\vec{\xi}_B) \quad (2.1.3)$$

which satisfies

$$H_\beta \phi_\beta = E_\beta \phi_\beta \quad (2.1.4)$$

where $E_\beta = E_d + E_B$.

In the following, the displacement variable $\vec{r}_\beta$ denotes the separation of the centers of mass of B and d. The variable $\vec{r}_\alpha$ is similarly defined. Internal coordinates will be denoted by $\vec{\xi}_\beta$ or $\vec{\xi}_\alpha$. Fig. 2.1.1 summarizes the definitions of the coordinates involved. The development which follows uses non-relativistic kinematics—a relativistic generalization will be presented later.

Now the total energy of the system may be expressed as

$$E = E_\beta + \frac{\hbar^2 k_\beta^2}{2\mu_\beta} \quad (2.1.5)$$

where μ_β is the reduced mass and $\hbar k_\beta$ is the relative momentum in partition β . The wavefunction ψ_β of relative motion in partition β may now be introduced by taking the overlap between the total wavefunction $\Psi_\alpha^{(+)}$

Fig. 2.1.1 Reaction coordinate definitions and labels for the constituents of the (p,d) reaction.

and the internal wavefunction ϕ_β ,

$$\psi_\beta = \langle \phi_\beta | \psi_\alpha^{(+)} \rangle \quad (2.1.6)$$

The integration in the bracket notation will be assumed to be over internal coordinates only unless otherwise stated. After multiplication from the left by ϕ_β and integration over $\vec{x}_\beta$, Eq. (2.1.1) becomes

$$(\epsilon_\beta - T_\beta) \psi_\beta = \langle \phi_\beta | V_\beta | \psi_\alpha^{(+)} \rangle \quad (2.1.7)$$

where

$$\epsilon_\beta = E - E_\beta = \frac{\hbar^2 k_\beta^2}{2\mu_\beta} \quad (2.1.8)$$

With the identification of

$$T_\beta = \frac{-\hbar^2}{2\mu_\beta} \nabla_{\vec{x}_\beta}^2 \quad (2.1.9)$$

Eq. (2.1.7) takes the form of the inhomogeneous Helmholtz wave equation (Jac 62),

$$(\nabla_{\vec{x}_\beta}^2 + k_\beta^2) \psi_\beta = \frac{2\mu_\beta}{\hbar^2} \langle \phi_\beta | V_\beta | \psi_\alpha^{(+)} \rangle \quad (2.1.10)$$

This well-known equation may be solved with the Green's function (Jac 62), (Aus 70)

$$G(\vec{r}_\beta, \vec{r}'_\beta) = \frac{2\mu_\beta}{\hbar^2} \frac{e^{ik_\beta |\vec{r}_\beta - \vec{r}'_\beta|}}{4\pi |\vec{r}_\beta - \vec{r}'_\beta|} \quad (2.1.11)$$

where

$$(\nabla_{\vec{r}_\beta}^2 + k_\beta^2) G(\vec{r}_\beta, \vec{r}'_\beta) = -\delta(\vec{r}_\beta - \vec{r}'_\beta) \quad (2.1.12)$$

The asymptotic solution of (2.1.10) may then be expressed in the limit $r_\beta \gg r'_\beta$ as (Aus 70)

$$\psi_\beta \rightarrow e^{i\vec{k}_\beta \cdot \vec{r}_\beta} \delta_{\alpha\beta} - \frac{\mu_\beta}{2\pi\hbar^2} \frac{e^{ik_\beta r_\beta}}{r_\beta} \langle \phi_\beta e^{i\vec{k}_\beta \cdot \vec{r}'_\beta} | V_\beta | \psi_\alpha^{(+)} \rangle \quad (2.1.13)$$

where the integration now includes r'_β as well as $\vec{\xi}_\beta$. Another expression for the asymptotic form of ψ_β may be obtained by invoking the boundary conditions of the problem discussed following Eq. (2.1.1),

$$\psi_\beta \xrightarrow{r_\beta \rightarrow \infty} e^{i\vec{k}_\alpha \cdot \vec{r}_\alpha} \delta_{\alpha\beta} + f(\vec{k}_\alpha, \hat{r}_\beta) \frac{e^{ik_\beta r_\beta}}{r_\beta} \quad (2.1.14)$$

In this expression, the amount of incident flux scattered into partition β is modulated by the scattering amplitude $f(\vec{k}_\alpha, \hat{r}_\beta)$, from which all observables (cross-sections, polarizations, etc.) are obtained. The scattering amplitude depends upon the incident momentum $\vec{k}_\alpha$, as well as the scattering angle $\hat{r}_\beta$. An

exact expression for $f(\vec{k}_\alpha, \hat{r}_\beta)$ may be obtained by a direct comparison of the two asymptotic formulae for ψ_β , Eqs. (2.1.13) and (2.1.14), yielding

$$f(\vec{k}_\alpha, \hat{r}_\beta) = \frac{-\mu_\beta}{2\pi\hbar^2} \langle \phi_\beta e^{i\vec{k}_\beta \cdot \vec{r}'_\beta} | V_\beta | \psi_\alpha^{(+)} \rangle \quad (2.1.15)$$

The transition amplitude $T_{\alpha\beta}$ is introduced to simplify algebraically the scattering amplitude, viz.,

$$T_{\alpha\beta} = \frac{-2\pi\hbar^2}{\mu_\beta} f(\vec{k}_\alpha, \hat{r}_\beta) \quad (2.1.16a)$$

$$= \langle \phi_\beta e^{i\vec{k}_\beta \cdot \vec{r}'_\beta} | V_\beta | \psi_\alpha^{(+)} \rangle \quad (2.1.16b)$$

This form of the transition amplitude contains the interaction V_β with a total scattering solution $\psi_\alpha^{(+)}$ containing the plane wave

$$e^{i\vec{k}_\alpha \cdot \vec{r}_\alpha}$$

as a boundary condition. This is referred to as the "post" form of $T_{\alpha\beta}$. An equivalent form of $T_{\alpha\beta}$ is obtained by using $\psi_\beta^{(-)}$, giving

$$T_{\alpha\beta} = \langle \psi_\beta^{(-)} | V_\alpha | e^{i\vec{k}_\alpha \cdot \vec{r}_\alpha} \phi_\alpha \rangle \quad (2.1.17)$$

This is referred to as the "prior" form. The choice of which form to use is generally dictated by

convenience, as will be discussed below.

The explicit form of V_α or V_β must now be examined. In analogy with (2.1.2),

$$V_\alpha = H - T_\alpha - H_\alpha \quad (2.1.18)$$

The Hamiltonian H_α describing the internal motion in the $p + A$ partition is simply

$$H_\alpha = H_B + V_{n,B} + T_{n,B} \quad (2.1.19)$$

The entire system is described by the Hamiltonian

$$H = H_B + T_{n,B} + T_{p,A} + V_{n,B} + V_{p,B} + V_{p,n} \quad (2.1.20)$$

which requires

$$V_\alpha = V_{p,B} + V_{p,n} \quad (2.1.21)$$

The form of $T_{\alpha\beta}$ may be altered by the introduction of a potential U_i which is arbitrary apart from the stipulation that it depend only on the channel displacement variable $\vec{r}_i$ (and hence cannot introduce rearrangement in channel i). Using the Gell-Mann, Goldberger transformation (GG 53), (Aus 70), (Mes 63), we can write

$$T_{\alpha\beta} = \langle \psi_\beta^{(-)} | V_\alpha - U_\alpha | \phi_\alpha \chi_\alpha^{(+)} \rangle \text{ (prior)} \quad (2.1.22a)$$

$$= \langle \phi_\beta \chi_\beta^{(-)} | V_\beta - U_\beta | \psi_\alpha^{(+)} \rangle \text{ (post)} \quad (2.1.22b)$$

where the distorted waves $\chi_i^{(+)}$ satisfy

$$(T_i + U_i) \chi_i^{(+)} = E_i \chi_i^{(+)} \quad (2.1.23)$$

with incoming wave boundary conditions. The distorted wave χ_i is observed to reduce to the plane-wave eigenfunction $\exp(i\vec{k}_i \cdot \vec{r}_i)$ in the limit $U_i \rightarrow 0$. The utility of this form of $T_{\alpha\beta}$ is that the effects of the nuclear distortion can be approximated by the potentials U_i as will be discussed in detail below.

The exact expressions (2.1.22) for $T_{\alpha\beta}$ are not useful since the total wavefunction $\psi_i^{(+)}$ cannot be calculated. However, an approximate expression which is numerically tractable is obtained by expanding the exact wavefunction ψ_i in a distorted wave series. Formally, this may be accomplished by substituting for ψ_i the formal solution of the Lippmann-Schwinger equation (Sto+ 67),

$$\langle \psi_\beta^{(-)} | = \langle \chi_\beta^{(-)} | \phi_\beta \{ 1 + (V_{nB} + V_{PB} - U_\beta)(E - H)^{-1} \} \quad (2.1.24)$$

The distorted wave Born approximation (DWBA) consists of using the first term above so that (Tob 61), (Aus 70)

$$\langle \Psi_{\beta}^{(-)} | \approx \langle \chi_{\beta}^{(-)} \phi_{\beta} | \quad (2.1.25)$$

which gives

$$T_{\alpha\beta}^{DWBA} \approx \langle \chi_d^{(-)} \phi_d \phi_B | V_{Pn} + V_{PB} - U_{PA} | \chi_P^{(+)} \phi_P \phi_A \rangle \quad (2.1.26)$$

in the prior form, where the bound state wavefunctions have been explicitly identified. This expression can be further simplified by choosing the potential U_{PA} so as to cancel as completely as possible V_{PB} . This gives the standard form of $T_{\alpha\beta}^{DWBA}$ for the $A(p,d)B$ reaction:

$$T_{\alpha\beta}^{DWBA} = \langle \phi_B \phi_d \chi_d^{(-)} | V_{Pn} | \chi_P^{(+)} \phi_P \phi_A \rangle \quad (2.1.27)$$

The utility of choosing the prior for of the transition amplitude is now readily seen. First, the V_{Pn} interaction can be approximated in zero range (as discussed in Sect. 2.3) which considerably simplifies (by one less three dimensional integration) expression (2.1.27). Additionally, in the post form the transition amplitude involves

$$V_{\beta} - U_{\beta} = V_{nB} + V_{PB} - U_{\beta} \quad (2.1.28)$$

The simplest approximation for U_{β} in (2.1.28) would make the transition amplitude vanish.

Implicit in Eq. (2.1.27) are two assumptions, namely (Sto+ 67)

$$||V_{PB} - U_\alpha|| \approx 0 \quad (2.1.29a)$$

and

$$||V_{PB} + V_{nB} - U_\beta|| \approx 0 \quad (2.1.29b)$$

which come from (2.1.24) and (2.1.26), respectively. Aside from the minor restrictions mentioned above, the potentials U_i are arbitrary. Hence they could be chosen so as to satisfy these conditions and thereby maximize the accuracy of the DWBA. The physical basis for the choice of the U_i 's will be addressed in detail in Sects. 2.5, 2.6, and 2.7.

Introducing the coordinates explicitly, the transition amplitude may now be written

$$T_{\alpha\beta}^{DW} = \langle \chi_d^{(-)}(\vec{k}_d, \vec{r}_d) \phi_B(\vec{\xi}_B) \phi_d(\vec{r}_{Pn}) | V_{Pn}(\vec{r}_{Pn}) | \phi_A(\vec{\xi}_B, \vec{r}_n) \phi_P(\vec{\xi}_P) \chi_P^{(+)}(\vec{k}_P, \vec{r}_P) \rangle \quad (2.1.30)$$

where the coordinates are those of Fig. 2.1.1.

The next step in the evaluation of the transition amplitude is to integrate over the internal coordinates of the target and residual nuclei. In

(2.1.30) the internal coordinates $\vec{\xi}_A$ of the target wavefunction ϕ_A have been broken up into two parts. Coordinates $\vec{\xi}_B$ represent the internal coordinates of the core, and coordinates $\vec{r}_n$ describe the position of the picked-up neutron with respect to the center of

mass of the core (see Fig. 2.1.1). The target wavefunction ϕ_A may now be expanded in terms of the core wavefunction and the wavefunction of the neutron which is to be picked up,

$$\phi_A(\vec{\xi}_A) = \phi_{M_A}^{J_A}(\vec{\xi}_B, \vec{r}_n) \quad (2.1.31)$$

$$= \sum_{J, J_n} \sqrt{S_{J_n J}^{J_A}} [\phi^J(\vec{\xi}_B) \phi^{J_n}(\vec{r}_n)]_{M_A}^{J_A}$$

where the square of the expansion coefficient $S_{J_n J}^{J_A}$ is called the spectroscopic factor, and represents the probability of finding the neutron in the core + neutron system with angular momentum quantum number J_n . The spectroscopic amplitude

$$\sqrt{S_{J_n J}^{J_A}}$$

is defined by (She 78)

$$\sqrt{S_{J_n J}^{J_A}} = \sqrt{n_1} \langle \psi_{J_1} || (a_j^\dagger \psi_{J_2})_{J_1} \rangle \quad (2.1.32)$$

where $J_1 = J_i$, $J_2 = J_f$ for pickup, and vice versa for stripping. $a_j^\dagger$ is the standard creation operator for a nucleon of angular momentum j . The double bars indicate a reduced matrix element defined by the Wigner-Ekhardt theorem in the notation of Rost (Ros 79)

$$\langle \phi_{M_1}^{J_1} | T_Q^K | \chi_{M_2}^{J_2} \rangle = \langle K J_2 Q M_2 | J_1 M_1 \rangle \langle \phi^{J_1} | [T^K \chi^{J_2}]^{J_1} \rangle \quad (2.1.33)$$

The spectroscopic factors used for the (p,d) studies reported here were obtained using a procedure due to Shepard (She 79), in which the coefficients of fractional parentage were computed from the wave functions of Norton and Goldhammer (NG 70), (Nor 70). Spectroscopic factors obtained using this procedure are compared to the spectroscopic factors of Cohen and Kurath (CK 67) in Table 2.1.1. The good agreement between the two predictions indicates that we are indeed testing the reaction mechanism for cases of known nuclear structure.

Now that the spectroscopic factors have been determined, the integration of (2.1.28) over $\vec{\xi}_B$ may be performed, yielding (Emi 76)

$$\begin{aligned} T_{\alpha\beta}^{DW} = & \sum_{J_n J_B} \sqrt{S_{J_n J_B}^{J_A}} \langle J_B J_n M_B M_A - M_B | J_A M_A \rangle \\ & \times \int d\vec{r}_n d\vec{r}_p \chi^{*(-)}(\vec{k}_d \vec{r}_d) \phi_d^*(\vec{r}_{pn}) V_{pn}(\vec{r}_{pn}) \\ & \phi_{M_A - M_B}^{J_n}(\vec{r}_n) \phi_p(\vec{\xi}_p) \chi^{(+)}(\vec{k}_p, \vec{r}_p) \end{aligned} \quad (2.1.34)$$

Table 2.1.1. $^{13}\text{C}(p,d)^{12}\text{C}$ Spectroscopic Factors

J_i	J_f	T_f	j_{tr}	$C^2S(N+G)\underline{a/}$	$C^2S(C.+K.)\underline{b/}$
$1/2^-$	0^+	0	p1/2	0.674	0.613
	2^+	0	p3/2	1.038	1.112
	1^+	0	p1/2	0.022	0.001
	1^+	0	p3/2	0.416	0.661
	1^+	1	p1/2	0.011	0.002
	1^+	1	p3/2	0.667	0.600
	2^+	1	p3/2	1.150	1.012
$3/2^-$	2^+	0	p1/2	0.765	
	1^+	0	p1/2	0.339	
	1^+	1	p1/2	0.210	
	2^+	1	p1/2	0.056	
	0^+	0	p3/2	0.133	
	2^+	0	p3/2	0.290	
	1^+	0	p3/2	0.000	
	1^+	1	p3/2	0.029	
	2^+	1	p3/2	0.051	
$5/2^-$	2^+	0	p1/2	0.310	
	2^+	1	p1/2	0.374	
	2^+	0	p3/2	0.048	
	1^+	0	p3/2	0.043	
	4^+	0	p3/2	1.358	
	1^+	1	p3/2	0.011	
	2^+	1	p3/2	0.003	

a/ Spectroscopic factors computed from the wavefunctions of Norton and Goldhammer (NGo 70), (Nor 70).

b/ Spectroscopic factors from the tables of Cohen and Kurath (CKu 67).

The differential cross section for the transition from the incident channel α to the outgoing channel β is the ratio of the outgoing flux in channel β per unit solid angle to the incident flux per unit area. In terms of the transition amplitude (Tob 61), (Aus 70),

$$\frac{d\sigma}{d\Omega}(\alpha \rightarrow \beta) = \frac{k_\beta}{k_\alpha} \frac{\mu_\alpha \mu_\beta}{(2\pi\hbar)^2} |T_{\alpha\beta}|^2 \quad (2.1.35)$$

Since the incident proton beam used for the (p,d) studies was unpolarized, and the polarization of the deuteron was not detected, the initial spins are averaged over, and the final spins summed over. The final result is (Emi 76)

$$\frac{d\sigma}{d\Omega}(\alpha \rightarrow \beta) = \frac{k_\beta}{k_\alpha} \frac{\mu_\alpha \mu_\beta}{(2\pi\hbar)^2} \frac{2J_B+1}{2J_A+1} \sum_{J_n, M} \frac{S_{J_n J_B}^{J_A}}{2(2J_n+1)} |B_M^{J_n}(\theta)|^2 \quad (2.1.36)$$

where

$$B_M^{J_n}(\theta) = \int d\vec{r}_n d\vec{r}_p \chi_d^{*(-)}(\vec{r}_d) \phi_d^*(\vec{r}_{pn}) v_{pn}(\vec{r}_{pn}) \phi_M^{J_n}(\vec{r}_n) \chi_p^{(+)}(\vec{r}_p) \quad (2.1.37)$$

It should be noted that 800 MeV protons are relativistic, having $\gamma = E_{\text{total}}/E_{\text{rest}} = 1.85$. Consequently, for all the calculations presented, relativistic kinematics were used, with the following

substitutions in the usual (nonrelativistic) Schroedinger equation (FGH 70), (C7 64)

$$\frac{2\mu E}{\hbar^2} = \frac{2E}{\hbar^2 c^2} \frac{1}{1 + \frac{E}{Amc^2}} \quad (2.1.38a)$$

and

$$k^2 + \frac{E^2 - m^2 c^4}{\hbar^2 c^2} \frac{1}{1 + \frac{E}{Amc^2}} \quad (2.1.38b)$$

where E is the total center of momentum energy of the proton-target nucleus system, m is the nucleon mass, and A is the atomic weight of the target nucleus.

Finally the complications of spin and Coulomb scattering at large distances were taken care of in the usual way.

PART B: Techniques of the DWBA

2.2 The PWBA Limit

The transition amplitude may be expressed in a simple, analytic form by ignoring the effects of distortion, thereby utilizing the plane wave Born approximation (PWBA). In this approximation, the physical roles played by the various factors appearing in the transition amplitude are simply delineated, making possible a clearer interpretation of the underlying physics.

In the previous section, just prior to the introduction of distortion via the optical potentials, the transition amplitude was shown to be

$$T_{\alpha\beta} = \langle \psi_{\beta}^{(-)} | V_{\alpha} | e^{i\vec{k}_{\alpha} \cdot \vec{r}_{\alpha}} \phi_{\alpha} \rangle \quad (2.2.1)$$

in the prior representation. This may be broken into two terms, reflecting the two constituents of V_{α} ,

$$T_{\alpha\beta} = \langle \psi_{\beta}^{(-)} | V_{PN} | e^{i\vec{k}_{\alpha} \cdot \vec{r}_{\alpha}} \phi_{\alpha} \rangle + \langle \psi_{\beta}^{(-)} | V_{PB} | e^{i\vec{k}_{\alpha} \cdot \vec{r}_{\alpha}} \phi_{\alpha} \rangle . \quad (2.2.2)$$

The PWBA consists of two drastic approximations.

First, the total wave function in the outgoing channel, $\langle \psi_{\beta}^{(-)} |$, is replaced by the first term in the Born series, analagous to (2.1.24),

$$\langle \psi_{\beta}^{(-)} | \approx \langle e^{+i\vec{k}_d \cdot \vec{r}_d} \phi_{\beta} | . \quad (2.2.3)$$

Here the wave function of relative motion in channel β is the plane wave $\exp(+i\vec{k}_d \cdot \vec{r}_d)$, instead of the distorted wave χ_{β} (as noted following (2.2.23) for the case of no distortion). The second approximation is to ignore the contributions from the second term of (2.2.2).

In the PWBA, the transition amplitude becomes

$$T_{\alpha\beta}^{Pw} = \langle e^{+i\vec{k}_d \cdot \vec{r}_d} \phi_d \phi_B | V_{Pn} | e^{i\vec{k}_p \cdot \vec{r}_p} \phi_p \phi_A \rangle . \quad (2.1.4)$$

This expression for $T_{\alpha\beta}$ is identical in form to the DWBA embodied in (2.1.26), except that plane waves have replaced the distorted waves χ_i .

The internal wavefunctions of the target and residual nucleus may be integrated out exactly as in (2.1.29), leaving

$$T_{\alpha\beta}^{Pw} = \sum_{J_n J_B} \langle J_B J_n M_B M_A - M_B | J_A M_A \rangle \sqrt{\frac{S_{J_B J_n}^{J_A}}{S_{J_B J_n}^{J_A}}} \\ \langle e^{i\vec{k}_d \cdot \vec{r}_d} \phi_d(\vec{r}_{Pn}) | V_{Pn}(\vec{r}_{Pn}) | \phi_n(\vec{r}_{nc}) \phi_p e^{i\vec{k}_p \cdot \vec{r}_p} \rangle . \quad (2.2.5)$$

Integrating out the proton spinor, one is left with an integral including two vector variables. The transition amplitude may be factorized by expressing r_p and r_d in terms of the variables r_{pn} and r_{nb} ,

$$\vec{r}_p = \vec{r}_{pA} = \vec{r}_{pn} + \frac{N-1}{N} \vec{r}_{nb} \quad (2.2.6)$$

$$\vec{r}_d = \vec{r}_{dB} = \vec{r}_{nb} + \frac{1}{2} \vec{r}_{pn}$$

where N ($N-1$) denotes the atomic weight of the target (residual) nucleus, respectively, and the

factor 1/2 is the ratio of the proton mass to the deuteron mass. The resulting factorized expression is

$$T_{\alpha\beta}^{PW} = \sum_{J_n J_B} \sqrt{S_{J_n J_B}^{J_A}} \langle J_B J_n M_B M_A - M_B | J_A M_A \rangle$$

$$\int d\vec{r}_{Pn} e^{i\vec{q} \cdot \vec{r}_{Pn}} V_{Pn}(\vec{r}_{Pn}) \phi_d(\vec{r}_{Pn}) \int d\vec{r}_{nB} e^{i\vec{Q} \cdot \vec{r}_{nB}} \phi_n(\vec{r}_{nB})$$

(2.2.7)

where the momentum transfers $\vec{q}$ and $\vec{Q}$ are

$$\vec{q} = \vec{k}_p - \frac{1}{2} \vec{k}_d$$

$$\vec{Q} = \frac{N-1}{N} \vec{k}_p - \vec{k}_d$$

(2.2.8)

The physical significance of the two integrals appearing in (2.2.7) is quite clear. The Fourier transform of $V_{np}\phi_d$ is identified as the reaction dynamics term, and corresponds to the upper vertex in the ONM diagram of Fig. 2.1. The lower vertex in the same figure contains all of the nuclear structure information, and is the Fourier transform of the neutron bound state wavefunction.

The following two sections will discuss the evaluation of $V_{np}\phi_d$ and ϕ_n .

2.3 Deuteron Wave Function

In the previous section, the plane wave transition amplitude was shown to be proportional to the product of two Fourier transforms. The first of these, the Fourier transform of $V_{pn}\phi_d$, will now be discussed in detail.

An estimate of the behavior of the Fourier transform of $V_{pn}\phi_d$ may be obtained by utilizing the Hulthén form for the S-state part of the deuteron wave function (Aus 70),

$$\phi_d = \left(\frac{\alpha\beta(\alpha+\beta)}{2\pi(\alpha-\beta)^2} \right)^{1/2} \left(\frac{e^{-\alpha\rho}}{\rho} - \frac{e^{-\beta\rho}}{\rho} \right), \quad (2.3.1)$$

where $\vec{\rho} = \vec{r}_p - \vec{r}_n$, the proton-neutron separation. The parameters α and β determine the long and short range behavior of ϕ_d , respectively. In the limit of large n - p separation, $V_{pn} \rightarrow 0$ and $\phi_d \rightarrow N \frac{e^{-\alpha\rho}}{\rho}$.

Insertion into the Schroedinger equation then yields

$$\frac{-\hbar^2 \alpha^2}{2\mu} = E_d = -2.225 \text{ MeV}. \quad (2.3.2)$$

This determines the value of α , namely $\alpha = 0.235 \text{ fm}^{-1}$. The parameter β is determined by fitting the r.m.s. charge radius of the deuteron,

$$\begin{aligned}
\langle r_p^2 \rangle_{\text{point}} &= \frac{1}{4} \frac{\int \rho^2 \phi_d^* \phi_d d\vec{\rho}}{\int \phi_d^* \phi_d d\vec{\rho}} \\
&= \frac{1}{2} \left(\frac{\frac{4}{(\alpha+\beta)^3} - \frac{\beta^3 + \alpha^3}{4\alpha^3\beta^3}}{\frac{2}{(\alpha+\beta)} - \frac{\beta+\alpha}{2\alpha\beta}} \right)^{\frac{1}{2}}, \quad (2.3.3)
\end{aligned}$$

where r_p is the separation of the proton from the deuteron center of mass: $\vec{r}_p = \frac{1}{2}\vec{\rho}$. The relationship between the r.m.s. charge radius and the r.m.s. point radius is

$$\langle r^2 \rangle_{\text{ch}}^{\frac{1}{2}} = \sqrt{\langle r^2 \rangle_{\text{proton}} + \langle r_p^2 \rangle_{\text{point}}} \quad (2.3.4)$$

Using $\langle r^2 \rangle_{\text{proton}}^{\frac{1}{2}} = 0.78$ fm, and $\langle r^2 \rangle_{\text{ch}}^{\frac{1}{2}} = 2.09$ fm, the value of β may be found from successive iteration of (2.3.8), yielding a value of $\beta = 1.61$ fm⁻¹.

The Fourier transform of this wavefunction may now be evaluated. Defining the quantity

$$D(\vec{r}_{pn}) = v_{pn}(\vec{r}_{pn}) \phi_d(\vec{r}_{pn}), \quad (2.3.5)$$

one has

$$\begin{aligned}
\tilde{D}(\vec{q}) &= \int e^{i\vec{q} \cdot \vec{r}} D(\vec{r}) d\vec{r} \\
&= \int e^{i\vec{q} \cdot \vec{r}} D(r) y_{00}(\hat{r}) d\vec{r}
\end{aligned}$$

$$\begin{aligned}
&= \sqrt{4\pi} \int r^2 dr D(r) \frac{\sin qr}{qr} \\
&= \sqrt{4\pi} \int r^2 \frac{\sin qr}{qr} \left(\frac{-\hbar^2}{2\mu} (\alpha^2 + \nabla^2) \phi_d(r) \right) dr \\
&= \frac{-\hbar^2 N_d \sqrt{4\pi}}{2\mu} \frac{\beta^2 - \alpha^2}{\beta^2 + q^2} \quad (2.3.6)
\end{aligned}$$

where N_d is the Hulthén normalization of (2.3.1). $\tilde{D}(q)$ is plotted vs. q in Fig. 2.3.1. This quantity is negative, reflecting the attractive nature of V_{pn} , has a (maximum) magnitude at $q = 0$ of $\tilde{D}(0) = -.24 \text{ MeV} - \text{fm}^{3/2}$ and drops asymptotically to zero for large q .

The Hulthén wave function is of a simple analytic form and has a purely $L = 0$ character. A more realistic wavefunction which also includes the D-state component of the deuteron (required to provide the observed non-zero static quadrupole moment of the deuteron), was calculated by Rost and Shepard (RS 75). The procedure used was to write

$$[V_{pn}(\vec{r}) \phi_d(\vec{r})]_M^1 = D_0(\vec{r}) [Y^0(\hat{r}) \chi^1]_M^1 + D_2(\vec{r}) [Y^2(\hat{r}) \chi^1]_M^1, \quad (2.3.7)$$

where the deuteron wavefunction has been broken up into its S and D states; χ denotes the deuteron spin function, and the square brackets imply vector

Fig. 2.3.1 Fourier transform of $V_{pn}(r_{pn})\phi_d(r_{pn})$ vs. momentum transfer q . The solid (dashed) line is the $\tilde{D}(q)$ obtained from Eq. (2.3.9a) (2.3.9b) corresponding to the deuteron S-state (D-state). The dash-dot line is the $\tilde{D}(q)$ obtained from the Hulthén wave function (S-state only) discussed in the text. The dotted line shows the zero range approximation. The energies indicate the region of q sampled by the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction at four bombarding energies.

FOURIER TRANSFORM
of $V_{pn} \phi_d$ vs q

coupling. The n - p potential was chosen to be of the Reid soft-core form

$$V_{pn}(\vec{r}) = V_C(\vec{r}) + V_T(\vec{r})S_{np} + V_{LS}(\vec{r}) \vec{L} \cdot \vec{S} \quad (2.3.8)$$

with central, tensor, and spin-orbit terms. In terms of the S state $[u(r)]$ and D state $[w(r)]$ deuteron radial wavefunctions,

$$D_0(\vec{r}) = V_C u + \sqrt{8} V_T w \quad (2.3.9a)$$

$$D_2(\vec{r}) = (V_C - 2V_T - 3V_{LS})w + \sqrt{8} V_T u \quad (2.3.9b)$$

where the coupled S and D state deuteron radial wavefunctions were calculated numerically (Ros 79).

The dependence of $\tilde{D}_L(q)$ on momentum transfer q may be seen from the Fourier transform of $D_L(r)$,

$$\tilde{D}_L(q) = 4\pi \int_0^\infty r^2 dr j_L(qr) D_L(r) \quad (2.3.10)$$

just as for the Hulthén deuteron wave function but with $L = 0$ or 2 . The results are shown in Fig. 2.3.1 along with those for the Hulthén wavefunction, for comparison. As expected, the $L = 0$ part of $\tilde{D}_L(q)$ is similar to the $\tilde{D}(q)$ obtained using the ($L = 0$) Hulthén wavefunction. Of note is the contribution from the deuteron D state to $\tilde{D}_L(q)$ which rises from zero at

$q = 0$ to exceed $\tilde{D}_0(q)$ for $q \gtrsim 2.0 \text{ fm}^{-1}$. The region of momentum transfer covered in the $^{12}\text{C}(p,d)^{11}\text{C}(\text{g.s.})$ reaction at various bombarding energies is indicated in the figure. At 800 MeV the effects of the deuteron D state are quite important, if not dominant.

At lower bombarding energies, it has been customary to make the zero range (ZR) approximation, to reduce the dimensionality of the plane wave transition amplitude integral from six to three. This approximation consists of replacing $D(r)$ with

$$D(\vec{r}) \approx D_0 \delta(\vec{r}) \quad (2.3.11)$$

where

$$D_0 = \int d\vec{r} \, v_{pn}(\vec{r}) \phi_d(\vec{r}) \quad . \quad (2.3.12)$$

Physically, this approximation means that the incoming proton absorbs a neutron at the same point at which the deuteron is emitted. The Fourier transform of $V_{pn}\phi_d$ then becomes a single constant, $D_0 = -124 \text{ MeV fm}^{3/2}$. Also shown in Fig. 2.3.1 is the zero range approximation for comparison to the other two approximations outlined above. Since these treat the full six dimensional integration in the PW transition amplitude, they are known as exact finite range (EFR) prescriptions. As shown by Fig. 2.3.1, in the region of momentum transfer appropriate to the 800 MeV (p,d) reaction, the zero range approximation is inapplicable.

2.4 Neutron Bound State Potentials

The second Fourier transform in the PW transition amplitude is that of the neutron bound state wavefunction. Since the momentum transfer for the 800 MeV $^{12}\text{C}(p,d)$ reaction ranges from 2.5 to over 5 fm^{-1} , the transition amplitude can be expected to be quite sensitive to the high momentum components of the neutron bound state wavefunction. In the ONM, this large momentum transfer must be supplied by the picked up neutron, in spite of the fact that the Fermi momentum of a bound nucleon is only about 1.4 fm^{-1} . (SF 74)

In the single particle model, the total nuclear wavefunction is an antisymmetrized product of single particle wavefunctions (ES 67)

$$\psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_A) = a \psi_1(\vec{r}_1) \dots \psi_A(\vec{r}_A) \quad (2.4.1)$$

which satisfy the Schroedinger equation

$$(T_i + V_i) \psi_n(\vec{r}_i) = E_n \psi_n(\vec{r}_i) \quad (2.4.2)$$

where n denotes the quantum numbers of nucleon i . The Hamiltonian which generates the wavefunction of the picked up neutron can be written

$$H = T_n + V(\vec{r}_n) \quad (2.4.3)$$

where $V(\vec{r}_n)$ represents the interaction of the neutron with the core. In this picture (ignoring antisymmetrization) the wavefunction is just

$$\psi_A = \psi(\vec{r}_n) \psi(\vec{\xi}_B) \quad . \quad (2.4.4)$$

The bound state potential $V(\vec{r}_n)$ was parameterized by choosing it to be of Woods-Saxon form,

$$V(\vec{r}_n) = V\{1 + \exp[(r-r_0 A^{1/3})/a]\}^{-1} \quad (2.4.5)$$

with a spin orbit potential given by

$$V_{SO}(r) = -0.55 \frac{1}{r} \frac{dV(r)}{dr} \vec{L} \cdot \vec{S} \quad . \quad (2.4.6)$$

The parameters of $V(\vec{r}_n)$ were then constrained in the following way (ES 67). First, the eigenenergy of the Schroedinger equation satisfied with the Hamiltonian of (2.4.3) was taken to be the separation energy as determined experimentally by (p,2p) experiments.

Secondly, the geometry (radius r_0 and diffuseness a) of $V(\vec{r}_n)$ was constrained so as to generate single particle wavefunctions which yielded a nuclear matter density consistent with high momentum transfer electron scattering data. The nuclear matter density

$$\rho(\vec{r}) = \frac{A}{\Sigma} |\psi_i(\vec{r})|^2 \quad (2.4.7)$$

was folded in with the proton charge density distribution to obtain the total nuclear charge density

$$\rho_{ch}(\vec{r}) = \int \sum_{i=1}^Z |\psi(\vec{r}_{iB})|^2 \rho_p(|\vec{r}-\vec{r}_{iB}|) d\vec{r}_{iB} d\hat{r} \quad (2.4.8)$$

The nuclear form factor is related to the charge density by

$$F(q) = \frac{1}{Ze q} \int_0^\infty r \sin(qr) \rho_{ch}(r) dr \quad (2.4.9)$$

This quantity is compared to the experimentally measured charge form factor given by (SM 70)

$$F^2 = \frac{d\sigma/d\Omega}{\sigma_M} \quad (2.4.10)$$

where the Mott cross section is

$$\sigma_M = \left(\frac{Ze^2}{2E} \right)^2 \frac{\cos^2 \theta/2}{\sin^4 \theta/2} \frac{1}{1 + \frac{2E}{Mc^2} \sin^2 \frac{\theta}{2}} \quad (2.4.11)$$

and the momentum transfer is

$$q = \frac{2E \sin \theta/2}{\sqrt{1 + \frac{2E}{Mc^2} \sin^2 \frac{\theta}{2}}} \quad (2.4.12)$$

In the above expressions, θ is the laboratory scattering angle, E the electron bombarding energy, and Z

and m are the target charge and mass, respectively.

It should be noted that this approach yields the single particle potential $V_p(\vec{r}_p)$ appropriate for describing the interaction of a proton with the core. Since the wavefunction of interest in the PW transition amplitude is that of a neutron generated by a single particle potential $V_n(\vec{r}_n)$, we have assumed that the single particle potentials appropriate for neutrons and protons are the same. There is ample evidence that this is a reasonable assumption, for example elastic π^+ and π^- scattering (Joh+ 79), (Mor 76), (Ive+ 78), or intermediate energy proton scattering (Ray+ 78), (RCH 78). In addition, most of the targets studied in this dissertation had isospin $T = 0$, for which $\rho_n \sim \rho_p$ is a very good approximation.

An example of the sensitivity of this procedure is shown in Fig. 2.4.1, where the square of the ^{16}O electron charge form factor (SM 70) measured at 375 MeV is compared to the predictions from (2.4.10) for three values of the radius (r_0) and diffuseness (a) of $V(r_n)$. Criteria in the selection of the proper potential were that the first minimum in the form factor be reproduced, and that the large momentum transfer behavior of the form factor be reproduced within reasonable limits. Although the first criterion was easily achievable (as demonstrated in

Fig. 2.4.1 Shown are 375 MeV $^{16}\text{O}(\text{e},\text{e},)$ data (open circles) (SM 70) presented as the square of the charge form factor vs. q^2 (momentum transfer squared). The three curves represent the predictions with neutron bound state geometries of $r = 1.40$, $a = 0.55$ (solid line), $r = 1.44$, $a = 0.55$ (dashed line), and $r = 1.48$, $a = 0.40$ (dash-dot line).

electron scattering
(charge form factor)² vs q^2

Fig. 2.4.1), the second was never fully realizable, implying that perhaps the Woods-Saxon form chosen for the potential was inadequate at large momentum transfers.

The results of this analysis on the nuclei ^{12}C , ^{16}O , and ^{40}Ca are presented in Table 2.4.1, along with the potentials used for the other relevant nuclei.

2.5 Empirical Optical Potentials

Now that the ingredients contained in the PW transition amplitude have been fully discussed, distortion may be introduced as outlined in Sect. 2.1, and accounted for by means of the optical potentials discussed therein. The distorted wave transition amplitude relevant to the discussions which follow is defined in Eq. (2.1.34), which contains the physical ingredients of the plane wave transition amplitude discussed in the previous two sections, as well as new effects arising from the distortion. It should be reiterated that the optical potentials of (2.1.23) are arbitrary, subject to the following restrictions: they must be functions of the channel radius variable alone, and they should satisfy conditions (2.1.29) as well as possible.

The first of conditions (2.1.29) requires that there be an approximate cancellation between the many body potential V_{PB} , which represents the interaction

Table 2.4.1. Neutron Bound State Potentials

Nucleus	V (MeV)	4 (fm)	a (fm)	λ^*
${}^7\text{Li}$	-39.3	1.82	0.55	0.0
${}^{12}\text{C}$	-50.8	1.44	0.38	25.0
${}^{13}\text{C}$	-33.74	1.44	0.38	25.0
${}^{16}\text{O}$	-47.20	1.44	0.55	25.0
${}^{28}\text{Si}$	-55.99	1.33	0.70	25.0
${}^{40}\text{Ca}$	-51.12	1.33	0.70	25.0

*Thomas spin orbit term

$$V_{\text{SO}} = \frac{-\lambda}{45.2} \frac{1}{r} \frac{dV(r)}{dr} \vec{L} \cdot \vec{S}$$

not used in deuteron optical potential.

of the projectile with all the nucleons comprising the core nucleus, and the potential U_α , which represents an average interaction of the projectile with all the nucleons in the core. This prescription is analagous to the situation of light propagating in a medium, where the microscopic interaction with the constituents of the medium is approximated by a (complex) index of refraction (Fes 58).

The form of the optical potential is taken to be

$$\begin{aligned}
 U(r) = & V_C(r) + V_R f(x_R) + iV_I f(x_I) + V_{RS} \frac{df(x_{RS})}{dx_{RS}} \\
 & + iV_{IS} \frac{df(x_{IS})}{dx_{IS}} - \left[V_{RSO} \frac{1}{r} \frac{df(x_{RSO})}{dx_{RSO}} \right. \\
 & \left. + iV_{ISO} \frac{1}{r} \frac{df(x_{ISO})}{dx_{ISO}} \right] \vec{L} \cdot \vec{S}
 \end{aligned} \tag{2.5.1}$$

where the Woods-Saxon shape function is

$$f(x_i) = (1 + \exp(x_i))^{-1} \tag{2.5.2}$$

and

$$x_i = \frac{r - r_i A^{1/3}}{a_i} \tag{2.5.3}$$

The Coulomb potential V_C is that of a uniformly charged sphere of radius $R_C = r_C A^{1/3}$,

$$\begin{aligned}
V_C(r) &= \frac{zZe^2}{2R_C} \left(3 - \frac{r^2}{R_C^2} \right) & r < R_C \\
&= \frac{zZe^2}{r} & r \geq R_C \quad . \quad (2.5.4)
\end{aligned}$$

The atomic numbers of the two constituents of the partition being considered are denoted by z and Z . The total number of parameters (strengths V_i , radii r_i , and diffusenesses a_i) in the full optical potential number 19, although since the spin-orbit potential was found to be unimportant at 800 MeV, this term was neglected. Additionally, the optical potential was usually chosen to be of volume form, and the geometry of the real and imaginary volume terms was taken to be the same. Thus there were really only 5 potential parameters: R_C , V_R , V_I , r_i , and a_i , separately for the proton and deuteron channels.

One of the great advantages of the optical model approach is that the optical potentials may often be determined "experimentally" via elastic scattering. The strengths, radii and diffusenesses of (2.5.1) are treated as parameters to be determined phenomenologically by fitting measured elastic scattering data. The five parameters mentioned above must reproduce the overall magnitude, slope, and minima exhibited in the experimental angular distributions.

An optical model search code (Ros 78) was used to generate the $p + A$ optical potentials at 800 MeV. This procedure was possible due to the availability of 800 MeV elastic scattering data for targets of ^{12}C , ^{13}C , and ^{58}Ni (Bla 78), (Bla+ 77). Fits to the ^{12}C (She 79) and ^{58}Ni (Ray 79) elastic scattering data were generated by minimizing the quantity

$$\chi^2 = \frac{1}{N} \sum_{i=1}^N \left(\frac{x_i^{\text{th}} - x_i^{\text{expt}}}{\Delta x_i^{\text{expt}}} \right)^2 \quad (2.5.5)$$

where x_i represents an elastic differential cross section at a particular scattering angle either measured experimentally or predicted from the optical potential of (2.5.1).

The proton optical potentials appropriate for ^{12}C and ^{58}Ni determined using the empirical approach of this section are tabulated in Table 2.6.1, along with the deuteron optical potentials discussed in the following two sections.

2.6 KMT Optical Potentials

Due to the dearth of deuteron elastic scattering data near 800 MeV, the deuteron optical potentials were constructed from the measured nucleon-nucleon phase shifts using the formalism developed by Kerman, McManus, and Thaler (KMT 59) and extended by Feshbach,

Gal, and Hüfner (FGH 70). In this approach (She 78) the scattering potential $V(\vec{q})$ is related to the nucleon-nucleon scattering amplitude $M(\vec{q})$, and the momentum-space nuclear density $F(\vec{q})$ by

$$V(\vec{q}) = \frac{-\hbar^2 c^2}{(2\pi)^2} \eta (A - 1) \frac{p_L}{E_L p_0} M(\vec{q}) F(\vec{q}) \quad . \quad (2.6.1)$$

In this expression, A denotes the atomic weight of the target, E_L is the total lab energy, p_L is the projectile lab momentum, p_0 is the momentum in the center of mass frame of the two nucleon system, and m is the nucleon mass. The quantity

$$\eta = \frac{1 + \frac{2A}{A^2 + 1} \frac{E_L}{m}}{1 + \frac{A}{A^2 + 1} \frac{E_L}{m} + \frac{m}{E_L}} \quad (2.6.2)$$

arises from relativistic corrections. The configuration space scattering potential $V(\vec{r})$ is obtained by taking the Fourier transform of Eq. (2.6.1). The observation customarily made at this point (FGH 70) is that $M(\vec{q})$ is a much more slowly varying function of q than $F(\vec{q})$; consequently $M(\vec{q})$ may be taken to be constant, equal to its value at $q = 0$, leaving

$$V(\vec{r}) = -(\hbar c)^2 \eta (A-1) \frac{p_L}{E_L p_0} M(q=0) \frac{2\pi \rho(\vec{r})}{A} \quad . \quad (2.6.3)$$

Employing the optical theorem (Mer 61),

$$\text{Im } M(q=0) = \frac{P_0 \sigma_T}{4\pi} \quad (2.6.4)$$

the imaginary part of the optical potential becomes

$$\text{Im } V(\vec{r}) = \frac{-(\hbar c)^2}{2E_L} \eta P_L \frac{A-1}{A} \sigma_T \rho(\vec{r}) \quad (2.6.5)$$

Note that this is of the volume absorptive form.

The nuclear density (to be used in constructing a deuteron nucleus optical potential) is obtained by folding the deuteron density with the target charge density,

$$\rho(\vec{r}) = \int \rho_d(\vec{r}') \rho_{ch}(\vec{r} - \vec{r}') d\vec{r}' \quad (2.6.6)$$

In this expression for $\rho(\vec{r})$ the charge density approximates a folding of the target point density and the nucleon matter density, so that the finite size of the nucleons is roughly accounted for in $\rho_{ch}(\vec{r}-\vec{r}')$. The radius and diffuseness of the target charge densities were taken from electron scattering tables (DDD 74). Implicit in this approach is the assumption that the neutron and proton distributions are the same. This point was discussed in Sect. 2.4.

The deuteron density was chosen to be of the Hulthén form, which has been discussed in detail in

Sect. 2.3. The folding integral (2.6.6) was evaluated numerically to give the configuration space density appropriate for the $d + A$ optical potential. In the present work the optical model geometry was parameterized by a Woods-Saxon potential of the form (2.5.2). In this manner, radii and diffusenesses for the various targets were obtained for the optical potential U_β . The geometry of the real part of U_β was assumed to be the same as for the imaginary part. The final values of the radius and diffuseness were found to depend only slightly on the number of points included in the numerical integration, although these differences are negligible when the densities are plotted vs. the nuclear radius. Densities were chosen from fits including approximately 50 points from the numerical integration of (2.6.6). These data sets emphasized the nuclear surface, where the density changes most rapidly, and therefore provided a more rigorous constraint on the parameters r and a .

The depth of $\text{Im } V(r)$ may easily be determined from (2.6.5) and the depth of the Woods-Saxon well resulting from the least squares fit described above, subject to the normalization N where

$$A = 4\pi N \int \rho(r) r^2 dr \quad . \quad (2.6.7)$$

The kinematic quantities, and total cross sections

from which the imaginary depths were computed (Bar+ 78) are tabulated in Table 2.6.1, for three proton kinetic energies. The total cross sections tabulated, appropriate for isospin zero nuclei, are

$$\sigma_T = \frac{1}{2}(\sigma_T^{pp} + \sigma_T^{pn}) \quad . \quad (2.6.8)$$

The depth of the real part of $V(r)$ may be found from $\text{Im } V(r)$ and the ratio of the real to imaginary parts of the nuclear scattering amplitude. Since only the forward ($\theta = 0^\circ$) scattering amplitude is relevant to this discussion, the spin flip and double spin flip terms are zero. Then the nucleon-nucleon scattering amplitude (Coulomb contributions excluded) may be written as

$$f = f_s + f_t = f_s + f_{00} + f_{11} + f_{-1-1} \quad (2.6.9)$$

where s denotes singlet, t denotes triplet, and the ij in f_{ij} denote the projections of the total spin angular momentum on the z -axis of the two nucleons. Examining the p - p amplitudes, and including the nuclear phase shifts explicitly (SYM 57),

$$f_s = \frac{2}{ik} \sum_{\substack{\ell \\ \text{even}}} P_\ell(\theta = 0) \left(\frac{2\ell + 1}{2} \right) \alpha_\ell \quad (2.6.10a)$$

and

Table 2.6.1. KMT Optical Potential Input Quantities (appropriate for protons of the specified kinetic energy)

T_p (MeV)	$\frac{P_L}{\hbar}(\text{fm}^{-1})$	E_L (MeV)	$\frac{P_0}{\hbar}(\text{fm}^{-1})$	E_0 (MeV)	$\eta(^{12}\text{C})$	$\sigma_T(\text{fm}^2)$	$\frac{\text{Re } f}{\text{Im } f}$
200	3.26	1135	1.550	983.7	1.028	3.20	+ .895
400	4.829	1335	2.192	1030.	1.051	2.95	+ .239
800	7.407	1735	3.099	1117.	1.091	3.90	-0.075*

*750 MeV phase shifts used.

$$f_t = \frac{2}{ik} \sum_{\substack{\ell \\ \text{odd}}} P_\ell(\theta=0) \left\{ \left(\frac{2\ell+3}{2} \right) \alpha_{\ell, \ell+1} + \left(\frac{2\ell+1}{2} \right) \alpha_{\ell, \ell} \right. \\ \left. + \left(\frac{2\ell-1}{2} \right) \alpha_{\ell, \ell-1} \right\} \quad (2.6.10b)$$

where

$$\alpha_\ell = e^{2i\delta_\ell} - 1, \quad (2.6.11a)$$

$$\alpha_{\ell\ell} = e^{2i\delta_{\ell\ell}} - 1, \quad (2.6.11b)$$

and

$$\alpha_{j\pm 1, j} = \cos 2E_j e^{2i\delta_{j\pm 1, j}} - 1. \quad (2.6.11c)$$

The δ_{ej} are the nuclear (bar) phase shifts (Coulomb contributions excluded) and the E_j are mixing parameters. The isospin averaged scattering amplitude is

$$\bar{F} = \frac{1}{4}(3\bar{F}_{T=1} + \bar{F}_{T=0}) \quad (2.6.12)$$

where

$$\bar{F}_{T=1} = \frac{1}{4}(3f_t^{PP} + f_s^{PP}) \quad (2.6.13a)$$

and

$$\bar{F}_{T=0} = \frac{1}{4}(3f_t^{np} + f_s^{np}). \quad (2.6.13b)$$

The ratio of $\text{Re } \bar{F}$ to $\text{Im } \bar{F}$ was computed using the phase shifts of (MAW 68a), (MAW 68b), (AHR 74) and (AHR 77). These ratios are tabulated in Table 2.6.1 for a number of different nucleon kinetic energies. Table 2.6.2

presents the results of the MNT analysis: the 800 MeV deuteron optical potentials appropriate for each of the nuclei studied.

Implicit in the construction of the deuteron optical potential from the nuclear optical potentials are two assumptions. The first is that the neutron and proton nuclear densities (and consequently the corresponding optical potentials) are equivalent. This point was addressed in Sect. 2.4 and was shown to be a reasonable assumption in the light of recent investigations. The second assumption inherent in the KMT approach is that the optical potential appropriate for the description of deuteron elastic scattering is constructed as the sum of the neutron and proton optical potentials at half the deuteron kinetic energy.

An alternative approach which yields the same result has been outlined by Johnson and Soper (JS 69). These workers have attempted to include deuteron breakup in their adiabatic theory, with the result being the replacement of their deuteron optical potential with the sum of proton-nucleus and neutron-nucleus optical potentials valid for half the kinetic energy of the deuteron. Although the adiabatic approximation (impulse approximation) becomes more reliable as the deuteron kinetic energy rises, the additional assumption of deuteron breakup into a relative S-state

Table 2.6.2. 800 MeV KMT Optical Potentials

Potential	V	r_R	a_R	W	r_I	a_I
p + ^{12}C phenomenological	+13.73	0.872	0.304	-68.93	0.948	0.472
p + ^{58}Ni phenomenological	+2.97	1.100	0.537	-108.62	0.933	0.634
d + ^6Li	-9.70	0.946	0.734	-40.52	0.946	0.734
d + ^{11}C	-14.02	0.896	0.725	-58.64	0.896	0.725
d + ^{12}C	-15.70	0.856	0.727	-65.70	0.856	0.727
d + ^{15}O	-16.02	0.910	0.767	-67.02	0.910	0.767
d + ^{27}Si	-15.44	0.945	0.795	-64.62	0.945	0.795
d + ^{39}Ca	-16.20	0.970	0.801	-67.76	0.970	0.801

(only) becomes untenable (RS 75).

The importance of the deuteron D state (see Sect. 2.3) for the high energy (p,d) reaction (RS 75) suggests that deuteron breakup into channels with non-zero relative angular momentum may become important in the large momentum transfer regime. If this is indeed the case, the choice of U_β becomes ambiguous, since it is not known at which (different) kinetic energies the (energy dependent) neutron and proton potentials V_{pB} and V_{nB} should be evaluated. The following section presents an attempt to deal with this problem.

2.7 Well-Matched Optical Potentials

The empirical and KMT optical potentials described in the previous sections do not necessarily satisfy the conditions (2.1.29)

$$||V_{pB} - U_\alpha|| \approx 0 \quad (2.1.29a)$$

and

$$||V_{pB} + V_{nB} - U_\beta|| \approx 0 \quad (2.1.29b)$$

which are referred to as well-matching conditions. In the analysis of some large momentum mismatch reactions at low energies, such as 35 MeV (p, α) (Smi 76), or 18 MeV ($^3\text{He},\alpha$) (Sto+ 67), it has been found that care in satisfying the above conditions is essential in obtaining good fits and consistent results in the

analysis of these reactions. Since the 800 MeV (p,d) reaction involves a large momentum mismatch ($2.0 - 5.5 \text{ fm}^{-1}$) the use of "well-matched" optical potentials may be important, as was found to be at low energies.

The first of approximations (2.1.29) simply neglects the intermediate states in the $p + A$ system, i.e. elastic scattering is assumed to be the dominant partition. However, inelastic excitation of rotational and vibrational states in the $^{12}\text{C}(p,p')$ reaction (Bla 78) can compete with, and even exceed the elastic scattering at some angles. When this situation arises, two-step, coupled-channels calculations (discussed in Sect. 2.8) must be employed as opposed to the one-step calculations considered so far.

The second of approximations (2.1.29) is ambiguous in the sense that the energy at which the potentials V_{pB} and V_{nB} are evaluated is ill-defined. One approach is to form the deuteron optical potential U_β from the sum of proton (V_{pB}) and neutron (V_{nB}) optical potentials evaluated at one half the deuteron kinetic energy. This was the approach outlined in the previous (KMT) section.

An alternative approach is to form the 800 MeV deuteron optical potential U_β from the sum of the 800 MeV proton optical potential (for V_{pB}) and the neutron bound state potential (for V_{nB}). This is the approach

which has proven useful in choosing among families of optical potentials in the above mentioned analysis of large momentum mismatch reactions at lower bombarding energies. Although these deuteron optical potentials are inappropriate for the description of 800 MeV deuteron elastic scattering, they provide the maximum cancellation in (2.1.29b) and may be more appropriate for DWBA calculations of the 800 MeV (p,d) reaction. Table 2.7.1 lists the 800 MeV well-matched deuteron optical potentials used for the calculations presented in Chapter VI.

2.8 Core Coupling

The approximation (2.1.29a) that the proton-core potential V_{pB} is adequately represented by the optical potential U_α , which describes the elastic scattering of protons on the target nucleus, may break down if other reaction channels can compete with the elastic scattering channel. The most obvious candidate for this competition is the lowest inelastic channel, where the target or residual nucleus is excited vibrationally or rotationally before or after the neutron pickup. Indeed, recent large momentum transfer 800 MeV (p,p') data on ^{12}C (Bla 78) have demonstrated that, as the scattering angle (momentum transfer) increases beyond about 38° ($q = 4.8 \text{ fm}^{-1}$), the only states appreciably excited are the 4.44 MeV 2^+

Table 2.7.1. 800 MeV Well-Matched Optical Potentials

Potential	V	r_R	a_R	w	r_I	a_I
p+ nucleus phenomenological	+13.73	0.872	0.304	-68.93	0.948	0.472
d + ^6Li	-39.3 +13.73	1.82 0.872	0.55 0.304	-68.93	0.948	0.472
d + ^{11}C	-50.8 +13.73	1.44 0.872	0.38 0.304	-68.93	0.948	0.472
d + ^{12}C	-33.74 +13.73	1.44 0.872	0.38 0.304	-68.93	0.948	0.472
d + ^{15}O	-47.20 +13.73	1.44 0.872	0.55 0.304	-68.93	0.948	0.472
d + ^{27}Si	-57.8 +13.73	1.27 0.872	0.60 0.304	-68.93	0.948	0.472
d + ^{39}Ca	-53.27 +13.73	1.27 0.872	0.60 0.304	-68.93	0.948	0.472

and 14.1 MeV 4^+ rotational levels, and the 9.6 MeV 3^- vibrational level. In light of this fact, calculations were performed which included the strongest inelastic excitations of the target and residual nucleus, before and after the neutron pickup. The formalism on which these calculations are based is a coupled channels (CC) one, as embodied in the computer code CHUCK (Kun 79c). Unfortunately, however, no finite range coupled channels code was available, so that the two-step (TS) calculations were all performed in zero range (see Sect. 2.2).

To demonstrate the effects of two-step, coupled channels pathways, a calculation for the excitation of the 4.44 MeV 2^+ level in ^{12}C from the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction will be presented. The pathways accounted for in these calculations are shown in Fig. 2.8.1, from which it is apparent that this procedure demands that the spectroscopic factors for all pickup steps be known, as well as the inelastic strength or deformation parameters in both nuclei. The deformation parameters were taken from Blanpied (Bla 78). The spectroscopic amplitudes were calculated as described in Sect. 2.1, and presented in Table 2.1.1.

Fig. 2.8.2 presents the results of a two-step, zero range coupled channels, a one-step, zero range, and a one-step exact finite range calculation, so that

TWO STEP PATHWAYS

Fig. 2.8.1 Two step pathways for $p_{3/2}$ (solid lines) and $p_{1/2}$ (dashed lines) pickup to the 4.44 MeV 2^+ state of ^{12}C from the $^{13}\text{C}(p,d)^{12}\text{C}$ reaction. The deformation parameters β_2 are also shown. Spectroscopic factors for the pickup legs are tabulated in Table 2.1.1.

Fig. 2.8.2 Calculations of the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ (4.44 MeV 2^+) reaction. The solid line is the EFR one-step result, normalized with a spectroscopic factor $C^2S = 1.038$. The ZR coupled channels prediction is shown as the dash-dot line, and the ZR one-step curve (dashed) is shown for comparison. Both ZR curves employ a zero range strength $D_0 = -80 \text{ MeV fm}^{3/2}$.

the effects of each are more easily seen. The effect of the two-step coupled channels calculation is to wash out the structure visible in the zero range calculation, and raise the overall normalization slightly, especially at forward angles. In Fig. 2.8.2, a D_0 strength (see Sect. 2.3) has been employed in the zero range one-, and two-step calculations which matches the 0° predictions between the $^{13}\text{C}(p,d)^{12}\text{C}(\text{g.s.})$ zero range one-step, and the exact finite range one-step calculations. This empirical value, $D_0 = 80 \text{ MeV} \cdot \text{fm}^{3/2}$, is significantly lower than the calculated value of $124 \text{ MeV} \cdot \text{fm}^{3/2}$, appropriate for lower energies (And+ 78).

2.9 Numerical Tests and Sensitivities

For all the 800 MeV (p,d) calculations, the number of partial waves in the proton and deuteron channels was chosen to be 50. This number is known to be adequate because calculations performed with 50, 60 and 70 partial waves gave essentially the same differential cross sections for $^{40}\text{Ca}(p,d)^{39}\text{Ca}(\text{g.s.})$, the worst case. Furthermore, the radial integrals have dropped typically by a factor of 1000 from their maximum by the fiftieth partial wave.

The integration step size and radial cutoffs for the numerical integrations are constrained by the computer code's requirement that the maximum number of

steps be less than 400. Since the form factor and radial wave functions had wavelengths of roughly one to two fermis, the step size varied from 0.025 fm for ^{12}C to 0.03 fm for ^{40}Ca , with the maximum radial integration cutoff varying from 10 fm to 12 fm for the same two nuclei, respectively. Calculations were performed for 800 MeV $^{13}\text{C}(\text{p},\text{d})$ in zero range with the following pairs of values for the step size and maximum cutoff: 0.04 and 16.0 fm, 0.04 and 12.0 fm, and 0.03 and 12.0 fm. The results of the three calculations were indistinguishable from one another, indicating the values used in this work were reasonable.

An important feature of the calculations is their sensitivity to the nuclear interior. This was tested with ZR calculations in which the lower cutoff for the radial integrations was moved from its normal location at the origin, out to the surface in 1 fm steps. Fig. 2.9.1 shows the results of these calculations for 800 MeV ^{40}Ca , $^{13}\text{C}(\text{p},\text{d})$ to the ground states of the residual nuclei. Apparent from these figures is the importance of the nuclear interior, the main effects being to lower the cross sections and increase the slope of the angular distributions as more and more of the nuclear interior is sampled. For reference purposes, the nuclear charge densities of ^{40}Ca and ^{13}C are shown in Fig. 2.9.2 to indicate where in

2.9.1 Sensitivity of the DWBA calculations
to the nuclear interior, demonstrated
for 800 MeV $^{13}\text{C}(p,d)$ and $^{40}\text{Ca}(p,d)$

Fig. 2.9.2 Nuclear charge densities of ^{13}C (solid line) and ^{40}Ca (dashed line) vs. the nuclear radius, for purposes of comparison to Fig. 2.9.1.

the nuclear volume each cutoff was made.

The effects arising from changes in the optical model parameters in the proton and deuteron channels were also investigated. While the differential cross sections are sensitive to almost every parameter, the largest effects were observed while varying the radius or diffuseness of the deuteron imaginary optical potential. The results of these changes are summarized in Table 2.8.1, for the ZR DWBA calculations of 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}(\text{g.s.})$. Since these parameter variations result in a lowering of the calculated differential cross sections by factors of 3 to 7, care was taken to choose consistent sets of optical potential parameters for all the targets.

The proton optical potentials generated from fits to measured elastic scattering on targets of ^{12}C and ^{58}Ni are tabulated in Table 2.6.2. Calculations performed in zero range for 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}(\text{g.s.})$ were done utilizing each of these proton potentials, and the KMT deuteron optical potential of Table 2.6.2. The results, presented in Fig. 2.9.3, indicate that no appreciable difference arises from using one potential over the other. Consequently, for the calculations presented in Chapter VI, the same (^{12}C) proton optical potential parameters were used for each target.

Table 2.8.1. Optical Potential Parameter
Sensitivities (Deuteron Imaginary
Radius and Diffuseness) 800 MeV
 $^{40}\text{Ca}(p,d)^{39}\text{Ca}(\text{g.s.})$ ZR DWBA

	$V = 13.73$	$w = -68.93$	
$U_p =$	$r = 0.872$	$r_i = 0.948$	
	$a = 0.304$	$a_i = 0.472$	
	$V = -8.0$	$w = -115.0$	
$U_d =$	$r = 0.96$	$r_i = \text{variable}$	
	$a = 0.41$	$a_i = \text{variable}$	
$\frac{d\sigma^*}{d\Omega}(6^\circ)_{\text{ZR}}$	$\frac{\frac{d\sigma}{d\Omega}(6^\circ)_{\text{ZR}}}{\frac{d\sigma}{d\Omega}(6^\circ)_{\text{expt}}}$	deuteron imaginary radius	deuteron imaginary diffuseness
2.5 b	30	1.0 fm	0.41 fm
1.8	21	1.2	0.41
0.70	9	1.4	0.41
2.5	30	1.0	0.41
0.85	10	1.0	0.70
0.34	4	1.0	1.00

*Note: $D_0 = 60 \text{ MeV fm}^{3/2}$

Fig. 2.9.3 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}(\text{g.s.})$ ZR DWBA predictions using best fit (see Sect. 2.5) proton optical potentials derived from 800 MeV proton elastic scattering on targets of ^{12}C (dashed line) and ^{58}Ni (solid line).

Fig. 2.9.4 Sensitivity of 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}(\text{g.s.})$
DWBA calculations to the choice of parameters for the neutron bound state:
 $r = 1.40$, $a = 0.45$ (dashed line) and
 $r = 1.44$, $a = 0.55$ (solid line).

$^{16}\text{O}(p,d)^{15}\text{O} \text{ (} p_{1/2} \text{ gs)}$

The sensitivity of the calculations to the neutron bound state potential, discussed in Sect. 2.4, was also investigated. Shown in Fig. 2.9.4 are EFR DWBA calculations for 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}(\text{g.s.})$ calculations, employing two different neutron bound state potentials. Slight differences in the overall magnitude, as well as differences in the location of the various (shallow) minima, are observed to exist between the two calculations. This difference suggests that, once the DWBA treatment is well understood for the (p,d) reaction in the 800 MeV regime, comparison with the experimental data may provide a means of determining the geometry of the neutron bound state potentials.

2.10 Angular Momentum Selectivity

Preferential population of final states with large angular momentum quantum numbers is a feature to be expected in large momentum transfer reactions (Aus 70), (BW 62). Maximal overlap of the radial integrals, and correspondingly greater cross sections, occurs for partial waves L_p and L_d when $L_p \approx k_p R$ and $L_d \approx k_d R$ (BW 62), where $k_p(k_d)$ is the wave number of the proton (deuteron), and R is the nuclear radius. For large momentum mismatch reactions such as 800 MeV (p,d), one has $k_d > k_p$. Consequently, the angular momentum transferred to the core $\vec{\ell} = \vec{L}_p - \vec{L}_d$, is also

large since $\ell = |\vec{k}_p - \vec{k}_d|R$. Transitions with $\ell \sim |\vec{k}_p - \vec{k}_d|R$ will be enhanced relative to transitions with $\ell < |\vec{k}_p - \vec{k}_d|R$ (BW 62), (Peh+ 65). This observation is indeed born out by the data presented in Chapter VI.

CHAPTER III

THEORY OF THE (π^+, p) REACTION

Another reaction with which to study large momentum transfer neutron pickup is the (π^+, p) reaction. The One Nucleon Model (ONM) interpretation of the reaction mechanism involves a positive pion (π^+) incident upon a nucleus consisting of A nucleons which comprise a core (B) and the neutron (n) to be transferred. The incident positive pion is absorbed by the neutron, the resulting reaction products being the proton (p) and the residual nucleus (A - 1 or B).

Large momentum transfers $|\vec{q}| = |\vec{k}_p - \vec{k}_\pi|$ are possible with this reaction even for pions of relatively little ($\lesssim 100$ MeV) kinetic energy because upon absorption, the rest mass of the pion ($m_{\pi^\pm} = 139.567$ MeV) (Bar+ 78) is converted into kinetic energy of the reaction products. Even for very slow pions ($T_\pi \approx 0$), momentum conservation requires that the transferred neutron have a momentum roughly equal to twice the typical nuclear Fermi momentum of ~ 250 MeV/c.

Another feature of the (π^+, p) reaction is the accessibility of $\Delta T = 3/2$ states. The pion exists as an isospin triplet (isospin $T = 1$) which can couple with the neutron isospin of $1/2$ to yield

isospin transfers of 3/2 or 1/2.

Various models for the (π^+, p) reaction mechanism have been proposed in the past. Fig. 3.1 illustrates the most common of these models. The ONM for the (π^+, p) reaction is analogous to the ONM for the (p, d) reaction discussed in the previous chapter. This single neutron pickup model requires the momentum mismatch be supplied by a single neutron. Figs. 3.1b and 3.1c depict the Two Nucleon Model (TNM) in its most common forms. These triangle diagrams allow the momentum mismatch to be supplied by two nucleons, or by a nucleon and a pion, thus decreasing the momentum to be supplied by any one participant.

Much discussion has been centered around the relative importance of these reaction mechanisms, especially in the case of the time-reversed reaction (p, π^+) , for which more data exist (Hoi 79), (Dah+ 74). The cross sections of the two reactions may be related using detailed balance, the result being (Per 72)

$$\sigma_{\pi A \rightarrow p B} = 2 \frac{2J_B + 1}{2J_A + 1} \frac{p_B^2}{p_{\pi A}^2} \sigma_{p B \rightarrow \pi A} , \quad (3.1)$$

where p_i denotes the center of mass relative momentum of particles i , J the nuclear angular momentum quantum numbers, and the cross sections are those at the same center of mass energy.

Fig. 3.1 Schematic (Feynmann-like) diagrams of the one nucleon model (ONM), the two nucleon model (TNM) and the one pion exchange model (OPEM).

In the analysis of (p, π) reactions, evidence gathered in favor of the ONM consists of the following. In the ONM, the (p, π^-) reaction can only take place when initial or final state interactions (charge exchange) are included. In the TNM, no such charge exchange interactions are necessary to produce negative pions, although π^+ production is expected to be more copious than π^- production. Experimentally, π^- production has been observed to be roughly twenty times less (for pion production on ^{13}C) (Dah+ 73) than π^+ production (Hoi 79). This fact, together with an observed nearly isotropic angular distribution for π^- production suggestive of multistep processes, has been used as evidence in favor of the ONM. Additionally, the (p, π^+) cross sections have been observed to scale (although there are exceptions) with the initial and final nuclear spins roughly in accordance with the factor expected from stripping: $S(2J_f + 1) / (2J_i + 1)$ (Nob 76) (Hoi 79).

Another way to examine the (π^+, p) reaction mechanism is by comparison of (π^+, p) data to low energy pickup and stripping experiments leading to the same residual nucleus. Ignoring the effects generated by two-step pathways, if the TNM (or ONM) is important, the relative intensities of final states populated in the (π^+, p) reaction should be similar to

those in the low energy stripping (or pickup) process. Inclusion of two step contributions can make states which are weakly populated in low energy pickup strongly populated in the (π^+, p) reaction. However, states which are strongly excited (relative to neighboring levels of comparable spin) in low energy stripping, and are only weakly excited in the (π^+, p) reaction, suggest that the TNM is not the dominant (π^+, p) reaction mechanism. This point is discussed more thoroughly in Sect. 6.2.

The (π^+, p) data presented in Chapter V will be examined with the above arguments in mind, to elucidate the dominant reaction mechanism in the (π^+, p) reaction. Meanwhile for the sake of consistency with the (p, d) theory presented earlier, the ONM will now be presented in detail.

3.1 One Nucleon Mechanism (π^+, p) Reaction Theory

The approach customarily used (LE 66) to obtain cross sections for the $A(\pi^+, p)B$ reaction consists of utilizing first-order, time-dependent perturbation theory. In this approach (the ONM), the problem is formulated in terms of a Hamiltonian H describing the absorption of a pion on a single nucleon. This Hamiltonian is obtained from a non-relativistic (Foldy-Wouthuysen) reduction of the pseudoscalar-pseudoscalar pion nucleon interaction (JE 70) and is

written (RK 73)

$$H = \frac{f}{\mu} \sum_{i=1}^A \left[\left(1 - \frac{\mu}{2M}\right) \vec{\sigma}_i \cdot \vec{\nabla}_\pi (\vec{\tau}_i \cdot \vec{\psi}_\pi) - \frac{\mu}{M} \vec{\tau}_i \cdot \vec{\psi}_\pi \vec{\sigma}_i \cdot \vec{\nabla} \right] \quad (3.1.1)$$

in the Galilean-invariant form (JE 70). An alternative expression casts the Hamiltonian in the "static" form (MP 74)

$$H = \sqrt{4\pi} \frac{f}{\mu} \vec{\sigma} \cdot \vec{\nabla}_\pi \vec{\tau} \cdot \vec{\psi}_\pi . \quad (3.1.2)$$

In these expressions, f denotes the pion-nucleon coupling constant ($f^2/4\pi = 0.083$), μ and M are the pion and nucleon masses, $\vec{\sigma}$ and $\vec{\tau}$ denote the nucleon spin and isospin Pauli matrices, the gradient $\vec{\nabla}$ operates only on the nuclear wavefunction, and the gradient $\vec{\nabla}_\pi$ acts only on the pion wavefunction $\vec{\psi}_\pi$. These equations assume that $\hbar = c = 1$. The cross section for the (π^+, p) reaction is then given by (LE 66)

$$\sigma(\theta, \phi) = \frac{M}{(2\pi)^2} E_\pi \frac{k}{q} \sum_f |\langle f | H | i \rangle|^2 , \quad (3.1.3)$$

where q and k are the asymptotic pion and nucleon momenta and $E_\pi = \sqrt{\mu^2 + q^2}$.

The nuclear overlap implicit in (3.1.3) is handled in exactly the same way as in the (p, d) case, outlined in Chapter II. The target internal

wavefunction is expanded in terms of the core wavefunction and the wavefunction of the transferred neutron, the expansion coefficients being the same spectroscopic amplitudes $S^{\frac{1}{2}}$ as used previously. This means that the same (well-known) nuclear structure information (embodied in the spectroscopic amplitudes) is put into the (π^+, p) calculations as is put into the (p, d) calculations.

The matrix element of the pion field operator between an initial state $|\vec{q}\rangle$ where one positive pion (of momentum $\vec{q}$) is present, and a final state consisting of no pions $\langle 0|$ is just (JE 70)

$$\langle 0 | \vec{\tau} \cdot \vec{\psi}_\pi | \vec{q} \rangle = \frac{-1}{\sqrt{2E_\pi}} \psi_\pi(\vec{q}, \vec{r}) \tau_+ , \quad (3.1.4)$$

where the projection operator τ_+ yields non-zero nuclear matrix elements only between initial neutron (nuclear) states and final proton states. In the plane wave approximation, ψ_π is just an incident plane wave $\exp(i\vec{q} \cdot \vec{r})$. However, since the effects of distortion are known to be important (Hoi 79), much attention has been paid to calculating the distorted wave form of ψ_π . Most workers (JE 70), (RK 73) have chosen to obtain the distorted pion field by using the Kisslinger model with the Lorentz-Lorenz (or Kroll) modification included. The Klein-Gordon equation for pion-nucleus scattering which results is (JE 70)

$$(-\nabla^2 + \mu^2) \psi_\pi = \{ (E_\pi - V_C)^2 + Ab_0 q^2 \rho(\vec{r}) - \vec{\nabla} \frac{Ab_1 \rho(\vec{r})}{1 + \frac{1}{3} \xi Ab_1 \rho(r)} \vec{\nabla} \} \psi_\pi \quad (3.1.5)$$

In this equation, V_C denotes the Coulomb potential presented by the nucleus to the incoming pion, and $\rho(\vec{r})$ is the nuclear density (normalized to unity). The complex constants b_0 and b_1 can be obtained directly from the experimentally measured pion-nucleon scattering amplitudes (JE 70), (Wat 58), (AFS 67), (RWF 65). With $\xi = 0$, equation (3.1.5) takes the form of the ordinary Kisslinger equation. The parameter ξ introduces a non-linear dependence of the potential on the scattering centers, which has an effect of softening the velocity-dependent part of the pion-nucleus potential. These points and others related to the determination of proper distorted pion waves have been discussed extensively by others (Ros 78), (MP 74) and will not be discussed further here.

Proper proton distorted waves for the exit channel may be obtained phenomenologically, or constructed from the KMT prescription of Sect. 2.6.

3.2 General Features of the Theory

The calculations of the (π^+, p) reaction were performed with the computer code CHOPER (Ros 79), which is a modified version of the zero range, coupled channels code CHUCK (Kun 79c) used for calculations

involving nuclear projectiles. The (π^+, p) calculations involve the same nuclear model as do the (p, d) calculations discussed in the previous chapter, in that the binding energy, bound state potential well, and spectroscopic amplitudes of the picked up neutron are the same in each case. This is easily seen from Figs. 3.1a and 2.1, which show the ONM reaction diagrams for the (π^+, p) and (p, d) reactions, respectively. The nuclear structure effects of the reaction are depicted in the lower half of these diagrams, and a comparison of these two figures reveals that the lower half of these diagrams is the same for (p, d) as for (π^+, p) . Consequently, in the ONM the same nuclear structure is probed by both reactions. Only the upper half of each figure differs, independent of the nuclear structure involved.

The pion optical potentials used for the (π^+, p) calculations were generated from the energy dependent parameters of Rosenthal (Ros 78). These parameters are tabulated in Table 3.2.1 for the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction, for pion bombarding energies of 90 MeV and 180 MeV. The pion optical potential which employs these parameters is (Dig+ 77)

$$u_{\text{opt}}(r) = u_K(r) + u_A(r) + V_C(r) \quad , \quad (3.2.1)$$

where the Kisslinger potential is

Table 3.2.1. Pion Optical Potentials for the
 $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ Reaction (see Eq.
(3.2.1))

	$p + ^{12}\text{C}$ $T_\pi = 90 \text{ MeV}$	$p + ^{12}\text{C}$ $T_\pi = 180 \text{ MeV}$
$b_0 (\text{fm}^3)$	$-0.7482 + 0.4077i$	$-0.5339 + 0.2772i$
$b_1 (\text{fm}^3)$	$7.1540 + 2.5495i$	$0.6367 + 7.3579i$

$$u_K(r) = -b_0 k^2 \rho(r) + \nabla \cdot b_1 \rho(r) \nabla , \quad (3.2.2)$$

the S-wave absorption term is

$$u_A(r) = -4\pi B_0 \rho^2(r) , \quad (3.2.3)$$

and the Coulomb potential $V_C(r)$ is as described in Sect. 2.5. The nuclear density $\rho(r)$ was taken from nuclear charge density tables (DDD 74) employing the results of electron scattering. For ^{12}C , a Gaussian (harmonic oscillator) density was chosen of the form

$$\rho(r) = \rho_0 (1 + \alpha(r/a)^2) \exp[-(r/a)^2] , \quad (3.2.4)$$

with ρ_0 determining by requiring

$$4\pi \int \rho(r) r^2 dr = A = 12. \quad (3.2.5)$$

The proton optical potentials were generated in a quasi-phenomenological fashion by adopting the Woods-Saxon geometry obtained from a best fit to 185 MeV $^{12}\text{C}(p,p)^{12}\text{C}$ (IHS 73) elastic scattering data. Then the depth of the imaginary volume optical potential was computed via the KMT procedure outlined in Sect. 2.6, by substituting the normalization ρ_0 obtained from (3.2.5), for $\rho(r)$ in Eq. (2.6.1). The real depth was obtained from the ratio of the real to imaginary forward scattering amplitudes as described

in Sect. 2.6. The resulting proton optical potentials appropriate for the 90 MeV and 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reaction are tabulated in Table 3.2.2.

The sensitivity of the (π^+,p) calculations to the choice of the proton optical potential u_p was investigated. A calculation employing the purely phenomenological proton optical potential of (IHS 73) appropriate for 185 MeV protons was only 10% less in magnitude than a calculation which used the phenomenological proton potential geometry and KMT scaled depths appropriate for 220 MeV protons (which is more nearly correct for the 90 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reaction). In addition, inclusion of the phenomenological spin orbit potential of (IHS 73) had no effect at all on the calculations for the differential cross sections. An alternative proton optical potential was constructed from the harmonic oscillator nuclear charge density of (3.2.4). The procedure followed to construct this potential was similar to the methods of Sect. 2.6. The geometry was taken to be exactly the same as the geometry of the nuclear charge density, and the depths were scaled in accordance with Eqs. (2.6.5) and (3.2.5). These results are also listed in Table 3.2.2. Calculations of the 90 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}(3/2^- \text{ g.s.})$ reaction which employed the KMT harmonic oscillator, and quasi-phenomenological proton

Table 3.2.2. Proton Optical Potentials for
the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ Reaction

	$p + ^{11}\text{C}$ $T_\pi = 90 \text{ MeV}$	$p + ^{11}\text{C}$ $T_\pi = 180 \text{ MeV}$
<u>Harmonic Oscillator</u>		
$V_R (\text{MeV})$	-25.4	-19.0
(fm)	1.082	1.082
$a (\text{fm})$	1.692	1.692
$V_I (\text{MeV})$	-28.3	-31.8
(fm)	1.082	1.082
$a (\text{fm})$	1.692	1.692
<u>Quasi-phenomenological</u>		
$V_R (\text{MeV})$	-9.02	-6.73
$r (\text{fm})$	1.24	1.24
$a (\text{fm})$	0.64	0.64
$V_I (\text{MeV})$	-10.1	-11.22
$r (\text{fm})$	1.29	1.29
$a (\text{fm})$	0.78	0.78

optical potentials were performed. The calculation employing the harmonic oscillator proton potential (notorious for its poor behavior at large q) is observed to be three times lower in magnitude than the quasi-phenomenological calculation, although of the same shape. While the (π^+, p) calculations are sensitive to the proper choice of u_p , this sensitivity is smaller than that to the pion optical potential, discussed next.

The sensitivity of the (π^+, p) calculations to the specific choice of the pion optical potential u_π was checked and found to be extreme. This is especially demonstrated in the predicted energy dependence of the (π^+, p) reaction. Shown in Fig. 3.2.1 are three calculations of the $^{12}\text{C}(\pi^+, p)^{11}\text{C}(\text{g.s.})$ reaction, employing the quasi-phenomenological proton optical potentials of Table 3.2.1, and pure Kisslinger pion potentials (defined in (3.2.2)). The predictions for $T_\pi = 180$ MeV are almost forty times smaller than the predictions for $T_\pi = 90$ MeV. The third calculation is exactly the same as the 180 MeV calculation, except that the 90 MeV pion optical potential was substituted for the 180 MeV pion potential. This substitution raises the cross sections by about a factor of forty, thus revealing that the source of the energy dependence in the (π^+, p) calculations is entirely in the pion

Fig. 3.2.1 Predicted energy dependence of the $^{12}\text{C}(\pi^+, p)^{11}\text{C}(3/2^- \text{ g.s.})$ reaction, at 90 MeV (solid curve) and 180 MeV (dashed curve). The dotted curve is the same calculation as performed to generate the dashed (180 MeV) curve, only with the 90 MeV pion optical potential used to generate the solid curve. This isolates the cause of the energy dependence, namely, the pion optical potential. The curves employ the quasi-phenomenological proton potential and pure Kisslinger potentials.

optical potential. Elastic scattering predictions employing the 180 MeV pion potential are compared to 162 MeV $^{12}\text{C}(\pi,\pi)$ data in Fig. 3.2.2.

The effect of the S-wave absorption term was also investigated and found to be negligible. The inclusion of the angle transformation (which transforms u_K from the pion-nucleon system to the pion-nucleus system) was found to drop the (π^+,p) predictions by a factor of 2.5 for the ^{11}C $3/2^-$ state.

For the (π^+,p) calculations presented with the data in Chapter VII, pure Kisslinger pion potentials (with no angle transformation), and quasi-phenomenological proton optical potentials were used. The parameters of the Kisslinger potential will be those appropriate to the beam energy.

Fig. 3.2.2 A comparison of 162 MeV $^{12}\text{C}(\pi, \pi)^{12}\text{C}$ elastic scattering data (Pif+ 77) with the predictions for 180 MeV $^{12}\text{C}(\pi, \pi)^{12}\text{C}$ obtained from the pion optical potential of Table 3.2.1.

The High Resolution Spectrometer

CHAPTER IV

EXPERIMENTAL METHODS FOR THE (p,d) STUDIES

The recent development of an accelerator providing 800 MeV proton beams allows us to study neutron pickup with the (p,d) reaction on nuclei in the range of momentum transfer between 2 and 5 fm⁻¹. In order to resolve discrete final states separated by as little as 200 KeV, one needs a spectrometer capable of delivering a momentum resolution of .01%. The High Resolution Spectrometer (HRS) at the Clinton P. Anderson Meson Physics Facility (LAMPF) is such a spectrometer (How 74).

4.1 Beam Preparation

LAMPF (see Fig. 4.1.1) utilizes 750 KeV Cockroft Walton injectors to feed beams of H⁺, and H⁻ or polarized H⁻, to the main accelerator. The first 100 MeV of acceleration is provided by a 201.25 MHz post coupled Alvarez drift tube linac. This section feeds an 805 MHz, side coupled cavity wave guide accelerator that brings the hydrogen ions to their final energy of 800 MeV in pulses 500 microseconds long with a duty factor of 6% (Liv 77) (All 77). The

Fig. 4.1.1 General layout of the Los Alamos Meson Physics Facility, showing the relative locations of the various experimental areas.

H^- ions are magnetically separated from the simultaneously accelerated H^+ ions at the switchyard (Fig. 4.1.2). The H^+ ions travel to Area A, and the H^- ions are sent down Line X. In the following, the right handed coordinate systems are defined such that the positive z-axis points in the beam direction, and the x-axis always points outward along the radius of curvature of the various bending magnets that are encountered in the beam line and spectrometer. In Line X, foil and wire strippers tailor the x, y, and p (momentum) spread of the beam, as well as the intensity (Zei 71), (Bla 78). The beam then enters Line C (see Fig. 4.1.3) which separates protons destined for HRS from those traveling to Area B. This is accomplished by bending magnets LCBM01-03 and quadrupole doublets LCQM01-02. Next, the beam is horizontally dispersed by the two 15 kG, 57° bending magnets LCBM04-05.

The dispersed beam (and the x-axis) is twisted through 90° to the vertical by quadrupoles LCQM04-08, and the dispersion is matched by quadrupoles LCQM03, 09-11 to conform to the spectrometer optics (Zei 71) (Bla 78). Dispersion matching is necessary because the incident proton beam is not monochromatic in energy. The value of $\Delta p/p = .26\%$ (How 74) corresponds to an uncertainty in energy of ± 4.2 MeV for 800 MeV

Fig. 4.1.2 Experimental areas A, B, and C at LAMPF, including the switchyard. The HRS is housed in area C.

Fig. 4.1.3 Closeup of beam area C, showing the locations of the HRS and the various magnetic elements of the beam line.

protons. Clearly an uncertainty in energy of this magnitude would defeat the purpose of a spectrometer designed to resolve discrete nuclear levels. Consequently the beam is dispersed, to correlate the precise energy of the beam with spatial coordinates. The dispersion of the incident proton beam is then matched so that this correlation between energy and x is exactly that which the HRS expects to find at the target. When this matching has been achieved, particles having the same energy loss at the target are focussed to a line at the spectrometer focal plane even if their absolute energies differ. The matching condition is given by (Zei 71)

$$R_{11} \times \frac{R_{36B}}{R_{16S}} = \left(\frac{P_{A0}}{P_{C0}} \right) \frac{\frac{\partial M}{\partial P_A}}{\partial M / \partial P_C} \quad (4.1)$$

where R_{11} = spectrometer magnification, R_{36B} = dispersion of the beam preparation system, R_{16S} = dispersion of the spectrometer, M = mass-energy of the residual nucleus, and the subscript "0" in P_{A0} refers to the "central" momentum of particle A in the reaction $B(A,C)D$. The dispersion matching, as well as the selection of beam intensity and spot size at the HRS target, is performed by skilled operators working in the Central Control Room (CCR).

4.2 HRS

The HRS is a QDD (quadrupole-dipole-dipole) (Fig. 4.2.1) which operates in a VHV (vertical dispersion, horizontal scattering, vertical analysis) mode (Zei 71). This means the beam is dispersed vertically (x-direction) at the target, undergoes a horizontal (y-direction) scattering, and is bent vertically (perpendicular to the plane of the floor) during magnetic analysis by the HRS. Protons incident to the HRS pass through an evacuated two meter diameter scattering chamber with a target at the center (see Fig. 4.2.2). At the target, protons are focussed in the y direction and dispersed in the x direction, yielding a beam spot approximately 3 mm by 20 mm (Zei 71). To minimize interactions between the beam halo and the target frame, the target size is a generous 5 cm by 10 cm. The targets are discussed in Sect. 4.6.

The non-reacting beam is collected in a shielded beam dump. Reaction products to be analyzed are first focussed in the y direction and defocussed in the x direction by quadrupole magnet HSQM01, in order to reduce the size in y of the beam (and the required gap in the large bending magnets which follow) (Zei 71). Following the quadrupole are two 3.5 meter radius dipole magnets HSBM01,02 which vertically bend the beam 150° to detectors consisting of wire chambers

Fig. 4.2.1 Side view of the HRS, showing the locations of the detectors, spectrometer dipoles, and scattering chamber.

Fig. 4.2.2 Overhead view of the HRS scattering chamber, showing the locations of the various beam monitors such as the ion chambers (IC1-4), secondary emission monitor (SEM), and the elements (D1-4) of the monitor telescopes (MT1-2). Other devices shown are explained in the legend.

SCATTERING CHAMBER SCHEMATIC

LEGEND

- c-Evacuated Beam Pipe
- b-Evacuated Scattering Chamber
- a-Rotatable Target Table
- d-Rotatable Target Wheel
- e-Profile Monitor
- f-Vacuum Coupling to HRS
- g-T.V. Camera
- h-Mylar Window

and scintillators. The detectors measure such quantities as pulse heights, arrival times, and positions in x and y at various locations in z, as discussed below in Sects. 4.3 and 4.5. These quantities are then analyzed in an on-line computer to give the desired particle identification and momentum specifications, as treated in Sects 4.4 and 4.9.

4.3 Scintillation Detectors

The detectors used at the HRS (see Fig. 4.3.1) for the (p,d) experiments consisted of scintillators and delay line readout multi-wire proportional chambers. Jaw and veto scintillation counters served to define the spectrometer solid angle and focal plane. Hardware interrupts, that is, the acceptance of an event as a candidate for a deuteron, occurred only for those particles having reliable trajectories, defined as a four fold coincidence in the large scintillators S2, S6, S3 and S4 ($S2 \cdot S6 \cdot S3 \cdot S4$). A check on the interrupt detection efficiency was provided by the small test counters T1-3, which were less vulnerable to rate problems due to their small size.

Due to the very high (1 to 100 na.) incident proton beam currents necessitated by the low cross sections in the 800 MeV (p,d) reaction, it was found that a third test counter located about midway between T1 and T3 was needed. The ensuing triple coincidence

HRS Detectors

Fig. 4.3.1 Detectors located atop the HRS, including jaw counters (J1-2), veto counters (V1-2), test counters (T1-3), trigger (scintillator) counters (S2-4, S6) and delay line proportional counters (DL1, DL6).

($T_1 \cdot T_2 \cdot T_3$) was sufficiently free from random coincidences (caused by the enormous neutron background) to verify that the interrupt efficiency defined as $\frac{T_1 \cdot T_2 \cdot T_3 \cdot \text{INT}}{T_1 \cdot T_2 \cdot T_3}$ was greater than 99%.

As shown in Fig. 4.3.2, when the interrupt signal is in coincidence with the beam gate signal (as provided by CCR) and in anti-coincidence with a computer-busy signal, a strobe is generated which signals the PDP 11-45 data acquisition computer to write the various parameters associated with that event to magnetic tape for future detailed analysis.

4.4 Particle Identification

Scintillators S2, S3 and S4 perform the additional function of defining a 1.5 meter time-of-flight (TOF) path ($(S_3 \cdot S_4) - S_2$), which when used in conjunction with the differential energy loss as provided by the geometric mean of the pulse heights in S3 and S4 ($\sqrt{S_3 \cdot S_4}$), forms an excellent particle identification (PID) system. Typical results are displayed as TOF vs. dE/dx in Fig. 4.4.1 for laboratory scattering angles of 12.5° and 29° . As this figure demonstrates, events associated with protons, deuterons, and tritons are cleanly separated from one another.

At large angles, the background of particles other than deuterons passing through the spectrometer

HRS TRIGGER ELECTRONICS

Fig. 4.3.2 Schematic of the HRS trigger electronics. Shown are the signals from each end (up (U) and down (D)) of the trigger scintillation counters, and how these signals are combined to generate a flag used to identify a valid event. Also shown is the monitor telescope electronics.

Fig. 4.4.1 Particle identification spectra taken at the HRS on a ^{12}C target at spectrometer angles of 12.5° (A) and 20° (B). The particle groups corresponding to deuterons are enclosed by boxes. Tritons are to be found to the upper right of the deuterons, while protons compose the group to the lower left. The axes are differential energy loss or scintillation pulse height (x-axis) vs. time of flight (y-axis).

is seen to be substantial. This is due to the lower spectrometer fields demanded by the kinematics, and also the increased proton beam current needed to compensate for the rapidly falling (p,d) cross sections with angle. Although it is easy to see why the less energetic continuum tritons survive passage through a field (approximately 18 kG) set up to pass deuterons, it is harder to identify the particles that fall in the "proton" group. This group consists of fast moving, minimum ionizing particles which could include electrons, pole-tip scattered pions and protons, or neutrons. Regardless of the identity of the particles constituting the "proton" group, they were always well separated from the deuteron group of interest. During analysis, computer gates were set around the deuteron group, and these gates were checked at every angle.

4.5 Wire Chambers

During the (p,d) experiments, positions and angles of reaction products at the HRS focal plane were measured with delay line multiwire proportional chambers (MWPC) (Mor 78). The anode (cathode) planes consisted of wires in the y(x) plane spaced 3 mm (1 mm) apart which tapped into a 2.5 ns/cm microstrip printed circuit delay line. The different planes were separated by grounded sheets of aluminized mylar. The

charge collected from the passage of a particle near a wire was then fed into the delay line; the difference in arrival times at each end of the delay line is proportional to the location of the wire which collected the charge. In the cathode plane, position resolution was nearly 0.5 mm (FWHM) (Mor 78). Fig. 4.5.1 shows signals from one end of the delay line vs. signals from the other end for the anode and cathode planes of chamber one (DL1). Fig. 4.5.1 may be understood as follows. The signals from either end of the anode planes are denoted by AP or AN. Since these signals represent arrival times at each end of a fixed length delay line, they obey the relationship

$$(AP) + (AN) = L_{DL} + f(x) + g(\theta) \quad (4.5.1)$$

where L_{DL} is the time difference due to the fixed length of the delay line, $f(x)$ is the correction due to the uncertainty in the position along the actual anode wire the charged particle passed, and $g(\theta)$ is the drift time correction. In the (unrealistic) limit where $f(x) = g(\theta) = 0$, Fig. 4.5.1A would consist of discrete points (one for each anode wire) along the line $(AP) = L_{DL} - (AN)$. The length of the lines in Fig. 4.5.1A in the direction parallel to $y = x$ is due to the $f(x)$ and $g(\theta)$ terms in Eq. 4.2. Also visible in Fig. 4.5.1A is the discrete nature of the anode

Fig. 4.5.1 Shown are the signals from each end of the anode (A) and cathode (B) wire planes, as described in the text.

wire plane, which measured the scattering angle (relative to the nominal angle setting of the spectrometer at the rate of 0.1° per wire). The effects of attenuation and dispersion in the delay line are visible as a broadening of the lines in the direction perpendicular to $y = x$. It is seen that these effects are most important near the edges of the delay line, where one of the signals has been attenuated. Consequently, overlapping delay lines were used to improve the angular resolution as much as possible away from the center of the chamber.

Fig. 4.5.1B shows the continuous nature of the spectrum from the more closely spaced cathode wires, which provided a measure of the x coordinate of the event at the focal plane; this may be related to the excitation energy.

The efficiency of the chambers varied from 75% to 95%, although it hovered at $90\% \pm 5\%$ for all but a few runs. The efficiency of chamber one was defined as

$$\text{CHEFF}(1) = \frac{\text{PID} \cdot \overline{V1} \cdot \overline{V2} \cdot \overline{J1} \cdot \overline{J2} \cdot \text{DL1} \cdot \text{DL2}}{\text{PID} \cdot \overline{V1} \cdot \overline{V2} \cdot \overline{J1} \cdot \overline{J2} \cdot \text{DL2}} \quad (4.5.2)$$

A similar expression for CHEFF(2) was multiplied by CHEFF(1) to give the chamber efficiency used (CHEFF). As can be seen from Fig. 4.4.1B, the event rate at

large angles for continuum tritons passing through the spectrometer was much larger than that for deuterons. Consequently, triton events were used as the PID element of the chamber efficiency calculation at these large ($>20^\circ_{\text{lab}}$) spectrometer settings. Chamber efficiencies calculated using tritons compared well with efficiencies calculated using deuterons at those spectrometer angles where both particles had good counting rates.

Although the chambers typically draw less than 2 μa of current in (p,p') experiments, during the (p,d) data acquisition, currents of up to 9 μa were tolerated at the normal operating voltage of approximately 2000 volts. This was necessary because of the small deuteron yields from the (p,d) reaction, requiring the use of large (up to 100 na.) beam currents, which in turn kept the chambers busy handling background events.

4.6 Targets

Since the (p,d) studies described here are survey experiments of a reaction with a very low ($\leq 1 \mu\text{b/sr}$) cross section, the ever-present compromise between resolution and counting rate tended to favor counting rate considerations.

The ^7Li targets were natural (93% ^7Li) Lithium with areal densities of 349 and 940 mg/cm^2 ($\pm 3\%$).

These thicknesses were measured with a micrometer at several locations on each target. To minimize surface contamination, the Lithium targets were stored in pump oil and wiped clean with Ketone just before placement in the scattering chamber.

The ^{12}C targets had total areal densities of 173 mg/cm^2 and 328 mg/cm^2 ($\pm 3\%$). They were composed of Pilot B scintillator material ($\text{C}_{10}\text{H}_{11}$) containing natural carbon (99% ^{12}C). Several rods were made of this material to serve as special targets for various tune up procedures for calibrating angles at the target. The areal densities were computed by measuring the target thicknesses with a micrometer.

The ^{13}C target had a thickness of 312 mg/cm^2 ($\pm 3\%$), and contained a 15% ($\pm 1\%$) ^{12}C impurity, as well as a 2% ^1H impurity (Bla 78). This target was self-supporting, consisting of approximately 92% isotopic purity ^{13}C powder held together with glue (Bla 78).

A special cell was constructed to contain the water target for ^{16}O . It consisted of a frame holding two thin mylar sheets. Distilled water was introduced between the mylar sheets in such a way that air bubbles were removed. The resulting target bowed out in the vacuum of the scattering chamber, presenting a target of variable density. The effective ^{16}O target

thickness, as measured using a procedure to be outlined in Sect. 4.12, was 458 mg/cm^2 ($\pm 8\%$).

The thin strip of 99% isotopically pure ^{25}Mg used as a target had a thickness of 101 mg/cm^2 as specified by the supplier. This target was smaller than the others and therefore the beam spot was monitored by means of TV cameras, phosphor targets, and a split ion chamber used as a beam profile monitor, to insure that all of the beam would fall on the target. In fact, it was standard procedure to check the beam profile on target at every angle, and tune the beam spot as needed.

The natural (92% ^{28}Si) silicon targets used were 86 mg/cm^2 , 306 mg/cm^2 and 475 mg/cm^2 ($\pm 3\%$). The thicknesses of the silicon targets were measured with a micrometer, as were the ^{40}Ca targets and the CD_2 target.

The ^{40}Ca targets were 48.4 mg/cm^2 (100% isotopic purity) and 235 mg/cm^2 or 470 mg/cm^2 (97% isotopic purity).

A CD_2 target with an areal density of $23 \text{ mg/cm}^2 \pm 3\%$ was used for normalization purposes.

All the targets could be mounted simultaneously on the 36-target capacity target wheel in the evacuated scattering chamber.

4.7 Monitors

Three independent methods of relative normalization are available at the HRS (Bla 78). Only one of these methods was found to be reliable over the entire range of beam currents used for the 800 MeV (p,d) reaction.

At spectrometer angles of less than 12° , unscattered protons from the incident beam collide with the support structure of the HRS. The large background which results requires that only small (~ 1 na.) beam currents be employed at these angles. Since the (p,d) differential cross sections are found to be very forward peaked, this requirement can be tolerated. By the same token, this rapidly falling cross section with scattering angle requires the usage of the largest beam currents (~ 100 na.) the HRS can tolerate at larger angles ($\geq 25^\circ_{\text{lab}}$). A study of the beam monitor rates vs. beam current revealed that only the monitor telescopes were reliable over this entire range of beam current.

The monitor telescopes consisted of two sets (see Fig. 4.2.2) of four carefully aligned small plastic scintillators operated in coincidence (see Fig. 4.3.2) and situated at fixed angles of $\pm 120^\circ$ with respect to the incident proton beam. During the ^7Li and ^{13}C data runs, these angles were $+80^\circ$ and -85° .

At these angles, the four-fold coincidences from the telescopes were sensitive only to beam-target interactions. The solid angle of 0.28 msr presented by the monitor telescopes was defined by the third element of the telescope which measured 1" x 1" and was located 152 cm from the target (Bla 78). These monitors were used to provide the relative normalization of the data.

The secondary emission monitor (SEM) located just downstream from the scattering chamber (see Fig. 4.2.2) was useful at spectrometer angles larger than 12° and at beam currents less than about 30-40 na. It provided a way to normalize the monitor telescopes to beam current for the different targets in an absolute fashion.

In addition, three ion chambers were available (see Fig. 4.2.2). However, the background associated with beam passage through IC1 and IC2 was so large that these ion chambers were swung out of the beam for most angles, allowing higher beam currents to be utilized.

Since the monitor telescopes (MT) provided the relative normalization of the (p,d) data, a study was made to insure that the target angles chosen did not present any blind spots to the MT. In this study, a plot of $\frac{MT \times \cos \theta_t}{SEM}$ vs. θ_t was found to be flat, where θ_t = target angle with respect to the incident beam.

The ${}^7\text{Li}$ and ${}^{13}\text{C}$ angular distributions were carried out at a fixed target angle of 12° , while all other targets were set to 40° with respect to the incoming beam (see Fig. 4.2.2).

4.8 Acceptance

Ideally, the acceptance of a detector remains constant over its active area, falling abruptly to zero at its ends. The relative variation of solid angle with focal plane position was carefully checked and corrected for by sweeping strongly populated peaks across the focal plane. During the first (p,d) experiment (denoted as LAMPF experiment number 265), which measured angular distributions for ${}^7\text{Li}$ and ${}^{13}\text{C}$, the 2.185 MeV 3^+ state of ${}^6\text{Li}$ from the ${}^7\text{Li}$ (p,d) reaction was used. Data on the other targets were taken six months later (Expt. 117) when the ground state and 6.48 MeV $7/2^-$ state of ${}^{11}\text{C}$ resulting from the ${}^{12}\text{C}$ (p,d) reaction were used to study the focal plane efficiency. In each case the spectrometer fields were dropped in steps of approximately $\frac{1}{2}\%$ in particle (deuteron) delta, where $\delta \equiv \frac{p - p_s}{p_s}$, p_s = central momentum of the spectrometer, and p = particle momentum. The fields were accurately set with nuclear magnetic resonance probes.

Whenever it was necessary to raise the fields in the HRS dipoles, the current was first dropped to zero, then raised to saturation, and finally set to

the desired value in order to follow the hysteresis loop.

The resulting relative efficiency curves are displayed in Fig. 4.8.1, where channel number (or δ) is plotted vs. yield where

$$\text{yield} = \frac{\text{Area} \times \text{LT} \times 10^3}{\text{MT} \times \text{CHEFF}} \quad (4.8.1)$$

Here, LT is the computer live time defined as the ratio of strobes to interrupts (see Fig. 4.3.2), MT is the number of counts in the monitor telescope, and CHEFF is the chamber efficiency as defined in Sect. 4.5. A line was fit to the data of Expt. 265, $y(x) = .986 \times -1360$, via a linear least squares method. This fit is shown in Fig. 4.8.1A. The focal plane efficiency for Expt. 265 was therefore taken to be

$$\text{FPEFF}(x) = (8.61 \times 10^{-5})x - .119 \quad (4.8.2)$$

where x is the channel number of the resultant 2.185 MeV 3^+ level in ${}^6\text{Li}$, and channel 13000 was arbitrarily chosen to have $\text{FPEFF}(13,000) = 1$. The absolute normalization is treated in Sect. 4.11.

Fig. 4.8.1A also shows the focal plane efficiency for two data acquisition conditions. As a reminder, the quantity x angle is defined as the angle between the momentum vector of the reaction product

Fig. 4.8.1 The focal plane relative efficiency data taken for experiment 265 (A) and experiment 117 (B). The solid line in (A) is the fit discussed in the text. Values obtained with wide and narrow cuts in X angle are contrasted in (A), while in (B) values for different final states in $11C$ are compared.

and the x-axis at the target. Fig. 4.8.1A shows that the focal plane efficiency calculated with a tight x angle gate agrees nicely with the results obtained via the wide x angle gate. The extent of these gates is portrayed in Fig. 4.8.2 which shows x angle (with cuts) vs. excitation energy. The wide x angle gate was that used to produce the final spectra which generated the angular distributions presented in Chapter VI.

The relative focal plane efficiency for the data of Expt. 117 was taken to be flat. Fig. 4.8.1B shows the focal plane efficiency for Expt. 117 as calculated from the ground state yield and the 6.48 MeV $7/2^-$ yield from the $^{12}\text{C}(p,d)^{11}\text{C}$ reaction. Possible reasons for the improved uniformity of acceptance in Expt. 117 include the following: the spectrometer was found to be misaligned with respect to the scattering chamber, and this problem was corrected during the period between the two experiments. During this period it was also discovered that the HRS was not level, and levelers were installed which were adjusted at every angle during Expt. 117.

4.9 Data Acquisition System

Data acquisition at the HRS is computer controlled. In addition, much of the processing of the parameters associated with an event is done within a

Fig. 4.8.2 Software gates set on x angle (y-axis) vs. focal plane x. Box 5 is the inner (central) box which constitutes the tight x angle cut. Box 4 is the wide x angle cut.

computer environment. Signals representing the raw event information from the detectors at the HRS focal plane (pulse heights, and timing signals from the scintillators and wire chambers) are processed electronically and then fed into two CAMAC crates in the counting house. One CAMAC crate contains time to digital converters (TDC's), analog to digital converters (ADC's), coincidence registers, and interrupt modules. The other crate contains scalers and a visual display unit. When a strobe (see Sect. 4.3) is sent to the Bi-Ra Microprogrammable Branch Driver (MBD), the approximately 40 data words associated with an event are read from the CAMAC crates into an 800 word software buffer. When this buffer is filled it is written out to a Kennedy 9100 9 track, 1600 bpi (bytes per inch) magnetic tape drive. These transfers are controlled with the data acquisition program "Q." (Sch 76) When the PDP-11/45 data acquisition computer (124 K words core memory) is not busy with these transfers, some of the events are processed (as time is available, e.g., between beam bursts) into time of flight between various scintillators, scintillator arithmetic and geometric means, chamber times, coordinates (x, y, x angle, y angle) at the focal plane and at the target, momentum loss, scattering angle, and excitation energy of the residual nucleus (Bla 78).

The various quantities may be viewed on line by the experimenter, either as histograms or as one of two coordinates in a two-dimensional dotplot on Tektronix 4010 graphic display terminals. Examples of the data monitored during the experiment have already been shown as Figs. 4.4.1, 4.5.1 and 4.8.2.

Tapes containing the event information may be replayed off line to process all of the data written to tape. This allows more deeply considered gates, cuts and conditions to be applied to the events after the run, and polynomial correction terms may be included in the arithmetic to optimize the final calculations.

4.10 Correlations

The utility of computer based data acquisition is demonstrated for the case of software corrections to observed correlations in the data between excitation energy and both particle delta and angles at the target. Since the spectrometer dipoles were operated near saturation (typical dipole field settings are 14 kG for (p,p') and 17.9 kG for (p,d)), all of the correlations observed previously in 800 MeV (p,p') were remeasured for the (p,d) reaction.

The various correlations were removed by including a polynomial series in the directly measured quantities from the wire chambers at the focal plane.

The coefficients of the polynomial series for each correlation were determined from a least squares fit to rays (500 rays for each of seven focal plane positions) representing the experimentally measured wire chamber quantities. The experimental resolution of the excitation energy spectra was extremely sensitive to the correlations in δ and x angle. The feasibility of a given experiment depends on the ultimate resolution which is achieved, consequently considerable time was invested in an attempt to remove these correlations. The polynomial series used to correct δ was

$$\begin{aligned} \delta = & a_0 + a_1\psi + a_2\psi^2 + a_3x + a_4y + a_5\theta + a_6\phi + a_7x^2 \\ & + a_8\theta^2 + a_9y^2 + a_{10}\phi^2 + a_{11}x\theta + a_{12}y\phi + a_{13}x^3 \\ & + a_{14}x^2\theta + a_{15}x\theta^2 + a_{16}x\theta^3 + a_{17}x^2\theta^2 \end{aligned} \quad (4.10.1)$$

where ψ = scattering angle at the target, θ = focal plane x angle, ϕ = focal plane y angle, and x and y are the focal plane x and y coordinates. The a_i are the best fit coefficients from the program COFEPX (Bou78). The excitation energy was computed from δ and angles at the target, and then corrected via

$$\begin{aligned} EE = & EE + a_0 + a_1x + a_2\theta + a_3x\theta + a_4x^2 + a_5\theta^2 + a_6x^2\theta \\ & + a_7x^4 \end{aligned} \quad (4.10.2)$$

Fig. 4.10.1 Focal plane θ vs. excitation energy with (A) and without (B) the polynomial correction terms discussed in the text.

An example of the effect of these corrections may be seen in Fig. 4.10.1, where excitation energy is plotted vs. focal plane θ (x angle) with the full corrections for δ (Fig. 4.10.1A) and with linear correction terms in the δ polynomial only (Fig. 4.10.1B). The inclusion of higher order correction terms substantially improves the resolution.

In addition to the correlations previously found for normal (p,p') scattering, a very strong correlation was observed between vertical angle at the target out of the scattering plane (x angle) and excitation energy. It was necessary to reject those events with vertical angles at the target outside of a specified range. This cut, made in the software, is displayed in Fig. 4.8.2, along with a tight cut used to obtain higher resolution spectra (but with poorer statistics).

4.11 Absolute Normalization

Absolute normalization was done extremely carefully for the (p,d) work, since the characteristics of the HRS were not known in regions so near saturation. The consistency of the methods used to normalize the data was checked by using three independent methods.

The 800 MeV p + H elastic scattering reaction has been carefully done at LAMPF by Willard et al. (Wil+ 76). At laboratory angles of 12.5° and 17.1°,

data at the HRS were taken for p + H elastic scattering as well as for $^{12}\text{C}(p,d)^{11}\text{C}$, on the same 125 mil $\text{C}_{10}\text{H}_{11}$ scintillator target. The $^{12}\text{C}(p,d)^{11}\text{C}$ cross section may then be expressed as

$$\sigma_{\text{Lab}}^{12\text{C}} = \sigma_{\text{Lab}}^{\text{H}} \left(\frac{Y_{12\text{C}}}{Y_{\text{H}}} \right) \left(\frac{MT_{\text{H}}}{MT_{12\text{C}}} \right) \frac{1}{.893} \quad (4.11.1)$$

where $\sigma_{\text{Lab}}^{\text{H}}$ is the known p + H elastic scattering cross section. The term Y_{H} is the number of counts detected from p + H elastic scattering at HRS multiplied by the correction factor CF where

$$\text{CF} = (\text{CHEFF} \times \text{CLT})^{-1} \quad (4.11.2)$$

and CHEFF is the chamber efficiency (see Sect. 4.5) and CLT is the computer live time (see Sect. 4.8). The MT values are the monitor telescope results for the two measurements. The number 0.893 is the ratio of ^{12}C atoms to H atoms in the scintillator target. While this method has been used previously at the HRS for 800 MeV (p,p') normalizations, its application to the (p,d) reaction must be made with caution. This is because of the assumption implicit in Eq. (4.11.1), that the solid angle for p + H elastic scattering is the same as that for the much higher fields demanded by the (p,d) reaction.

In order to test this assumption, a second normalization was performed with a thick graphite slit of dimensions 1/2" by 1/2" placed at the entrance to the spectrometer. The .26 msr solid angle defined by this aperture was much smaller than the standard and nominal value of 3 msr (How 74). Agreement with the $^{12}\text{C}(p,d)$ cross sections obtained without the aperture was within 3%.

As a third method of normalization, the $^2\text{H}(p,d)^1\text{H}$ cross section was measured by detecting the deuterons using a 23 mg/cm^2 CD_2 target at a spectrometer angle of 12.5°_{Lab} ($= 28^\circ_{\text{C.m.}}$). The $^{12}\text{C}(p,d)^{11}\text{C}$ cross sections were then measured using the same target and computed as in Eq. (4.11.1), by comparing to the back angle neutron-deuteron elastic scattering data of Bonnet et al. (Bon+ 78) at a center of mass angle of $(\pi - 28^\circ)$. This method makes use of the fact that for this kinematical situation, the differences between $p + d$ and $n + d$ processes due to Coulomb effects are negligible (Bon+ 78). The results of this approach compare well with the other normalization methods, as shown in Table 4.11.1.

4.12 Cross Sections

The differential cross sections were computed via

Table 4.11.1. Absolute Normalization Results for the 800 MeV
(p,d) Data

	wide x angle cut		tight x angle cut	
	$\left. \frac{d\sigma}{d\Omega} \right _{\text{g.s.}}$	$\left. \frac{d\sigma}{d\Omega} \right _{6.48 \text{ MeV}}$	$\left. \frac{d\sigma}{d\Omega} \right _{\text{g.s.}}$	$\left. \frac{d\sigma}{d\Omega} \right _{6.48 \text{ MeV}}$
$^{12}\text{C}(p,d)^{11}\text{C}$ cross sections ($\mu\text{b/sr}$)				
12.5° normalization $\text{H}(p,p)\text{H}$	0.365	1.30	.386	1.58
12.5° normalization $^2\text{H}(p,d)^1\text{H}$	0.365	1.34	.343	1.52
12.5° normalization $\text{H}(p,p)\text{H}$ small aperture	0.369	1.35	.378	1.13

$$\frac{d\sigma}{d\Omega} \left(\frac{\mu b}{sr} \right)_{cm} = \frac{k \times J \times A \times Y \times \cos \theta_t}{T \times MT \times F_T} \frac{LT}{(CHEFF)(FPEFF)} \quad (4.12.1)$$

where J = the Jacobian relating the laboratory and center of mass solid angles ($J = d\Omega_{Lab}/d\Omega_{cm}$), A = target atomic weight (Sect. 4.7), T = target thickness in mg/cm^2 (see Sect. 4.6), MT = number of counts in the monitor telescopes (see Sect. 4.7), LT = computer live time (see Sect. 4.8), $CHEFF$ = chamber efficiency (see Sect. 4.5), and $FPEFF$ = focal plane efficiency (see Sect. 4.8).

The factor k is determined by comparing the $^{12}C(p,d)^{11}C$ cross sections obtained via (4.12.1) with $k = 1$, to those of Table 4.11.1.

The factor F_T is necessary to obtain absolute normalizations for the targets other than ^{12}C . This is because the measure of the incident proton beam current, made with the monitor telescopes, is target dependent. Consequently, for each target, runs were selected at spectrometer angles and proton beam currents such that the SEM was well within its range of reliability (see Sect. 4.7). The factor F_T was then taken to be the ratio of SEM to MT1 averaged over these selected runs. Thus the target dependence in the monitor telescope normalization was removed.

Eq. (4.12.1) was used to calculate the differential cross sections for all the targets. However, measurement of the ^{16}O target thickness presented special problems, since the areal density was non-uniform. This problem was solved by taking advantage of the hydrogen present in the liquid H_2O target. At laboratory angles of 12.5° and 17.1° , $p + \text{H}$ elastic scattering was measured on the hydrogen present in the water target, and compared to the known $p + \text{H}$ elastic scattering cross sections of (Wil+ 76), using the analogy of Eq. (4.11.1). The appropriately averaged ^{16}O target thickness obtained via this procedure was $458 \text{ mg/cm}^2 \pm 8\%$. Then, normalization of all subsequent ^{16}O data with the monitor telescopes eliminated any worries about non-uniform areal density or water leakage from the target, as well as the unlikely formation of bubbles within the target.

4.13 Experimental Uncertainties

The breakdown of experimental uncertainties can be made into three separate categories: errors associated with the absolute normalization of the cross sections, with the cross sections associated with a given state, and with the cross sections associated with a given angle.

The error in the absolute normalization was $\pm 8\%$ for Expt. 117. This number contains a 4%

uncertainty in the $p + H$ elastic scattering cross sections (Bla 78), a 3% error in target thickness (see Sect. 4.6), and a 4% error associated with the monitors (see Sects. 4.7 and 4.12). In addition, there is a generous 5% error allowed for the solid angle uncertainty (see Sect. 4.8).

The absolute normalization error for Expt. 265 was 12%. The error was larger than for Expt. 117 because the Expt. 265 data were normalized to ${}^7\text{Li}$ and ${}^{13}\text{C}$ runs taken during Expt. 117.

The variation of solid angle with focal plane position presents an error dependent on focal plane position. Since usually a given state was placed at about the same focal plane position at every angle, this error is reflected as an uncertainty in the absolute normalization of one state relative to another. Typical values for this error are ~5% for a state in the center of the focal plane (roughly at 15 MeV excitation energy) and increasing 1-2% every 5 MeV in each direction. This uncertainty is not included in the error bars shown on the final resulting data points.

Associated with every point in an angular distribution is a statistical error which usually is about 7%. In addition, there is the focal plane solid angle error and the error from the monitors. Each

point in an angular distribution represents data within an angular bin of 2.5° , except for some points which were split into bins $.6^\circ$ wide.

The ^{16}O data have a larger error associated with them of $\pm 12\%$, due to the way in which the ^{16}O target thickness was calculated (see Sect. 4.12).

The data taken during the (p,d) experiments are presented in Chapter VI.

A pictorial presentation of the EPICS spectrometer.

CHAPTER V

EXPERIMENTAL METHODS FOR THE (π^+, p) STUDIES

In order to study large ($2-5 \text{ fm}^{-1}$) momentum transfer neutron pickup leading to discrete final states with the (π^+, p) reaction, an intense, high quality, pion beam needs to be incident on a target viewed by a spectrometer capable of delivering an energy resolution of several hundred kilovolts or less. The Energetic Pion Channel and Spectrometer (EPICS) at LAMPF was designed to deliver an average flux of $4 \times 10^8 \pi^+/\text{sec}$ (at 1 ma primary proton beam current) to a pion spectrometer capable of 100 KeV resolution. The EPICS channel is able to deliver pions of kinetic energies between 50 and 300 MeV, and the spectrometer can detect particles of momentum up to about 800 MeV/c, corresponding to 295 MeV protons. This system was used to study the (π^+, p) reaction on some of the targets where the neutron pickup has also been studied with 800 MeV protons (see Chapters IV, VI).

5.1 Pion Production

The method used at LAMPF to create pions is, ironically, the inverse of the (π ,p) reaction which is to be studied. An extremely intense beam (0.5 ma) of 800 MeV protons is directed to a graphite target, where the pions are produced from the nucleons which comprise the graphite. In this process, pions of kinetic energies from 0 to about 600 MeV are produced and emitted in every direction, as are many other particles such as muons, electrons, kaons, neutrinos, and other nucleons.

The immense power (0.4 Megawatts) contained in the primary proton beam requires a production target of careful design indeed (Dun 76). Target A-1 (see Fig. 5.1.1) consists of a spoked graphite wheel which spins rapidly within a special target cell. The portion of the target wheel not in the beam is water cooled, providing adequate heat dissipation. Carbon graphite was chosen because of its low density (1.73 g/cm^3) and good thermal characteristics (Thi 70).

The target spot produced by the incoming (focussed) proton beam is $2 \text{ mm} \times 2 \text{ cm}$, with the short dimension in the vertical direction. This small vertical target size (focus) is needed to give the EPICS channel its good momentum resolution. The length of the production target in the direction

Fig. 5.1.1 Experimental Area A showing the location of the EPICS, A-1 production target, and other meson lines.

parallel to the primary beam is 3 cm, corresponding to an 11 MeV energy loss and an 8% reduction in the primary proton beam intensity. The remaining 92% of the beam continues into Area A, where it encounters two more meson production targets and finally a water cooled, cylindrical copper beam dump (see Fig. 5.1.1).

5.2 Pion Channel

The design of the EPICS channel involves a complicated trade-off between many factors. The short halflife (2.6030×10^{-8} sec) of the charged pion (Bar+ 78) precludes the luxury of an extensive beam preparation system such as that for the HRS discussed in Sect. 4.1, in spite of the requirement that variable energy pions be delivered. For a 50 MeV pion, for example, only 23% of the pions entering the 15.24 m long EPICS channel survive passage to the target. Additionally, those pions which do decay contaminate the pion beam with muons and electrons. The muons come from $\pi^{\pm} \rightarrow \mu^{\pm} \nu$ with a 2.6×10^{-8} sec halflife and almost a 100% branching ratio. The electrons come from the subsequent decay of the muons via $\mu^{\pm} \rightarrow e^{\pm} \nu \bar{\nu}$ with a halflife of 2.197134×10^{-6} seconds and a branching ratio of 98.6% (Bar+ 78). The compromise effected here was to use a relatively short (15.24 m) channel which is efficient over a limited range of

pion kinetic energies (100-300 MeV).

Another compromise which has been made at the EPICS concerns the selection of the angle at which the channel views the pion production target. Fig. 5.2.1 suggests that a 0° production angle is optimal. However, this implies that no production targets could be utilized downstream of target A-1. A small production angle also foils the option of two-armed coincidence experiments as well as providing a conflict between the shielding wall for the primary proton beam and the physical dimensions of the spectrometer. A production angle of 35° is used at the EPICS.

The EPICS channel is depicted schematically in Fig. 5.2.2, along with the optical functions of the various magnetic elements comprising the channel. Since the momentum acceptance of the channel is $\Delta p/p = \pm 2\%$, a means of reducing the momentum measuring accuracy of the EPICS to the design value of 2×10^{-4} (fwhm) is required. The intensity of pions is critical for reactions with low ($\leq 10 \mu\text{b}$) cross sections such as (π^+, p) . Consequently, the pion beam containing a broad range of incident energies is dispersed on the target, correlating pion momentum with target position, and incorporating the full momentum acceptance of the channel.

Fig. 5.2.1 Plot of the pion production cross section vs. production angle at the A-1 pion production target.

Fig. 5.2.2 Schematic of the EPICS channel and its associated optics in the horizontal plane (dashed line) and the vertical plane (solid line).

This dispersion is accomplished by bending magnets BM01-04 (see Fig. 5.2.2), which bend and disperse the particles in a vertical plane. A crossed E and B field velocity separator for removing protons from the pion beam is located between BM01 and BM02, although this was not used for the (π^+, p) experiments. Three multipole focusing magnets constructed at the University of Colorado (FM01-03) and consisting of quadrupole, sextupole (top) and sextupole (bottom) windings are located in the channel to optimize the resolution (Pet+ 75). There is a steering magnet (SM01) located just before BM04. Focusing in the vertical plane is point to point to point to point and in the horizontal plane it is point to parallel to point to parallel as may be seen in Fig. 5.2.2. The function of FJ04, located at a common focus of x and y, is to define the channel acceptance (vertical jaws) and the beam purity in terms of pion-proton separation resulting from the separator (horizontal jaws). Jaws FJ03 clean up the slit scattering from the previous jaws. The vertical FJ01 jaws control the vertical divergence of the beam, providing the main control for the intensity of pions delivered to the target.

The channel solid angle of 3.4 msr illuminates a reaction target of dimensions 20 cm $\times$ 8 cm. This enormous target size is the price paid for the

increased pion flux gained by dispersing the beam. While a target this size presents no drawbacks for some isotopes, its inconvenience was appreciated when well over \$20,000 had to be spent to obtain a high purity ^{13}C target for the $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ reaction.

5.3 Spectrometer

The EPICS spectrometer consists of five magnetic elements, three quadrupoles followed by two dipoles, operating in the VHV mode as explained in Sect. 4.2. The spectrometer is depicted schematically in Fig. 5.3.1. The channel delivers a dispersed beam to a target centrally located inside a 1 meter radius evacuated scattering chamber. Reaction products then pass through a Mylar foil of thickness 0.01 inch stretched 120° across the exit of the scattering chamber, as shown in Fig. 5.3.2. After passage through about a foot of air, particles falling within the solid angle of the spectrometer pass through a 0.03 inch thick Mylar foil and enter a vacuum system which extends the length of the spectrometer, ending just after the second dipole BM06, and before the rear detectors.

The first magnetic elements encountered by reaction products entering the spectrometer are the three quadrupoles QM01-03. The purpose of this quadrupole triplet is to provide a focus at the front

Fig. 5.3.1 Schematic representation of the magnetic elements comprising the EPICS spectrometer.

Fig. 5.3.2 Schematic diagram of the EPICS scattering chamber, its various elements, and relevant angles.

detectors located just upstream of dipole magnet BM05. This focus was designed to allow measurements of x and y at the front detectors independent of multiple scattering. The x coordinate (see Fig. 5.3.1), in the direction of the spectrometer bend, is used for projection back to the target and thus provides a measure of the incident pion beam momentum. The y coordinate, into the plane of Fig. 5.3.1, is proportional to ϕ (the scattering angle) at the target. An additional function of the quadrupole triplet is to reduce the phase space of the beam presented to the dipoles, as well as to provide some momentum selectivity to reduce the singles rate in the front detectors.

The measurement of the momentum of reaction products is performed by the two large 1.3 m radius magnetic dipoles BM05-06, which vertically bend the beam 120° to a series of detectors located at the exit of the spectrometer. The difference in x between the first (front) detectors and the second (rear) detectors located after BM06 determines the radius of curvature and hence the momentum of the reaction product. The y coordinate measured in the rear chambers determines y at the source. The solid angle resulting from this arrangement is 15 msr, with a momentum acceptance $\Delta p/p \approx 20\%$.

Although the spectrometer magnets were designed for a maximum momentum of 700 MeV/c, the $^{13}\text{C}(\pi^+, p)$ reaction on the (3,3) resonance produces protons of maximum momentum 800 MeV/c. Consequently, the cooling system for the spectrometer quadrupole and dipole magnets was upgraded. In spite of the extra efforts made for proper cooling, the power supplies ran extremely hot, and prompted the choice of pion energies slightly below the (3,3) resonance energy of 190 MeV which we would have preferred to use.

5.4 Scintillation Detectors

Just as in the HRS detection system, scintillators and multiwire proportional chambers were used for detection and localization of charged particles passing through the EPICS.

The jaw (J1N and J1P) and veto (V4N and V4P) plastic scintillation detectors helped to define the solid angle of the EPICS (see Fig. 5.4.1). Particles passing through these paddles were rejected because their trajectories took them too close to the edges of the (sometimes saturated) dipole magnets BM05 and BM06.

The focal plane scintillators S2P, S2N, S3P, and S3N were used to define a (~1.5 meter) time-of-flight path as well as to provide pulse height information useful in particle identification. Excellent

EPICS Detector Arrangement

Fig. 5.4.1 The EPICS detector arrangement showing the relative locations of the wire chambers, scintillators, and magnetic elements of the spectrometer.

separation of pions and protons was obtained by placing software gates on the particle groups observed in dotplots of S3-S2 TOF vs. $\sqrt{S2 \cdot S3}$ pulse heights. This was the same technique employed for particle identification in the (p,d) reaction and discussed in Sect.

4.4. Examples of PID dotplots are exhibited in Fig. 5.4.2 for a forward angle (typical case) and also for a backangle (worst case) situation. The extremely high proton momenta resulting from the conversion of the pion's rest mass into proton kinetic energy in the (π^+, p) reaction ensured the clean separation of reaction products after magnetic analysis by the EPICS.

In addition to providing PID, scintillators S2 and S3 formed event triggers when in coincidence with signals from the (FRONT) front chambers (see Fig. 5.4.3). Event triggers were $\text{EVENT} \cdot \overline{\text{BUSY}}$ where $\text{EVENT} \equiv S2 \cdot S3 \cdot \text{FRONT}$ and $\overline{\text{BUSY}}$ means beam gates (provided by the CCR) were in coincidence with computer-not-busy signals. Formation of an event trigger meant the particle responsible had an acceptable trajectory through the spectrometer, and the data acquisition system was notified that an event suitable for writing to magnetic tape was available.

Scintillator S1 was removed for the (π^+, p) studies to help improve the resolution. It was normally used to provide a long TOF path to the focal

Fig. 5.4.2 Particle identification dotplots of
(^{12}C target) TOF vs. dE/dx as defined in
Sect. 4.4, at spectrometer angles of 20°
(A) and 100° (B).

Fig. 5.4.3. Diagram of the fast electronics used at EPICS for this experiment. The event circuit is at the top, and chamber circuits at the bottom. Raw signals enter from the left. Arrows represent the various scaler, ADC, TDC, and trigger inputs. (Figure taken from the thesis of Iversen (Ive 78))

plane scintillators S2 and S3 for PID, however, the large spectrometer fields used for the (π^+, p) studies eliminated most μ and e contamination so that use of S1 was unnecessary. In addition, the freon Cherenkov detector was not used for the same reason.

5.5 Wire Chambers

Positions and angles were measured with delay line readout multiwire proportional chambers at the EPICS. These chambers are similar in construction to those used at the HRS and described in Sect. 4.5. However, they differ in that improved resolution is obtained at the EPICS by utilizing the drift time information available in the time sum $t_S = t_A + t_B$ where t_A and t_B are the arrival times of signals from each end of the delay line relative to a common (scintillator) start signal. When a charged particle passes an anode wire, the time it takes for the electrons created by the passage of the charged particle to drift to the wire is proportional to the distance from the wire to the point of closest approach of the charged particle. Once the anode wire closest to where the charged particle passed has been identified (with $t_D = t_A - t_B$ as described in Sect. 4.5), the distance from this anode wire is determined from t_S . The decision whether to add or subtract this drift distance from the position obtained via the time

difference t_D is made by comparison to positions measured by an identical chamber offset from the first by half the anode wire spacing, and closely placed directly downstream of the first chamber. Consequently the improvement in resolution costs the experimenter a factor of two increase in the number of chambers. Position resolution obtained with this method has been reported to be as good as .34 mm (FWHM) (Mor 78).

The front chambers, F1, F2, F3, and F4 (see Fig. 5.4.1) were double drift chambers in the sense that drift information from both the anode and cathode planes is utilized. Since two pairs of drift chambers were used here, the front chambers were able to measure positions and angles independently of the rear chambers. Efficiency of the front chambers was taken to be

$$FCHEFF = \frac{ALL \cdot ALL \cdot ALL \cdot ALL \cdot ALL \cdot ALL \cdot ALL \cdot ALL}{\overline{1A} \cdot \overline{1C} \cdot \overline{2A} \cdot \overline{2C} \cdot \overline{3A} \cdot \overline{3C} \cdot \overline{4A} \cdot \overline{4C}} \quad (5.5.1)$$

where ALL is the logical "and" of all chambers,

$$ALL = 1A \cdot 1C \cdot 2A \cdot 2C \cdot 3A \cdot 3C \cdot 4A \cdot 4C \cdot 5 \cdot 6 \cdot 9 \cdot 10 \quad (5.5.2)$$

and $\overline{1A}$ means ALL except for chamber one anode and $\overline{1C}$ means ALL except for chamber one cathode.

Chambers 5, 6, 9, and 10 were the rear chambers (see Fig. 5.4.1). These chambers utilized drift information in the x-direction only, where the momentum dispersion is made. Again, two pairs of drift chambers R1, R2, R5, and R6 are required for angle measurements. A third, redundant pair of rear chambers R3 and R4, were not used for the (π^+, p) measurements. The rear chamber efficiency was defined as

$$RCHEFF = \frac{ALL \cdot ALL \cdot ALL \cdot ALL}{5 \cdot 6 \cdot 9 \cdot 10} \quad (5.5.3)$$

where ALL is defined as in Eq. (5.5.2) and $\bar{5}$ means ALL except for chamber five.

A further efficiency calculation was performed for the drift chambers to eliminate events whose position as measured with the first member of a drift chamber pair differed radically from the position as obtained with the second element of the drift chamber pair. The drift efficiency was defined as

$$DREFF = \frac{P \cdot GATE}{FRONT} \quad (5.5.4)$$

where P denotes a good proton event as determined by the PID system outlined in Sect. 5.4, and FRONT means a good hit in all the front chambers, i.e., $FRONT = 1 \cdot 2 \cdot 3 \cdot 4$. The quantity GATE is a logical AND of three software gates G1, G2, and G3. Each gate

consisted of a check on the sum of the drift position differences as measured in the anode or cathode planes of the front or rear chambers. For example, the front anode drift difference gate G1 was placed on the calculated quantity

$$FADD = (F1A - F2A)^2 + (F3A - F4A)^2$$

where F1A and F2A were signals from the first front drift chamber pair of anode planes, etc. Similar expressions hold for G2 (front cathode drift difference) and G3 (rear anode drift difference). The widths of these gates were 6.5 mm (G1), 3 mm (G2), and 7.8 mm (G3).

The total efficiency of the chambers was taken to be the product

$$CHEFF \equiv FCHEFF \cdot RCHEFF \cdot DREFF \quad (5.5.5)$$

5.6 Targets

The targets chosen for the (π^+, p) studies were isotopes of Carbon and Lithium. The ${}^6\text{Li}$ target consisted of 98.7% isotopically enriched ${}^6\text{Li}$, with an areal density of 314 mg/cm^2 . The natural Li targets were 93% ${}^7\text{Li}$ and had thicknesses of 88 mg/cm^2 and 245 mg/cm^2 . The thin ${}^7\text{Li}$ target was only used at a spectrometer angle of 20° (lab). All three Lithium

targets were stored in pump oil and wiped clean prior to installation in the 4-target capacity target ladder. All targets were mounted in large aluminum target frames.

The natural C targets were 99% ^{12}C with areal densities of 228 mg/cm^2 and 476 mg/cm^2 . In addition, a CD_2 target was used for some runs, with an areal density of 194 mg/cm^2 . The ^{13}C targets consisted of a 90% isotopically enriched ^{13}C target which had a thickness of 430 mg/cm^2 and a special 99% isotopically pure ^{13}C target which had a thickness of 210 mg/cm^2 .

Errors in the target thicknesses were estimated to be approximately 3%.

5.7 Monitors

A variety of means of obtaining relative normalization at the EPICS is available. An ion chamber located near the Al production target measured the relative amount of primary proton beam on target (BOT), which is proportional (assuming constant optics) to the number of pions which pass through the channel and survive to interact with the target in the EPICS scattering chamber. Since residual radioactivity in the region of the production target can trigger the BOT ion chamber when the primary proton beam is off, care was taken not to accept counts from the chamber during these "beam-off" periods. This

technique for relative normalization may not be used to relate runs taken at different channel settings. Consequently, each channel setting was absolutely normalized by means of a technique described in Sect. 5.10, and the BOT provided the relative normalization for runs taken at the same channel setting.

Two ion chambers located in the EPICS scattering chamber provided a means of checking the normalizations obtained with the BOT method. These chambers were located on a rotatable target table at the exit (for undeflected projectiles) of the scattering chamber (see Fig. 5.3.2). Since these chambers are triggered by protons, muons, and electrons as well as pions, they must be absolutely calibrated at each channel setting just like the BOT.

Another means of relative normalization was provided by toroidal current monitor LACM01, which measured absolutely the current of the primary proton beam at the Al production target.

During the acquisition of the (π^+, p) data, ratios of each monitor to BOT were plotted for every run, to insure the quality of the BOT monitor (which was ultimately used as a measure of the incident pion flux).

5.8 Acceptance

The acceptance of the EPICS spectrometer was measured by varying the spectrometer field in steps of approximately 300 gauss while looking at the 180 MeV ${}^2\text{H}(\pi^+,p){}^1\text{H}$ reaction on a CD_2 target (see Sect. 5.6) at a spectrometer angle of 20° . This spectrometer field variation swept the proton peak across the focal plane in steps of 2% in particle delta (see Sect. 4.8). The resulting acceptance curve is shown in Fig. 5.8.1. As this figure shows, the acceptance of the focal plane is uniform to within about 10% for $-4\% < \delta < 4\%$, where most of the discrete final states were placed. An empirical correction based on this acceptance curve was applied to all the (π^+,p) data using a method similar to that described in Sect. 4.8.

5.9 Data Acquisition and Correlations

The data acquisition system is similar to that used at HRS and described in Sect. 4.9, and will not be discussed here.

Correlations such as those observed for the (p,d) reaction and detailed in Sect. 4.10 were seen in the (π^+,p) data as well. The polynomial correction function used to eliminate correlations in delta was (without entering the coefficients)

Focal Plane Efficiency

Fig. 5.8.1 Acceptance curve for the EPICS spectrometer obtained using the 180 MeV ${}^2\text{H}(\pi^+, p){}^1\text{H}$ reaction.

$$\begin{aligned}
\delta = & \text{XF} + \text{TF} + \text{XR} + \text{YF} + \text{PF} + (\text{XF})^2 + (\text{XF})(\text{TF}) \\
& + (\text{XF})(\text{XR}) + (\text{TF})^2 + (\text{TF})(\text{XR}) + (\text{YF})^2 + (\text{YF})(\text{PF}) \\
& + (\text{PF})^2 + (\text{XR})^2 + (\text{TF})(\text{XF})^2 + (\text{XF})(\text{TF})^2 \\
& + (\text{XF})(\text{TF})(\text{XR}) + (\text{TF})^3 + (\text{TF})(\text{YF})^2 + (\text{TF})(\text{PF})^2 \\
& + (\text{XR})(\text{TF})^2 + (\text{TF})(\text{XR})^2 + (\text{XR})^3
\end{aligned} \tag{5.9.1}$$

where XF = x coordinate as measured by the front chambers, TF = theta as measured by the front chambers (x angle), YF = y coordinate as measured by the front chambers, PF = phi as measured by the front chambers (y angle), and XR = x coordinate as measured by the rear chambers. The coefficients of the various terms in (5.9.1) were determined exactly as described in Sect. 4.10.

Resolution in the (π^+, p) studies was so poor (~800 KeV) due to target thickness and straggling that further correlations were obscured, so no polynomial correction to excitation energy was made.

5.10 Absolute Normalization

The method used to normalized absolutely the (π^+, p) data is analagous to the procedure with which the 800 MeV(p,d) data were normalized, as outlined in Sect. 4.11. At each channel setting (that is, each

incident pion energy) the ${}^2\text{H}(\pi^+, p){}^1\text{H}$ yield was measured and compared to the known $\pi^+ d \rightarrow pp$ cross section. In this way, all quantities such as spectrometer solid angle, pion constituency of the incident beam, target thickness, etc. were automatically accounted for.

The measured $\pi^+ d \rightarrow pp$ cross sections were taken from the compilation of C. Richard-Serre et al. (Ric+70). This reference parameterizes the angular dependence of the center-of-mass differential cross sections for the $\pi^+ d \rightarrow pp$ reaction via the following formula

$$\frac{d\sigma}{d\Omega}_{\text{c.m.}} = K(A + \cos^2 \theta - B \cos^4 \theta) \quad (5.10.1)$$

where θ is the center of mass scattering angle and the constants K , A , and B are listed in Table 5.10.1 for the energies used in the (π^+, p) studies reported here.

The (π^+, p) cross section σ_T (in mb/sr) on a target of thickness x_T mg/cm² and atomic mass A_T can be related to the measured $\pi^+ d \rightarrow pp$ cross section σ_D (in mb/sr) with the formula

$$\sigma_T = \sigma_D \frac{x_D}{x_T} \frac{4}{16} \frac{A_T}{A_D} \frac{N_T}{N_D} \frac{CF_T}{CF_D} \quad (5.10.2)$$

Here $x_D = 196$ mg/cm² is the areal density of the CD_2

Table 5.10.1 $\pi^+d \rightarrow pp$ Differential Cross Section
Parameterizations

$$\left. \frac{d\sigma}{d\Omega} \right)_{\text{c.m.}} = K(A + \cos^2 \theta - B \cos^4 \theta)$$

(Ric+ 70)

E_π	$K \left(\frac{\text{mb}}{\text{sr}} \right)_{\text{cm}}$	A	B
75	2.31	.220	.06
88 ^{a/}	2.57	.240	.2
91 ^{a/}	1.95	.27	0
165 ^{b/}	3.82	.224	.43
174 ^{b/, c/}	3.41	.22	.34
185 ^{c/}	3.10	.210	.15
187 ^{c/}	2.92	.236	.442

^{a/} 88 MeV and 91 MeV averaged to give 90 MeV normalization.

^{b/} 165 MeV and 174 MeV averaged to give 170 MeV normalization.

^{c/} 175 MeV averaged to the average of 185 MeV and 187 MeV to give 180 MeV normalization.

target used to measure the $\pi^+d \rightarrow pp$ cross section at the same laboratory spectrometer angle and channel setting as used to obtain N_T , the number of counts of the $A_T(\pi^+,p)A_T-1$ reaction that is to be calibrated. The factor $4/16$ accounts for the deuterium constituency of the CD_2 target, $A_D = 2$, and N_D is the number of counts detected from the ${}^2H(\pi^+,p){}^1H$ reaction. The correction factor CF contains efficiencies, target angle, and normalization to BOT, and is given by

$$CF = \frac{\cos \theta_t}{BOT \cdot DOMEGA \cdot CLT \cdot CHEFF} \quad (5.10.3)$$

The computer live time CLT is given by the ratio

$$CLT = \frac{EVENTS \cdot \overline{BUSY}}{EVENTS}$$

where EVENTS and $\overline{BUSY}$ have been defined in Sect. 5.4. The chamber efficiency is the product of the efficiencies of the front and rear chambers, and the drift efficiency. All these quantities have been defined in Sect. 5.5. The focal plane efficiency DOMEGA is an empirical correction as discussed in Sect. 5.8. The normalization to BOT is explained in Sect. 5.7. The angle θ_t is the target angle (see Fig. 5.3.2).

5.11 Experimental Uncertainties

Just as in the (p,d) situation outlined in Sect. 4.13, the experimental uncertainties in the (π^+ ,p) reaction may be broken up into three categories. There are errors associated with: (a) absolute normalization, (b) a given state, and (c) a given angle (or data run).

a. Absolute normalization:

The data of reference (Ric+ 70) are reported to be good to $\pm 4\%$, however, the parameterizations have a larger uncertainty associated with them. A generous average of the uncertainties for several energies yields $\pm 15\%$. Target thickness uncertainties are about 3% (see Sect. 5.6). A generous 2% error for the BOT monitor and a 5% error for the focal plane efficiency correction (see Sect. 5.8) must be included. This amounts to a total of $\pm 16\%$ error in absolute normalization.

b. Individual states:

The focal plane efficiency correction is, of course, more uncertain for states falling near the limits of the spectrometer acceptance (see Fig. 5.8.1). However, care was taken to place the discrete final states of interest from the (π^+ ,p) reaction in the region of uniform spectrometer acceptance, where the estimated scatter in yield is estimated to be 5%.

The statistical errors in the angular distributions are variable, and they range anywhere from 3% to 100%. This error is partially reflected in the error bars shown in the angular distributions which result from a gaussian peak fit using the peak fitting routine SPECTR (Pru 79). This includes an uncertainty in the background estimate by taking the error to be $(\text{total counts} + (2 \times \text{background counts}))^{1/2}$.

The angular uncertainty of $\pm 1.5^\circ$ reflects the 3° angular binning used to collect all of the (π^+, p) spectra, to be presented in Chapter VII.

CHAPTER VI

FEATURES OF THE 800 MeV (p,d) REACTION

Presented in this chapter are the results of the 800 MeV (p,d) studies on targets of ${}^7\text{Li}$, ${}^{12}\text{C}$, ${}^{13}\text{C}$, ${}^{16}\text{O}$, ${}^{25}\text{Mg}$, ${}^{28}\text{Si}$, and ${}^{40}\text{Ca}$. Previous (p,d) data published at proton bombarding energies greater than 200 MeV are limited to 770 MeV ${}^4\text{He}(p,d)$ (Bau+ 77) and 700 MeV ${}^{12}\text{C}(p,d)$ studies (Bak+ 74), both done at Saturne. For this reason, the targets in this study were chosen to span a range of A sufficient to reveal mass dependent systematics of the (p,d) reaction. In addition, targets were selected which would lead to residual nuclei with widely separated (≥ 500 KeV) final states whenever possible, since the experimental resolution averaged about 400 KeV in most cases.

Energy calibration of the various spectra was accomplished by assuming the HRS focal plane response was linear in excitation energy. Indeed, in the case of ${}^{13}\text{C}(p,d){}^{12}\text{C}$, where known levels of ${}^{12}\text{C}$ or ${}^{11}\text{C}$ (from the ${}^{12}\text{C}$ contaminant in the target) are populated over a 25 MeV range in excitation energy, the linear energy calibration yielded a better χ^2 (see Sect. 2.5) than the quadratic, cubic, or quartic linear least squares fits to the known levels. Consequently, the

procedure followed was to base the energy calibration of each spectrum on the ground state channel number, and the number of MeV per channel. This second parameter was obtained from the separation of any two strongly excited, known levels in the spectrum to be calibrated, when two such levels existed. If, as was the case for ^{28}Si and ^{40}Ca , no such known levels other than the ground states could be located, the MeV per channel parameter was obtained from a $^{12}\text{C}(p,d)^{11}\text{C}$ or $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ spectrum accumulated at the same angle as was the spectrum to be calibrated. Results of the energy calibrations are presented in the following sections for each of the targets studied.

Also presented in the following sections are the results of DWBA and CCBA calculations, discussed in Chapter II, for all of the ground state angular distributions, as well as for some transitions to excited levels. The exact finite-range (EFR) DWBA calculations were performed with the computer code DWUCK5 (Kun 79a). All two-step coupled-channels calculations (CCBA) were performed with the computer code CHUCK (Kun 79c). The one-step zero range calculations utilized the code DWUCK4 (Kun 79b).

Two sets of deuteron-nucleus optical potentials were used in the calculations presented in this chapter, corresponding to either the "KMT folded" or

the "well-matched" prescriptions defined in Chapter II. The proton potentials were fit to experimentally measured 800 MeV proton elastic scattering data on ^{12}C (Bla 78). The proton potentials and both sets of deuteron potentials are presented in Tables 2.6.2 and 2.7.1.

The data will be presented for each target in turn, and theoretical predictions provided as appropriate. The summary discussion, treating all these results, will be found in the final chapter of this thesis.

6.1 $^7\text{Li}(p,d)^6\text{Li}$

The first of the 800 MeV (p,d) results to be presented are the $^7\text{Li}(p,d)^6\text{Li}$ data. A typical spectrum at 9.5° (laboratory angle) is presented in Fig. 6.1.1. This figure displays counts per channel (proportional to the cross section) vs. channel number (proportional to excitation energy in the residual nucleus). Of note in the spectrum is the excitation of the $J^\pi = 1^+$, $T = 0$ ^6Li ground state, and the 3^+ , $T = 0$ first excited state at 2.185 MeV, which has about five or six times the strength of the ground state. No appreciable excitation of the 0^+ level at 3.56 MeV is observed. A broad, weakly excited structure near 5.5 MeV is seen which probably includes contributions from both the 5.37 MeV 2^+ $T = 1$ and 5.65 MeV 1^+ $T = 0$

Fig. 6.1.1 A spectrum of the 800 MeV ${}^7\text{Li}(p,d){}^6\text{Li}$ reaction at $\theta_L = 9.5^\circ$.

unbound levels. The assignment of these excitation energies is based on a scale determined by the ^{13}C (p,d) ^{12}C reaction, for which the many more known states make possible a superior calibration.

The relative intensities observed at lower proton bombarding energies are quite different from the situation just described (Kal+ 74), (Dev+ 69). At lower energies the 3^+ strength in the (p,d) reaction is nearly the same as that of the 1^+ , a result which is in agreement with spectroscopic strengths calculated from reasonable nuclear models (CK 67), (NG 70), (Nor 70). However, as the scattering angle, and with it the momentum transfer, is increased the strength of the 3^+ level relative to the 1^+ level is observed to increase (Kal+ 74). Another difference is that at lower bombarding energies, the 0^+ level is populated with half as much strength as the ground state (Kal+ 74).

The feature exhibited here by the comparison with lower energy (p,d) studies is the first example of several preferential populations of high angular momentum states as the bombarding energy is raised. This feature, discussed in Sect. 2.10, is a consequence of the large momentum transfer involved at 800 MeV in the (p,d) reaction, and was observed in all of the targets studied.

The angular distributions of the differential cross sections for the 1^+ and 3^+ levels of ${}^6\text{Li}$ are presented in Fig. 6.1.2. Cross sections were measured from 5° to 40° (lab frame) and were observed to fall from a maximum of several microbarns/sr to a minimum of several hundred picobarns/sr. The angular distributions are smooth and structureless.

Calculations employing the EFR DWBA have been performed for both the 1^+ and 3^+ levels in ${}^6\text{Li}$. Such calculations have proven to give reasonable results for the case of the 700 MeV ${}^{12}\text{C}(p,d){}^{11}\text{C}$ reaction to the ground state (RS 75). For the case of ${}^7\text{Li}(p,d){}^6\text{Li}$, all angular momentum transfer contributions ($p_{1/2}$ and $p_{3/2}$) have been combined incoherently. The spectroscopic amplitudes, taken from pre-existing theory, are tabulated in Fig. 6.1.3. The parameters of the $n + {}^6\text{Li}$ Woods-Saxon potential well have been adjusted (She 79) to give the correct r.m.s. charge radius for ${}^7\text{Li}$ of 2.35 fm (BBD 72). These parameters are tabulated in Table 2.4.1. Figure 6.1.4 shows the ${}^7\text{Li}(p,d){}^6\text{Li}(1^+\text{g.s.})$ data compared to calculations employing the KMT (see Sect. 2.6) and the well-matched (see Sect. 2.7) prescriptions for the deuteron optical potential. These potentials are tabulated in Tables 2.6.2 and 2.7.1. The KMT result lies slightly closer in absolute magnitude to the data, although

Fig. 6.1.2 Angular distributions for the 1^+ and 3^+ levels of ${}^6\text{Li}$ populated in the 800 MeV ${}^7\text{Li}(p,d){}^6\text{Li}$ reaction.

two step pathways
and spectroscopic amplitudes

Fig. 6.1.3 Two step pathways, deformations and spectroscopic amplitudes used as ingredients in the ${}^7\text{Li}(p, d){}^6\text{Li}$ calculations (CCBA) including pickup of a $p_{3/2}$ (solid line) and $p_{1/2}$ (dashed line) neutron.

Fig. 6.1.4 The 800 MeV ${}^7\text{Li}(p,d){}^6\text{Li}(1^+ \text{ g.s.})$ angular distribution is compared to EFR DWBA calculations employing KMT folded (solid line) and well-matched (dashed line) deuteron optical potentials with $\Sigma C^2S=0.83$.

each calculation fails to reflect the true strength of the data. Each calculation also exhibits a node near $\theta_{\text{c.m.}} = 20^\circ$ which is not reflected in the experimental angular distributions. The general shape of the well-matched calculation is superior to the KMT calculation. A possible reason for the poorer shape of the KMT calculation with respect to the data is that the two-parameter Fermi parameterization of the $d + {}^6\text{Li}$ folded density (see Sect. 2.6) becomes unreasonable for light nuclei such as ${}^6\text{Li}$, whose densities are usually parameterized in three parameter Fermi or harmonic oscillator forms (DDD 74). In fact, the χ^2 (goodness of fit parameter, see Sect. 2.5) of the fit to the results of the numerical integration of the $d - {}^6\text{Li}$ folding integral (see Sect. 2.6), was about twice as large as the value of χ^2 resulting from the fits to the folded densities of the other (higher mass) nuclei.

This same problem with the shape of the calculations (especially for the KMT calculation) also manifests itself for the 3^+ angular distribution, as demonstrated by Fig. 6.1.5. Also apparent in this figure is the gross under prediction of the strength of the 3^+ data by more than an order of magnitude.

The calculations predict a 3^+ level populated with approximately the same intensity relative to the

Fig. 6.1.5 Data for the 800 MeV ${}^7\text{Li}(p,d){}^6\text{Li}$ (2.185 MeV 3^+) level are shown, with four calculations: EFR DWBA predictions employing ($C^2S = 0.47$) KMT (solid line) and well-matched (dashed line) deuteron optical potentials, and ZR calculations ($D_0 = 80$ MeV - fm $^{3/2}$) utilizing one step (dotted line) and the two step (dot-dash line) pathways of Fig. 6.1.3. The ZR calculations demonstrate the unimportance of two step pathways for this excitation. The underprediction of the data by the EFR DWBA is not understood.

1^+ ground state as observed experimentally at lower bombarding energies (Kal+ 74). The enhancement of the 3^+ level relative to the 1^+ level, seen experimentally at 800 MeV, is not described by the EFR DWBA.

In an attempt to understand this discrepancy (KO 77), zero-range coupled channels (CC) calculations were performed for the 3^+ level, employing the coupling scheme, rotational model spectroscopic factors (She 79), and deformation parameter (She 79) of Fig. 6.1.3. Two calculations were performed. First, a CC calculation in zero-range which included only the pathway from the ${}^7\text{Li } 3/2^-$ ground state directly (p3/2 pickup) to the 3^+ level of ${}^6\text{Li}$. The normalization for this calculation was $80 \text{ MeV} \cdot \text{fm}^{3/2}$, which is the zero-range strength required to make ${}^{13}\text{C}(p,d){}^{12}\text{C}(\text{g.s.})$ EFR and ZR predictions agree with each other at 0° . Then, employing the same normalization, a two-step calculation was performed which also included pickup from the inelastically excited $7/2^-$ level of ${}^7\text{Li}$, as shown in Fig. 6.1.3. The results of these two calculations, also presented in Fig. 6.1.5, reveal that two-step processes are unimportant for the excitation of the 3^+ level. The reason for the strong excitation of this level relative to the ground state is not understood at this time.

6.2 $^{13}\text{C}(\text{p},\text{d})^{12}\text{C}$

Shown in Fig. 6.2.1 is a sample spectrum at 12° laboratory angle for the 800 MeV $^{13}\text{C}(\text{p},\text{d})^{12}\text{C}$ reaction. The energy calibration of this spectrum was particularly easy due to the well-known and widely separated levels of ^{12}C . For the calibration of levels populated at excitation energies greater than 16.1 MeV, contaminant peaks from the $^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ reaction (arising from the 15% ^{12}C impurity in the ^{13}C target) were used to extend the spectrum of known calibration levels up to about 23 MeV of excitation in ^{12}C .

Features of interest in the spectrum include the strong excitation of the 14.1 MeV $4^+ T = 0$ level, which is excited more strongly relative to the ground state than is seen at lower bombarding energies (Tak+ 68), (Sco+ 70), (Par+ 74). A similar statement applies to the 4.44 MeV $2^+ T = 0$ level. All levels populated in lower energy (p,d) studies are populated at 800 MeV. In addition, levels at 20.61 ± 0.04 and 25.4 ± 0.1 MeV are seen strongly at 800 MeV; these are only weakly populated, if at all, at lower bombarding energies (Par+ 74).

The lack of appreciable strength to the $3^- T = 0$ level at 9.6 MeV has interesting implications. The two-nucleon model (see Fig. 6.2.2, and the discussion pertaining to the ONM in the introduction to this

Fig. 6.2.1. Sample spectrum of the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction at 12° .

Fig. 6.2.2 Schematic (Feynmann-like) diagram of the two nucleon model (TNM) representation of the (p,d) reaction mechanism.

chapter predicts a cross-section which is roughly proportional to the stripping cross-section that leads to the same residual nucleus as populated in the (p,d) reaction. In terms of the triangle diagram shown in Fig. 6.2.2, the cross-sections of the TNM are proportional to the lower-right hand vertex which represents the reabsorption of a nucleon from the intermediate (d + nucleon) projectile. This same vertex also describes the stripping reaction, wherein a nucleon from the projectile is stripped off and captured by the target nucleus. The diagram representing such a process is the same as for the ONM in the (p,d) reaction, shown in Fig. 2.1, only reversed in time (read from right to left).

In the case of $^{13}\text{C}(p,d)^{12}\text{C}$, the one nucleon model pickup of a p-shell neutron from the $1/2^-$ ground state of ^{13}C must always lead to a positive parity level. Population of a negative parity level such as the 3^- can be understood by including the possibility of inelastic (vibrational) excitation in the target or residual nucleus, as discussed in Sect. 2.8. Alternatively, the 3^- could be populated in the TNM (or in the stripping reaction) if the nucleon which was stripped out of the projectile found itself in the next highest allowed orbital, the $d_{5/2}$ orbital. As a consequence of these last two statements, the following

may be said: If a level which is strongly populated in the (p,d) reaction is also strongly populated in the stripping reaction on the target two holes distant (such as $^{11}\text{B}(^3\text{He},d)^{12}\text{C}$, in this example), then either two-step processes are important in the excitation of the level, or the TNM is important, or both. If, on the other hand, a level is very weakly populated (relative to pure single neutron pick-up levels of nearly the same angular momentum quantum number) in the (p,d) reaction, which is strongly excited with the stripping reaction, then both two-step processes and the TNM are unimportant (or possibly both are important and interfere destructively to such an extent that they cancel—an unlikely event).

Shown in Fig. 6.2.3A is a spectrum of 43 MeV (bombarding energy) $^{11}\text{B}(^3\text{He},d)^{12}\text{C}$ taken at the University of Colorado 52" cyclotron facility with solid state detectors in order to observe a large range of excitation energy in the residual nucleus. Prominent in this spectrum is the 9.6 MeV 3^- level, which is excited strongly relative to the other levels, and can be understood as stripping of a proton into the $d_{5/2}$ orbital. The 3^- level at 9.6 MeV is only very weakly excited relative to the other levels in the 43 MeV $^{13}\text{C}(^3\text{He},\alpha)^{12}\text{C}$ reaction, as demonstrated in Fig. 6.2.3B from another spectrum taken under the same

Fig. 6.2.3 Shown are sample spectra of the 43 MeV $^{11}\text{C}(^3\text{He}, d)^{12}\text{C}$ reaction (A) and the 43 MeV $^{13}\text{C}(^3\text{He}, \alpha)^{12}\text{C}$ reaction (B) at $\theta_L = 30^\circ$. Note the enhanced excitation of the 9.64 MeV 3^- relative to the 4.44 MeV 2^+ level in the stripping reaction (A) as opposed to the pickup reaction (B).

conditions as part A. In fact, in the 43 MeV $^{13}\text{C}(^3\text{He},\alpha)^{12}\text{C}$ reaction, the ratio of the pickup strength to the 9.6 MeV 3^- state and another high spin state such as the 4.44 MeV 2^+ is less than one-tenth. Any enhancement of this ratio at 800 MeV could be interpreted as due to TS processes or the TNM or both. Fig. 6.2.1 shows that, in spite of the preferential population of high spin states previously noted for the 800 MeV (p,d) reaction, the ratio of the 3^- strength to the 2^+ strength is also very small for 800 MeV pickup (about 1:12 at 12°).

If the remote possibility of almost complete cancellation between TS and the TNM is overlooked, the conclusion that follows from the above observations on a critical test case transition is that the TNM is not dominating in the $^{13}\text{C}(p,d)$ reaction even at 800 MeV.

Angular distributions for most of the levels populated in the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction are presented in Fig. 6.2.4. Apparent from this figure are slight changes in the shapes of the angular distributions for different levels. The 14.1 MeV 4^+ level, which cannot be excited via a single step, has a much smoother angular distribution than the other levels. The 4.44 MeV 2^+ level also has a smooth angular distribution. Due to the rotational character

Fig. 6.2.4 Angular distributions for some of the levels excited in the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction.

of this state, TS contributions are expected here as well as for the 4^+ .

Presented in Fig. 6.2.5 is the $^{13}\text{C}(p,d)^{12}\text{C } 0^+$ ground state angular distribution with the results of EFR DWBA calculations employing KMT and well-matched deuteron optical potentials. Although the results of the two calculations are similar, the KMT prescription is seen to give a superior account of the data. The spectroscopic factors used to normalize absolutely the calculations in this section are those computed from the wavefunctions of Norton and Goldhammer (NG 70), (Nor 70) and presented in Table 2.1.1. The agreement of the KMT DWBA calculation with the experimental data is almost perfect in shape and magnitude: the calculation predicts the observed forward angle maximum (near $\theta_{\text{cm}} \approx 8^\circ$) as well as the minimum (near $\theta_{\text{cm}} \approx 18^\circ$). These wiggles in the angular distribution are attributable to the nodes in the momentum space neutron bound state wavefunction. Apparently, some high- q dependence of $\tilde{\phi}_n(q)$ is showing up in the 800 MeV (p,d) reaction, and is, in fact, correctly described by the DWBA.

Compared to the data in Fig. 6.2.6 are the results of KMT EFR DWBA calculations for $^{13}\text{C}(p,d)^{12}\text{C}$ to the two 1^+ states at 12.7 MeV ($T = 0$) and 15.11 MeV ($T = 1$), as well as the 16.1 MeV 2^+ $T = 1$ state.

Fig. 6.2.5 The 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}(0^+ \text{ g.s.})$ angular distribution is compared to EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials. Note the excellent agreement with the data of the KMT curve.

Fig. 6.2.6 Angular distributions of the 12.7 MeV 1^+
 $T = 0$, 15.1 MeV 1^+ $T = 1$, and 16.1 MeV 2^+
 $T = 1$ levels excited in the 800 MeV
 $^{13}\text{C}(p,d)^{12}\text{C}$ reaction are compared to EFR
DWBA calculations with the KMT deuteron
optical potential used in Fig. 6.2.4.

Reasonable agreement with the experimental data is observed, although the calculations appear to have too much structure.

The effects of TS pathways to the 4.44 MeV 2^+ $T = 0$ level were investigated, using the pathways and deformations (Bla 78), (Bla+ 77) shown previously in Fig. 2.8.1, the spectroscopic amplitudes of Table 2.1.1, and a $D_0 = 80 \text{ MeV} - \text{fm}^{3/2}$. The inelastic $3/2^-$ and $5/2^-$ levels at 3.68 MeV and 7.55 MeV, respectively, in ^{13}C are the strongest inelastic levels observed in the 800 MeV $^{13}\text{C}(p,p')$ reaction at back angles (Bla 78), (Bla+ 77). Fig. 6.2.7 presents the results of the investigation of TS contributions to the 4.44 MeV level of ^{12}C . Shown in this figure are the results of three calculations. The dot-dash curve depicts an EFR DWBA calculation employing the KMT deuteron optical potential of Table 2.6.1. The dashed curve is a zero-range one-step calculation presented to separate the effects due to ZR from those due to TS. The ZR TS calculation is the solid curve, normalized with the same D_0 as used to normalize the ZR one-step calculation. This $D_0 = 80 \text{ MeV} - \text{fm}^{3/2}$ was empirically determined by requiring the $^{13}\text{C}(p,d)^{12}\text{C}(0^+ \text{g.s.})$ predictions in ZR and EFR to agree. Comparison to the calculated value of $D_0 = 124 \text{ MeV} - \text{fm}^{3/2}$ (see Sect. 2.3) emphasizes the importance of performing the calculations in

Fig. 6.2.7 The effect of the coupled channels pathways (shown in Fig. 2.8.1) is demonstrated for the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ (4.44 MeV 2^+) angular distribution, by comparing the results of EFR DWBA (dot-dash line), ZR one step (dashed line) and ZR two step (solid line) calculations ($D_0 = 80 \text{ MeV-fm}^{3/2}$).

EFR. A comparison of the ZR one-step and two-step calculations reveals that the effect of two-step processes is to raise the predicted cross sections slightly and to wash out the structure. This kind of effect would serve to improve the agreement between the one-step EFR calculation and the data, suggesting that further improvement would be achieved if the EFR calculations could include TS processes.

Presented in Fig. 6.2.8B are the results of a two-step calculation to the 14.1 MeV 4^+ $T = 0$ level, employing the pathway (path 1) shown in Fig. 6.2.8A, and a zero range normalization of $D_0 = 80 \text{ MeV} \cdot \text{fm}^{3/2}$. The data are underpredicted by the calculation. However, the coupling scheme of Fig. 6.2.8A is incomplete. Due to the limitations of the CCBA program, only the few simple pathways of Fig. 6.2.8A could be included. Including the other pathways might increase the calculated cross sections, improving agreement with the data.

An estimate of the contribution from the only other pathway of major significance may be obtained from the strong coupling model. This pathway (path 2) includes quadrupole inelastic excitation of the $5/2^-$ level in ^{13}C , followed by $p_{3/2}$ neutron pickup to the 4^+ level of interest in ^{12}C . In the strong coupling (rotational) model, the spectroscopic factor for such

Fig. 6.2.8 Shown is the deformation, spectroscopic amplitude, and pathway (A) for the two step calculation of $^{13}\text{C}(p,d)^{12}\text{C}$ (14.1 MeV 4^+). The results of the TS calculation employing this pathway are compared to the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}(4^+)$ data in (B), with a $D_0 = 80 \text{ MeV-fm}^{3/2}$. Note that an estimate (given in the text) of the contribution from $p_{3/2}$ pickup to the 4^+ from the $5/2^-$ in ^{13}C raises the cross sections by a factor of 2.25, bringing the forward angle calculation into agreement with the data.

a process is given by (Sat 58)

$$S = \frac{2J_f + 1}{2J_i + 1} (J_f j K_f \Delta K | J_i K_i)^2, \quad (6.2.1)$$

where J_i (J_f) denotes the initial (final) state before (after) the pickup, j is the transferred angular momentum (3/2 in this example), K_i (K_f) is the spin of the bandhead of the initial (final) nucleus, and ΔK is the difference $K_i - K_f$. Therefore, in this crude picture, inclusion of path 2 should increase the predicted cross sections to the 4^+ by the factor

$$\left| \frac{S_{\text{path 1}}^{1/2} + S_{\text{path 2}}^{1/2}}{S_{\text{path 1}}^{1/2}} \right|^2 = 2.25. \quad (6.2.2)$$

This factor is just what is needed to bring the forward angle differential cross sections into agreement with the data, as may be seen from Fig. 6.2.8B. However, the calculation still underpredicts the data at back angles.

6.3 $^{12}\text{C}(p,d)^{11}\text{C}$

Shown in Fig. 6.3.1 are spectra of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction collected at laboratory angles of 5° and 19° . Many discrete levels are discernible up to an excitation energy of 10.6 MeV, after which a mysterious broad region of enhanced excitation near

Fig. 6.3.1 Spectra collected at 5.2° (A) and 19° (B) of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction.

13.2 MeV is observed. The well known and widely separated level structure of ^{11}C makes possible the positive identification of most levels. Consequently, the calibration procedures of this chapter were tested by comparison to these known levels. Linear calibrations were performed based upon two levels in a $5^\circ \text{ }^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ spectrum, upon two levels in a $5^\circ \text{ }^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}$ spectrum, and upon the average separation of two levels in $^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ spectra at five different angles. The results of these calibrations are presented in Table 6.3.1, from which it may be observed that agreement of the results of each calibration technique with the known excitation energies is within the errors assigned to each excitation energy. In fact, the results of each calibration agree well enough that confidence in each technique is established.

The relative intensities of the levels populated in the 800 MeV $^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ reaction are vastly different from those populated at lower energies (<200 MeV) (Bac+ 69), (Lee+ 68). The $6.48 \text{ MeV } 7/2^-$ state dominates at 800 MeV, but is only weakly seen at lower bombarding energies. Additionally, the ground state strength no longer dominates the spectrum as it does at lower energies. The excitation of high spin states such as $7/2^-$ is expected to be favored in the

Table 6.3.1. 800 MeV $^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ Calibration

Peak #	Excitation Energy ^{a/}	Excitation Energy ^{b/}	Excitation Energy ^{c/}	Excitation Energy ^{d/}	J
1	0.000 ± .1	0.000 ± .1	0.000 ± .1	0.000	3/2 ⁻
2	1.968 ± .1	1.970 ± .1	1.961 ± .1	2.000	1/2 ⁻
3	4.285 ± .1	4.289 ± .1	4.269 ± .1	4.319	5/2 ⁻
4	4.779 ± .1	4.783 ± .1	4.761 ± .1	4.804	3/2 ⁺
5	6.480 ± .1	6.485 ± .1	6.456 ± .1	6.480	7/2 ⁻
6	8.193 ± .2	8.199 ± .2	8.163 ± .2	8.100	3/2 ⁻
7	8.592 ± .2	8.600 ± .2	8.561 ± .2	8.70, 8.40, 8.66	5/2 ⁺ 5/2 ⁻ 7/2 ⁺
8	9.981 ± .2	9.989 ± .2	9.944 ± .2	10.08	7/2 ⁺
9	10.56 ± .20	10.57 ± .20	10.52 ± .20	10.68	9/2 ⁺
10	13.22 ± .25	13.23 ± .25	13.17 ± .25		

^{a/}Present work: linear calibration of 5° spectrum based upon ground state and 6.48 MeV levels.

^{b/}Present work: linear calibration of 5° spectrum based upon ground state and calibration constant from $^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}(1.37 \text{ MeV}, 6.01 \text{ MeV})$ at 5°.

^{c/}Present work: linear calibration based upon 5 angles, using the ground state and 6.48 MeV levels.

^{d/}Levels identified from the compilations of (Ajz 75).

large momentum transfer regime (see Sect. 2.10).

Also, TS processes may play a more visible role at these energies, as has been pointed out previously (KO 77). Many of the observed levels are known to have appreciable parentage in the rotational and vibrational states of ^{12}C . For example, the pickup of a $p_{3/2}$ neutron preceded by inelastic excitation of the 4.44 MeV 2^+ of ^{12}C yields $(2^+ \times (p_{3/2})^{-1})$ states of angular momentum $1/2^-$, $3/2^-$, $5/2^-$, and $7/2^-$ which may be identified at excitation energies of 2.00 MeV, 4.804 MeV, 4.319 MeV, and 6.48 MeV, respectively. The intensities of levels populated in ^{11}C relative to known pure hole levels excited via a single-step process (SS) such as the $3/2^-$ ground state, have been used to gauge the importance of two step mechanisms. A comparison (KO 77) of 185 MeV and 700 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reveals that the intensity of the 2.00 MeV $1/2^-$ level relative to the ground state is nearly the same at the two energies, suggesting that the 2.00 MeV level is mainly a $(p_{1/2})^{-1}$ hole state. However, the intensities of the $5/2^-$, $7/2^-$, $3/2^-$ and $(5/2^-, 5/2^+, \text{ or } 7/2^+)$ levels at excitation energies of 4.33 MeV, 6.48 MeV, 8.11 MeV and about 8.66 MeV relative to the single step (SS) ground state excitation are much greater at 700 MeV (and 800 MeV) than at 185 MeV. This difference has been (KO 77) ascribed to the rising

importance of TS contributions as the beam energy goes up. The reasons quoted by authors Källne and Obst are twofold. First, the experimentally measured proton elastic scattering cross sections rise a factor of four to five from 185 MeV to 700 MeV for the excitations of the 4.44 MeV 2^+ and 9.6 MeV 3^- levels of ^{12}C . This increase is expected in the deuteron channel as well, and thus a reaction mechanism dependent on inelastic excitations is favored at higher bombarding energies. The second reason may be seen by examining the momentum transfers in each mechanism. In the SS model, the momentum transfer is $q = |\vec{p}_p - \vec{p}_d|$. In the TS model, however, at least one more momentum $\vec{q}'_p$ is involved which represents the momentum of the inelastically scattered proton in the process $p + ^{12}\text{C} \rightarrow p' + ^{12}\text{C}^* \rightarrow d + ^{11}\text{C}$. Then the momentum transfer is $q = |\vec{p}_p - \vec{p}'_p + \vec{p}'_p - \vec{p}_d|$. The cross sections in the TS model are enhanced because the wavefunction of the picked up neutron is sampled at lower momenta whenever the inequality $|\vec{p}'_p - \vec{p}_d| < |\vec{p}_p - \vec{p}_d|$ is satisfied. The fact that this results in larger cross sections may be seen from Eq. (2.2.7) in the PWBA, where the transition amplitude is shown to be proportional to the momentum space wavefunction of the picked-up neutron. This wavefunction has been plotted vs. q in Figure 1.1, from which it may be seen that

lower q implies larger $\phi_n(q)$ (if the effects of the minima are ignored). The greater momentum sharing allowed in multistep processes should enhance the yield to levels populated in such a fashion.

Angular distributions for the levels populated in ^{11}C are presented in Figs. 6.3.2 and 6.3.3. Except for the presence of a slight minimum near $\theta_{\text{cm}} = 15^\circ$ for some levels (mainly those levels which are expected to be of predominantly single hole character), the angular distributions are featureless, dropping about five orders of magnitude in forty degrees.

In order to establish that the angular bin of 2° used to collect the $^{12}\text{C}(p,d)^{11}\text{C}$ data was not too coarse, the angular distribution of the 6.48 MeV $7/2^-$ level of ^{11}C was plotted in bins of only 0.5° . The results of this binning are presented in Fig. 6.3.4, from which it may be seen that the angular binning of 2° is sufficient to resolve the structure present in the data.

Additional evidence that the binning is sufficiently fine may be obtained from a perusal of the only other medium energy (p,d) data extant, 770 MeV $^4\text{He}(p,d)$ done at Saturne (Bau+ 77), and 700 MeV $^{12}\text{C}(p,d)$ performed at Saclay (Bak+ 74). Because the 700 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ data comprise the only other work which overlaps the work of this dissertation, these

Fig. 6.3.2 Angular distributions of the differential cross sections for some of the levels excited in the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction.

Fig. 6.3.3 Angular distributions for some of the levels excited in the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction.

Fig. 6.3.4 The angular distribution of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ (6.48 MeV $7/2^-$) reaction is presented with an angular binning of only 0.46° , as opposed to the normal bin of 1.8° used in the rest of the figures.

are compared to the 800 MeV $^{12}\text{C}(\text{pd},)^{11}\text{C}$ data by plotting the distributions of the measured differential cross sections of the two experiments vs. momentum transfer $q = |\vec{k}_p - \vec{k}_d|$. The results of this comparison appear in Figs. 6.3.5, 6.3.6, and 6.3.7 for the excitation of the ^{11}C ground state, 2.00 MeV, and 6.48 MeV levels, respectively. It should be noted that in Figs. 6.3.5, 6.3.6, and 6.3.7, only every second or third cross section in the angular distributions measured by the French group has been plotted in order to make the figures more presentable. These figures show that, in general, good agreement exists in the region of overlap between the measurements performed at the separate facilities.

Shown in Fig. 6.3.8 is the 800 MeV $^{12}\text{C}(\text{p,d})^{11}\text{C}(\text{g.s.})$ angular distribution with the results of EFR DWBA calculations employing the KMT folded, and well-matched deuteron optical potentials of Tables 2.6.2 and 2.7.1. Both calculations describe the magnitude of the data very well, but the well-matched deuteron optical potential generates a better fit to the shape of the data. The KMT calculation exhibits too much structure. The spectroscopic factor used to normalize absolutely both calculations is 3.2, a value determined from the p-shell wavefunctions of Norton and Goldhammer (NG 70), (Nor 70). This value

Fig. 6.3.5 The differential cross sections of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction to the $3/2^-$ ground state of ^{11}C (open circles) are compared to those of the 700 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction to the same state (solid circles) (Bak+ 74) vs. momentum transfer q .

Fig. 6.3.6 Same as for Fig. 6.3.5 only for the 2.0 MeV $1/2^-$ state of ^{11}C .

Fig. 6.3.7 Same as for Fig. 6.3.6 only for the 6.48 MeV $7/2^-$ state of ^{11}C .

Fig. 6.3.8 The 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}(\text{g.s.})$ data are compared to EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials.

may be compared with the value obtained by Cohen and Kurath (CK 67) of 2.80.

The origin of the 2.0 MeV wide region of enhanced excitation near 13.2 MeV in ^{11}C was also investigated. The importance of TS contributions to the excitations of some of the levels in ^{11}C has already been discussed. In the extreme case where the TS mechanism dominates the excitations of levels in ^{11}C , a picture such as that depicted in Fig. 6.3.9 emerges, where all the levels except the ground state are pictured as being core-excited levels based on the first 2^+ , 3^- , or 4^+ states of ^{12}C . Within the framework of this extreme (TS) situation, it is natural to ascribe the parentage of the region in ^{11}C near 13.2 MeV to the 14.1 MeV 4^+ level in ^{12}C . If this inelastic excitation is followed by pickup of a $p_{3/2}$ valence neutron, then levels of angular momentum and parity $[4^+ \times (p_{3/2})^{-1}] = 11/2^-, 9/2^-, 7/2^-,$ and $5/2^-$ are to be expected. The inelastic excitation of the 14.1 MeV 4^+ level in ^{12}C has been shown to be very strong in 800 MeV $^{12}\text{C}(p,p')$ work (Bla 78), (Bla+77), especially for increasing momentum transfer. Additionally, the angular distribution of the inelastic scattering to this level cannot be properly accounted for unless the TS pathway $0^+ \rightarrow 2^+ \rightarrow 4^+$ is included along with the SS pathway $0^+ \rightarrow 4^+$ (Bla 78),

Possible core excited configurations

Fig. 6.3.9 Parentage of levels in ^{11}C populated by the $^{12}\text{C}(p, d)^{11}\text{C}$ reaction, in the (extreme) weak coupling model.

(Ray+ 78). Consequently, a zero range three-step calculation was performed to attempt to describe the 13.2 MeV bump in $^{12}\text{C}(p,d)^{11}\text{C}$. The pathways, deformation parameters β_2 and β_4 (Bla 78), and pickup spectroscopic amplitudes are shown in Fig. 6.3.10. The spectroscopic amplitudes are computed from the weak coupling model, where the strength to each member of the $[4^+ \times (p3/2)^{-1}]$ multiplet scales as $2j_f + 1$, subject to the normalization that the total pickup strength is equal to the sum rule for p3/2 pickup, i.e. 4. Then the spectroscopic factor is given by

$$S = \frac{n}{2j + 1} \frac{2J + 1}{2J_c + 1} ,$$

where n is the number of particles in the shell. In this example, J_c is 4, j is 3/2, and J is 11/2, 9/2, 7/2 or 5/2. The zero range strength parameter used in the CCBA calculation was $D_0 = 80 \text{ MeV} - \text{fm}^{3/2}$. The results of the calculation to the $7/2^-$ member of the multiplet, presented in Fig. 6.3.11, reveals that this mechanism is unable to account for the observed strength at 13.2 MeV in ^{11}C , if only this one expected path is included.

A second calculation was performed to explore the possibility that the 13.2 MeV bump represents a signature of inner shell (s1/2) pickup. The EFR DWBA

Multiple step pathways to the 13.2 MeV region of ^{11}C

Fig. 6.3.10 Two-step pathways, deformations, and spectroscopic amplitudes for p_{3/2} (solid line) and p_{1/2} (dashed line) neutron pickup to the 7/2⁻ member of a (suspected) multiplet located at 13.2 MeV in ^{11}C , populated by the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction.

Fig. 6.3.11 Shown is the angular distribution of the 13.2 MeV bump populated in the 800 MeV $^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ reaction, and the results of two calculations. The solid line shows the results of a three step ZR CCBA calculation ($D_0 = 80 \text{ MeV} \cdot \text{fm}^{3/2}$) utilizing the pathways of Fig. 6.3.10, which severely underpredicts the data. The dashed curve is the result of an EFR DWBA calculation of inner shell ($0s_{1/2}$) neutron pickup. The good agreement of this calculation with the data (see discussion in text) is suspected to be fortuitous.

calculation assumed the picked up neutron was in the $0s_{1/2}$ shell when picked up from the ^{12}C ground state. A spectroscopic factor of 2.0 (sum rule) was used to normalize the prediction. The results, presented in Fig. 6.3.11, reveal that the magnitude of the observed strength is roughly accounted for. However, the inner shell strength has been shown (Tyr+ 66) to lie higher in excitation energy, near 19.5 MeV, and to be much broader (9.2 MeV) than the width of 2 MeV observed in the 800 MeV $^{12}\text{C}(p,d)$ reaction. In addition, the same calculation performed at 121 MeV predicts a strength of about one-tenth the strength to the ground state. In fact, no such strength is seen in this region at any other lower bombarding energy (Bac+ 69), (Lee+ 68). Consequently the identification of the 13.2 MeV strength in ^{11}C with inner shell pickup becomes untenable, in spite of the rough agreement of the prediction with the present data. It is, of course, possible that some as yet unaccounted for mechanism suppresses the inner shell pickup at lower bombarding energies, and changes its location and width slightly from what is found in $(p, 2p)$ measurements.

The 13.2 MeV strength in ^{11}C is more likely due to unresolved high spin states. Such states have been observed in ^{11}B , the mirror nucleus of ^{11}C , at

excitation energies of 13.12 MeV ($9/2^-$), 13.16 MeV ($5/2^+$, $7/2^+$), and 14.02 MeV ($11/2^+$) in the $^{10}\text{B} + n$ system (HNL 73). In fact, levels are predicted at 13.25 MeV ($9/2^+$) and 13.52 MeV ($7/2^+$) in the tabulations of Norton (Nor 70). The wavefunctions predicted for these levels are predominantly $L = 4$, $S = 1/2$, and $T = 1/2$ (Nor 70) suggesting that their parentage is indeed in the (14.1 MeV) 4^+ of ^{12}C , followed by p shell neutron pickup.

6.4 $^{16}\text{O}(p,d)^{15}\text{O}$

Sample spectra of the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction are shown in Fig. 6.4.1, for laboratory scattering angles of 5.2° and 17.1° . The linear energy calibration of the $^{16}\text{O}(p,d)^{15}\text{O}$ spectra was accomplished utilizing the ground state and 6.176 MeV levels of ^{15}O in spectra collected at laboratory angles of 5° , 10° , 12.5° , 15° , and 17° . The results were averaged, and are presented in Table 6.4.1. Remarkable in the spectra presented in Fig. 6.4.1 are several features. First, the strongest states populated are at large (>10 MeV) excitation energies, where little strength has been seen at lower energies (Roo+ 75), (Ger+ 75). The smoothness of the angular distributions (presented in Figs. 6.4.2 and 6.4.3) for these states relative to the ground state may suggest that two step mechanisms contribute to the population of these levels.

Fig. 6.4.1 Spectra of the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction at 5° and 17°.

Table 6.4.1. 800 MeV $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$
Calibration

Peak #	Excitation Energy ^{a/}	Standard Deviation ^{b/}	Excitation Energy ^{c/}	J ^{πc/}
1	0.000 $\pm$.1	--	0.000	1/2 ⁻
2	5.219 $\pm$.1	.048	5.241	5/2 ⁺
3	6.176 $\pm$.1	--	6.176	3/2 ⁻
4	7.203 $\pm$.1	.033	7.276	7/2 ⁺
5*	8.841 $\pm$.15	.158		
6	10.421 $\pm$.15	.10		
7**	10.869 $\pm$.15	.067		
8	12.210 $\pm$.15	.056		
9	13.593 $\pm$.15	.140		
10	19.02 $\pm$.20	--		
11	21.11 $\pm$.20	--		

^{a/} Present work: linear calibration based upon ground state and 6.176 MeV levels at 5 angles.

^{b/} Standard deviation based upon excitation energies calculated in the present work for laboratory scattering angles of 5°, 10°, 12.5°, 15°, and 17°.

^{c/} Levels identified from the compilations of (Ajz 76).

*15° point excluded.

**5° and 10° point excluded.

Fig. 6.4.2 Angular distributions of some of the levels excited in the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction.

Fig. 6.4.3 Angular distributions of some of the levels excited in the 800 MeV $^{16}\text{O}(\text{p},\text{d})^{15}\text{O}$ reaction.

Also visible in the spectra are two broad regions of enhanced excitation at 19 MeV and 21 MeV. The nature of these newly observed levels is not understood at present.

The excitation of the $5/2^+$ and $7/2^+$ levels at 5.22 MeV and 7.20 MeV has interesting consequences. At lower bombarding energies, these levels are populated only very weakly relative to the very pure single neutron hole states at 0.00 MeV ($p_{1/2}$ hole) and 6.18 MeV ($p_{3/2}$ hole) (Roo+ 75). Several workers recently published (Bac+ 77) a study of the population of the $(p_{1/2})^{-1}$ and $(p_{3/2})^{-1}$ single neutron hole states in ^{15}O via the 66 MeV $^{16}\text{O}(\pi^+, p)^{15}\text{O}$ reaction. Strength observed in a 3 MeV wide region centered at 6.18 MeV was ascribed to the $(p_{3/2})^{-1}$ hole state, on the basis of the low energy observations that the $5/2^+$ and $7/2^+$ states are only weakly populated. To their surprise, they found that the ratio of the $(p_{3/2})^{-1}$ strength to that of the $(p_{1/2})^{-1}$ varied between 6 and 13. Since these levels are such pure hole states, a ratio of 2 is expected in the one nucleon model, simply from the occupation numbers of the corresponding shell-model orbitals. Indeed, this ratio of two has always been observed at lower bombarding energies. In an attempt to explain the anomalous ratios of their experiment, it was proposed that:

(a) the $p_{1/2}$ and $p_{3/2}$ neutron wave functions behaved differently in the previously unaccessed large momentum regime, and/or (b) that some characteristic of the (π^+, p) reaction mechanism was responsible for the anomaly.

The 800 MeV $^{16}\text{O}(p, d)^{15}\text{O}$ spectra solve the mystery of the seemingly anomalous ratio of the $(p_{3/2})^{-1}$ to $(p_{1/2})^{-1}$ hole strength, because, as will be shown in Chapter VIII, the relative intensities of final states populated in the (π^+, p) reaction are almost identical with the relative intensities observed in the (p, d) reaction at the same momentum transfer. Under the assumption that this similarity holds the 800 MeV $^{16}\text{O}(p, d)^{15}\text{O}$ spectra may be viewed as a high resolution glimpse of the $^{16}\text{O}(\pi^+, p)^{15}\text{O}$ reaction. It is immediately observed from Fig. 6.4.1 that the positive parity levels one MeV on either side of the $(p_{3/2})^{-1}$ hole state are as strongly excited as the $(p_{3/2})^{-1}$ hole state itself.

The sum of the strength to the $5/2^+$, $3/2^-$, and $7/2^+$ levels (all located within the experimental resolution of (Bac+ 77)) is roughly a factor of 6-13 times the strength to the ground state, which verifies the observations of (Bac+ 77) and simultaneously explains their "anomalous" values for the $(p_{3/2})^{-1}$ to $(p_{1/2})^{-1}$ ratio. Indeed, the ratio of the 6.18 MeV $3/2^-$ level

to the $1/2^-$ ground state in the 800 MeV $^{16}\text{O}(p,d)$ data is roughly the expected ratio of two.

Identification of the higher lying levels in ^{15}O populated in the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction is difficult due to the high density of states (Ajz 76). The weakly excited 8.84 MeV level is closest to a known level at 8.74 MeV with $J^\pi = 1/2^+$. Many states lie within the experimental resolution of the doublet observed in the 800 MeV (p,d) work with excitation energies of 10.42 and 10.87 MeV. Between 10.3 MeV and 10.94 MeV are six levels with assigned spins and parities of $5/2^\pm$, $3/2^\pm$, $7/2^+$ and $1/2^+$. Two states with angular momentum 5/2 are tabulated near the strongly excited 12.21 MeV level seen at 800 MeV. Between 13.4 MeV and 13.8 MeV are tabulated five levels with $J^\pi = 3/2^\pm$ and $5/2^+$, any of which could be contributing to the strongest level excited in the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction, the 13.59 MeV state. Two broad levels are tabulated at 19.03 MeV (1 MeV width) and 20.5 and 22.0 MeV (2 MeV width) which could be associated with the 19.0 and 21.1 MeV (broad) levels observed in the 800 MeV $^{16}\text{O}(p,d)$ reaction.

The angular distributions of most of the levels populated in ^{15}O by the 800 MeV (p,d) reaction are presented in Figs. 6.4.2 and 6.4.3. It is apparent from these figures that no information regarding the

angular momentum or parity of the levels is discernible from the shapes of the angular distributions, which are smooth and structureless. This is in contrast to the situation for the (p,d) reaction at lower bombarding energies, where the location of minima in the angular distributions has long been a useful tool for extracting the angular momentum quantum numbers of levels in the residual nucleus.

There is a slight difference in the shapes of the angular distributions between known pure hole states such as the $(p1/2)^{-1}$ ground state, and levels which probably involve collective TS processes, such as the 7.2 MeV level or more distinctly, the strongest levels (probably of high spin) at 12.2 MeV and 13.6 MeV. This difference manifests itself as a minimum at about 12° for the hole states, which is completely absent for the states above 6.2 MeV in excitation energy.

Shown in Figs. 6.4.4 and 6.4.5 are the results of EFR DWBA calculations for the excitation of the ground state $1/2^-$ (Fig. 6.4.4) and 6.18 MeV $3/2^-$ (Fig. 6.4.5) levels of ^{15}O . The spectroscopic factors used to normalize these calculations were 2.0 and 4.0, respectively, which represent the spectroscopic sum rule limit for $p1/2$ and $p3/2$ pickup. The solid lines are the predictions employing the KMT folded deuteron

Fig. 6.4.4 The angular distribution of the $1/2^-$ ^{15}O ground state from the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction is presented along with EFR DWBA calculations employing KMT (solid curve) and phenomenological (dashed curve) (Bru 78) deuteron optical potentials.

Fig. 6.4.5 Data from the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ (6.48 MeV $3/2^-$) reaction are presented with the results of EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials.

optical potentials of Table 2.6.2. In Fig. 6.4.4, the dashed line demonstrates the sensitivity of the calculations to the folding prescription of Sect. 2.6. This dashed line was generated using a phenomenological deuteron optical potential with the following parameters: $V_R = 0.0$ MeV, $V_I = -75.85$ MeV, $r_I = 0.96$ fm, and $a_I = 0.537$ fm. This phenomenological potential was obtained from a fit employing the methods of Sect. 2.5 to 700 MeV $^{16}\text{O}(d,d)^{16}\text{O}$ data from Saturne (Bru 78), considered to be the only reliable deuteron elastic scattering data extant at energies greater than 200 MeV. The elastic scattering cross sections predicted by the two calculations are compared to the 700 MeV $^{16}\text{O}(d,d)^{16}\text{O}$ data in Fig. 6.4.6. As can be seen from Fig. 6.4.4, usage of the KMT folded deuteron optical potential results in improved agreement with the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}(\text{g.s.})$ data in both shape and magnitude.

Fig. 6.4.5 contrasts calculations using the KMT folded and the well-matched deuteron optical potentials of Tables 2.6.2 and 2.7.1, respectively, for the excitation of the 6.18 MeV $3/2^-$ level. Both calculations have roughly the correct magnitude, but oscillations are evident in the KMT calculation which are not reflected by the data.

Fig. 6.4.6 Saclay data (Bru 78) of the 700 MeV $^{16}\text{O}(d,d)^{16}\text{O}$ reaction are presented along with the elastic scattering predictions using a "best-fit" phenomenological (dashed line) (see Sect. 2.5), and a KMT (solid line) deuteron optical potential.

Two-step calculations were performed for the excitation of the $5/2^+$ and $7/2^+$ levels at 5.24 MeV and 7.28 MeV, employing the pathways, deformation parameter ($\beta_3 = -0.3$), and weak coupling spectroscopic amplitudes of Fig. 6.4.7. The CCBA results are shown in Fig. 6.4.8 for the $5/2^+$ and $7/2^+$ excitations. These calculations were normalized by requiring the predictions to the 3^- level reproduce the 800 MeV experimentally measured $^{16}\text{O}(p,p')^{16}\text{O}(6.13 \text{ MeV } 3^-)$ data at the first maximum, 27 mb/sr at $\theta_{\text{c.m.}} = 10.7^\circ$ (Ada+79). With this normalization, and a $D_0 = 80 \text{ MeV} \cdot \text{fm}^{3/2}$, the magnitude of the $7/2^+$ prediction accounts for the data fairly well. The $5/2^+$ data are slightly underpredicted, and both calculations exhibit nodes which are not present in the experimental data. This is especially the case for the $5/2^+$ prediction which underpredicts the $\theta_{\text{c.m.}} = 6^\circ$ data point by almost two orders of magnitude because of a node in the calculation. However, the agreement of the CCBA calculations with the $5/2^+$ and $7/2^+$ angular distributions is good enough to support the hypothesis that TS is mainly responsible for the excitation of these two positive parity levels.

$^{16}\text{O}(p, d)^{15}\text{O} \left(\frac{5}{2}^+, \frac{7}{2}^+ \right)$
 two step pathways and spect. amplitudes
 $p_{1/2}$ pickup

Fig. 6.4.7 Two step pathways, spectroscopic amplitudes, and deformation parameter for $^{16}\text{O}(p, d)^{15}\text{O}$ to the $5/2^+$ and $7/2^+$ levels at 5.2 MeV and 7.2 MeV.

Fig. 6.4.8 Angular distributions of the 800 MeV $^{16}\text{O}(p,d)^{15}\text{O}$ reaction to the $5/2^+$ (A) and $7/2^+$ (B) levels. Compared to the data are CCBA calculations employing the pathways of Fig. 6.4.7. The calculations have been scaled to reproduce the observed $^{16}\text{O}(p,p)^{16}\text{O}$ (6.18 MeV 3^-) strength (Ada+79), and employ a $D_0 = 80 \text{ MeV} \cdot \text{fm}^{3/2}$.

6.5 $^{25}\text{Mg}(p,d)^{24}\text{Mg}$

Many discrete levels are observed in the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction. Sample spectra collected at lab angles of 5.2° and 12.5° are presented in Fig. 6.5.1. Unambiguous identification of levels is possible at excitation energies corresponding to the 0^+ ground state, 1.37 MeV 2^+ , 5.24 MeV 3^+ , and 6.01 MeV 4^+ states. A level observed at 4.12 MeV probably includes contributions from both the 4.12 MeV 4^+ and 4.24 MeV 2^+ levels.

Calibration of the $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ spectra was simplified due to the identification of these levels. The separation of the 1.37 MeV 2^+ and 6.01 MeV 4^+ levels, both strongly excited, provided the basis for the linear calibration. The results of this calibration at laboratory angles of 5° , 10° , and 12° were averaged, and the results are presented in Table 6.5.1. The calibration constants (at a given angle) were considered reliable enough to calibrate spectra from other targets. Results of this technique compared to two others were presented in Table 6.3.1 for the case of $^{12}\text{C}(p,d)^{11}\text{C}$.

Angular distributions for most of the levels populated in ^{24}Mg are presented in Figs. 6.5.2, 6.5.3, and 6.5.4. Once again the lack of J^π information in the 800 MeV angular distributions is clearly

Fig. 6.5.1 Spectra of the 800 MeV $^{25}\text{Mg(p,d)}^{24}\text{Mg}$ reaction at 5.2° and 12.5°.

Table 6.5.1. 800 MeV $^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}$
Calibration

Peak #	Excitation Energy ^a / MeV	Standard Deviation ^b / MeV	Excitation Energy ^c / MeV	J ^π
1	0.052 ± .10	.127	0.00	0 ⁺
2	1.370 ± .10	--	1.37	2 ⁺
3	4.124 ± .10	.022		
4	5.266 ± .10	.029	5.24	3 ⁺
5	6.01 ± .10	--	6.01	4 ⁺
6	7.664 ± .10	.006		
7	8.361 ± .10	.054		
8	9.441 ± .10	.044		
9	10.12 ± .15	.040		
10	10.63 ± .15	.055		
11	11.05 ± .15	.062		
12	11.99 ± .18	.125		
13	12.39 ± .18	.106		
14	13.30 ± .18	.084		
15	14.13 ± .20	.118		
16*	15.17 ± .20	.153		
17*	15.68 ± .20	.148		
18*	16.56 ± .20	.163		

^a/ Present work: linear calibration based upon 1.37 MeV and 6.01 MeV levels at 3 angles.

^b/ Standard deviations based upon excitation energies calculated in the present work for laboratory scattering angles of 5°, 10°, and 12°.

^c/ Levels identified from the compilations of (EV 78).

*12° point excluded.

Fig. 6.5.2 Angular distributions for some of the levels populated in the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction.

Fig. 6.5.3 Angular distributions for some of the levels populated in the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction.

Fig. 6.5.4 Angular distributions for some of the levels populated in the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction.

apparent. Nothing except differences in overall magnitude distinguishes one angular distribution from the next.

Many of the levels populated in the $^{25}\text{Mg}(p,d)$ ^{24}Mg reaction are near to known (EV 78) high spin states. The level density of ^{24}Mg is such that the 6.01 MeV level observed in the 800 MeV $^{25}\text{Mg}(p,d)$ ^{24}Mg reaction can be positively identified as a 4^+ $T = 0$ level. More 4^+ strength is known to lie near 9.5 MeV, where a state is strongly populated (at 9.44 MeV) in the 800 MeV (p,d) reaction. Negative parity levels at 7.62 MeV (3^-), 10.03 MeV (5^-) and 12.45 MeV (7^-) are located close to levels seen in the 800 MeV (p,d) reaction at excitation energies of 7.66 MeV, 10.12 MeV, and 12.39 MeV. Although the level density of ^{24}Mg and poor experimental resolution of the 800 MeV (p,d) data do not permit the unambiguous association of the observed levels with levels of high spin, the correlation is suggestive of a striking preference for the population of high spin levels.

The 0^+ ground state of ^{24}Mg is only very weakly excited (100 nb/sr maximum measured cross section at 800 MeV). However, as shown in Fig. 6.5.5, this weak strength is as predicted by the EFR DWBA. The spectroscopic factor used was 0.33, which was obtained from the Nilsson model (Elb+ 75), and compares well

Fig. 6.5.5 Data for the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ reaction to the 0^+ ^{24}Mg ground state are presented with the results of an EFR DWBA calculation employing a KMT deuteron optical potential.

with the measured value of 0.270 in the 18 MeV $^{25}\text{Mg}(h,\alpha)^{24}\text{Mg}$ reaction (Elb+ 75). For the 800 MeV $^{25}\text{Mg}(p,d)^{24}\text{Mg}$ calculation, the KMT deuteron optical potential and neutron bound state parameters used were those appropriate for ^{28}Si .

No calculations are presented for other known levels because of the complications introduced by the multiple angular momentum transfers possible from the $d_{5/2}$ ground state of ^{25}Mg to levels other than 0^+ (^{24}Mg ground state).

6.6 $^{28}\text{Si}(p,d)^{27}\text{Si}$

Many previously unobserved levels are excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction. Sample spectra are exhibited in Fig. 6.6.1 at laboratory angles of 5.2° and 12.5° . Little is known of ^{27}Si above 7 MeV excitation energy, yet the greatest strength in the 800 MeV (p,d) reaction feeds discrete sharp levels at excitation energies greater than 7 MeV. In particular, levels at 7.18 MeV, 8.32 MeV, and 11.65 MeV are very strongly excited at 800 MeV, although these levels have not been observed at other (lower) bombarding energies (Bac+ 69), (Koz 68), (Ger+ 75).

The linear calibration of the ^{27}Si spectra was accomplished in the following way. At each of four separate angles (5° , 10° , 12° , and 17°), the ^{27}Si ground state was used to predict the excitation

Fig. 6.6.1 Sample spectra of the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction at 5.2° and 12.5° .

energies of other ^{27}Si levels based upon the separation of the ^{11}C ground state and 6.48 MeV levels populated in $^{12}\text{C}(\text{p},\text{d})^{11}\text{C}$ at that same angle. The results of the four calibrations were averaged, and the results along with the standard deviation of the averaging are presented in Table 6.6.1. The (generous) errors in excitation energies are based on the standard deviations and on the quality of peak fits used to obtain centroids for the calibration. The resolution of the 800 MeV $^{28}\text{Si}(\text{p},\text{d})^{27}\text{Si}$ data, and density of states in ^{27}Si is such that only the ground state may be positively identified.

Remarkable in the ^{27}Si spectra is the excitation of several strongly excited levels (relative to the ground state) at excitation energies greater than 7 MeV. The highest known $T = 1/2$ level in ^{27}Si quoted in the compilations (EV 78) is only 6.34 MeV. Recent $^{28}\text{Si}(\text{p},\text{d})^{27}\text{Si}$ work carried out at $T_p = 121$ MeV at Indiana, however, (Mil 79) reveals the existence of levels within the energy errors quoted in Table 6.6.1 for the 7.18 MeV, 8.32 MeV, 9.59 MeV, and 11.65 MeV states. Additionally, the IUCF data reveal that the angular distributions for these levels are anomalously isotropic (relative to the angular distributions of states at lower excitation), suggesting that these levels are of high spin. This suggestion

Table 6.6.1. 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$
Calibration

Peak #	Excitation Energy ^{a/}	Standard Deviation ^{b/}	Excitation Energy ^{c/}	J ^{π}
1	0.000 $\pm$.10	--	0.000	5/2 ⁺
2	0.797 $\pm$.10	.032		
3	2.460 $\pm$.25	.199		
4	2.888 $\pm$.10	.081		
5	4.201 $\pm$.15	.119		
6	4.574 $\pm$.20	.155		
7	5.372 $\pm$.15	.077		
8*	6.566 $\pm$.30	.186		
9	7.183 $\pm$.15	.104		
10	8.324 $\pm$.15	.138		
11	9.590 $\pm$.18	.139		
12	11.65 $\pm$.20	.173		
13**	14.79 $\pm$.25	.209		
14**	15.283 $\pm$.25	.204		
15**	15.735 $\pm$.25	.222		
16**	16.250 $\pm$.25	.246		

^{a/} Present work: linear calibration based upon the ground state at 4 angles, using the calibration constant from $^{12}\text{C}(p,d)^{11}\text{C}$ (0.00 MeV, 6.48 MeV) at that angle.

^{b/} Standard deviations based upon excitation energies in the present work for laboratory scattering angles of 5°, 10°, 12°, and 17°.

^{c/} Levels identified from the compilations of (EV 78).

*12° point excluded.

**5° point excluded.

is again in agreement with the observed preferential population of high spin states at 800 MeV (see Sect. 2.10).

In the 15 MeV region of excitation in ^{27}Si is a bump reminiscent of the broad levels observed in 800 MeV (p,d) to ^{11}C and ^{15}O . The bump in ^{27}Si is, however, clearly fragmented into at least four separate levels (each separated by about 0.5 MeV). The clear identification of four separate levels in this region is possible at almost every angle due to the usage of an especially thin target for most of the $^{28}\text{Si}(p,d)^{27}\text{Si}$ data collection.

In addition to the $5/2^+$ ground state, many other levels in ^{27}Si are excited which have been seen before. The 0.80 MeV level probably includes contributions from both the $1/2^+$ and $3/2^+$ levels at 0.78 MeV and 0.96 MeV, respectively. The energy levels of ^{27}Si are too dense to permit much more to be said about the other observed levels below 6.34 MeV, and nothing is known of ^{27}Si above this excitation. However, known states of high spin are located at 2.91 MeV ($9/2^+$) and 4.45 ($11/2^+$) in ^{27}Si (EV 78), tantalizingly close to the levels observed at $2.89 \pm .1$ MeV, and $4.20 \pm .15$, $4.57 \pm .20$ MeV in this work.

The nucleus ^{27}Al is the mirror nucleus of ^{27}Si , and may be expected to have a level structure very

similar to ^{27}Si . Much more is known of ^{27}Al in the high excitation region (EV 78). A $9/2^+$ level is known at 5.67 MeV in ^{27}Al which could be associated with the 5.37 MeV level observed in ^{27}Si . High spin states of $J^\pi = 9/2^+$ and $9/2^-$ are located in ^{27}Al at excitation energies of 7.17 and 7.23 MeV, well within the error assigned to the 7.18 MeV level strongly excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction. A $3/2^-$ level at 8.60 MeV in ^{27}Al is close to the observed 8.32 MeV state in ^{27}Si observed in this work. Three levels are tabulated within 100 KeV of the 9.59 state observed in ^{27}Si , all of which have tentative spin assignments of $1/2^-$ and $3/2^-$. No strength above this excitation energy has been reported.

A fascinating correlation also exists between the strongest levels observed in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction, and the levels of high spin and negative parity in ^{28}Si . This correlation is shown in Fig. 6.6.2, where all the negative parity levels of ^{28}Si below 12 MeV excitation energy are shown (plus a 6^- $T = 1$ level at 14.36 MeV) along with the strongest levels observed in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction. The 3^- , 4^- , 5^- , and 6^- levels in ^{28}Si shown line up so nicely with the strongest levels observed in ^{27}Si that one is tempted to submit Fig. 6.6.2 as circumstantial evidence for weak

Fig. 6.6.2 Correlation between the negative parity levels of ^{28}Si , and the strongest states excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction, suggesting the importance of weak coupling in the 800 MeV (p,d) reaction. Several negative parity levels between 12 MeV and 14 MeV (in ^{28}Si) have been omitted for clarity.

coupling in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction.

Angular distributions of all the levels observed are presented in Figs. 6.6.3, 6.6.4, and 6.6.5. Very little structure is seen in any of the angular distributions, although the ground state and 0.80 MeV levels have a hint of a minimum near 12° (lab). No unusual variations are observed for any of the levels, including the four fragments of the "bump" observed near 15 MeV.

Two EFR DWBA calculations were performed for the only level of known spin and parity populated in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction. Fig. 6.6.6 presents the results of calculations of the ^{27}Si ground state utilizing the KMT and well-matched deuteron optical potentials of Tables 2.6.2 and 2.7.1. Both calculations (especially the KMT) account for the magnitude of the data well, although too much structure is again evident in the KMT calculation. The spectroscopic factor used to normalize these EFR DWBA calculations was 3.4, and comes from a rotational model calculation (Deh 72) using a $\beta = +0.30$ or $\beta = -0.27$. The measured value is $C^2S = 3.45$ (Koz 68).

The EFR DWBA accounts for the only transition (to the ^{27}Si ground state) about which enough is known to trust the calculation. However, the excellent agreement with the data achieved for this case

Fig. 6.6.3 Angular distributions for some of the levels excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction.

Fig. 6.6.4 Angular distributions for some of the levels excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction.

Fig. 6.6.5 Angular distributions for some of the levels excited in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction.

Fig. 6.6.6 Angular distribution for excitation of the ^{27}Si $5/2^+$ ground state, excited with the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ reaction. Compared to the data are EFR DWBA calculations utilizing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials.

suggests that the intriguing spectroscopy evident in the 800 MeV $^{28}\text{Si}(p,d)^{27}\text{Si}$ data might be unraveled if more J^π and C^2S information were known or suspected. More investigations of ^{27}Si should be performed with reactions which are known to populate high spin states, and which yield more J^π information. Such reactions are (p,α) , (α,p) , and (α,d) at low (~ 50 MeV) bombarding energies (Smi 76).

6.7 $^{40}\text{Ca}(p,d)^{39}\text{Ca}$

The highest mass target used in the study of the systematic A-dependence of the 800 MeV (p,d) reaction was ^{40}Ca . Sample spectra taken at laboratory angles of 5.2° and 12.5° are presented in Fig. 6.7.1.

Calibration of the $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ spectra was complicated due to the use of a thick target to offset the diminished cross section, and the high level density of ^{39}Ca . Only the ground state could be positively identified. Excitation energies for the remaining levels were based upon the ^{39}Ca ground state location and the (linear) calibration constant from $^{25}\text{Mg}(p,d)^{24}\text{Mg}(1.37 \text{ MeV}, 6.01 \text{ MeV})$. The results of this technique at laboratory angles of 10° , 12° , and 17° were averaged, and the results are presented in Table 6.7.1. The effect of the thick target is reflected in the larger assigned errors in these excitation energies.

Fig. 6.7.1 Spectra of the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction at 5.2° and 12.5° .

Table 6.7.1. 800 MeV $^{40}\text{Ca}(\text{p},\text{d})^{39}\text{Ca}$
Calibration

Peak #	Excitation Energy ^{a/}	Standard Deviation ^{b/}	Excitation Energy ^{c/}	J ^π
1	0.000 ± .10	--	0.000	3/2 ⁺
2	2.678 ± .20	.141		
3	3.792 ± .20	.147		
4	5.400 ± .20	.152		
5	6.391 ± .25	.203		
6	7.260 ± .25	.162		
7	7.775 ± .25	.169		
8	10.92 ± .30	.184		
9	17.83 ± .50	--		

^{a/} Present work: linear calibration based upon the ground state at three angles, using the calibration constant from $^{25}\text{Mg}(\text{p},\text{d})^{24}\text{Mg}$ (1.37 MeV, 6.0. MeV) at that angle.

^{b/} Standard deviations based upon excitation energies in the present work for laboratory scattering angles of 10°, 12°, and 17°.

^{c/} Levels identified from the compilations of (EV 78).

Most of the levels seen in the ^{39}Ca spectra undoubtedly contain contributions from at least two levels. The widths and shapes of the peaks in the ^{39}Ca spectra differ from the cleanly separated $3/2^+$ ground state shape, indicating multiple contributions to these levels. Levels of high spin in ^{39}Ca are tabulated at energies of 2.80 MeV ($7/2$), 3.64 MeV ($9/2^-$ ($3/2^-$, $5/2^-$)), and 5.49 MeV ($3/2^+$, $5/2^+$) which are within the vicinity of the 2.68 MeV, 3.79 MeV, and 5.40 MeV levels observed in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction. In the mirror nucleus ^{39}K , levels with spins as high as $19/2^-$ are tabulated at excitation energies up to 8 MeV, but the density of states is such that no correlations can be made with the present data.

The most remarkable feature in the ^{39}Ca spectra is the existence of two very broad (~ 2 MeV) bumps high in excitation energy at 10.92 MeV and 17.8 MeV. Most of the (discrete) strength in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction is to these levels, whose nature is unknown. No discrete strength has been observed in this region previously (Koz 68), (Roo+ 75).

The angular distributions of the levels observed in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction are shown in Figs. 6.7.2 and 6.7.3. Again, the data are featureless in shape, although the magnitude of the

Fig. 6.7.2 Angular distributions for some of the levels excited in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction.

Fig. 6.7.3 Angular distributions for some of the levels excited in the 800 MeV $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction.

cross-sections is smaller than for the other targets studied.

The results of EFR DWBA calculations to the $^{39}\text{Ca } 3/2^+$ ground state are presented in Fig. 6.7.4. The usage of a well-matched deuteron optical potential is seen to result in a superior fit to the data. The KMT calculation overpredicts the data by roughly a factor of five. The spectroscopic factor used to normalize both calculations was 3.0 (Kun 79). This compares with a measured value of 3.70 (Koz 68) obtained from the $^{40}\text{Ca}(p,d)^{39}\text{Ca}$ reaction at 33 MeV.

Fig. 6.7.4 Data for the excitation of the $3/2^+$ ^{39}Ca ground state, populated in the 800 MeV $^{40}\text{Ca}(p, d)^{39}\text{Ca}$ reaction, are presented with EFR DWBA calculations employing KMT (solid curve) and well-matched (dashed curve) deuteron optical potentials.

CHAPTER VII

PRESENTATION OF THE (π^+, p)

DATA AND CALCULATIONS

Very few data on the (π^+, p) reaction have been collected to date. The data that have been taken suffer from extremely poor energy resolution and insufficient statistics (Ama+ 74), (Ama+ 78), (Bac+ 77). More plentiful data have been taken on the (p, π^+) reaction, however (Hoi 79), (Ben+ 78), (Bor+ 76), (Hoi 79b), which is related to the (π^+, p) reaction by detailed balance (see Chapter III). The (π^-, n) reaction has been studied on ^3He and ^4He in the region of the (3,3) resonance (Kal+ 78), but otherwise the 90 MeV and 180 MeV (π^+, p) reactions span a region (corresponding via detailed balance to the region $245 \text{ MeV} < T_p < 350 \text{ MeV}$) where no previous data exist.

The linear energy calibration of the (π^+, p) data presented in this chapter was simplified due to the fact that the energy levels of the residual nuclei are so well known. In addition, the similarity of the (π^+, p) spectra to those of the 800 MeV (p, d) reaction shown in Chapter VI aided in the identification of levels. Since the final levels populated in

the (π^+, p) reaction are so well known (due to the above reasons) no detailed discussion is needed for the linear energy calibration of the (π^+, p) spectra.

The resolution (≈ 800 keV) obtained in the (π^+, p) spectra was sufficient to resolve unambiguously many discrete levels, and the spectra and angular distributions which follow represent the first high resolution measurement of the (π^+, p) reaction.

7.1 ${}^6\text{Li}(\pi^+, p){}^5\text{Li}$

The prominent feature in the spectrum of the ${}^6\text{Li}(\pi^+, p){}^5\text{Li}$ reaction is the background characteristic of all the (π^+, p) data, namely the contribution from pion absorption on two nucleons. Two nucleon absorption yields protons with a continuous distribution in energy from about 260 MeV down to almost zero, for 180 MeV pions incident on ${}^{12}\text{C}$. This is to be compared to protons of roughly 290 MeV which emerge from the 180 MeV ${}^{12}\text{C}(\pi^+, p){}^{11}\text{C}$ reaction to the ground state of ${}^{11}\text{C}$. In most cases, however, the discrete levels of interest are not seriously affected by the large two nucleon absorption background.

This is especially the case for the ${}^6\text{Li}(\pi^+, p){}^5\text{Li}$ reaction, as demonstrated in Fig. 7.1.1, which shows a 20° spectrum at a pion bombarding energy of 175 MeV. Inset in this figure is the region near the ${}^5\text{Li } 3/2^-$ ground state, which has a natural width of ~ 1.5 MeV

Fig. 7.1.1 Sample spectrum of the 175 MeV ${}^6\text{Li}(\pi^+, p){}^5\text{Li}$ reaction at $\theta_L = 20^\circ$.

(Ajz 79). The first excited state of ${}^5\text{Li}$ is a 5 MeV wide state ($J^\pi = 1/2^-$) located between 5 MeV and 10 MeV of excitation. Counts are seen in the ${}^6\text{Li}(\pi^+, p){}^5\text{Li}$ spectra in this region, but it is unclear whether or not they are associated with this very broad level. The only other states reported in the literature for ${}^5\text{Li}$ are at excitation energies of 16.6 MeV and higher, where the two nucleon background is rising. It should be noted that the apparent peak in the $\pi\text{NN} \rightarrow \text{Np}$ background is an artifact of the decreasing focal plane efficiency near its edges. The actual $\pi\text{NN} - \text{Np}$ background on ${}^6\text{Li}$ at $T_\pi = 175$ MeV (studied by lowering the spectrometer fields in successive 50 MeV steps) begins rising near $T_p = 270$ MeV, reaches a maximum near $T_p = 200$ MeV, and tapers off gradually for lower proton energies.

Angular distributions of the 75 MeV and 175 MeV ${}^6\text{Li}(\pi^+, p){}^5\text{Li}(\text{g.s.})$ reactions are presented in Fig.

7.1.2. The striking feature of these angular distributions is their energy dependence. The 75 MeV angular distribution exhibits a broad minimum near 90° , while the 175 MeV angular distribution is featureless. Also, the magnitude of the 175 MeV forward angle cross sections is only about five times less than the 75 MeV cross sections. This is in contrast to the predicted energy dependence of the ${}^{12}\text{C}(\pi^+, p){}^{11}\text{C}$ reaction, as

Fig. 7.1.2 Angular distributions of the 75 MeV and 175 MeV $^{6}\text{Li}(\pi^{+}, p)^{5}\text{Li}(\text{g.s.})$ reaction.

presented in Sect. 3.2. No specific DWBA predictions have been made in this thesis for the (π^+, p) reaction on ${}^6\text{Li}$.

7.2 ${}^7\text{Li}(\pi^+, p){}^6\text{Li}$

A sample spectrum of the 175 MeV ${}^7\text{Li}(\pi^+, p){}^6\text{Li}$ reaction, collected at 20° , is shown in Fig. 7.2.1. Again, the $\pi\text{NN} \rightarrow \text{Np}$ background is prominent in this spectrum, and the comments made in the previous section with regard to this background apply equally well here.

Inset in Fig. 7.2.1 is the ${}^6\text{Li}$ ground state region. The ${}^6\text{Li}$ 1^+ ground state is excited, as well as the 2.185 MeV 3^+ state. The spectrum is similar to the 800 MeV ${}^7\text{Li}(p, d){}^6\text{Li}$ spectra, presented in Sect. 6.1, in that the 3^+ level is excited more strongly than the 1^+ ground state (roughly a factor of 2.5:1 in the ${}^7\text{Li}(\pi^+, p){}^6\text{Li}$ reaction, and a factor of about 5.5:1 in the 800 MeV ${}^7\text{Li}(p, d){}^6\text{Li}$ reaction). As was mentioned in Sect. 6.1, low momentum transfer neutron pickup experiments find the 1^+ and 3^+ strengths more nearly the same.

The angular distributions of the ${}^7\text{Li}(\pi^+, p){}^6\text{Li}$ reaction to the 1^+ and 3^+ levels are presented in Fig. 7.2.2, for pion bombarding energies of 75 MeV (A) and 175 MeV (B). The shapes of the angular distributions of the 1^+ and 3^+ levels are identical at a given

Fig. 7.2.1 Sample spectrum of the 175 MeV ${}^7\text{Li}(\pi^+, p){}^6\text{Li}$ reaction at $\theta_L = 20^\circ$.

Fig. 7.2.2 Angular distributions for the excitation of the 1^+ ground state and 2.185 MeV 3^+ levels of ${}^6\text{Li}$, excited in the ${}^7\text{Li}(\pi^+, p){}^6\text{Li}$ reaction at 75 MeV (A) and 175 MeV (B).

incident pion energy, however, the 1975 MeV angular distributions drop more uniformly than the 75 MeV angular distributions, which show a broad minimum near 100°_{cm} . The energy dependence in the magnitude of the differential cross sections at 75 MeV and 175 MeV remains puzzling, in that the observed forward angle cross sections are nearly of the same magnitude, in contrast to the predictions of the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ energy dependence presented in Sect. 3.2.

Calculations are compared to the data in Fig. 7.2.3. At both bombarding energies the calculations underpredict the data, but have approximately the correct shape. The 75 MeV calculations (A) are too low by a factor of about 20 at forward angles. The 175 MeV predictions underpredict the data by a factor of about 500. Clearly, a serious normalization problem exists in the calculations.

More interesting than the overall normalization problem is the predicted energy dependence. It was shown in Sect. 3.2 that a similar energy dependence was predicted for the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction. Moreover, it was shown that if the low energy pion optical potential was substituted for the high energy pion optical potential, then the high energy (π^+, p) calculations are of nearly the same shape and magnitude as the low energy calculations. The data show very little energy

Fig. 7.2.3 Shown are the 75 MeV (A) and 175 MeV (B) angular distributions of the ${}^7\text{Li}(\pi^+, p){}^6\text{Li}(1^+, 3^+)$ reaction, with the results of calculations described in Chapter III for both the 1^+ (solid curve) and 3^+ (dashed curve) levels.

dependence in the magnitude of the differential cross sections, possibly a manifestation of the fact that the pion propagates considerably off shell ($E^2 \neq p^2 + m^2$) in the (π^+, p) reaction. In other words, the pion optical potential which describes the elastic scattering of pions on the nucleus is inappropriate for the description of the (π^+, p) reaction.

Another interesting feature of the calculations (which survives in spite of the overall normalization problem) is the relative strengths of the 3^+ and 1^+ predictions at both energies. Just as was found to be the case for the 800 MeV ${}^7\text{Li}(p, d){}^6\text{Li}$ calculations, the magnitude of the 3^+ calculation is less than that of the 1^+ calculation. Although this correctly describes the relative intensity of the 1^+ and 3^+ levels observed in low momentum transfer neutron pickup experiments, it does not describe the enhanced excitation of the 3^+ to the 1^+ observed in the ${}^7\text{Li}(\pi^+, p)$ reaction, and the 800 MeV ${}^7\text{Li}(p, d){}^6\text{Li}$ reaction. The spectroscopic factors used to normalize these (π^+, p) calculations are the same ones used to normalize the 800 MeV (p, d) calculations presented in Sect. 6.1 (Nor 70), (NG 70).

7.3 ${}^{12}\text{C}(\pi^+, p){}^{11}\text{C}$

A typical spectrum from the 180 MeV ${}^{12}\text{C}(\pi^+, p){}^{11}\text{C}$ reaction (at $\theta = 20^\circ$) is presented in Fig. 7.3.1.

Fig. 7.3.1 Representative spectrum of the 180 MeV $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction collected at $\theta_L = 20^\circ$.

Just as for the ${}^6,{}^7\text{Li}(\pi^+,p)$ spectra, the $\pi\text{NN} \rightarrow \text{Np}$ background dominates. However, many discrete levels are observed whose angular distributions were easily extracted.

The $3/2^-$ ground state and 2.00 MeV $1/2^-$ state of ${}^{11}\text{C}$ are cleanly separated from neighboring levels. Although these levels dominate the spectra of low energy neutron pickup experiments on ${}^{12}\text{C}$ (see Sect. 6.3), they are excited with the same (or less) strength relative to the other levels in the ${}^{12}\text{C}(\pi^+,p){}^{11}\text{C}$ reaction, or in the 800 MeV ${}^{12}\text{C}(p,d){}^{11}\text{C}$ reaction. In fact, the spectra of the ${}^{12}\text{C}(\pi^+,p){}^{11}\text{C}$ reaction are remarkably similar to those of the 800 MeV ${}^{12}\text{C}(p,d){}^{11}\text{C}$ reaction. This point will be elaborated upon in Chapter VIII.

The doublet at 4.32 MeV ($5/2^-$) and 4.80 MeV ($3/2^-$) was not sufficiently resolved in this work to allow extraction of both angular distributions, although at some angles the two states were clearly separated, and in the same ratio as observed in the 800 MeV ${}^{12}\text{C}(p,d){}^{11}\text{C}$ reaction reported in Sect. 6.3. The 6.48 MeV $7/2^-$ level is seen to dominate the spectrum as it did in the 800 MeV ${}^{12}\text{C}(p,d){}^{11}\text{C}$ reaction. In addition, levels are seen at 8.66 MeV, 9.90 MeV and 10.57 MeV which are also seen in the 800 MeV (p,d) work. But the most remarkable similarity in the two

reactions is the excitation of a broad level (or levels) near an excitation of 13.2 MeV, which has not been observed in any other reaction at any other bombarding energy (see Sect. 6.3). Yet this level is seen in both the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction at 90 MeV and 180 MeV, and the 800 MeV $^{12}\text{C}(p, d)^{11}\text{C}$ reaction. This correlation, along with the observed similarity in relative intensities of all the other levels, strongly suggests the two reactions sample the same nuclear structure. This may be pictured by comparing the ONM or TNM diagrams for each reaction: the lower half of each diagram (which is the part sensitive to nuclear structure) is exactly the same for the (π^+, p) and (p, d) reactions. The similarity in the spectra of the two reactions thus strongly suggests a corresponding similarity in the reaction mechanisms.

Notably absent in the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ spectra is the 12.5 MeV $1/2^-$, $T = 3/2$ state reported by previous investigators (Ama+ 78) studying the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction at 49 MeV. This state is particularly interesting since its excitation utilizes the unique isovector nature of the pion, which allows $\Delta T = 3/2$ transfers. The previous experiments have suffered from extremely poor energy resolution and large background, thus it seems likely that this level has been misidentified in the earlier work, and is probably the

13.2 MeV level seen in the 90 MeV and 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ spectra in this work. However, the observed width of the 13.2 MeV level is approximately 2 MeV (FWHM). Therefore the possibility cannot be ruled out that the 12.5 MeV $T = 3/2$ level is partially responsible for the excitation of the (possible unresolved multiplet) 13.2 MeV level seen in the (π^+,p) reaction. This possibility does not seem likely, however, in light of the fact that the same bump at 13.2 MeV, with the same width of ~ 2 MeV, is seen in the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ reaction which cannot excite a $T = 3/2$ level (in the ONM).

The angular distributions extracted for the 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reaction are presented in Fig. 7.3.2. While the statistical errors shown in these angular distributions might seem large to the reader accustomed to angular distributions in low energy nuclear physics, it should be remembered that these cross sections (in the microbarn region) are obtained with an incident pion flux of roughly $10^7 \pi^+/\text{sec}$, which corresponds to less than one picoampere of incident beam. All the angular distributions exhibit roughly the same featureless shape. Apparently, little J^π information is available in the angular distributions of the (π^+,p) reaction.

Fig. 7.3.2 Angular distributions of some of the levels excited in the 180 MeV $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction.

The energy dependence of the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ reaction is shown for $T_\pi = 90$ MeV and 180 MeV in Fig. 7.3.3, for the transition to the ^{11}C ground state. The 90 MeV data are only 20-30% higher at forward angles than the 180 MeV data. In addition, the 90 MeV data show a minimum near 90° which is not seen in the 180 MeV data. This is the same situation presented by the $^6, ^7\text{Li}(\pi^+, p)$ data. Direct comparison of the energy dependence observed in the data, with the calculations presented in Sect. 3.2, again suggests that the pion is propagating off-shell in the (π^+, p) reaction.

Presented in Fig. 7.3.4 are the 180 MeV $^{12}\text{C}(\pi^+, p)^{11}\text{C}(\text{g.s.})$ data, with the predictions of Sect. 3.2 for 180 MeV and 90 MeV incident pions. Although the magnitude of the 90 MeV prediction is too low only a factor of roughly two at forward angles, the slope of the prediction is much too small compared to the 180 MeV data (which are similar in shape and magnitude to the 90 MeV data at forward angles). The 180 MeV prediction not only exhibits the same shape problem, but is too low compared to the data by a factor of about 25 at forward angles.

7.4 $^{13}\text{C}(\pi^+, p)^{12}\text{C}$

Shown in Fig. 7.4.1 is a sample spectrum of the 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reaction, collected at $\theta = 20^\circ$. The $\pi\text{NN} - \text{NP}$ continuum again dominates the

Fig. 7.3.3 Energy dependence of the $^{12}\text{C}(\pi^+, p)^{11}\text{C}$ (g.s.) reaction, sampled at 90 MeV (closed circles) and 180 MeV (open circles).

Fig. 7.3.4 The 180 MeV $^{12}\text{C}(\pi^+, p)^{11}\text{C}(\text{g.s.})$ data are presented with the results of calculations for the 90 MeV (solid curve) and 180 MeV (dashed curve) $^{12}\text{C}(\pi^+, p)^{11}\text{C}(\text{g.s.})$ reaction.

Fig. 7.4.1 Typical spectrum of the 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reaction at $\theta_L = 20^\circ$.

spectrum. The cleanly separated levels are the same ones observed in the 800 MeV $^{13}\text{C}(p,d)^{11}\text{C}$ reaction, and include the 0^+ ground, state, 4.44 MeV 2^+ $T = 0$, 9.64 MeV 3^- , 12.71 MeV 1^+ $T = 1$, 14.08 MeV 4^+ , 15.11 MeV 1^+ $T = 1$, and the 16.1 MeV 2^+ $T = 1$ states. Also seen higher in excitation are levels at 20.6 MeV and 25.6 MeV, which are only weakly seen (if at all) in lower energy neutron pickup studies on ^{13}C , but are excited strongly in the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction. Contaminant peaks from the $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reaction, arising from the ^{12}C contaminant in the target, appear at the higher excitation energies. The similarity of the $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ spectra and those of the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction (see Sect. 6.2) is again striking.

An important element of this similarity is the weak excitation of the 9.64 MeV 3^- level relative to the 4.44 MeV 2^+ level in the $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ spectra. In an argument developed in Sect. 6.2 for the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction (and which is entirely applicable to the $^{13}\text{C}(\pi^+,p)^{12}\text{C}$ reaction) it was shown that this feature suggests that the TNM is not the dominating reaction mechanism. In light of the remarkable similarity observed in spectra of the (π^+,p) and 800 MeV (p,d) reactions on all of the targets in common, it would not be surprising to find that the two reactions

are governed by similar reaction mechanisms.

The excitation of the 14.1 MeV 4^+ level in ^{12}C is noteworthy. Since excitation of this level cannot take place via a simple (single step) neutron pickup, its excitation may be considered as evidence of the importance of two step (coupled channels) mechanisms in the (π^+, p) reaction. A similar conclusion was reached concerning the excitation of this level in the 800 MeV $^{13}\text{C}(p, d)^{12}\text{C}$ reaction, where (see Sect. 6.2) a CCBA calculation was found to be in qualitative agreement with the data.

Angular distributions of some of the levels excited in the 180 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reaction are displayed in Fig. 7.4.2. Nor surprises exist in these structureless angular distributions. Note the good agreement of data collected at $\theta = 20^\circ$, for runs taken at the beginning and end of the experiment.

The energy dependence of the $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reaction is illustrated in Fig. 7.4.3, which shows data for the 90 MeV and 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ (4.44 MeV 2^+) reaction. The 90 MeV data are about two times higher than the 170 MeV data. In addition, the 90 MeV data show a minimum near 100° . The collection of data on the 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reaction stopped at 80° , however, so no comparison can be made pertaining to the shapes of the angular distributions at the two

Fig. 7.4.2 Angular distributions of some of the levels excited in the 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reaction.

Fig. 7.4.3 Energy dependence of the $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ (4.44 MeV) reaction, sampled at 90 MeV (closed circles) and 170 MeV (open circles).

energies, except to note that the 170 MeV data fall off at a slightly steeper rate.

Calculations for the $^{13}\text{C}(\pi^+, p)^{12}\text{C}(\text{g.s.})$ reaction are presented in Fig. 7.4.4, along with the 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}(0^+ \text{ g.s.})$ data. The magnitude of the forward angle prediction for 90 MeV is about two times lower than the 90 MeV data (not shown). The magnitude of the 180 MeV forward angle predictions is more than fifty times too low. Both calculations fail to reproduce the shape of the data—the predicted slope of the angular distributions is much too small.

More detailed comparison of these results to the (p,d) results will be found in Chapter VIII, and general conclusions about the reaction mechanisms will be presented.

Fig. 7.4.4 The 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}(\text{g.s.})$ data are presented with the results of calculations for the 90 MeV (solid curve) and 170 MeV (dashed curve) $^{13}\text{C}(\pi^+, p)^{12}\text{C}(\text{g.s.})$ reaction.

CHAPTER VIII

COMPARISONS AND CONCLUSIONS

8.1 The 800 MeV (p,d) Reaction

This dissertation has presented the results of the 800 MeV (p,d) reaction on targets of ${}^7\text{Li}$, ${}^{12}\text{C}$, ${}^{13}\text{C}$, ${}^{16}\text{O}$, ${}^{25}\text{Mg}$, ${}^{28}\text{Si}$, and ${}^{40}\text{Ca}$. This list constitutes a fair sampling of mass number shape (including spherical, and deformed nuclei of positive and negative deformation), neutron excess (0 or 1), binding energy, and valence neutron orbitals ($s_{1/2}$, $p_{3/2}$, $p_{1/2}$, $d_{5/2}$ and $d_{3/2}$). The good experimental resolution achieved in these studies permitted the identification of many discrete levels in the residual nuclei.

The one nucleon (pickup) model in the form of the exact finite range DWBA has been applied to the ground state angular distributions of all the targets. The agreement is quite good in each case, over the entire range of mass number, although the calculations for the population of the ${}^6\text{Li}$ ground state are somewhat too low. Both methods for obtaining deuteron optical potentials account for the general features of the data. The only exception occurs for the ${}^{40}\text{Ca}(p,d){}^{39}\text{Ca}$ reaction, where usage of the KMT

deuteron optical potential (Sect. 2.6) gives rise to an overprediction of the data by a factor of about five. In addition, too much structure is evident in some of the calculations employing the KMT folded deuteron optical potential.

The pickup model also accounts reasonably well for the population of excited states in the residual nuclei, with one glaring exception. This exception is the 3^+ level of ${}^6\text{Li}$, which is underpredicted by an order of magnitude. Since coupled channel calculations indicate the unimportance of two-step contributions to this particular level, the reason for this failure remains a mystery.

The importance of two step mechanisms for the excitation of some transitions in the 800 MeV (p,d) reaction has been demonstrated (KO 77). The excitations of levels which cannot be populated via single step neutron pickup (such as the 14.08 MeV 4^+ level of ${}^{12}\text{C}$, and the 5.24 MeV $5/2^+$ and 7.28 MeV $7/2^+$ levels of ${}^{15}\text{O}$) are reasonably well described in coupled channels calculations employing simple two-step paths despite the fact that the observed strength of these levels is comparable to, or greater than, the observed strength to pure neutron hole levels. More evidence suggesting the importance of two step mechanisms may be understood by recognizing that the ground states of

the residual nuclei usually exhaust at least 50% of the spectroscopic strength available for pickup of valence neutrons of that particular j-value. Nevertheless, most of the strength observed in the 800 MeV (p,d) reaction is found at higher excitation energies.

Evidence that high spin states are preferentially populated in the 800 MeV (p,d) reaction has been found. The 3^+ level of ${}^6\text{Li}$, 2^+ and 4^+ levels of ${}^{12}\text{C}$, $5/2^-$ and $7/2^-$ levels of ${}^{11}\text{C}$, and $5/2^+$ and $7/2^+$ levels of ${}^{15}\text{O}$ are all examples of this preferential population in the high momentum transfer (p,d) reaction. Circumstantial evidence has also been presented which suggests a correlation between several strongly excited levels in ${}^{27}\text{Si}$ (above 7 MeV in excitation energy) and high spin states. A similar correlation was presented for levels observed in ${}^{11}\text{C}$.

Certainly an important result of this work is the population of many new, previously unobserved high-lying levels in almost every one of the target nuclei studied. These new levels, are, for the most part, more strongly excited than the known low-lying levels. The nature of most of these new levels is probably one of high angular momentum. In several of the nuclei studied, broad regions of enhanced excitation whose nature is unknown appear at high excitation energies.

A disappointing aspect of the 800 MeV (p,d) reaction is that the featureless angular distributions contain no information on the spin or parity of the final states. Thus, although many new levels are populated, no indication of their angular momentum or parity is evident, other than the suggestion that they involve large angular momentum transfers. However, the existence of these states at 800 MeV suggests that they be studied at lower bombarding energies (200 - 400 MeV), where their quantum numbers may be more easily established.

Although the shapes of the angular distributions reveal little spin or parity information, there does seem to be some sensitivity to the details of the neutron bound state wavefunction, especially evident in the lighter targets. Calculations have been presented which support the claim that the slight structure observed in the angular distributions reflects the high Fourier components of the neutron bound state wavefunction.

Some evidence has been found which suggests the two nucleon model does not dominate in the 800 MeV (p,d) reaction to low-lying states. The weakness of the 9.64 MeV 3^- level populated in the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ reaction, contrasted with its strong excitation in low energy stripping reactions, is

difficult to reconcile with the two nucleon model which predicts a strong excitation of this 3^- level in the pickup reaction.

Thus it has been shown that the ONM which works so well for the description of the low energy (p,d) reaction is adequate for the description of the 800 MeV (p,d) reaction, if done in exact finite range and, in some cases, extended to second order.

8.2 The (π^+, p) Reaction Near 90 MeV and 180 MeV

The first high resolution measurements of the (π^+, p) reaction have been performed on targets of ^6Li , ^7Li (at 75 MeV and 175 MeV), ^{12}C (at 90 MeV and 180 MeV), and on ^{13}C (at 90 MeV and 170 MeV). Many discrete levels were resolved in the spectra of the carbon isotopes, in spite of the large yield from the background $\pi\text{NN} \rightarrow \text{Np}$ process.

Agreement of the theoretical curves with the ground state angular distributions is poor. This may be due to either an improper treatment of ingredients in the calculation, or a fundamental inadequacy of the pickup model applied to the (π^+, p) reaction.

The energy dependence of the (π^+, p) reaction was studied at pion bombarding energies on and off the (3,3) resonance. The low energy (π^+, p) angular distributions exhibit a broad minimum near 90° not

seen in the high energy angular distributions. The magnitudes of the on-resonance data are typically a factor of about two less than the off-resonance data. The calculations, however, predict a much greater energy dependence, which was shown to be due to the different pion optical potentials used on and off the resonance. This suggests that the (π^+, p) reaction is extremely sensitive to the off-shell behavior of the pion optical potential—a quantity not measured in elastic scattering. Therefore the (π^+, p) reaction may prove to be a useful tool in the determination of the off-shell behavior of the pion optical potential, once the remaining ambiguities in the calculations are removed.

The two step mechanism contributes to the (π^+, p) reaction, as demonstrated by the strong excitation of the 14.08 MeV 4^+ level of ^{12}C . In addition, high spin states appear to be preferentially excited, just as they were for the 800 MeV (p, d) reaction.

The weak excitation of the 9.64 MeV 3^- level in ^{12}C suggests that the two nucleon model is not dominating the (π^+, p) reaction mechanism at these bombarding energies, just as was found to be the case for the 800 MeV (p, d) reaction. However, it is interesting to note that the intensity of the 3^- level relative to the 4.44 MeV 2^+ level is greater in the

(π^+, p) reaction than it is in the 800 MeV (p, d) reaction. While the ratio of the 3^- strength to the 2^+ strength is still much less than that observed in low-energy stripping reactions, the difference observed in this ratio for the (π^+, p) and 800 MeV (p, d) reactions may suggest that a two nucleon mechanism while still not dominating is relatively more important in the (π^+, p) reaction than it is in the 800 MeV (p, d) reaction.

8.3 Comparison of the (p, d) and (π^+, p) Reactions

The (p, d) and (π^+, p) reactions have been studied on common targets at nearly the same momentum transfers. The spectra of each reaction are observed to be almost identical at equivalent momentum transfers, on a given target. This near equivalence is demonstrated in Fig. 8.3.1 for the 800 MeV (p, d) and 180 MeV (π^+, p) reactions on ^{12}C , and in Fig. 8.3.2 for the 800 MeV (p, d) and 170 MeV (π^+, p) reactions on ^{13}C . These figures compare reactions at the same momentum transfer, with the same scale in excitation energy. The similarity of the two reactions is striking. The same 2 MeV wide bump at 13.2 MeV in ^{11}C is seen in both reactions. The 20.6 MeV and 25.4 MeV levels in ^{12}C are also seen by both reactions. All three of these levels are unique to these high momentum transfer reactions (except the 20.6 MeV level is

Fig. 8.3.1 Spectra of the 800 MeV $^{12}\text{C}(p,d)^{11}\text{C}$ and 180 MeV $^{12}\text{C}(\pi^+,p)^{11}\text{C}$ reactions are compared at a momentum transfer of 2.8 fm^{-1} .

Fig. 8.3.2 Spectra of the 800 MeV $^{13}\text{C}(p,d)^{12}\text{C}$ and 170 MeV $^{13}\text{C}(\pi^+, p)^{12}\text{C}$ reactions are compared near a momentum transfer of 2.6 fm^{-1} .

very weakly excited at lower energies). In addition, nearly the same relative intensities are observed in each reaction, in spite of the fact that the relative intensities observed in low energy neutron pickup experiments are entirely different from those of the present work.

A simple pickup model describes the 800 MeV (p,d) data adequately, and the weak excitation of the $^{12}\text{C } 3^-$ level is well described with this mechanism. It seems reasonable that the one nucleon mechanism is an appropriate model for the (π^+,p) reaction, as well as for the (p,d) reaction, although inadequate pion distorting potentials prevent a quantitative description of the (π^+,p) cross sections. The remarkable similarity in the spectra of the (π^+,p) and 800 MeV (p,d) reactions suggests a corresponding similarity in the reaction mechanisms.

ACKNOWLEDGMENTS

It has been my privilege to work at several facilities, and with many excellent physicists during the course of my graduate studies. There is a long list of those who I wish to thank.

First of all, I am deeply indebted to my advisor, Jerry Peterson, and to the intermediate energy physics program which he has established. I was given every opportunity and every resource to do research in Boulder and Los Alamos. Jerry's eagerness to supply these items, as well as exciting discussions about new ideas and current projects, made my life as an experimental physicist very enjoyable and fruitful. He was an ideal advisor, and I urge other graduate students to take advantage of the opportunity to work with him in the medium energy program.

I also owe a great deal of thanks to Jim Shepard, who played the dual role of friend and advisor, as well as theorist and experimentalist. Jim had the formidable task of turning a hard core experimentalist fresh from the data farm at LAMPF into a physicist to whom the word "theory" was not completely foreign. Not only did I really enjoy

working with Jim, but also I expect his influence and attitudes about physics will stay with me for a long time.

I also wish to thank Ernie Rost for teaching me nuclear physics in class as well as out of the classroom, and for straightening me out on so many theoretical aspects of this work.

I owe Dale Prull a great deal of thanks for guiding me in my computer education. Computers were essential in every phase of this work, and without Dale's help and guidance I would never have mastered them.

I also owe Dick Boudrie thanks for my education at LAMPF. He helped me to become familiar with HRS and EPICS, and aided me in software development, as well as providing friendship in Los Alamos.

I benefited immensely from my associations with Tom Bauer, Chris Morris, and Arch Thiessen, and I wish to thank each of them for the many things they have taught me.

I also wish to thank the staffs of both the University of Colorado Cyclotron Lab and LAMPF, for providing so much assistance to me over the years.

REFERENCES

- Ada+ 79 G. S. Adams, Th. S. Bauer, G. Igo, G. Pauletta, C. A. Whitten, Jr., A. Wriekat, G. W. Hoffman, G. R. Smith, M. Gazzaly, L. Ray, W. G. Love, and F. Petrovich, Phys. Rev. Lett. 43, 421 (1979).
- And+ 78 R. E. Anderson, J. J. Kraushaar, J. R. Shepard, and J. R. Comfort, Nucl. Phys. A311, 93 (1978).
- AFS 67 E. H. Auerbach, D. M. Fleming and M. M. Sternheim, Phys. Rev. 162, 1683 (1967).
- AHR 74 R. A. Arndt, R. H. Hackman, and L. D. Roper, Phys. Rev. C9, 555 (1974).
- AHR 77 R. A. Arndt, R. H. Hackman, and L. D. Roper, Phys. Rev. C15, 1002 (1977).
- Ajz 75 F. Ajzenberg-Selove, Nucl. Phys. A248, 1 (1975).
- Ajz 76 F. Ajzenberg-Selove, Nucl. Phys. A268, 1 (1976).
- Ajz 79 F. Ajzenberg-Selove, Nucl. Phys. A320, 1 (1979).
- All 77 J. C. Allred, LAMPF: The Meson Factory, LASL Report LA-0590, 1977.
- Ama+ 74 J. Amato, R. L. Burman, R. Macek, J. Oostens, W. Shlaer, E. Arthur, S. Sobottka, and W. C. Lam, Phys. Rev. C9, 501 (1974).
- Ama+ 78 J. Amann, P. Barnes, K. Doss, S. Dytman, R. Eisenstein, J. Sherman, and W. Wharton, Phys. Rev. Lett. 40, 758 (1978).
- Aus 70 N. Austern, Direct Nuclear Reaction Theories, Vol. XXV, John Wiley & Sons, N.Y., 1970.

- Bac+ 69 D. Bachelier, M. Bernas, I. Brissaud, C. Detraz, and P. Radvanyi, Nucl. Phys. A126, 60 (1969).
- Bac+ 77 D. Bachelier, J. L. Boyard, T. Hennino, J. C. Jourdain, P. Radvanyi, and M. Roy-Stéphan, Phys. Rev. C15, 2139 (1977).
- Bak+ 74 S. D. Baker, R. Bertini, R. Beurtey, F. Brochard, G. Euge, H. Catz, A. Chaumeaux, G. Cvijanovich, J. M. Durand, J. C. Faivre, J. M. Fontaine, D. Garreta, F. Hibou, D. Legrand, J. C. Lugol, J. Saudinos, J. Thirion, and E. Rost, Phys. Lett. 52B, 57 (1974).
- Bart+ 78 N. Barash-Schmidt, A. Barbaro-Galtieri, C. Bricman, R. L. Crawford, C. Dionisi, R. J. Hemmingway, C. P. Horne, R. L. Kelly, M. L. Losty, M. Mazzucato, L. Montanet, A. Rittenberg, M. Roos, T. G. Trippe, G. P. Yost, and F. E. Armstrong, Phys. Lett. 75B, 1 (1978).
- Bau+ 77 T. Bauer, A. Boudard, H. Catz, A. Chaumeaux, P. Couvert, M. Carbon, J. Guyot, D. Legrand, J. C. Lugol, M. Matoba, B. Mayer, J. P. Tabet, and Y. Terrien, Phys. Lett. 67B, 265 (1977).
- BBD 72 F. A. Bumiller, F. R. Buskirk, and J. N. Dyer, Phys. Rev. C5, 391 (1972).
- Ben+ 78 R. D. Bent, P. T. Debevec, P. H. Pile, R. E. Pollack, R. E. Marrs, and M. C. Green, Phys. Rev. Lett. 40, 495 (1978).
- Bla+ 77 G. S. Blanpied, W. R. Coker, R. P. Liljestrang, L. Ray, G. W. Hoffman, D. Madland, C. L. Morris, J. C. Pratt, J. E. Spencer, H. A. Thiessen, N. M. Hintz, G. S. Kyde, M. A. Oothoudt, T. S. Bauer, J. S. Fong, G. Igo, R. J. Ridge, C. A. Whitten, T. Kozlowski, D. K. McDaniels, P. Varghese, P. M. Lang, H. Nann, K. K. Seth, and C. Glashausser, Phys. Rev. Lett. 39, 1447 (1977).

- Bla 78 G. S. Blanpied, Ph.D. thesis, University of Texas, 1978, Los Alamos Scientific Laboratory Report LA-7262-T.
- Bon+ 78 B. E. Bonner, J. E. Simmons, M. L. Evans, G. Glass, J. C. Hiebert, M. Jain, L. C. Northcliffe, C. W. Bjork, P. J. Riley, and C. G. Cassapakis, Phys. Rev. C17, 671 (1978).
- Bor+ 76 Y. LeBornec, B. Tatischeff, L. Bimbot, I. Brissaud, H. D. Molmgren, J. Källne, F. Reide, and N. Willis, Phys. Lett. 61B, 47 (1976).
- Bou 78 R. L. Boudrie, private communication, 1976.
- Bru 78 G. Bruge, Saclay internal report DPh-N/ME/78-1, 1978 (unpublished).
- CK 67 S. Cohen and D. Kurath, Nucl. Phys. A101, 1 (1967).
- CW 69 N. S. Cragie and C. Wilkin, Nucl. Phys. B14, 477 (1969).
- Dah+ 73 S. Dahlgren, P. Grafström, B. Hoistad and A. Asberg, Phys. Lett. 47B, 439 (1973).
- Dah+ 74 S. Dahlgren, P. Grafström, B. Hoistad and A. Asberg, Nucl. Phys. A227, 245 (1974).
- DDD 74 C. W. DeJager, H. DeVries, and C. DeVries, Atomic Data and Nuclear Data Tables, Vol. 14, 479, Academic Press, N.Y., 1974.
- Deh 72 D. Dehnhard, Phys. Lett. 38B, 389 (1972).
- Dev+ 69 D. W. Devins, S. M. Bunch, H. H. Forster, J. Hoxhikian, and C. C. Kim, Nucl. Phys. A126, 261 (1969).
- DG 66 L. R. Dodd and K. R. Greider, Phys. Rev. 146, 675 (1966).
- Dig+ 77 N. DiGiacomo, A. S. Rosenthal, E. Rost, and D. A. Sparrow, Phys. Lett. 66B, 421 (1977).

- DiG 79 N. J. DiGiacomo, Ph.D. thesis, University of Colorado, 1979.
- DSB 63 R. M. Drisko, G. R. Satchler, and R. H. Bassel, Phys. Lett. 5, 347 (1963).
- Dun 76 E. Dunn, Medium Energy Physics Program Progress Report, LASL Report LA-6678-PR, Aug.-Oct. 1976.
- EHJ 53 J. P. Elliott, J. Hope, and H. A. Jahn, Phil. Trans. A246, 241 (1953).
- Elb+ 75 F. El-Bedewi, M. Shalaby, A. Khazbak, and F. Raoof, J. Phys. G1, 749 (1975).
- Emi 76 R. A. Emigh, Ph.D. thesis, University of Colorado, 1976.
- ES 67 L. R. B. Elton and A. Swift, Nucl. Phys. A94, 52 (1967).
- EV 78 P. M. Endt and C. Van Der Leun, Nucl. Phys. A310, 1 (1978).
- Fes 58 H. Feshbach, Ann. Rev. Nucl. Sci., Vol. 8, Annual Reviews, Inc., Palo Alto, Cal., 1958, p. 49.
- FGH 70 H. Feshbach, A. Gal, J. Hüfner, Ann. Phys. 66, 20 (1971).
- Ger+ 75 E. Gerlic, H. Langevin-Joliot, P. Roos, J. Van de Wiele, J. P. Didelez, and G. Duhamel, Phys. Rev. C12, 2106 (1975).
- GG 53 M. Gell-Mann and M. L. Goldberger, Phys. Rev. 91, 398 (1953).
- GW 64 M. L. Goldberger and K. M. Watson, Collision Theory, John Wiley & Sons, N.Y., 1964.
- HNL 73 S. L. Hausladen, C. E. Nelson, and R. O. Lane, Nucl. Phys. A217, 563 (1973).
- Hoi 79 B. Hoistad, "Advances in Nuclear Physics," Vol. 11, Plenum Press, N.Y. (1979).

- Hoi 79b B. Hoistad, T. Johansson, and O. Jonsson, Phys. Lett. B79, 385 (1979).
- How 74 H. Howard, LAMPF Users Handbook, LASL Report MP-DO-1-UHB, 1974.
- IHS 73 A. Ingemarsson, E. Hagberg and H. Sherif, Nucl. Phys. A216, 271 (1973).
- Ive+ 78 S. Iversen, A. Obst, K. K. Seth, H. A. Thiessen, C. L. Morris, N. Tanaka, E. Smith, J. F. Amann, R. Boudrie, G. Burleson, M. Devereux, L. W. Swenson, P. Varghese, K. Boyer, W. J. Braithwaite, W. Cottingham, and C. F. Moore, Phys. Rev. Lett. 40, 17 (1978).
- Jac 62 J. D. Jackson, Classical Electrodynamics, John Wiley & Sons, N.Y., 1962.
- JE 70 W. B. Jones and J. M. Eisenberg, Nucl. Phys. A154, 49 (1970).
- Joh+ 79 R. R. Johnson, T. Masterson, B. Bassalleck, W. Gyles, T. Marks, K. L. Erdman, A. W. Thomas, D. R. Gill, E. Rost, J. J. Kraushaar, J. Alster, C. Sabev, J. Arvieux, and M. Krell, Phys. Rev. Lett. 43, 844 (1979).
- JS 69 R. C. Johnson, and P. J. R. Soper, Phys. Rev. C1, 976 (1970).
- JVa 51 H. A. Jahn and H. Van Wieringen, Proc. Roy. Soc. (London) A209, 502 (1951).
- Käl+ 74 J. Källne, B. Fagerstrom, O. Sundberg, and G. Tibell, Phys. Lett. 52B, 313 (1974).
- Käl+ 78 J. Källne, D. A. Hutcheon, W. J. McDonald, A. N. Anderson, J. L. Beveridge, and J. Rogers, Phys. Rev. Lett. 41, 1638 (1978).
- Käl+ 78 J. Källne, H. A. Thiessen, C. L. Morris, S. L. Verbeck, G. R. Burleson, M. J. Devereaux, J. S. McCarthy, J. E. Bolger, C. F. Moore, and C. A. Goulding, Phys. Rev. Lett. 40, 378 (1978).

- KMT 59 A. K. Kerman, H. McManus, and R. M. Thaler, Ann. Phys. (N.Y.) 8, 551 (1959).
- KO 77 J. Kallne and A. Obst, Phys. Rev. C15, 477 (1977).
- Koz 68 R. L. Kozub, Phys. Rev. 172, 1078 (1968).
- Kun 79 P. D. Kunz, University of Colorado, private communication (1979).
- Kun 79a P. D. Kunz, the Program DWUCK4, University of Colorado, private communication.
- Kun 79b P. D. Kunz, The Program DWUCK5, University of Colorado, private communication.
- Kun 79c P. D. Kunz, The Program CHUCK, University of Colorado, private communication.
- LE 66 J. Letourneux and J. M. Eisenberg, Nucl. Phys. 87, 331 (1966).
- Lee+ 68 J. K. P. Lee, S. K. Mark, P. M. Portner, and R. B. Moore, Nucl. Phys. A106, 357 (1968).
- Liv 77 M. S. Livingston, LAMPF: A Nuclear Research Facility, LASL Report LA-6878-MS, 1977.
- MAW 68a M. MacGregor, R. A. Arndt, and R. M. Wright, Phys. Rev. 169, 1149 (1968).
- MAW 68b M. MacGregor, R. A. Arndt, and R. M. Wright, Phys. Rev. 173, 1272 (1968).
- Mer 61 E. Merzbacher, Quantum Mechanics, John Wiley & Sons, N.Y., 1961.
- Mes 63 Albert Messiah, Quantum Mechanics, Wiley, N.Y., 1963.
- Mil 79 D. Miller, Indiana University Cyclotron Facility, private communication (1979).
- Mor 76 C. L. Morris, Phys. Rev. C13, 1755 (1976).
- Mor 78 C. L. Morris, H. A. Thiessen, G. W. Hoffmann, IEEE Trans. on Nucl. Science, Vol. NS-25, no. 1, Feb. 1978.

- MP 74 G. A. Miller and S. C. Phatak, Phys. Lett. 51B, 129 (1974).
- NG 70 J. L. Norton and P. Goldhammer, Nucl. Phys. A165, 33 (1970).
- Nob 76 J. V. Noble, A. I. P. Conference Proceedings No. 33 of the Carnegie-Mellon Conference on Meson-Nuclear Physics (1976).
- Nor 70 J. Norton, Ph.D. thesis, University of Kansas, 1970.
- Par+ 74 L. J. Parish, R. A. Brown, K. A. Eberhard, A. Richter, and W. von Witsch, Phys. Rev. C9, 876 (1974).
- Peh+ 65 R. H. Pehl, J. Cerny, E. Rivet, and B. G. Harvey, Phys. Rev. B140, 605 (1965).
- Pet+ 75 R. J. Peterson, J. J. Kraushaar, R. A. Ristinen, H. A. Thiessen, and M. Rickey, Nucl. Inst. Meth. 129, 47 (1975).
- Pif+ 77 J. Piffaretti, R. Corfu, J. P. Egger, P. Gretillat, C. Lunke, E. Schwarz, C. Perrin, and B. M. Freedom, Phys. Lett. 71B, 324 (1977).
- Pru 79 D. Prull, the RSX11M version of SPECTR, University of Colorado, private communication, 1979.
- Ray+ 78 L. Ray, G. W. Hoffman, G. S. Blanpied, W. R. Coker, and R. P. Liljestrang, Phys. Rev. C18, 1756 (1978).
- Ray+ 78 L. Ray, G. S. Blanpied, W. R. Coker, R. P. Liljestrang, and G. W. Hoffman, Phys. Rev. Lett. 40, 1547 (1978).
- Ray 79 M. Rayman, private communication, 1979.
- RCH 78 L. Ray, W. R. Coker, and G. W. Hoffman, Phys. Rev. C18, 2641 (1978).
- Ric+ 70 C. Richard-Serre, W. Hirt, D. F. Measday, E. G. Michaelis, M. J. M. Saltmarsh, and P. Skarek, Nucl. Phys. B20, 413 (1970).

- RK 73 E. Rost and P. D. Kunz, Phys. Lett. 43B, 17 (1973).
- Roos+ 75 P. G. Roos, S. M. Smith, V. K. C. Cheng, G. Tibell, A. A. Cowley, and R. A. J. Riddle, Nucl. Phys. A255, 187 (1975).
- Ros 78 A. Rosenthal, Ph.D. thesis, University of Colorado, 1978.
- Ros 78 E. Rost, The Program OPTIMR, University of Colorado, private communication, 1978.
- Ros 79 E. Rost, University of Colorado, private communication, 1979.
- RS 75 E. Rost and J. R. Shepard, Phys. Lett. 59B, 413 (1975).
- RSS 78 E. Rost, J. R. Shepard, and D. A. Sparrow, Phys. Rev. C17, 1513 (1978).
- RWF 65 L. D. Roper, R. M. Wright, and B. T. Feld, Phys. Rev. 138, 190 (1965).
- Sat 58 G. R. Satchler, Ann. of Phys. 3, 275 (1958).
- Sch 76 S. Schlaer, Data Acquisition Program Q, LAMPF, 1976 (unpublished).
- Sco+ 70 D. K. Scott, P. M. Portner, J. M. Nelson, A. C. Shotter, A. J. Mitchell, N. S. Chant, D. G. Montague, and K. Ramavataram, Nucl. Phys. A141, 497 (1970).
- SF 74 A. deShalit and H. Feshbach, Theoretical Nuclear Physics, Volume 1: Nuclear Structure, Wiley, N.Y., 1974.
- She 78 J. R. Shepard, University of Colorado, private communication, 1978.
- She 79 J. R. Shepard, University of Colorado, private communication, 1979.
- SM 70 I. Sick and J. S. McCarthy, Nucl. Phys. A150, 631 (1970).

- Smi 76 P. A. Smith, Ph.D. thesis, University of Michigan, 1976..
- Sto+ 67 R. Stock, R. Bock, P. David, H. H. Duhm, and T. Tamura, Nucl. Phys. A104, 136 (1967).
- SYM 57 H. P. Stapp, T. J. Ypsilantio, and N. Metropolis, Phys. Rev. 105, 302 (1957).
- Tak+ 68 H. Taketani, J. Muto, H. Yamaguchi, and J. Kokame, Phys. Lett. 27B, 625 (1968).
- Thi 70 H. A. Thiessen, A Proposal for EPICS, LASL Report LA-4534-MS, 1970.
- Thi+ 77 H. A. Thiessen, J. C. Kallne, J. F. Amann, R. J. Peterson, S. J. Greene, S. L. Verbeck, G. R. Burleson, S. G. Iverson, A. W. Obst, K. K. Seth, C. F. Moore, J. E. Bolger, W. J. Braithwaite, D. C. Slater, and C. L. Morris, Epics Pion Channel Performance, LASL Report LA-6663-MS, March 1976.
- Thi 78 H. A. Thiessen, MP-10 Progress Report, private communication, April 1978.
- Tob 61 W. Tobocman, Theory of Direct Nuclear Reactions, Oxford University Press, London, 1961.
- Tyr+ 66 H. Tyren, S. Kullander, O. Sundberg, R. Ramachandran, P. Isacson, and T. Berggren, Nucl. Phys. 79, 321 (1966).
- Wat 58 K. M. Watson, Rev. Mod. Phys. 30, 565 (1958).
- Wil+ 76 H. B. Willard, B. D. Anderson, H. W. Baer, R. J. Barrett, P. R. Bevington, A. N. Anderson, H. Willmes, and N. Jarmie, Phys. Rev. C14, 1545 (1976).
- Zei 71 B. Zeidman, The HRS Spectrometer System: Concepts and Design, LASL Report LA-4773-MS Part 1, 1977.
- Zur 78 D. H. Zurstadt, The Program OPTIMR, University of Colorado, private communication.

APPENDIX

Tables of the (π^+ ,p) and (p,d) Data

(π^+ ,p) DATA TABLE

TARGET MASS = 6.00

PION ENERGY = 75.0

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
21.0	0.324E+02	0.26E+01
41.9	0.167E+02	0.16E+01
62.6	0.740E+01	0.60E+00
83.0	0.419E+01	0.53E+00
103.0	0.486E+01	0.51E+00
118.2	0.128E+02	0.88E+00

(PI+,P) DATA TABLE

TARGET MASS = 6.00
PION ENERGY = 175.0

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
21.6	0.243E+02	0.27E+01
37.6	0.619E+01	0.85E+00
53.5	0.431E+01	0.48E+00
53.5	0.359E+01	0.34E+00
69.1	0.196E+01	0.20E+00
84.4	0.912E+00	0.10E+00
99.4	0.419E+00	0.56E-01

(PI+,P) DATA TABLE

TARGET MASS = 7.00
PION ENERGY = 75.0

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
20.9	0.139E+02	0.16E+01
41.7	0.991E+01	0.11E+01
62.2	0.328E+01	0.39E+00
82.5	0.170E+01	0.28E+00
117.8	0.124E+01	0.22E+00

(PI+,P) DATA TABLE

TARGET MASS = 7.00
PION ENERGY = 75.0

DATA FOR PEAK AT 2.20 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
20.9	0.537E+02	0.33E+01
41.7	0.316E+02	0.20E+01
62.2	0.113E+02	0.71E+00
82.5	0.388E+01	0.40E+00
117.8	0.321E+01	0.36E+00

(PI+,P) DATA TABLE

TARGET MASS = 7.00

PION ENERGY = 175.0

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
21.3	0.146E+02	0.84E+00
37.2	0.572E+01	0.11E+01
53.0	0.186E+01	0.15E+00
68.5	0.513E+00	0.58E-01
83.8	0.169E+00	0.17E+00
98.8	0.114E+00	0.11E+00
98.8	0.131E+00	0.38E-01

(PI+,P) DATA TABLE

TARGET MASS = 7.00

PION ENERGY = 175.0

DATA FOR PEAK AT 2.20 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
21.3	0.361E+02	0.14E+01
37.2	0.123E+02	0.18E+01
53.0	0.346E+01	0.22E+00
68.5	0.119E+01	0.87E-01
83.8	0.507E+00	0.34E+00
98.8	0.341E+00	0.11E+00
98.8	0.432E+00	0.56E-01

(PI+,P) DATA TABLE

TARGET MASS = 12.00

PION ENERGY = 180.0

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.388E+02	0.29E+01
15.6	0.321E+02	0.28E+01
20.8	0.243E+02	0.31E+01
20.8	0.240E+02	0.20E+01
20.8	0.239E+02	0.13E+01
24.9	0.161E+02	0.15E+01
31.1	0.975E+01	0.11E+01
31.1	0.774E+01	0.44E+00
31.1	0.342E+01	0.15E+00
36.3	0.509E+01	0.48E+00
41.5	0.337E+01	0.29E+00
51.8	0.186E+01	0.15E+00
62.0	0.606E+00	0.58E-01
72.1	0.316E+00	0.50E-01
92.3	0.585E-01	0.13E-01
102.2	0.731E-01	0.12E-01
117.1	0.501E-01	0.15E-01

(PI+,P) DATA TABLE

TARGET MASS = 12.00

PION ENERGY = 180.0

DATA FOR PEAK AT 2.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.154E+01	0.10E+01
15.6	0.393E+01	0.12E+01
20.8	0.265E+01	0.88E+00
20.8	0.285E+01	0.55E+00
24.9	0.113E+01	0.65E+00
31.1	0.104E+01	0.52E+00
31.1	0.629E+01	0.42E+00
31.1	0.739E+00	0.49E-01
36.3	0.641E+00	0.26E+00
41.5	0.136E+01	0.20E+00
51.8	0.311E+00	0.81E-01
62.0	0.821E-01	0.26E-01
72.1	0.488E-01	0.33E-01
92.3	0.803E-01	0.13E-01
102.2	0.305E-01	0.12E-01

(PI+,P) DATA TABLE

TARGET MASS = 12.00
PION ENERGY = 180.0

DATA FOR PEAK AT 4.56 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.274E+02	0.30E+01
15.6	0.235E+02	0.23E+01
20.8	0.233E+02	0.21E+01
20.8	0.189E+02	0.12E+01
24.9	0.126E+02	0.16E+01
31.2	0.120E+02	0.12E+01
31.2	0.832E+01	0.13E+01
31.2	0.455E+01	0.14E+00
36.3	0.894E+01	0.67E+00
41.5	0.113E+02	0.87E+00
51.8	0.206E+01	0.23E+00
62.0	0.514E+00	0.50E-01
72.2	0.968E-01	0.32E-01
92.3	0.929E-01	0.18E-01
102.2	0.429E-01	0.12E-01

(PI+,P) DATA TABLE

TARGET MASS = 12.00

PION ENERGY = 180.0

DATA FOR PEAK AT 6.48 MEV

THETA(CH) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
-----	-----	-----
10.4	0.426E+02	0.40E+01
15.6	0.406E+02	0.31E+01
20.8	0.285E+02	0.29E+01
20.8	0.294E+02	0.15E+01
24.9	0.233E+02	0.22E+01
31.2	0.194E+02	0.19E+01
31.2	0.173E+02	0.15E+01
31.2	0.672E+01	0.14E+00
36.3	0.123E+02	0.99E+00
41.5	0.182E+02	0.11E+01
51.8	0.323E+01	0.29E+00
62.0	0.763E+00	0.64E-01
72.2	0.768E+00	0.80E-01
92.3	0.194E+00	0.26E-01
102.3	0.740E-01	0.18E-01

(PI+,P) DATA TABLE

TARGET MASS = 12.00

PION ENERGY = 180.0

DATA FOR PEAK AT 8.40 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.164E+02	0.63E+01
15.6	0.191E+02	0.79E+01
20.8	0.125E+02	0.24E+01
20.8	0.135E+02	0.13E+01
24.9	0.841E+01	0.17E+01
31.2	0.176E+02	0.16E+01
36.3	0.364E+01	0.70E+00
41.5	0.892E+01	0.14E+01
51.8	0.191E+01	0.26E+00
62.0	0.652E+00	0.13E+00
72.2	0.191E+00	0.48E-01
92.3	0.159E+00	0.27E-01
102.3	0.699E-01	0.19E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.154E+01	0.27E+00
15.5	0.272E+01	0.38E+00
20.7	0.966E+00	0.58E+00
20.7	0.138E+01	0.29E+00
20.7	0.165E+01	0.18E+00
31.0	0.133E+01	0.20E+00
41.3	0.622E+00	0.70E-01
61.7	0.139E+00	0.60E-01
82.0	0.127E-01	0.13E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 4.40 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.121E+02	0.70E+00
15.5	0.106E+02	0.89E+00
20.7	0.651E+01	0.93E+00
20.7	0.832E+01	0.47E+00
20.7	0.700E+01	0.37E+00
31.0	0.517E+01	0.38E+00
41.3	0.245E+01	0.14E+00
61.8	0.781E+00	0.48E-01
82.0	0.121E+00	0.21E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 9.60 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.198E+01	0.64E+00
15.5	0.313E+01	0.39E+00
20.7	0.108E+01	0.36E+00
20.7	0.154E+01	0.21E+00
20.7	0.199E+01	0.58E+00
31.0	0.786E+00	0.28E+00
41.3	0.295E+00	0.74E-01
61.8	0.462E-01	0.12E-01
82.0	0.335E-01	0.71E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 12.70 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.115E+02	0.98E+00
15.5	0.125E+02	0.99E+00
20.7	0.722E+01	0.11E+01
20.7	0.766E+01	0.58E+00
20.7	0.743E+01	0.43E+00
31.0	0.375E+01	0.43E+00
41.3	0.276E+01	0.15E+00
61.8	0.427E+00	0.12E+00
82.0	0.715E-01	0.29E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 14.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.250E+02	0.16E+01
15.5	0.259E+02	0.15E+01
20.7	0.210E+02	0.21E+01
20.7	0.173E+02	0.89E+00
20.7	0.171E+02	0.67E+00
31.0	0.111E+02	0.79E+00
41.3	0.430E+01	0.20E+00
61.8	0.987E+00	0.36E+00
82.0	0.283E+00	0.46E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 15.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
10.4	0.214E+02	0.16E+01
15.5	0.199E+02	0.14E+01
20.7	0.941E+01	0.17E+01
20.7	0.128E+02	0.86E+00
20.7	0.116E+02	0.62E+00
31.0	0.550E+01	0.78E+00
41.3	0.270E+01	0.19E+00
61.8	0.677E+00	0.37E+00
82.0	0.135E+00	0.38E-01

(PI+,P) DATA TABLE

TARGET MASS = 13.00

PION ENERGY = 170.0

DATA FOR PEAK AT 16.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
10.4	0.256E+02	0.16E+01
15.5	0.223E+02	0.14E+01
20.7	0.171E+02	0.17E+01
20.7	0.154E+02	0.86E+00
20.7	0.138E+02	0.66E+00
31.0	0.663E+01	0.81E+00
41.3	0.327E+01	0.20E+00
61.8	0.144E+01	0.14E+00
82.0	0.187E+00	0.42E-01

800 MEV (P,D) DATA TABLE

TARGET MASS = 7.00

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.8	0.101E+01	0.45E-01
6.8	0.100E+01	0.51E-01
6.8	0.866E+00	0.52E-01
10.0	0.627E+00	0.34E-01
12.7	0.433E+00	0.18E-01
15.9	0.209E+00	0.51E-02
19.8	0.699E-01	0.17E-02
19.8	0.880E-01	0.38E-02
23.7	0.305E-01	0.19E-02
23.7	0.199E-01	0.13E-02
23.7	0.217E-01	0.18E-02
27.6	0.638E-02	0.80E-03
32.7	0.140E-02	0.37E-03
39.1	0.439E-03	0.10E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 7.00

DATA FOR PEAK AT 2.19 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.8	0.546E+01	0.11E+00
6.8	0.540E+01	0.12E+00
6.8	0.529E+01	0.13E+00
10.0	0.311E+01	0.82E-01
12.7	0.232E+01	0.43E-01
15.9	0.102E+01	0.11E-01
19.8	0.292E+00	0.35E-02
19.8	0.369E+00	0.78E-02
23.7	0.154E+00	0.47E-02
23.7	0.130E+00	0.29E-02
23.7	0.139E+00	0.39E-02
27.6	0.401E-01	0.15E-02
32.7	0.119E-01	0.74E-03
39.1	0.232E-02	0.20E-03
46.0	0.640E-03	0.25E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.456E+01	0.14E+00
12.3	0.124E+01	0.45E-01
13.7	0.716E+00	0.23E-01
15.2	0.346E+00	0.14E-01
17.0	0.302E+00	0.10E-01
18.5	0.268E+00	0.34E-02
20.6	0.126E+00	0.54E-02
20.6	0.152E+00	0.59E-02
22.9	0.832E-01	0.24E-02
25.4	0.342E-01	0.10E-02
25.4	0.327E-01	0.15E-02
30.3	0.761E-02	0.44E-03
34.5	0.226E-02	0.16E-03
38.0	0.146E-03	0.44E-04

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 2.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.548E+00	0.50E-01
12.3	0.123E+00	0.17E-01
13.7	0.115E+00	0.10E-01
15.2	0.113E+00	0.87E-02
17.0	0.110E+00	0.64E-02
18.5	0.765E-01	0.44E-02
20.6	0.235E-01	0.25E-02
20.6	0.292E-01	0.28E-02
22.9	0.120E-01	0.96E-03
25.4	0.651E-02	0.48E-03
25.4	0.439E-02	0.60E-03
30.3	0.174E-02	0.25E-03
34.5	0.353E-03	0.88E-04
38.0	0.583E-04	0.29E-04

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 4.32 MEV

THETA(CH) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.204E+01	0.10E+00
12.3	0.924E+00	0.48E-01
13.7	0.798E+00	0.26E-01
15.2	0.379E+00	0.18E-01
17.0	0.233E+00	0.11E-01
18.5	0.127E+00	0.69E-02
20.6	0.358E-01	0.33E-02
20.6	0.469E-01	0.35E-02
22.9	0.247E-01	0.15E-02
25.4	0.138E-01	0.96E-03
25.4	0.126E-01	0.14E-02
30.3	0.444E-02	0.41E-03
34.5	0.144E-02	0.17E-03
38.0	0.160E-03	0.44E-04

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00
DATA FOR PEAK AT 4.80 MEV

THETA(CH) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.6	0.527E+00	0.65E-01
12.3	0.158E+00	0.33E-01
13.7	0.791E-01	0.18E-01
15.2	0.843E-01	0.12E-01
17.0	0.604E-01	0.81E-02
18.5	0.457E-01	0.52E-02
20.6	0.196E-01	0.27E-02
20.6	0.184E-01	0.26E-02
22.9	0.683E-02	0.11E-02
25.4	0.423E-02	0.72E-03
25.4	0.515E-02	0.10E-02
30.3	0.108E-02	0.28E-03
34.5	0.147E-03	0.11E-03
38.0	0.175E-03	0.44E-04

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 6.48 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.360E+01	0.16E+00
12.3	0.278E+01	0.88E-01
13.7	0.209E+01	0.47E-01
15.2	0.125E+01	0.37E-01
17.0	0.723E+00	0.20E-01
18.5	0.381E+00	0.12E-01
20.6	0.116E+00	0.66E-02
20.6	0.132E+00	0.80E-02
22.9	0.683E-01	0.27E-02
25.4	0.344E-01	0.12E-02
25.4	0.327E-01	0.18E-02
30.3	0.571E-02	0.41E-03
34.5	0.215E-02	0.20E-03
38.0	0.830E-03	0.12E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 8.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.735E+00	0.12E+00
12.3	0.314E+00	0.52E-01
13.7	0.192E+00	0.39E-01
15.2	0.879E-01	0.26E-01
17.0	0.463E-01	0.19E-01
18.5	0.411E-01	0.81E-02
20.6	0.145E-01	0.43E-02
20.6	0.250E-01	0.49E-02
22.9	0.143E-01	0.18E-02
25.4	0.633E-02	0.93E-03
25.4	0.504E-02	0.13E-02
30.3	0.174E-02	0.39E-03
34.5	0.569E-03	0.15E-03
38.0	0.874E-04	0.73E-04

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 8.60 MEV

THETA(CH) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.245E+01	0.17E+00
12.3	0.125E+01	0.75E-01
13.7	0.870E+00	0.50E-01
15.2	0.435E+00	0.30E-01
17.0	0.286E+00	0.24E-01
18.5	0.138E+00	0.11E-01
20.6	0.393E-01	0.52E-02
20.6	0.629E-01	0.64E-02
22.9	0.228E-01	0.21E-02
25.4	0.803E-02	0.10E-02
25.4	0.921E-02	0.15E-02
30.3	0.130E-02	0.36E-03
34.5	0.569E-03	0.16E-03
38.0	0.175E-03	0.73E-04

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00
DATA FOR PEAK AT 9.90 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.813E+00	0.50E-01
12.3	0.765E+00	0.50E-01
13.7	0.514E+00	0.30E-01
15.2	0.291E+00	0.34E-01
17.0	0.200E+00	0.15E-01
18.5	0.114E+00	0.90E-02
20.6	0.371E-01	0.41E-02
20.6	0.361E-01	0.40E-02
22.9	0.239E-01	0.20E-02
25.4	0.760E-02	0.72E-03
25.4	0.888E-02	0.15E-02
30.3	0.190E-02	0.36E-03
34.5	0.756E-03	0.17E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00

DATA FOR PEAK AT 10.60 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.507E+00	0.10E+00
12.3	0.574E+00	0.41E-01
13.7	0.445E+00	0.30E-01
15.2	0.281E+00	0.29E-01
17.0	0.199E+00	0.15E-01
18.5	0.121E+00	0.90E-02
20.6	0.443E-01	0.50E-02
20.6	0.474E-01	0.40E-02
22.9	0.249E-01	0.20E-02
25.4	0.110E-01	0.10E-02
25.4	0.806E-02	0.14E-02
30.3	0.331E-02	0.44E-03
34.5	0.795E-03	0.17E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 12.00
DATA FOR PEAK AT 13.20 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.6	0.537E+01	0.16E+00
12.3	0.463E+01	0.12E+00
13.7	0.300E+01	0.66E-01
15.2	0.124E+01	0.51E-01
17.0	0.112E+01	0.25E-01
18.5	0.651E+00	0.17E-01
20.6	0.273E+00	0.11E-01
20.6	0.327E+00	0.15E-01
22.9	0.175E+00	0.40E-02
25.4	0.525E-01	0.21E-02
25.4	0.691E-01	0.32E-02
30.3	0.113E-01	0.10E-02
38.0	0.166E-02	0.29E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 13.00

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.3	0.290E+00	0.32E-01
6.0	0.390E+00	0.38E-01
8.9	0.347E+00	0.34E-01
11.3	0.122E+00	0.14E-01
11.3	0.193E+00	0.15E-01
14.2	0.271E-01	0.31E-02
17.7	0.117E-01	0.15E-02
21.2	0.100E-01	0.11E-02
24.6	0.525E-02	0.56E-03
29.2	0.143E-02	0.32E-03
35.0	0.429E-03	0.14E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 13.00

DATA FOR PEAK AT 4.44 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.3	0.132E+01	0.61E-01
6.0	0.157E+01	0.68E-01
8.9	0.169E+01	0.67E-01
11.3	0.863E+00	0.31E-01
11.3	0.865E+00	0.30E-01
14.2	0.314E+00	0.97E-02
17.7	0.131E+00	0.46E-02
21.2	0.543E-01	0.21E-02
24.6	0.236E-01	0.99E-03
29.2	0.447E-02	0.43E-03
35.0	0.103E-02	0.13E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 13.00

DATA FOR PEAK AT 12.70 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.3	0.118E+01	0.12E+00
6.0	0.129E+01	0.12E+00
8.9	0.875E+00	0.93E-01
11.3	0.380E+00	0.44E-01
11.3	0.424E+00	0.41E-01
14.2	0.115E+00	0.12E-01
17.7	0.876E-01	0.81E-02
21.2	0.476E-01	0.29E-02
24.6	0.130E-01	0.11E-02
29.3	0.403E-02	0.72E-03
35.0	0.122E-02	0.20E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 13.00

DATA FOR PEAK AT 14.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.3	0.148E+01	0.16E+00
6.0	0.128E+01	0.13E+00
9.0	0.144E+01	0.14E+00
11.3	0.778E+00	0.65E-01
11.3	0.776E+00	0.67E-01
14.2	0.372E+00	0.12E-01
17.7	0.167E+00	0.11E-01
21.2	0.686E-01	0.35E-02
24.6	0.238E-01	0.15E-02
29.3	0.463E-02	0.93E-03
35.0	0.105E-02	0.36E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 13.00

DATA FOR PEAK AT 15.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.3	0.165E+01	0.15E+00
6.0	0.139E+01	0.14E+00
9.0	0.114E+01	0.70E-01
11.3	0.540E+00	0.58E-01
11.3	0.570E+00	0.69E-01
14.2	0.147E+00	0.29E-01
17.7	0.130E+00	0.30E-01
21.2	0.721E-01	0.90E-02
24.6	0.165E-01	0.14E-02
29.3	0.431E-02	0.97E-03
35.0	0.128E-02	0.40E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 13.00

DATA FOR PEAK AT 16.10 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
4.3	0.242E+01	0.17E+00
6.0	0.233E+01	0.16E+00
9.0	0.170E+01	0.13E+00
11.3	0.602E+00	0.57E-01
11.3	0.684E+00	0.54E-01
14.2	0.181E+00	0.16E-01
17.7	0.146E+00	0.11E-01
21.2	0.822E-01	0.53E-02
24.6	0.247E-01	0.49E-02
29.3	0.956E-02	0.96E-03
35.0	0.905E-03	0.20E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.441E+00	0.19E-01
11.8	0.898E-01	0.10E-01
14.7	0.797E-01	0.51E-02
17.8	0.555E-01	0.19E-02
19.9	0.178E-01	0.12E-02
22.0	0.876E-02	0.85E-03
24.4	0.351E-02	0.44E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 5.22 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.760E+00	0.41E-01
11.8	0.151E+00	0.38E-01
14.7	0.922E-01	0.72E-02
17.8	0.535E-01	0.31E-02
19.9	0.260E-01	0.14E-02
22.0	0.856E-02	0.85E-03
24.4	0.689E-02	0.18E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 6.18 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.142E+01	0.56E-01
11.8	0.287E+00	0.58E-01
14.7	0.159E+00	0.92E-02
17.8	0.936E-01	0.41E-02
19.9	0.347E-01	0.16E-02
22.0	0.138E-01	0.11E-02
24.4	0.113E-01	0.25E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00
DATA FOR PEAK AT 7.20 MEV

THETA(CN) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.650E+00	0.40E-01
11.8	0.325E+00	0.51E-01
14.7	0.140E+00	0.10E-01
17.8	0.487E-01	0.32E-02
19.9	0.158E-01	0.14E-02
22.0	0.739E-02	0.94E-03
24.4	0.610E-02	0.17E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00
DATA FOR PEAK AT 10.42 MEV

THETA(CN) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.325E+00	0.61E-01
11.8	0.279E+00	0.58E-01
14.7	0.130E+00	0.14E-01
17.8	0.554E-01	0.42E-02
19.9	0.255E-01	0.23E-02
22.0	0.112E-01	0.22E-02
24.4	0.411E-02	0.83E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 10.87 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.357E+00	0.62E-01
11.8	0.241E+00	0.49E-01
14.7	0.149E+00	0.15E-01
17.8	0.265E-01	0.35E-02
19.9	0.128E-01	0.19E-02
22.0	0.202E-02	0.15E-02
24.4	0.206E-02	0.70E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 12.21 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.800E+00	0.79E-01
11.8	0.631E+00	0.52E-01
14.7	0.254E+00	0.18E-01
17.8	0.824E-01	0.84E-02
19.9	0.254E-01	0.24E-02
22.0	0.127E-01	0.19E-02
24.4	0.443E-02	0.70E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 13.59 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.4	0.180E+01	0.11E+00
11.8	0.942E+00	0.56E-01
14.7	0.418E+00	0.24E-01
17.8	0.122E+00	0.11E-01
19.9	0.307E-01	0.27E-02
22.0	0.177E-01	0.22E-02
24.4	0.754E-02	0.89E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 16.00

DATA FOR PEAK AT 8.84 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
11.8	0.507E-01	0.12E-01
14.7	0.388E-01	0.85E-02
17.8	0.134E-01	0.39E-02
19.9	0.430E-02	0.12E-02
22.0	0.314E-02	0.79E-03
24.4	0.953E-03	0.32E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
6.1	0.845E-01	0.11E-01
11.3	0.182E-01	0.31E-02
14.0	0.448E-02	0.12E-02
19.0	0.120E-02	0.37E-03
21.1	0.586E-03	0.20E-03
23.4	0.408E-03	0.10E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 1.37 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.293E+00	0.21E-01
11.3	0.920E-01	0.65E-02
14.0	0.376E-01	0.32E-02
19.0	0.516E-02	0.64E-03
21.1	0.254E-02	0.33E-03
23.4	0.168E-02	0.18E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 4.12 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.290E+00	0.21E-01
11.3	0.103E+00	0.69E-02
14.0	0.433E-01	0.32E-02
19.0	0.107E-01	0.92E-03
21.1	0.364E-02	0.39E-03
23.4	0.247E-02	0.23E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 5.27 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.581E-01	0.11E-01
11.3	0.336E-01	0.41E-02
14.0	0.202E-01	0.22E-02
19.0	0.295E-02	0.55E-03
21.1	0.111E-02	0.20E-03
23.4	0.306E-03	0.10E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 6.01 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.182E+00	0.16E-01
11.3	0.923E-01	0.69E-02
14.0	0.393E-01	0.32E-02
19.0	0.442E-02	0.64E-03
21.1	0.189E-02	0.26E-03
23.4	0.102E-02	0.15E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 7.66 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
6.1	0.317E+00	0.32E-01
11.3	0.203E+00	0.10E-01
14.0	0.105E+00	0.62E-02
19.0	0.159E-01	0.13E-02
21.1	0.690E-02	0.59E-03
23.4	0.260E-02	0.31E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 8.36 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
6.1	0.208E+00	0.29E-01
11.3	0.906E-01	0.82E-02
14.0	0.276E-01	0.45E-02
19.0	0.562E-02	0.10E-02
21.1	0.338E-02	0.52E-03
23.4	0.158E-02	0.28E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 9.44 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.515E+00	0.45E-01
11.3	0.260E+00	0.13E-01
14.0	0.128E+00	0.80E-02
19.0	0.238E-01	0.18E-02
21.1	0.931E-02	0.98E-03
23.4	0.425E-02	0.41E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 10.12 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.169E+00	0.42E-01
11.3	0.371E-01	0.82E-02
14.0	0.241E-01	0.19E-01
19.0	0.571E-02	0.11E-02
21.1	0.319E-02	0.65E-03
23.4	0.150E-02	0.31E-03
27.9	0.262E-03	0.15E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 10.63 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.269E+00	0.50E-01
11.3	0.122E+00	0.12E-01
19.0	0.105E-01	0.17E-02
21.1	0.514E-02	0.85E-03
23.4	0.222E-02	0.51E-03
27.9	0.931E-03	0.23E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 11.05 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.644E+00	0.63E-01
11.3	0.191E+00	0.13E-01
14.0	0.762E-01	0.87E-02
19.0	0.135E-01	0.27E-02
21.1	0.651E-02	0.12E-02
23.4	0.107E-02	0.48E-03
27.9	0.553E-03	0.20E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 11.99 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.364E+00	0.61E-01
11.3	0.129E+00	0.16E-01
14.0	0.448E-01	0.65E-02
19.0	0.103E-01	0.18E-02
21.1	0.241E-02	0.65E-03
23.4	0.196E-02	0.38E-03
27.9	0.320E-03	0.17E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00

DATA FOR PEAK AT 12.39 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.261E+00	0.53E-01
11.3	0.893E-01	0.13E-01
14.0	0.368E-01	0.10E-01
19.0	0.111E-01	0.18E-02
21.1	0.605E-02	0.11E-02
23.4	0.150E-02	0.54E-03
27.9	0.815E-03	0.20E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 13.30 MEV

THETA(CM) (DEGREES) =====	CROSS SECTION (UB/SR) =====	ERROR (UB/SR) =====
6.1	0.369E+00	0.48E-01
11.3	0.141E+00	0.12E-01
14.0	0.545E-01	0.85E-02
19.0	0.150E-01	0.19E-02
21.1	0.586E-02	0.10E-02
23.4	0.186E-02	0.56E-03
27.9	0.582E-03	0.26E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 14.13 MEV

THETA(CM) (DEGREES) =====	CROSS SECTION (UB/SR) =====	ERROR (UB/SR) =====
6.1	0.211E+00	0.48E-01
11.3	0.135E+00	0.13E-01
14.0	0.615E-01	0.92E-02
19.0	0.151E-01	0.20E-02
21.1	0.371E-02	0.10E-02
23.4	0.380E-02	0.51E-03
27.9	0.728E-03	0.26E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 15.17 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.1	0.264E+00	0.45E-01
11.3	0.166E+00	0.12E-01
14.0	0.757E-01	0.62E-02
19.0	0.110E-01	0.26E-02
21.1	0.108E-01	0.14E-02
23.4	0.155E-02	0.38E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 15.68 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.1	0.333E+00	0.50E-01
11.3	0.106E+00	0.11E-01
14.0	0.550E-01	0.60E-02
19.0	0.133E-01	0.27E-02
21.1	0.690E-02	0.14E-02
23.4	0.183E-02	0.43E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 25.00
DATA FOR PEAK AT 16.56 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.1	0.224E+00	0.50E-01
11.3	0.119E+00	0.13E-01
14.0	0.724E-01	0.10E-01
19.0	0.176E-01	0.25E-02
21.1	0.840E-02	0.16E-02
23.4	0.214E-02	0.46E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.901E+00	0.77E-01
11.2	0.252E+00	0.21E-01
13.9	0.193E+00	0.13E-01
15.7	0.788E-01	0.66E-02
16.9	0.481E-01	0.39E-02
18.9	0.209E-01	0.28E-02
20.9	0.130E-01	0.19E-02
23.2	0.579E-02	0.69E-03
27.7	0.965E-03	0.21E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 0.80 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.256E+00	0.46E-01
11.2	0.159E+00	0.18E-01
13.9	0.122E+00	0.10E-01
15.7	0.572E-01	0.56E-02
16.9	0.312E-01	0.34E-02
18.9	0.168E-01	0.25E-02
20.9	0.117E-01	0.19E-02
23.2	0.385E-02	0.56E-03
27.7	0.965E-03	0.21E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 2.46 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.246E+00	0.61E-01
11.2	0.410E-01	0.96E-02
13.9	0.142E-01	0.43E-02
15.7	0.141E-01	0.38E-02
16.9	0.658E-02	0.21E-02
18.9	0.226E-02	0.75E-03
20.9	0.361E-02	0.11E-02
23.2	0.212E-02	0.35E-03
27.7	0.678E-03	0.13E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 2.89 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.1	0.123E+01	0.12E+00
11.2	0.521E+00	0.33E-01
13.9	0.260E+00	0.18E-01
15.7	0.121E+00	0.88E-02
16.9	0.938E-01	0.53E-02
18.9	0.450E-01	0.50E-02
20.9	0.225E-01	0.31E-02
23.2	0.119E-01	0.74E-03
27.7	0.229E-02	0.52E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 4.20 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
6.1	0.348E+00	0.92E-01
11.2	0.546E-01	0.18E-01
13.9	0.101E+00	0.22E-01
15.7	0.844E-01	0.72E-02
16.9	0.147E-01	0.75E-02
18.9	0.166E-01	0.33E-02
20.9	0.583E-02	0.19E-02
23.2	0.221E-02	0.69E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 4.57 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.532E+00	0.97E-01
11.2	0.317E+00	0.30E-01
13.9	0.115E+00	0.22E-01
15.7	0.522E-01	0.59E-02
16.9	0.348E-01	0.79E-02
18.9	0.118E-01	0.30E-02
20.9	0.361E-02	0.17E-02
23.2	0.182E-02	0.69E-03
27.7	0.756E-03	0.21E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 5.37 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.553E+00	0.72E-01
11.2	0.226E+00	0.23E-01
13.9	0.962E-01	0.12E-01
15.7	0.488E-01	0.12E-01
16.9	0.344E-01	0.41E-02
18.9	0.249E-01	0.33E-02
20.9	0.777E-02	0.17E-02
23.2	0.307E-02	0.69E-03
27.7	0.965E-03	0.23E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 6.57 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.374E+00	0.92E-01
11.2	0.169E+00	0.38E-01
15.7	0.422E-01	0.11E-01
16.9	0.288E-01	0.58E-02
18.9	0.241E-01	0.45E-02
20.9	0.694E-02	0.22E-02
23.2	0.679E-02	0.17E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 7.18 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
6.1	0.112E+01	0.14E+00
11.2	0.106E+01	0.70E-01
13.9	0.544E+00	0.41E-01
15.7	0.256E+00	0.14E-01
16.9	0.162E+00	0.10E-01
18.9	0.688E-01	0.11E-01
20.9	0.211E-01	0.28E-02
23.2	0.329E-02	0.61E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 8.32 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
6.1	0.162E+01	0.16E+00
11.2	0.522E+00	0.57E-01
13.9	0.229E+00	0.33E-01
15.7	0.123E+00	0.12E-01
16.9	0.666E-01	0.83E-02
18.9	0.274E-01	0.10E-01
20.9	0.119E-01	0.28E-02
23.2	0.735E-02	0.74E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 9.59 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
6.1	0.353E+00	0.11E+00
11.2	0.239E+00	0.51E-01
13.9	0.186E+00	0.33E-01
15.7	0.847E-01	0.15E-01
16.9	0.630E-01	0.88E-02
18.9	0.314E-01	0.45E-02
20.9	0.944E-02	0.31E-02
23.2	0.549E-02	0.74E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 11.65 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
6.1	0.211E+01	0.20E+00
11.2	0.151E+01	0.92E-01
13.9	0.983E+00	0.60E-01
15.7	0.462E+00	0.28E-01
16.9	0.288E+00	0.16E-01
18.9	0.129E+00	0.70E-02
20.9	0.500E-01	0.42E-02
23.2	0.895E-02	0.13E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00
DATA FOR PEAK AT 14.79 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
*****	*****	*****
11.2	0.372E+00	0.44E-01
13.9	0.224E+00	0.29E-01
15.7	0.149E+00	0.19E-01
16.9	0.628E-01	0.98E-02
18.9	0.261E-01	0.53E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 15.28 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
11.2	0.311E+00	0.43E-01
13.9	0.197E+00	0.27E-01
15.7	0.934E-01	0.17E-01
16.9	0.468E-01	0.86E-02
18.9	0.246E-01	0.50E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 15.74 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
11.2	0.346E+00	0.45E-01
13.9	0.260E+00	0.28E-01
15.7	0.105E+00	0.18E-01
16.9	0.451E-01	0.85E-02
18.9	0.269E-01	0.53E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 28.00

DATA FOR PEAK AT 16.25 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
11.2	0.350E+00	0.49E-01
13.9	0.170E+00	0.26E-01
15.7	0.434E-01	0.16E-01
16.9	0.331E-01	0.81E-02
18.9	0.138E-01	0.50E-02

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00
DATA FOR PEAK AT 0.00 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
5.9	0.853E-01	0.97E-02
11.0	0.471E-01	0.53E-02
13.6	0.294E-01	0.40E-02
15.4	0.170E-01	0.19E-02
18.5	0.693E-02	0.75E-03
20.4	0.316E-02	0.43E-03
22.7	0.137E-02	0.16E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00
DATA FOR PEAK AT 2.68 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
5.9	0.134E+00	0.18E-01
11.0	0.836E-01	0.77E-02
13.6	0.481E-01	0.52E-02
15.4	0.321E-01	0.24E-02
18.5	0.889E-02	0.10E-02
20.4	0.524E-02	0.10E-02
22.7	0.188E-02	0.21E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 3.79 MEV

THETA(CM) (DEGRFES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
5.9	0.132E+00	0.77E-01
11.0	0.725E-01	0.91E-02
13.6	0.422E-01	0.36E-02
15.4	0.308E-01	0.31E-02
18.5	0.721E-02	0.75E-03
20.4	0.337E-02	0.93E-03
22.7	0.177E-02	0.26E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 4.44 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
5.9	0.757E-01	0.76E-01
11.0	0.279E-01	0.77E-02
13.6	0.637E-02	0.20E-02
15.4	0.872E-02	0.23E-02
18.5	0.449E-02	0.66E-03
20.4	0.932E-03	0.79E-03
22.7	0.264E-03	0.24E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 5.40 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
5.9	0.332E+00	0.47E-01
11.0	0.205E+00	0.13E-01
13.6	0.108E+00	0.80E-02
15.4	0.612E-01	0.34E-02
18.5	0.150E-01	0.17E-02
20.4	0.516E-02	0.79E-03
22.7	0.177E-02	0.29E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 6.39 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
5.9	0.214E+00	0.31E-01
11.0	0.129E+00	0.12E-01
13.6	0.469E-01	0.64E-02
15.4	0.349E-01	0.36E-02
18.5	0.108E-01	0.10E-02
20.4	0.481E-02	0.86E-03
22.7	0.201E-02	0.34E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 7.26 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
5.9	0.106E+00	0.19E-01
11.0	0.370E-01	0.82E-02
13.6	0.374E-01	0.72E-02
15.4	0.108E-01	0.95E-02
18.5	0.384E-02	0.94E-03
20.4	0.301E-02	0.10E-02
22.7	0.122E-02	0.42E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 7.78 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
=====	=====	=====
5.9	0.222E+00	0.37E-01
11.0	0.170E+00	0.45E-01
13.6	0.513E-01	0.80E-02
15.4	0.245E-01	0.98E-02
18.5	0.393E-02	0.10E-02
20.4	0.179E-02	0.10E-02
22.7	0.114E-02	0.42E-03

800 MEV (P,D) DATA TABLE

TARGET MASS = 40.00

DATA FOR PEAK AT 10.92 MEV

THETA(CM) (DEGREES)	CROSS SECTION (UB/SR)	ERROR (UB/SR)
5.9	0.240E+01	0.32E+00
11.0	0.104E+01	0.12E+00
13.6	0.538E+00	0.74E-01
15.4	0.219E+00	0.62E-01
18.5	0.277E-01	0.51E-02
22.7	0.927E-02	0.31E-02