Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

TITLE: NUCLEAR MODELS RELEVANT TO EVALUATION

AUTHOR(S): E. D. Arthur, T-DC
M. B. Chadwick, T-2 Postdoc
G. M. Hale, T-2
P. G. Young, T-2

SUBMITTED TO The International Conference on Nuclear Data for Science and
Technology, Julich, Germany, May 13-17, 1991

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

Los Alamos Los Alamos National Laboratory
Los Alamos, New Mexico 87545

LA-UR-91-1662

NUCLEAR MODELS RELEVANT TO EVALUATION

E. D. Arthur, M. B. Chadwick, G. M. Hale, and P. G. Young

Theoretical Division, MS B243
Los Alamos National Laboratory
Los Alamos, New Mexico 87545, USA

Abstract: The widespread use of nuclear models continues in the creation of data evaluations. The reasons include extension of data evaluations to higher energies, creation of data libraries for isotopic components of natural materials, and production of evaluations for radioactive target species. In these cases, experimental data are often sparse or nonexistent. As this trend continues, the nuclear models employed in evaluation work move towards more microscopically-based theoretical methods, prompted in part by the availability of increasingly powerful computational resources.

Advances in nuclear models applicable to evaluation will be reviewed. These include advances in optical model theory, microscopic and phenomenological state and level density theory, unified models that consistently describe both equilibrium and nonequilibrium reaction mechanisms, and improved methodologies for calculation of prompt radiation from fission.

(Keywords: Nuclear theory, nuclear models, R-matrix, optical model, statistical theory, fission theory, preequilibrium theory, nuclear data evaluation)

Introduction

Reviews by Uhl [1], Gardner [2], and Young [3] at earlier international nuclear data conferences have stressed the increased and important role that nuclear models play in the provision of data for a variety of applications. This trend continues with theoretical calculations representing a major component required for development of evaluated data files for ENDF-VI, JENDL, JEF, and the BROND libraries. These data systems have expanded their scope beyond the incorporation of neutron-induced reaction data for energies below 20 MeV. Extensions include provision of reaction data for incident charged-particles and higher energy data. In these extensions, nuclear models have played a significant and major role. Applications such as nuclear waste transmutation, radiation protection for manned space exploration, and radiotherapy involving energetic neutron beams all require expanded nuclear data bases [4-6]. Nuclear models will be used to produce major parts of the databases required for these new applications.

The application of nuclear models to data evaluation has largely involved use of a collection of models—optical, statistical, and preequilibrium—where the user compromises between the complexity of microscopically-based formulations and the simplicity (and hence calculational speed) of more phenomenological approaches. Much of the nuclear model effort directed towards evaluation uses results and insights from fundamental nuclear theory development to guide the choice of approximations required for applications. Determination of parameter systematics is a major effort area that aims at improving the interpolative or predictive capability of phenomenologically-based models. Finally, it is now a universal practice in model applications to simultaneously analyze data for a variety of reaction channels for the system of interest, as well as data that may be available for nearby nuclei or for other incident particles. Achieving an overall consistency in calculated results helps ensure the reliability of extrapolated results and guarantees that fundamental conservation principles are adhered to in the resulting evaluated data files.

This paper briefly reviews the main nuclear models used for applications and will indicate areas of major development and application activity. Discussion of more fundamental theoretical development is covered in other review papers at this conference [7-8]. Also the primary emphasis will be on model development and application during the period since the 1988 Mito Conference.

The areas to be covered include examples of models appropriate for the calculation of properties of reactions

involving light nuclei as targets along with models describing reactions involving medium to heavy nuclei. This includes optical model development of interest for application along with models that make up components of statistical reaction theories (level density and gamma-ray strength formalisms). Nonequilibrium reaction theories (classical preequilibrium and quantum-mechanical models) will be discussed along with efforts aimed at creation of unified reaction models for applications. Finally, a brief description of recent developments in models used for calculation of prompt radiation emission from fission will be provided.

Methods for Light-Element Evaluations

The most successful and frequently used method for light-element evaluations is R-matrix theory. The theory is ideally suited for describing the resonances that are usually seen in light-element reactions, and at the same time it builds in the correct energy dependence of the transition (T) matrix elements at low energies by making explicit use of the solutions for the external (long-ranged) parts of the interaction (Coulomb and angular momentum barriers, etc.). Thus, the method gives reliable extrapolations to low energies for both neutron- and charged-particle-induced reactions.

Applications of the theory usually involve parametrizing the R matrix for a light system as the channel-space spectral expansion,

$$R_{cc'}^B(E) = \sum_{\lambda} \frac{\gamma_{c\lambda} \gamma_{c'\lambda}^T}{E_{\lambda} - E} \quad (1)$$

in terms of the reduced-width amplitudes $\gamma_{c\lambda}$ and eigenenergies E_{λ} . The $\gamma_{c\lambda}$ are channel-surface projections of eigenfunctions $|\lambda\rangle$ that are regular solutions of the Schrödinger equation governing the motion of the particles at small distances and satisfying at the channel radius $r_c = a_c$ real, energy-independent boundary values B_c . Using standard methods, it is convenient to transform Eq. (1) to the outgoing-wave R matrix, $R_{cc'}^L(E)$, having boundary values given by the logarithmic derivatives L_c of the external outgoing wave solutions O_c at $r_c = a_c$. From the elements of the multichannel partial-wave transition matrix,

$$T_{cc'} = O_c^{-1} R_{cc'}^L O_{c'}^{-1} = O_c^{-1} \text{Im}(O_c) \delta_{cc'}, \quad (2)$$

can be calculated observables (cross sections, polarizations, etc.) for any two-body reaction in the system described by the R matrix in Eq. (1). Its parameters, $\gamma_{\alpha\lambda}$, E_{α} , and sometimes a_{α} , are usually determined by fitting experimental data for all the reactions in the system. Some evaluators consider only a subset of the possible reactions by using a variation of the method that involves a complex, reduced R matrix.

The full method was used in several of the light-element evaluations for ENDF/B-VI, including the standards evaluations for ^1H , ^3He , ^6Li , and ^{10}B . Other light-element ENDF evaluations that used R-matrix methods were those for ^{12}C [9], ^{14}N , and ^{16}O [10].

An example that illustrates the wide-ranging utility of this approach and the ability to incorporate fundamental theoretical constraints is the R-matrix description of the $A=4$ reactions being done at Los Alamos. All possible reactions involving the channels $n+^3\text{H}$, $n+^3\text{He}$, $p+^3\text{H}$, $p+^3\text{He}$, and $d+d$ are described using a single set of Coulomb-corrected, charge-independent R-matrix parameters. Neutron cross sections from this analysis have been used in the ^3He evaluation and will be used in a future update of the ^3He evaluation where the previous low-energy total cross sections differ notably from the present calculations [11] and data [12].

A novel application of the analysis has been to compare its results [13] with measurements being done for the $d+d$ muon-catalyzed fusion (μCF) reactions. Room temperature μCF measurements [14] give the surprising result that the branching ratio for the P-wave part of the $d+d$ reactions favors the $n+^3\text{He}$ branch over that for $p+^3\text{H}$ by about 40%. At lower temperatures, where the molecular transitions allow increasing amounts of S-wave formation between the two deuterons, the branching ratio has recently been observed [15] to decrease toward unity. The P-wave branching ratio calculated from the analysis is 1.43, while that for the S-waves is 0.886, giving excellent agreement with the measured room-temperature number, and also accounting qualitatively for the decrease in the branching ratio toward unity as the S-wave admixture increases at lower temperatures. Thus, the analysis successfully accounts for all the new experimental information (including the surprises) about the $d+d$ reactions that has come from this exciting new field, confirming by individual partial waves the reliability of R-matrix extrapolations to low energies.

One would like to connect the parameters obtained in R-matrix analyses with more microscopic models of nuclear reactions that start, e.g. from the nucleon-nucleon interaction. An effort to make this connection with shell-model (SM) quantities was reported by Knox [16]; this work has been continued by Ressler [17]. An unproved prescription that relates the SM energies to the correct boundary conditions and takes into account the renormalization and nonorthogonality of the SM states due to the finite extent of the nuclear region has been given by Hale [18]. Bolsterli [19] has developed a continuum SM approach that has many similarities to R-matrix theory, especially when the interactions are approximated by delta functions at the nuclear surface. By adjusting the levels and interactions phenomenologically he is able to match closely with fewer parameters the S-wave T-matrix elements that Hale *et al.* [10] obtained from their R-matrix description of the $n+^{14}\text{N}$ reactions.

Optical Model

Good optical model analyses are essential for thorough nuclear data evaluations. In the period since the Mito Conference, much of this activity has dealt with the use of dispersion relationships in optical model analyses. Because a review paper on the dispersive optical model is being given at this conference [8], we will limit our discussion to three examples of optical model application in current evaluations.

A number of optical model studies have been performed at Argonne National Laboratory (ANL) that utilize dispersion relationships. Isotopes studied include ^{51}V ,

^{59}Co , ^{89}Y , ^{93}Nb , ^{115}In , and ^{209}Bi , and several of the analyses and accompanying data sets form the basis for ENDF/B-VI evaluations. In such analyses the real (V) and imaginary (W) optical model potentials are related through a dispersive correction term [20],

$$\Delta V(r, E) = \frac{1}{\pi} P \int_{-\infty}^{\infty} dE' \frac{W(r, E')}{E' - E}, \quad (3)$$

where P signifies the principal-value integral. The ANL studies usually involve fitting experimental neutron total and scattering data in the energy range $E_n \sim 0.5 - 10$ MeV, either with a spherical optical potential that includes a dispersion relation [21], or a standard spherical potential that is then interpreted using dispersion relationships and the method of moments [22].

Several of these analyses, when coupled with bound-state data, give clear evidence of Fermi surface anomalies, as illustrated by the $n + ^{51}\text{V}$ [23] example in Fig. 1. Here the volume integral/nucleon of the real potential strength obtained using dispersion relations (curve) is compared to J_v values derived from experimental data (circles). While the data for $E_n > 0$ are consistent (approximately) with linear behavior, the values of J_v needed to reproduce the single-particle and hole bound-state energies shows characteristic anomalous behavior.

Fig. 1. The volume integral per nucleon of the real potential strength vs neutron energy for $n + ^{51}\text{V}$ reactions.

The ANL optical-statistical analyses also highlight the difficulties associated with use of global and even regional optical model potentials at lower incident energies. The strength and/or diffuseness parameters associated with the imaginary potential are found to be strongly influenced by the level structure and properties of individual nuclei.

Coupled-channel optical model analyses for permanently deformed nuclei continue to be an essential tool in data evaluations. An example of coupled-channel calculations of ground state rotational bands is offered by the ENDF/B-VI analyses of neutron reactions on ^{235}U , ^{238}U , ^{237}Np , and ^{239}Pu [24]. For these calculations earlier "SPRT" analyses [25] were adjusted to improve agreement with σ_{tot} data up to 20 MeV and were then used to calculate

neutron transmission coefficients for detailed Hauser-Feshbach statistical theory analyses as well as shape elastic and direct (n,n') cross sections. Very similar potentials were utilized for the four target nuclei. The ^{237}Np potential was obtained prior to release of new total cross section data [26] and is based on a simple interpolation of the neighboring potentials.

A third example of optical model phenomenology illustrates the movement to better characterize optical model potentials at higher energies. In this study Kozack and Madland [27] adapted and combined Dirac phenomenology [28] with a relativistic generalization [29] of the Lane model [30] to successfully fit ^{208}Pb neutron total cross sections and proton elastic scattering data between 95 and 300 MeV. This analysis used standard Lorentz potentials (scalar and vector, real and imaginary). Several different forms for the energy dependence of the potential strengths and two forms for the isoscalar and isovector potentials were tested in the analysis. Best results were obtained with strengths parameterized in the form

$$U_0(T) = A + \alpha \ln(T) \pm \eta B \quad (4)$$

where T is the projectile kinetic energy, $\eta = (N-Z)/A$, the plus (minus) sign refers to neutrons (protons), and the parameters A , B , and α were determined by the data fitting procedure. A total of 20 parameters were determined in the analysis, which is similar to the number of parameters involved in more usual Schrödinger analyses. Better fits were obtained to the higher energy data with the Dirac formalism. Kozack and Madland's results for the $p + ^{208}\text{Pb}$ reaction cross section and for the $n + ^{208}\text{Pb}$ total cross section compare well with experimental data. One outcome of this analysis is the observation that dramatic differences occur between the behavior of protons and neutrons elastically scattered from ^{208}Pb at incident energies in the neighborhood of 100 MeV, especially differential scattering cross sections and analyzing powers [31]. This prediction awaits experimental verification.

Nuclear Level Densities

Nuclear level densities are central to Hauser-Feshbach [32] and Weisskopf-Ewing [33] models used to determine equilibrium contributions to cross sections. The importance of realistic level density information in applied and basic calculations is reflected in the continuing body of literature aimed at both phenomenological and microscopic models. Application calculations continue to rely principally on simple phenomenological models, e.g., the Gilbert-Cameron (GC) [34] or the Back Shifted Fermi Gas (BSFG) [35]. These models feature unphysical assumptions and uncertainties in parametrization in areas pertaining to shell effects, equidistant spacing of particle levels, excitation energy dependence of the spin cutoff parameter and parity factors. For this reason much recent effort has focused on use of more sophisticated phenomenological models [36] or to use of microscopic Fermi gas models employing realistic single particle schemes [37].

Arthur and Guenther [38] explored the sensitivity of calculated particle emission spectra to level density models and associated parameters used. The $A=60$ region was chosen because of the existence of experimental spectral data for neutron, proton, and alpha-particle emission. BSFG model results were calculated with recent parameter improvements due to Ivascu *et al.* [39] and Zhuang *et al.* [40] while GC level densities were determined using systematics for the level density parameter [41] coupled with adjustment of the constant temperature portion of the GC model to fit low-lying discrete level data. Figure 2-a illustrates differences in ^{60}Ni level densities calculated with the models and parameters shown, while Fig. 2-b illustrates the rather negligible effect of these level density differences on calculated neutron emission spectra for $^{60}\text{Ni}(n,n')$. More sensitivity was found for emission channels (proton and alpha particles) that are minor

in magnitude (as compared with neutron emission). This occurs because in Hauser-Feshbach calculations the cross section is dominated by the large s-wave partial width for particle emission to the final width, where the neutron emission channel dominates calculation of both partial and total widths; thus, large level density changes there produce relatively little impact on the result.

Fig. 2-a. Level densities for ^{60}Ni computed using the GC model with Cook parameters (solid curve), GC with Zhuang parameters (dotted curve), and the BSFG with Ivascu parameters (dashed curve).

Fig. 2-b. Calculated $^{60}\text{Ni}(n,n')$ spectra using these level densities.

In many application calculations, emphasis is placed on the reliability of level density extrapolations to mass regions where little or no data exist. Analyses by Reffo [42] and Iijima [43], e.g., have produced sets of level density parameters optimized using relevant experimental data over a limited range of masses. These parameters exhibit significant differences when compared to straightforward applications of Gilbert-Cameron systematics for the Fermi gas parameter a and U_x , the matching energy between constant temperature and Fermi gas forms used in this model. These parameters exhibit correlations with shell effects, not accounted for in more general parameter systematics. Such studies yield

uncertainty estimates of the general magnitude $\Delta a = \pm 15\%$, $\Delta U_x = \pm 15\%$, and uncertainties in the spin cutoff parameter σ^2 of $\pm 40\%$. Calculated (n,γ) cross sections (the area of emphasis in Ref. [42]) are most sensitive to changes in the FG and matching energy. Uncertainties in these parameters translate into cross section variations about 2 to 3 times larger.

The GC and BSFG level density models commonly used in data application calculations do not properly account for the damping out of shell effects at higher excitation energies. Microscopic calculations predict such effects; they are included in more modern phenomenological level density expression, e.g., such as those due to Schmidt [37] and Ignatyuk [36]. The model derived by Schmidt is used in the STAPRE model code version described in Ref. [44] and results from microscopic calculations that employ realistic single particle levels and pairing corrections. These produce a simple relation between the entropy of a system, S , and the level density

$$\omega(E) = 1/(2\pi)^{3/2} (D)^{-1/2} \exp(S), \quad (5)$$

where

$$S = 2[a(E) + \delta U_f(E) \delta P_h(E)]^{1/2} \text{ and } D = 18\pi^{-1/4} a^{1/2} E^{5/2}. \quad (6)$$

The high energy asymptotic level density parameter is determined from an expression for a Fermi gas with a diffuse surface. The difference between experimental and the liquid drop masses is used to determine the sum of δU plus δP , while δP is determined from standard pairing energy assumptions. The energy dependences of the shell effects and effective pairing corrections are described by the weighting functions $f(E) = (1 - \exp(-\gamma E))$ where $\gamma = 2.5aA^{-1/3}$. The function $h(E)$ describes the transition occurring in the pairing energy at the critical energy E_c at which a phase transition occurs from a superconducting state to a normal one.

The phenomenological level density model of Ignatyuk also accounts for disappearance of shell effects through use of an energy dependent Fermi gas parameter given by

$$a(E) = a_{LDM} (1 + \delta W(1 - \exp(-\gamma E)/E)), \quad (7)$$

where the high energy asymptotic value

$$a_{LDM} = (0.154 + 6.3 \times 10^{-5} A) \text{ and } \gamma = 0.054. \quad (8)$$

The experimental shell correction δW is again given by the difference between experimental and liquid drop masses.

The Ignatyuk level density model has been implemented into two codes, STAPRE [45] and GNASH [46], which are often used in data evaluation calculations. In the STAPRE implementation, the lower excitation energy region is described using a BSFG and a logarithmic interpolation is made to join results from this model with those from the Ignatyuk expressions at higher energies. In the GNASH implementation, the GC constant temperature form is used to describe low-lying regions of excitation energies. This expression is then joined to the Ignatyuk expression at a matching energy U_x in a manner described earlier.

Results calculated using the Ignatyuk and GC level density models have recently been compared [47] with $^{204-208}\text{Pb}(n,x\gamma)$ data. The data were measured for incident neutron energies up to 100 MeV by Haight [48]. The comparison, made for specific gamma rays from $^{208}\text{Pb}(n,x\gamma)$ measurements, provides a unique test of shell effects and dissipation at higher excitation energies. Figure 3 compares the calculated excitation function for the $^{208}\text{Pb}(n,3n\gamma)^{206}\text{Pb}$ reaction using these two level density models. Because of the dissipation of shell effects, increasing the level density in the residual system of interest (here ^{206}Pb) and in competing residual systems (^{205}Pb), the Ignatyuk level density results peak earlier and drop more rapidly than the GC results. This behavior appears to be in better agreement with these data. For nuclei farther away from shell closures, calculated differences in such excitation functions are smaller.

Fig. 3. Excitation functions for the $^{208}\text{Pb}(n,3n\gamma)^{206}\text{Pb}$ reactions using the Ignatyuk (solid curve) and GC (dashed curve) level density models.

Gamma Ray Strength Functions

The use of gamma-ray strength functions provides the preferred mechanism for determination of gamma-ray transmission coefficients in statistical model calculations. Recent studies of gamma-ray strength function systematics include the work of Gardner, Gardner, and Hoff [49], which extend earlier investigations centered primarily around rare earth nuclei to the actinide mass region. They modeled $f_{E1}(E_\gamma)$ using an energy dependent Breit-Wigner line shape, while $f_{M1}(E_\gamma)$ was represented by a constant value as suggested by the Weisskopf single-particle approximation. Comparisons were made with partial gamma-ray width data for ^{232}Th , ^{237}Np , and ^{238}U in statistical model calculations that employed large sets of modeled, discrete sets (> 100 levels) for the daughter nuclei. They found that the $M1/E1$ ratio was the same as earlier values they determined for the rare-earth mass region, but that for these actinide nuclei, the total transition strength was almost thirty percent greater.

Kopecky and Uhl [50] have also studied the impact of models for $E1$ and $M1$ strength functions on the results of statistical model calculations of average radiation widths, radiative neutron capture cross sections, and gamma ray spectra. Several models were compared that reproduced photoabsorption and other data but differed significantly at low gamma-ray energies. They found strong evidence to support an $E1$ strength function model, including an energy dependent spreading width in a Lorentzian form, as well as a non-zero, temperature-dependent limit for the strength function as E_γ approaches zero. The specific form is given by

$$f_{E1}(E_\gamma, T) = 8.68 \times 10^{-8} (\text{mb}^{-1} \text{MeV}^{-1}) \left[\frac{E_\gamma \Gamma(E_\gamma)}{(E_\gamma^2 - E^2)^2 + E_\gamma^2 \Gamma(E_\gamma)^2} + \frac{0.7 \Gamma 4 \pi^2 T^2}{E^3} \right] \sigma_0 \Gamma, \quad (9)$$

where $\Gamma(E_\gamma) = \Gamma[(E_\gamma^2 + 4\pi^2 T^2)/E^2]$.

The rightmost portion of the expression is a term describing $f_{E1}(E_\gamma=0, T)$, as proposed by Kadmenakij [51]. The overall expression is correct in the low-energy form and exhibits little difference with a Lorentzian near the neutron binding energy. It is, however, not valid near the giant resonance peak. Because of the temperature dependence of the expression, it partially violates the Brink-Axel model, [52] which calculates photoabsorption for excited states in the same fashion as for the ground state by assuming a giant resonance built on each excited state.

Data for ^{197}Au , ^{143}Nd , ^{105}Pd , and $^{93}\text{Nb}(n,\gamma)$ reactions were compared with predictions using this gamma-ray strength function form. Such data do not extend to low enough gamma-ray energies to unambiguously support use of this non-zero gamma-ray strength function in the limit of $E_\gamma \rightarrow 0$. Comparisons with data for the total gamma ray width are sensitive to behavior of the gamma-ray strength function in the energy range between zero and a few MeV. In comparisons of gamma-ray strength models to these data, overprediction of total widths was found when the standard Lorentzian strength function was used. The best agreement resulted from use of the generalized Lorentzian form given above. Likewise use of this expression provided consistently better agreement with (n,γ) excitation functions as well as spectral data resulting from capture reactions.

These analyses indicate a preference for the generalized Lorentzian form for spherical nuclei. This form is significantly different than the standard Lorentzian often used in statistical model analyses. Likewise its use implies a partial breakdown of the Brink hypothesis--a principal assumption underlying most statistical model calculations. Because of the potentially important impact on applied calculations, similar analyses should be extended to other mass regions, particularly those involving deformed nuclei.

Preequilibrium Model Development

The study of preequilibrium particle and gamma ray emission continues to be an area of significant development for applied calculations. Here we choose several example areas to illustrate progress and development directions occurring during the interval since Mito. This period has seen the continued development and application of quantum mechanical theories such as those of Feshbach, Kerman, Koonin [53] (FKK model), Tamura, Udagawa, Lenske [54] (TUL model), and currently the multistep reaction model of Nishioka, Verbaarschot, Weidenmüller, and Yoshida [55] (NVW model). Recently Hodgson [56] reviewed the status of multistep compound analyses of (n,nx) , (r,px) , and (p,nx) reactions on medium weight nuclei for incident energies around 14 MeV. There, concurrent analyses of all open channels provided rather stringent tests of the theory as well as important parameters such as optical potential and the strength of the effective interaction used in the FKK model. This analysis illustrated that the FKK theory provides reliable spectral and angular distribution information for important reaction channels in this mass and energy range.

Comparisons of the multistep direct formalisms to higher energy reaction data (spectra and angular distributions) have been made. Examples include analyses of neutron and proton nonelastic scattering data by Marcinkowski [57], Kalka [58], Cowley [59], and recently Scobel [60]. Scobel found that the FKK theory was able to describe (p,n) angular distributions for incident energies up to 160 MeV, whereas the classical hybrid preequilibrium model failed to predict angular distributions correctly because of an overestimate of the first step cross section. An example of these results is presented in Fig. 4.

MSD analyses over extended incident energy ranges provide increased information concerning the energy dependence of the effective interaction strength, V_0 . This parameter is fixed by fitting it to cross section data. Several models are used to describe the interaction potential, with the most realistic involving use of a Yukawa two-body interaction. A series of analyses using this Yukawa potential form has resulted in an energy dependent form given by $V = 31 - 0.15 E$ [56,60]. This form agrees with standard optical model results and applies to both MSC and MSD calculations within the FKK model framework.

Analyses using the other formalisms listed above have been performed. These efforts are often aimed at providing an integrated perspective on the similarities and differences of models for multistep emission processes. For example, in [Ref. 61] Herman and Reffo compare content and assumptions inherent in the FKK and exciton emission

Fig. 4. Comparison of neutron angular distributions for 120- and 160-MeV $^{208}\text{Pb}(p,xn)$ measured by Scobel. Curves are obtained from statistical multistep direct emission calculations.

models. In Ref. [62], Köning and Akkermans explore different nonequilibrium reaction models (FKK, TUL, NVW) and classical exciton models, and they classify them in terms of the assumption of leading particle or residual system statistics. More information concerning quantum mechanical non-equilibrium reaction models can be found in the invited paper by Weidenmüller at this conference.

A second area of activity involves efforts to introduce parameter consistency between equilibrium and nonequilibrium portions of unified reaction models. A focus has been consistent treatments between exciton state densities used in preequilibrium models and total level densities used in equilibrium reaction calculations. Examples include work by Avrigeanu [63] to achieve consistent exciton state and level densities and a two fermion state density analysis by Hennan, Reffo, and Costa [64] using realistic single-particle levels. Similarly, Fu [65] has extended his earlier analysis of unification and simplification of total and partial level densities performed in the framework of the one-fermion model to the case of two-fermion expressions. Pairing interaction effects and spin cutoff expressions were the main focus of the

work. The effect of using the two-fermion model results in determination of preequilibrium state densities is illustrated in Fig. 5 where the ratios $\rho_1(n=2)/\rho_2(n=2)$ and $\rho_1(\text{all } n)/\rho_2(\text{all } n)$ are shown. Here the subscripts refer to the one- and two-fermion expressions and n refers to the exciton configuration. These ratios exhibit very different behavior as a function of the excitation energy U —a situation that is, quite difficult to approximate in preequilibrium Hauser-Feshbach calculations that attempt simple corrections for one- and two-fermion model results. Use of such formalisms achieves consistency in level density models required for compound and precompound reactions description.

Fig. 5. Ratio of one-fermion and two-fermion level and state densities calculated for ^{41}Ca .

Non Equilibrium Gamma-Ray Emission

The emission of gamma-rays via preequilibrium mechanisms has been the subject of much recent work within both semi-classical and quantum-mechanical preequilibrium models. For gamma-rays that are emitted with energies below about 30 MeV, the dominant mechanism is a direct/semi-direct process via the excitation of the Giant Dipole Resonance (GDR). Akkermans and Gruppelaar [66] presented a description of such a process within the exciton model, based upon the application of detailed balance using the Brink-Axel hypothesis, which yields correct emission cross sections in the equilibrium limit. The exciton model description of gamma-ray emission in the GDR region is easy to implement and yields emission cross sections which generally describe the data well; for a full review see [67]. Oblozinsky has proposed [68] a hybrid model for GDR gamma-ray emission that is consistent with Akkerman's exciton model in its application of detailed balance. The exciton model description of GDR gamma rays has also been applied with some success by Cvelbar *et al* [69] to calculate (n,γ) and (p,γ) activation cross sections.

There have been some recent theoretical developments into gamma-ray emission in the GDR region using quantum-mechanical reaction models. The FKK multistep compound theory has been developed by Oblozinsky and Chadwick [70] to include gamma-ray emission. This theory yields the correct Hauser-Feshbach spectrum in its equilibrium limit and is able to account for a considerable fraction of the high-energy gamma-ray cross section. Multistep compound and direct gamma-ray theories are also being developed within the NVW multistep formalism [53].

For photon energies above 30 MeV the two-body emission mechanism becomes important. Two nucleons collide, de-accelerate, and a bremsstrahlung photon is emitted. The two-body nature of this emission led Oblozinsky to present a preequilibrium model of hard photon emission using the quasi-deuteron concept [71]. As in all semi-classical preequilibrium models, this model uses the inverse photoabsorption cross section, applies detailed balance, and is easy to implement in a reaction model code. The success and simplicity of the quasi-deuteron approach to hard photon emission has led to an a collaboration between Los Alamos, Bratislava, Bologna, and Oxford to develop and implement this model in nucleon-induced and heavy-ion reaction codes. Since a thorough understanding of the photoabsorption mechanism is essential for a preequilibrium model of hard photon emission, a photoabsorption model has been developed [72] that describes the effects of Pauli-blocking theoretically and accounts for the quasi-deuteron photoabsorption data well. Only final states that can be reached via linear momentum, as well as energy conservation, were included in the accessible state densities. More refined hard-photon emission quasi-deuteron models are presently being developed by Chadwick and Oblozinsky. Hard photons from the first n - p collision (the $n=1$ preequilibrium stage) dominate the high-energy emission spectrum, and in one such model the $n=1$ theoretical emission rate is given by

$$\lambda_\gamma(\epsilon, \epsilon_\gamma) = \frac{e_\gamma^2 \sigma_{qd}(\epsilon_\gamma)}{4\pi^3 \hbar^3 c^2} \frac{\int \rho(2p1h, \epsilon - \epsilon_\gamma, \mathbf{k} - \mathbf{k}_\gamma) d\Omega_\gamma}{\rho(2p2h, \epsilon_\gamma, \mathbf{k}_\gamma)} \quad (10)$$

where ϵ , $\mathbf{k}$, ϵ_γ and $\mathbf{k}_\gamma$ are the nuclear excitation energy, the incident nucleon momentum, the emitted photon energy and the emitted photon momentum respectively. The $2p1h$ and $2p2h$ state densities are determined within a Fermi-gas model and include linear momentum conservation, and the inverse quasi-deuteron cross section, $\sigma_{qd}(\epsilon_\gamma)$, was taken from the photoabsorption model of Ref [72]. In the numerator we integrate the residual nucleus $2p1h$ state density over all directions of the $2p1h$ momentum, determined by the various photon emission directions Ω_γ . The $2p2h$ density in the denominator is independent of the direction of $\mathbf{k}_\gamma$. In Fig. 6 the photon emission spectrum obtained in this model is compared with data for the reaction $^{197}\text{Au}(p,\gamma)$ induced by 72 MeV protons. It is evident that this model describes the data well except for the lowest energies where photons from more complex preequilibrium stages and the GDR-tail photons (which we have not included) will contribute. It should be noted that a calculation that used the full Fermi-gas state densities, and that ignored linear momentum effects, led to an overprediction of the data by a factor of 2-3 [73].

Fig. 6. Comparison of energetic gamma-ray emission from 72-MeV $\text{Au}(p,\gamma)$ predictions of quasi-deuteron photon emission theory.

Prompt Radiations from Fission

Recent actinide data evaluations for libraries such as ENDF include descriptions of prompt radiation from fission (neutrons, specifically) based upon models such as those developed by Madland and Nix [74]. Likewise, recent higher-energy evaluated data libraries [75] have included prompt neutron emission contributions to total neutron emission spectra for neutron and proton interactions with ^{238}U . In this latter case, prompt neutron and gamma-ray emission were determined using cascade evaporation model calculations of the decay of fission fragments (distributions thereof) produced during nucleon-induced fission [76]. Figure 7 illustrates a reaction chain appropriate for the case proton-induced fission of ^{238}U . Here several points are important. Statistical model calculations of multi-chance fission components of the total fission cross section included both (p,xn) and (p,pxn) paths that populated both uranium and neptunium compound systems. For each compound system undergoing fission, separate model calculations were then performed for deexcitation of the binary distribution of fragments produced during fission of a specific compound system. This schematic representation indicates the complexity of such calculations and provides an introduction to models that can be used for determination of prompt fission radiation emission.

Fig. 7. Reaction chain used for calculation of $^{237}\text{Np} + p$ total fission cross sections up to 100 MeV.

Recent reviews by Madland [77] and Märtén [78] summarize models used in calculation of prompt fission emissions. A short classification of these models is presented here. The first category is that of the temperature distribution model as typified by the Madland-Nix [74] and Dresden [79] formulations. These models address both neutron-induced and spontaneous fission and account for the distribution of fission fragment excitation energy, the energy dependence of the cross section for compound nucleus formation, the center-of-mass motion of the fission fragments, and multiple-chance fission at higher neutron energies. In these formulations, a triangular approximation to the corresponding fission fragment residual nuclear temperature distribution simulates the initial distribution of fission-fragment excitation energies and subsequent cooling as neutrons are emitted. A second approach, the cascade evaporation model (CEM [80]) is based upon the Weisskopf evaporation theory with factors applied to account for neutron-gamma-ray competition. This model, in addition to the features described with the temperature distribution model, includes the distribution of fission-fragment excitation energy in each step of the cascade process, the anisotropy of the center-of-mass neutron spectrum, complete fission-fragment mass and kinetic energy distributions, and fission-fragment nuclear level densities. The cascade-evaporation model has also been formulated to explicitly include effects of competition

between neutron and gamma-ray emission as described by Bozoian [81]. This later formulation provides a crude basis for calculation of prompt gamma-ray emission during deexcitation of fission fragment pairs. The final, and most sophisticated approach to calculation of prompt fission neutron, and particularly prompt gamma-ray, emission involves Hauser-Feshbach calculation of the deexcitation of the fission fragments. This model allows explicit competition between neutron and gamma-ray emission in the deexcitation of a given fission fragment, neutron transmission coefficients from optical model potentials, gamma-ray transmission coefficients, and angular momentum distributions for each fragment considered. Examples of this approach can be found in Ref. [82].

Important technical issues that impact modeling of prompt emissions from excited fission fragments include determination of the fission fragment mass distribution along with implementation of sampling procedures for fragment selection, determination of the total energy release in fission and its dependence on masses (experimental and theoretical) used to determine it, fission fragment kinetic energies, and, significantly, the partition of total fission fragment excitation energy between light and heavy fragments. This latter area is particularly important in calculations using the models described above. Several methods can be utilized in the energy partitioning process. The simplest assumption scales the total fragment excitation energy by the masses of fragments involved; $E_L^H = E^*/(1+A_L/A_H)$ and $E_H^L = E^*/(1+A_H/A_L)$. A similar approach partitions the excitation energy using the respective Fermi gas level density parameter a : $E_L^H = E^*/(1+a_L/a_H)$ and $E_H^L = E^*/(1+a_H/a_L)$. A standard approach for verifying choices for fragment excitation energy partitioning is by comparing calculations with data for neutron multiplicity as a function of fragment mass. Figure 8 illustrates such a comparison using partitioning based on the two mass and Fermi gas parameters. The example illustrates difficulties occurring when simple energy partition models are used. Better agreement can be achieved through adjustment of Fermi gas parameters [81] or through more detailed models for energy partitioning [83].

The question of better methods for partitioning of fission fragment energy at scission is a major uncertainty that restricts the applicability of fission neutron theory. Recent conclusions from specialists' workshops (e.g., Vienna, 1990 [84]) recommend that substantial effort be devoted towards improvement of models for energy partitioning.

Fig. 8. Calculation of the average number of neutrons emitted per fragment using two methods for fragment excitation energy partitioning.

Summary

This paper has reviewed the application of nuclear models in current evaluation efforts. Selected examples were given for principal models and their components. Significant progress continues in model development that is directly translated into improved calculations for data libraries. This trend will continue into the future and is driven by factors such as increased computational capabilities and expanded requirements for evaluated data files. Model improvement will also occur with general trends being direct use of more microscopic models specifically in data evaluation and/or use of microscopic treatments to guide phenomenology developed for application uses.

References

1. M. Uhl, Proc. Int. Conf. Nuclear Data and Technology, May 1988, S. Igarasi, Ed, p. 435.
2. M. A. Gardner, Proc. Int. Conf. on Nuclear Data for Basic and Applied Science, Santa Fe, May 1985, Eds., P. Young, R. Brown, G. Auchampaugh, P. Lisowski, L. Stewart, p. 1481.
3. P. G. Young, Proc. Int. Conf. Nuclear Data and Technology, Sept. 1982, Antwerp, Ed., K. H. Bockhoff, p. 506.
4. P. W. Lisowski, C. D. Bowman, E. D. Arthur, and P. G. Young, Nuclear Physics Information Needed for Accelerator Driven Transmutation of Nuclear Waste, contributed paper, this conference.
5. S. Howe, Nuclear Data Needs for the Space Exploration Initiative, invited paper, this conference.
6. A. Wambersie, Nuclear Data Relevant to Cancer Therapy with Fast Neutrons and Charged-Particles, invited paper this conference.
7. H. A. Weidenmüller, Recent Progress in the Statistical Theory of Nuclear Reactions, invited paper, this conference.
8. P. E. Hodgson, Dispersion Relations in Optical Model Development, invited paper, this conference.
9. C. Y. Fu, Nucl. Sci. Eng. **106**, 489 (1990).
10. G. M. Hale, P. G. Young, M. Chadwick, and Z. P. Chen, New Evaluations for ^{14}N and ^{16}O , invited paper, this conference.
11. G. M. Hale, D. Dodder, J. D. Seagrave, B. Berman, and T. Phillips, Phys. Rev. **C42**, 438 (1990).
12. T. W. Phillips, B. L. Berman, and J. D. Seagrave, Phys. Rev. **C22** (1980).
13. G. M. Hale, Proc. Int. Conf. on Muon Catalyzed Fusion, Vienna, May 1990.
14. D. V. Balin *et al.*, Phys. Rev. Lett. **141B**, 173 (1984); JETP Lett. **40**, 112 (1984).
15. D. V. Balin *et al.*, Proc. Int. Conf. on Muon Catalyzed Fusion, Vienna, 1990.
16. H. D. Knox, Proc. Int. Conf. on Nuclear Data for Basic and Applied Science, Santa Fe, NM, May 1985, Eds., P. G. Young, R. E. Brown, G. F. Auchampaugh, P. W. Lisowski, and L. Stewart, p. 1007.
17. D. A. Ressler, Advanced Modeling of Reaction Cross Sections for Light Nuclei, Proc. Wkshp. on Strong, Weak, and Electromagnetic Interactions in Nuclei and Astrophysics, Livermore, Calif., Jan. 1991; Lawrence Livermore Nat. Lab. Report, UCRL-JC106542.
18. G. M. Hale, Proc. Symp. in Honor of Akito Arima: Nuclear Physics in the 1990s, Santa Fe, N. M., May 1990, p. 8.
19. M. Bolsterli, A Scattering Theory for the Shell Model, Los Alamos Nat. Lab. report LA-UR-90-2506.
20. G. R. Satchler, *Direct Nuclear Reactions* (Clarendon Press, Oxford, 1983).
21. R. D. Lawson, P. T. Guenther, and A. B. Smith, Phys. Rev. **C34**, 1599 (1986).
22. See, e.g., S. Chiba, P. Guenther, R. Lawson, and A. B. Smith, Phys. Rev. **C42**, 2487 (1990) and refs.
23. R. D. Lawson, P. T. Guenther and A. B. Smith, Nucl. Phys. **A493**, 267 (1989).
24. P. G. Young and E. D. Arthur, Proc. Int. Conf. Nuclear Data For Science and Technology, Jülich, Germany, 13-17 May 1991 (to be issued).
25. G. Haouat, *et al.*, Nucl. Sci. Eng. **31**, 491 (1982).
26. P. Lisowski, personal communication, Los Alamos National Laboratory (1989).
27. R. Kozack and D. G. Madland, Phys. Rev. **C39**, 1461 (1990).
28. For example, B. C. Clark, R. L. Mercer, and P. Schwandt, Phys. Lett. **122B**, 211 (1983).
29. B. C. Clark *et al.*, Phys. Rev. **C30**, 314 (1984).
30. A. M. Lane, Nucl. Phys. **35**, 676 (1962).
31. R. Kozack and D. G. Madland, Nucl. Phys. **A509**, 664 (1990).
32. W. Hauser and H. Feshbach, Phys. Rev. **87**, 366 (1952).
33. V. W. Weisskopf and D. Ewing, Phys. Rev. **57**, 472 (1940).
34. A. Gilbert and A. G. W. Cameron, Can. J. Phys. **43**, 1446 (1965).
35. W. Dilg *et al.*, Nucl. Phys. **A217**, 269 (1973).
36. A. V. Ignatyuk *et al.*, Sov. J. Nucl. Phys. **21**, 450 (1979).
37. K. H. Schmidt *et al.*, Zeits. für Physik **A308**, 215 (1982).
38. E. D. Arthur *et al.*, Applied Uses of Level Density Models, Specialists' Meeting on Nuclear Level Densities, Bologna, 1989.
39. M. Ivascu *et al.*, Rev. Roum. Phys. **32**, 697 (1987).
40. Zhuang Youxiang *et al.*, Chinese J. of Nucl. Phys. **8**, 199 (1986).
41. J. L. Cook *et al.*, Aust. J. Phys. **20**, 477 (1967).
42. G. Reffo *et al.*, Nucl. Ins. Methods **A267**, 408 (1988).
43. S. Iijima *et al.*, Nucl. Science Technology **20**, 77 (1983).
44. M. Avrig *et al.*, Zeits. für Physik **A335**, 299 (1990).
45. M. Uhl and B. Strohmaier, Institut für Radiumforschung report, IRK 76/01, Vienna, 1976.
46. P. G. Young and E. D. Arthur, GNASH: A Preequilibrium Statistical Nuclear Model Code, Los Alamos Nat. Lab. Report LA-6947 (1977).
47. P. G. Young *et al.*, Calculation of (n,xy) Cross Sections Between Threshold and 100 MeV, Los Alamos Nat. Lab. report LA-UR-90-2129 (1990).
48. R. C. Haight *et al.*, Bull. Am. Phys. Soc. **35**, 1038 (1990).
49. D. G. Gardner, M. A. Gardner, and R. W. Hoff, Livermore Nat. Lab. report UCRL-95966 (1987).
50. J. Kopecky and M. Uhl, Phys. Rev. **C41**, 1941 (1990).
51. S. G. Kadenskij *et al.*, Sov. J. Nucl. Phys. **37**, 165 (1983).
52. D. M. Brink, Ph. D. Thesis, Oxford University, 1955.
53. H. Feshbach, A. Kerman, and S. Koonin, Ann. of Phys. **125**, 429 (1980).
54. T. Tamura, T. Udagawa, and H. Lenske, Phys. Rev. **C26**, 379 (1982).
55. H. Nishioka, J. J. M. Verbaarschot, H. A. Weidenmüller, and S. Yoshida, Ann. of Phys. **172**, 67 (1986).
56. P. E. Hodgson, Analysis of Multistep Reaction Cross Sections with the FKK Theory, Oxford Univ. (1990).
57. A. Marcinkowski *et al.*, Nucl. Phys. **A501**, 1, 1989.
58. H. Kalka, Proc. Specialists' Meeting on Preequilibrium Nuclear Reactions (sponsored by NEANDC/OECD), Semmering, Austria, 10 Feb. 1988, Ed., B. Strohmaier, p. 305.
59. W. Scobel *et al.*, Phys. Rev. **C41**, 2010 (1990).
60. A. A. Cowley *et al.*, Phys. Rev. **C43**, 678 (1991).
61. M. Herman and G. Reffo, Nuovo Cimento **A102**, 577 (1990).

62. A. J. Köning and J. M. Akkermans, Randomness in Multi-Step Direct Reactions, ECN Netherlands report, ECN-RX-90-065 (1990).
63. M. Avrigeanu and V. Avrigeanu, J. Phys. G15, L261, (1989).
64. M. Herman, G. Reffo, and C. Costa, Phys Rev C39, 1269 (1989).
65. C. Y. Fu, Pairing Corrections and Spin Cutoff Factors in Exciton Level Densities for Two Kinds of Fermion, Oak Ridge Nat. Lab. Preprint (1990).
66. J.M. Akkermans and H. Gruppelaar: Phys. Lett. 157B, 95 (1985).
67. P. Oblozinsky, Proc. Specialists' Meeting on Preequilibrium Nuclear Reactions, (sponsored by NEANDC/OECD), Semmering, Austria, Feb.10, 1988. Ed., B. Strohmaier, p. 157.
68. Oblozinsky: Phys. Lett. B215, 597 (1988).
69. F. Cvelbar, E. Betak, and J. Merhar: J. Phys. G17, 113 (1991).
70. P. Oblozinsky and M. B. Chadwick: Phys. Rev. C 42, 1652 (1990).
71. P. Oblozinsky: Phys. Rev. C 40, 1591 (1989).
72. M. B. Chadwick, P. Oblozinsky, P.E. Hodgson and G. Reffo, submitted to Phys. Rev. C.
73. M. B. Chadwick and G. Reffo, accepted for publication by Phys. Rev. C.
74. D. G. Madland and J. R. Nix, Nucl. Sci. Eng. 81, 213 (1982).
75. P. G. Young *et al.*, Transport Data Libraries for Incident Proton and Neutron Energies to 100 MeV, Los Alamos Nat. Lab. report, LA-11753-MS (1990).
76. E. D. Arthur and P. G. Young, Proc. Conf 50 Years with Fission, Gaithersburg, 1989, Eds., J. W. Behrens and A. D. Carlson, p. 931.
77. D. G. Madland, Theoretical Description of Neutron Emission in Fission, Los Alamos Nat. Lab. report, LA-UR-91-437 (1991).
78. H. Mårten *et al.*, Proc. Conf 50 Years with Fission, Gaithersburg, 1989, Eds., J. W. Behrens and A. D. Carlson, p. 743.
79. H. Mårten and D. Seeliger, J. Phys. G10, 349 (1984).
80. K. Arnold *et al.*, Nucl Phys A502, 325 (1989).
81. M. Bozoian *et al.*, Proc. Conf. 50 Years with Fission, Gaithersburg, 1989, Eds., J. W. Behrens and A. D. Carlson, p. 624.
82. J. C. Browne and F. S. Dietrich, Phys Rev C10, 2545 (1974).
83. A. Ruben *et al.*, Zeit. für Physik A338, 67 (1991).
84. LAEA Consultants' Meeting on Nuclear Data for Neutron Emission in the Fission Process, Vienna, October 1990.