THE UTILITY OF SPECTRAL MEASUREMENTS OF SECONDARY REACTION PRODUCTS

W.W. HEIDBRINK

*Plasma Physics Laboratory, Princeton University
Princeton, New Jersey, USA*

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

ABSTRACT. The spectra of 15 MeV protons and 14 MeV neutrons produced in the burnup of 0.8 MeV ^3He ions and 1 MeV tritons through the $d(^3\text{He},p)\alpha$ and $d(t,n)\alpha$ fusion reactions contain information on the velocity distributions of the energetic ^3He ions and tritons.

In a magnetically confined deuterium plasma, the behavior of alpha-like particles can be studied by measuring the number of 1 MeV tritons and 0.8 MeV ^3He ions (produced in the d-d fusion reactions) that undergo subsequent $d(t,n)\alpha$ and $d(^3\text{He},p)\alpha$ fusion reactions. "Burnup" measurements of 14 MeV neutrons and of 15 MeV protons have been reported for the PLT [1,2] and PDX tokamaks [2,3] and are in progress on the TFTR [4], JET [5], and Frascati [6] tokamaks. The burnup fluence depends on both the energetic ion confinement and slowing down, while burnup flux measurements are particularly useful for studying the rate of slowing down. The purpose of this note is to point out that spectral measurements of the triton and ^3He burnup also provide useful information about the behavior of alpha-like particles.

Consider a plasma with moderately low impurity levels (Z_{eff}) and electron temperature T_e . The tritons and ^3He ions slow down primarily on electrons [7] so pitch-angle scattering of the energetic ions may be neglected. We also assume that the bulk deuterium temperature T_i is small compared to the $d(t,n)\alpha$ and $d(^3\text{He},p)\alpha$ resonant energies so that the burnup reactions can be treated as beam-target reactions. Then the energy distribution of 14 MeV neutrons or 15 MeV protons $F(E_3)$ is given by [8]

$$F(E_3) = \frac{1}{2} \int \sin \chi_1 d\chi_1 \int dE_1 \sigma(E_1) E_1 f_1(E_1) \hat{F}(E_3, E_1, \chi_1), \quad (1)$$

where $f_1(E_1)$ is the triton or ^3He energy distribution function, σ is the $d(t,n)\alpha$ or $d(^3\text{He},p)\alpha$ fusion cross section, and $\hat{F}(E_3, E_1, \chi_1)$ is the reduced distribution function defined in eq. (7) of Ref. [8]. For fast ions that slow down on electrons, $f_1(E_1) \propto E_1^{-3/2}$. If all of velocity space is confined, the integration over triton or ^3He pitch angle χ_1 is from $\chi_1 = 0$ to $\chi_1 = \pi$.

The utility of spectral measurements of the fusion-product burnup is that the spectrum contains information about the velocity-space distribution of the tritons and ^3He ions. As an illustration, imagine a vertically

viewing neutron spectrometer, or a collimated 15 MeV proton detector that measures protons with velocities perpendicular to the magnetic field [9]. If the triton or ^3He confinement is isotropic in velocity space, numerical integration of eq. (1) gives the broad neutron and proton spectra shown in Fig. 1. If, however, the velocity space confinement is anisotropic, deviations from this shape are predicted. For example, if a physical mechanism such as ripple losses [10] or fishbones [11] expel energetic ions in the trapped region in velocity space but do not affect the confinement of circulating ions, then more narrow spectra are expected (Fig. 1).

The perpendicular detection geometry illustrated in Fig. 1 is probably the easiest to implement experimentally. Greater sensitivity to energetic ion anisotropy is available with other viewing angles, however. In Fig. 2, this increased sensitivity is illustrated by plotting the contributions to the burnup spectrum from ^3He ions in various portions of velocity space for a 15 MeV proton detector that measures protons with pitch angle $\chi_3 \approx \pi/4$. For this detection geometry, the contributions of co-circulating, perpendicular, and counter-circulating ^3He ions to the energy spectrum are separated by more than 1 MeV.

In summary, collimated spectral measurements of the triton and ^3He burnup are useful for detecting possible anisotropies in the velocity-space distributions of these energetic ions.

This work was supported by U.S. Department of Energy Contract No. DE-AC02-76-CHO-3073.

References

- [1] P. Colestock, J.D. Strachan, M. Ulrickson, R. Chrien, Phys. Rev. Lett. **43** (1979) 768.
- [2] W.W. Heidbrink, R.E. Chrien, J.D. Strachan, Nucl. Fusion **23** (1983)

- [3] W.W. Heidbrink, Ralph Hay, J.D. Strachan, Phys. Rev. Lett. **53** (1984) 1905.
- [4] J.D. Strachan *et al.*, in Proc. of Workshop on Basic Physical Processes of Toroidal Fusion Plasmas, (Varennna, 1985).
- [5] G. Sadler *et al.*, Bull. Am. Phys. Soc. **30** (1985) 1585.
- [6] E. Bittoni *et al.*, Proc. of the Europ. Conf. on Cont. Fusion and Plasma Physics (Budapest, 1985).
- [7] The condition that the critical energy (for which ion drag equals electron drag) is less than the energy at the peak of the cross section is $T_e Z_{eff} \lesssim 30$ for the triton and $T_e Z_{eff} \lesssim 120$ for the ^3He ion.
- [8] W.W. Heidbrink, Nucl. Instrum. Meth. **A236** (1985) 380.
- [9] W.W. Heidbrink, Nucl. Fusion **24** (1984) 636.
- [10] R.J. Goldston, R.B. White, A.H. Boozer, Phys. Rev. Lett. **47** (1981) 647.
- [11] K. McGuire *et al.*, Phys. Rev. Lett. **50** (1983) 891.

Figure Captions

Fig. 1 (a) Spectrum of perpendicular 14 MeV neutrons ($\chi_3 = \pi/2$) produced by 1.0 MeV tritons that slow down on electrons in a cold deuterium plasma. The tritons are assumed to populate velocity space uniformly (solid curve); the dashed curve is for a loss cone in the perpendicular part of velocity space ($|\chi_1 - \pi/2| < 0.63$). (o) Spectrum of perpendicular 15 MeV protons ($\chi_3 = \pi/2$) produced by 0.8 MeV ^3He ions that slow down on electrons in a cold deuterium plasma. The ^3He ions are assumed to populate velocity space uniformly (solid curve); the dashed curve is for a loss cone in the perpendicular part of velocity space ($|\chi_1 - \pi/2| < 0.63$).

Fig. 2 Spectrum of 15 MeV protons ($\chi_3 = \pi/4$) produced by 0.8 MeV ^3He ions that slow down on electrons in a cold deuterium plasma. The solid curve is for an isotropic ^3He velocity space distribution, the dashed curve is the spectrum produced by co-circulating ions ($\chi_1 < \pi/4$) and the dot-dash curve is the spectrum if only counter-circulating ions ($\chi_1 > 3\pi/4$) are confined.

Fig. 1

Fig. 2

EXTERNAL DISTRIBUTION IN ADDITION TO UC-20

Plasma Res Lab, Austro Nat'l Univ, AUSTRALIA
 Dr. Frank J. Paoloni, Univ of Wollongong, AUSTRALIA
 Prof. I.R. Jones, Flinders Univ., AUSTRALIA
 Prof. M.H. Brennan, Univ Sydney, AUSTRALIA
 Prof. F. Cap, Inst Theo Phys, AUSTRIA
 Prof. Frank Verheest, Inst theoretische, BELGIUM
 Dr. D. Balumbo, Dg XII Fusion Prog, BELGIUM
 Ecole Royale Militaire, Lab de Phys Plasmas, BELGIUM
 Dr. P.H. Sakanaka, Univ Estadual, BRAZIL
 Dr. C.R. James, Univ of Alberta, CANADA
 Prof. J. Teichmann, Univ of Montreal, CANADA
 Dr. H.M. Skarsgard, Univ of Saskatchewan, CANADA
 Prof. S.R. Greenivasan, University of Calgary, CANADA
 Prof. Tudor W. Johnston, INRS-Energie, CANADA
 Dr. Hannes Barnard, Univ British Columbia, CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Chalk River, Nucl Lab, CANADA
 Zhenggu Li, SW Inst Physics, CHINA
 Library, Tsing Hua University, CHINA
 Librarian, Institute of Physics, CHINA
 Inst Plasma Phys, Academia Sinica, CHINA
 Dr. Peter Lukac, Komenského Univ, CZECHOSLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Prof. Schatzman, Observatoire de Nice, FRANCE
 J. Radet, CEN-BP6, FRANCE
 AM Dupas Library, AM Dupas Library, FRANCE
 Dr. Tom Mial, Academy Bibliographic, HONG KONG
 Preprint Library, Cent Res Inst Phys, HUNGARY
 Dr. R.K. Chhajlani, Vikram Univ, INDIA
 Dr. B. Deygupta, Saha Inst, INDIA
 Dr. P. Kaw, Physical Research Lab, INDIA
 Dr. Phillip Rosenau, Israel Inst Tech, ISRAEL
 Prof. S. Cuperman, Tel Aviv University, ISRAEL
 Prof. G. Rostagni, Univ Di Padova, ITALY
 Librarian, Int'l Ctr Theo Phys, ITALY
 Miss Clelia De Falo, Assoc EURATOM-ENEA, ITALY
 Biblioteca, del CNR EURATOM, ITALY
 Dr. H. Yamato, Toshiba Res & Dev, JAPAN
 Direc. Dept. Ig. Tokamak Dev. JAERI, JAPAN
 Prof. Nobuyuki Inoue, University of Tokyo, JAPAN
 Research Info Center, Nagoya University, JAPAN
 Prof. Kyoji Nishikawa, Univ of Hiroshima, JAPAN
 Prof. Sigeru Mori, JAERI, JAPAN
 Prof. S. Tanaka, Kyoto University, JAPAN
 Library, Kyoto University, JAPAN
 Prof. Ichiro Kawakami, Nihon Univ, JAPAN
 Prof. Satoshi Itoh, Kyushu University, JAPAN
 Dr. D.I. Choi, Adv. Inst Sci & Tech, KOREA
 Tech Info Division, KAERI, KOREA
 Bibliothek, Pom-Inst Voor Plasma, NETHERLANDS
 Prof. B.S. Wiley, University of Waikato, NEW ZEALAND
 Prof. J.A.C. Cabral, Inst Superior Tecn, PORTUGAL
 Dr. Octavian Petrus, ALI CUZA University, ROMANIA
 Prof. M.A. Hellberg, University of Natal, SO AFRICA
 Dr. Johan de Villiers, Plasma Physics, Nucor, SO AFRICA
 Fusion Div. Library, JEN, SPAIN
 Prof. Hans Wilhelmson, Chalmers Univ Tech, SWEDEN
 Dr. Lennart Stenflo, University of UMEA, SWEDEN
 Library, Royal Inst Tech, SWEDEN
 Centre de Recherches, Ecole Polytech Fed, SWITZERLAND
 Dr. V.T. Tolok, Kharkov Phys Tech Ins, USSR
 Dr. D.D. Ryutov, Siberian Acad Sci, USSR
 Dr. G.A. Eliseev, Kurchatov Institute, USSR
 Dr. V.A. Glukhikh, Inst Electro-Physical, USSR
 Institute Gen. Physics, USSR
 Prof. T.J.M. Boyd, Univ College N Wales, WALES
 Dr. K. Schindler, Ruhr Universitat, W. GERMANY
 Nuclear Res Estab, Jülich Ltd, W. GERMANY
 Librarian, Max-Planck Institut, W. GERMANY
 Bibliothek, Inst Plasmaforschung, W. GERMANY
 Prof. R.K. Janev, Inst Phys, YUGOSLAVIA