DOE/ER/60675--5

DOE/ER/60675--5

DE92 008799

**Progress Report for the Continuation Application
DE-FG02-88ER60675**

Title: Molecular Mechanisms in Radiation Damage to DNA

Roman Osman, P.I. and Harel Weinstein, Co-P.I.
Department of Physiology and Biophysics
Mount Sinai School of Medicine, New York, NY 10029

The objectives of this work are to elucidate the molecular mechanisms that are responsible for radiation-induced DNA damage. The overall goal is to understand the relationship between the chemical and structural changes produced by ionizing radiation in DNA and the resulting impairment of biological function expressed as carcinogenesis or cell death. The studies are based on theoretical explorations of possible mechanisms that link initial radiation damage in the form of base and sugar damage to conformational changes in DNA. These mechanistic explorations should lead to the formulation of testable hypotheses regarding the processes of impairment of regulation of gene expression, alteration in DNA repair, and damage to DNA structure involved in cell death or cancer.

Last Year Accomplishments:

In the past year we have dedicated a major effort in two directions. One direction was to investigate the properties of base radicals with quantum mechanical methods; the other direction was to explore the conformational properties of radiation damaged DNA in water.

Properties of Base Radicals. In preparation for the investigation of the mechanism of H-abstraction by base radicals from the sugar we have expanded our studies from the hydroxyuracilyl and hydroxythyminylyl radicals to those of cytosine and adenine. The structures of the radicals produced by the addition of OH $\cdot$ to the base were optimized with high quality basis sets. The results indicate that the differences between the isomers of hydroxycytosinylyl radicals are small, both in energies and in structures, with the 6-OH adduct more stable than the 5-OH adduct. The behavior of the hydroxycytosinylyl radicals is very much like that of the hydroxyuracilyl and hydroxythyminylyl radicals. Both the energy and the structures of the different isomers of the hydroxyadeninylyl radicals are very different from each other. The most stable is the 8-OH adduct, which is nearly planar, while the 4-OH adduct is highly distorted and its energy is much higher than the 8-OH adduct. The charge distribution and the propensity of these radicals to abstract a hydrogen atom from sugars is now under investigation.

We have studied the mechanism of hydrogen abstraction by a hydroxyl radical. In model systems, such as methanol and ethanol, we were able to reproduce the experimentally observed rate constants for hydrogen abstraction in gas phase. We were able to predict for the first time the proportion of H-abstraction from the carbon and the oxygen at a range of temperatures covering a wide range from 200-2000 $^{\circ}$ K. A paradoxical behavior is observed, in which the ratio of the rates of the two processes (C-H abstraction and O-H abstraction) is diminishing as the temperature increases. This suggests a different tunneling behavior of the two processes. These predictions must await experimental confirmation.

Conformational Properties of Radiation Damaged DNA. The other major direction explored this year was the changes in conformational properties induced in radiation damaged DNA. These changes were studied by conducting molecular dynamics simulations of intact and radiation damaged DNA in water. The molecular dynamics simulations of the intact DNA, d(CGCGAATTCGCG) $_2$, show that DNA in aqueous solution is very flexible. The flexibility is expressed by a multitude of conformations with similar energies that can coexist in solution. While generally the DNA remains in a B-DNA form it undergoes severe unwinding of the helix and distortion of the helical axis. The average structure of the DNA in solution is in good agreement with that inferred from NMR measurements.

Radiation damaged DNA, i.e., a structure in which the thymine in position seven was replaced

MASTER

by a radiation damaged thymine, shows a different behavior than the intact DNA. Under similar simulation conditions the radiation damaged DNA shows a considerable rigidity in the area of the damage. The damage itself introduces a considerable distortion in the structure, however the DNA does not exhibit as extensive a flexibility as the intact structure. An analysis of the sources of this difference is now under investigation.

Plans for next year

Our plans for next year are as outlined in the renewal application from last year. We will explore the mechanism of H-abstraction from sugars by base radicals and radical cations. We will conduct molecular dynamics simulation on other radiation damaged DNA, such as those with cytosine or adenine radicals. These explorations will serve in formulating hypotheses about how the possible changes in damaged DNA produce a signal for the cellular repair machinery.

Published Articles and Conferences:

1. L. Pardo, R. Osman, J. Banfelder, A. P. Mazurek and H. Weinstein, Molecular Mechanisms of Radiation Induced DNA Damage: H-Abstraction and β -Cleavage. *Free Radical Res. Comms.*, **12-13**, 461-463, 1991.
2. R. Osman, L. Pardo, J. Banfelder, A. P. Mazurek, L. Shvartzman, R. Strauss and H. Weinstein, Molecular Mechanisms of Radiation Induced DNA Damage: H-Addition to Bases, Direct Ionization and Double Strand Breaks. *Free Radical Res. Comms.*, **12-13**, 465-467, 1991.
3. Weinstein, H., Osman, R., Mercier, G.A., Mazurek, P.A., Pardo, L., Rubenstein, L.A. Theory and computation of molecular mechanisms in biological processes. In *Computer Assisted Analysis And Modeling on the IBM 3090*. H. U. Brown, III, ed., 1989 IBM 3090 Supercomputing Competition, MIT Press, 1990, pp. 629-673.
4. L. Pardo, J. Banfelder and R. Osman. Theoretical Studies of the Kinetics, Thermochemistry, and Mechanism of H-Abstraction from Methanol and Ethanol, *J. Am. Chem. Soc.*, submitted.
5. R. Osman, K. Miaskiewicz, and H. Weinstein. Structure-Function Relations in Radiation Damaged DNA. Proceedings of DOE Meeting at Woods Hole, September 1990.
6. K. Miaskiewicz, R. Osman and H. Weinstein. Molecular Dynamics Simulations of the Hydrated d(CCGGAATTCGCG)₂ Dodecamer. *J. Am. Chem. Soc.*, submitted.

Drs. Karol Miaskiewicz, a research associate, and Roman Osman, the P.I., presented results of this work at the Seventh International Conversation in Biomolecular Structure and Dynamics at Albany, NY, June 18-21, 1991.

Dr. Harel Weinstein presented an invited lecture at the 9th International Congress of Radiation Research in Toronto, Canada, July 6-9, 1991.

Dr. Roman Osman presented a poster on this work at the 15th International Union of Biochemistry Meeting in Jerusalem, Israel, August 4-9, 1991.

Dr. Roman Osman presented an invited plenary lecture at the Molecular Graphics Society Meeting which took place at Oxford, England, Sep. 15-18, 1991.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

**DATE
FILMED**

4 / 15 / 92