TECHNICAL REPORT

September 1 Through November 30, 1991

Project Title: Microbial Strain Improvement For Organosulfur
Removal From Coal **DE-FG 22-91 PC 91334**

Principal Investigator: John J. Kilbane II, Institute of Gas Technology

Project Manager: Dr. Daniel D. Banerjee, CRSC

ABSTRACT

IGT has developed a microbial culture of Rhodococcus rhodochrous, designated as IGTS8, that is capable of specifically cleaving carbon-sulfur bonds in a range of organosulfur model compounds and is capable of removing organic sulfur from coal and petroleum without sacrificing the calorific value of the fuel. Although IGTS8 possesses the ability to specifically remove organic sulfur from coal, a major research need is to develop improved strains of microorganisms that possess higher levels of desulfurization activity and therefore will permit more favorable biodesulfurization process conditions: faster rates, more complete removal, and smaller reactor size. Strain improvement is the single most important aspect to the development of a practical coal desulfurization process. Microbial strain improvement can be accomplished using genetic engineering recombinant DNA techniques as well as by non-recombinant techniques of microbiology and genetics. Both approaches will be employed in the proposed research, with the objective of developing derivatives of IGTS8 with markedly enhanced levels of desulfurization activity and evaluating the ability of improved cultures to remove organic sulfur from coal. During the current quarter, desulfurization-deficient mutants of IGTS8 have been obtained and characterized, improved cloning vectors have been constructed, the construction and screening of genomic libraries has been initiated, and a derivative of IGTS8 with enhanced desulfurization ability has been obtained.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISSEMINATION OF THIS DOCUMENT IS UNLIMITED

This project is funded by the U. S. Department of Energy (PETC) and by the Illinois Department of Energy and Natural Resources as part of their cost-shared program. **CB**

EXECUTIVE SUMMARY

The objective of the proposed research is to develop derivatives of Rhodococcus rhodochrous IGTS8 that possess greatly enhanced levels of desulfurization activity and to evaluate the ability of improved cultures to remove organic sulfur from coal.

The approach taken in this project to achieve that objective is to use genetic engineering/recombinant DNA techniques as well as non-recombinant methodologies of microbiology and genetics to produce/isolate strains of IGTS8 with improved desulfurization activity. IGTS8 has been shown to be capable of specifically cleaving carbon-sulfur bonds in a range of organosulfur model compounds and in coal. The challenge now is to develop a practical, economical coal biodesulfurization process. While coal has been successfully desulfurized using IGTS8, the shortest treatment time and minimum ratio of coal-to-bacterial culture obtained in experiments with solid coal so far is three weeks and one gram-to seven liters, respectively. It is not only desirable, but absolutely essential, that improved derivatives of IGTS8 with greatly enhanced levels of desulfurization activity should be developed in order to allow briefer treatment times, more complete removal, smaller reactor volumes, and more favorable coal-to-bacterial culture ratios.

The development of improved cultures with enhanced levels of desulfurization activity is critical to the successful development of a coal biodesulfurization process. Until recently, techniques for the genetic analysis and modification of Rhodococcus rhodochrous were lacking. However, research at IGT has resulted in the development of tools and techniques for genetic studies of IGTS8 as well as the development of several assays for the quantification of the levels of desulfurization activity of IGTS8 or derivative cultures.

It is not known how many desulfurization genes exist but a good way to investigate this question is to obtain mutants of the desulfurization trait and then after characterizing them determine how many groupings/categories of similar mutations exist. Over 25,000 individual colonies of mutagenized bacterial cultures were examined using a variety of techniques: only one category of desulfurization-deficient mutation was observed. While it is difficult to envision a single enzyme capable of selectively cleaving carbon-sulfur bonds in a wide array of chemical structures the data obtained thus far are consistent with the possibility that a single structural gene and a single enzyme may be responsible for the desulfurization ability of IGTS8. The collection of desulfurization-deficient mutants obtained thus far will also be helpful in screening genomic libraries for the presence of desulfurization genes.

Previously a plasmid/replicon capable of functioning in IGTS8 was identified but a convenient cloning vector for use in IGTS8 did not exist prior to this project. During the current quarter, several cloning vectors were constructed. Genomic libraries were prepared using these vectors and the screening of those libraries is currently in progress.

Tolyldisulfide is not only not metabolized by IGTS8 but appears to be a specific inhibitor of the desulfurization trait of IGTS8. However, a mutant/derivative culture of IGTS8, designated IGTS86, has been obtained that

is capable of utilizing tolyldisulfide as its sole source of sulfur for growth. Preliminary characterization of IGTS86 indicates that it is about 25% more active than IGTS8 in desulfurization assays.

OBJECTIVES

The objective of this research is to develop improved derivatives of Rhodococcus rhodochrous IGTS8 with greatly enhanced levels of desulfurization activity and to evaluate the ability of improved cultures to remove organic sulfur from coal. Specifically, recombinant DNA techniques will be used to identify DNA fragments that encode desulfurization genes which will be subsequently manipulated to achieve enhanced expression of these genes. Additionally, non-recombinant DNA techniques of mutagenesis, selection, and screening will be used to obtain improved bacterial cultures.

INTRODUCTION AND BACKGROUND

There are numerous physical, chemical, and microbiological techniques that can effectively remove inorganic sulfur from coal prior to combustion. Moreover, there are physical and chemical techniques for pyrite/ash removal that have been successfully commercialized and are routinely employed in the coal industry. However, while there are technologies capable of removing organic sulfur from coal prior to combustion no commercially viable technology currently exists. The removal of organically bound sulfur from coal by physical/chemical techniques requires harsh conditions as compared with microbiological techniques; therefore, the microbiological approach to the process with more favorable economics than currently available technologies.

Unfortunately, with nearly all Illinois coals physical cleaning alone will not produce a coal that complies (if burned without additional sulfur removal from the combustion gases) with New Source Performance Standards (NSPS). To remove the additional organic sulfur, another method such as microbiological treatment that breaks the chemical bonds binding the sulfur to the carbon must be developed. A coal cleaning strategy employing only physical or chemical cleaning methods to remove as much of the pyrite and other mineral matter as economically possible will be unlikely to produce a product capable of meeting emission standards. A separate step such as microbial cleaning will likely be needed that focuses on the removal of the organic sulfur. A microbial culture, Rhodococcus rhodochrous IGTS8, has been identified that is capable of selectively cleaving carbon-sulfur bonds in model compounds and in coal; however, the level of expression of desulfurization enzymes by IGTS8 is low. Improved bacterial cultures are needed if a practical coal biodesulfurization process is to be developed. The goal of this project is to use recombinant and non-recombinant DNA techniques to develop microbial cultures with enhanced desulfurization abilities.

RESULTS AND DISCUSSION

The complementation of desulfurization-deficient (dsz⁻) mutants is the main strategy employed in this project to identify cloned DNA fragments that encode desulfurization genes. Accordingly the characterization of dsz⁻ cultures is rather important. It is crucial to utilize stable dsz⁻ cultures; that is, cultures that don't readily revert to the dsz⁺ phenotype. Moreover it is possible that more than one kind/category of dsz⁻ mutants exist reflecting mutations in different genes. To address these issues seven of the most stable dsz⁻ mutants in our collection (designated IGTS8M1 thru IGTS8M7)

as well as two newly derived cultures obtained by UV mutagenesis (UV23 and UV24) were tested for their ability to grow with various organosulfur compounds and the Gibbs assay was used to assess the desulfurization activity of these cultures. The results of these growth and activity tests are reported in Tables 1 and 2. The dsz⁻ mutants were expected to grow with sulfate but not with DBT as sulfur sources. The ability of dsz⁻ mutants to grow with DBT₂SO₂, DMSO, thiamine, or methionine was investigated to allow the detection of different categories of dsz⁻ mutants. The results shown in Tables 1 and 2 confirm that the control cultures IGTS8 and IGTS85 both utilize DBT and DBT₂SO₂ as sulfur sources and score positively in the Gibbs assay. Moreover these results demonstrate differences between IGTS8 and IGTS85 in that IGTS8 yielded positive Gibbs assay results only when grown on DBT or DBT₂SO₂ and that growth of IGTS8 using DMSO and thiamine was sparse. In comparison, IGTS85 grew well with DMSO and with thiamine and yielded positive Gibbs assay values when grown with either DBT, DBT₂SO₂, DMSO, or with thiamine. Of the mutant cultures tested IGTS8M2, IGTS8M5, and IGTS8M6 failed to grow under any circumstances and IGTS8M4 gave variable results so that these cultures do not appear to be good candidates for use in the detection of cloned dsz genes. IGTS8M1, IGTS8M3, and UV23 appear to be good dsz⁻ cultures in that they grow with sulfate, DMSO, and thiamine as sulfur sources but not with DBT or DBT₂SO₂, and they fail to yield a positive Gibbs assay value when grown under any circumstances. IGTS8M7 yielded confusing results in that it grew only with sulfate and DBT₂SO₂ but it did not score positively in the Gibbs assay. More testing is needed to determine if IGTS8M7 is a dsz⁻ mutant distinct from IGTS8M1, IGTS8M3, and UV23. The UV24 culture shows a level of desulfurization activity comparable to IGTS85 when grown with either sulfate, DBT, DBT₂SO₂, DMSO, or with thiamine. This examination of dsz⁻ mutants of IGTS8 failed to detect different categories of mutants based on the range of organosulfur compounds metabolized. Moreover GC/MS studies of dsz⁻ mutants exposed to DBT failed to detect intermediate metabolites. All of the dsz⁻ mutants examined appear to belong to a single class. This is consistent with the hypothesis that a single desulfurization gene exists. To lend further support to this supposition dsz⁻ mutants identified because of their inability to utilize DBT are similarly unable to utilize other structurally unrelated compounds such as trithiane yet if revertants are obtained that have regained their ability to utilize trithiane they simultaneously regain their ability to utilize DBT. Likewise revertants of dsz⁻ mutants that are selected for their ability to utilize DBT simultaneously regain their ability to utilize trithiane (and other organosulfur compounds). It is difficult to envision a single enzyme encoded for by a single gene being capable of cleaving carbon-sulfur bonds in a broad array of chemical structures; nonetheless, the data obtained thus far in the characterization of IGTS8 is consistent with that possibility.

The plasmid pRF29 was selected for research concerning the genetic analysis of the dsz trait of IGTS8 because it was found to be essentially the only plasmid known to be capable of replicating in IGTS8. Unfortunately pRF29 is not a particularly convenient cloning vector as it contains few unique restriction sites that can be used for the insertion of chromosomal DNA fragments in the construction of genomic libraries. Moreover the ampicillin resistance gene encoded by pRF29 is expressed in E. coli but not in IGTS8 and the chloramphenicol resistance gene is expressed in IGTS8 but not in E. coli. Most commonly used cloning vectors allow the positive selection of derivatives that contain inserted/cloned DNA fragments or allow derivatives

Table 1. GROWTH OF IGTS8 AND DESULFURIZATION DEFICIENT MUTANTS OF IGTS8 ON VARIOUS ORGANOSULFUR COMPOUNDS

<u>Sample</u>	<u>Bacterial Culture</u>	<u>Sulfur Source</u>	<u>(Klett Units)</u>	<u>ppm 2-HBP/10⁸ Cells</u>
1	IGTS8	None	15	0
2	IGTS8	Sulfate	405	0
3	IGTS8	DBT	310	1.30
4	IGTS8	DBTSO ₂	232	1.73
5	IGTS8	DMSO	7	0
6	IGTS8	Thiamine	56	0
7	IGTS8	Methionine	370	0
8	IGTS8M1	None	42	0
9	IGTS8M1	Sulfate	570	0
10	IGTS8M1	DBT	69	0
11	IGTS8M1	DBTSO ₂	48	0
12	IGTS8M1	DMSO	510	0
13	IGTS8M1	Thiamine	325	0
14	IGTS8M1	Methionine	600	0
15	IGTS8M2	None	4	0
16	IGTS8M2	Sulfate	7	0
17	IGTS8M2	DBT	8	0
18	IGTS8M2	DBTSO ₂	9	0
19	IGTS8M2	DMSO	10	0
20	IGTS8M2	Thiamine	10	0
21	IGTS8M2	Methionine	10	0
22	IGTS8M3	None	13	0
23	IGTS8M3	Sulfate	246	0
24	IGTS8M3	DBT	14	0
25	IGTS8M3	DBTSO ₂	22	0
26	IGTS8M3	DMSO	170	0
27	IGTS8M3	Thiamine	460	0
28	IGTS8M3	Methionine	103	0
29	IGTS8M4	None	30	0
30	IGTS8M4	Sulfate	474	0
31	IGTS8M4	DBT	105	0
32	IGTS8M4	DBTSO ₂	44	0
33	IGTS8M4	DMSO	206	0
34	IGTS8M4	Thiamine	325	0
35	IGTS8M4	Methionine	335	0
36	IGTS8M5	None	9	0
37	IGTS8M5	Sulfate	5	0
38	IGTS8M5	DBT	5	0
39	IGTS8M5	DBTSO ₂	8	0
40	IGTS8M5	DMSO	11	0
41	IGTS8M5	Thiamine	8	0
42	IGTS8M5	Methionine	7	0

Table 2. EXAMINATION OF DESULFURIZATION DEFICIENT MUTANTS OF IGTS8

<u>Sample</u>	<u>Bacterial Culture</u>	<u>Sulfur Source</u>	<u>(Klett Units)</u>	<u>ppm 2-HBP/10⁸ Cells</u>
1	IGTS85	None	48	0
2	IGTS85	Sulfate	410	0
3	IGTS85	DBT	147	1.77
4	IGTS85	DBTSO ₂	410	2.14
5	IGTS85	DMSO	440	0.15
6	IGTS85	Thiamine	234	0.75
7	IGTS8M1	None	55	0
8	IGTS8M1	Sulfate	195	0
9	IGTS8M1	DBT	52	0
10	IGTS8M1	DBTSO ₂	75	0
11	IGTS8M1	DMSO	99	0
12	IGTS8M1	Thiamine	225	0
13	IGTS8M2	None	31	0
14	IGTS8M2	Sulfate	25	0
15	IGTS8M2	DBT	19	0
16	IGTS8M2	DBTSO ₂	25	0
17	IGTS8M2	DMSO	22	0
18	IGTS8M2	Thiamine	23	0
19	IGTS8M3	None	73	0
20	IGTS8M3	Sulfate	175	0
21	IGTS8M3	DBT	55	0
22	IGTS8M3	DBTSO ₂	89	0
23	IGTS8M3	DMSO	113	0
24	IGTS8M3	Thiamine	280	0
25	IGTS8M4	None	110	0
26	IGTS8M4	Sulfate	68	0
27	IGTS8M4	DBT	37	0
28	IGTS8M4	DBTSO ₂	41	0
29	IGTS8M4	DMSO	63	0
30	IGTS8M4	Thiamine	340	0

Table 2, Cont. EXAMINATION OF DESULFURIZATION DEFICIENT MUTANTS OF IGTS8

<u>Sample</u>	<u>Bacterial Culture</u>	<u>Sulfur Source</u>	<u>(Klett Units)</u>	<u>ppm 2-HBP/10⁸ Cells</u>
31	IGTS8M5	None	13	0
32	IGTS8M5	Sulfate	13	0
33	IGTS8M5	DBT	13	0
34	IGTS8M5	DBTSO ₂	19	0
35	IGTS8M5	DMSO	15	0
36	IGTS8M5	Thiamine	138	0
37	UV24	None	52	0
38	UV24	Sulfate	250	0.24
39	UV24	DBT	284	3.07
40	UV24	DBTSO ₂	288	3.03
41	UV24	DMSO	93	1.11
42	UV24	Thiamine	365	1.09
43	UV23	None	71	0
44	UV23	Sulfate	420	0
45	UV23	DBT	82	0
46	UV23	DBTSO ₂	105	0
47	UV23	DMSO	193	0
48	UV23	Thiamine	320	0
49	IGTS8M6	None	9	0
50	IGTS8M6	Sulfate	9	0
51	IGTS8M6	DBT	8	0
52	IGTS8M6	DBTSO ₂	25	0
53	IGTS8M6	DMSO	11	0
54	IGTS8M6	Thiamine	11	0
55	IGTS8M7	None	19	0
56	IGTS8M7	Sulfate	258	0
57	IGTS8M7	DBT	3	0
58	IGTS8M7	DBTSO ₂	185	0
59	IGTS8M7	DMSO	9	0
60	IGTS8M7	Thiamine	9	0

that contain inserted DNA fragments to be identified in some convenient fashion such as the inactivation of an antibiotic resistance gene. No such possibility exists with pRF29. The best straightforward use of pRF29 as a cloning vector is to utilize the XhoI sites, which occur in non-essential regions of pRF29 as regards replication and antibiotic resistance genes, for insertion of XhoI and/or SalI fragments of IGTS8 chromosomal DNA. To facilitate this approach the two smallest XhoI fragments of pRF29 were deleted to yield pIGTS1 which has a single XhoI site. Genomic libraries prepared using chromosomal DNA from dsz⁺ IGTS8 cells were prepared and the libraries were introduced by electroporation into the dsz⁻ strains IGTS8M1 and IGTS8M3. Chloramphenicol resistant transformants were selected and individual colonies were examined using the spray-plate fluorescence assay to detect the restoration of the dsz⁻ phenotype to dsz⁺ as a consequence of genetic complementation. Similarly chloramphenicol resistant transformants were pooled and used to inoculate Sulfur Bioavailability Assays thereby selecting for dsz⁺ microorganisms. No complementation was detected using either approach so far. The screening of these genomic libraries in E. coli was also attempted using both approaches described above; however, no dsz⁺ clones were detected.

Various fragments of pRF29 or pIGTS1 were subcloned into pBR322 or pBR325 to either aid in the analysis of pRF29 or to create a more convenient cloning vector. Most notably the Hind III fragment of pRF29 that encodes its replication functions (ori) was cloned into pBR325 to yield a plasmid designated pIGTS2. pIGTS2 was shown to be capable of successfully transforming IGTS8 to chloramphenicol resistance. The construction and screening of genomic libraries prepared in pIGTS2 will be included in the next report.

In other work, non-recombinant DNA techniques were used to obtain a derivative of IGTS8 that appears to be about 25% more active than IGTS8 in the Gibb's desulfurization assay. A wide range of organosulfur compounds are metabolized by IGTS8 and will serve as sole sources of sulfur in the IGT Sulfur Bioavailability Assay; however, some compounds, tolyldisulfide in particular, appear to be specific inhibitors of the desulfurization trait. IGTS8 is unable to utilize tolyldisulfide as a source of sulfur. IGTS8 can grow well when sulfate and tolyldisulfide are both present but can not grow on organosulfur compounds such as DBT when tolyldisulfide is also present. This suggests that tolyldisulfide may be a specific inhibitor of the desulfurization trait. Mutagenized IGTS8 cells were eventually obtained that can grow with tolyldisulfide as their sole source of sulfur. This culture, designated IGTS86, appears to express the desulfurization trait at higher levels than IGTS8 as measured by the Gibbs assay. Preliminary tests indicate IGTS86 is about 25% more active than IGTS8. During the next quarter, IGTS86 will be characterized more thoroughly and additional derivatives of IGTS8 capable of utilizing tolyldisulfide will be sought and examined.

CONCLUSIONS AND RECOMMENDATIONS

Good progress has been made in the isolation and characterization of desulfurization-deficient mutants of IGTS8 and in the construction of improved cloning vectors and genomic libraries. The construction and screening of

genomic libraries to identify DNA fragments that encode desulfurization genes will be the focus of continuing research. The development of cultures with enhanced desulfurization activity as a consequence of adaptation to tolyl disulfide appears to be quite promising and will be examined further.

END

DATE
FILMED

4 102 192

I