Progress Report May, 1991 - April 30, 1992
PARTIAL-WAVE ANALYSES OF HADRON SCATTERING BELOW 2 GeV

Richard A. Arndt and L. David Roper

DOE/ER/40454--T1

Department of Physics
Virginia Polytechnic Institute and State University
Blacksburg, Virginia 24061-0435

DE92 005468

Prepared for the U.S. Department of Energy
Under Contract Number DE-FG05-88ER40454

ABSTRACT

The Center for Analysis of Particle Scattering (CAPS) in the Department of Physics at Virginia Polytechnic Institute and State University has analyzed basic two-body hadron reactions below 2 GeV for the last two decades. Reactions studied were nucleon-nucleon, pion-nucleon, K^+ -nucleon and pion photoproduction systems. In addition to analyses of these systems, a computer graphics system (SAID) has been developed and disseminated to over 250 research institutions using VAX computers. The computer-interactive system for disseminating information on basic scattering reactions is also accessible to the physics community through TELNET on the VPI&SU physics department VAX.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

Nucleon-Nucleon Analyses:

Over the last year, much work has been done with the nucleon-nucleon scattering data base, which consists at present of about 12,000 p - p scattering data and 7000 n - p scattering data below $T_{lab}=1600$ MeV. Work has been done through BITNET with Mike McNaughton of LAMPF to "clean-up" the data base by eliminating redundant data sets, replacing preliminary measurements with final values, ensuring proper referencing and re-working the polarization data from LAMPF to correct for a miscalculated beam polarization. This latter effort has affected data accumulated over the last decade at Los Alamos. We have now completed this effort and have prepared a rough draft of our present analyses for submission to Physical Review. Mike McNaughton is joining us as a co-author on this publication and is presently writing a section describing the data-base changes.

In May we presented the results of our current analyses¹ to a meeting of "NEANDC Specialists on Neutron Cross Section Standards for the Energy Region above 20 MeV" held in Uppsala, Sweden. This is an international group of individuals doing radiation studies with nucleons who depend for calibration of their experiments upon n - p differential cross sections. At lower energies (below 20 MeV) they rely upon a "standard" (ENDF-VI) developed, and periodically revised, by a group at Los Alamos who do an R-matrix fit to low-energy NN data. They now require a similar standard at higher energies and have requested our VPI&SU group to provide one for energies up to 400 MeV. In response to this request we have developed a fit to elastic NN data below 400 MeV and have created a subroutine which precisely duplicates the n - p cross section predictions of this fit; this subroutine will be circulated for use by groups requiring calibration information. We also have done a substantial amount of work to determine what appropriate accuracy should be ascribed to this "standard." It is our belief that, given the present state of experimental data, n - p differential cross sections are reliable only at the few-percent level; this is often the limiting factor upon the precision with which the NEANDC group can do their studies. This suggests the need for further direct measurements of cross sections and other observables required to "pin-down" the asymmetries in the n - p system.

REFERENCES

- [1] "Nucleon-Nucleon Elastic Scattering Analyses at VPI&SU," Richard Arndt, Conference on Neutron Cross Section Standards for the Energy Region above 20 MeV, Uppsala, Sweden, May 1991.

Pion-Nucleon Analyses:

Our analyses of pion-nucleon elastic scattering data to 2 GeV have been published¹ this year. This work covers the majority of nucleon resonance states listed by the Particle Data Group. A complimentary study of the photon-decay amplitudes for these resonance states is also nearing completion. (See the pion photoproduction section below for details.) While there is qualitative agreement with the Karlsruhe-Helsinki and Carnegie-Mellon-Berkeley analyses, one feature of our analyses has generated considerable controversy. It is our reduced value of the fundamental pion-nucleon coupling constant. We have commented on the consequences of this result² and have repeated our determination using the Bugg, Carter and Carter method³, interior dispersion relations⁴ and the Goldberger-Miyazawa-Oehme sum rule⁵. The sum rule technique gives an essentially model-independent value for the coupling and also predicts a value below the "canonical" result of Koch and Pietarinen⁶. We have made some preliminary studies of the effect of this result on the pion-nucleon scattering lengths and sigma term. Work on this topic is continuing.

REFERENCES

- [1] Pion-nucleon partial-wave analysis to 2 GeV, R. A. Arndt, Z. Li, L. D. Roper, R. L. Workman and J. M. Ford, Phys. Rev. D43, 2131 (1991).
- [2] Confirmation and consequences of a reduced pion-nucleon coupling constant, R. A. Arndt and R. L. Workman, Phys. Rev. C43, 2436 (1991).
- [3] The pion-nucleon coupling constant, R. A. Arndt, Z. Li, L. D. Roper and R. L. Workman, Phys. Rev. D44, 289 (1991).
- [4] R. A. Arndt, unpublished.
- [5] R. L. Workman, R. A. Arndt and M. Pavan, On the Goldberger-Miyazawa-Oehme sum rule, submitted to Phys. Rev. Lett.
- [6] R. Koch and E. Pietarinen, Nucl. Phys. A336, 331 (1980).

K^+ -Nucleon Analyses:

The analysis of K^+ -nucleon scattering has now been completed and a preliminary draft of a paper for publication, based upon the thesis of John Hyslop¹ has been completed. These analyses are for K^+ - p scattering to a laboratory kinetic energy of 2650 MeV and K^+ -deuteron elastic and inelastic scattering to a laboratory kinetic energy of 1100 MeV.

REFERENCES

[1] "An Analysis of K^+ -Nucleon Scattering," VPI&SU Ph.D. Thesis of John Smith Hyslop III.

Pion-Photoproduction Analyses:

We have extended our analyses of pion photoproduction to a photon laboratory energy of 1.8 GeV. In addition, a new experiment¹ has provided us with total cross-section and differential cross-section measurements near the $p\pi^0$ threshold. This has allowed us to examine² the result³ of Mazzucato et al., which suggested a possible violation of the low-energy theorem. The results of the analyses to 1.8 GeV were also used in a re-examination of the Drell-Hearn-Gerasimov sum rule⁴.

We have extracted coupling constants for all seventeen resonances found within our energy range. We have given the resonance amplitudes a rating on a scale of one to three stars, according to their shape and resonance/background composition. Resonances with a two-star or three-star rating give a consistent and reliable result for coupling constants. This has been discussed⁵ and presented at the Hadron '91 conference. A further refinement of the analysis result, including a critical examination of the data base, is underway.

Zhujun Li spent this summer at CEBAF as a SURA/CEBAF fellowship recipient. There she worked on the modeling of pion electroproduction experiments using a non-relativistic quark model and studied the possibility of the Roper resonance being a hybrid state⁶. Pion photoproduction is her Ph.D. thesis topic.

- [1] R. Beck, J. D. Kellie and S. J. Hall, Phys. Rev. Lett. 65, 1841 (1990).
- [2] R. Workman, R. A. Arndt and Z. Li, Mod. Phys. Lett. A6, 1757 (1991).
- [3] E. Mazzucato et al., Phys. Rev. Lett. 57, 3144 (1986).
- [4] "Saturation of the Drell-Hearn-Gerasimov Sum Rule Revisited," R. L. Workman and R. A. Arndt, submitted to Phys. Rev. D.
- [5] "Resonance Photon-Decay Amplitudes," Z. Li, R. A. Arndt, L. D. Roper and R. L. Workman, contributed paper to Hadron '91 Conference.
- [6] "Electroproduction of the Roper Resonance as a Hybrid State," Zhenping Li, Volker Burkert and Zhujun Li, submitted to Phys. Rev. Lett., CEBAF-PR-91-032.

**DATE
FILMED
02/13/92**

I