Progress Report for Second Grant Year

DOE/ER/45076--2

DE92 007959

Inelastic Electron Scattering from Surfaces

Abstract

This report contains highlights of accomplishments of the past year, for the University of California, Irvine and the University of Wisconsin, Milwaukee collaboration on surface excitations, and their interactions with low energy electrons. In addition, we present a summary of future research to be carried out in the coming grant year.

Prof. S. Y. Tong
University of Wisconsin, Milwaukee

Prof. D. L. Mills
University of California, Irvine

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. Highlights of the Last Grant Period

The California-Wisconsin joint research program continues to produce exciting and leading-edge results. In the past grant year, our new results are published in *Phys. Rev. Lett.*, *Science*, *Surface Science Lett.*, etc. Mills and Tong finished an invited review article on high-resolution electron-energy-ion spectroscopy for the *Springer Series in Surface Sciences*. The editor of the prestigious Landoldt-Boernstein Volume has requested that Mills and Tong write separate chapters for its *Volume on the Physics of Solid Surfaces*.

The thrust of our research is to make quantitative analysis and predictions of the structural and dynamical properties of a clean surface or a surface with adsorbed atoms or molecules on it. Our research program is unique in the U. S. in that the theory we developed has a high degree of precision so that direct and quantitative contact can be made with the experimental measurements on the final (scattering) state of the outgoing inelastically scattered electron. While our present efforts are focused on electron energy losses through coupling to vibrational and magnetic degrees of freedom, it is the case that our theoretical program is directly relevant to measurements in a host of other important areas, such as synchrotron radiation based photoemission. In view of increased commitment to fund new synchrotron-radiation based photoemission experiments, it is anticipated that the new data will need quantitative theoretical interpretation. Our theoretical effort can provide such highly precise interpretation of new data because we directly calculate the outgoing photoexcited electron flux, which includes the excitation matrix element effect as well as final state scattering effects. The formalism we have developed and implemented, which now includes a fully relativistic description of the electron-substrate interaction, can be applied to diverse aspects of the electron spectroscopy of materials, including those with magnetic order.

Our program is a coordinated joint effort between researchers at the University of California, Irvine and the University of Wisconsin-Milwaukee. The two principal investigators have collaborated actively during the past ten years, and the joint research program, carried out in parallel with innovative experiments in the U. S. and Europe, is a leading effort in this important area. Our joint program has been responsible for a number of major advances made in recent years. The close collaboration of the two programs can be seen by tracing the methods used in the past few years. The highly precise inelastic electron scattering method was developed at Wisconsin based on the generalized Green's function approach (Tong *et al.* *Phys. Rev. B* **16**, 1459 (1977)). This approach is very versatile and is applicable to complex surface unit cells of clean and adsorbate covered systems. In the last grant period, Mills and coworkers expanded the theory to include spin-dependent diffraction. Now that the California program is working out well, Tong and coworkers will develop a spin-dependent photoemission method based partly on the California approach. This rapid expertise transfer cannot take place without the direct and strong link between the two groups.

II. Work Finished at the University of California, Irvine

a. Spin Dependent Electron Scattering from Ferromagnetic Surfaces; The ΔT Expansion

During the first grant year, we developed and implemented the first spin dependent electron scattering calculations from magnetically ordered substrates, based on *ab initio* crystal potentials (provided kindly by Prof. A. J. Freeman and his colleagues) that have been subject to no adjustment whatsoever. These calculations provided a remarkably quantitative account of spin polarized electron diffraction data (SPLEED) reported by Waller and Gradmann, over a wide range of energies. This analysis was described in last year's progress report, and in our view constitutes a major advance. We are now able, for the first time, to carry through quantitative calculations of spin dependent electron/surface interactions, in a parameter free manner (see item (b) below). Earlier work by R. Feder employed potentials with adjustable parameters, and as a consequence were often confined to selected energies. Analyses by H. L. Davis are impressive in the account they provide of spin orbit scattering from non-magnetic substrates.

The calculations, however, are quite costly in computer time, if one wishes to explore the scattering produced by various models of the magnetic moment distribution near the surface.

During the past grant year, we have developed a new perturbation theoretic approach that appears very accurate, and enables us to explore a wide range of parameter space quickly and accurately.

The approach is to carry out full calculations for the case where the magnetic moments assume their bulk magnitude and orientation everywhere, including the surface. Then if Δt_ℓ is the change in single site t matrix produced by changing moments in layer ℓ by $\Delta\mu_\ell$,

$$f = f_\infty + \sum_\ell \left(\frac{\partial f}{\partial t_\ell} \right) \Delta t_\ell$$

where f_∞ is the scattering amplitude for the "bulk moment everywhere" model.

The above scheme has been demonstrated to reproduce the results of full calculations very accurately. Furthermore, it is far faster. We can now scan a wide range of models of surface magnetic structure quickly and efficiently, as well as wide ranges of energy and angle. We have obtained superb and complete accounts of the extensive data set generated by Waller and Gradmann by this means.

Our earlier study of SPLEED data by Waller and Gradmann examined only four beam energies, while the full data set contains information on dozens of energies. We found earlier, unfortunately, that only one of the four energies chosen by us contains intensities sensitive to the magnetic moment magnitudes near the surface. We were interested in locating other "windows of sensitivity", and our new ability to perform spin polarized scattering calculations quickly has allowed us to explore sensitivity of the entire data set to moment distribution.

Our results are described in a forthcoming paper.⁽¹⁾

1) Alim Ormeci, Burl M. Hall, and D. L. Mills, "Sensitivity of SPLEED Exchange Asymmetries to Surface Magnetic Moment; The Case of Fe (110)", Phys. Rev. B (in press).

b. Theory of Spin Asymmetries in the Transmissivity of Electrons through Ultra Thin Films; The Question of the Spin Dependence of the Electron Mean Free Path in Ferromagnets

A major goal of our present program is to develop a fully quantitative theory of spin dependent electron scattering from magnetic surfaces, and surfaces of ultra thin magnetic films. Spin dependent losses and elastic scatterings are to be studied.

If our program is to be successful, and if we are to have the capability of analyzing data within models that contain few adjustable parameters, substantial improvements in the state of the art are required. Earlier workers employ crystal potentials whose spin dependent pieces are adjusted, with parameters that depend on beam energy. Our work, as described in subsection (a), utilizing excellent potentials supplied by the Freeman group, has demonstrated one may make contact with data in a fully quantitative manner with parameter free potentials, over a wide range of beam energy and angle.

There is another issue. Earlier workers have argued that a critical ingredient that must be present in the theory of spin dependent electron-surface scattering from ferromagnetic surfaces is a spin dependent mean free path. This is another source of uncertainty in the inputs required before theory can confront data.

An experimental group at Irvine headed by Prof. Herbert Hopster has claimed to measure the variation with energy, and the spin asymmetry of the electron mean free path in ferromagnetic Fe. This is done by depositing an ultra thin film of Fe (four atomic layers) on Cu(100), and photoemitting electrons from the Cu 3d bands. These must then propagate through the Fe film before reaching the detector.

We have analyzed this experiment within the framework of our spin dependent multiple scattering theory, and we find, contrary to the interpretation offered by Hopster *et al.*, it is not necessary to invoke any spin dependence in the mean free path to interpret the data. A fully quantitative account may be obtained *by using standard spin independent electron attenuation lengths known from earlier LEED analyses*. The effect observed by Hopster *et al.* instead has its origin in the spin asymmetry of the elastic forward scattering amplitude, which describes electron propagation through the ultra thin film.⁽¹⁾

These results imply that one may account for spin dependent electron-surface interactions in a fully quantitative manner, through use of (i) unadjusted potentials, and (ii) complex inner potentials (spin independent) taken from the LEED literature. This means we are now in a position to carry out quantitative studies of electron probes of magnetic surfaces and ultra thin films, within a framework where the only adjustable parameters relate to the nature of magnetism at the surface.

1) M. P. Gokhale and D. L. Mills, Phys. Rev. Letters **66**, 2251 (1991).

c. Theory of Surface Plasmon Excitation on Simple Metals, in Electron Energy Loss

In the past two years, the electron energy loss method has proved of central importance in the study of the dynamical response of electrons in the outermost atomic layers of metals. The experiments probe certain collective modes of electrons localized at surfaces, known as surface plasmons.

Recent data from Rocca and his colleagues in Genoa calls into question the standard theory of dipole excitation of plasmons. At issue is a rather complex dependence of the excitation cross section on wave vector transfer.

During the last grant year, we have completed the first quantitative study of the energy and angle variation of the surface plasmon excitation cross section, to obtain a fully quantitative account of the data. A key to our analysis is full inclusion of an interference effect in the scattering *amplitude* associated with surface plasmon excitation. As the electron approaches the crystal surface, it may excite the plasmon before striking the crystal surface, thus reflecting off at the scattered energy. Conversely, it may reflect from the surface at the incident energy, then excite the mode as it recedes from the surface, on the outgoing leg of its trajectory. The resulting excitation cross section depends on the relative *phase* as well as amplitude of the two electron surface reflection coefficients just described.

Our results are described in a paper that appeared recently.⁽¹⁾

1) Burl M. Hall and D. L. Mills, Phys. Rev. B**44**, 1202 (1991).

III. Joint Paper Between UC Irvine and UW Milwaukee

a. The Study of Surface Phonons by Electron Energy Loss Spectroscopy: Theoretical and Experimental Considerations

Mills, Tong and Black have reviewed the state-of-the-art applications of the HR-EELS technique.⁽¹⁾ The development of this important area is closely linked to the theoretical predictions made in our earlier publications based on research supported by this DOE joint program. High-resolution EELS is a powerful and versatile technique. It has already provided modern finger-printing services to organic and molecular chemists, as well as to researchers in the field of heterogeneous catalysis.

1) D. L. Mills, S. Y. Tong and J. E. Black, *Springer Series in Surface Sciences*, eds. W. Kress *et al.*, vol. **21**, 193 (1991).

IV. Work Finished at UW Milwaukee

a. Auger Electron Angular Distributions from Surfaces: Forward Focusing or Silhouettes?

We have studied the behavior of Auger and photoexcited electrons⁽¹⁾ as they are forward-scattered by atoms in the surface region. The scattering produces a forward-focusing effect, which acts as a search-light seeking out crystallographic directions inside a solid. This technique provides a rare worm's eye-view of buried atoms in the interface region. In this paper, we pointed out the difference between forward-focusing (the correct interpretation of data) and forward-shadowing, which was suggested by another research group.

1) X. D. Wang, Z. L. Han, B. P. Tonner, Y. Chen, S. Y. Tong in *Science*, vol. 248, p. 1129 (1990).

b. Surface Phonons and Structure of Epitaxial Nickel Layers on Cu(001)

New vibrational modes in epitaxially grown thin films on a Cu(001) substrate are predicted and their existence are confirmed by experimental measurements.⁽¹⁾ The new measurements and accompanying theoretical interpretation also elucidate the novel properties of these localized modes.

1) Y. Chen, S. Y. Tong, J. S. Kim, M. H. Mohamed and L. L. Kesmodel, *Phys. Rev. B* **43**, 6788 (1991).

c. Surface Acoustic Waves and Phonons

We have written a chapter⁽¹⁾ for the Landolt-Boernstein Volume on *The Physics of Solid Surfaces*. The volume is an important reference book in which the best measurements in the subject area are included and critically reviewed. Also, we provide a succinct description of the progress of the field to date.

1) A. A. Maradudin, R. F. Wallis, S. Y. Tong in *Physics of Solid Surfaces*, ed. G. Chiarotti, Landolt-Boernstein, Berlin (1992).

d. High-Resolution Electron Energy-Loss Spectroscopy Analysis of Ag(001): Discovery of a New Surface Longitudinal Mode Using First-Principles Phonon Calculations

We introduce the first analysis of high-resolution electron energy loss spectroscopy (HREELS) of vibrational modes of Ag(001) using normal mode frequencies and displacements obtained from first-principles microscopic calculations.⁽¹⁾ The first-principles phonon calculations include the full microscopic electron screening at the surface and contain no adjustable parameters. The combined method of using eigensolutions from first-principles calculations and fully multiple scattering inelastic cross section calculations represents the most sophisticated

theoretical treatment of this kind to date. The use of accurate theoretical treatment and experimental data, taken at a range of incident energies, is responsible for the discovery of a new localized longitudinal surface phonon at $\bar{M}$. This new mode has a large cross section at specific incident energies, and this energy range is determined.

1) Y. Chen, S. Y. Tong, M. Rocca, P. Moretto, U. Valbusa, K. P. Bohnen and K. M. Ho, *Surface Sci. Lett.* **250**, L389 (1991).

e. Forward-Focusing Photoelectron Diffraction—A New Structural Tool for Surfaces and Interfaces

We present the use of forward-focusing in *electrons* and forward-shadowing in *positrons* in Cu and Co films.⁽¹⁾ This technique allows layer-by-layer growth to be monitored. The presence of forward-focusing peaks in a system is evidence that an emitting atom is buried one, two or more atomic layers below the surface of a film. The technique thus offers a view of the surface structure from the vantage point of a buried atom.

1) S. Y. Tong, Y. Chen, Hua Li and C. M. Wei, in *Surface Physics*, Gordon and Breach, London (1991).

f. Characterization of Surface Phonons on Cu(001) and Ag(001): First-Principles Phonon Calculations With Experimental and Theoretical Studies of High-Resolution Electron-Energy Loss Spectra

Parameter-free first-principles phonon calculation is combined with multiple scattering electron-energy loss spectra calculation to characterize surface phonons of Cu(001) and Ag(001).⁽¹⁾ Again, the more accurate theory leads to the discovery of two new surface modes on these surfaces, and the inelastic electron scattering cross sections of these modes are investigated. With the discovery of the new modes, and knowledge about their inelastic cross section behavior, data of He-atom scattering and inelastic electron scattering are very satisfactorily explained. A comprehensive picture is established for the surface phonons on Cu(001) and Ag(001).

1) Y. Chen, S. Y. Tong, J. S. Kim, L. L. Kesmodel, T. Rodach, K. P. Bohnen and K. M. Ho, *Phys. Rev. B* **44**, Nov. 15 (1991).

g. Bonding and Structure on Semiconductor Surfaces

We have reviewed the nature of surface bonds on semiconductor surfaces.⁽¹⁾ Bulk-terminated semiconductor surfaces are invariably unstable because of the large number of unpaired dangling bonds which result from the creation of a sharp, idealized vacuum-solid interface (i.e., an ideal surface). To reduce the number of such unpaired bonds, surface atoms often rearrange in position to form new bonds having a higher electron density per bond. In many instances, elastic stress is introduced in the surface region. The balance between electronic and stress energies in the near surface region gives rise to the array of superlattices found

on reconstructed semiconductor surfaces of different materials: e.g., (2×1) , (2×2) , $(\sqrt{3} \times \sqrt{3})R30^\circ$, (7×7) , $c(2 \times 8)$, etc.

1) S. Y. Tong, H. Huang and C. M. Wei, in *Chemistry and Physics of Solid Surfaces*, vol. 8, eds. R. Vanselow *et al.*, Springer-Verlag, Berlin, P. 395 (1990).

V. Projects for the Coming Year *(removed)*

VI. Papers and Articles Published During The Second Grant Year

1. Alim Ormeci, Burl M. Hall and D. L. Mills, "Sensitivity of SPLEED Exchange Asymmetries to Surface Magnetic Moment; The Case of Fe(110)", *Phys. Rev. B* (in press).
2. M. P. Gokhale and D. L. Mills, *Phys. Rev. Letters* **66**, 2251 (1991).
3. Burl M. Hall and D. L. Mills, *Phys. Rev. B* **44**, 1202 (1991).
4. D. L. Mills, S. Y. Tong and J. E. Black, *Springer Series in Surface Sciences*, eds. W. Kress *et al.*, Vol. **21**, 193 (1991).
5. X. D. Wang, Z. L. Han, B. P. Tonner, Y. Chen, and S. Y. Tong, *Science*, **248**, 1129 (1990).
6. Y. Chen, S. Y. Tong, J. S. Kim, M. H. Mohamed, and L. L. Kesmodel, *Phys. Rev. B* **43**, 6788 (1991).
7. A. A. Maradudin, R. F. Wallis and S. Y. Tong, in *Physics of Solid Surfaces*, edited by G. Chiarotti, Landoldt-Boernstein, Berlin (1992).
8. Y. Chen, S. Y. Tong, M. Rocca, P. Moretto, U. Valbusa, K. P. Bohnen and K. M. Ho, *Surface Sci. Lett.* **250**, L389 (1991).
9. S. Y. Tong, Y. Chen, Hua Li and C. M. Wei, in *Surface Physics*, Gordon and Breach, London (1991).
10. Y. Chen, S. Y. Tong, J. S. Kim, L. L. Kesmodel, T. Rodach, K. P. Bohnen and K. M. Ho, *Phys. Rev. B* **44**, Nov. 15 (1991).

END

**DATE
FILMED**

3 / 23 / 92