DOE/

DOE/ER/13468--T1

DE92 007115

Stanford
Progress Report
DE-F603-86ER13468

**THE DYNAMICS OF ADSORPTION ON CLEAN AND ADSORBATE-MODIFIED
TRANSITION METAL SURFACES**

1989-1991

**A Proposal for Continued Support
to the U.S. Department of Energy
Division of Chemical Sciences
Office of Basic Energy Sciences**

from
**Robert J. Madix
Department of Chemical Engineering
Stanford University
Stanford, CA 94305**

(415) 723-2402

Start Date: January 1, 1992

Submitted June 1991

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

85

TABLE OF CONTENTS

Abstract	iii
I. Introduction	1
II. Summary of Previous Work Accomplished	2
II. A. Previous Work Accomplished: Overview	2
II. A.1. Molecular Adsorption of Alkanes on Pt(111)	2
II. A.2. Adsorption of Xenon on Pt(111)	3
II. A.3. Adsorbate-Assisted Adsorption	4
II. A.4. Activated Dissociative Adsorption of Alkanes on Pt(111)	5
II. B. The Unique Role of Molecular Beams	7

ABSTRACT

Continuation of support is requested for a three year period for research on both nondissociative and dissociative adsorption of gases on transition metal surfaces, with emphasis on alkanes. Particular attention will be placed on understanding the effects of adsorbed species on adsorption probabilities and on clarifying the role of energy exchange processes at the surface in both dissociative and nondissociative adsorption. Molecular beam methods will be coupled with methods of surface science to gain the greatest control of the variables of the problem, including the incident kinetic energy, vibrational energy and angle of incidence of the incoming molecules. The dynamics of both direct and precursor-influenced dissociative adsorption routes will be examined. Surface intermediates formed by activated adsorption will be identified by vibrational spectroscopy and temperature programmed reaction. Emphasis will be on comparative studies of low molecular weight alkanes in order to understand the origin of the differences in their reactivities on different metals.

I. INTRODUCTION

The work proposed herein is directed toward understanding the dynamics of bond activation in reactive collisions at surfaces. Such processes are fundamental to heterogeneous catalysis. Bond activation can occur either by formation of bonds between the incident molecule and the surface during impact (direct activation) or via trapping into a molecularly adsorbed state on the surface^{1,2} which acts as a precursor for dissociation.^{3,4} Activated adsorption is currently poorly understood, yet it is a crucial step in many heterogeneously catalyzed reactions important in the energy field, including the activation of CO and CO₂ in methanation on nickel,^{5,6,7,8} the activation of alkanes and H₂O in steam reforming⁹ and N₂ activation in the ammonia synthesis over iron based catalysts.¹⁰ Even though catalytic methanation over nickel catalysts involves the reaction of CO with dihydrogen, CO is not easily dissociated on the low index planes of nickel at room temperature¹¹ as it is on iron,¹² for example. However, dissociation is seen at elevated temperatures on polycrystalline and single crystal nickel,^{13,14} indicating that activated adsorption occurs to produce the dissociated state. Further, alkanes do not dissociatively adsorb on clean nickel at low kinetic energies under ultrahigh vacuum conditions, but dissociation is probably required for the formation of dihydrogen and carbon dioxide in the nickel-catalyzed steam reforming process.

Since activated adsorption is fundamental to heterogeneous catalysis, it is important to gain an understanding of its mechanisms on clean surfaces and the way in which it is affected by adsorbed species. In principle, all modes of energy in the molecule, including translation, vibration and rotation, and the vibrational modes of the surface and near-surface region can influence the probability of direct activation or the probability of adsorption into the precursor state. The purpose of this research is to advance our understanding of the role and

significance of these dynamical factors in adsorption in both clean and adsorbate-covered metal surfaces. Specific studies to be performed concern alkane activation on clean and modified metal surfaces.

II. SUMMARY OF PREVIOUS WORK ACCOMPLISHED

II. A. Previous Work Accomplished: Overview

Since collisional bond activation can proceed either upon impact with the surface or via a weakly adsorbed molecular (precursor) state, it is important to understand the dynamics of both direct collisional dissociative adsorption and molecular (nondissociative) adsorption. The overall scientific purpose of the work proposed here is to understand directly activated and precursor-assisted adsorption by coupling the modern tools of surface science with molecular beam technology to study gas-surface reactions, including alkane activation on clean and modified well-defined single crystal metal surfaces. Since this proposal involves, in part, a continuation and extension of research conducted under the previous DOE grant, a brief summary of the work accomplished during the last grant period in my laboratory is presented next. A more extended presentation of these results and a list of the publications arising from the work during the previous DOE grant period is given in Appendix 1. The proposed work follows in section III.

II. A.1. Molecular Adsorption of Alkanes on Pt(111)

Since precursor state kinetics may be important in the activation of alkanes on metal surfaces,^{15,16} we have focused attention on the dynamics of molecular adsorption of methane,¹⁷ ethane,¹⁸ and propane¹⁹ on Pt(111). Moreover, Pt(111) has been a surface of choice in trajectory simulations for nonreactive scattering events, and computer codes exist that are potentially

useful for trajectory simulations of alkane adsorption. At moderate kinetic energies the nondissociative adsorption probabilities of ethane and propane on Pt(111) were found to decrease monotonically as $E_T \cos^{0.6} \theta$, where E_T is the total kinetic energy of the molecules and θ is the angle of incidence measured from the surface normal. This result implies substantial involvement of momentum exchange parallel to the surface in the adsorption process due to significant roughness (corrugation) of the alkane-Pt(111) surface potential parallel to the surface plane. Similar results have been reported for ethane adsorption on Ir(110).²⁰ This result is a marked deviation from the predictions of simple one-dimensional theories and makes it clear that the prediction of adsorption probabilities requires a more sophisticated theoretical treatment that accurately incorporates a realistic gas-surface potential energy function.

The nondissociative adsorption probabilities of methane,²¹ ethane²² and propane²³ are nearly 1.0 on clean Pt(111) at low energies and decrease with increasing translational energy. At a given translational energy the adsorption probabilities increase with molecular weight of the alkane. This trend can be rationalized on the basis of increasing loss of translational energy of the alkane in the gas-surface collision due to both the increasing molecular weight and the increasing attractive potential between the alkane and the surface.

II. A.2. Adsorption of Xenon on Pt(111)

The adsorption probability of xenon on Pt(111), chosen as a model system, clearly shows a dependence on parallel momentum at hyperthermal energies. Experiments were performed with xenon because xenon scattering could be simulated with trajectory calculations.²⁴ In order to fit the dependence of the adsorption probability on incident energy and angle, trajectory simulations required a three parameter Xe-Pt potential function to obtain the

correct combination of surface corrugation and steepness of the repulsive interaction.²⁵ Thus, even for a large rare gas, one of the simplest adsorption systems imaginable, we observed a measurable corrugation in the potential parallel to the surface. Others have observed slightly different behavior in detail,²⁶ but the general conclusion regarding surface corrugation is the same.

II. A.3. Adsorbate-Assisted Adsorption

In heterogeneously catalyzed reactions occurring in the steady state under practical conditions the surface may be partially covered with molecular fragments as well as by the adsorbed reactant itself. The presence of such species may lead to efficient trapping of incident molecules on the surface and the formation of extrinsic precursor states - commonly viewed as molecules which are induced to weakly adsorb by the adsorbate, and may move easily over the surface to reactive centers. For example, it has been shown that ethylidyne is readily formed from ethylene on Pt(111)²⁷ and supported platinum catalysts²⁸ but that it is not readily hydrogenated to ethane.²⁹ Activation of species via adsorption into the precursor state may thus occur either on clean surfaces or surfaces covered with adsorbed species. Precursor dominated adsorption causes strong deviations from Langmuir kinetics and can be quite important even for weakly adsorbed molecules.

We have found that the nondissociative adsorption probabilities of ethane,³⁰ propane³¹ and xenon³² increase significantly on Pt(111) with surface coverage (figure 1). Qualitatively, this difference can be understood by simple classical mechanics. Due to the better mass matching between the colliding partners, ethane exchanges momentum and energy much better with itself than with the heavier platinum atom. We have observed that the enhancement in adsorption caused by the adsorbed species depends linearly on the surface coverage. For ethane³³

Figure 1. The initial adsorption probability for ethane on clean and ethane-covered Pt(111) at 95 K plotted against the incident translational energy of the molecules in the molecular beam (beam at normal incidence).

and xenon³⁴ the coverage dependence of the net adsorption probability is given by

$$S_T = S_0 \theta + S_0' (1 - \theta)$$

where S_T , S_0 and S_0' are the net, clean and covered surface adsorption probabilities and θ is the fractional surface coverage. This relationship is a limiting case of a more complex form derived on the basis of the existence of extrinsic precursor states, but in this simple form it accounts well for the basic observations and gives an excellent fit to the results. Similar results have been obtained by others for ethane on Ir(110).³⁵

When the surface is covered with a monolayer of adsorbate, the sticking probabilities decrease with increasing total translational energy, E_T , and are independent of the angle of incidence, θ , suggesting the presence of a strong static and/or "dynamic" corrugation in the scattering potential.³⁶ These experiments probe dynamical effects in adsorption on surfaces covered by molecules or molecular fragments and are important to the understanding of the dynamical basis of extrinsic precursor states, which may be very important in the kinetic description of catalytic reactions proceeding on surfaces partially covered with adsorbed molecular fragments.

II. A.4. Activated Adsorption of Alkanes on Pt(111)

Since alkane activation on platinum is a problem of substantial importance in catalysis, in the 1978 grant period we have studied methane and ethane activation on Pt(111). The long range goal of this work is the understanding of differences in reactivity of alkanes of differing molecular weight and structure. Both methane and ethane are readily dissociatively adsorbed on Pt(111) by direct collisional activation; reaction probabilities of

the incident alkanes increase markedly with increasing kinetic energy. In the range of kinetic energy studied the probability of dissociative adsorption depends exponentially on the "normal" kinetic energy, $E_T \cos^2 \theta$. Unit changes in the kinetic and vibrational energies produce nearly equal changes in the reaction probability for ethane at the lower kinetic energies studied. Though the reaction probabilities of the two alkanes overlap over the range of kinetic energy from 70 to 90 kJ/mol, the apparent barrier height for methane dissociation is lower than that for ethane, and the sensitivity of the reactivity of methane to increasing translational energy is larger than for ethane (figure 2).^{37,38} Our results for methane agree well with concurrent studies of others over the common energy range.³⁹ This behavior is qualitatively similar to that observed for the dynamics of methane activation on Ni(111) and W(110) (see below) where it has been proposed that tunneling of hydrogen through the potential barrier to dissociation is involved. Overall, however, the differences in the reactivities between methane and ethane are not understood and form the basis for a portion of the proposed work.

In addition to the fundamental insights gained from this research, general observations of potential use in catalysis emerge from the studies of the previous grant period. First, the adsorption probabilities of alkanes at kinetic energies below 5 kJ/mol are greater than 0.5, at least for low surface temperatures. Even at kinetic energies nearly as large as the binding energy of the molecule to the surface, the adsorption probabilities are greater than one tenth. Second, weakly adsorbed species greatly enhance adsorption. Combined, the observations indicate that adsorption probabilities of moderately large molecules should be high over a wide range of kinetic energies on metal surfaces. One of the objectives of the proposed research is to test the applicability of these trends under conditions more characteristic of the catalytic steady state.

Figure 2. Dissociative adsorption probability of methane and ethane on Pt(111) at 500 K plotted against the normal kinetic energy. The nozzle temperatures were 993 K and 947 K, respectively.

II. B. The Unique Role of Molecular Beams

Molecular beam techniques were essential to the discoveries outlined above. In each of these studies a nearly monoenergetic molecular beam of the desired gas was directed onto the single crystal surface, which had been previously cleaned and characterized by low energy electron diffraction and Auger electron spectroscopy.⁴⁰ The adsorption probability was determined from the change in partial pressure of the gas of interest in the scattering chamber when the crystal was suddenly exposed to the beam and adsorption commenced (see Appendix).^{41,42}

Nozzle beams allow the effects of translational and vibrational energies of the adsorbing gas on adsorption or surface reaction to be studied separately. Kinetic energies between about 2 and 400 kJ/mol can be obtained by heating the beam source and/or seeding the desired gas in dilute concentration into a light carrier gas,⁴³ depending on the molecule of choice and the carrier gas. Vibrational energy is controlled by changing the temperature of the beam source in order to vary the Boltzmann population of vibrational levels in the molecule. Independent variation of the population of the vibrational states of the molecules in the beam and the translational energy is effected by holding the nozzle temperature constant and varying the concentration of the desired molecule in the carrier gas.

The angular distribution of the molecules reflected from the surface can also be utilized to obtain important information concerning adsorption. Generally, on clean surfaces with the surface in which the molecules rebound, exchanging energy with the surface but without adsorbing and accommodating to the surface temperature (direct inelastic collisions), give angular distributions which peak near the specular angle of scattering (figure 3, 45 degrees). Adsorbed species which are desorbing in the steady state give angular distributions in flux peaked at the

surface normal (figure 3, zero degrees) due to the loss of memory of the incident momentum of the desorbing molecule.⁴⁴ The results depicted on figure 3 clearly show that at a kinetic energy of 15 kJ/mol ethane exhibits both adsorption-desorption and direct-inelastic scattering, whereas at 78 kJ/mol there is very little, if any, adsorption-desorption, and the collisional encounter is dominated by direct-inelastic scattering from the surface.⁴⁵ In favorable cases adsorption probabilities can be estimated from angular distributions.⁴⁶

Figure 3. Angular distributions of ethane scattered from Pt(111) at 500 K for beams of 15 and 78 kJ/mol incident kinetic energies.

END

**DATE
FILMED**

3 / 10 / 92