Lawrence Livermore Laboratory

TRENDS IN OTHER SPECIES

Joyce E. Penner

March 1979

NOTE: This is a position paper prepared for the NASA Stratospheric Workshop to be held in Reston, Virginia, June 4-8, 1979.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. Introduction

The predictions of stratospheric models depend on accurate knowledge of the background concentrations of many different species and of the processes which determine their concentrations in the stratosphere. It is important to try to test stratospheric models by careful comparison of predicted and measured changes. Of primary interest here is the verification of the effects of increases in the concentrations of chlorofluoromethanes on ozone. While the predicted changes in total ozone are probably too small to detect at this time, changes at higher altitude (around 40 km) are expected to be larger, and one would hope to detect the first sign of predicted decreases in ozone there.

Unfortunately, the effects of other processes operating on the ozone concentration at high levels may hinder our ability to detect the CFM-caused changes. One such process is long-term solar flux variations in the 1800-3400 Å region. Only a few sets of measurements showing a variation are available and these need to be confirmed (Heath and Thekackara, 1977; Simon, 1978), but model calculations of the resulting variations in ozone and temperature appear to be fairly consistent with measurement (Penner and Chang, 1978; Callis et al., 1979).

It is desirable to try to confirm the effects of chlorofluoromethanes and solar variability by looking at the long-term trends expected for trace species other than ozone. Unfortunately, data which might support long-term changes in stratospheric species other than ozone either do not exist or are difficult to interpret. Few measurements of any kind are available at high altitude in the stratosphere, and in the lower

Trends in Other Species

Joyce E. Penner

Lawrence Livermore Laboratory, University of California
Livermore, California 94550

ABSTRACT

Predicted and measured trends in trace species other than ozone are compared and current knowledge regarding man's role in the budget for several species is discussed. Possible trends in stratospheric species caused by long-term solar uv variability are compared to trends detected to date or to predicted trends caused by anthropogenic emissions.

This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore Laboratory under contract number W-7405-ENG-48.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

209

stratosphere concentrations of many constituents are thought to be controlled to a large extent by dynamics, so that changes from time to time or place to place may only indicate the presence of air masses originating from different regions at the times of measurement, and not a true trend in species concentration.

In this paper we review our knowledge regarding possible and expected trends in stratospheric trace species and compare model predictions with measurement. We first consider the halocarbons, since the observed increase in tropospheric concentrations should have led to an increase in the stratospheric burden of chlorine species. CO and CO₂ are also expected to have increased due to man's increasing use of fossil fuels and we discuss the evidence and theories for trends in these species concentrations next. Then we briefly review current thinking on possible trends in N₂O, CH₄ and H₂O whose concentrations may have increased in the past. Finally, we discuss the trends expected for species (other than ozone) from solar flux variations. As we shall see the predicted changes from uv flux variations appear to dominate the other trends discussed herein at high altitude.

II. CFCl_3 and CF_2Cl_2

Given the emission rates for CFCl_3 and CF_2Cl_2 (McCarthy et al., 1977), one expects increasing concentrations for these species both in the troposphere and stratosphere.

Tropospheric measurements of CFCl_3 date from 1971 (Lovelock et al., 1973). Its concentration has apparently been increasing steadily since then. Pack et al. (1977) have analyzed the CFCl_3 data of Lovelock and coworkers and deduce a growth rate of 14.6%/year in the southern hemisphere and 13.2%/year in the northern hemisphere. Until 1974 the emission rate for CFCl_3 was increasing by approximately 13.9%/year (NAS, 1976). Thus the rate of increase of CFCl_3 as determined from Lovelock's data appears to be consistent with the emission data. The absolute concentrations of Lovelock, however, are smaller than those measured by several other groups (e.g. Rasmussen, et al., 1976; Singh, et al., 1979), and the inferred inter-hemispheric gradient is larger. Rasmussen and coworkers have been measuring CFCl_3 since early 1975 and Singh began taking measurements in November 1975. The data of Lovelock imply a "best estimate" for the atmospheric lifetime of CFCl_3 of 15-20 years. The data of Rasmussen imply a lifetime of 50-60 years, consistent with the lifetime determined by photolysis in the stratosphere. (Compare Rowland and Molina, 1976; Sze and Wu, 1976; Chang and Penner, 1976; Meakin et al., 1978.) The discrepancies have complicated the problem of determining whether any tropospheric sinks, which so far have not been included in the models, might exist. The estimates of lifetime are of course very crude and one needs good global coverage in order to obtain accurate estimates. For these reasons experiments continue with

the Atmospheric Lifetime Experiments sponsored by the M&A in cooperation with both Rasmussen and Lovelock's groups.

Tropospheric measurements of CF_2Cl_2 were first made in late 1973 by Lovelock (1974). The stated uncertainties for measurement of CF_2Cl_2 are generally larger than for CFCl_3 . Thus the problems of determining accurate global atmospheric concentrations are compounded for this species. The tropospheric lifetime estimated by Singh et al. (1979) from their measurements is 65 to 70 years, consistent with that estimated by models assuming that stratospheric photolysis is the only sink.

In situ stratospheric measurements of CFCl_3 and CF_2Cl_2 date from 1973 (Heidt et al., 1975). The available measurements are fewer than in the troposphere and the natural variability larger, making global abundance estimates even more difficult. Calculated model abundances appear to be reasonably consistent with the measured concentrations (see, for example, Logan et al., 1978), and a retroactive comparison of 1968 infrared data with a detection made in 1975 showed that both CFCl_3 and CF_2Cl_2 increased by more than a factor of 2 in the stratosphere in that period (Murray, et al., 1975).

III. CCl_4

According to the analysis of Pack et al., (1977), the average rate of growth of emissions of CCl_4 was about 3.9%/year. Their measured concentrations of CCl_4 , which date from January 1972 to the end of 1975, increased by 2.2%/year in the southern hemisphere and by 10.5%/year in the northern hemisphere. The tropospheric concentrations measured by Singh et al. (1979), which date from late 1976 increased by about 2%/year. According to the analysis of Singh et al. (1979), the CCl_4 distribution and budget is consistent with the man-made source though their atmospheric burden (121 ppt) is somewhat larger than that derived from all known sources (see also Singh et al., 1976; Altshuller, 1976; Galbali, 1976).

Much of the data for CCl_4 shows a considerable amount of scatter both from time to time, due possibly to measurement in polluted air masses (see Pack et al. 1977), and from researcher to researcher, indicating possible calibration problems. Thus the budget for CCl_4 needs to be better established.

CH_3CCl_3

The use and emission of CH_3CCl_3 has been increasing at an average rate of about 19%/year according to the analysis of emission rates by Chang and Penner (1978). CH_3CCl_3 is thought to be removed by reaction with OH in the troposphere, and by photolysis and reaction with OH in the stratosphere. The contribution of CH_3CCl_3 to the stratospheric burden of CLO_x depends on how efficiently CH_3CCl_3 is removed by reaction with OH in the troposphere. Estimates of the tropospheric lifetime from budget

considerations yield lifetimes ranging from 5 to 11 years (Lovelock, 1977; Singh, 1977; Chang and Penner, 1978). This implies average tropospheric OH concentrations which are consistent with some model estimates but are smaller than previously thought, (Crutzen & Fishman 1977). However, detailed comparison of model predicted OH measurements with measured concentrations are not always consistent (Philen et al., 1978). Thus, further study is needed to clarify the role of OH removal of CH_3CCl_3 in the troposphere.

IV. CO₂

The increased concentrations of CO₂ are well documented. Systematic measurements taken at Mauna Loa and the South Pole are available which date from 1958 (Keeling et al., 1976a; Keeling et al., 1976b), and some earlier measurements date from about 1870 (see summary by Callender, 1958). The SCEP and SMIC Reports deduce a rate of increase in atmospheric CO₂ of approximately 0.2%/year, or 0.64 ppm out of 320 ppm during the early 1960's. The excess of the annual average concentration of CO₂ over the preindustrial value has grown at the rate of 4%/year, which approximately equals the rate of increase of the cumulative amount of CO₂ produced by the burning of fossil carbon. The total amount of CO₂ accumulated to date in the atmosphere, however, is only half as large as the amount that has been released by combustion of fossil fuels. Apparently, some of the released CO₂ has been stored in the oceans and/or the land biota. The quantification of the ocean or biota sink is necessary in order to calculate atmospheric levels of CO₂ in the future, but several studies predict concentrations rising to between 375 and 400 ppm by the year 2000. Predictions beyond the year 2000 are quite uncertain but concentrations may increase to 4 to 8 times their present levels over the next 200 years.

Increases in stratospheric CO₂ are estimated to cause decreases in stratospheric temperature and therefore increases in O₃. Doubling CO₂ may increase column O₃ by between 1.2 and 6% according to several estimates (Luther et al., 1977; Groves et al., 1978; Boughner, 1978). The models predict an increase in O₃ to date of less than 2% over preindustrial values.

CO

If one assumes the emissions of CO from fossil fuel combustion have been increasing at the same rate as emissions of CO₂, the increase in global average CO concentrations near the surface should have been about 40% since 1940 (Sze, 1977; Fenner et al., 1977; Logan et al., 1978). It is known, however, that CO is highly variable with latitude (Seiler, 1974), so that detection of such a small global increase in CO is difficult. In fact the currently measured CO concentrations appear to be consistent with those measured by Seiler in 1968 (Heidt and Krassec, 1978).

V. H_2O

All of the species trends discussed thus far have been associated with tropospheric concentrations. While long-term tropospheric trends should be associated with similar trends in the stratosphere, direct evidence for trends in stratospheric concentrations is much more difficult to establish because of the remoteness of the stratosphere, because of the short time over which many stratospheric species have been measured, and because many species exhibit strong concentration gradients at the tropopause and have long lifetimes in the lower stratosphere leading to a high degree of spatial variability. There is a large amount of uncertainty in stratospheric concentrations of H_2O due to scatter in the measured concentrations of H_2O (Harries, 1976). In addition concentrations of H_2O in the lower stratosphere apparently vary with latitude (Harries, 1976; Hilsenrath et al., 1977) and time. Mastenbrook (1971, 1974) made a series of measurements of stratospheric H_2O which suggested both a seasonal variation and a secular increase of about 33% for the period 1964 to 1973 with an abrupt drop thereafter (Penndorf, 1978). The physical mechanism responsible for these changes has not been established, though Mastenbrook concludes that there appears to be a correspondence between the observed stratospheric midlatitude mixing ratio and the temperature or saturation mixing ratio at the tropical tropopause.

N_2O

There has been a great deal of concern recently regarding the possible increase of nitrous oxide due to anthropogenic nitrogen fixation. A review

of measurements of N_2O has been given by Pierotti and Rasmussen (1977). Measurements before 1966 were scattered and imprecise, so that these data cannot be used to establish a trend. In 1966, however, Goody (1969) began systematic measurements of the total N_2O column over Milton, Massachusetts. The N_2O concentrations remained roughly the same through the end of 1967, but then increased by about 17% during 1968. Junge's laboratory in Germany made concurrent ground level measurements from November 1966 to May 1969 in Mainz and from June 1967 to December 1968 at Schauinsland (Schütz et al., 1970) and they report a similar increase. The trend measured for 1967-1968 did not apparently continue, but a search for sources and sinks which might explain the measured behavior has been undertaken by several groups. The distribution of N_2O today appears to be fairly constant in both latitude and time (Rasmussen and Perotti, 1978), leaving the 1967-1968 measurements unexplained.

CH₄

Recently Seiler et al., (1978) have reported measured methane concentrations in the northern hemisphere ranging from 1.64 to 1.82 ppmv, with an average value of 1.7 ppmv. These concentrations were measured on aircraft flights over Europe in the 50°-60° latitude belt in November and December 1976. They are 20-30% higher than the methane concentrations reported previously. For example, Ehhalt and Heidt (1973) obtained an average value of 1.41 ppmv in aircraft samples over North America in 1966 and 1967, while Heidt and Pollack report concentrations in excess of 1.5 ppmv for measurements made from 1974 to 1976 (see discussion in Ehhalt, 1978).

Ehhalt and Schmidt (1976) and Ehhalt (1974) have reviewed the budget for atmospheric methane and estimate the anthropogenic component of CH_4 production to be less than 20% of the total from all sources. Thus unless CH_4 release has increased at a rate close to 10%/year over the last 10 years, it is difficult to explain the measured increase as directly attributable to man's release of the gas. The trend needs further confirmation and study.

VI. Trends from UV Flux Variations

Figs 1 to 15 show expected changes in trace concentrations associated with variations in the solar uv flux. The calculation used the LLL 1-D radiative equilibrium—species concentration model, and perturbed the solar flux according to the fit of Penner and Chang (1978) model C to the data of Heath and Thekaekara (1977). The model is discussed in Penner and Chang (1979). As shown there, the predicted changes are consistent with the few stratospheric measurements that are presently available. Since the predicted changes in O_3 at high altitudes are large compared to CFM-caused changes, it is important to try to confirm the uv flux perturbations by examining changes in other species. These calculations can be used to help identify those species which will provide the clearest signal should uv flux variations occur over long (11-year) time scales. Clearly, it is desirable to confirm the solar uv variability, since the changes shown in Figs. 1-15 could easily mask the detection of stratospheric trends due to anthropogenic emissions for most of the species discussed in this paper.

References

- Altshuller, A. P., Average Tropospheric Concentration of Carbon Tetrachloride Based on Industrial Production, Usage, and Emissions, Environ. Sci. Technol. 10, 596-598, 1976.
- Boughner, R. E., The Effect of Increased Carbon Dioxide Concentrations on Stratospheric Ozone, J. Geophys. Res. 83, 1326-1332, 1978.
- Callender, G. S., On the Amount of Carbon Dioxide in the Atmosphere, Tellus 10, 243-248, 1958.
- Callis, L. B., N. Natarajan, and J. E. Nealy, Ozone and temperature trends associated with the eleven year solar cycle, Science, 1979 (in press).
- Chang, J. S. and J. E. Penner, Analysis of Global Budgets of Halocarbons, Atmos. Environ. 12, 1867-1873, 1978.
- Crutzen, P. J. and J. Fishman, Average Concentrations of OH in the Northern Hemisphere, and the Budgets of CH_4 , CO, H_2 , and CH_3CCl_3 , Geophys. Res. Lett. 4, 321-324, 1977.
- Ehhalt, D. H., The Atmospheric Cycle of Methane, Tellus 26, 58-70, 1974.
- Ehhalt, D. H., The CH_4 Concentration Over the Ocean and its Possible Variation with Latitude, Tellus 30, 169-176, 1978.
- Ehhalt, D. H. and L. E. Heidt, Vertical Profiles of CH_4 in the Troposphere and Stratosphere, J. Geophys. Res. 78, 5265-5271, 1973.
- Ehhalt, D. H. and U. Schmidt, Sources and Sinks of Atmospheric Methane, Pure and App. Geophys. 116, 452-464, 1978.
- Galbali, I. E., Man-made CCl_4 in the Atmosphere, Science 193, 573-576, 1976.

- Goody, R. M., Time Variations in Atmospheric N_2O in Eastern Massachusetts, *Planet. Space Sci.* 17, 1319-1320, 1969.
- Groves, K. S., S. R. Mattingly, and A. F. Tuck, Increased Atmospheric Carbon Dioxide and Stratospheric Ozone, *Nature* 273, 711-715, 1978.
- Harries, J. E., Distribution of Water Vapor in the Stratosphere, *Rev. Geophys. Space Phys.* 14, 565-575, 1976.
- Heath, D. F. and M. P. Thekaekara, The Solar Spectrum Between 1200 and 3000 Å in *The Solar Output and its Variation*, O. R. White, Ed. (Colorado Assoc. Univ. Press, Boulder, Colorado, 1977) p. 193.
- Heidt, L. E. and J. P. Krasnec, Tropospheric Distributions of CO , CH_4 , CO_2 and N_2O in the Northern and Southern Hemispheres, presented at the AGU Fall Meeting, San Francisco, California, 1978.
- Heidt, L. E., R. Lueb, W. Pollock, and D. H. Ehhalt, Stratospheric Profiles of CCl_3F and CCl_2F_2 , *Geophys. Res. Lett.* 2, 455-447, 1975.
- Hilsenrath, E., B. Guenther, and P. Dunn, Water Vapor in the Lower Stratosphere Measured from Aircraft Flight, *J. Geophys. Res.* 82, 5453-5458, 1977.
- Jesson, J. P., P. Meakin, and L. C. Glasgow, The Fluorocarbon-ozone Theory II: Tropospheric Lifetimes. An Estimate of the Tropospheric Lifetime of CCl_3F , *Atmos. Environ.* 11, 499-508, 1977.
- Keeling, C. D., J. A. Adams, Jr., C. A. Ekdahl, Jr., and P. R. Guenther, Atmospheric CO_2 Variations at the South Pole, *Tellus* 28, 552-564, 1975a.
- Keeling, C. D., R. B. Bacaston, A. E. Bainbridge, C. A. Ekdahl, Jr., P. R. Guenther, L. S. Waterman, and J. F. S. Chin, Atmospheric CO_2 Variations at Mauna Loa Observatory, Hawaii, *Tellus* 28, 538-551, 1976b.

- Logan, J. A., M. J. Prather, S. C. Wofsy, and M. B. McElroy, Atmospheric Chemistry: Response to Human Influence, Phil Trans. Roy. Soc. London 290, 187-234, 1975.
- Lovelock, J. E., Atmospheric Halocarbons and Stratospheric Ozone, Nature 252, 292-294, 1974.
- Lovelock, J. E., Methylchloroform in the Troposphere as an Indicator of OH Radical Abundance, Nature 267, 33, 1977.
- Lovelock, J. E., R. J. Maggs, and R. J. Wade, Halogenated Hydrocarbons in and over the Atlantic, Nature 241, 194-196, 1973.
- Luther, F. M., D. J. Wuebbles, and J. S. Chang, Temperature Feedback in a Stratospheric Model, J. Geophys. Res. 82, 4935-4942, 1977.
- Mastenbrook, H. J., The Variability of Water Vapor in the Stratosphere, J. Atmos. Sci. 28, 1495-1501, 1971.
- Mastenbrook, H. J., Water-Vapor Measurements in the Lower Stratosphere, Can. J. Chem. 52, 1527-1531, 1974.
- McCarthy, R. L., F. A. Bower, and J. P. Jesson, The Fluorocarbon-ozone Theory I: Production and release. World production and release of CCl_3F and CCl_2F_2 (Fluorocarbons 11 and 12) through 1975, Atmos. Environ. 11, 491-497, 1977.
- Meakin, P., Gunerman, P. S., Glasgow, L. C., and Jesson, J. P., The Fluorocarbon-ozone Theory III: Fluorocarbon Mixing and Photolysis. The Effects of Eddy Diffusion and Tropospheric Lifetime on CCl_3F and CCl_2F_2 Tropospheric Mixing Ratios, Atmos. Env. 12, 1271-1285, 1978.
- Murcay, D. G., F. S. Bonomo, J. N. Brooks, A. Goldman, F. H. Murcay, and W. J. Williams, Detection of Fluorocarbons in the Stratosphere, Geophys. Res. Lett. 2, 109-112, 1975.

- National Academy of Sciences, Panel on Atmospheric Chemistry, Halocarbons: Effects on Stratospheric Ozone, 1976.
- Pack, D. H. J. E. Lovelock, G. Cotton, and C. Curthoys, Halocarbon Behavior from a Long Time Series, Atmos. Environ. 11, 329-344, 1977.
- Penndorf, R. Analysis of Ozone and Water Vapor Field Measurement Data, Final Report to the U. S. Department of Transportation, Federal Aviation Administration, FAA-EE-78-29, November, 1978.
- Penner, J. E. and J. S. Chang, Possible variations in atmospheric ozone related to the eleven year solar cycle, Geophys. Res. Lett. 5, 817-820, 1978.
- Penner, J. E. and J. S. Chang, manuscript in preparation, 1979.
- Penner, J. E., M. B. McElroy, and S. C. Wofsy, Sources and Sinks for Atmospheric H_2 : A current analysis with projections for the influence of anthropogenic activity, Planet Space Sci. 25, 521-540, 1977.
- Pierotti, D. and R. A. Rasmussen, The Atmospheric Distribution of Nitrous Oxide, J. Geophys. Res. 82, 5823-5832, 1977.
- Philen, C., W. Heaps, and D. D. Davis, Boundary Layer and Free Tropospheric OH Measurements at Tropical and Sub-tropical Latitudes in the Northern and Southern Hemispheres, paper presented at AGU Fall Meeting San Francisco, California, December 1978.
- Rasmussen, R. A. and D. Pierotti, Global and Regional H_2O Measurements, Pure and App. Geophys. 116, 405-413, 1978.
- Rasmussen, R. A. Pierotti, D., Krasnec, J., and B. Holter, Trip report of the cruise of the Alpha Helix research vessel from San Diego, California to San Martin, Peru, March 5-20, 1976.

- Rowland, F. S. and M. J. Molina, Chlorofluoromethanes in the Environment, *Rev. Geophys. and Space Phys.* 13, 1-35, 1975.
- Rowland, F. S. and M. J. Molina, Estimated Future Atmospheric Concentrations of CCl_2F (Fluorocarbon-11) for Various Hypothetical Removal Rates, *J. Phys. Chem.* 80, 2049-2056, 1976.
- SCEP Report, C. L. Wilson and W. H. Mathews, Eds., Man's Impact on the Global Environment (The M.I.T. Press, Cambridge, MA, 1970).
- Schütz, K., C. Junge, R. Beck, and B. Albrecht, Studies of Atmospheric N_2O , *J. Geophys. Res.* 75, 2230-2246, 1970.
- Seiler, W., The Cycle of Atmospheric CO, *Tellus* 26, 116-135, 1974.
- Seiler, W., F. Müller, and H. Oeser, Vertical Distribution of Chlorofluoromethanes in the Upper Troposphere and Lower Stratosphere, *Pure and Appl. Geophys.* 116, 554-556, 1978.
- Simon, P. C., Irradiation Solar Flux Measurements Between 120 and 400 nm Current Position and Future Needs, *Planet. Space Sci.* 26, 355-365, 1978.
- Singh, H. B., Atmospheric Halocarbons: evidence in favor of reduced average hydroxyl concentration in the troposphere, *Geophys. Res. Lett.* 4, 101-104, 1977.
- Singh, H. B., D. P. Fowler and T. O. Peyton, Atmospheric Carbon Tetrachloride: Another Man-Made Pollutant, *Science* 192, 1231-1234, 1976.
- Singh, H. B., L. J. Salas, H. Shigeishi, and E. Scribner, Atmospheric Halocarbons, Hydrocarbons, and Sulfur Hexafluoride: Global Distributions, Sources, and Sinks, *Science* 203, 899-903, 1979.
- SMIC Report, C. L. Wilson and W. H. Mathews, Eds., Inadvertant Climatic Modification (The M.I.T. Press, Cambridge, MA, 1971).

- Sze, N. D., Anthropogenic CO Emissions: Implications for Atmospheric CO-H₂-CH₄ Cycle, Science, N.Y., 195, 673-674, 1977.
- Sze, N. D. and M. F. Wu, Measurements of Fluorocarbons 11 and 12 and Model Validation: An Assessment, Atmos. Environ. 10, 1117-1125, 1976.

Reference to a company or product names does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

NOTICE

"This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights."

Figures 1 - 15 show calculated percentage change for 15 species from maximum solar flux to minimum solar flux ($(\text{max}-\text{min})/\text{min} \times 100$) (after Penner and Chang, 1979).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Fig. 10

Fig. 11

Fig. 22

Fig. 13

Fig. 14

Fig. 15